══ R E V I E W C O M M O N S M A N U S C R I P T ══

IMPORTANT:

- Manuscripts submitted to Review Commons are peer reviewed in a journal-agnostic way.
- Upon transfer of the peer reviewed preprint to a journal, the referee reports will be available in full to the handling editor.
- The identity of the referees will NOT be communicated to the authors unless the reviewers choose to sign their report.
- The identity of the referee will be confidentially disclosed to any affiliate journals to which the manuscript is transferred.

GUIDELINES:

- For reviewers: https://www.reviewcommons.org/reviewers
- For authors: https://www.reviewcommons.org/authors

CONTACT:

The Review Commons office can be contacted directly at: office@reviewcommons.org

Development of an *in vivo* electroporation-based chromosomal engineering

technique in the murine uterine epithelium

Satoru Iwata[1, 2, 3, 4, *], Yumi Miura[1], and Takashi Iwamoto[1, 2]

[1]Center for Education in Laboratory Animal Research, Chubu University, Kasugai,

Aichi 487-8501, Japan

[2]Department of Biomedical Sciences, College of Life and Health Sciences, Chubu

University, Kasugai, Aichi 487-8501, Japan

[3]College of Bioscience and Biotechnology, Chubu University, Kasugai, Aichi 487-8501,

Japan

[4]Center for Mathematical Science and Artificial Intelligence, Chubu University, Kasugai,

Aichi 487-8501, Japan

Lead Contact

Satoru Iwata

* Correspondence: satoru_iwata@fsc.chubu.ac.jp

Abstract

Clustered regularly interspaced palindromic repeat (CRISPR)/CRISPR-associated

protein (Cas)-based *in vivo* chromosomal rearrangements are a promising approach for

generating model organisms with specific chromosomal abnormalities. However,

conventional *in vivo* methods rely on viral vectors, which are expensive, require

specialized equipment, and pose potential safety risks, thereby limiting their widespread

application. To overcome the limitations above, we developed a novel, efficient, and

cost-effective *in vivo* chromosomal engineering strategy using CRISPR

ribonucleoprotein electroporation for the murine uterine epithelium. Our method

successfully induced translocations at multiple loci and repaired a 57.8-Mb inversion.

The findings of the present study establish *in vivo* electroporation as a practical

alternative to traditional chromosomal engineering methods and provide a foundation

for its broader application in genome editing technologies.

Keywords

chromosomal rearrangements, *in vivo* electroporation, CRISPR/Cas9

Introduction

Chromosomal abnormalities, particularly inversions and translocations, play a crucial role in the development of various cancers and hereditary disorders. Chromosomal rearrangements not only alter gene expression and promote tumorigenesis by dysregulating oncogenes and tumor suppressor genes (Mitelman et al., 2007; Chen et al., 2010) but also induce mutations in genes involved in essential physiological functions, thereby causing infertility and various hereditary diseases (Lee et al., 2006; Feuk, 2010; Pellestor et al., 2011; Zepeda-Mendoza et al., 2019). The precise and efficient repair of such chromosomal abnormalities is essential for treatment, and the clustered regularly interspaced palindromic repeat (CRISPR)/CRISPR-associated protein (Cas) system is widely used for this process (Katti et al., 2022; Nambiar et al., 2022).

The CRISPR/Cas system has been used successfully to generate model organisms with specific chromosomal abnormalities (Blasco et al., 2014; Maddalo et al., 2014; Reimer et al., 2017). These models have contributed significantly to the elucidation of various disease mechanisms and the development of novel therapeutic strategies. Furthermore, newly developed prime editing techniques, such as WT-PE and PETI, exhibit high precision in inducing inversions and translocations in cultured human cells (Tao et al., 2022; Kweon et al., 2023). These techniques have great potential for the treatment of diseases caused by chromosomal rearrangements.

However, *in vivo* approaches of inducing chromosomal rearrangements using the CRISPR/Cas system often require a combination of viral vectors, which entails considerable effort, cost, specialized experimental facilities, and potential safety risks, thereby limiting their widespread application. Moreover, to the best of our knowledge,

no study has demonstrated successful induction of chromosomal rearrangements via

prime editing *in vivo* to date. Therefore, the development of simple, safe, and efficient

in vivo chromosomal rearrangement techniques is necessary to treat various genetic

disorders and cancers.

We have previously performed *in vivo* electroporation using RecQ-like helicase 5

(*Recql5*) mutations to efficiently induce complex chromosomal rearrangements (CCRs)

by delivering CRISPR ribonucleoproteins (RNPs) into fertilized mouse embryos in the

oviduct (Iwata et al., 2024). Nevertheless, the widespread application of our approach is

limited by restrictions on the use of fertilized embryos and ethical regulations

surrounding human embryo editing (Greely, 2019). In addition, our *Recql5*-based

approach revealed the involvement of replication-based complex rearrangement

mechanisms, such as chromoanasynthesis-like events, in the repair of double-strand

breaks (DSBs) associated with chromosomal rearrangements. Chromoanasynthesis is

characterized by extensive structural alterations, including localized duplications and

triplications, caused by replication fork stalling, template switching (FoSTeS), and

microhomology-mediated break-induced replication (MMBIR) (Pellestor & Gatinois,

2018). Such mechanisms increase the risk of unintended genomic alterations, thereby

limiting their application in chromosomal engineering.

In this study, we extended the scope of chromosomal engineering from fertilized

embryos to organ tissues using *in vivo* electroporation. Specifically, we modified the

CRISPR RNP electroporation method developed by Kobayashi et al. (2023) and

assessed its safety and efficiency in inducing chromosomal rearrangements in the

murine uterine epithelium. We also examined the presence of FoSTeS/MMBIR during

chromosomal repair. Gynecological cancers, particularly uterine cancer, are among the

most frequently diagnosed malignancies worldwide, with chromosomal abnormalities

such as translocations and inversions being common hallmark features of tumorigenesis

(Lu et al., 2021). The present study demonstrates the feasibility of *in vivo*

electroporation as a practical and efficient method for inducing chromosomal

rearrangements in the uterine epithelium.

Results

Establishment of an *in vivo* editing technique for CCR in the mouse uterine epithelium

Several clinical trials have demonstrated the potential of CRISPR-based genome editing in the treatment of various genetic disorders and cancers (Morshedzadeh et al., 2024). However, their potential for *in vivo* chromosomal rearrangements remains to be elucidated. To address this, we modified the CRISPR RNP electroporation method developed by Kobayashi et al. (2023) and applied it to induce chromosomal rearrangements in the murine uterine epithelium (Figure 1A; Figure S1). CRISPR RNPs were first injected into the uterine lumen via microinjection and subsequently introduced into uterine epithelial cells through electroporation. The use of a microinjector allowed for more precise and controlled reagent delivery, reducing mechanical stress on the tissue and minimizing leakage, which is expected to facilitate a more localized and consistent distribution of RNPs within the uterine lumen.

As an initial trial, we tested 1.1-Mb CCRs within mouse chromosome 10, encompassing the *Hmga2*, *Wif1*, and *Rassf3*, which are implicated in various human cancers and are mapped in a configuration similar to that of human chromosome 12 (Pareja et al., 2019) (Figure 1B). This region was previously targeted in fertilized embryos (Iwata et al., 2024), and was selected for gene editing in somatic tissues to enable direct comparison of editing efficiencies. On postoperative day 7, genomic DNA was extracted from the tissue and the achievement of the intended chromosomal rearrangements was assessed via PCR and Sanger sequencing.

Sequencing of the genomic DNA isolated from the uterus revealed that chromosomal

rearrangements were successfully induced at three different breakpoints (Figure 1C).

Sequence analysis confirmed that HR, nonhomologous end-joining (NHEJ), and

microhomology-mediated end-joining (MMEJ) were involved in breakpoint repair

(Figure 1C). Although the results suggest relatively high occurrence of genome-editing

events compared to previous reports in fertilized embryos (Figure 1D; Iwata et al.,

2024), direct comparisons should be interpreted with caution because of the intrinsic

differences between embryonic and somatic tissues. In fertilized embryos, genome

editing occurs at the single-cell stage, enabling precise determination of the proportion

of successfully edited embryos. In contrast, somatic tissues consist of a mosaic

population of edited and unedited cells, making absolute quantification of editing

efficiency more complex. Nevertheless, consistent detection of chromosomal

rearrangements at multiple breakpoints suggests that *in vivo* electroporation effectively

facilitates genome editing in somatic tissues.

Efficient induction of interchromosomal translocations in the mouse uterine

epithelium

Next, we evaluated the feasibility of inducing translocations, in which segments from

the two chromosomes were exchanged. The functional intergenic repeating RNA

element (*Firre*) gene located on the X chromosome was selected as the target locus. The

Firre locus plays a central role in interchromosomal interactions by facilitating gene

activation through three-dimensional spatial interactions with regulatory regions on

other chromosomes (Maass et al., 2019) (Figure 2A). Therefore, *Firre* serves as a key

hub for transcriptional regulation and is expected to provide a structural basis for

efficient translocation induction. We designed experiments to induce translocations

between the *Eef1a1* neighborhood locus (*Eef1a1N*) on chromosome 9 and the *Atf4*

neighborhood locus (*Atf4N*) on chromosome 15, both of which interact with the *Firre*

loci. Additionally, we targeted the translocations between *Ypel4* neighborhood locus

(*Ypel4N*) on chromosome 2 and *Atf4N* on chromosome 15, as these loci are also

physically linked to the *Firre* locus (Figure 2A, B). On postoperative day 7, the

predicted translocations were detected in uterine samples (Figure 2C, D). The observed

translocation efficiencies were 33.3 and 66.7% for *Eef1a1N–Atf4N* and 40 and 30% for

Ypel4N–Atf4N, respectively (Figure 2E). These findings suggest a high efficiency of

translocation induction in this experimental setting and support the potential

effectiveness of this *in vivo* electroporation-based approach.

Repair of the large-scale 57.8-Mb chromosomal inversion, *In(6)1J*, in the mouse

uterine epithelium

CRISPR/Cas9 technology and its derivatives, including base and prime editors, offer

new therapeutic avenues for cancer and genetic disorders (Porto et al., 2020; Katti et al.,

2022; Chen et al., 2023). Here, we explored the feasibility of repairing chromosomal

abnormalities in the mouse uterine epithelium via CRISPR/Cas9-based *in vivo*

electroporation. We aimed to repair the large-scale 57.8-Mb inversion *In(6)1J* on

chromosome 6, which was first identified in the C3H/HeJ strain in 2006 (Akeson et al.,

2006; Ackert-Bicknell et al., 2007) and whose breakpoints were previously mapped in

2021 (Iwata et al., 2021). This inversion spans approximately 40% of the total length of

chromosome 6, and is implicated in specific phenotypes, including low bone mass and

increased bone marrow adiposity (Ackert-Bicknell et al., 2007). In the present study, we

designed gRNAs near breakpoints and employed a strategy to repair chromosomal

inversions using ssODNs corresponding to normal genomic sequences (Figure 3A).

Repair efficiencies of 18.2 and 36.4% were observed at the left and right breakpoints,

respectively (Figure 3B, C). The successful repair of this 57.8-Mb inversion

demonstrated that this *in vivo* approach is applicable to the repair of large-scale

chromosomal abnormalities. To the best of our knowledge, this is the largest

chromosomal rearrangement targeted for genome editing.

In vivo chromosomal engineering of the mouse uterine epithelium by targeting

multiple genetic loci

Based on these findings, a broad investigation of locus-dependent differences in editing

efficiency is crucial. Therefore, we evaluated the ability of *in vivo* electroporation to

induce translocations and inversions at other genetic loci. To introduce oncogenic fusion

genes, translocations were induced between *Ncoa2* and *Greb1*, as well as between

Ywhae and *Nutm2*, both of which are associated with uterine sarcomas (Figure 4A, B).

GREB1–NCOA2 fusion integrates the estrogen-responsive growth-promoting function

of GREB1 with the nuclear receptor coactivator function of NCOA2, thereby enhancing

hormone-responsive cancer cell proliferation, tumor initiation, and progression

(Brunetti et al., 2018). YWHAE–NUTM2 fusion promotes oncogenesis by inducing

cyclin D1 overexpression via the regulation of the RAF/mitogen-activated protein

kinase and Hippo signaling pathways, driving the proliferation of high-grade

endometrial stromal sarcoma cells (Ou et al., 2021). Additionally, we re-examined the

7.67-Mb inversion previously reported for fertilized embryos (Iwata et al., 2024)

(Figure S2A). This targeted locus exhibits FoSTeS/MMBIR in fertilized embryos

carrying *Recql5* mutations (Iwata et al., 2024).

Successful chromosomal engineering was observed at all the tested genetic loci (Figure

4C, D, E; Figure S2B, C). Notably, no FoSTeS/MMBIR-associated events were detected

at any of the loci analyzed in this study. Although the fusion genes were successfully

induced during the translocations, no tumors were detected during the observation

period. Given that tumor development often requires an extended timeframe (Klein,

2009), further long-term investigations are required. Collectively, our findings suggest

that *in vivo* electroporation effectively induced chromosomal rearrangements at multiple

genetic loci.

Discussion

Using CRISPR RNP electroporation, we successfully induced interchromosomal

translocations and repaired a large-scale 57.8-Mb inversion, demonstrating the

feasibility of this approach for correcting chromosomal abnormalities. Given that

large-scale chromosomal rearrangements are frequently associated with genetic

disorders and cancers (Popova et al., 2012; Pellestor, 2019; Comaills et al., 2023), the

ability to efficiently correct them *in vivo* presents a promising avenue for therapeutic

genome editing.

Previous studies, such as those by Reimer et al. (2017), utilized lentiviral vectors to

induce chromosomal translocations, including the t(11, 19)/MLL-ENL translocation.

While effective, lentiviral-based methods pose risks, such as insertional mutagenesis

and prolonged Cas9 expression, increasing the likelihood of off-target effects and

genomic instability (Bushman, 2020). Additionally, their production is costly, complex,

and requires biosafety containment. Adeno-associated virus-based genome editing has

been widely adopted because of its low immunogenicity and stable transgene expression

(Verdera et al., 2020). However, its small packaging capacity (~4.7 kb) restricts the

delivery of large genome-editing components such as Cas9 and multiple guide RNAs.

Adenoviral vectors offer an alternative viral delivery system with high packaging

capacity (~36 kb) and robust transgene expression. Unlike lentiviral vectors, AdVs do

not integrate into the host genome, thereby minimizing the risk of insertional

mutagenesis (Li and Lieber, 2019). Nevertheless, their high immunogenicity often

triggers strong inflammatory responses, leading to reduced efficacy *in vivo* (Asmamaw

Mengstie, 2022). In contrast, our CRISPR RNP electroporation approach eliminates

these risks by enabling transient genome editing, thereby reducing the potential for

genomic instability. Additionally, electroporation does not require viral vector

production, simplifies the experimental workflow, and significantly reduces costs. These

advantages highlight the safety and practicality of *in vivo* electroporation for

chromosomal engineering, particularly for applications requiring precise genome

editing in somatic tissues.

In addition to these advantages, a key observation in this study was that targeted

chromosomal rearrangements were successfully induced and consistently detected in

somatic tissues at a higher frequency than in fertilized embryos. To the best of our

knowledge, CRISPR RNP electroporation has not been reported to successfully induce

translocation in fertilized embryos. This discrepancy may be attributed to several factors,

including differences in the number of available target cells, cell cycle duration, and

DNA repair mechanisms, which are predominant in each cellular environment. Unlike

fertilized embryos, in which genome editing occurs at the one-cell stage, somatic tissues

consist of numerous target cells, increasing the absolute number of successfully edited

cells and enabling the consistent detection of chromosomal rearrangements. However,

in fertilized embryos, a rapid cell cycle may constrain the time window available for

DNA repair (Khokhlova et al., 2020), particularly in the context of large-scale

chromosomal rearrangements. These rearrangements can induce genomic instability,

reduce cell viability, and limit the efficiency of chromosomal engineering. Previous

studies have reported that the efficiency of CRISPR-based genome editing is

significantly influenced by the cell cycle phase (Abe et al., 2020), and that rapid

embryonic cell division may leave insufficient time for precise repair. In contrast, the

prolonged cell cycle in somatic tissues provides more time for accurate DNA repair

(Kermi et al., 2019), which may have facilitated the successful induction of

translocation in the present study. Additionally, uterine epithelial cells, particularly those

in the metestrus phase, exhibit more stable cell cycle regulation (Baranda-Avila et al.,

2009), potentially providing a favorable environment that supports efficient

chromosomal engineering while mitigating the risk of genomic instability-induced cell

death.

Overall, the findings highlight the feasibility of CRISPR RNP-based *in vivo*

electroporation as a viable approach for chromosomal engineering of somatic tissues.

This approach offers a transient and flexible alternative to virus-based methods, thereby

avoiding the concerns related to stable genome integration. The successful repair of a

57.8-Mb chromosomal inversion further underscores its potential for large-scale

genome correction, offering new opportunities in cancer research, congenital disorder

treatment, and regenerative medicine. Further optimization and long-term validation

studies are essential to fully utilize this approach for therapeutic applications.

Limitations

In the present study, chromosomal rearrangements were efficiently induced in uterine

epithelial samples, although variability in genome-editing efficiency was observed.

Therefore, further optimization of the electroporation parameters, including voltage and

pulse duration, is necessary to improve the effectiveness of this technique. In addition,

its feasibility in other tissues and cell types should be explored to enhance its

applicability.

Future studies with extended observation periods and comprehensive biological

evaluations, including tumorigenic risk assessments, are required, to determine the

long-term biological effects of this method. This study provides important insights into

the feasibility of *in vivo* electroporation-based chromosomal engineering. However,

further rigorous validation is essential to establish its safety and efficacy for future

clinical applications.

Materials and methods

Experimental animals

Wild-type (WT) mice (C57BL/6NCrSlc; Japan SLC, Shizuoka, Japan) and *In(6)1J*

inversion-carrying mice (C3H/HeJJcl; CLEA Japan, Tokyo, Japan) were used in this

study. Sexually mature female mice aged 8 weeks or older were selected for

experiments. All animals were maintained under controlled environmental conditions at

a constant temperature of 22 ± 2 °C and humidity of 50 ± 10% under a 12/12-h

light/dark cycle. All animal experiments were approved by the Institutional Animal Care

and Use Committee of Chubu University (permit number #202410012; Kasugai, Aichi,

Japan) and adhered to the guidelines.

CRISPR RNP and single-stranded oligodeoxynucleotide (ssODN) preparation

CRISPR guide RNAs (gRNAs) were designed using CHOPCHOP

(http://chopchop.cbu.uib.no/; Table S1) (Labun et al., 2019). The CRISPR RNP

complex was assembled using Alt-R S.p. Cas9 Nuclease 3NLS (Integrated DNA

Technologies, Coralville, IA, USA) and a custom-designed gRNA consisting of a

CRISPR RNA (crRNA):trans-activating CRISPR RNA (tracrRNA) duplex (Integrated

DNA Technologies). After resuspending in the nuclease-free duplex buffer (Integrated

DNA Technologies) at a final concentration of 4,000 ng/μL, crRNA and tracrRNA were

mixed at equimolar ratios, heated at 95 °C for 10 min, and gradually cooled to 25 °C to

facilitate duplex formation. Then, the crRNA:tracrRNA duplex was incubated with

Alt-R S.p. Cas9 Nuclease 3NLS at 25 °C for 10 min to assemble the RNP complex.

Specific concentrations are indicated in the following section: *in vivo* genome editing of

the murine endometrium. Subsequently, ssODNs were synthesized by Eurofins

Genomics (Tokyo, Japan) to facilitate the seamless joining of the two DNA sequences,

ensuring that the junction was positioned at the center of the predicted cleavage sites

located within 3 bp of the protospacer adjacent motif (PAM) sequences (Table S2).

Moreover, 5′ and 3′ ends of ssODNs were modified with two consecutive

phosphorothioate (*) linkages to enhance the HR efficiency (Renaud et al., 2016) (Table

S2).

In vivo **genome editing of the murine endometrium**

Female metestrus or diestrus mice were subjected to *in vivo* genome editing of the

murine endometrium using a modified version of the method previously described by

Kobayashi et al. (2023). To generate complex CCRs, the following CRISPR solution

was used: 520 ng/μL Alt-R S.p. Cas9 Nuclease 3NLS, 30 μM crRNA:tracrRNA duplex

each for the left, middle, and right targets, 158 ng/μL ssODN each for the left, middle,

and right targets, and 0.02% Fast Green FCF (Wako, Osaka, Japan) marker diluted in

the Opti-MEM (Thermo Fisher Scientific, Waltham, MA, USA). To generate

chromosomal inversions and translocations, the following CRISPR solution was used:

520 ng/μL Alt-R S.p. Cas9 Nuclease 3NLS, 30 μM crRNA:tracrRNA duplex each for

the left and right targets in inversions and for each breakpoint on the respective

chromosomes in translocations, 158 ng/μL ssODN each for the left and right targets in

inversions and for each breakpoint on the respective chromosomes in translocations,

and 0.02% Fast Green FCF (Wako) marker diluted in the Opti-MEM (Thermo Fisher

Scientific). Prior to electroporation, the female mice were anesthetized with a mixture of

medetomidine (0.75 mg/kg, Nippon Zenyaku Kogyo, Fukushima, Japan), midazolam (4

mg/kg, Sandoz, Tokyo, Japan), and butorphanol (5 mg/kg, Meiji Seika Pharma, Tokyo,

Japan).

Glass micropipettes were first pulled using a vertical capillary puller (NARISHIGE,

Tokyo, Japan) to create fine-tipped capillaries suitable for microinjection. Before use,

the tip of each capillary was cut diagonally to a length of 0.5 mm using micro-scissors

to facilitate smooth reagent flow. The cut capillary was then connected to a syringe via a

silicone tube and filled with the CRISPR solution, ensuring complete reagent loading.

Once filled, the capillary was attached to a capillary holder and secured in place using a

flexible stand. The CRISPR solution (2 μL) was then introduced into the uterus via

microinjection from the downstream part of the oviduct using the prepared glass

micropipette and the FemtoJet 4i microinjector (Eppendorf, Hamburg, Germany) under

the following parameters: injection pressure (pi) = 100 hPa, injection duration (ti) = 0.2

s, and compensation pressure (pc) = 0 hPa. To prevent reagent backflow, the oviduct

near the ovary was clamped using a hemostatic clip (Natsume Seisakusho, Tokyo,

Japan). The uterus was clamped to the cervical side using a hemostatic clip (Natsume

Seisakusho) to prevent leakage. Following the injection, the uterine horns were clamped

using tweezer electrodes (LF650P5; BEX, Tokyo, Japan), and electroporation was

performed using a CUY21EDIT II electroporator (BEX). The electrode orientation was

adjusted and three additional sets of electric pulses were applied. The following

electroporation parameters were used: primary large pulse (PLP): 20 V, 30.0-ms pulse

duration, 50.0-ms pulse interval, 3 pulses, 10% decay (± pulse orientation). Secondary

decay pulse (Pd): 10 V, 50.0-ms pulse duration, 50.0-ms pulse interval, 3 pulses, 40%

decay (± pulse orientation). After electroporation, the uterine horns were repositioned

and the incisions were sutured. To reverse the effects of medetomidine, atipamezole

hydrochloride (0.75 mg/kg, Nippon Zenyaku Kogyo, Fukushima, Japan) was

intraperitoneally administered postoperatively.

Analysis of the CRISPR/Cas9-engineered mice

To screen for CRISPR/Cas9-induced mutations, genomic DNA was extracted from the

uterine horns of founder mice using a DNeasy Blood & Tissue Kit (Qiagen, Hilden,

Germany) and subjected to polymerase chain reaction (PCR) amplification using

EmeraldAmp PCR Master Mix (Takara Bio, Shiga, Japan). For nested PCR, the first

round consisted of 40 cycles of denaturation at 98 °C for 10 s, annealing at 60 °C for 30

s, and extension at 72 °C for 1 min, followed by a final indefinite hold at 10 °C. The

second round of PCR consisted of 15 cycles of denaturation at 98 °C for 10 s, annealing

at 63 °C for 30 s, and extension at 72 °C for 30 s, with a final indefinite hold at 10 °C.

The resulting PCR products were purified using a NucleoSpin Gel and PCR Cleanup

Kit (Takara Bio) and cloned into the pTAC-1 vector (Biodynamics, Tokyo, Japan).

Individual clone sequences were determined using Sanger sequencing (Eurofins

Genomics). All PCR primers used for genotyping are listed in Table S3.

Acknowledgments

The authors thank the laboratory members for their support with animal care and the experiments. The authors would like to thank Editage (www.editage.com) for English language editing.

Funding

This study was supported by JSPS KAKENHI Grant Number 23K21287 (to SI), the Science Research Promotion Fund from the Promotion and Mutual Aid Corporation for Private Schools of Japan (to SI), and a Chubu University Grant (S) (to SI).

Author contributions

SI designed and performed the experiments and drafted the manuscript. YM performed animal experiments. TI supervised the study and revised the manuscript. All the authors have read and approved the final manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

References

Abe T, Inoue KI, Furuta Y, & Kiyonari H. 2020. Pronuclear microinjection during
S-phase increases the efficiency of CRISPR-Cas9-assisted knockin of large DNA
donors in mouse zygotes. *Cell Rep* **31**: 107653.

Ackert-Bicknell CL, Salisbury JL, Horowitz M, DeMambro VE, Horton LG, Shultz KL,
Lecka-Czernik B, Rosen CJ. 2007. A chromosomal inversion within a quantitative trait
locus has a major effect on adipogenesis and osteoblastogenesis. *Ann N Y Acad Sci*
1116: 291–305.

Akeson EC, Donahue LR, Beamer WG, Shultz KL, Ackert-Bicknell C, Rosen CJ,
Corrigan J, Davisson MT. 2006. Chromosomal inversion discovered in C3H/HeJ mice.
Genomics **87**:311–313.

Asmamaw Mengstie M. 2022. Viral vectors for the in vivo delivery of CRISPR
components: advances and challenges. *Front. bioeng. biotechnol.* **10**: 895713.

Baranda-Avila N, Mendoza-Rodríguez CA, Morimoto S, Langley E, & Cerbón M. 2009.
Molecular mechanism of cell proliferation in rodent uterus during the estrous cycle. *J*
Steroid Biochem Mol Biol **113**: 259-268.

Bushman FD. 2020. Retroviral Insertional Mutagenesis in Humans: Evidence for Four
Genetic Mechanisms Promoting Expansion of Cell Clones. *Mol. Ther* **28**: 352–356.

Blasco RB, Karaca E, Ambrogio C, Cheong TC, Karayol E, Minero VG, Voena C,
Chiarle R. 2014. Simple and rapid in vivo generation of chromosomal rearrangements
using CRISPR/Cas9 technology. *Cell Rep* **9**:1219–1227.

Brunetti M, Panagopoulos I, Gorunova L, Davidson B, Heim S, Micci F. 2018. RNA-sequencing identifies novel GREB1-NCOA2 fusion gene in a uterine sarcoma with the chromosomal translocation t(2;8)(p25;q13). *Gene Chromosome Cancer* **57**: 176–181.

Chen JM, Cooper DN, Férec C, Kehrer-Sawatzki H, Patrinos GP. 2010 Genomic Rearrangements in Inherited Disease and Cancer. *Semin. Cancer Biol* **20**: 222–233.

Chen PJ, Liu DR. 2023. Prime editing for precise and highly versatile genome manipulation. *Nat Rev Genet* **24**: 161–177.

Comaills V, & Castellano-Pozo M. 2023. Chromosomal instability in genome evolution: From cancer to macroevolution. *Biology* **12**: 671.

Feuk L. 2010 Inversion variants in the human genome: role in disease and genome architecture. *Genome Med* **2**: 11.

Greely HT. 2019. CRISPR'd babies: human germline genome editing in the 'He Jiankui affair'. *J Law Biosci* **6**:111–83.

Iwata S, Sasaki T, Nagahara M, Iwamoto T. 2021. An efficient *i*-GONAD method for creating and maintaining lethal mutant mice using an inversion balancer identified from the C3H/HeJJcl strain. *G3* **11**: jkab194.

Iwata S, Nagahara M, Ido R, Iwamoto T. 2024. A *Recql5* mutant facilitates complex CRISPR/Cas9-mediated chromosomal engineering in mouse zygotes. *Genetics* **227**;

iyae054

Katti A, Diaz BJ, Caragine CM, Sanjana NE, Dow LE. 2022. CRISPR in cancer biology
and therapy. *Nat Rev Cancer* **22**: 259–279.

Kermi C, Aze A, Maiorano D. 2019. Preserving Genome Integrity During the Early
Embryonic DNA Replication Cycles. *Genes* **10**: 398.

Khokhlova EV, Fesenko ZS, Sopova JV, & Leonova EI. 2020. Features of DNA repair
in the early stages of mammalian embryonic development. *Genes* **11**: 1138.

Klein CA. 2009. Parallel progression of primary tumours and metastases. *Nat Rev*
Cancer **9**: 302–312.

Kobayashi R, Kawabata-Iwakawa R, Sugiyama M, Oyama T, Ohtsuka M, Horii T,
Morita S, Nishiyama M, Hatada I. 2023. Multiplexed genome editing by in vivo
electroporation of Cas9 ribonucleoproteins effectively induces endometrial carcinoma in
mice. *Int J Cancer* **152**: 2331–2337.

Kweon J, Hwang HY, Ryu H, Jang AH, Kim D, Kim Y. 2023. Targeted genomic
translocations and inversions generated using a paired prime editing strategy. *Mol Ther*
31:249–259.

Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E. 2019.
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic*
Acids Res **47**:W171–W174.

Lee JA, Lupski JR. 2006. Genomic rearrangements and gene copy-number alterations
as a cause of nervous system disorders. *Neuron* **52**:103-121

Li C, and Lieber A. 2019. Adenovirus Vectors in Hematopoietic Stem Cell Genome
Editing. *FEBS Lett.* **593**: 3623–3648.

Lu B, Jiang R, Xie B, Wu W, Zhao Y. 2021. Fusion Genes in Gynecologic Tumors: The
Occurrence, Molecular Mechanism and Prospect for Therapy. *Cell Death Dis* **12**: 783.

Maddalo D, Manchado E, Concepcion CP, Bonetti C, Vidigal JA, Han YC, Ogrodowski
P, Crippa A, Rekhtman N, de Stanchina E, et al. 2014. *In vivo* engineering of oncogenic
chromosomal rearrangements with the CRISPR/Cas9 system. *Nature* **516**: 423–427.

Maass PG, Barutcu AR, Rinn JL. 2019. Interchromosomal interactions: a genomic love
story of kissing chromosomes. *J Cell Biol* **218**: 27–38.

Mitelman F, Johansson B, Mertens F. 2007. The impact of translocations and gene
fusions on cancer causation. *Nat Rev Cancer* **7**: 233–245.

Morshedzadeh F, Ghanei M, Lotfi M, Ghasemi M, Ahmadi M, Najari-Hanjani P, Sharif
S, Mozaffari-Jovin S, Peymani M, Abbaszadegan MR. 2024. An update on the
application of CRISPR technology in clinical practice. *Mol Biotechnol* **66**:179–197.

Nambiar TS, Baudrier L, Billon P, Ciccia A. 2022. CRISPR-based genome editing
through the lens of DNA repair. *Mol Cell* **82**: 348–388.

Ou WB, Lundberg MZ, Zhu S, Bahri N, Kyriazoglou A, Xu L, Chen T,

Mariño-Enriquez A, Fletcher JA. 2021. YWHAE-NUTM2 oncoprotein regulates
proliferation and cyclin D1 via RAF/MAPK and Hippo pathways. *Oncogenesis* **10**:37.

Pareja F, Geyer FC, Brown DN, Sebastiao APM, Gularte-Merida R, Li A, Edelweiss M,
Da Cruz Paula A, Selenica P, Wen HY, et al. 2019. Assessment of *HMGA2* and *PLAG1*
rearrangements in breast adenomyoepitheliomas. *NPJ Breast Cancer* **5**:6.

Pellestor F, Anahory T, Lefort G, Puechberty J, Liehr T, Hedon B, Sarda P. 2011.
Complex chromosomal rearrangements: Origin and meiotic behavior. *Hum Reprod*
Updat **17**: 476–494.

Pellestor F, Gatinois V. 2018. Chromoanasynthesis: another way for the formation of
complex chromosomal abnormalities in human reproduction. *Hum Reprod* **33**:1381–
1387.

Pellestor F. 2019. Chromoanagenesis: cataclysms behind complex chromosomal
rearrangements. *Molecular Cytogenetics* **12**:1–12.

Popova T, Manié E, Rieunier G, Caux-Moncoutier V, Tirapo C, Dubois T, ... & Stern M
H. 2012. Ploidy and large-scale genomic instability consistently identify basal-like
breast carcinomas with BRCA1/2 inactivation. Cancer research 72, 5454-5462.

Porto EM, Komor AC, Slaymaker IM. & Yeo GW. 2020. Base editing: advances and
therapeutic opportunities. *Nat Rev Drug Discov* **19**: 839–859.

Reimer J, Knoess S, Labuhn M, Charpentier EM, Gohring G, Schlegelberger B,
Klusmann JH, Heckl D. 2017. CRISPR–Cas9-induced t(11;19)/MLL-ENL
translocations initiate leukemia in human hematopoietic progenitor cells *in vivo*.

Haematologica **102**:1558–1566.

Renaud JB, Boix C, Charpentier M, De Cian A, Cochennec J, Duvernois-Berthet E,

Perrouault L, Tesson L, Edouard J, Thinard R, et al. 2016. Improved genome editing

efficiency and flexibility using modified oligonucleotides with TALEN and

CRISPR-Cas9 nucleases. *Cell Rep* **14**:2263–2272.

Tao R, Wang Y, Hu Y, Jiao Y, Zhou L, Jiang L, Li L, He X, Li M, Yu Y, et al. 2022

WT-PE: Prime editing with nuclease wild-type Cas9 enables versatile large-scale

genome editing. *Signal Transduct Target Ther* **7**:108.

Verdera HC, Kuranda K, and Mingozzi F. 2020. AAV Vector Immunogenicity in

Humans: A Long Journey to Successful Gene Transfer. *Mol. Ther.* **28**: 723–746.

Zepeda-Mendoza CJ, Morton CC. 2019. The iceberg under water:

unexploredcomplexity of chromoanagenesis in congenital disorders. *Am J Hum Genet*

2019:**104**:565–77.

Figure Legends

Figure 1.

CCRs in the uterine epithelium induced via *in vivo* electroporation. (A) Schematic

overview of the experimental procedure for CCR induction by *in vivo* electroporation.

(B) Diagram illustrating the CCRs among *Hmga2*, *Wif1*, and *Rassf3* on chromosome 10.

(C) Sequence alignment of PCR products corresponding to the genomic breakpoint

junctions, *Hmga2–Wif1*, *Hmga2–Rassf3*, and *Wif1–Rassf3*. Boxed regions indicate the

microhomology sequences. (D) Comparative analysis of CCR induction efficiency via

in vivo electroporation of within-oviduct fertilized embryos, fertilized embryos, and

uterine epithelium.

Figure 2.

Translocations at the *Firre* locus in the uterine epithelium induced via *in vivo*

electroporation. (A) Schematic representation of the translocation between the

Eef1a1N on chromosome 9 and *Atf4N* on chromosome 15. (B) Sequence alignment of

the PCR products corresponding to the genomic breakpoint junctions of *Eef1a1N* and

Atf4N. (C) Schematic representation of the translocation between the *Ypel4N* on

chromosome 2 and *Atf4N* on chromosome 15. (D) Sequence alignment of the PCR

products corresponding to the genomic breakpoint junctions of *Ypel4N* and *Atf4N*.

Boxed regions indicate the microhomology sequences. (E) Overall translocation

efficiency in the uterine epithelium.

Figure 3.

Repair of a large-scale chromosomal inversion in the uterine epithelium via *in vivo*

electroporation. (A) Schematic representation of the CRISPR/Cas9-based strategy to

repair the 57.8-Mb inversion, *In(6)1J*, on chromosome 6. (B) Sequence alignment of the

PCR products corresponding to the left and right breakpoint junctions after repair. (C)

Overall repair efficiency in the uterine epithelium.

Figure 4.

Oncogenic translocations in the uterine epithelium induced via *in vivo*

electroporation. (A) Schematic representation of the translocation between the *Ncoa2*

on chromosome 1 and *Greb1* on chromosome 12. (B) Sequence alignment of the PCR

products corresponding to the genomic breakpoint junctions of *Ncoa2* and *Greb1*.

Boxed regions indicate the microhomology sequences. (C) Schematic representation of

the translocation between *Ywhae* on chromosome 11 and *Nutm2* on chromosome 13. (D)

Sequence alignment of the PCR products corresponding to the genomic breakpoint

junctions of *Ywhae* and *Nutm2*. (E) Overall translocation efficiency in the uterine

epithelium.

Cas9 targets	Distance (Mb)	Sample Types		Strains	n	Chromosomal rearrangements		
						Left breakpoints Detected (%)	Middle breakpoints detected (%)	Right breakpoints detected (%)
Hmga2 Wif1 Rassf3	1.01	In-oviduct fertilized egg (Iwata et al. 2024)		WT	28	1 (3.57)	1 (3.57)	3 (10.7)
				Recql5em1/em1	10	1 (10)	3 (30)	3 (30)
		Fertilized egg (Iwata et al. 2024)		WT	15	1 (6.67)	0 (0)	0 (0)
				Recql5em1/em1	36	4 (11.1)	1 (2.78)	4 (11.1)
		Uterus (In this study)		WT	10	3 (30)	3 (30)	3 (30)

A

In(6)1J

57.8 Mb inversion

DSB PAM

In(6)1J gDNA

Homology arm

ssODN
(repair template)

Homology arm

Repair

Normal chr. 6

B

Left breakpoints

Predicted GTCCTTTAATATATGTGTGTTGGAGGCCTCATATCAGTTGGTATATGCTGCCTGGTTGGTGGTCCAGCGCCTGAGAGATCTTGGGGTTCCAGG

#L1 GTCCTTTAATATATGTGTGTTGGAGGCCTCAT--------------------GGCCTGGTTGGTGGTCCAGCGCCTGAGAGATCTTGGGGTTCCAGG NHEJ

#R1 GTCCTTTAATATATGTGTGTTGGAGGCCTCATATCAGTTGGTATATGCTGCCTGGTTGGTGGTCCAGCGCCTGAGAGATCTTGGGGTTCCAGG HR

Right breakpoints

Predicted ACTGACAGGTCCTCGAGAAGAAGAAGAAGGCTCTTAACAACTGC————————AGCCTCCAGATGTACTACTATGATAATCCAATCCT

#L1 ACTGA---(Δ81 bp   )//AGGAAATG NHEJ

#R1 ACTGACAGGTCCTCGAGAAGAAGAAGAAGGCTCTTAACAACTGCtcATACCAACTGAT--CTCCAGATGTACTACTATGATAATCCAATCCT NHEJ

#L5 ACTGACAGGTCCTCGAGAAGAAGAAGAAGGCTCTTAACAACTGC————AGCCTCCAGATGTACTACTATGATAATCCAATCCT HR

#L6 ACTGACAGGTCCTCGAGAAGAAGAAGAAGGCTCTTAACAACTGC————AGCCTCCAGATGTACTACTATGATAATCCAATCCT HR

C

Cas9 targets	Distance (Mb)	*n*	Chromosomal rearrangements	
			Left breakpoints detected (%)	Right breakpoints detected (%)
In(6)1J left breakpoint *In(6)1J* right breakpoint	57.8	11	2 (18.2)	4 (36.4)