Golgi-Localized Mannanases Sustain Hemicellulose Biosynthesis

Talia Jacobson[1,2], Mair Edwards[1,2], Moni Qiande[1], Madalen Robert[1,3], Julia Moncrieff[1], Cătălin Voiniciuc[1,2,3]*

[1]Department of Horticultural Sciences, University of Florida, Gainesville, FL 32611, USA

[2]Plant Molecular and Cellular Biology Graduate Program, University of Florida, Gainesville, FL 32609, USA

[3]Independent Junior Research Group–Designer Glycans, Leibniz Institute of Plant Biochemistry, 06120 Halle (Saale), Germany

* Corresponding Author: Cătălin Voiniciuc

Email: cvoiniciuc@ufl.edu

Classification: Biological Sciences, Biochemistry

This PDF file includes:

Summary

- Mannans with β-1,4-linked backbones are common cell wall components of algae
and land plants. Prior challenges to enhance β-mannan content in plants point to
unclear metabolic bottlenecks and the potential for hidden biosynthetic players.

- Endo-β-MANNANASEs (MANs) are the main glycosyl hydrolases that mobilize
extracellular β-mannans during seed germination. However, we found that
Arabidopsis *man2 man5* seeds resemble β-mannan biosynthetic mutants.
CELLULOSE SYNTHASE-LIKE A (CSLA) overexpression restored β-mannan
synthesis in the *man* double mutant and increased the distribution of crystalline
polymers, but impaired the release of other mucilaginous polysaccharides.

- Using yeast synthetic biology, we dissected the functional interplay of MAN
enzymes with CSLAs. Intracellular MAN2 and MAN5 reduced the quantity of
insoluble β-mannan but elevated the content of water-soluble carbohydrates.

- We propose that Arabidopsis MAN2/5, and orthologous crop enzymes with a
transmembrane domain, sustain hemicellulose production in the Golgi apparatus
by cleaving insoluble β-mannan polymers into hydrophilic counterparts.

Keywords: plant cell wall, hemicellulose, polysaccharide biosynthesis, glycosyl
hydrolases, synthetic biology

Introduction

Among carbohydrate-active enzymes, glycosyl transferase (GT) and glycosyl hydrolase (GH) enzymes generally play antagonistic roles on plant cell wall polysaccharides. Hemicelluloses can cross-link cellulose microfibrils and interact with pectin domains to shape plant biology, and to define the structure of traits, dietary fibers, materials, and chemical feedstocks for industrial applications. The *Arabidopsis thaliana* (hereafter Arabidopsis) seed coat epidermal cells offer an excellent genetic model system to untangle cell wall polysaccharide biosynthesis and function. Upon contact with water, dry seeds release copious amounts of mucilaginous polysaccharides (Voiniciuc *et al.*, 2015c) with smaller amounts of hemicelluloses such as branched β-mannans (Voiniciuc *et al.*, 2015b) and crystalline cellulose (Griffiths *et al.*, 2015). Plant β-mannan backbones are defined by their β-1,4-linked mannose (Man) residues (Voiniciuc, 2022), but can also contain β-1,4-glucose (Glc). In Arabidopsis mucilage galactoglucomannan (GGM), the Man units are substituted with single α-1,6-linked galactosyl (Gal) residues (Voiniciuc *et al.*, 2015b; Yu *et al.*, 2018), or β-1,2-Gal-α-1,6-Gal disaccharides (Yu *et al.*, 2022). Highly branched GGM contributes to mucilage density and crystalline cellulose deposition (Voiniciuc *et al.*, 2015b), while another hemicellulose, branched xylan, is important for the adherence of pectic rhamnogalacturonan-I (~90% of Arabidopsis mucilage) to the seed surface (Voiniciuc *et al.*, 2015a; Ralet *et al.*, 2016). Other seed-extracted galactomannans (e.g. guar gum and locust bean gum) are frequently added as thickening agents, stabilizers, and emulsifiers in numerous food and household products, or serve as dietary fiber supplements to control appetite and reduce blood Glc levels in patients with type 2 diabetes (Juhász *et al.*, 2023).

Since their backbone and branches are composed entirely of hexose sugars (Scheller & Ulvskov, 2010), plant mannans have desirable properties for a circular bioeconomy and could support the bioengineering of new products or sustainable feedstocks for biomanufacturing (Voiniciuc, 2022, 2023; Jacobson *et al.*, 2024). Despite long-standing demands to elevate hexose sugar content in lignocellulosic biomass (Pauly & Keegstra, 2008), β-mannan content has been challenging to elevate in plants. For instance, seed-specific expression of a guar mannan synthase in the legume *Medicago*

truncatula severely disrupted metabolism and unexpectedly decreased galactomannan
levels (Naoumkina *et al.*, 2008).

As recently reviewed (Voiniciuc, 2022), β-mannan biosynthesis is orchestrated by
several classes of Golgi-localized glycosyl transferase (GTs) with a suite of secreted
glycosyl hydrolases (GHs) expected to modify polysaccharides after their deposition in the
cell wall. CELLULOSE SYNTHASE-LIKE A2 (AtCSLA2) is the chief catalyst for β-
mannan elongation in the Arabidopsis seed coat (Yu *et al.*, 2014). During mucilage GGM
elongation, MUCILAGE-RELATED10 (AtMUCI10 also known as AtMAGT1) decorates
Man residues with single α-1,6-Gal side chains (Voiniciuc *et al.*, 2015b; Yu *et al.*, 2018),
which can be further substituted with β-1,2-Gal residues by AtMBGT1 (Yu *et al.*, 2022).
Plant mannans are often galactosylated (Scheller & Ulvskov, 2010) or acetylated (Zhong
et al., 2019) because *in vitro* assays show that β-mannan oligosaccharides crystallize as
they elongate (Grimaud *et al.*, 2019). In the yeast *Pichia pastoris* (hereafter *Pichia*),
MANNAN-SYNTHESIS RELATED1 (AtMSR1) acts as a co-factor that enables
AtCSLA2 to produce glucomannan instead of relative pure mannan (Voiniciuc *et al.*,
2019). Since *Amorphophallus konjac* AkCSLA3 can produce glucomannan even without
MSR1 in yeast, it has been used to investigate the protein regions important for substrate
specificity (Robert *et al.*, 2021), to illuminate the localization and bottlenecks for
hemicellulose biosynthesis (Grieß-Osowski *et al.*, 2025).

Once β-mannans are deposited in the cell wall, endo-β-1,4-MANNANASES
(MANs) become the chief catalysts for the backbone modification and degradation. While
the biological roles of different mannan structures remains remain ambiguous (Boerjan *et*
al., 2024; Guevara-Rozo *et al.*, 2025), MANs are involved in wall softening and
carbohydrate release during plant development (Rodríguez-Gacio *et al.*, 2012). MAN
hydrolytic activity increases in a wide range of plant species during seed germination,
presumably to mobilize carbon reserves and to promote growth (Downie *et al.*, 1997;
Nonogaki *et al.*, 2000; Wang *et al.*, 2005; Iglesias-Fernández *et al.*, 2011, 2020; De Farias
et al., 2015; Carrillo-Barral *et al.*, 2018). Four of the seven *Arabidopsis MAN* genes are
expressed in dry seeds and three different single mutants (*man5*, *man6,* and *man7*, but not

man2) decreased or delayed germination (Iglesias-Fernández *et al.*, 2011). Mannan oligosaccharides released by AtMAN6 are proposed to activate signaling for xylem thickening, while AtMAN3 and AtMAN7 are induced by cadmium to increase apoplastic Man and promote heavy metal tolerance (Chen *et al.*, 2015; Wu *et al.*, 2023; Zhang *et al.*, 2024). Some MANs also have transglycosylation activity to essentially 'cut-and-paste' polymers, e.g. for fruit ripening (Schröder *et al.*, 2009). Intriguingly, some *MANs* are co-expressed with *CSLAs* during seed development in Arabidopsis (Voiniciuc, 2022) and in crops like coffee (Joët *et al.*, 2014), but their impact on hemicellulose accumulation remains unknown (Voiniciuc, 2022). Here, we apply plant and yeast bioengineering strategies to demonstrate that Arabidopsis MAN2 and MAN5 have non-canonical roles in the synthesis of soluble β-mannans.

Materials and Methods

The following sections describe the major materials and methods, with additional details for plant cultivation conditions, DNA construct assembly, yeast growth, biochemical analyses, cytometry and microscopy in the supplemental information (Methods **S1)**.

Genetic Materials and Analyses

Arabidopsis T-DNA insertion mutant seeds were obtained from the ABRC or NASC stock centers, *man2-1* (SALK_126628c.20.a), *man2-2* (SALK_016450), *man5-1* (707G06.01.a), *man5-2* (SALK_015220), *man5-3* (SALK_068367), *man2-1 man5-1* (2716/1a1.01), *muci10-1* (SALK_061576), and *csla2-3* (SALK_149092). *Agrobacterium*-mediated transformations of *Arabidopsis* plants, and transient assays in *Nicotiana benthamiana* leaves were conducted using established protocols (Yang *et al.*, 2022). For co-localization experiments, plasmid DNA for the Golgi yellow fluorescent protein (YFP) marker G-yk, previously validated for Arabidopsis (Nelson *et al.*, 2007), was obtained from ABRC (CD3-965).

Table **S1** lists the genotyping primers for the plant Touch-and-Go Method (Berendzen *et al.*, 2005) or for bacteria or yeast colony polymerase chain reaction (PCR). Cloning was mainly performed Golden Gate assembly (Engler *et al.*, 2014; Marillonnet &

Grützner, 2020), or NEB HiFi cloning (Table **S2**), as noted in Methods **S1**. Transcript levels in mature green siliques were analyzed using prior methods (Voiniciuc *et al.*, 2015b) and normalized to the geometric mean of two housekeeping genes (Table **S1**).

Biochemical Analysis of Mucilage and Yeast Cell Wall Fractions

Total mucilage was extracted for monosaccharide profiling via High-Performance Anion-Exchange Chromatography with Pulsed Amperometric Detection (HPAEC-PAD) as specified by (Voiniciuc *et al.*, 2015b; Voiniciuc & Günl, 2016), using an established Metrohm HPAEC-PAD system and eluent gradient (Robert *et al.*, 2021). *Pichia* alkaline-insoluble (AKI) polymers were isolated as previously described (Robert *et al.*, 2021), followed by the separation of alkaline-soluble fractions (Voiniciuc *et al.*, 2019). Water-soluble glycans were extracted from *Pichia* cells as previously described (Cocuron *et al.*, 2007) with an incubation step in 70% ethanol for 5 min and a subsequent incubation in water for 5 min at 100 C without Triton X-100. After centrifugation, the supernatant was removed, hydrolyzed, and injected for HPAEC-PAD analysis. The remaining AKI was incubated in NaOH for 90 min followed by hydrolysis. Water-soluble glycans from Arabidopsis dry seeds were analyzed in a similar manner. Seeds were first ground and then incubated in ethanol and then water. The water-soluble and alcohol-insoluble residue (AIR) fractions were then hydrolyzed with trifluoroacetic acid and analyzed via HPAEC-PAD.

Light Microscopy of Plant Materials and Yeast

Seed mucilage area was stained with ruthenium red (RR) (Voiniciuc *et al.*, 2015b) and quantified using an AXIO Vert A1 or a stereoscope equipped with an Axiocam 208 color camera or a 202 monochrome camera (all from Zeiss). Seed birefringence in water was captured with an adjustable polarizer/analyzer module, and intensity plots were generated as previously described (Voiniciuc *et al.*, 2016). Confocal images were captured on a Leica Stellaris 8 at the University of Florida Interdisciplinary Center for Biotechnology Research (UF/ICBR) Cytometry Core Facility (RRID: SCR_019119). Unless otherwise noted in figure legends, the 20x (0.75 NA, numerical aperture) objective was used with additional zoom in the Leica LAS X software. Calcofluor white (CF)

staining of seeds (Voiniciuc *et al.*, 2015b) or yeast (Robert *et al.*, 2021) was performed using 0.01% Fluorescent Brightener 28 (Sigma Aldrich). Mucilage immunolabeling was followed an established protocol (Voiniciuc, 2017), using four monoclonal antibodies CCRC-M36, CCRC-M70, CCRC-M75, and CCRC-M170 (CarboSource Services, University of Georgia) and a goat anti-mouse Alexa Fluor 488 secondary antibody (ThermoFisher Scientific, A28175). Confocal images were acquired with the following (Excitation | Emission) wavelengths: CF or turquoise protein fluorescence (405 nm | 450-470 nm), green or yellow protein fluorescence (500 nm | 505-550 nm), red fluorescence (550 nm | 600-640 nm). All plant and *Pichia* images were processed and analyzed with the Fiji distribution of ImageJ (Schindelin *et al.*, 2012).

Application of MannanTrack Probe in Yeast

The MannanTrack probe used here is corresponds to + pFDH1:PaCBM35:pmScarlet-I-2A-pSmTq2(stop)ScCYC1tt αMF-PaCBM35-mScarlet-P2A-mTurquoise2 and is introduced in the Results section, and is part of an upcoming collection of genetically encoded probes for plant mannan synthases (Robert *et al*, manuscript in preparation). This MannanTrack coding sequences were codon optimized for *Pichia pastoris*, synthesized and assembled using Golden Gate cloning in the GoldenPiCS vector BB3rN_14, under the control of the pFDH1 methanol-inducible promoter and the ScCYC1tt transcriptional terminator (Prielhofer *et al.*, 2017) Competent *Pichia* cells harboring the *AkCSLA3-Venus* cassette were transformed with the MannanTrack construct, integrated into the *RGI2* locus. After PCR genotyping and growth in YPM + G medium as previously described (Robert *et al.*, 2021), a representative colony was selected and made competent (Lin-Cereghino *et al.*, 2005). *MAN* episomal vectors were then introduced into the background. All colonies were verified via PCR genotyping before growth and further analyses specified in **Methods S1**.

Statistical Analyses

Statistically significant changes were identified one-way ANOVA analyses performed in PAST 5 software (Hammer *et al.*, 2001), or using Microsoft Excel's t.test function assuming two-tailed distribution and equal variance of two samples.

Results

***man2 man5* Seeds Resemble Mannan-Deficient *csla2* and *muci10* Mutants**

We focused on deciphering the roles of *MAN2* and *MAN5* (Fig. **1a**), two closely related paralogs with overlapping expression profiles (Iglesias-Fernández *et al.*, 2011). We obtained an uncharacterized *man2-1 man5-1* double mutant from the GABI-DUPLO collection (Bolle *et al.*, 2013), which targeted unlinked gene pairs in Arabidopsis. Real-time PCR confirmed that single and double mutants significantly reduced the transcript levels of the respective genes in developing siliques, at the stage of mucilage production (Fig. **1b**). The *man2-1 man5-1* seeds had 30% smaller mucilage capsules than the wild type (WT), (Fig. **1c,d**), while single *man2/5* mutants were not significantly different. Such defects are observed in several mucilage mutants, including *csla2* (Yu *et al.*, 2014) and *muci10* seeds with arrested β-mannan synthesis (Voiniciuc *et al.*, 2015b). In contrast to the reduced *MAN2* and *MAN5* transcript levels, the expression of *CSLA2* or *MUCI10* was not significantly altered in the *man2 man5* double mutant (Fig. **1b**). Additional insertions in *MAN5* could not be verified (Table **S1**), so we crossed an independent *man2-2* allele to *man5* plants and isolated a novel *man2-2 man5* double mutant, which also had compact mucilage compared to WT, without altering the seed dimensions (Fig. **1c-e**). Consistent with prior *man5* data (Iglesias-Fernández *et al.*, 2011), only 50% of *man2 man5* seeds germinated after 48 h after stratification, compared to 80% of WT seeds (Fig. **2b**). The double mutant seeds had delayed germination across growth batches (Figs. **2b**, **S1c**), but caught up with WT and *csla2* seeds by 96-120 h.

Fig. 1. Mannanase double mutant Arabidopsis seeds have decreased mucilage accumulation. (a) T-DNA insertions in *AtMAN2* and *AtMAN5* genes, showing exon and intron structure. Red arrows indicate positions of primer binding utilized for quantitative PCR (Scale bar, 200 bp). (b) Fold changes in *AtMAN2* and *AtMAN5* or *AtCSLA2* and *AtMUCI10* transcription relative to WT in mature green siliques. Data show mean + SD of three biological replicates normalized to the geometric mean of *UBQ5* and *GAPC1*. Asterisks indicate significant decreases relative to WT (Student's *t* test, $P < 0.01$). (c) Pectin release of WT and mutant seeds stained with RR. (d) Projected mucilage and (e) seed surface area. Data shows mean + SD of three biological replicates (around 20 seeds each). Asterisks indicate significant decreases relative to WT (Student's *t* test, $P < 0.05$). Bars: (c) 200 μm.

To understand why mutations in β-mannan synthases and hydrolases resulted in

similar staining defects, despite their antagonistic biochemical activities, we profiled total

mucilage composition with HPAEC-PAD. Compared to *man* single mutants and WT,

man2 man5 mucilage contained significantly less Gal and Man, the most characteristic

monosaccharides of branched β-mannans (Fig. **2c**; Table **S3a**). Glc, the third component

of GGM, showed similar trends but had a high coefficient of variation because it can be

derived from several cell wall polymers as well as starch. Although pectin immunolabeling

with CCRC-M36 showed results consistent with the RR staining defects, β-mannan

antibodies (M70, M75, or M170) in the CCRC collection (Thorne *et al.*, 2023) could not

bind WT or *man2 man5* mucilage (Fig. **S1e**). Nevertheless, *man2 man5* mucilage

phenotypes and chemotypes approached the severity of *csla2* and *muci10*, with up to 50%
reductions in Gal and Man compared to WT across multiple growth batches (Figs. **2c, 3c,**
S1). These data are consistent with the participation of *MAN2* and *MAN5* in β-mannan
production, rather than its catabolism in the seed coat.

Fig. 2. The *man2 man5* mutant seeds are impaired in β-mannan elongation. (a) Schematic of mucilage mannan biosynthesis and MAN2/5 protein domains. (b) Mean + SD germination rate for three biological replicates. For each time point, asterisks indicate significant decreases relative to WT (Student's *t* test, $P < 0.05$). (c) Absolute monosaccharide levels of three biological replicates. Different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$).

AtCSLA2 Overexpression Rescues *csla2* Defects, but Impairs *man2 man5* Mucilage

Despite the yeast activities described later in the Results section, CSLAs and MANs
cDNA sequences (Fig. **S1a,b**) have been difficult to express in plants. In this study, we
found that *csla2* mucilage β-mannan deficiency (Table **S3b**) can be rescued with a seed
coat-specific *pTBA2:CSLA2* construct, featuring the *TESTA ABUNDANT2* promoter
(McGee *et al.*, 2021). The *CSLA2* transgene restored the RR staining of pectin (Fig. **S2a)**
and CF staining of cellulosic rays (Fig. **3a**) around *csla2* seeds. In contrast, when
introduced in the *man2 man5* background, *pTBA2:CSLA2* led to longer yet disordered rays
(Fig. **3a,b**) and a patchy mucilage capsule (Fig. **S2a**). The trends in cellulose staining were
supported by imaging of crystalline polymer birefringence under polarized light. The *man2*
man5 double mutants had an intermediate reduction of birefringence compared to *csla2*
seeds (Fig. **S2b-d**). Although *CSLA2* overexpression in the *man2 man5* seed coat elevated
Man and Gal content to nearly WT levels (**Fig. 3c**), it decreased the amount of total

extractable mucilage by nearly 60% (Fig. **3d,** Table **S3b**). This was likely due to reduced mucilage release/extraction, rather than impaired secretion, because scanning electron microscopy (SEM) did not reveal major changes in the surface morphology of the mucilage pockets or the columellae (Fig. **S1d**).

Fig. 3. Golgi-localized MAN enzymes are important for mucilage structure. (a) CF staining of β-glucans in seed mucilage. Arrows mark disordered rays. (b) Length of CF-stained rays. (c) *AtCSLA2* overexpression affects Gal and Man content, and (d) total mucilage release. Biochemical (e) and mucilage RR staining (f) complementation of *man2 man5* by AtMAN gDNA constructs. Different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$). (g) Optical sections of *N. benthamiana* leaves expressing a Golgi-YFP marker and AtMAN2-mCherry or an mCherry control. Arrows mark co-localization. Bars: (a) 100 μm, (f) 200 μm, (g) 30 μm.

Non-Canonical MAN2 and MAN5 Localize in the Golgi Apparatus

Unlike most plant and microbial MANs, AtMAN2 and AtMAN5 are predicted to have an N-terminal transmembrane domain instead of a signal peptide (Figs. **2a, S3**). However, determining their subcellular localization *in planta* was not straightforward and required extensive troubleshooting (summarized in Table **S4**). *AtMAN2/5* cDNA sequences

did not show any detectable protein expression/function *in planta* using *pTBA2*, the putative *pMAN2* promoter, or the viral *p35S* promoter (Fig. **S1a-d**). Similar vectors worked for fluorescent protein controls and cDNA sequences for other enzymes (e.g. CSLA2), but plant gene expression may require distal or intron-localized enhancers (Vollen *et al.*, 2024), which are difficult to predict. To circumvent these unexpected challenges, we amplified *AtMAN*[gDNA] genomic sequences for C-terminal *mCherry* fusions driven by the Arabidopsis *UBIQUITIN10* promoter (*pUBQ10)*. At long last, the *MAN2*[gDNA]*-mCherry* combination showed detectable fluorescence after transient expression in *N. benthamiana*. Unlike the cytosolic mCherry control, MAN2-mCherry co-localized with the known Golgi-YFP marker G-yk (Nelson *et al.*, 2007), in small punctae (Fig. **3g**) and showed similar trajectories (Video **S1**), consistent with roles in mannan biosynthesis**.** Since *MAN5*[gDNA]*-mCherry* did not show any detectable fluorescence, we cloned an even longer genomic *MAN5* construct that features its putative promoter, native stop codon, and its 3' untranslated region. Eight *pMAN5:MAN5*[gDNA] and four *pUBQ10:MAN2*[gDNA]*-mCherry* transformants rescued the *man2 man5* biochemical defects and compact mucilage phenotype (Fig. **3e,f**, Table **S3c**).

Intracellular MAN2 and MAN5 Reduce Insoluble β-Mannan Accumulation in Yeast

MAN[cDNA]*-mCherry* constructs were expressed in a modular *Pichia* platform via an episomal vector compatible with Plant Modular Cloning (MoClo) standards (Grieß-Osowski *et al.*, 2025). AtMAN2 and AtMAN5 proteins were localized in intracellular compartments (Figs. **4b**, **S4d**) and did not alter the glycan composition of WT X-33 cells (Fig. **4a**). While episomal expression of AtMSR1 enhanced glucomannan production in CSLA yeast strains (Fig. **4a**, **S6a**), co-expression of AtMAN2/5 with either AkCSLA3 or AtCSLA2 reduced insoluble β-mannan content by 30–40% (Figs. **4a**, **S4a**). To ensure that GH activity led to these differences, AtMAN2's catalytic nucleophile (Wang *et al.*, 2015) was mutated to an alanine (Figs. **2a**, **S3a-c**). AtMAN2[E335A] had similar fluorescence intensity and subcellular localization to its WT sequence (Fig. **S3d**) but no longer altered mannan content (Fig. **4a**, **S4a**). Similar biochemical changes to the episomal constructs were observed for integrative *AtMAN-mRuby2* constructs driven by a weaker promoter.

280　After alkaline treatment, the integrative *AtMAN2/5* constructs reduced the content of

281　insoluble mannans (Fig. **S5**), without significant changes in the alkaline-soluble fractions

282　that are rich in mannosylated yeast proteins.

Fig. 4. MAN2/5 enzymes reduce insoluble mannan content in yeast. (a) Relative Man content of insoluble yeast polymers. Box plots show three biological replicates. (b) Arrows mark the co-localization of AkCSLA3-Venus and AtMAN2/5-mCherry and counterstained with CF in confocal micrographs acquired with an 86x (1.2 NA) objective. (c) AKI polymers made by AtCSLA2 plus AtMAN2 or the fungal AaMAN5a. (d) AZCL-galactomannan digestion assays using *Saccharomyces* cell mixtures and supernatant, alongside commercial MAN controls (Megazyme, E-BMANN). Different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$). Magenta labels indicate a C-terminal protein fusion to mCherry. Bars: (b) 5 μm.

AtMAN2 Co-Localizes with AkCSLA3 and Cannot Digest Extracellular β-Mannans

Since AkCSLA3 tolerates C-terminal fluorescent tags (Grieß-Osowski *et al.*, 2025),

we co-expressed AtMAN2-mCherry with AkCSLA3-Venus in *Pichia* cells. Although only

a subset of AkCSLA3-Venus cells showed mCherry fluorescence due to the greater

heterogeneity of episomal plasmid expression, the two proteins co-localized in the

endomembrane system (Fig. **4b**, **S4c,d**). Furthermore, AkCSLA3-Venus fluorescence

remained constant with either mCherry or MAN-mCherry protein (Fig. **S4c**). We

hypothesized that unbranched glucomannan production by CSLAs would perturb yeast cell

ultrastructure, and that AtMAN2/5 enzymes could alleviate such defects. However,

transmission electron micrographs of 3-day-old *Pichia* cells showed enlarged vacuoles and

signs of autophagy, even without CSLAs (Fig. **S7a,b**). Although the same harvest

timepoint was suitable for live cell imaging and biochemical analyses in this study and

prior work (Grieß-Osowski *et al.*, 2025), we could not observe Golgi bodies in thin sections

(100 nm) of cells preserved via high-pressure freezing or chemical fixation. However, we

noticed that AkCSLA3 cells (accumulating glucomannan) had 30% thicker cell walls

compared to the negative control (**Fig. S7c,d**). Co-expression of AkCSLA3 with the wild-

type AtMAN2 (but not the AtMAN2^{E335A} variant) restored the wild-type thickness.

Plant MANs are reported to have low catalytic activity (Wang *et al.*, 2015) relative

to microbial MANs (Ishii *et al.*, 2016), such as *AaMAN5a* from the plant pathogen

Aspergillus aculeatus. While AtMAN2 reduced insoluble Man levels in the *Pichia*

AtCSLA2 strain by 46%, fungal AaMAN5a led to a 66% decrease (Fig. **4c**), resulting in

WT-like yeast cell walls. Since AaMAN5a has been used to ferment β-mannans in

Saccharomyces cerevisiae (Ishii *et al.*, 2016), we also introduced *AtMANs* in this host and

tested their ability to digest exogenous Azurine Cross-Linked (AZCL)-galactomannan.

Blue-dyed β-mannan oligosaccharides were released by the supernatant fraction of cells

secreting AaMAN5a (Fig. **4d**), or by minute amounts of a fungal MAN purified from

Aspergillus niger. Neither AtMAN2 nor AtMAN5 could digest extracellular

galactomannan (Fig. **4d**), despite replacing their N-terminal transmembrane regions (Fig.

S3a) with the yeast α-galactosidase (αGal; enabling AaMAN5a secretion) or the α-mating

factor (αMF) signal peptide. This is likely due to a lack of AtMAN protein secretion,

because fluorescently tagged AtMAN2/5 proteins were retained inside *Pichia* cells despite

the addition of the αMF signal peptide (Fig. **S3d**) or if their transmembrane domains were

simply removed (Fig. **S4b**). However, when removing the first 34 amino acids without

adding a signal peptide, MAN2(34)-mRuby2 proteins became cytosolic and no longer

altered the production of insoluble mannans (Fig. **S6a,b**), which occurs in the Golgi lumen.

Tracking Mannan Accumulation and Hemicellulose Solubility

320 To further quantify how AtMANs influence hemicellulose accumulation in yeast,

321 we applied a novel, genetically encoded probe featuring a carbohydrate-binding module

(CBM) derived from a microbial MAN to specifically track β-mannans. The potential utility of CBM-based probes for live cell imaging was previously reviewed (Voiniciuc *et al.*, 2018b), and the development of a collection of MannanTrack probes to monitor the activity of mannan synthases in yeast and plants will be fully characterized in a separate manuscript (Robert *et al*, in preparation). Here, we briefly introduce and demonstrate the utility of one MannanTrack probe that consists of the *Podospora anserina* CBM35, which is involved in mannan-binding based on structural and biochemical analyses (Couturier *et al.*, 2013). An N-terminal αMF signal peptide targets the PaCBM35 for secretion, and a C-terminal tags containing mScarlet-I (Bindels *et al.*, 2017) and a superfolder mTurquoise2 ox (Meiresonne *et al.*, 2019) as ratiometric reporters for red and blue fluorescence (Fig. **5a**). The two fluorescent proteins are separated by a "self-cleaving" P2A linker from the porcine teschovirus-1 2A, which can result in ribosome skipping during translation to generate separate polypeptides from bi-cistronic vectors (Liu *et al.*, 2017). We observed successful skipping and recommencement of translation for our MannanTrack probe resulting in two separatee fluorescent reporters: the mannan-binding αMF-PaCBM35-mScarlet was fully secreted from *Pichia* cells without AkCSLA3 (**Fig. 5a,** additional images in **Fig. S8**), while the mTurquoise2 (downstream of P2A) remained in the cytosol.

In contrast to yeast cells with native polysaccharide composition, the αMF-PaCBM35-mScarlet probe was largely retained inside cells (ie. reducing the RFP secretion) producing insoluble mannan (Fig. **5a,b**) and co-localized with AkCSLA3-Venus (Figs. **5a**, **S8**). Based on plate reader measurements (Fig. **5b**), confocal microscopy (Fig. **S8**), and flow cytometry (Fig. **5c**), the introduction of *MAN2/5* significantly increased MannanTrack secretion. For instance, *MAN2/5* enhanced mScarlet fluorescence in the culture supernatant by 20% relative to empty vector or *MAN2^{E335A}* controls (Fig. **5b**), which prompted us to evaluate β-mannan solubility. Using hot water to directly extract soluble β-1,4-linked glucans from ground yeast cells (Cocuron *et al.*, 2007), we isolated 30–60% more Man from cells co-expressing *AkCSLA3* with *AtMAN2/5*, compared to *AkCSLA3* alone or the non-functional *MAN2^{E335A}* (Fig. **5d, S9**). To determine whether MANs function similarly in Arabidopsis, we ground dry seeds and isolated alcohol-insoluble residue (AIR) as well as water-soluble fractions for HPAEC-PAD analysis (see extraction scheme in **Fig. S9**).

The *man2 man5* mutants contained significantly less Man in both fractions compared to WT, and *CSLA2* overexpression increased Man content primarily in the insoluble polymer fraction (Fig. **5e**). Therefore, we propose that MAN enzyme activity in the endomembrane system converts insoluble mannan polysaccharides into shorter and more hydrophilic molecules.

Fig. 5. *MAN2* and *MAN5* promote mannan solubility and secretion. (a) MannanTrack probe components in *Pichia* control strains to validate the blue (BFP), yellow (YFP) and red (RFP) protein fluorescence patterns. (b) Relative secretion (supernatant RFP / cell mixture RFP) of the PaCBM35-mScarlet, normalized to the BFP cell fluorescence relative to empty vector (EV). (c) Number of cells with detectable AkCSLA3-Venus and PaCBM35a-mScarlet fluorescence quantified via flow cytometry. (d) Man content of hot water (H_2O-soluble) and AKI polymers in *Pichia* extracts. (e) Man content of hot water (H_2O-soluble) and AIR polymers in Arabidopsis ground seeds. Box plots show three biological replicates and different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$). Magenta labels indicate a C-terminal protein fusion to mCherry. (f) Updated model of β-mannan biosynthesis in the Golgi apparatus (based on Voiniciuc, 2022) with AtMAN2/5 role(s). Bars: (a) 5 μm.

Discussion

Breaking A Few Bricks to Build a Wall?

Thus far, plant MANs are reported to degrade or cut-and-paste extracellular β-mannans (Bewley, 1997; Moreira & Filho, 2008; Schröder *et al.*, 2009), and typically have antagonistic consequences to Golgi-localized (gluco)mannan synthases (Voiniciuc, 2023). In this study, we discovered that Arabidopsis seeds have reduced mannan production without *MAN2* and *MAN5* (Fig. **1d**, **2c, 5e**), despite wild-type *CSLA2* and *MUCI10* transcript levels (**Fig. 1b**). The significant loss of mannan accumulation in seeds (**Fig. 5e**) despite the normal secretion of other mucilage polysaccharides (**Fig. S1d,e**) implies that *man2 man5* double mutants primarily disrupt mannan elongation. Therefore, we hypothesize that intracellular MAN2/5 enzymes cooperate with Golgi-localized GTs to sustain mannan production (**Fig. 5f**). Although this represents a novel role for endo-MANs, an endo-glucanase is known to participate in cellulose biosynthesis. KORRIGAN1 (KOR1) is a plasma membrane-localized cellulase that associates with the Cellulose Synthase Complex (Vain *et al.*, 2014). While genetic studies demonstrate its importance for cellulose synthesis, KOR1's precise mode of action remains elusive. For instance, KOR1 is proposed to relieve mechanical stress placed on the cell as crystalline polymers are deposited into the wall (Mielke *et al.*, 2021). Since unbranched β-mannans also form crystalline structures as they elongate (Grimaud *et al.*, 2019), MAN2/5 could be needed to reduce intracellular polymer aggregates and mechanical strain (Fig. **5f**). This raises the question: why would plant cells build polysaccharides only to immediately degrade them? One possible explanation is that CSL polysaccharide synthases may lack an intrinsic mechanism to stop polymer elongation, making controlled cleavage by GHs essential for regulating fiber length and preventing aggregation.

Although *AtCSLA2* overexpression using *pTBA2* forced the elongation of mannans, it could not fully restore the content of water-soluble mannans in seeds (**Fig. 5e**) and disrupted the organization of other mucilage polymers (**Fig. 3a,b**, **S2a,d**). In the *csla2* background, *pTBA2:CSLA2* nearly doubled the amount of mannan compared to WT plants (**Fig. 3c**). In contrast, *pTBA2:CSLA2* increased the mannan content of *man2 man5* mucilage extracts only to WT levels but significantly disrupted the release of pectin (**Fig. 3d, S2**).

The *man2 man5* + *CSLA2* seeds could have reduced secretion of rhamnogalacturonan-I to the cell wall and/or impaired mucilage release due to altered matrix composition and elevated content of birefringent polymers. The second option appears more likely as the SEM of *man2 man5* seeds resembled the WT (Fig. **S1d**), instead of the severe defects of seeds deficient in pectin synthesis (Voiniciuc *et al.*, 2018a). Mutants such as *mum2* are known to release minimal mucilage due to elevated polysaccharide cross-linking rather than a reduction in pectin content (Dean *et al.*, 2007; Voiniciuc *et al.*, 2015c). Therefore, our results reinforce the importance of mannans for matrix polysaccharide architecture as well as cellulose deposition (Voiniciuc *et al.*, 2015b; Yu *et al.*, 2018; Voiniciuc, 2022). MAN2/5 also have functions beyond the seed coat, with similar biosynthetic roles proposed by an independent study of *man2 man5* xylem vessels (Kikuchi *et al.*, 2025). We also found that *man2 man5* double mutants had delayed germination (Fig. **2c**), consistent with *MAN* gene expression in the endosperm (Iglesias-Fernández *et al.*, 2011).

How Hemicellulose Structure Impacts Matrix Polysaccharide Secretion

Despite increasing similarities between branched β-mannan and xyloglucan biosynthesis (Yu *et al.*, 2022; Grieß-Osowski & Voiniciuc, 2023), major questions remain about the precise functions of different hemicellulose structures in plants (Boerjan *et al.*, 2024). Xyloglucan mutants lacking Gal decorations have severe growth penalties and secretion defects (Hoffmann & McFarlane, 2024), yet mutations in the five *CSLC* glucan synthases in Arabidopsis leads to relatively mild defects (Kim *et al.*, 2020). Similarly, poplar *csla* mutants lacking detectable mannans in the xylem were shown to grown normally (Guevara-Rozo *et al.*, 2025). We anticipate that galactosylation and/or acetylation are critical substitutions to maintain β-mannan solubility in the Golgi and matrix polysaccharide secretion to the cell wall. The lack of enzymes or donor substrates for mannan or glucan substitution in native *Pichia* cells could explain the intracellular retention of the MannanTrack probe in glucomannan-producing yeast cells (Fig. **5a-c**, **S8**). The modularity of this platform offers further opportunities to decipher how plant GTs and GHs cooperate to define hemicellulose content and properties.

Unraveling CSLA and MAN Roles through Yeast Synthetic Biology

While *in vitro* GH assays yield useful catalytic benchmarks and are helpful to improve enzymes for industrial use (Jacobson *et al.*, 2024), they could not have predicted the unexpected biological roles of AtMAN2/5. Yeast species, including *Pichia* and *Saccharomyces* (Grieß-Osowski *et al.*, 2025), can serve as surrogate hosts to characterize plant carbohydrate-active enzymes (Pauly *et al.*, 2019). Intracellular MAN2/5 enzymes reduced insoluble β-mannan production by CSLAs through a variety of biochemical and microscopic assays (Fig. **4**, **5**). Surprisingly, yeast cells showed no evidence of MAN2/5 secretion even when the transmembrane domains were replaced with αMF or αGal secretion signals that worked for the MannanTrack probe and the fungal AaMAN5a enzyme (Figs. **4d**, **S3**, **S4**). Previously, MAN2 was affinity purified from *Pichia* cultures for *in vitro* assays using a similar N-terminal truncation (Wang *et al.*, 2015); however, our experiments differ in the choice of the recombinant protein tags and our application of the stronger *pCAT1* promoter in episomal vectors (Vogl *et al.*, 2016). Recent examples of human GTs involved in *N*-glycosylation indicate that the N-terminal transmembrane stem as well as the carbohydrate-active domains can define Golgi localization (Yagi *et al.*, 2024). Amino acids beyond the predicted N-terminal membrane stem may thus contribute to the intracellular retention of MAN2/5 in yeast, although the truncation of the first 34 AtMAN2 amino acids were sufficient for its redistribution to the cytosol (Fig. **S6**), where it could no longer reduce the accumulation of insoluble mannans. The co-expression of AtMAN2/5 with CSLA elevated the content of water-soluble Man (Fig. **5d**), likely representing β-mannan oligosaccharides isolated from the endomembrane system. Using the same extraction method, *CSLC4* expression in *Pichia* produced water-soluble β-1,4-linked glucan oligosaccharides with an estimated length of four to six residues (Cocuron *et al.*, 2007), while co-expression with *XXT1* (a xylosyltransferase enzyme belonging to the GT34 family that includes *MUCI10/MAGT1*) boosted insoluble glucan production (Robert *et al.*, 2021).

Potential Modes of Action for Golgi-Localized MANs

Finally, we consider three (not mutually exclusive) mechanisms by which MAN2/5 enzymes could function together with CSLAs to benefit β-mannan synthesis (Fig. **5f**): i)

hydrolysis of insoluble molecules, ii) transglycosylation of oligosaccharides, and/or iii) protein interactions with CSLAs. In our yeast data, MAN2/5 enzymes partly hydrolyzed insoluble mannans made by CSLAs (Fig. **4a, 5d**), so we favor the first hypothesis. Transglycosylation, the second mode of action, seems unlikely because purified MAN2 enzyme acted solely as a hydrolase *in vitro* (Wang *et al.*, 2015). Cellulose synthases and KORRIGAN1 are known to interact (Vain *et al.*, 2014), so future studies could investigate if CSLAs and MANs form protein complexes in the Golgi apparatus. Nevertheless, the catalytically inactive AtMAN2^{E355A} largely lost its hydrolytic effects despite having no obvious changes in protein expression level or subcellular localization (Figs. **S3**, **S4**), and therefore its potential for interactions with CSLA. Since crops have close orthologs of AtMAN2/5, the unexpected roles of intracellular MANs in CSLA-driven hemicellulose biosynthesis open up new paths to go glyco (Voiniciuc, 2023) and to enhance β-mannan fibers for the bioenergy and bioproducts. Even for plant MANs with characterized biochemical activities, experimental evidence for their subcellular localization is lacking (Table **S8**). Therefore, we anticipate that co-expressing CSLAs with intracellular MANs could alleviate historic bottlenecks to bioengineering plants with designer β-mannans.

Acknowledgments

We thank the following individuals for their technical assistance: Bo Yang (initial *man2 man5* screening), Hannah Zheng (seed phenotyping), and Karen Kelley (sample preparation at the University of Florida ICBR Electron Microscopy Core Facility, RRID:SCR_019146). Sylvestre Marillonnet provided expert advice and materials for Plant MoClo assembly. Doug Smith and Mariza Miranda provided technical advice for confocal microscopy and flow cytometry at the University of Florida ICBR Cytometry Core Facility, RRID:SCR_019119. This work was supported by startup funding from UF/IFAS, the Horticultural Sciences Department, and the USDA National Institute of Food and Agriculture, Research Capacity Fund (Hatch) project 7004470. CV was also supported by core funding (Leibniz Association) from the Federal Republic of Germany and the state of Saxony-Anhalt, and by DFG grants (414353267). TJ and ME thank the PMCB program and College of Agricultural and Life Sciences (CALS) Dean's Awards for graduate research funding. The research visit of MR was supported by a DAAD (German Academic

478 Exchange Service) scholarship, while JM was supported by the UF University Scholars
479 program. Finally, we thank Toshihisa Kotake for helpful discussions about MAN function
480 since learning about our independent approaches in the summer of 2024.
481

482 **Competing Interest Statement:** The authors declare no competing interest.
483
484 **Author Contributions:** TJ and CV designed the work, analyzed data and wrote the
485 manuscript. TJ acquired most of the data, with additional contributions from ME, MQ and
486 JM. MR and CV developed the MannanTrack yeast strain.

487
Data availability
Data supporting the findings of this work are available in the main text and in the figures
and tables in Supporting Information. The genetic materials generated in this current study
are available from the corresponding author upon request. Popular plasmids and seeds will
be donated to Addgene and ABRC, respectively. High-resolution transmission electron
micrographs for yeast are on FigShare: dx.doi.org/10.6084/m9.figshare.30369646

References
Berendzen K, Searle I, Ravenscroft D, Koncz C, Batschauer A, Coupland G, Somssich IE, Ülker B.
2005. A rapid and versatile combined DNA/RNA extraction protocol and its application to the
analysis of a novel DNA marker set polymorphic between Arabidopsis thaliana ecotypes Col-0
and Landsberg erecta. *Plant Methods* **1**: 4.

Bewley JD. **1997**. Breaking down the walls — a role for endo-β-mannanase in release from seed
dormancy? *Trends in Plant Science* **2**: 464–469.

Bindels DS, Haarbosch L, van Weeren L, Postma M, Wiese KE, Mastop M, Aumonier S,
Gotthard G, Royant A, Hink MA, *et al.* 2017. mScarlet: a bright monomeric red fluorescent
protein for cellular imaging. *Nature Methods* **14**: 53–56.

Boerjan W, Burlat V, Cosgrove DJ, Dunand C, Dupree P, Haas KT, Ingram G, Jamet E, Mohnen
D, Moussu S, *et al.* 2024. Top five unanswered questions in plant cell surface research. *The Cell*
Surface **11**: 100121.

Bolle C, Huep G, Kleinbölting N, Haberer G, Mayer K, Leister D, Weisshaar B. **2013**. GABI-
DUPLO: a collection of double mutants to overcome genetic redundancy in Arabidopsis thaliana.
The Plant Journal **75**: 157–171.

Carrillo-Barral N, Matilla AJ, Rodríguez-Gacio MDC, Iglesias-Fernández R. **2018**. Mannans and
endo-β-mannanase transcripts are located in different seed compartments during Brassicaceae
germination. *Planta* **247**: 649–661.

Chen J, Yang L, Gu J, Bai X, Ren Y, Fan T, Han Y, Jiang L, Xiao F, Liu Y, *et al.* 2015. *MAN 3* gene
regulates cadmium tolerance through the glutathione-dependent pathway in *Arabidopsis*
thaliana. *New Phytologist* **205**: 570–582.

Cocuron J-C, Lerouxel O, Drakakaki G, Alonso AP, Liepman AH, Keegstra K, Raikhel N,
Wilkerson CG. **2007**. A gene from the cellulose synthase-like C family encodes a β-1,4 glucan
synthase. *Proceedings of the National Academy of Sciences* **104**: 8550–8555.

Couturier M, Roussel A, Rosengren A, Leone P, Stålbrand H, Berrin J-G. **2013**. Structural and
Biochemical Analyses of Glycoside Hydrolase Families 5 and 26 β-(1,4)-Mannanases from
Podospora anserina Reveal Differences upon Manno-oligosaccharide Catalysis *. *Journal of*
Biological Chemistry **288**: 14624–14635.

De Farias ET, Da Silva EAA, Toorop PE, Bewley JD, Hilhorst HWM. **2015**. Expression studies in
the embryo and in the micropylar endosperm of germinating coffee (Coffea arabica cv. Rubi)
seeds. *Plant Growth Regulation* **75**: 575–581.

Dean GH, Zheng H, Tewari J, Huang J, Young DS, Hwang YT, Western TL, Carpita NC, McCann
MC, Mansfield SD, *et al.* 2007. The Arabidopsis MUM2 Gene Encodes a β-Galactosidase
Required for the Production of Seed Coat Mucilage with Correct Hydration Properties. *The Plant*
Cell **19**: 4007–4021.

Downie B, Hilhorst HWM, Bewley JD. **1997**. Endo-β-mannanase activity during dormancy
alleviation and germination of white spruce (*Picea glauca*) seeds. *Physiologia Plantarum* **101**:
405–415.

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ, Marillonnet S.
2014. A Golden Gate Modular Cloning Toolbox for Plants. *ACS Synthetic Biology* **3**: 839–843.

Grieß-Osowski A, Robert M, Qiande M, Clauss S, Voiniciuc C. **2025**. Illuminating Glucomannan
Synthases To Explore Cell Wall Synthesis Bottlenecks. *ACS Synthetic Biology* **14**: 2472–2479.

Grieß-Osowski A, Voiniciuc C. **2023**. Branched mannan and xyloglucan as a dynamic duo in plant
cell walls. *The Cell Surface* **9**: 100098.

Griffiths JS, Šola K, Kushwaha R, Lam P, Tateno M, Young R, Voiniciuc C, Dean G, Mansfield SD,
DeBolt S, *et al.* 2015. Unidirectional Movement of Cellulose Synthase Complexes in Arabidopsis
Seed Coat Epidermal Cells Deposit Cellulose Involved in Mucilage Extrusion, Adherence, and Ray
Formation. *Plant Physiology* **168**: 502–520.

Grimaud F, Pizzut-Serin S, Tarquis L, Ladevèze S, Morel S, Putaux J-L, Potocki-Veronese G.
2019. *In Vitro* Synthesis and Crystallization of β-1,4-Mannan. *Biomacromolecules* **20**: 846–853.

Guevara-Rozo S, de Vries L, Yoshimi Y, Eli E, Dupree P, Mansfield SD. **2025**. Unraveling mannan
biosynthesis in poplar and spruce: functional characterization of Cellulose Synthase-Like family A
(CSLA) genes. *New Phytologist* **n/a**: 10.1111/nph.7068.

Hammer Ø, Harper DAT, Ryan PD. **2001**. PAST: Paleontological statistics software package for
education and data analysis. *Palaeontologia Electronica* **4**.

Hoffmann N, McFarlane HE. **2024**. Xyloglucan side chains enable polysaccharide secretion to the
plant cell wall. *Developmental Cell*: S1534580724003848.

Iglesias-Fernández R, Pastor-Mora E, Vicente-Carbajosa J, Carbonero P. **2020**. A Possible Role of
the Aleurone Expressed Gene HvMAN1 in the Hydrolysis of the Cell Wall Mannans of the Starchy
Endosperm in Germinating Hordeum vulgare L. Seeds. *Frontiers in Plant Science* **10**: 1706.

Iglesias-Fernández R, Rodríguez-Gacio MC, Barrero-Sicilia C, Carbonero P, Matilla A. **2011**.
Three endo-β-mannanase genes expressed in the micropylar endosperm and in the radicle
influence germination of Arabidopsis thaliana seeds. *Planta* **233**: 25–36.

**Ishii J, Okazaki F, Djohan AC, Hara KY, Asai-Nakashima N, Teramura H, Andriani A, Tominaga
M, Wakai S, Kahar P, *et al.* 2016**. From mannan to bioethanol: cell surface co-display of β-
mannanase and β-mannosidase on yeast Saccharomyces cerevisiae. *Biotechnology for Biofuels*
9: 188.

Jacobson T, Edwards M, Voiniciuc C. **2024**. Sharp solutions to cleave plant fibers. *Current
Opinion in Biotechnology* **90**: 103219.

Joët T, Laffargue A, Salmona J, Doulbeau S, Descroix F, Bertrand B, Lashermes P, Dussert S.
2014. Regulation of galactomannan biosynthesis in coffee seeds. *Journal of Experimental Botany*
65: 323–337.

**Juhász AE, Greff D, Teutsch B, Gede N, Hegyi P, Horváth EM, Deák PÁ, Nyirády P, Ács N, Juhász
R**. **2023**. Galactomannans are the most effective soluble dietary fibers in type 2 diabetes: a
systematic review and network meta-analysis. *The American Journal of Clinical Nutrition* **117**:
266–277.

**Kikuchi A, Sato E, Yoshimi Y, Takasaki H, Nishigaki N, Atsuzawa K, Kaneko Y, Yamaguchi M,
Takahashi D, Dupree P, *et al.* 2025**. Atypical endo-β-1,4-mannannases are necessary for normal
glucomannan synthesis in Arabidopsis.

**Kim S-J, Chandrasekar B, Rea AC, Danhof L, Zemelis-Durfee S, Thrower N, Shepard ZS, Pauly M,
Brandizzi F, Keegstra K**. **2020**. The synthesis of xyloglucan, an abundant plant cell wall
polysaccharide, requires CSLC function. *Proceedings of the National Academy of Sciences* **117**:
20316–20324.

Lin-Cereghino J, Wong WW, Xiong S, Giang W, Luong LT, Vu J, Johnson SD, Lin-Cereghino GP.
2005. Condensed protocol for competent cell preparation and transformation of the
methylotrophic yeast Pichia pastoris. *BioTechniques* **38**: 44–48.

Liu Z, Chen O, Wall JBJ, Zheng M, Zhou Y, Wang L, Ruth Vaseghi H, Qian L, Liu J. **2017**.
Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector. *Scientific*
Reports **7**: 2193.

Marillonnet S, Grützner R. **2020**. Synthetic DNA Assembly Using Golden Gate Cloning and the
Hierarchical Modular Cloning Pipeline. *Current Protocols in Molecular Biology* **130**: e115.

McGee R, Dean GH, Wu D, Zhang Y, Mansfield SD, Haughn GW. **2021**. Pectin Modification in
Seed Coat Mucilage by *In Vivo* Expression of Rhamnogalacturonan-I- and Homogalacturonan-
Degrading Enzymes. *Plant and Cell Physiology* **62**: 1912–1926.

Meiresonne NY, Consoli E, Mertens LMY, Chertkova AO, Goedhart J, den Blaauwen T. **2019**.
Superfolder mTurquoise2ox optimized for the bacterial periplasm allows high efficiency in vivo
FRET of cell division antibiotic targets. *Molecular Microbiology* **111**: 1025–1038.

Mielke S, Zimmer M, Meena MK, Dreos R, Stellmach H, Hause B, Voiniciuc C, Gasperini D.
2021. Jasmonate biosynthesis arising from altered cell walls is prompted by turgor-driven
mechanical compression. *Science Advances* **7**: eabf0356.

Moreira LRS, Filho EXF. **2008**. An overview of mannan structure and mannan-degrading enzyme
systems. *Applied Microbiology and Biotechnology* **79**: 165–178.

Naoumkina M, Vaghchhipawala S, Tang Y, Ben Y, Powell RJ, Dixon RA. **2008**. Metabolic and
genetic perturbations accompany the modification of galactomannan in seeds of Medicago
truncatula expressing mannan synthase from guar (Cyamopsis tetragonoloba L.). *Plant*
Biotechnology Journal **6**: 619–631.

Nelson BK, Cai X, Nebenführ A. **2007**. A multicolored set of *in vivo* organelle markers for co-
localization studies in Arabidopsis and other plants. *The Plant Journal* **51**: 1126–1136.

Nonogaki H, Gee OH, Bradford KJ. **2000**. A Germination-Specific Endo-β-Mannanase Gene Is
Expressed in the Micropylar Endosperm Cap of Tomato Seeds. *Plant Physiology* **123**: 1235–1246.

Pauly M, Gawenda N, Wagner C, Fischbach P, Ramírez V, Axmann IM, Voiniciuc C. **2019**. The
Suitability of Orthogonal Hosts to Study Plant Cell Wall Biosynthesis. *Plants* **8**: 516.

Pauly M, Keegstra K. **2008**. Cell-wall carbohydrates and their modification as a resource for
biofuels. *The Plant Journal* **54**: 559–568.

Prielhofer R, Barrero JJ, Steuer S, Gassler T, Zahrl R, Baumann K, Sauer M, Mattanovich D,
Gasser B, Marx H. **2017**. GoldenPiCS: a Golden Gate-derived modular cloning system for applied
synthetic biology in the yeast Pichia pastoris. *BMC Systems Biology* **11**: 123.

Ralet M-C, Crépeau M-J, Vigouroux J, Tran J, Berger A, Sallé C, Granier F, Botran L, North HM.
2016. Xylans Provide the Structural Driving Force for Mucilage Adhesion to the Arabidopsis Seed
Coat. *Plant Physiology* **171**: 165–178.

Robert M, Waldhauer J, Stritt F, Yang B, Pauly M, Voiniciuc C. **2021**. Modular biosynthesis of
plant hemicellulose and its impact on yeast cells. *Biotechnology for Biofuels* **14**: 140.

Rodríguez-Gacio MDC, Iglesias-Fernández R, Carbonero P, Matilla ÁJ. **2012**. Softening-up
mannan-rich cell walls. *Journal of Experimental Botany* **63**: 3975–3988.

Scheller HV, Ulvskov P. **2010**. Hemicelluloses. *Annual Review of Plant Biology* **61**: 263–289.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,
Saalfeld S, Schmid B, *et al.* 2012. Fiji: an open-source platform for biological-image analysis.
Nature Methods **9**: 676–682.

Schröder R, Atkinson RG, Redgwell RJ. **2009**. Re-interpreting the role of endo-β-mannanases as
mannan endotransglycosylase/hydrolases in the plant cell wall. *Annals of Botany* **104**: 197–204.

Thorne K, Urbanowicz BR, Hahn MG. **2023**. Plant Cell Wall Glycan-Directed Monoclonal
Antibodies. In: Plant Cell Walls. Boca Raton: CRC Press, 206–236.

Vain T, Crowell EF, Timpano H, Biot E, Desprez T, Mansoori N, Trindade LM, Pagant S, Robert S,
Höfte H, *et al.* 2014. The Cellulase KORRIGAN Is Part of the Cellulose Synthase Complex. *Plant*
Physiology **165**: 1521–1532.

Vogl T, Sturmberger L, Kickenweiz T, Wasmayer R, Schmid C, Hatzl A-M, Gerstmann MA, Pitzer
J, Wagner M, Thallinger GG, *et al.* 2016. A Toolbox of Diverse Promoters Related to Methanol
Utilization: Functionally Verified Parts for Heterologous Pathway Expression in Pichia pastoris.
ACS Synthetic Biology **5**: 172–186.

Voiniciuc C. **2017**. Whole-seed Immunolabeling of Arabidopsis Mucilage Polysaccharides. *BIO-*
PROTOCOL **7**: 1–10.

Voiniciuc C. **2022**. Modern mannan: a hemicellulose's journey. *New Phytologist* **234**: 1175–1184.

Voiniciuc C. **2023**. It's time to go glyco in cell wall bioengineering. *Current Opinion in Plant*
Biology **71**: 102313.

Voiniciuc C, Dama M, Gawenda N, Stritt F, Pauly M. **2019**. Mechanistic insights from plant
heteromannan synthesis in yeast. *Proceedings of the National Academy of Sciences* **116**: 522–
527.

Voiniciuc C, Engle KA, Günl M, Dieluweit S, Schmidt MH-W, Yang J-Y, Moremen KW, Mohnen
D, Usadel B. **2018a**. Identification of Key Enzymes for Pectin Synthesis in Seed Mucilage. *Plant*
Physiology **178**: 1045–1064.

Voiniciuc C, Günl M. **2016**. Analysis of Monosaccharides in Total Mucilage Extractable from
Arabidopsis Seeds. *BIO-PROTOCOL* **6**: 1–12.

Voiniciuc C, Günl M, Schmidt MH-W, Usadel B, Guenl M, Schmidt MH-W, Usadel B. **2015a**.
Highly Branched Xylan Made by IRREGULAR XYLEM14 and MUCILAGE-RELATED21 Links Mucilage
to Arabidopsis Seeds. *Plant Physiology* **169**: 2481–95.

Voiniciuc C, Pauly M, Usadel B. **2018b**. Monitoring polysaccharide dynamics in the plant cell
wall. *Plant Physiology* **176**: 2590–2600.

**Voiniciuc C, Schmidt MH-W, Berger A, Yang B, Ebert B, Scheller HV, North HM, Usadel B, Günl
M**. **2015b**. MUCILAGE-RELATED10 Produces Galactoglucomannan That Maintains Pectin and
Cellulose Architecture in Arabidopsis Seed Mucilage. *Plant Physiology* **169**: 403–420.

Voiniciuc C, Yang B, Schmidt MH-W, Günl M, Usadel B. **2015c**. Starting to gel: how Arabidopsis
seed coat epidermal cells produce specialized secondary cell walls. *International Journal of
Molecular Sciences* **16**: 3452–3473.

Voiniciuc C, Zimmermann E, Schmidt MH-W, Günl M, Fu L, North HM, Usadel B. **2016**.
Extensive Natural Variation in Arabidopsis Seed Mucilage Structure. *Frontiers in Plant Science* **7**.

Vollen K, Zhao C, Alonso JM, Stepanova AN. **2024**. Sourcing DNA parts for synthetic biology
applications in plants. *Current Opinion in Biotechnology* **87**: 103140.

Wang Y, Azhar S, Gandini R, Divne C, Ezcurra I, Aspeborg H. **2015**. Biochemical characterization
of the novel endo-β-mannanase AtMan5-2 from Arabidopsis thaliana. *Plant Science* **241**: 151–
163.

Wang A, Wang X, Ren Y, Gong X, Bewley JD. **2005**. Endo-β-mannanase and β-mannosidase
activities in rice grains during and following germination, and the influence of gibberellin and
abscisic acid. *Seed Science Research* **15**: 219–227.

Wu Q, Meng YT, Feng ZH, Shen RF, Zhu XF. **2023**. The endo-beta mannase MAN7 contributes to
cadmium tolerance by modulating root cell wall binding capacity in *Arabidopsis thaliana*. *Journal
of Integrative Plant Biology* **65**: 1670–1686.

Yagi H, Tateo S, Saito T, Ohta Y, Nishi E, Obitsu S, Suzuki T, Seetaha S, Hellec C, Nakano A, *et al.*
2024. Deciphering the sub-Golgi localization of glycosyltransferases via 3D super-resolution
imaging. *Cell Structure and Function* **49**: 47–55.

Yang B, Stamm G, Bürstenbinder K, Voiniciuc C. **2022**. Microtubule-associated IQD9
orchestrates cellulose patterning in seed mucilage. *New Phytologist* **235**: 1096–1110.

Yu L, Lyczakowski JJ, Pereira CS, Kotake T, Yu X, Li A, Mogelsvang S, Skaf MS, Dupree P. **2018**.
The Patterned Structure of Galactoglucomannan Suggests It May Bind to Cellulose in Seed
Mucilage. *Plant Physiology* **178**: 1011–1026.

Yu L, Shi D, Li J, Kong Y, Yu Y, Chai G, Hu R, Wang J, Hahn MG, Zhou G. **2014**. CELLULOSE
SYNTHASE-LIKE A2, a Glucomannan Synthase, Is Involved in Maintaining Adherent Mucilage
Structure in Arabidopsis Seed. *Plant Physiology* **164**: 1842–1856.

Yu L, Yoshimi Y, Cresswell R, Wightman R, Lyczakowski JJ, Wilson LFL, Ishida K, Stott K, Yu X,
Charalambous S, *et al.* 2022. Eudicot primary cell wall glucomannan is related in synthesis,
structure and function to xyloglucan. *The Plant Cell*: koac238.

Zhang R, Li B, Zhao Y, Zhu Y, Li L. **2024**. An essential role for mannan degradation in both cell
growth and secondary cell wall formation. *Journal of Experimental Botany* **75**: 1407–1420.

Zhong R, Cui D, Ye Z. **2019**. Evolutionary origin of O-acetyltransferases responsible for
glucomannan acetylation in land plants. *New Phytologist* **224**: 466–479.

Supporting Information

Additional Supporting Information may be found online:

Methods S1 Supplemental Methods for Plant and Yeast Experiments

Fig. S1 Additional phenotyping of *man2 man5* seeds.

Fig. S2 Mucilage architecture after *CSLA2* overexpression in the seed coat.

Fig. S3 MAN2 and MAN5 protein structure and expression in yeast.

Fig. S4 Polymer composition following CSLA/MAN expression in *Pichia*.

Fig. S5 Fractionation of alkaline-soluble and insoluble cell wall glycans.

Fig. S6 AtMAN2 truncation alters its localization and its hydrolytic effects.

Fig. S7 Transmission electron microscopy of *Pichia* cells.

Fig. S8 Confocal microscopy of MannanTrack in yeast cells.

Fig. S9 Workflow of water-soluble mannan extraction from yeast and plant samples.

Table S1 Sequences of primers used for genotyping and quantitative PCR.

Table S2 Sequences of primers used for cloning.

Table S3 Monosaccharide composition of total mucilage extracts.

Table S4 Plant constructs and functional evaluation.

Table S5 Yeast strains used in this study.

Table S6 Summary of characterized plant MANs.

Video S1 Timelapse of Golgi-YFP and MAN2-mCherry in Tobacco

none712 **Figure Legends**

Fig. 1. Mannanase double mutant Arabidopsis seeds have decreased mucilage accumulation. (a) T-DNA insertions in *AtMAN2* and *AtMAN5* genes, showing exon and intron structure. Red arrows indicate positions of primer binding utilized for quantitative PCR (Scale bar, 200 bp). (b) Fold changes in *AtMAN2* and *AtMAN5* or *AtCSLA2* and *AtMUCI10* transcription relative to WT in mature green siliques. Data show mean + SD of three biological replicates normalized to the geometric mean of *UBQ5* and *GAPC1*. Asterisks indicate significant decreases relative to WT (Student's *t* test, *P* < 0.01). (c) Pectin release of WT and mutant seeds stained with RR. (d) Projected mucilage and (e) seed surface area. Data shows mean + SD of three biological replicates (around 20 seeds each). Asterisks indicate significant decreases relative to WT (Student's *t* test, *P* < 0.05). Bars: (c) 200 μm.

Fig. 2. The *man2 man5* mutant seeds are impaired in β-mannan elongation. (a) Schematic of mucilage mannan biosynthesis and MAN2/5 protein domains. (b) Mean + SD germination rate for three biological replicates. For each time point, asterisks indicate significant decreases relative to WT (Student's *t* test, $P < 0.05$). (c) Absolute monosaccharide levels of three biological replicates. Different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$).

Fig. 3. Golgi-localized MAN enzymes are important for mucilage structure. (a) CF staining of β-glucans in seed mucilage. Arrows mark disordered rays. (b) Length of CF-stained rays. (c) *AtCSLA2* overexpression affects Gal and Man content, and (d) total mucilage release. Biochemical (e) and mucilage RR staining (f) complementation of *man2 man5* by AtMAN gDNA constructs. Different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$). (g) Optical sections of *N. benthamiana* leaves expressing a Golgi-YFP marker and AtMAN2-mCherry or an mCherry control. Arrows mark co-localization. Bars: (a) 100 µm, (f) 200 µm, (g) 30 µm.

Fig. 4. MAN2/5 enzymes reduce insoluble mannan content in yeast. (a) Relative Man content of insoluble yeast polymers. Box plots show three biological replicates. (b) Arrows mark the co-localization of AkCSLA3-Venus and AtMAN2/5-mCherry and counterstained with CF in confocal micrographs acquired with an 86x (1.2 NA) objective. (c) AKI polymers made by AtCSLA2 plus AtMAN2 or the fungal AaMAN5a. (d) AZCL-galactomannan digestion assays using *Saccharomyces* cell mixtures and supernatant, alongside commercial MAN controls (Megazyme, E-BMANN). Different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$). Magenta labels indicate a C-terminal protein fusion to mCherry. Bars: (b) 5 µm.

Fig. 5. *MAN2* and *MAN5* promote mannan solubility and secretion. (a) MannanTrack probe components in *Pichia* control strains to validate the blue (BFP), yellow (YFP) and red (RFP) protein fluorescence patterns. (b) Relative secretion (supernatant RFP / cell mixture RFP) of the PaCBM35-mScarlet, normalized to the BFP cell fluorescence relative to empty vector (EV). (c) Number of cells with detectable AkCSLA3-Venus and PaCBM35a-mScarlet fluorescence quantified via flow cytometry. (d) Man content of hot water (H_2O-soluble) and AKI polymers in *Pichia* extracts. (e) Man content of hot water (H_2O-soluble) and AIR polymers in Arabidopsis ground seeds. Box plots show three biological replicates and different letters denote significant changes with one-way ANOVA (Tukey's post-hoc test, $P < 0.01$). Magenta labels indicate a C-terminal protein fusion to mCherry. (f) Updated model of β-mannan biosynthesis in the Golgi apparatus (based on Voiniciuc, 2022) with AtMAN2/5 role(s). Bars: (a) 5 μm.