A TNL receptor mediates microbiome feedbacks in Arabidopsis

Authors:

Henry Janse van Rensburg[1*], Niklas Schandry[2,3*§], Jan Waelchli[1], Katja Stengele[1], Selma Cadot[1],

Katharina Jandrasits[2], Claude Becker[2,3,§], Klaus Schlaeppi[1,§]

[1] Department of Environmental Sciences, University of Basel, Basel, Switzerland

[2] Gregor Mendel Institute of Molecular Plant Biology GmbH, Austrian Academy of Sciences, Vienna

BioCenter (VBC), Vienna, Austria

[3] LMU Biocenter, Faculty of Biology, Ludwig-Maximilians-Universität Munich, Martinsried,

Germany

[*] Contributed equally

[§] **Corresponding authors:**

Niklas Schandry: n.schandry@bio.lmu.de

Claude Becker: claude.becker@lmu.de

Klaus Schlaeppi: klaus.schlaeppi@unibas.ch

ORCID identifiers:

H. J.v.R.:   0000-0002-5632-8422

N. S.:   0000-0003-3099-7860

K. St.:   0009-0008-3843-2868

J. W.:   0000-0002-4384-6570

K. J.:   0000-0002-0161-582X

C. B.:   0000-0003-3406-4670

K. S.:   0000-0003-3620-0875

Author contributions:

H.J.v.R., N.S., C.B., and K.S., designed the research; H.J.v.R. and K. St. performed greenhouse and

S.C. the field soil conditionings; K.J and N.S. performed the phenotyping of accessions and GWA

analysis; H.J.v.R. performed GWA candidate gene selection, MBOA sensitivity assays, genetic

characterisation of Arabidopsis mutants, Arabidopsis mutant feedback, microbiome and transcriptome

experiments and analysis; J.W., performed bioinformatic analysis for the microbiome and transcriptome

analyses; C.B., provided funding and access to high-throughput phenotyping infrastructure; H.J.v.R.

and N.S. analysed the data; H.J.v.R., N.S. and K.S. wrote the manuscript. All authors revised the paper.

Manuscript information:

Number of Figures: 5 (all colour)

Number of Supplementary Materials figures: 9

Number of Supplementary Materials tables: 3

Declarations of interest:

none

Abstract

44
Plant performance depends on the soil microbiome. While microbiome feedbacks are well documented,
the mechanisms by which plants perceive and mediate these feedbacks remain unclear. We established
a framework using two distinct microbiomes in the same soil that differently affect plant growth.
Screening 410 *Arabidopsis thaliana* accessions revealed substantial variation in growth feedbacks,
which we used for genome-wide association mapping to identify Mediator of Microbiome Feedback 1
(MMF1), an immune receptor. Characterisation in the reference accession Col-0 revealed that *mmf1*
mutants lack the differential growth feedbacks, have an altered root bacterial community, and fail to
induce a defence-related transcriptional response observed in wild-type plants. The discovery of MMF1
implies that integration of microbial effector signals optimises host microbiome composition and
immune status to enhance growth.

Keywords

Growth feedback, GWAS, Arabidopsis, soil microbiome, TNL receptors

Main

Microbiomes are widely recognized for supporting their hosts with many vital services leading to improved host performance. Similar to the gut microbiome of humans, plant microbiomes contribute to nutrition homeostasis, stress tolerance and host health (*1*). A large knowledge gap in all fields of host-microbiome interactions, is how hosts genetically control their microbiome. For plants, the performance depends to a large extent on below-ground bacteria, fungi and other microbes that collectively function as a soil microbiome (*1*, *2*). Soil microbiomes consist of functionally diverse members providing beneficial services to plants, including nutrient acquisition, growth promotion, stress tolerance, and suppression of plant pathogens, but soil microbiomes can also contain pathogenic members that reduce plant fitness (*3*, *4*). Plant responses, which are causally linked to a soil microbiome, are referred to as microbiome feedbacks (*5*). This term is akin to plant-soil feedbacks describing the ecology of soil properties (more broadly, but including the microbiome) that influence plant performance (*6*, *7*). Microbiome feedbacks are highly relevant in agricultural contexts, as soil microbiome composition has been linked to agronomic traits of crops such as maize and potato (*8*, *9*), to field productivity (*10*), and to over- and under yielding areas in fields (*11*, *12*). While different soil microbiomes have often been seen as drivers of differential plant responses (*13*, *14*), little is known about the mechanisms by which plants perceive complex soil microbiomes and mediate microbiome feedbacks.

Plants have evolved a sophisticated immune system that regulates interactions with beneficial microbes while defending against pathogens (*15–17*). Plant immunity operates through two key pathways: pattern-triggered immunity (PTI) and effector-triggered immunity (ETI). PTI is initiated at the cell surface by pattern-recognition receptors that recognize microbe-associated molecular patterns and provides basal defence to deter non-adapted microbes. In contrast, ETI is activated within the cytoplasm by nucleotide-binding leucine-rich repeat receptors (NLRs), which detect pathogen effectors or their activities, leading to definite immune responses (*18–20*). NLRs are divided into Toll/interleukin-1 receptor (TIR-NLRs; TNLs) and coiled-coil NLRs (CNLs). *Arabidopsis thaliana* (hereafter: Arabidopsis) encodes 54 TNLs and 88 CNLs, highlighting the complexity of the plant immune system (*21*). Both PTI and ETI depend on signalling molecules such as salicylic acid (SA) and reactive oxygen species (ROS) (*19*, *20*). Although these pathways were primarily studied in the context of binary interactions with pathogenic or beneficial microbes, recent evidence suggests that PTI and ETI also influence plant-associated microbiomes. These immune pathways shape microbiome composition, modulating microbial interactions and contributing to overall plant health and resilience (*16*, *22–24*).

To study microbiome feedbacks, we developed an experimental framework based on two distinct soil microbiomes in the same soil type. These soil microbiomes are generated by growing wild-type maize, which produces benzoxazinoids (BXs), or BX-deficient mutant maize. BXs are bioactive compounds of *Poaceae* (*25*) that, among multiple functions, shape the root and rhizosphere microbiomes of plants (*14*, *26*, *27*). After three months of growth, the wild-type and mutant plants are

removed, leaving behind BX-conditioned (BX_{plus}) and its control (BX_{minus}) soil microbiomes,

respectively. We have described growth and defence feedbacks of maize and wheat in response to

BX_{minus} and BX_{plus} soil microbiomes (*14*, *28*), and we have proven agronomic relevance of these

microbiome feedbacks (*28*, *29*). Through sterilization and microbiome complementation experiments,

we demonstrated that the feedbacks are BX-dependent but that the soil microbiomes drive the

differential plant responses (*14*, *28*, *29*). Recently, we found that the model plant Arabidopsis also

expressed microbiome feedbacks (*30*). The reference accession Columbia-0 (Col-0) responded with

improved growth and enhanced resistance when grown on the BX_{plus} compared to the BX_{minus} soil

microbiome. As plausible mechanistic explanation for this apparent lack of a growth-defence trade-off,

the plants growing on the BX_{plus} soil microbiome were found to be primed for their SA defences (*31*).

Microbiome feedbacks are often influenced by the genotype of the response plant. For instance,

rhizobacteria-induced growth promotion varied significantly across Arabidopsis accessions (*32*).

Similarly, genotype-specific resistance against Fusarium has been observed in tomato, where the

outcome of the interaction depends on the tomato genotype and the composition of the native soil

microbiome (*13*). In our work with maize and wheat responding to BX_{minus} and BX_{plus} soil microbiomes,

we also observed a genetic component in microbiome feedbacks (*14*, *28*). Here, we harnessed the

genetic diversity among Arabidopsis accessions to study their growth feedbacks as a function of the

plant genotype and the soil microbiome (**fig. S1**).

Results

Arabidopsis displays substantial phenotypic variation in microbiome feedbacks

We grew 410 natural, genome-sequenced Arabidopsis accessions (*33*) in both control (BX_{minus}) and BX-conditioned (BX_{plus}) soil microbiomes in a climate chamber equipped with an automated top-view imaging system. The process from imaging to determination of rosette area and feedback calculation is illustrated with the two contrasting accessions IP-Hue-3 and Hadd-3, which grew better and worse on the BX_{plus} soil microbiome, respectively (**Fig. 1ab**). Plants were imaged twice a day and plant morphometric traits extracted using ARADEEPOPSIS (*34*). We focussed on the first 20 days after germination (DAG) to avoid confounding phenotyping effects such as early flowering and overlapping leaves. We used rosette area as a measurement of plant size to calculate the daily growth feedback as the difference in rosette area between BX_{plus} and BX_{minus} microbiomes, i.e. the accessions IP-Hue-3 and Hadd-3 exhibited positive and negative feedbacks, respectively (**Fig. 1b**).

The screening of accessions revealed substantial variation in growth feedbacks to BX_{plus} and BX_{minus} microbiomes (**Fig. 1c; Dataset 1** and **Suppl. File 1**). Across the 410 accessions, 48 % (196 accessions) displayed positive feedbacks, while 52 % (214 accessions) showed negative feedbacks. At the two extremes of the distribution, the two soil microbiomes accounted for ~2-fold (IP-Hue-3) and >3-fold (Hadd-3) differences in rosette area. Consistent with earlier work, Col-0 expressed a subtle positive feedback (*30*). Feedback responses were independent of the accessions' geographical origin (**Fig. 1d, fig. S2a**).

Since BXs are allelopathic to plants (*35*, *36*), we tested whether the accession's growth differences could be due to varying sensitivity to BXs. BXs differ between the two soil variants, with MBOA as the most abundant BX compound (*29*, *37*). We screened the subset of accessions with the strongest opposing growth feedbacks for sensitivity to MBOA. All accessions exhibited shorter roots in the presence of MBOA (**fig. S3a, Dataset 2** and **Suppl. File 2**), which is indicative of phytotoxicity. However, there was no correlation between feedback response and MBOA sensitivity (**fig. S3b**). These results indicate that the observed growth feedbacks are not linked to varying sensitivity to the BXs present in the soil.

In conclusion, we found substantial phenotypic variation in Arabidopsis growth feedbacks to two soil microbiomes, providing a powerful resource for association studies on genetics that drive the differences in feedbacks.

Figure 1. **Natural variation in growth feedbacks among Arabidopsis accessions**. (**a**) Growth of the two *A. thaliana* accessions (IP-Hue-3 and Hadd-3) that exhibited the largest opposing growth feedbacks in response to the BX_{minus} and BX_{plus} microbiome at the conclusion of the experiment (20 days after germination (DAG)). (**b**) The daily rosette area measured in pixels extracted by ARADEEPOPSIS (left), and the daily microbiome growth feedback ($BX_{plus} - BX_{minus}$) (right) for the first 20 DAGs for IP-Hue-3 and Hadd-3 grown on the two soil microbiomes. Shaded areas (left) and coloured bars (right) indicate positive (blue, IP-Hue-3) or negative (orange, Hadd-3) microbiome feedbacks in response to the BX_{plus} microbiome. (**c**) Microbiome feedback of the 410 *A. thaliana* accessions used in the genome-wide association study at the conclusion (DAG 20) of the experiment. The position of IP-Hue-3, Hadd-3 and Col-0 are indicated on the graph. Data is from two phenotyping experiments (experiment I and II, **Table S1**), each consisting of a subset of 210 accessions (3 replicates per accessions per soil), with 10 overlapping accessions in each of the experiments. (**d**) The map shows the geographic distribution of these accessions, coloured by their microbiome feedback (blue for positive; orange for negative).

Candidate gene loci are involved in plant growth and defence

We performed a genome-wide association study (GWAS) using the phenotypic variation in growth

feedbacks of the 410 accessions (**Fig. 1c**) and employed a linear mixed model accounting for population

structure among the used Arabidopsis accessions (*38*). Noting temporal variation in daily growth

feedbacks (**fig. S2b**), we ran separate GWAS on daily data, and calculated associations between single

nucleotide polymorphisms (SNPs) and the phenotypes for each of the 20 DAGs (**Dataset 3**).

Across daily GWAS runs, we identified three genomic regions with significant associations to

the growth feedback (**Fig. 2a; Dataset 3** and **Suppl. File 3**). Genes in the regions of the three peaks are

related to plant growth and plant defence. We then focused on SNPs that were significant for multiple

consecutive days in the repeated GWAS (see Methods). The top SNP (chromosome 3, position

16203882) was in peak #2 (**Fig. 2b**), of which an alternative allele was found in 11 accessions (**Fig.**

2c,d). These accessions expressed better growth in response to the BX_{plus} microbiome compared to

accessions with the reference allele of the SNP (**Fig. 2c**). This SNP is in close proximity to the TNL

receptor gene AT3G44630 (position 16200156; 3.7kb upstream) and a more distal second TNL

(AT3G44670; position 16218427; 14.5kb downstream; **Fig. 2b**). We focussed on the uncharacterized

AT3G44630 because of the closer proximity to the top SNP, and its differential expression in roots of

Arabidopsis grown on BX_{minus} and BX_{plus} soils (*30*). We named this gene *Mediator of Microbiome*

Feedback 1 (*MMF1*). Taken together, GWAS identified a genomic locus on chromosome 3 that includes

MMF1 as a candidate gene that could explain the differential growth feedbacks of Arabidopsis

accessions in response to the BX microbiomes.

Figure 2. **Genetic association mapping of growth feedbacks in response to the BX$_{plus}$ microbiome**. (**a**) Manhattan plot showing the -log$_{10}$-transformed *p*-values for the top 500 single nucleotide polymorphisms (SNPs) across the 5 chromosomes of *A. thaliana*. SNPs that are significantly associated with growth feedbacks after Bonferroni correction are shown in red. The biological functions of genes adjacent to the top three SNPs are indicated. (**b**) The genomic region (50 kb) flanking the SNP at position 16,203,882, including the two TNL genes AT3G44630 and AT3G44670. (**c**) Average growth feedback among accessions carrying the alternative allele of the SNP at position 16,203,882 (positive feedback) compared to those with the reference allele (negative feedback). (**d**) Overall positive microbiome feedback distribution of accessions carrying the alternative allele of the SNP at position 16,203,882 (black lines and filled circles) on the growth feedback distribution curve at the conclusion of the experiment.

MMF1 mediates differential microbiome feedbacks

To test if MMF1 functions in growth feedbacks, we phenotyped two T-DNA insertion lines in the Col-

0 background for their response to the BX_{minus} and BX_{plus} microbiomes. The *mmf1-1* and *mmf1-2* lines

are knockdown and knockout mutants, respectively (**fig. S4**). Both mutants grew similarly to wild type,

albeit slightly smaller, and showed no obvious disease symptoms (**Fig. 3a; Dataset 4** and **Suppl. File**

4). Consistent with earlier work (*30*) and the GWAS experiment (**Fig. 1c**), wild- type plants expressed

subtle positive growth feedbacks in response to the BX_{plus} microbiome, with higher fresh weight and

larger rosette area compared to those growing on the BX_{minus} microbiome (**Fig. 3b**).

203
Figure 3. **MMF1 is involved in growth feedback responses to the BX_{plus} microbiome**. (**a**) Phenotypes of 6-
week-old Col-0 wild-type (WT) and two *mmf1* T-DNA insertion mutants (AT3G44630) grown on the BX_{minus} and
BX_{plus} soil microbiomes (experiment IV, **Table S1**). (**b**) Fresh weight (left) and rosette area (right; in pixels) of
the indicated genotypes grown for 6 weeks on either a BX_{minus} or BX_{plus} microbiome. Sample sizes are indicated
at the bottom of each plot. Boxplots illustrate the median, standard error, and interquartile range, and the whiskers
extend from the box to the smallest and largest values. Output of the ANOVA and the FDR corrected *p*-values
are indicated in each plot. Results were confirmed in three independent repetition experiments (experiment III, IV
and V) on two independently conditioned soil batches.

In contrast, both *mmf1* mutants lost the positive growth feedback, reflected in fresh weight and rosette area. This was confirmed in two additional experiments using independent soil batches (**fig. S5a,b**). This functional validation allowed to conclude that, in Col-0, a functional MMF1 is required for the differential growth feedback in response to the two soil microbiomes.

The plant immune system relies on two TNL signalling pathways: Enhanced Disease Susceptibility 1 (EDS1) together with Phytoalexin Deficient 4 (PAD4) lead to resistance, and EDS1 jointly with Senescence-Associated Gene 101 (SAG101) induce hypersensitive responses and cell death (*39*). To test the involvement of the two TNL signalling pathways in microbiome feedbacks, we performed feedback assays with *eds1*, *pad4* and *sag101* mutant plants (*40*, *41*). These mutants showed no altered growth phenotypes or obvious disease symptoms in response to BX_{minus} and BX_{plus} soil microbiomes (**fig. S5c**). Similar to wild type, the three immune signalling mutants expressed positive growth feedbacks in response to the BX_{plus} soil microbiome (**fig. S5d**). Hence, single gene knockouts in EDS1, PAD4 and SAG101 do not disrupt immune signalling that leads to growth feedbacks, or alternatively, suggest that this process may be independent of these two classical pathways.

MMF1 affects root bacterial community assembly sourced from BX_{plus} soil microbiome

In earlier work, we found differential microbiomes on roots of plants with differential growth feedbacks (*14*, *30*). We therefore determined the root bacterial, fungal and oomycete communities that wild-type and *mmf1* mutant plants source from BX_{minus} and BX_{plus} soils. The detailed microbiome analysis is documented in **Dataset 5** and **Suppl. File 5**. Shannon diversity did not differ between the factors *microbiome* (BX_{minus} and BX_{plus}) and *line* (WT and *mmf1-2*; **fig. S6a**). We used Permutational Multivariate Analysis of Variance (PERMANOVA) to assess the impact of these factors on root community composition. *Microbiome* alone accounted for little variation in bacterial ($R^2 = 3.9\%$, $p = 0.09$), fungal ($R^2 = 3.7\%$, $p = 0.25$) and oomycete communities ($R^2 = 5.5\%$, $p < 0.01$; **Fig. 4a**). However, we found significant variation in bacterial – but not fungal or oomycete – communities between wild type and *mmf1* mutants ($R^2 = 5.7\%$, $p < 0.01$) and in the interaction of the factors *line* with *microbiome* ($R^2 = 4.2\%$, $p < 0.05$). The interaction effect indicates that the root microbiome of the mutant differed from wild type in only one soil. Reduced pairwise PERMANOVA confirmed this with more and significant variation between wild type and mutants in the BX_{plus} ($R^2 = 15.7\%$, $p < 0.01$) compared to the BX_{minus} ($R^2 = 4.6\%$, $p = 0.605$) soil microbiome. In contrast, fungal and oomycete communities showed no differences between *lines* and *microbiomes* (**fig. S6b**). Constrained Analysis of Principal Coordinates (CAP) confirmed these findings: root bacterial communities clearly separated between wild type and mutant on the BX_{plus} but not BX_{minus} soil microbiome (**Fig. 4b**), while fungal and oomycete communities showed minimal, non-significant separation (**fig. S6b**). Taken together, the *mmf1* mutant assembled a distinct bacterial root microbiome relative to wild-type plants, but only on the BX_{plus} source microbiome.

Figure 4. **Root microbiome profiling of wild-type and *mmf1-2* mutants grown on the BX$_{plus}$ or BX$_{minus}$ soil**.
(**a**) The output of PERMANOVA on Bray-Curtis (BC) dissimilarities of bacterial, fungal and oomycete root
microbiome of Arabidopsis wild-type (WT; Col-0) and *mmf1-2* mutants grown on the BX$_{minus}$ or BX$_{plus}$ soils
(sampled from experiment IV, **Table S1**). The R^2 and *p*-values indicate the *line* and *microbiome* (conditioning)
effects and their interaction. Pairwise PERMANOVA for bacterial communities compares WT and *mmf1-2*
mutants on each soil condition; significant effects ($p < 0.05$) are indicated in bold. (**b**) Constrained Analysis of
Principal Coordinates (CAP) for WT (filled circles) and *mmf1-2* mutants (open triangles) grown on the BX$_{minus}$
(green) or BX$_{plus}$ (yellow) soils confirming the effects of the soil conditioning and lines observed in the
PERMANOVA analysis. Ellipses represent 95% confidence intervals for plants grown on BX$_{minus}$ (green) or
BX$_{plus}$ (yellow) soils. Axes show the percentage of variance explained by the first two principal coordinates.

MMF1 is required for a defence-related transcriptional response

To obtain first insights into how MMF1 contributes to positive growth feedbacks, we analysed the root
transcriptomes of wild-type and *mmf1-2* mutants in response to both soil microbiomes (detailed
transcriptome analysis documented in **Dataset 6** and **Suppl. File 6**). Analogous to the microbiome
analysis, we first examined the contributions of the factors *microbiome* (conditioning) and *line* on the
transcriptome using PERMANOVA. *Microbiome* accounted for 29.0% (based on R^2, $p < 0.01$), *lines*
for 33.7% ($p < 0.01$) and their interaction for 12.4% ($p < 0.01$) of the transcriptomic variation (**Fig. 5a**).
Again, the significant interaction term revealed that mutant and wild type responded differently to the
two soil microbiomes. Reduced pairwise PERMANOVA revealed again greater variation in wild-type
($R^2 = 79.7\%$, $p < 0.01$) compared to mutant ($R^2 = 10.8\%$, $p = 0.349$) plants in response to the two
microbiomes. Principal Component (PC) analysis confirmed a clear separation between the two source
microbiomes (PC1, 48% of variation) and plant lines (PC2, 16% of variance; **Fig. 5b**). Overall, the
strongest separation was between wild-type plants in response to the two microbiomes, whereas the
mmf1 mutant transcriptomes clustered together.

Figure 5. **Plants lacking MMF1 fail to mount a transcriptional response to a BX microbiome**. (**a**) The output
of PERMANOVA on Bray-Curtis dissimilarities of the root transcriptome of wild-type (WT) and *mmf1-2* mutant
plants in response to the BX_{minus} and BX_{plus} microbiomes (sampled from experiment IV, **Table S1**). The R^2 and *p*-
values summarize the effects of *line* (plant genotype) and *microbiome* (soil condition) along with their interaction.
Pairwise PERMANOVA results compare the transcriptomes of WT and *mmf1-2* mutant plants when grown on
the BX_{minus} soil microbiome compared to the BX_{plus} soil microbiome; significant effects are indicated in bold. (**b**)
Principal component analysis (PCA) of the same root transcriptome confirms the *line* (WT, closed circles; *mmf1-*
2, open triangles) and *microbiome* (BX_{minus}, green; BX_{plus}, yellow) effect of the PERMANOVA results. The axes
display the percentage of variance explained by each principal component. (**c**) MA plots illustrating differentially
expressed genes (DEGs) in WT between BX_{minus} and BX_{plus} (left), and the absence of DEGs in *mmf1-2* (right).
Genes upregulated on the BX_{plus} microbiome (yellow) or BX_{minus} microbiome (green) are colour coded. Genes
with a $\log_2$ fold-change ≥ 1 and false discovery rate < 0.05 (empirical Bayes' shrinkage; DESeq2) are considered
significant.

Next, we analysed the transcriptomes at gene level determining differentially expressed genes (DEGs;

$\log_2$ fold-change ≥ 1; empirical Bayes shrinkage, *p* values adjusted for false discovery rate < 0.05).

Aligned with the microbiome analysis (**Fig. 4**), we found more DEGs between the lines in response to the BX_{plus} (347 total; 230 up and 117 down in wildtype) compared to the BX_{minus} soil microbiome (128 total; 68 up and 60 down; **fig. S7**). Wild-type plants had 563 DEGs in response to the two microbiomes (107 down and 456 up, **Fig. 5c**). Surprisingly, not a single microbiome specific DEG was found in mutant plants (**Fig. 5d**). Taken together, a functional MMF1 leads to a strong transcriptional reprogramming, mostly in response to the BX_{plus} soil microbiome.

Finally, we performed co-expression and gene ontology (GO)-term enrichment analysis (underlying genes documented in **Dataset 6**) to characterise the transcriptional changes of wild-type plants in response to the two microbiomes (**Fig. 5c**). Hierarchical clustering identified clusters of co-expressed genes: four up- and three downregulated (**fig. S8**). Upregulated processes included hypoxia (clusters #2 and #3) and immune responses (clusters #2 to #5). Processes linked to hypoxia included the GO terms response to hypoxia and cellular response to hypoxia (**Dataset 6**). Upregulated immune processes included GO terms such as response to ethylene and systemic acquired resistance (cluster #2), defence response to insects, and response to bacterial molecules (cluster #3), indole metabolism and response to salicylic acid (SA; cluster #4), and hypersensitive response, programmed cell death, and interaction with symbionts (cluster #5; see **Dataset 6** for the genes in each cluster). Of note, several of these genes are typically involved in TNL-triggered immunity (*39*, *42*). Downregulated processes included fatty acid biosynthesis, secondary metabolite biosynthesis and malate transport (clusters #6 to #8; **Dataset 6**). In conclusion, wild-type plants, which achieved better growth in response to the BX_{plus} soil microbiome, exhibited a strong transcriptional reprogramming, most noticeably an upregulation of defence-related genes.

Discussion

While genotype-dependent microbiome feedbacks have been observed in many plant systems (*14*, *28*, *29*, *32*), the mechanisms underlying plant responses to complex microbiomes remain largely unresolved. Here, we used natural variation in growth feedbacks to BX_{plus} and BX_{minus} soil microbiomes among Arabidopsis accessions (**Fig. 1c**) to uncover candidate loci through which plants respond to different microbiomes in natural soils. We focussed on the gene AT3G44630, which was in the proximity of the top SNP in GWAS (**Fig. 2**), named this TNL receptor gene *MMF1* and characterized it in the reference accession Col-0. Arabidopsis *mmf1* mutants lack three hallmarks of wild-type plants in response to the BX_{plus} soil microbiome: they lose the positive growth feedback (**Fig. 3**), have an altered root bacterial community (**Fig. 4**), and fail to trigger the defence-related transcriptional response (**Fig. 5**). Given the molecular function of TNLs - to recognise microbial effectors intracellularly and trigger ETI (*20*) – our results support the following mechanistic model (**fig. S9**): When grown on the BX_{plus} soil microbiome, wild-type plants show enhanced shoot growth resulting from a defence-related transcriptional response (**fig. S8**). This MMF1-dependent transcriptional response, together with the typical intracellular localisation of TNLs, imply that a bacterial (see below) effector of the BX_{plus} microbiome is perceived inside the root cells. This model is consistent with the reduced shoot growth of wild-type plants on the BX_{minus} microbiome (in the absence of the effector) as well as *mmf1* mutants on the BX_{plus} microbiome (lack of the receptor), and is further supported by the lack of a defence-related transcriptome in *mmf1* mutants.

The plant responses, including a defence-related transcriptome and enhanced shoot growth, coincided with an altered root microbiome (**fig. S8**). While overall root microbiomes were similar between soil sources, the bacterial communities of wild-type and *mmf1* mutant plants differed significantly on the BX_{plus} but not on the BX_{minus} microbiome (**Fig. 4**). These data do not allow to establish a causal link between the altered root microbiome and the differential shoot growth of the genotypes. However, the observation suggests that a bacterial effector in the BX_{plus} microbiome, which presumably triggers the defence-related transcriptome via the TNL MMF1, may be responsible. Identification of the bacterial species and the actual effector are obvious future research targets.

As part of the immune system, TNLs act as gatekeepers against opportunistic pathogens, maintaining host health and microbiome homeostasis (*20*). Earlier work showed that Arabidopsis with impaired ETI undergo microbiome dysbiosis in their leaves, and ultimately these plants have reduced fitness (*24*, *43*). Analogously, in roots, we report microbiome dysbiosis in the ETI mutant *mmf1* and demonstrate that MMF1 is required for growth promotion in wild-type plants. Both examples imply that TNLs integrate microbiome signals inside host cells for maintaining microbiome homeostasis and host health. While TNLs have been predominantly studied in single-pathogen and gene-for-gene interactions, they face a multiplicate challenge at the microbiome level, with many more effectors being present in a diverse microbiome (*16*). Thus, plant immune responses rely on more complex signal integration when interacting with microbial communities compared to single isolates (*16*, *43*).

Moreover, there is evidence that signal integration may further depend on TNL receptor dosage:
recently, high TNL copy numbers within the *RPP-1*-like gene cluster among Arabidopsis accessions
were linked with reduced bacterial diversity in their phyllosphere (*44*). Of note, the closest homolog of
MMF1 is Recognition of Peronospora Parasitica 1 (RPP1) (*45*), and additional TNLs are in the
proximity of the SNP most strongly associated with the variation in growth feedback among the tested
accessions (**Fig. 2b**). Here, we have shown that lack of MMF1 is sufficient for the loss of growth
feedback, but future work is needed to test whether additional members of the TNL family or their
dosage also contribute to feedback in other accessions.

We set out to identify a first genetic component through which plants respond to different
microbiomes in natural soils. The substantial variation in growth feedbacks among Arabidopsis
accessions (**Fig. 1c**) implies that there are additional components to discover. The TNL MMF1 presents
a first genetic component for such feedbacks, but for now it only explains the enhanced growth and
increased defence of the reference accession Col-0 responding to the BX_{plus} microbiome. MMF1 and
the top SNP in its proximity alone cannot explain the massive variation in growth feedbacks among the
other accessions. For instance, Hadd-3, which shares the reference allele of the top SNP with Col-0,
grew more poorly on the BX_{plus} microbiome, while IP-Hue-3 harboured the alternative allele and grew
much better than Col-0. Solving this puzzle requires further analysis of the complex locus around the
top SNP (*44*), and investigating the MMF1 alleles of the extreme accessions such as Hadd-3 and IP-
Hue-3. Together with the other GWAS candidate loci (**Fig. 2a**), additional genetic components likely
contribute to explaining the massive variation in growth feedbacks among the accessions.

In conclusion, the identified TNL receptor provides a relatively straightforward explanation for
the enhanced shoot growth of Col-0 on the BX_{plus} soil microbiome: MMF1 perceives an effector of the
BX_{plus} microbiome, triggers defences, and alters the root microbiome, which collectively coincide with
better shoot growth by yet unknown mechanisms. This work contributes to understanding host genetic
pathways by which plants perceive complex soil microbiomes and mediate microbiome feedbacks.
Such knowledge may pave the way towards crops that leverage beneficial microbiome feedbacks,
ultimately improving yield in a more sustainable manner.

Materials and methods

Plant material

The maize (*Zea mays* L.) genotypes B73 and *bx1* (B73) transposon insertion mutant (*46*) were used for soil conditioning (see below). The 410 *Arabidopsis thaliana* (hereafter: Arabidopsis) accessions used in the GWAS (see below) were obtained from the 1001 genomes project (The 1001 Genomes Consortium, 2016). For a full list of the accessions used, see **Dataset 1**. Arabidopsis T-DNA insertion mutants (AT3G44630) for *mmf1-1* (SALK_144159C) and *mmf1-2* (SALK_002318C) are in the Col-0 background and were obtained from the Nottingham Arabidopsis Stock Centre (NASC). Mutants were confirmed for homozygosity and characterised by qPCR (**fig. S4;** primers in **Table S3**). The mutant lines *eds1-12* (*40*), *pad4-1* (*47*), and *sag101-3* (*48*) in Col-0 background were kindly provided by Prof. Jane Parker.

Generation of BX$_{minus}$ and BX$_{plus}$ soil microbiomes

We used soil from an agricultural field site of Agroscope in Changins, Switzerland (46°24'00.0"N6°14'22.8"E; three adjacent plots, Parcel 29, Parcel 30, or Parcel 31) and we generate the distinct soil microbiomes either in the field or in the greenhouse using pots filled with that soil. Starting from replicates of the same batch of unconditioned field soil, the two distinct microbiomes are generated by growing of either wild-type maize (B73), which secrete benzoxazinoids (BXs) to the surrounding soil, or BX-deficient mutant maize (*bx1* (B73)). After three months of growth, the plants and roots are removed and the two maize lines "leave behind" BX-conditioned (BX$_{plus}$) and control (BX$_{minus}$) microbiomes in the soil. We then harvest either replicate soil cores (20 × 20 × 20 cm, conditioning in the field) or soil from replicate pots (3 L, 13.1 cm deep and 20.2 cm diameter; conditioning in the greenhouse), sieve the soil through a 1 cm^2 sieve and stored it at 4°C until the subsequent feedback experiments. We described in detail the approach of generating BX$_{minus}$ and BX$_{plus}$ soil microbiomes (we often referred to this as 'soil conditioning') including the physicochemical characteristics of the soil (*14, 28, 37*), management practices and cropping history of the field site (*14, 26*) and the conditions for greenhouse conditioning (*14, 28, 30*). The batches of conditioned soils for this study, and the experiments linked to soil batches are listed in **Table S1** and **S2**.

Microbiome feedback experiments

We list the setups and purposes of different experiments (Exp. I to Exp. VI) with Arabidopsis grown on BX$_{minus}$ and BX$_{plus}$ soil microbiomes in **Table S2**. For feedback experiments, the replicate soil samples containing the BX$_{plus}$ and BX$_{minus}$ microbiomes were individually amended with sand, by manually mixing in 20% (v/v) autoclaved quartz sand (*14*). Pots were equally filled by weight and randomized across trays. Below we detail the specific setups and conditions for the high-throughput phenotyping of the different Arabidopsis accessions (Exp. I and II) and the characterisation of wild-type and mutant lines in Col-0 background (Exp. III - VI).

High-throughput phenotyping of Arabidopsis accessions: We performed two microbiome

feedback experiments; each experiment was done with 210 accessions (10 accessions repeated in both

runs) resulting in a total of 410 tested Arabidopsis genotypes (**Dataset 1**). In each experiment, every

accession was seeded on both BX_{minus} and BX_{plus} soils with 3 replicates (2 experiments of 210 accessions

x 2 soils x 3 replicates = 1260 pots). Plant cultivation and phenotyping was previously described and

the images of the first experiment were utilized to develop the phenotyping software ARADEEPOPSIS

(without comparing BX_{minus} and BX_{plus} soils) (*34*). In brief, plants were grown at the Gregor Mendel

Institute of Molecular Plant Biology, Vienna, Austria, in a custom-built growth chamber (YORK

Refrigeration), equipped with an environmental control unit (Johnson Controls) and a climate control

unit (Incotec) under long-day conditions (16h light [21°C], 8h dark [16°C], 140 $\mu E/m^2 s$) with 60 %

relative humidity and watered twice per week by flooding. Plants were imaged twice daily, and plant

morphometric phenotypes extracted using the phenotyping software ARADEEPOPSIS (*34*). The

phenotypes were converted into "days after germination" phenotypes by taking the average of the two

measurements that happened on the same day. **Dataset 1** contains the source data and **Suppl. File 1**

documents the analysis.

Feedback experiments with Arabidopsis mutants: For microbiome feedback experiments

performed on Arabidopsis Col-0 mutant lines, we used greenhouse conditioned soils prepared as

described before. Seeds were stratified for 3 days in the dark at 4 °C and sowed on pots (5.5 × 5.5 × 5

cm) filled by weight. Plants were grown in a plant growth chamber (Percival, CLF Plant Climatics,

Wertingen, Germany) under short days with 10h light (100 $\mu mol\,m^{-2}\,s^{-1}$) (21 °C) and 14h darkness (18

°C) at 60% relative humidity. Plants were fertilized at day 14 and 21 after germination using 10 mL of

1:3 times diluted ½ strength Hoagland solution (*49*), watered 3 times per week by weight, and

randomized weekly. Analysis was blinded by adding barcodes to the pots for phenotyping which was

used to trace back sample names after analysis. Plants were photographed weekly and phenotyped using

the ARADEEPOPSIS pipeline to obtain the rosette area (plant region area) (*34*). Plants were grown for

6 weeks before the rosettes and roots were harvested for downstream analysis. **Dataset 4** contains the

source data and **Suppl. File 4** documents the analysis.

Genome-wide association study

Genome-wide association studies (GWAS) search for associations between single nucleotide

polymorphisms (SNPs) in a genome and a phenotype (*50*). As phenotype we took the differential growth

feedback of the accessions on the two soil microbiomes, i.e. the mean growth on BX_{minus} was subtracted

from the mean growth on BX_{plus} soil microbiomes [$BX_{plus} - BX_{minus}$]. We focussed on the first 20 days

after germination (DAG) to avoid confounding effects of phenotypes like early flowering and

overlapping leaves. GWA was performed using an in-house Nextflow pipeline

(https://gitlab.lrz.de/beckerlab/gwas-nf) that wraps limix (*38*) and applies the same multi-trait model

used previously in Huether et al. 2020. Only SNPs with a minor allele frequency of at least 1% were

included. The model estimates each SNP's "specific" effect on the trait "differential growth feedback". In this study, we used plant_region_area as a trait (**Dataset 1**), and the specific effect reflects the difference in that measurement between the soil conditions. The analysis was conducted independently for each day because we noted temporal variation in daily growth feedbacks, and the resulting tables were combined without re-adjusting the corrected p-values. SNPs were retained and visualized if they had a $-\log_{10}$(adjusted p-value) of at least 5 on at least three consecutive days and a minor allele count of at least 5. From these plots, we identified SNPs with a robust multi-day signal and analysed them further using gwaR (https://github.com/Gregor-Mendel-Institute/gwaR). This allowed us to investigate the genomic neighbourhood of those SNPs and pinpoint genes likely involved in the microbiome feedback phenotype. Detailed approach to select candidate genes is explained in **Suppl. Text**. The GWAS data are in **Dataset 3** and their analysis is documented in the **Suppl. File 3**. For computational analyses, we used the BioHPC-Genomics compute cluster at the Leibniz Rechenzentrum and the CLIP cluster at Vienna BioCenter (VBC).

GWAS candidate gene selection

To identify candidate genes associated with BX-growth feedbacks, we performed the genome wide association study (GWAS) with the 410 genome-sequenced accessions using their phenotypic diversity in growth feedbacks (**Fig. 1c**). To account for population structure within the *Arabidopsis* accessions, a linear mixed model was employed (*38*). We used the daily growth feedback data, which includes the temporal dynamics (**fig. S2a**), and calculated associations between single nucleotide polymorphisms (SNPs) and phenotypes for each of the 20 DAGs (**Dataset 3**). In essence, we performed the GWAS using each day as a covariable in the analysis.

To impose stringency in SNP identification, we performed FDR correction and Bonferroni correction of the *p*-values of SNPs that associated with the growth feedbacks (**Fig. 2a**, **Dataset 3**). Next, we filtered the SNPs for those with a minor allele frequency of higher than 2.5 % to also include rare SNPs associated with local and environmental adaptation (*51*), and those SNPs with significance (Bonferroni correction) for at least three consecutive days. To isolate candidate genes, we used the "get_nearest_genes" function from the gwaR package in R (https://github.com/Gregor-Mendel-Institute/gwaR) to identify genes located near each candidate SNP. Finally, to select candidate genes for functional validation, we screened the genes identified in the GWAS for overlap with differentially expressed genes in the roots and shoots of Arabidopsis plants grown on a BX_{minus} and BX_{plus} microbiome (*30*).

MBOA allelopathic growth assays

To test for the allelopathic effect of MBOA on plant growth, we grew different Arabidopsis accessions on ½ MS (Duchefa, Haarlem, the Netherlands) plates containing 1.5% phytoagar (Duchefa, Haarlem, the Netherlands) supplemented with 0.25 mM MBOA (Sigma Aldrich, St. Louis, USA) dissolved in

DMSO or on control plates supplemented with the same concentration (500 μL per L) of DMSO (*52*).

Seeds were stratified by incubating plates for 3 days at 4 °C. The plates were placed upright and grown

in a controlled growth chamber (Percival, CLF Plant Climatics, Wertingen, Germany) in short days

with 10 hours light (100 μmol m^{-2} s^{-1}) (21 °C) and 14 hours darkness (18 °C) at 60 % relative humidity.

After 10 days, the lateral root length was measured using the image software ImageJ (*53*).

Root microbiome profiling

DNA extraction: For root microbiome analysis, 10 randomly selected *Arabidopsis* root systems were

removed from the pots, and excess soil carefully removed by hand. The root system was rinsed with

sterile ddH$_2$O to remove additional soil. The entire root system was flash frozen in liquid nitrogen and

stored at -80 °C until further processing. The roots were lyophilized for 24 h and grinded in 1.5 mL

Eppendorf tubes using a Retsch mill (Retsch GmBH, Haan; Germany) and 2 mm glass beads at 30 Hz

for 1 min. Approximately 20 mg of dry root material was used as the input for DNA extractions. DNA

extractions were performed in 96-well plates using the QIAGEN® MagAttract® PowerSoil® DNA KF

Kit (Qiagen, Germany) and the KingFisher™ Flex™ Purification System (ThermoFisher Scientific,

Waltham, MA) following the manufacturers guidelines. For all samples, the DNA concentration was

determined using the AccuClear® Ultra High Sensitivity dsDNA Quantitation Kit (Biotum, Fremont,

United States). For library preparation, DNA concentration was adjusted to 2 ng/μL with a Myra Liquid

Handler (Bio Molecular Systems, Upper Coomera; Australia), and 10 ng used for library preparation.

Library preparation: Library preparation for bacteria, fungi and oomycetes was performed

using an adapted two-step barcoded host-associated microbiome hamPCR protocol (*54*). Primers used

are documented in **Table S3**. For bacteria, the first PCR was performed using a combination of the

CS1-*fs*-799-F and CS2-*fs*-1193R frameshift (fs) (pooled in 5 equimolar concentrations) in

combination with the host specific *GIGANTEA* CS1-At-GI-F and CS2-At-GI-R primers in a single well.

We followed earlier work PCR settings for annealing temperature (*55*) and cycling conditions (*54*).

Briefly, the PCR settings were 94 °C for 2 min, and 10 cycles of 94 °C for 30 s, 55 °C for 50 s, 72 °C

for 30 s and finally 72 °C for 5 min. For fungi, the first PCR was performed with the internal transcribed

spacer (ITS) region primers ITS1fF and ITS2R with standard temperature (*56*) and cycle (*54*) settings.

The PCR settings were 94 °C for 3 min, and 10 cycles of 94 °C for 45 s, 50 °C for 60 s, 72 °C for 90 s,

and finally 72 °C for 10 min. For oomycetes, we adapted the ITS1-O_F1_G-46636 and 5.8s-O_R1_G-

46637 primers (*54*) with the CS1 and CS2 adapter sequences and utilised the same PCR settings as for

the fungi. PCR amplicons were purified using the ChargeSwitch® PCR Clean-Up Kit (Invitrogen,

Waltham, MA, USA) on a KingFisher™ Flex™ Purification System (ThermoFisher Scientific,

Waltham, MA, USA). The second PCR added unique barcodes to 5 μL of purified DNA using the

Fluidigm Access Array barcoding primers (Standard BioTools, California, USA). The cycling

conditions were 94 °C for 2 min, 25 cycles of 94 °C for 30 s, 60 °C for 30 s, 72°C for 60 s, and finally

extension 72°C for 10 min. Samples were again purified as before and quantified and pooled by adding

80 ng of each sample using a Myra Liquid Handler to create sub-libraries for bacteria, fungi and oomycetes. The pooled bacterial sub-library was subsequently separated on an 1.5 % agarose gel, the bacterial and GIGANTEA fragments excised, and gel purified using the Wizard® SV Gel and PCR Clean-Up System (Promega, Madison, Wisconsin, USA). The final bacterial sub-library was created by scaling the purified bacterial and GIGANTEA amplicons in a 24:1 ratio (bacteria:GIGANTEA) to reduce the amount of plant reads during sequencing. Five random samples were duplicated during library preparation and left unscaled to bioinformatically rescale the scaled samples (see ref (*54*) for details). The final sequencing library was created by pooling the sub-libraries in a DNA ratio of 87.5% (963 ng) scaled bacteria, 5.5 % (60 ng) unscaled bacterial samples and 3.5 % (39 ng) each for fungal and oomycetes. The library was paired end sequenced on an Illumina MiSeq instrument (v3 chemistry, 300 bp paired end) at the NGS platform (https://www.ngs.unibe.ch) of the University of Bern.

Bioinformatics: MiSeq reads were quality checked with FastQC v0.11.8 (Babraham Institute, Cambridge, United Kingdom). Barcodes were removed previously by the NGS platform and written to the sequence headers. With this information the data was demultiplexed and primers were removed using cutadapt v3.4 (*57*). Following the methods from (*29*), quality filtering, read merging and amplicon sequence variant (ASV) infering was performed with the dada2 package v1.26.0 (*58*) in R v4.2.0 (R Core Team 2022). Taxanomic assignment were done using naïve Bayesian classifier with a DADA2 formatted version from the SILVA v. 132 database (*59*) for bacteria and the FASTA general release from UNITE v8.3 for fungi. To improve taxonomy assignment, the taxonomy was assigned again with the IDTAXA classifier using the SILVA r138 and UNITE 8.2 databases from DECIPHER v2.26.0 (*60*) and replaced wherever IDTAXA assigned more taxonomic levels than the naïve Bayesian classifier. Bacterial ASVs assigned as Eukarya, Archaea, Cyanobacteria or Mitochondria and fungal ASVs belonging to Protista, Plantae, Protozoa or Animalia were discarded. Calculations were performed at the sciCORE (http://scicore.unibas.ch/) scientific computing center of the University of Basel.

Statistical analysis: Statistical analyses were performed with the vegan package v.2.6.4 (*61*) in R v4.2.0. Following the strategies described before (*62*), we normalised the data for bacteria, fungi and oomycetes by rarefication. Rarefaction thresholds per sample were defined as 6.02×10^4 for bacteria, 800 for fungi and 5700 for oomyctes (*62*). Outlier detection was performed with CLOUD using Bray-Curtis (BC) distances from the rarefied data for each condition and each treatment individually (*63*). The nearest number of neighbours were set to 60% of the sample size and we chose an empirical outlier percentile of 0.1. As described before (*54*), microbe-to-host ratios were reconstructed and the microbial load of each sample was quantified. Using a count table scaled by the microbial load, we performed ordinations based on Bray-Curtis dissimilarity matrices using R package phyloseq (*64*) and visualised it in a Constrained analysis of principle coordinates (CAP). We performed Permutational Multivariate Analysis of Variance (PERMANOVA) with 999 permutations with the adonis function from the vegan package to quantify the effects of our experiment. The microbiome data is in **Dataset 5** and its analysis is documented in **Suppl. File 5**.

Transcriptome analysis

RNA extraction, library preparation and sequencing: Roots from 6-week-old Arabidopsis plants were carefully removed from the soil. Excess soil was removed by briefly rinsing the roots in sterile ddH$_2$O and dried using a paper towel. The roots were then flash frozen in liquid nitrogen and homogenized in a 1.5 mL Eppendorf tube using a plastic micro pestle (Carl Roth, Karlsruhe, Germany). Total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) with on-column DNase treatment with the RNase-Free DNase kit (Qiagen, Hilden, Germany) following the manufacturer's protocol. RNA integrity was assessed using a Fragment Analyzer (Agilent Technology, Santa Clara, California, USA). RNA concentration was determined fluorometrically using the Qubit RNA HS Assay Kit (Thermo Fisher, Waltham, Massachusetts, USA). Libraries were prepared using the TruSeq Stranded mRNA kit (Illumina, SanDiego, California, USA) with 200 ng total RNA as input. Library preparation and sequencing was performed at the Genomics Facility (https://bsse.ethz.ch/genomicsbasel) of the University of Basel. Sequencing (51 bp paired end) was performed on an Illumina Novaseq® 6000 System (Illumina, SanDiego, California, USA) with an SP slow cell.

Bioinformatics and statistics: Reads were aligned to the *Arabidopsis thaliana* TAIR10 reference genome and read counts determined using STAR 2.7.9a (*65*). The raw data processing was performed at the scientific computing facility of the University of Basel (http://scicore.unibas.ch/). Sample outliers were identified using a combination of generalized principal component analysis and by analysing Poisson distances across samples. Sample clustering was performed using variance stabilizing transformations on the normalized gene counts. Differential gene expression analysis was performed in R using DESeq2 (*66*), a method for differential analysis of count data using a **negative binomial** generalized linear model **with shrinkage estimation of dispersions** and fold changes. Co-expression analysis was performed with the R package coseq v1.18.0 (*67*). Clustering of the co-expressed genes was performed by running a Gaussian mixture model and arcsine transformation of normalized expression profiles 20 times to identify the optimal number of clusters (*67*). The transcriptomic data is in **Dataset 6** and the documented analysis in **Suppl. File 6**.

Validation of single T-DNA insertion in mmf1-2

We validated that *mmf1-2* carries a single T-DNA insertion by long-read whole genome sequencing and reference-guided assembly. High molecular weight DNA was extracted from leaves using the Machery-Nagel NucleoBond HMW DNA kit according to manufacturer's instructions. Long-read sequencing was carried out on an Oxford Nanopore Technologies promethion 24 instrument in a multiplexed run on a R10.4 flow cell. After demultiplexing with dorado (https://github.com/nanoporetech/dorado) adapters were trimmed using porechop, and the genome was assembled using flye (*68*), polished with medaka (https://github.com/nanoporetech/medaka) and scaffolded against Col-CEN (*69*) using ragtag (*70*) and annotations (araport11) from Col-CEN were lifted over using liftoff (*71*). The T-DNA insertion

in AT3G44630 was identified by aligning to the reference genome. The T-DNA sequence was used as

a query for BLAST against the *mmf1-2* assembly, returning only the insertion in AT3G44630 (**fig. S4d**).

Statistical analysis

We used R (R core Team, 2016) for statistical analysis and visualization of data. The R version and

package version used for each step of the analysis is documented at the end of the markdown files. All

the code required to reproduce the statistical analysis and graphing can be found on the GitHub

repository of this article. Figures were exported from R and combined and formatted using Adobe

Illustrator. The appropriate statistical test used for each analysis is indicated in the corresponding

figure legends and is annotated in the available code.

Data availability

We have deposited all source data together with all analysis scripts underlying all figures on GitHub

(https://github.com/PMI-Basel/JansevanRensburg_et_al_GWAS_MMF1). The GitHub repository's

structure follows the structure of this manuscript with separate folders of source data and analysis scripts

on i) Accession growth feedback (Dataset 1), ii) MBOA sensitivity (Dataset 2), iii) GWAS statistics

(Dataset 3), iv) Col-0 and mutant growth feedback (Dataset 4), v) microbiome (Dataset 5) and vi)

transcriptome (Dataset 6). Each dataset and its analysis documentation (R-markdown outputs) is shared

with this article as **Dataset 1** to **6** and the corresponding **Suppl. Files 1** to **6**. The raw sequencing data

are stored at the European Nucleotide Archive (http://www.ebi.ac.uk/ena). The microbiome profiling is

stored under project number PRJEB85607, the transcriptome data is available under PRJEB85599 and

the genome of the *mmf1-2* mutant under PRJEB85789. The bioinformatic code of all analyses, including

information on barcodes, primers and sample assignments is provided on GitHub. Raw images are

available upon request.

Code availability

All the code required to reproduce the statistical analysis and graphing can be found on GitHub https://github.com/PMI-Basel/JansevanRensburg_et_al_GWAS_MMF1.

Acknowledgements

We specifically thank Nicolas Widmer from Agroscope Changins (Switzerland) for field access and assistance. We thank Prof. Jane Parker for providing Arabidopsis mutant seeds. We thank the team at the Plant Sciences Facility at Vienna BioCenter Core Facilities GmbH (VBCF), a member of the Vienna BioCenter (VBC), Austria, for their help with Arabidopsis image acquisition. We are very grateful to Dr. Pamela Nicholson and her team at the Next Generation Sequencing Platform in Bern for excellent technical support. We thank Dr. Phillipe Demougin and Dr. Christian Beisle (both University of Basel, Switzerland) for their support with RNAseq. Finally, we thank Duncan Crosbie (LMU Munich) and Helmut Blum and Stefan Krebs (Gene Center, LMU Munich) for support with genome sequencing of *mmf1-2*.

Funding

This project was funded by the Swiss National Foundation (Grant No. 189249 to K.S.), the Austrian Academy of Sciences (ÖAW, to C.B.), by the European Union's Horizon 2020 research and innovation programme by the European Research Council (ERC) (Grant Agreement No. 716823 'FEAR-SAP' to C.B.). N.S. received funding from the Deutsche Forschungsgemeinschaft (DFG) in the frame of TRR356 project A05 (project No. 491090170). For parts of the computational analyses, we used the BioHPC-Genomics compute cluster at the Leibniz Rechenzentrum Munich (DFG; grant no. 450674345) and the CLIP batch environment at the Vienna Biocenter.

Ethics declarations

None

References

1. P. Trivedi, J. E. Leach, S. G. Tringe, T. Sa, B. K. Singh, Plant–microbiome interactions: from community assembly to plant health. *Nature Reviews Microbiology* **18**, 607–621 (2020).

2. R. L. Berendsen, C. M. J. Pieterse, P. A. H. M. Bakker, The rhizosphere microbiome and plant health. *Trends in Plant Science* **17**, 478–486 (2012).

3. R. Mendes, P. Garbeva, J. M. Raaijmakers, The rhizosphere microbiome: Significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiology Reviews* **37**, 634–663 (2013).

4. W.-J. Li, X.-Y. Zhou, X.-L. An, L.-J. Li, C.-S. Lin, H. Li, H.-Z. Li, Enhancement of beneficial microbiomes in plant–soil continuums through organic fertilization: Insights into the composition and multifunctionality. *Soil Ecol. Lett.* **6**, 230223 (2024).

5. H. Janse van Rensburg, K. Stengele, K. Schlaeppi, Understanding plant responsiveness to microbiome feedbacks. *Current Opinion in Plant Biology* **81**, 102603 (2024).

6. T. M. Bezemer, C. S. Lawson, K. Hedlund, A. R. Edwards, A. J. Brook, J. M. Igual, S. R. Mortimer, W. H. Van Der Putten, Plant species and functional group effects on abiotic and microbial soil properties and plant-soil feedback responses in two grasslands. *Journal of Ecology* **94**, 893–904 (2006).

7. W. H. Van der Putten, R. D. Bardgett, J. D. Bever, T. M. Bezemer, B. B. Casper, T. Fukami, P. Kardol, J. N. Klironomos, A. Kulmatiski, J. A. Schweitzer, K. N. Suding, T. F. J. Van de Voorde, D. A. Wardle, Plant-soil feedbacks: The past, the present and future challenges. *Journal of Ecology* **101**, 265–276 (2013).

8. M. A. Meier, G. Xu, M. G. Lopez-Guerrero, G. Li, C. Smith, B. Sigmon, J. R. Herr, J. R. Alfano, Y. Ge, J. C. Schnable, J. Yang, Association analyses of host genetics, root-colonizing microbes, and plant phenotypes under different nitrogen conditions in maize. *eLife* **11** (2022).

9. Y. Song, E. Atza, J. J. S. Gil, D. Akkermans, R. de Jonge, P. G. H. de Rooij, D. Kakembo, P. A. H. M. Bakker, C. M. J. Pieterse, N. V. Budko, R. L. Berendsen, Seed tuber microbiome is a predictor of next-season potato vigor. *bioRxiv*, 2024.04.18.590071 (2024).

10. M. Wang, A.-H. Ge, X. Ma, X. Wang, Q. Xie, L. Wang, X. Song, M. Jiang, W. Yang, J. D. Murray, Y. Wang, H. Liu, X. Cao, E. Wang, Dynamic root microbiome sustains soybean productivity under unbalanced fertilization. *Nat Commun* **15**, 1668 (2024).

11. R. C. Wilhelm, H. M. van Es, D. H. Buckley, Predicting measures of soil health using the microbiome and supervised machine learning. *Soil Biology and Biochemistry* **164**, 108472 (2022).

12. X. Qiao, S. Bei, G. Wang, C. Li, H. Li, J. Zhang, F. Zhang, Soil biota is decisive for overyielding in intercropping under low phosphorus conditions. *Journal of Applied Ecology* **59**, 1804–1814 (2022).

13.   M. Chialva, A. Salvioli di Fossalunga, S. Daghino, S. Ghignone, P. Bagnaresi, M.
Chiapello, M. Novero, D. Spadaro, S. Perotto, P. Bonfante, Native soils with their
microbiotas elicit a state of alert in tomato plants. *New Phytologist* **220** (2018).

14.   L. Hu, C. A. M. Robert, S. Cadot, X. Zhang, M. Ye, B. Li, D. Manzo, N. Chervet, T.
Steinger, M. G. A. Van Der Heijden, K. Schlaeppi, M. Erb, Root exudate metabolites
drive plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota.
Nature Communications **9**, 1–13 (2018).

15.   P. J. P. Teixeira, N. R. Colaianni, C. R. Fitzpatrick, J. L. Dangl, Beyond pathogens:
microbiota interactions with the plant immune system. *Current Opinion in Microbiology*
49, 7–17 (2019).

16.   K. Yu, C. M. J. Pieterse, P. A. H. M. Bakker, R. L. Berendsen, Beneficial microbes
going underground of root immunity. *Plant, Cell & Environment* **42**, 2860 (2019).

17.   J. Bergelson, J. Mittelstrass, M. W. Horton, Characterizing both bacteria and fungi
improves understanding of the Arabidopsis root microbiome. *Scientific Reports 2019*
9:1 **9**, 1–11 (2019).

18.   M. Hu, J.-M. Zhou, Research on ADR1s helps understanding the plant immune
network. *Stress Biology* **2**, 12 (2022).

19.   J. Bigeard, J. Colcombet, H. Hirt, Signaling mechanisms in pattern-triggered immunity
(PTI). *Molecular Plant* **8**, 521–539 (2015).

20.   F. Locci, J. E. Parker, Plant NLR immunity activation and execution: a biochemical
perspective. *Open Biology* **14** (2024).

21.   F. Monteiro, M. T. Nishimura, Structural, functional, and genomic diversity of plant
NLR proteins: An evolved resource for rational engineering of plant immunity. *Annual
Review of Phytopathology* **56**, 243–267 (2018).

22.   G. Castrillo, P. J. P. L. Teixeira, S. H. Paredes, T. F. Law, L. De Lorenzo, M. E.
Feltcher, O. M. Finkel, N. W. Breakfield, P. Mieczkowski, C. D. Jones, J. Paz-Ares, J.
L. Dangl, Root microbiota drive direct integration of phosphate stress and immunity.
Nature **543**, 513–518 (2017).

23.   S. L. Lebeis, S. H. Paredes, D. S. Lundberg, N. Breakfield, J. Gehring, M. McDonald, S.
Malfatti, T. G. Del Rio, C. D. Jones, S. G. Tringe, J. L. Dangl, Salicylic acid modulates
colonization of the root microbiome by specific bacterial taxa. *Science* **349**, 860–864
(2015).

24.   S. Pfeilmeier, G. C. Petti, M. Bortfeld-Miller, B. Daniel, C. M. Field, S. Sunagawa, J. A.
Vorholt, The plant NADPH oxidase RBOHD is required for microbiota homeostasis in
leaves. *Nature Microbiology* **6**, 852–864 (2021).

25.   M. Frey, K. Schullehner, R. Dick, A. Fiesselmann, A. Gierl, Benzoxazinoid
biosynthesis, a model for evolution of secondary metabolic pathways in plants.
Phytochemistry **70**, 1645–1651 (2009).

26. S. Cadot, H. Guan, M. Bigalke, J. C. Walser, G. Jander, M. Erb, M. G. A. van der
Heijden, K. Schlaeppi, Specific and conserved patterns of microbiota-structuring by
maize benzoxazinoids in the field. *Microbiome* **9**, 1–19 (2021).

27. T. E. A. Cotton, P. Pétriacq, D. D. Cameron, M. A. Meselmani, R. Schwarzenbacher, S.
A. Rolfe, J. Ton, Metabolic regulation of the maize rhizobiome by benzoxazinoids.
ISME Journal **13**, 1647–1658 (2019).

28. S. Cadot, V. Gfeller, L. Hu, N. Singh, A. Sánchez-Vallet, G. Glauser, D. Croll, M. Erb,
M. G. A. van der Heijden, K. Schlaeppi, Soil composition and plant genotype determine
benzoxazinoid-mediated plant–soil feedbacks in cereals. *Plant Cell and Environment*
44, 3502–3514 (2021).

29. V. Gfeller, J. Waelchli, S. Pfister, G. Deslandes-Hérold, F. Mascher, G. Glauser, Y.
Aeby, A. Mestrot, C. A. M. Robert, K. Schlaeppi, M. Erb, Plant secondary metabolite-
dependent plant-soil feedbacks can improve crop yield in the field. *eLife* **12** (2023).

30. K. Stengele, L. Stauber, L. Thoenen, H. Janse van Rensburg, V. D'Adda, K. Schlaeppi,
Benzoxazinoid-mediated microbiome feedbacks enhance Arabidopsis growth and
defense. bioRxiv [Preprint] (2024). https://doi.org/10.1101/2024.10.21.619081.

31. A. Martinez-Medina, V. Flors, M. Heil, B. Mauch-Mani, C. M. J. Pieterse, M. J. Pozo, J.
Ton, N. M. van Dam, U. Conrath, Recognizing Plant Defense Priming. *Trends in Plant*
Science **21**, 818–822 (2016).

32. P. C. A. Wintermans, P. A. H. M. Bakker, C. M. J. Pieterse, Natural genetic variation in
Arabidopsis for responsiveness to plant growth-promoting rhizobacteria. *Plant*
Molecular Biology **90**, 623–634 (2016).

33. The 1001 Genomes Consortium, 1,135 Genomes Reveal the Global Pattern of
Polymorphism in Arabidopsis thaliana. *Cell* **166**, 481–491 (2016).

34. P. Hüther, N. Schandry, K. Jandrasits, I. Bezrukov, C. Becker, ARADEEPOPSIS, an
automated workflow for top-view plant phenomics using semantic segmentation of leaf
States. *Plant Cell* **32**, 3674–3688 (2020).

35. R. G. Belz, K. Hurle, Differential exudation of two benzoxazinoids - One of the
determining factors for seedling allelopathy of Triticeae species. *Journal of Agricultural*
and Food Chemistry **53**, 250–261 (2005).

36. N. Schandry, C. Becker, Allelopathic Plants: Models for Studying Plant–Interkingdom
Interactions. *Trends in Plant Science* **25**, 176–185 (2020).

37. V. Gfeller, S. Cadot, J. Waelchli, S. Gulliver, C. Terrettaz, L. Thönen, P. Mateo, C. A.
M. Robert, F. Mascher, T. Steinger, M. Bigalke, M. Erb, K. Schlaeppi, Soil chemical
and microbial gradients determine accumulation of root-exuded secondary metabolites
and plant–soil feedbacks in the field. *Journal of Sustainable Agriculture and*
Environment **2**, 173–188 (2023).

38. C. Lippert, F. P. Casale, B. Rakitsch, O. Stegle, LIMIX: genetic analysis of multiple
traits. doi: 10.1101/003905 (2014).

39. J. A. Dongus, J. E. Parker, EDS1 signalling: At the nexus of intracellular and surface
772   receptor immunity. *Current Opinion in Plant Biology* **62**, 102039 (2021).

773   40. J. Ordon, J. Gantner, J. Kemna, L. Schwalgun, M. Reschke, J. Streubel, J. Boch, J.
774   Stuttmann, Generation of chromosomal deletions in dicotyledonous plants employing a
775   user-friendly genome editing toolkit. *Plant Journal* **89**, 155–168 (2017).

776   41. R. N. Pruitt, F. Locci, F. Wanke, L. Zhang, S. C. Saile, A. Joe, D. Karelina, C. Hua, K.
777   Fröhlich, W. L. Wan, M. Hu, S. Rao, S. C. Stolze, A. Harzen, A. A. Gust, K. Harter, M.
778   H. A. J. Joosten, B. P. H. J. Thomma, J. M. Zhou, J. L. Dangl, D. Weigel, H. Nakagami,
779   C. Oecking, F. E. Kasmi, J. E. Parker, T. Nürnberger, The EDS1–PAD4–ADR1 node
780   mediates Arabidopsis pattern-triggered immunity. *Nature* **598**, 495–499 (2021).

781   42. J. Lang, B. Genot, J. Bigeard, J. Colcombet, MPK3 and MPK6 control salicylic acid
782   signaling by up-regulating *NLR* receptors during pattern- and effector-triggered
783   immunity. *Journal of Experimental Botany* **73**, 2190–2205 (2022).

784   43. T. Chen, K. Nomura, X. Wang, R. Sohrabi, J. Xu, L. Yao, B. C. Paasch, L. Ma, J.
785   Kremer, Y. Cheng, L. Zhang, N. Wang, E. Wang, X. F. Xin, S. Y. He, A plant genetic
786   network for preventing dysbiosis in the phyllosphere. *Nature* **580**, 653–657 (2020).

787   44. P. Runge, "Structural variation of a polymorphic RPP1-like cluster in local Arabidopsis
788   thaliana populations alters phyllosphere microbiome in Gorzów Wielkopolski," thesis,
789   University of Tübingen, Germany (2022).

790   45. S. Goritschnig, A. D. Steinbrenner, D. J. Grunwald, B. J. Staskawicz, Structurally
791   distinct Arabidopsis thaliana NLR immune receptors recognize tandem WY domains of
792   an oomycete effector. *New Phytologist* **210**, 984–996 (2016).

793   46. D. Maag, A. Köhler, C. A. M. Robert, M. Frey, J. L. Wolfender, T. C. J. Turlings, G.
794   Glauser, M. Erb, Highly localized and persistent induction of Bx1-dependent herbivore
795   resistance factors in maize. *The Plant Journal* **88**, 976–991 (2016).

796   47. D. Jirage, T. L. Tootle, T. L. Reuber, L. N. Frost, B. J. Feys, J. E. Parker, F. M. Ausubel,
797   J. Glazebrook, Arabidopsis thaliana PAD4 encodes a lipase-like gene that is important
798   for salicylic acid signaling. *Proceedings of the National Academy of Sciences* **96**,
799   13583–13588 (1999).

800   48. X. Sun, D. Lapin, J. M. Feehan, S. C. Stolze, K. Kramer, J. A. Dongus, J.
801   Rzemieniewski, S. Blanvillain-Baufumé, A. Harzen, J. Bautor, P. Derbyshire, F. L. H.
802   Menke, I. Finkemeier, H. Nakagami, J. D. G. Jones, J. E. Parker, Pathogen effector
803   recognition-dependent association of NRG1 with EDS1 and SAG101 in TNL receptor
804   immunity. *Nat Commun* **12**, 3335 (2021).

805   49. C. Cardoso, T. Charnikhova, M. Jamil, P. M. Delaux, F. Verstappen, M. Amini, D.
806   Lauressergues, C. Ruyter-Spira, H. Bouwmeester, Differential activity of Striga
807   hermonthica seed germination stimulants and Gigaspora rosea hyphal branching factors
808   in rice and their contribution to underground communication. *PLoS ONE* **9** (2014).

809   50. M. Togninalli, Ü. Seren, D. Meng, J. Fitz, M. Nordborg, D. Weigel, K. Borgwardt, A.
810   Korte, D. G. Grimm, The AraGWAS Catalog: A curated and standardized Arabidopsis
811   thaliana GWAS catalog. *Nucleic Acids Research* **46**, D1150–D1156 (2018).

51. Á. Ferrero-Serrano, S. M. Assmann, Phenotypic and genome-wide association with the
local environment of Arabidopsis. *Nature Ecology and Evolution* **3**, 274–285 (2019).

52. E. Knoch, J. Kovács, S. Deiber, R. Shanmuganathan, N. Serra Serra, C. Becker, N.
Schandry, N. Serra, S. C. Becker, Transcriptional response of a target plant to
benzoxazinoid and diterpene allelochemicals highlights commonalities in detoxification.
bioRxiv, 2022.02.21.480921 (2022).

53. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of
Image Analysis. *Nat Methods* **9**, 671–675 (2012).

54. D. S. Lundberg, P. P. N. Ayutthaya, A. Strauß, G. Shirsekar, W. S. Lo, T. Lahaye, D.
Weigel, Host-associated microbe pcr (Hampcr) enables convenient measurement of both
microbial load and community composition. *eLife* **10** (2021).

55. Y. Bai, D. B. Müller, G. Srinivas, R. Garrido-Oter, E. Potthoff, M. Rott, N.
Dombrowski, P. C. Münch, S. Spaepen, M. Remus-Emsermann, B. Hüttel, A. C.
McHardy, J. A. Vorholt, P. Schulze-Lefert, Functional overlap of the Arabidopsis leaf
and root microbiota. *Nature 2015 528:7582* **528**, 364–369 (2015).

56. K. L. McGuire, S. G. Payne, M. I. Palmer, C. M. Gillikin, D. Keefe, S. J. Kim, S. M.
Gedallovich, J. Discenza, R. Rangamannar, J. A. Koshner, A. L. Massmann, G. Orazi,
A. Essene, J. W. Leff, N. Fierer, Digging the New York City Skyline: Soil Fungal
Communities in Green Roofs and City Parks. *PLOS ONE* **8**, e58020 (2013).

57. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**, 10 (2011).

58. B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, S. P.
Holmes, DADA2: High-resolution sample inference from Illumina amplicon data.
Nature Methods **13**, 581–583 (2016).

59. C. Quast, E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, F. O.
Glöckner, The SILVA ribosomal RNA gene database project: Improved data processing
and web-based tools. *Nucleic Acids Research* **41**, D590 (2013).

60. E. S. Wright, Using DECIPHER v2.0 to analyze big biological sequence data in R. *R
Journal* **8**, 352–359 (2016).

61. J. Oksanen, G. L. Simpson, F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B.
O'Hara, P. Solymos, M. H. H. Stevens, E. Szoecs, H. Wagner, M. Barbour, M.
Bedward, B. Bolker, D. Borcard, G. Carvalho, M. Chirico, M. D. Caceres, S. Durand,
H. B. A. Evangelista, R. FitzJohn, M. Friendly, B. Furneaux, G. Hannigan, M. O. Hill,
L. Lahti, D. McGlinn, M.-H. Ouellette, E. R. Cunha, T. Smith, A. Stier, C. J. F. T.
Braak, J. Weedon, vegan: Community Ecology Package, version 2.6-6.1 (2024);
https://cran.r-project.org/web/packages/vegan/index.html.

62. S. Weiss, Z. Z. Xu, S. Peddada, A. Amir, K. Bittinger, A. Gonzalez, C. Lozupone, J. R.
Zaneveld, Y. Vázquez-Baeza, A. Birmingham, E. R. Hyde, R. Knight, Normalization
and microbial differential abundance strategies depend upon data characteristics.
Microbiome **5**, 1–18 (2017).

63. E. Montassier, G. A. Al-Ghalith, B. Hillmann, K. Viskocil, A. J. Kabage, C. E.
McKinlay, M. J. Sadowsky, A. Khoruts, D. Knights, CLOUD: A non-parametric
detection test for microbiome outliers. *Microbiome* **6**, 1–10 (2018).

64. P. J. McMurdie, S. Holmes, phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

65. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M.
Chaisson, T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29, 15–21 (2013).

66. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

67. A. Rau, C. Maugis-Rabusseau, Transformation and model choice for RNA-seq co-
expression analysis. *Briefings in Bioinformatics* **19**, 425–436 (2018).

68. M. Kolmogorov, J. Yuan, Y. Lin, P. A. Pevzner, Assembly of long, error-prone reads
using repeat graphs. *Nat Biotechnol* **37**, 540–546 (2019).

69. M. Naish, M. Alonge, P. Wlodzimierz, A. J. Tock, B. W. Abramson, A. Schmücker, T.
Mandáková, B. Jamge, C. Lambing, P. Kuo, N. Yelina, N. Hartwick, K. Colt, L. M.
Smith, J. Ton, T. Kakutani, R. A. Martienssen, K. Schneeberger, M. A. Lysak, F.
Berger, A. Bousios, T. P. Michael, M. C. Schatz, I. R. Henderson, The genetic and
epigenetic landscape of the Arabidopsis centromeres. *Science* **374**, eabi7489 (2021).

70. M. Alonge, L. Lebeigle, M. Kirsche, K. Jenike, S. Ou, S. Aganezov, X. Wang, Z. B.
Lippman, M. C. Schatz, S. Soyk, Automated assembly scaffolding using RagTag
elevates a new tomato system for high-throughput genome editing. *Genome Biology* **23**,
258 (2022).

71. A. Shumate, S. L. Salzberg, Liftoff: accurate mapping of gene annotations.
Bioinformatics **37**, 1639–1643 (2021).

Supplementary Materials

Title:

A TNL receptor mediates microbiome feedbacks in Arabidopsis

Authors:

Henry Janse van Rensburg[1*], Niklas Schandry[2,3*§], Jan Waelchli[1], Katja Stengele[1], Selma Cadot[1], Katharina Jandrasits[2], Claude Becker[2,3,§], Klaus Schlaeppi[1,§]

[1] Department of Environmental Sciences, University of Basel, Basel, Switzerland

[2] Gregor Mendel Institute of Molecular Plant Biology GmbH, Austrian Academy of Sciences, Vienna BioCenter (VBC), Vienna, Austria

[3] LMU Biocenter, Faculty of Biology, Ludwig-Maximilians-Universität Munich, Martinsried, Germany

The file includes:

Figures S1 to S9

Tables S1 to S3

Other Supplementary Materials for this manuscript include

Dataset S1 to S6 (Input data to reproduce figures)

Supplementary Files 1 to 6 (R markdown files to reproduce analysis)

Supplementary Figures

Figure S1. **Overview of the experimental workflow**. To condition the soil microbiomes, maize exuding BXs (B73) and a BX-deficient mutant (*bx1*) was grown for 12 weeks on field soil, resulting in a BX_{plus} and a BX_{minus} microbiome. The soil core surrounding the roots was removed and sieved to produce the soil used for growth-feedback assays. 410 Arabidopsis accessions were grown on the two conditioned soil microbiomes and phenotyped using a high-throughput phenotyping pipeline. Growth feedback response as the rosette size (BX_{plus} – BX_{minus}) on the two soil microbiomes was used as the phenotype for the genome-wide association study (GWAS). *Mediator of microbiome feedbacks 1* (*MMF1*) identified in the GWAS was functionally validated for growth feedback assays using T-DNA insertion mutant lines. The cause for lack of feedback in mutants were further dissected by analysing the microbiome and transcriptome of wild-type and mutant plants grown on the two microbiomes. Created with BioRender®.

Figure S2. **Microbiome feedback of Arabidopsis accessions**. (**a**) Relationship between growth feedbacks and geographical location of the phenotyped accessions. Scatter plot illustrating the relationship between latitude or longitude and the microbiome feedback of accessions at 15 days after germination. Each data point is coloured according to its microbiome feedback response, with orange and blue denoting a negative or positive microbiome feedback, respectively. The plot includes a linear regression line (in red), highlighting the relationship between location and phenotype. The regression statistics, including the R-squared value (R^2) and the p-value of the F-test, are displayed in the top right corner of the plot. The R-squared value quantifies the proportion of variance in the phenotype explained by either latitude or longitude, while the p-value refers to testing the null hypothesis that the slope of the regression line is zero. (**b**) The daily microbiome feedback of Col-0 and the top 10 accessions with the strongest positive (blue) or negative (orange) BX-growth feedback when grown on the BX_{plus} and BX_{minus} soils. The heatmap colours are the pseudo-log transformed microbiome feedbacks (rosette area on BX_{plus} − rosette area on BX_{minus}). Accessions are clustered on the y-axis based on agglomerative hierarchical clustering using Ward's method performed using Dynamic Time Warping (DTW) distances and Ward's method to identify groups with similar response patterns over time. The dendrogram on the right side of the heat map illustrates the clustering structure, grouping accessions that exhibit similar trajectories in their growth feedback **Dataset 1**.

934
Figure S3. Direct allelopathic effect of benzoxazinoids on the growth of Arabidopsis. (**a**) Sensitivity to
MBOA in terms of root length response (root length on control – root length on MBOA) of Col-0 and the
accessions with the strongest positive or negative microbiome feedback. Accessions are colour-coded as to
whether they showed a positive (blue) or negative (orange) microbiome feedback on the BX-conditioned soil
microbiome in the GWAS experiment. Boxplot insert displays the root length of Col-0 on the control or MBOA
treatment as an example. Plants were grown for 10 days on ½ MS media in the presence of 0.25 mM MBOA or
control (DMSO). (**b**) Pearson's correlation analysis between MBOA sensitivity and microbiome feedback at 15
days after germination across the different accessions. The scatter plot displays individual data points coloured by
microbiome feedback, with positive responses shown in blue and negative responses in orange. A linear regression
line (in red) illustrates the trend between the two variables. The correlation coefficient (R^2) and *p*-value are
annotated on the plot, indicating the strength and statistical probability of the relationship. **Dataset 2.**

955
Figure S4. **Genetic characterization of *mmf1* T-DNA insertion lines.** (**a**) Genomic location of the T-DNA
insertion sites for *mmf1-1* and *mmf1-2* in the open reading frame of *MMF1*. Exons and introns are shown in colour,
and the T-DNA insertion position as triangles. (**b**) Genotyping results of *mmf1-1* (SALK_144159C) and *mmf1-2*
(SALK_002318C) by PCR to confirm the presence of the T-DNA insertion. The gels represent genotyping of 6
mutant plants for the T-DNA insertion allele (left, LBb1.3 and RP primers) and the wild-type allele (right, LP and
RP primers). (**c**) Relative expression levels of *MMF1* determined by qRT-PCR in wild-type (Col-0) and the two
MMF1 mutant lines. (**d**) Mapping of genomic long-reads from *mmf1-2* onto Col-CEN v1.2. Top panel shows the
genomic coordinates on chromosome 3, middle panel shows the annotation of the coding sequence of *mmf-*
1, bottom panel shows the aggregated pile-up of mapped reads, confirming the T-DNA insertion in the third exon.
965

Figure S5. **Growth feedback in a BX_{plus} microbiome remains intact in effector-triggered immunity**
signalling mutants. (**a** and **b**) Repeated growth feedbacks (Experiment II and IV) of Col-0 wild-type (WT) and
two *mmf1* T-DNA insertion mutants (*At3g44630*) grown on the BX_{minus} and BX_{plus} soil microbiomes. Feedback
experiments were performed on greenhouse-conditioned soil batches (**a**) BS04 and (**b**) BS09. Graphs illustrate
the rosette area (in pixels) of 6-week-old plants. Sample sizes are indicated at the bottom of each plot. Output of
the ANOVA and FDR corrected *p*-values are indicated in each plot. (**c**) Phenotypes of 6-week-old WT, and
mutants (*eds1-12*, *pad4-1* and *sag101-3*) affected in effector-triggered immunity signalling pathways grown on
the BX_{minus} and BX_{plus} soils. (**b**) Fresh weight, and dry weight of WT, *eds1-12*, *pad4-1* and *sag101-3* illustrate the
positive growth feedback on the BX_{plus} soil microbiome. Boxplots illustrate the median, standard error, and
interquartile range, and the whiskers extend from the box to the smallest and largest values. Sample numbers are
indicated at the bottom of each plot. ANOVA output and FDR-corrected *p*-values comparing the growth feedback
of each line are shown above the graph. **Dataset 4.**

979
Figure S6. **Root microbiota profiling of WT and *mmf1-2* mutants grown on the BX$_{minus}$ or BX$_{minus}$ soils**. (**a**)
Bacterial, fungal and oomycete richness in the roots of wild-type (WT) and *mmf1-2* mutants grown on BX$_{plus}$ and
BX$_{minus}$ soil displayed as Shannon indices. Individual samples are shown as dots. Comparisons between the two
lines are shown as *p*-values based on Kruskal-Wallis Test. (**b**) The output of PERMANOVA on Bray-Curtis
dissimilarities of the bacterial, fungal and oomycete root microbiota of Arabidopsis WT (filled circles) and *mmf1-*
2 mutants (open triangles) grown on the BX$_{minus}$ (green) and BX$_{plus}$ (yellow) soils. The R^2 and *p* values indicate
the line and conditioning effect and their interaction on the two soils. Constrained Analysis of Principal
Coordinates (CAP) visually illustrating the effects found in the PERMANOVA analysis. Plants grown on a BX$_{plus}$
(yellow) and BX$_{minus}$ (green) soil microbiome are grouped based on a 95% confidence interval. The axes show the
percentage of explained variance for the first two principal components.
990

Figure S7. **Root transcriptome of WT and *mmf1-2* mutants grown on the BX$_{minus}$ and BX$_{plus}$ soils.** MA plot of differentially expressed genes between *mmf1-2* mutants and WT plants in response to the (**a**) BX$_{minus}$ and (**b**) BX$_{plus}$ soil microbiome. Significantly differentially expressed genes were calculated using DESeq2 (log$_2$fold change $\geq$ 1; empirical Bayes statistic false discovery rate < 0.05).

997
998
Figure S8. **Heatmap illustrating the co-expression analysis of the Arabidopsis WT root transcriptome**
grown on the BX$_{minus}$ and BX$_{plus}$ soil microbiome. Heatmap showing the co-expressed gene clusters (Benjamini-
Hochberg-correction, p-adjust < 0.01) as identified by coseq for plants grown on BX$_{minus}$ (green) or BX$_{plus}$ (yellow)
soils. Colours of the heatmap indicates the expression profiles in terms of normalized gene counts for each gene.
The log2Fold change and adjusted *p* values (empirical Bayes shrinking) for each gene inferred from DESeq2 is
shown on the right. Cluster labels show the top 5 enriched GO terms for biological function within each cluster.
GO text colour is for better readability only. Full tables of all enriched GO terms in each cluster are provided in
Dataset 6.
1007
1008

1009

Figure S9. **Proposed model of microbiome feedbacks through MMF1 in Col-0 plants**. Plants with a functional MMF1 receptor perceive a putative effector molecule that is present in the BX_{plus} soil microbiome, mounting a defence response that coincides with an altered root microbiome and promoted plant growth. The absence of this effector molecule in the BX_{minus} microbiome results in the lack of defence responses in plants grown on this soil microbiome. Col-0 plants lacking the MMF1 receptor fail to perceive the effector molecule from the BX_{plus} microbiome and do not mount defence responses; they have a different microbiome, and no enhanced shoot growth is observed. Figure created with BioRender®.

1017
1018

Supplementary Tables

Table S1: Setup and growth conditions of the different experiments performed in this study.
For each of the microbiome feedback experiments in the manuscript, the soil batch, measurement in terms of data generated, and time from start to conclusion is indicated.

Experiment	Plants	Soil batch	Measurement	Growth (weeks)
Exp. I	Accessions run 1	1	Growth (210 accessions)	12
Exp. II	Accessions run 2	2	Growth (210 accessions)	9
Exp. III	*mmf1* rep. 1	3	Growth	6
Exp. IV	*mmf1* rep. 2	3	Growth, microbiota, transcriptome	6
Exp. V	*mmf1* rep. 3	4	Growth	6
Exp. VI	ETI mutants	4	Growth	6

Table S2: **Set up of the different microbiome conditioning experiments**.
Each "soil batch" denotes an independent conditioning experiment from soils collected from three adjacent Agroscope research fields in Changins, Nyon, Switzerland (46°24'00.0"N, 6°14'22.8"E). The "scollection" column specifies when and from which field (Parcel) the soil was collected. For fertilization in the greenhouse, pots were treated weekly with the indicated solution. Abbreviations: "rel. hum." = relative humidity; "int." = intensity.

Soil batch	Collection	Conditioning	Fertilization	Growth conditions
Batch 1	April 2018 (Parcel 30/31)	Field[1]	NA[3]	Conventional farming
Batch 2	Dec. 2019 (Parcel 30/31)	Field[1]	NA[3]	Conventional farming
Batch 3	Aug. 2020 (Parcel 30/31)	Greenhouse	4 w. low iron,[4] 8 w. high iron[5]	16 h day at min. 23°C, 8 h night at min. 19°C, >500 μmol m^{-2} s^{-1} light int.
Batch 4	Sept. 2022 (Parcel 29/30)	Phytotron[2]	12 w. low iron[4]	14 h day at 22°C, 10 h night at 18°C, 60% rel. hum., ~550 μmol m^{-2} s^{-1} light int.

[1]Soil conditioning was performed in the field, and soil cores harvested at the conclusion (collection) of the conditioning.
[2]Phytotron Facility, University of Basel, Switzerland
[3]Fields were managed according to conventional Swiss farming practices, including the use of agrochemicals.
[4]100 mL of 0.2% Plantaaktiv Typ K (Hauert HBG Duenger AG, Grossaffoltern, Switzerland) and 0.0001% Sequestrene Rapid (Maag, Westland Schweiz GmbH, Dielsdorf, Switzerland)
[5]200 mL of 0.2% Plantaaktiv Typ K, 0.02% Sequestrene Rapid.

Table S3: Primer names and sequences used in this study

Primer name	Purpose	Primer sequence
*mmf1-1*_LP	Genotyping	GAAGGCTTTTCCAAATTCACC
*mmf1-2*_RP	Genotyping	ACCAAAATTCCATTAAACCCG
*mmf1-2*_LP	Genotyping	TAAGGATAATCGGGATTTGGG
*mmf1-2*_RP	Genotyping	TCCTGAAACATGAGCCAAAAG
LBb1.3	Genotyping	ATTTTGCCGATTTCGGAAC
*mmf1*_Fw	*qPCR*	TCGCATGTGAGGCATACGAA
*mmf1*_Rev	*qPCR*	GCGGTGATGAACGAATTGCT
PP2A_Fw	*qPCR*	CCTGCGGTAATAACTGCATCT
PP2A_Rev	*qPCR*	CTTCACTTAGCTCCACCAAGCA
CS1-*fs*-799-F	Microbiome	<u>ACACTGACGACATGGTTCTACA</u>*fs*AACMGGATTAGATACCCKG
CS2-*fs*-1193R	Microbiome	<u>TACGGTAGCAGAGACTTGGTCT</u>*fs*ACGTCATCCCCACCTTCC
CS1-At.GI-F	Microbiome	<u>ACACTGACGACATGGTTCTACA</u>CTGTAAAGATAAATGGGTCATCTAA
CS2-At-GI-R	Microbiome	<u>TACGGTAGCAGAGACTTGGTCT</u>AAGGGTTCAGCTTTGTCAACAA
ITS1-F	Microbiome	<u>ACACTGACGACATGGTTCTACA</u>CTTGGTCATTTAGAGGAAGTAA
ITS2-R	Microbiome	<u>TACGGTAGCAGAGACTTGGTCT</u>GCTGCGTTCTTCATCGATGC
ITS1-O_F1_G-46636	Microbiome	<u>ACACTGACGACATGGTTCTACA</u>CGGAAGGATCATTACCAC
5.8s-O_R1_G-46637	Microbiome	<u>TACGGTAGCAGAGACTTGGTCT</u>AGCCTAGACATCCACTGCTG