Deletion of core septin gene *aspB* in *Aspergillus fumigatus* results in fungicidal activity of

caspofungin

Rebecca Jean Busch[1], Carson Doty[2], C. Allie Mills[3], Flutur Latifi[4], Laura E. Herring[3], Vjollca

Konjufca[4], and José M Vargas-Muñiz[1,5,6*]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, United States

[2]School of Biological Sciences, Southern Illinois University-Carbondale, Carbondale, Illinois,

United States

[3]Michael Hooker Metabolomics and Proteomics Core Facility, Department of Pharmacology,

University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States

[4]Microbiology Program, Southern Illinois University-Carbondale, Carbondale, Illinois, United

States

[5]Fralin Life Science Institute, Virginia Tech, Blacksburg, VA

[6]Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Tech, Blacksburg, VA

Running title: Septin response to caspofungin

* Corresponding author

E-mail: j.vargas.muniz@vt.edu (JVM)

Abstract:

Septins are a family of GTP-binding proteins found in many eukaryotic lineages. Although highly conserved throughout many eukaryotes, their functions vary across species. In *Aspergillus fumigatus*, the etiological agent of invasive aspergillosis, septins participate in a variety of processes such as cell wall organization of conidia, septation, and response to cell wall stress. Previous studies determined that the Δ*aspB* strain had a greater sensitivity to anti-cell wall drugs, especially the echinocandin caspofungin, yet mechanisms behind this augmented sensitivity are unknown. We performed cell viability staining of the deletion strains post-caspofungin exposure and found that the Δ*aspA*, Δ*aspB*, and Δ*aspC* strains have significantly lower cell viability. Concomitant with the reduced viability, deletion strains are more susceptible to caspofungin on solid media. These results indicate that the septin cytoskeleton is important for *A. fumigatus* survival in the presence of caspofungin. Due to the potential of improved therapeutic outcome, we followed up using a neutropenic murine model of invasive aspergillosis. Animals infected with the Δ*aspB* strain and treated with caspofungin showed improved survival compared to the animals infected with *akuB*^KU80 wild-type or complemented strains. Additionally, histological analysis showed reduced fungal burden and inflammation in the Δ*aspB* infected, caspofungin-treated group. Affinity purification coupled with quantitative proteomics identified proteins involved in the septin-dependent response to caspofungin, includng four candidate interactors involved in cell wall stress response. Deletion of these candidate genes resulted in increased susceptibility to caspofungin and moderately reduced viability post-drug exposure. Taken together, these data suggest that septin AspB contributes to the fungistatic response to caspofungin.

Author Summary:

Invasive aspergillosis is a pulmonary disease caused by the fungus *Aspergillus fumigatus* that primarily occurs in immunocompromised patients. Invasive aspergillosis has a high mortality rate, ranging from 50-90%. Therapy options are limited due to few available drugs with fungicidal activity and growing global drug resistance. Treatment typically starts with triazoles, which target the fungal cell membrane. If unsuccessful, an echinocandin, which targets the cell wall, is given as a salvage therapy. Echinocandins, including caspofungin, are fungistatic against *A. fumigatus*, slowing growth of the fungus rather than killing it. Due to this, echinocandins have a high therapeutic failure rate. Previous work suggests that deletion of the cytoskeletal septin genes increases sensitivity to caspofungin. Here we describe our finding that the septin genes *aspA, aspB*, and *aspC* are involved in the fungal response to caspofungin. Additionally, the deletion of *aspB* results in fungicidal activity of this otherwise fungistatic drug. These findings show promise for novel therapy options that block the septin-mediated response to caspofungin.

Introduction:

Aspergillus fumigatus is a ubiquitous environmental mold responsible for a wide range of opportunistic systemic and allergic pathologies [1]. One such pathology is invasive aspergillosis (IA), an invasive systemic infection most common in immunocompromised patients [2]. Over 300,000 cases of IA are reported yearly, leading to a mortality rate ranging between 30-90% [1,3]. The at-risk patient population for IA is increasing due to increasing numbers of immunomodulating therapies, as well as emerging global diseases [4–6]. Additionally, IA has the highest per-patient cost of any invasive fungal disease, costing the United States an estimated $1.3 billion per year [7].

IA begins when the conidia, or asexual spores, are inhaled by an individual. The conidia are easily aerosolized and are prevalent in both indoor and outdoor environments [8,9]. Due to their small size (~2.5 μm diameter) and hydrophobic nature, the conidia can travel down the airway into the terminal alveoli [10]. In healthy individuals, the mucociliary escalator removes conidia [8]. However, any remaining conidia can then germinate within the lung. Epithelial cells, alveolar macrophages, neutrophils, and other immune cells can detect the germinating conidia, which leads to the release of cytokines and phagocytosis of conidia by immune cells [8]. In immunocompromised patients, the lack of immune responses leads to the progression of IA. Germinated conidia can then establish polarity, leading to the formation of hyphae that invades host tissue. IA can disseminate within the host via the release of hyphal fragments into the bloodstream. Neutropenic patients, such as those on anti-rejection medications post-transplant, are at high risk for IA [11]. In neutropenic patients, disease often presents with abundant hyphal growths, angioinvasion, and intra-alveolar hemorrhage [12,13]. Patients undergoing corticosteroid-induced immunosuppression are also at risk of IA, as glucocorticoids lead to a reduction of pattern recognition receptor (PRR) signaling and inhibit lymphocyte activation [14,15]. Pneumonia, inflammatory necrosis, and minimal hyphal growths are seen in cases of IA in patients under corticosteroid-induced immunosuppression [12,13]. Therapeutic options to effectively treat IA are limited. The frontline treatment for IA is the triazole class of drugs [16]. Triazole treatments are often prolonged and are now common to use as antifungal prophylaxis in at-risk populations. Nonetheless, cases of triazole-resistant *A. fumigatus* infections are becoming more common. This increase in the incidence of azole resistance led to the inclusion of *A. fumigatus* as a critical priority in the World Health Organization 2022 Fungal Priority Pathogens List [17].

Echinocandins are a class of antifungals that target the cell wall via β-glucan synthase [18]. In

fungi such as *C. albicans*, echinocandins have a fungicidal effect [19]. In contrast, the

echinocandins are fungistatic against *A. fumigatus* [18]. For immunocompromised patients, the

echinocandins are often used as a salvage or secondary therapy in conjunction with triazoles as

they cannot reap the benefits of solely fungistatic medications [16]. Slowing the growth of the

fungi through fungistatic therapies then requires either the immune system or fungicidal drugs to

kill the fungi and clear disease [20,21]. Because of its fungistatic nature, caspofungin use as a

salvage treatment had a favorable response rate of only 45% [22]. Understanding the

mechanisms within fungi that render these medications fungistatic rather than fungicidal can aid

in developing new therapies that improve the efficacy of existing antifungal drugs.

Previous studies have shown that septins AspA, AspB, and AspC contribute to the *Aspergillus*

spp. response to the echinocandin caspofungin [23,24]. Septins are a family of highly conserved

eukaryotic GTP-binding proteins. In *A. fumigatus,* septins are primarily involved in septation,

conidiation, and response to cell wall stress [24]. *A. fumigatus* has five septins: AspA, AspB,

AspC, AspD, and AspE [25]. AspA-D are core mitotic septins involved in polymerizing into

hexameric and octameric complexes. They are orthologs of *Saccharomyces cerevisiae*'s

Cdc11, Cdc3, Cdc12, and Cdc10, respectively [26]. AspE is a group 5 septin that is present in

filamentous fungi and other eukaryotic lineages [27]. Septins also can scaffold proteins involved

in stress response pathways. *Candida albicans* septins and phospholipid PI(4,5)P_2 are

regulated in response to cell wall stress induced from caspofungin exposure [28]. They

relocalize at specific foci on the plasma membrane to activate the MAPK Mkc1, which then

relocalizes chitin synthase and results in chitin deposition in the cell wall [28–30]. Similarly,

Aspergillus nidulans septins have been associated with the cell wall integrity (CWI) pathway

[23]. Double deletion strains of *aspB* and CWI pathway MAPK *mpkA* showed a novel phenotype

when grown on caspofungin compared to the respective single deletion strains [23]. These data

suggest AspB may have a role in pathways that maintain CWI.

In this study, we found that septins AspA, AspB, and AspC contribute to fungal viability post-

caspofungin exposure in *A. fumigatus*. The deletion of *aspB* in particular produced a fungicidal

response. We also observed a higher chance of survival, reduced lung inflammation, and

reduced fungal burden was associated with Δ*aspB* infected mice treated with caspofungin than

Δ*aspB* strain treated with saline or *akuB*^KU80 and Δ*aspB::aspB* strains with either treatment in

our neutropenic murine model of IA. To gain deeper mechanistic insights, we conducted a

quantitative mass spectrometry-based proteomics analysis to identify candidate AspB-

interacting proteins during exposure to caspofungin. Based on our proteomics analysis, we

narrowed down six possible proteins involved in the fungal response to caspofungin that were

significantly increased (by at least two-fold) upon caspofungin exposure. Gene ontology (GO)

analysis indicated that these candidate protein interactors might have a role in cell wall function

and organization, suggesting that AspB may mediate cell wall responses through its interactions

with these candidates. Deletion strains for each candidate were generated and characterized to

obtain an understanding of their biology at basal conditions. Strains were then tested against

caspofungin, and four out of the six genes, *bgt1, gel2, nsdD*, and *mapA*, were implicated in

fungal viability after caspofungin exposure.

Results:

Deletion of aspA, aspB*, and aspC Reduces Viability Post-Caspofungin Exposure*

Previous work demonstrated the hypersensitivity of *A. fumigatus* Δ*aspB* strain and sensitivity of

Δ*aspC* strain to caspofungin on solid media [24]. In order to further confirm and characterize the

response of septins to echinocandins, we conducted a spore dilution assay (10^4-10^1 conidia, 1

μg/mL caspofungin), minimum effective concentration assay (MEC), and E-strip test (10^6

conidia) to visualize the susceptibility to caspofungin. Deletion of the core septin genes, *aspA,*

aspB, and *aspC*, resulted in increased sensitivity to caspofungin compared to the *akuB*[KU80] wild-

type strain, Δ*aspB::aspB* complemented strain, and other septin deletion strains (Fig. 1A,

Supplementary Table 4). Similarly, the Δ*aspA*, Δ*aspB*, and Δ*aspC* strains had a clearer and

larger zone of effect on the E-strip test (Fig. 1D). To explore whether this effect was exclusive to

caspofungin, we conducted an E-strip test to visualize the susceptibility to another drug in the

echinocandin class, micafungin. The Δ*aspA and* Δ*aspC* strains showed a slight increase in

sensitivity to micafungin, while Δ*aspB* had a more noticeable decrease in growth in the zone of

effect (Fig. 1D). Thus, this increase in susceptibility of the Δ*aspA,* Δ*aspB*, and Δ*aspC* strains is

not limited to caspofungin, albeit it is more pronounced.

AspB is Required for Fungistatic Response to Caspofungin in vitro

Since we observed an increase in susceptibility in all three of our drug susceptibility assays, we

hypothesized that the absence of AspA, AspB, and AspC leads to a fungicidal effect of

caspofungin. To determine viability of the deletion strains against caspofungin, we utilized

CFDA to determine cell viability post-caspofungin exposure. In basal conditions, 10^4 conidia

were grown on coverslips in GMM liquid media for 24 hours, followed by 1 hour incubation in 1

μg/mL CFDA. All strains are equally viable, indicating that the increased clearance is not due to

a reduction in viability of septin deletion strains (Fig. 2A). Strains were then grown for 48 hours

in the presence of 1 μg/mL caspofungin, followed by incubation with CFDA. The *aspA, aspB*,

and *aspC* deletion strains were significantly less viable (p<0.0001) after 48 hours of caspofungin

exposure, with only 0.92% of Δ*aspB* (p<0.0001) being viable (Fig. 1B, 2B). Δ*aspB* also does not

form many microcolonies in the presence of caspofungin. Δ*aspA* and Δ*aspC* form microcolonies

in the presence of caspofungin; however, only 16.3% and 18.5% of the colonies, respectively,

were viable at the time of microscopic examination (p<0.0001)(Fig. 1B, 2B). We repeated this

assay with the echinocandin micafungin in order to test if this reduction in viability was specific

to caspofungin or a general response to echinocandin exposure. In contrast to caspofungin

treatment, the majority of colonies from all strains were viable after micafungin treatment (Fig.

1C). This indicates that the reduction in viability is specific to caspofungin.

AspB is Involved in Response to Caspofungin in Mature Mycelium

It is possible that most of these phenotypes that we observe are due to caspofungin acting as

the mycelia germinates, as exemplified by the small microcolonies in the Δ*aspB* strain. Mature

mycelia would be a more clinically relevant growth stage, and for this reason, we decided to

determine the effect of caspofungin on mature mycelium. To test this, we used propidium iodide

(PI) stain to determine hyphal damage. Strains were grown in GMM+UU for 24 hours at 37°C

before being stained with PI solution and visualized. No hyphal damage was seen under basal

conditions (Fig. 3A). Following the same procedure, we grew strains then incubated them with

caspofungin (1 µg/ml) for 2 hours. Plates were washed and stained as before, then visualized.

Similar to our viability assays, only the Δ*aspB* strain exhibits extensive hyphal damage after 2

hours of exposure to caspofungin (Fig. 3B). Taken together, these results suggest that AspB is

needed for fungal response to caspofungin.

Deletion of aspB *improves caspofungin treatment efficiency in a neutropenic murine model of*

invasive aspergillosis

As we observed a strong fungicidal effect against the Δ*aspB,* we decided to determine if

deletion of *aspB* led to improved survival in our murine model of invasive aspergillosis. As the

Δ*aspB* strain has no discernable difference in virulence compared to the *akuB*[KU80] and

Δ*aspB::aspB* strains with respect to fungal burden and lung inflammation, any effect on survival

would be attributed to the role that AspB plays in fungal response to caspofungin [24]. To test

this hypothesis, neutropenia was induced in 6-week old male CD-1 mice using 175 mg/kg

cyclophosphamide and 40 mg/kg triamcinolone acetonide. Neutropenic mice were then

intranasally infected with 4×10^6 conidia of *akuB*[KU80], Δ*aspB*, and Δ*aspB::aspB* strains. They

were subsequently treated with either 2 mg/kg caspofungin or an equivalent volume of the

saline vehicle. All groups of mice treated with saline all had the first death on day 3 post-

infection (Fig. 4A). Mice inoculated with Δ*aspB* strain and treated with caspofungin had a 70%

probability of survival (p<0.0001)(Fig. 4B). Additionally, in this group the first death did not occur

until day 12 post-infection. Compared with the next highest groups, Δ*aspB* strain treated with

saline and *akuB*[KU80] treated with caspofungin had a 30% probability of survival (Fig. 4A). Lung

histology was then performed to visualize inflammation and fungal lesions. Concurrent with the

survival graph, animals infected with the Δ*aspB* strain treated with caspofungin had reduced

inflammation and stunted fungal growth compared to animals from the other groups (Fig. 4D).

Previous infection models also showed no difference in inflammation or fungal lesions between

untreated *akuB*[KU80], Δ*aspB*, and Δ*aspB::aspB* strains [24]. This suggests the caspofungin

treatment reduces growth or potentially has a fungicidal effect on the fungus without AspB *in*

vivo, as well.

AspB Interactome Changes Post Caspofungin Exposure

Previous work determined that septins' localization is altered by exposure to caspofungin in *C.*

albicans and *A. fumigatus* [24,28]. Additionally, the protein interactome of septin AspB was

altered after exposure to caspofungin in a qualitative proteomic experiment [31]. This work only

detected for the presence or absence of interactant proteins between basal and caspofungin

conditions, potentially missing interactions that occur in both conditions but change in

abundance [31]. To gain a more mechanistic insight into how AspB contributes to the fungal

response to caspofungin, we applied affinity purification coupled with quantitative proteomics.

An AspB-eGFP expressing strain was grown in GMM and GMM supplemented with 1 µg/mL

caspofungin for 24 hours. AspB-GFP was then purified using a GFP-Trap® affinity matrix, and

proteins bound to AspB-GFP were prepared and analyzed by LC-MS/MS to identify protein

interactors in each condition. Principal Component Analysis (PCA) indicates that the GMM- and

caspofungin-AspB interactome are distinctive from each other (Fig. S1A). A total of 226 proteins

were significantly decreased (fold change (FC) < -2, $p < 0.05$) and 106 proteins were

significantly increased (FC > 2, $p < 0.05$) upon caspofungin treatment (Fig. S1B). Among the

proteins increased upon caspofungin exposure was PpoA, a characterized fatty acid

monooxygenase. PpoA was also identified by a previous study investigating the AspB

interactome post-caspofungin exposure, indicating that our analyses were able to confirm

previously described caspofungin-specific AspB interactions [31]. Candidate protein interactors

that met the screening criteria were analyzed using FungiFun, a gene ontology tool, to assess

changes in biological processes [32]. Proteins that are known to be involved in hyphal growth

were overrepresented among the list of proteins increased upon caspofungin treatment (Fig.

S1C). In contrast, proteins that are known to be involved in protein folding were overrepresented

in the list of proteins decreased upon caspofungin treatment (Fig. S1D). Using this information,

we selected six genes that were significantly increased by at least two-fold (FC > 2, $p < 0.05$)

after caspofungin exposure and had a role or putative role in cell-wall related functions for

further investigation (Table S3).

Δbgt1, Δgel2, ΔnsdD*, and* ΔmapA *Have a Reduction in Conidiation, While* ΔnsdD *and* ΔpunA

Have a Growth Defect

Based on our proteomics and gene ontology analyses, we performed the deletion of all six

candidate genes (Table S3) to determine their role in septin-related phenotypes. We first

characterized each deletion strain to understand how the gene affects growth during basal

conditions. Major and biologically relevant radial growth defects are seen in Δ*nsdD* (37.8 mm

average diameter compared to the 85 mm diameter of the *akuB*[KU80] strain) and Δ*punA* (24.3 mm

average diameter compared to the 57.3 mm average diameter of the *akuB*[KU80] strain at

30°C)(Fig. 5A, 5D, Fig.S2). The slight reduction of radial growth in Δ*aspB* was also seen in a

previous study and deemed not biologically relevant, which is also the case for the Δ*gel2* strain

[24]. Similarly, a previously noted defect in conidiation was seen in Δ*aspB* [24]. Δ*bgt1,* Δ*gel2,*

Δ*nsdD*, and Δ*mapA* also have reduced conidiation, but not as severely as Δ*aspB* (Fig. 5B, 5E).

As prior work noted that Δ*aspB* strain had delayed septation, we were interested in determining

if any of the candidate gene deletion strains were also defective in septation [24]. To test this,

we measured apical compartment length as an indirect method of measuring potential septation

defects. Δ*bgt1* mutants have a similar increase in apical compartment length to Δ*aspB*,

suggesting a delay in septation (Fig. 5C). In contrast, Δ*gel1* mutants have apical compartments

similar to the *akuB*[KU80] wild-type strain (Fig. 5C), suggesting that Gel1 is dispensable for septa

formation. Δ*gel2* (70.4 μm) and Δ*mapA* (67.7 μm) mutants exhibit hyperseptation. Δ*nsdD* (37.7

μm) has a more drastic reduction in apical compartment length, but this phenotype may be due

to its growth defect (Fig. 5C).

Δbgt1, Δgel2, ΔnsdD*, and* ΔmapA *Have Increased Sensitivity to Caspofungin Exposure*

Since the candidate proteins showed increased interaction with AspB during caspofungin

exposure, we explored their potential role in mediating the AspB-dependent fungal response to

caspofungin. To test this, we conducted a spore dilution assay on GMM agar supplemented with

1 µg/mL caspofungin. 10^4, 10^3, 10^2, and 10^1 spores were plated and incubated at either 30°C or

37°C for 2 days. Deletion of the *aspB* gene displayed colony-level growth defects in the 10^3,

10^2, and 10^1 concentrations (Fig. 6A, 6C). No other deletion strains displayed as severe of a

sensitivity to caspofungin. Δ*bgt1,* Δ*gel2,* Δ*nsdD,* and Δ*mapA* lost full colony growth at 10^2 and

10^1 concentrations (Fig. 6A). In contrast, Δ*gel1* and Δ*punA* are similar to the *akuB*^KU80 wild-type

and Δ*aspB::aspB* complemented strains (Fig. 6A, 6C). The minimum effective concentration

(MEC) of caspofungin for Δ*bgt1,* Δ*gel2,* Δ*nsdD,* and Δ*mapA* was determined to be lower than

that of the wild-type, and equal to or lower than that of Δ*aspB* (Table S4). Δ*punA* was not

determined as CLSI standards require 37°C, and this strain cannot grow at that temperature.

Reduced Viability of Δbgt1, Δgel2, ΔnsdD*, and* ΔmapA *Post-Caspofungin Exposure*

Since *aspB* deletion leads to a fungicidal response to caspofungin, we investigated whether any

candidate gene deletion strains also exhibited reduced viability. All strains showed equal

viability in basal conditions (Fig. 7A, 7C). We then grew all strains for 48 hours in the presence

of 1 µg/mL caspofungin and determined their viability with CFDA. The *aspB* deletion strain

showed nearly no viability at both 30°C and 37°C, as previously shown in Figure 1B (Fig. 6B, 6D

and Fig. 7B, 7D). No other strain demonstrated the complete loss of viability phenotype

observed in the Δ*aspB*. The Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains form microcolonies in the

presence of caspofungin like the *akuB*^KU80 strain, but they had a reduction in viability

(Δ*bgt1*=37.4%, Δ*gel2*=39.5%, Δ*nsdD*=27.8%, Δ*mapA*=26.1% compared to *akuB*^KU80

=90.4%)(Fig 6B and Fig.7B). Δ*gel1* and Δ*punA* did not have a reduction in cell viability when

grown in caspofungin (Fig. 6B, 6D and Fig. 7B, 7D).

Bgt1, Gel2, NsdD, and MapA are Involved in the Response to Caspofungin in Mature Mycelium

Next, we determined if any of our candidate gene deletion strains were susceptible to hyphal

damage when exposed to caspofungin as mature mycelia. To visualize hyphal damage, we

utilized a propidium iodide (PI) stain. No hyphal damage was observed in basal conditions,

indicating that there is no defect in the cell wall at basal conditions (Fig. 8A, 8C). The

experiment was then repeated with a 2-hour incubation in GMM supplemented with 1 µg/mL

caspofungin prior to PI staining. Extensive hyphal damage was seen in Δ*bgt1*, Δ*gel2*, Δ*nsdD*,

and Δ*mapA* strains similar to Δ*aspB* after exposure to caspofungin (Fig. 8B). Taken together,

Bgt1, Gel2, NsdD, and MapA are involved in the fungal response to caspofungin in earlier

stages of growth and mature hyphae.

Candidate Genes are not involved in the response to other Cell Wall Disrupting Agents or in the

Caspofungin Paradoxical Effect

We further characterized the candidate gene deletion strains sensitivities to other cell wall

disturbing agents. To test this, we plated conidia on GMM agar and GMM supplemented with

either 1 µg/mL caspofungin, 100 µg/mL Congo red, 2 µg/mL nikkomycin Z, 5 µg/mL calcofluor

white, or 10 µg/mL calcofluor white and incubated for three days. Additionally, we plated 4

µg/mL caspofungin and incubated for five days to observe whether the deletion strains were still

capable of the caspofungin paradoxical effect. After incubation, Δ*aspB* shows increased

susceptibility to caspofungin and Congo red, and slight increase in susceptibility to nikkomycin Z

and calcofluor white (Fig. S3). We observed Δ*gel1* had a mild increase in susceptibility to Congo

red compared to the *akuB*[KU80] wild-type (Fig. S3). Δ*bgt1*, Δ*gel2*, and Δ*nsdD* had a mild increase

in susceptibility to 10 µg/mL calcofluor white, but not as pronounced as Δ*aspB* (Fig. S3). Δ*punA*

visually appeared to have a reduction in conidiation and abnormal colony morphology when

grown on nikkomycin Z, but no differences in growth when compared to the wild type strain (Fig.

S4). Taken together, these candidate genes might mediate fungal response to caspofungin

while not being a part of a general fungal cell wall stress response.

Discussion:

The septins are a highly conserved family of cytoskeletal proteins with a variety of cellular roles.

This includes roles in cell division, stress response, cytoskeleton organization, and scaffolding

[24,33–35]. Previous work noted the role of septins in response to cell wall stress. In *C.*

albicans, septins are required for proper cell wall regulation and mislocalize during exposure to

caspofungin [29,36]. Additionally, septins also mislocalized during basal conditions in deletion

strains of protein kinases involved in cell wall regulation, including Gin4, Kin3, Vps34, and Cbk1

[36]. In *A. nidulans*, *C. albicans*, and *S. cerevisiae*, chitin synthases are dependent on septins

for proper localization [29,37,38]. Previous work in *A. fumigatus* determined that the septins

have a role in response to cell wall stress, notably showing that Δ*aspB* exhibits a hypersensitive

response to caspofungin exposure on solid media [24]. Here, we aimed to further investigate the

septin response to caspofungin. We found that deletion of *aspA, aspB*, and *aspC* showed a

reduction in fungal viability post-caspofungin exposure, while deletion of *aspB* elicited a

fungicidal response. The deletion of *aspB* also increased hyphal damage in mature mycelia.

The Δ*aspB* response to caspofungin was also recapitulated in an *in vivo* murine model, with

Δ*aspB* infected mice showing higher survival rates, reduced inflammation, and delayed mortality

upon treatment with caspofungin.

The *Aspergillus spp.* septins form heteropolymers consisting of hexamers (AspA-C) and

octamers (AspA-D) [39]. Deletion of *aspD* still allows proper formation of hexamers but not

octamers. In our work, we found that *aspA, aspB*, and *aspC* deletion strains had reduced

viability to caspofungin (Fig. 1). While the *aspD* deletion strain was slightly more sensitive than

the *akuB*^{KU80} wild-type, Δ*aspB::aspB*, or Δ*aspE* strains, its increased susceptibility was not as

pronounced as that of the *aspA, aspB*, and *aspC* deletion strains. This suggests that the

presence of the hexamer is at least sufficient, if not necessary, for proper fungal response to

caspofungin. It is unknown if only the sole presence of the octamer would also be sufficient, as

deletion of *aspA-C* prevents the formation of both structures. Similarly, Δ*aspB* strain had a

hypersensitive to fungicidal response to clinically relevant doses of caspofungin (Fig. 1), more

drastic than the one observed in the Δ*aspA* or Δ*aspC* strains. It is possible that AspB

specifically is needed to scaffold necessary proteins in response to caspofungin, but the loss of

both the hexamer and octamer causes general dysfunction that would lead to the phenotypes

seen in Δ*aspA* or Δ*aspC*. Proteomics analysis in *Cryptococcus neoformans* also supports the

possibility of subunit-specific protein interactions, where the *cdc3*^{aspB} and *cdc10*^{aspD} septins are

shown to have differences in interactomes after exposure to cellular stressors [40]. All of the

septin deletion strains retain the caspofungin paradoxical effect, suggesting that individual

septins and their heteropolymers act independently of the pathways that contribute to the

caspofungin paradoxical effect.

We found that the septin response to echinocandins is not limited to caspofungin, but the

fungicidal response of Δ*aspB* is specific to caspofungin. The echinocandin class includes four

clinically approved drugs: anidulafungin, micafungin, caspofungin, and most recently approved,

rezafungin [41,42]. These drugs all work through noncompetitive inhibition of the β-1,3-D-glucan

synthase, but alterations in their side chains comprise the major differences between each one

[18,43]. These changes in the side chains can lead to changes in how the drug impacts fungal

physiology. For instance, caspofungin is the only echinocandin to induce the paradoxical effect

[43]. Unlike micafungin, high concentrations of caspofungin increase the levels of cytosolic

calcium [44,45]. This activates calmodulin-calcineurin signaling and leads to the caspofungin

paradoxical effect [45]. Thus, our findings add to the growing evidence that even though the

different echinocandins have a shared mechanism of action, how the fungus responds to

echinocandins is drug specific.

Hyphal damage was not observed in any of our strains, indicating that there is no underlying cell

wall defect that would cause damage in basal conditions. However, after 2-hour exposure to

caspofungin, the ΔaspB strain showed significant hyphal damage. This suggests that AspB is

involved in the fungal cell wall response to caspofungin in mature mycelium. In *C. albicans*,

caspofungin has a fungicidal effect. Phosphatidylinositol-(4,5)-bisphosphate and septin

regulation are needed for proper cell wall stress response [28]. Disruption of either party causes

hypersusceptibility to caspofungin [28,29]. Based on our analyses, AspB is needed for proper

cell wall response to caspofungin in *A. fumigatus*. Previous proteomic analyses and our own

proteomics experiments show that MpkA, MkkA, and other cell wall integrity kinases co-

immunoprecipitated with AspB [31]. It is possible that AspB interacts with the cell wall integrity

pathway in *A. fumigatus* to facilitate fungal response to caspofungin [31]. In *A. nidulans,* deletion

of *aspB* partially rescues both the growth defect and susceptibility to caspofungin of the Δ*mpkA*

strain, further hinting a possible crosstalk between the septin and the CWI [23]. Further work will

be needed to investigate the possible role of AspB in the cell wall integrity pathway.

Animals infected with the Δ*aspB* strain and treated with caspofungin had a 70% chance of

survival, which is higher than that of animals infected with *akuB*[KU80] strain (30%). The first death

of Δ*aspB* strain infected mice treated with caspofungin was recorded on day +12 post infection,

compared to Δ*aspB* strain infected mice treated with saline or the other strains and conditions

with first deaths on day +3 or day +4. It is possible that colonies that persist under caspofungin

treatment are capable of growing and establishing infection after the drug pressure is released.

Our histological analyses showed reduced inflammation and reduced fungal lesions in the lungs

of Δ*aspB* infected mice treated with caspofungin. However, this assay is done on day +3 after

infection when mice are still being treated with caspofungin. Nonetheless, this increase in

survival makes AspB a prospective target for drug therapy developments, as use in conjunction

with caspofungin elicits a fungicidal response in *A. fumigatus*. Currently, only the plant cytokinin

forchlorfenuron (FCF) is known to disrupt septin organization [46]. This compound disrupts

septins within both fungal and mammalian cells by interfering with their ability to bind and

hydrolyze GTP [46–48]. It is possible that FCF could be modified and refined to target fungal-

specific septins. FCF analogues have previously been developed to better target ovarian and

endometrial cancers [49]. Thus, FCF analogues could be used in conjunction with caspofungin

to create a new fungicidal therapy for treatment of IA. Further work is required to better

understand the mechanisms of AspB's response to caspofungin and to develop novel clinical

therapies.

Our proteomic analysis uncovered six candidate protein interactors of septin AspB that were

significantly increased upon caspofungin treatment and that have a potential role in cell wall

functions. Bgt1, Gel1, and Gel2 are glucanosyltransferases in *A. fumigatus* [50]. These

glucanosyltransferases work to remodel unorganized β-1,3-glucan chains in the periplasmic

space to stabilize and modify the cell wall as needed [50]. Due to their role in cell wall

homeostasis, previous work identified these three glucanosyltransferases as potential drug

targets that required further investigation [50]. Bgt1 works by hydrolyzing a β-1,3-linked

oligosaccharide and placing it on another molecule of β-1,3-glucan [51]. In *C. albicans*, deletion

of *BGL2*, the *bgt1* homologue, resulted in an increase in susceptibility to nikkomycin Z, a chitin

synthase inhibitor [52]. In contrast, *A. fumigatus* Δ*bgt1* only has a slight increase in susceptibility

to nikkomycin Z (Fig. S3). Deletion of *BGL2* in *S. cerevisiae* increased the cellular chitin content

[53]. In *A. fumigatus,* though, Δ*bgt1*Δ*bgt2* double mutants do not have a change in chitin content

[51]. A slight reduction in conidiation was observed in *A. fumigatus* Δ*bgt1* (Fig. 5A, 5B). *A.*

fumigatus Δ*bgt1* susceptibility to calcofluor white and Congo red were similar to the *akuB*[KU80]

wild-type, although compared to basal conditions there was a reduction in growth (Fig. S3) [54].

A potential mechanism for the increased susceptibility in *A. fumigatus* Δ*bgt1* and Δ*aspB* to

caspofungin is that hyphal damage may be exacerbated by the lack of interaction between

AspB and Bgt1 when either party is missing (Fig. 6B). Due to the role of septins in septation and

scaffolding, the loss of AspB in particular may be more deleterious than just the loss of Bgt1 and

thus result in the observed fungicidal effect (Fig. 1B). Further work is needed to uncover the

relationship between glucanosyltransferase Bgt1 and the septins in septation and cell wall

stress responses.

Gel1 and Gel2 are two GPI-anchored glucanosyltransferase members of the Glycoside

Hydrolase Family 72 (GH72) [55]. Despite having the same enzymatic activity, deletion of *gel1*

did not result in a unique phenotype, but reduced mycelial growth and abnormal cell wall

architecture was observed in Δ*gel2* during basal conditions (Fig. 5A-B) [50,51,56]. An increase

in chitin is associated with deletion of *gel2*, a compensatory mechanism similar to deletion of

bgt1 [56]. *A. fumigatus* Gel1 is dispensable for proper fungal response to caspofungin (Fig. 6 A-

B). In contrast, Gel2 does appear to be involved in the response to caspofungin due to

increased susceptibility to caspofungin and reduced viability in caspofungin compared to

akuB^KU80 wild-type (Fig. 6A-B). Our proteomics analysis noted an increased interaction between

Gel2 and AspB, but cellular levels of Gel2 were not explored. It is unknown if cellular levels of

Gel2 are increased in *A. fumigatus* post-caspofungin exposure similar to *C. albicans*.

Additionally, further work exploring overexpression of Gel2 may uncover a new mechanism of

resistance to caspofungin in *A. fumigatus*.

NsdD (**n**ever in **s**exual **d**evelopment) is a putative GATA-type transcriptional activator with roles

in *A. nidulans* for sexual development [57]. NsdD is responsible for activating sexual

development [58]. Deletion of *nsdD* results in the inability to produce cleistothecia (fruiting

bodies), while overexpression allows production of Hülle cells even in conditions which typically

block sexual development [58]. Despite being a primarily asexual fungus, *A. fumigatus* is

capable of and still holds a variety of functional sexual reproductive genes [59]. NsdD has been

implicated in cell wall remodeling and hyphal fusion [60]. Deletion of *nsdD* results in reduced

hyphal growth on minimal media (Fig. 5B) [57,60]. Additionally, mating done with Δ*nsdD*

resulted in impaired heterokaryon formation [60]. Drug challenges show some sensitivity of

Δ*nsdD* to Congo red and caspofungin (Fig. 6A-B, Fig. S3) [60]. Previous work also noted

weakened hyphal tips in Δ*nsdD*, which is in accordance with the increased hyphal damage

during cell wall stress seen via PI staining (Fig. 8B) [60]. We suspect that while NsdD may have

independent roles in sexual development, it is possible that previous cell wall remodeling

functions of NsdD seen during hyphal fusion for heterokaryon formation may have alternative

functions in *A. fumigatus* cell wall remodeling outside of sexual development. Further work is

required to understand the NsdD-mediated response to caspofungin exposure.

MapA (gene *Afu4g06930*) is a previously uncharacterized protein in *A. fumigatus*. In *S.*

cerevisiae and *C. albicans*, cytosolic protein MAP2 is an ortholog that functions as a methionine

aminopeptidase. Deletion of MAP2 results in slightly slower growth rate and some chemical

sensitivities in *S. cerevisiae* and *C. albicans* [61,62]. It is possible that this methionine

aminopeptidase can remove the methionine to promote maturation of protein involved in

response to caspofungin.

PunA (gene *Afu6g07470*) is a previously uncharacterized protein in *A. fumigatus*. In yeast,

PUN1 is a Sur7 family tetraspan that localizes in eisosomes/MCC (membrane compartment of

Can1) [63,64]. These punctate membrane domains have roles in stable furrowing and promote

stress resistance, including detecting and responding to changes in cell wall integrity [63].

Deletion of *PUN1* resulted in decreased thermotolerance, abolished filamentous growth,

reduced cell wall components, and decreased metal ion tolerance [65–67]. This decreased

thermotolerance was also observed in our *A. fumigatus* Δ*punA* strain, thus requiring the growth

of Δ*punA* and comparison strains at 30°C. Yeast deletion strains of *SUR7*, a closely related

protein, saw little sensitivity to caspofungin but high sensitivity to the triazole fluconazole [68].

Interestingly, a previous study in *A. fumigatus* noted an increase in gene expression of *punA*

post-exposure to the triazole voriconazole [69]. *A. fumigatus* Δ*punA* had a similar phenotype to

the *akuB*KU80 wild-type when exposed to caspofungin or other cell wall stressors (Fig 6C-D, Fig.

S4). It is possible that *A. fumigatus* PunA contributes to the fungal response to non-cell wall

stressors, such as metal ion stress or membrane targeting drugs. It is also possible that PunA

might contribute with septin localization post-caspofungin exposure; however, this possible role

in septin localization does not contribute to *A. fumigatus* susceptibility to caspofungin.

There is increasing evidence that the septins are involved in the mediation of cell wall stress

[23,24,28,29,31]. In this work, we demonstrate that the septin hexamer is sufficient for response

to caspofungin and the deletion of the septin gene *aspB* results in a fungicidal effect. This

phenotype continues *in vivo*, suggesting potential therapeutic applications. We also identified

four AspB interactors that are involved, but not required, in the fungal response to caspofungin.

Overall, we suggest that the deletion of *aspB*, and thus loss of all AspB's interactions, results in

the fungicidal effect of caspofungin demonstrated. Future studies directed towards developing

drugs specific to fungal septins or in determining additional septin interactants that have key

roles in the caspofungin response can help to better understand ways in which caspofugnin

therapy can be improved for use against invasive aspergillosis.

Materials and Methods:

Strains, media, and conditions:

A. fumigatus akuB[KU80] served as the wild-type and control strain. Δ*aspB::aspB* complemented

strain served as an additional control to determine that phenotypes were due to the deletion of

the *aspB* gene. Septin deletion strains are described in (Table S1, [24,70]). The *aspB-egfp*

strain from [24], with *aspB-egfp* expressed by the *aspB* promoter from the native loci, was used

to pulldown AspB interactants. All cultures were plated on glucose minimal media (GMM)

supplemented with 5 mM uracil and 5 mM uridine (GMM+UU) and incubated at 37°C, unless

otherwise specified.

Drug susceptibility assays:

Spores were counted, diluted, and inoculated in rows onto agar plates containing caspofungin

(1 µg/mL) in 10µL quantities with a total of 10^4, 10^3, 10^2, and 10^1 conidia. Plates were incubated

for 48 hours at 37°C unless otherwise noted. For the minimum effective concentration assay, a

quantity of 2.5×10^4 spores were added to RMPI media with different concentrations of

caspofungin and incubated for 48 hours at 37°C [71]. We were not able to test the *aspD* and

punA deletion strains according to CLSI guidelines as Δ*aspD*, which is an uracil/uridine

auxotroph, could not grow in RMPI, and Δ*punA*, which has a temperature sensitivity, could not

grow at 37°C.

Fungal Viability Assay:

Conidia of *akuB*^{KU80}, Δ*aspA-E*, and Δ*aspB::aspB* were diluted to 10^4 spores and cultured on

coverslips immersed in 4 mL of GMM+UU broth and GMM+UU+caspofungin (1 µg/ml) and

incubated at 37°C for 24 and 48 hours, respectively. To examine cell viability, coverslips were

incubated in 5-carboxyfluorescein diacetate (CFDA) (50 µg/mL 0.1 M MOPS pH 3) for 1 hour at

37°C and 70 rpm. Slides were prepared for microscopy and image on an Inverted Leica DMi8

with Leica K5 Microscope Camera using a 10x objective. Images were analyzed using imageJ

[72]. Viable colonies were quantified by counting colonies manually and dividing fluorescent

colonies over total colonies counted. A minimum of 50 colonies per strain per replicate were

counted.

Hyphal damage assay:

Conidia (10^4) of *akuB*[KU80], Δ*aspA-E,* and Δ*aspB::aspB* were cultured on coverslips immersed in

4 mL of GMM+UU broth and incubated for 24 hours at 37°C. Coverslips were then incubated

with GMM+UU or GMM+UU+caspofungin (1 μg/ml) for 2 hours at 37°C. To examine the

damage of mature hyphae, coverslips were washed with 4 mL PIPES (pH 6.7) for 5 minutes.

PIPES was then removed and washed again with 4 mL PIPES for 5 minutes. After removing the

second PIPES wash, 500 uL of propidium iodide (PI) solution (12.5 ug/mL in 50 mM PIPES)

was added on top of the coverslip and let incubate in the dark for 5 minutes. Slides were

washed in 4 mL PIPES twice as described previously. They were then prepared for microscopy

and imaged on an Inverted Leica DMi8 with Leica K5 Microscope Camera using a 10x objective.

Images were analyzed in imageJ [72].

Neutropenic murine model of Invasive Aspergillosis:

Murine experiments followed previously established intranasal neutropenic models of IA [73,74].

Sixty 6-week old male CD1 mice (Charles River Laboratories, Raleigh, NC) were injected via

intraperitoneal route with 175 mg/kg cyclophosphamide on days −2 and +3 and 40mg/kg

triamcinolone acetonide subcutaneously on days −1 and + 6. Twenty mice per strain were

infected intranasally with 40 μl of 10^8 spores/ml conidia suspension of the *akuB*[KU80], Δ*aspB*, or

Δ*aspB::aspB* strains on day +0. On days +1 through +4, mice were injected via intraperitoneal

route with either 2 mg/kg caspofungin or saline. Mice were monitored until day +14 and

humanely euthanized if they showed severe symptoms. Survival was plotted on a Kaplan-Meier

curve with a log rank pair-wise comparison. Murine experiments were conducted in compliance

with the SIU Institutional Animal Care and Use Committee Protocol 20-034.

Histopathology Analysis:

Mice were immunocompromised and treated as described above in [Section: *Neutropenic*

murine model of Invasive Aspergillosis]. Lungs were harvested on day +3 after infection and

tissue sections were stained using hematoxylin and eosin (H&E) stains to visualize inflammation

and Gomori's methenamine silver stain to visualize fungal hyphae.

Protein extraction, AspB-eGFP fusion protein purification, and LC-MS/MS Analysis

The *aspB-egfp* strain from [24] was grown in GMM liquid media and GMM liquid media

supplemented with 1 μg/mL of caspofungin for 24 hours at 37°C. Protein extraction and

pulldown were completed as described by [31,75]. Fungal mycelia were homogenized to obtain

total cell lysate and AspB complexes were purified using the GFP-Trap® affinity purification

(Chromotek), as described [75]. GFP-Trap® magnetic beads were equilibrated by washing

beads three times in 500 μl ice-cold dilution buffer then resuspended in 100 μl ice cold dilution

buffer. The resin suspension is then mixed with total cell lysate (10 mg total protein) and

incubated at 4°C with gentle agitation for 2 hours. Beads were collected using a magnetic stand.

Beads were washed in 500 μl ice-cold lysis buffer and five times with 500 μl of wash buffer.

Beads were suspended in 50 μl wash buffer.

Samples were digested on-bead with trypsin followed by C18 desalting. Samples were analyzed

via LC-MS/MS on a Thermo Easy nLC 1200-QExactive HF in technical duplicate. All mass

spectra data was processed using MaxQuant (ver. 1.6.12.0) and searched against the Uniprot

Aspergillus fumigatus proteome reviewed database (Proteome ID UP000002530). MaxQuant

output was further processed via Perseus with filtering at 1% false discovery rate (FDR). Only

proteins with >1 peptide were reported.

Prioritization of candidate genes and generation of deletion mutants

Candidate genes were chosen by prioritizing proteins with at least a two-fold increase, statistical

significance, and known or putative roles in cell wall functions as listed in FungiFun GO term

search for biological processes (Table S3) [32]. Deletions of *bgt1* (*Afu1g11460*; fungidb.org),

gel1 (*Afu2g01170*; fungidb.org), *gel2* (*Afu6g11390*; fungidb.org), *nsdD* (*Afu3g13870*;

fungidb.org), *mapA* (*Afu4g06930*; fungidb.org), and *punA* (*Afu6g07470*; fungidb.org) genes

were obtained by replacing the gene with the 2.4 kb *pyrG* gene from *Aspergillus parasiticus.*

Approximately 1 kb of promoter and terminator region of each gene were PCR-amplified from

AF293 genomic DNA. Deletion constructs were generated by overlap fusion PCR and

subsequently transformed into *akuB*[KU80] *pyrG–* strain, all as previously described by [73].

Primers used in transformation of all candidate strains are found in (Table S2). Transformants

were validated via PCR screening.

Acknowledgments:
592

593   R.J.B. was supported by the Southern Illinois University Carbondale Master's Fellowship. R.J.B.

594   and J.V.M. were supported in part by 1R01AI165656-01A1. We also want to thank the members

595   of the Vargas-Muñiz group for critical reading of the manuscript. We want to thank Stacey

596   Mcgee (histology core facility) and Dr. Jennifer Harris (SIU Laboratory Animal Program Director)

597   for providing guidance and support during animal experiments.

References:

1. Bongomin F, Gago S, Oladele R, Denning D. Global and Multi-National Prevalence of Fungal Diseases—Estimate Precision. Journal of Fungi. 2017;3: 57. doi:10.3390/jof3040057

2. Latgé J-P, Chamilos G. Aspergillus fumigatus and Aspergillosis in 2019. Clin Microbiol Rev. 2019;33. doi:10.1128/CMR.00140-18

3. Rybak JM, Fortwendel JR, Rogers PD. Emerging threat of triazole-resistant Aspergillus fumigatus. Journal of Antimicrobial Chemotherapy. 2019;74: 835–842. doi:10.1093/jac/dky517

4. Li X, Lau SK, Woo PC. Fungal infection risks associated with the use of cytokine antagonists and immune checkpoint inhibitors. Exp Biol Med. 2020;245: 1104–1114. doi:10.1177/1535370220939862

5. Baddley JW. Clinical risk factors for invasive aspergillosis. Med Mycol. 2011;49: S7–S12. doi:10.3109/13693786.2010.505204

6. Gouzien L, Cocherie T, Eloy O, Legriel S, Bedos J-P, Simon C, et al. Invasive Aspergillosis associated with Covid-19: A word of caution. Infect Dis Now. 2021;51: 383–386. doi:10.1016/j.idnow.2020.12.008

7. Benedict K, Jackson BR, Chiller T, Beer KD. Estimation of Direct Healthcare Costs of Fungal Diseases in the United States. Clinical Infectious Diseases. 2019;68: 1791–1797. doi:10.1093/cid/ciy776

8. Dagenais TRT, Keller NP. Pathogenesis of Aspergillus fumigatus in Invasive Aspergillosis. Clin Microbiol Rev. 2009;22: 447–465. doi:10.1128/CMR.00055-08

9.   Shlezinger N, Irmer H, Dhingra S, Beattie SR, Cramer RA, Braus GH, et al. Sterilizing

immunity in the lung relies on targeting fungal apoptosis-like programmed cell death.

Science (1979). 2017;357: 1037–1041. doi:10.1126/science.aan0365

10.   Latgé J-P. Aspergillus fumigatus and Aspergillosis. Clin Microbiol Rev. 1999;12: 310–

350. doi:10.1128/CMR.12.2.310

11.   Morgan J, Wannemuehler KA, Marr KA, Hadley S, Kontoyiannis DP, Walsh TJ, et al.

Incidence of invasive aspergillosis following hematopoietic stem cell and solid organ

transplantation: interim results of a prospective multicenter surveillance program. Med

Mycol. 2005;43: 49–58. doi:10.1080/13693780400020113

12.   Balloy V, Huerre M, Latgé J-P, Chignard M. Differences in Patterns of Infection and

Inflammation for Corticosteroid Treatment and Chemotherapy in Experimental Invasive

Pulmonary Aspergillosis. Infect Immun. 2005;73: 494–503. doi:10.1128/IAI.73.1.494-

503.2005

13.   Stergiopoulou T, Meletiadis J, Roilides E, Kleiner DE, Schaufele R, Roden M, et al. Host-

Dependent Patterns of Tissue Injury in Invasive Pulmonary Aspergillosis. Am J Clin

Pathol. 2007;127: 349–355. doi:10.1309/UJRV9DLC11RM3G8R

14.   Coutinho AE, Chapman KE. The anti-inflammatory and immunosuppressive effects of

glucocorticoids, recent developments and mechanistic insights. Mol Cell Endocrinol.

2011;335: 2–13. doi:10.1016/j.mce.2010.04.005

15.   Cain DW, Cidlowski JA. Immune regulation by glucocorticoids. Nat Rev Immunol.

2017;17: 233–247. doi:10.1038/nri.2017.1

16.   Patterson TF, Thompson GR, Denning DW, Fishman JA, Hadley S, Herbrecht R, et al.

Practice Guidelines for the Diagnosis and Management of Aspergillosis: 2016 Update by

the Infectious Diseases Society of America. Clinical Infectious Diseases. 2016;63: e1–

e60. doi:10.1093/cid/ciw326

17. World Health Organization. WHO fungal priority pathogens list to guide research,

development and public health action. 2022. Available:

https://www.who.int/publications/i/item/9789240060241

18. Denning DW. Echinocandin antifungal drugs. The Lancet. 2003;362: 1142–1151.

doi:10.1016/S0140-6736(03)14472-8

19. Dudiuk C, Berrio I, Leonardelli F, Morales-Lopez S, Theill L, Macedo D, et al. Antifungal

activity and killing kinetics of anidulafungin, caspofungin and amphotericin B against

Candida auris. Journal of Antimicrobial Chemotherapy. 2019;74: 2295–2302.

doi:10.1093/jac/dkz178

20. Lewis JS, Graybill JR. Fungicidal versus Fungistatic: what's in a word? Expert Opin

Pharmacother. 2008;9: 927–935. doi:10.1517/14656566.9.6.927

21. Graybill JR, Burgess DS, Hardin TC. Key issues concerning fungistatic versus fungicidal

drugs. European Journal of Clinical Microbiology & Infectious Diseases. 1997;16: 42–50.

doi:10.1007/BF01575120

22. Hiemenz JW, Raad II, Maertens JA, Hachem RY, Saah AJ, Sable CA, et al. Efficacy of

caspofungin as salvage therapy for invasive aspergillosis compared to standard therapy

in a historical cohort. European Journal of Clinical Microbiology & Infectious Diseases.

2010;29: 1387–1394. doi:10.1007/s10096-010-1013-0

23. Mela A, Momany M. Septins coordinate cell wall integrity and lipid metabolism in a

sphingolipid-dependent process. J Cell Sci. 2022;135. doi:10.1242/jcs.258336

24.   Vargas-Muñiz JM, Renshaw H, Richards AD, Lamoth F, Soderblom EJ, Moseley MA, et

al. The Aspergillus fumigatus septins play pleiotropic roles in septation, conidiation, and

cell wall stress, but are dispensable for virulence. Fungal Genetics and Biology. 2015;81:

41–51. doi:10.1016/j.fgb.2015.05.014

25.   Juvvadi PR, Fortwendel JR, Rogg LE, Steinbach WJ. Differential localization patterns of

septins during growth of the human fungal pathogen Aspergillus fumigatus reveal novel

functions. Biochem Biophys Res Commun. 2011;405: 238–243.

doi:10.1016/j.bbrc.2011.01.017

26.   Bertin A, McMurray MA, Grob P, Park S-S, Garcia G, Patanwala I, et al. Saccharomyces

cerevisiae septins: Supramolecular organization of heterooligomers and the mechanism

of filament assembly. Proceedings of the National Academy of Sciences. 2008;105:

8274–8279. doi:10.1073/pnas.0803330105

27.   Shuman B, Momany M. Septins From Protists to People. Front Cell Dev Biol. 2022;9.

doi:10.3389/fcell.2021.824850

28.   Badrane H, Nguyen MH, Blankenship JR, Cheng S, Hao B, Mitchell AP, et al. Rapid

Redistribution of Phosphatidylinositol-(4,5)-Bisphosphate and Septins during the Candida

albicans Response to Caspofungin. Antimicrob Agents Chemother. 2012;56: 4614–4624.

doi:10.1128/AAC.00112-12

29.   Badrane H, Nguyen MH, Clancy CJ. Highly Dynamic and Specific Phosphatidylinositol

4,5-Bisphosphate, Septin, and Cell Wall Integrity Pathway Responses Correlate with

Caspofungin Activity against Candida albicans. Antimicrob Agents Chemother. 2016;60:

3591–3600. doi:10.1128/AAC.02711-15

30.   Wagner AS, Lumsdaine SW, Mangrum MM, Reynolds TB. Caspofungin-induced β(1,3)-

689   glucan exposure in Candida albicans is driven by increased chitin levels. mBio. 2023.

690   doi:10.1128/mbio.00074-23

691   31.   Vargas-Muñiz JM, Renshaw H, Waitt G, Soderblom EJ, Moseley MA, Palmer JM, et al.

692   Caspofungin exposure alters the core septin AspB interactome of Aspergillus fumigatus.

693   Biochem Biophys Res Commun. 2017;485: 221–226. doi:10.1016/j.bbrc.2017.02.116

694   32.   Priebe S, Linde J, Albrecht D, Guthke R, Brakhage AA. FungiFun: A web-based

695   application for functional categorization of fungal genes and proteins. Fungal Genetics

696   and Biology. 2011;48: 353–358. doi:10.1016/j.fgb.2010.11.001

697   33.   HARTWELL L. Genetic control of the cell division cycle in yeast *1IV. Genes controlling

698   bud emergence and cytokinesis. Exp Cell Res. 1971;69: 265–276. doi:10.1016/0014-

699   4827(71)90223-0

700   34.   Gladfelter A. The septin cortex at the yeast mother-bud neck. Curr Opin Microbiol.

701   2001;4: 681–689. doi:10.1016/S1369-5274(01)00269-7

702   35.   McMurray MA, Thorner J. Septins: molecular partitioning and the generation of cellular

703   asymmetry. Cell Div. 2009;4: 18. doi:10.1186/1747-1028-4-18

704   36.   Blankenship JR, Fanning S, Hamaker JJ, Mitchell AP. An Extensive Circuitry for Cell Wall

705   Regulation in Candida albicans. PLoS Pathog. 2010;6: e1000752.

706   doi:10.1371/journal.ppat.1000752

707   37.   DeMarini DJ, Adams AEM, Fares H, Virgilio C De, Valle G, Chuang JS, et al. A Septin-

708   based Hierarchy of Proteins Required for Localized Deposition of Chitin in the

709   Saccharomyces cerevisiae Cell Wall. J Cell Biol. 1997;139: 75–93.

710   doi:10.1083/jcb.139.1.75

38. Fukuda K, Yamada K, Deoka K, Yamashita S, Ohta A, Horiuchi H. Class III Chitin

Synthase ChsB of Aspergillus nidulans Localizes at the Sites of Polarized Cell Wall

Synthesis and Is Required for Conidial Development. Eukaryot Cell. 2009;8: 945–956.

doi:10.1128/EC.00326-08

39. Hernández-Rodríguez Y, Masuo S, Johnson D, Orlando R, Smith A, Couto-Rodriguez M,

et al. Distinct Septin Heteropolymers Co-Exist during Multicellular Development in the

Filamentous Fungus Aspergillus nidulans. PLoS One. 2014;9: e92819.

doi:10.1371/journal.pone.0092819

40. Martinez Barrera S, Hatchell E, Byrum SD, Mackintosh SG, Kozubowski L. Quantitative

analysis of septin Cdc10 & Cdc3-associated proteome during stress response in the

fungal pathogen Cryptococcus neoformans. PLoS One. 2024;19: e0313444.

doi:10.1371/journal.pone.0313444

41. Ben-Ami R. Systemic Antifungal Therapy for Invasive Pulmonary Infections. Journal of

Fungi. 2023;9: 144. doi:10.3390/jof9020144

42. Syed YY. Rezafungin: First Approval. Drugs. 2023;83: 833–840. doi:10.1007/s40265-

023-01891-8

43. Perlin DS. Current perspectives on echinocandin class drugs. Future Microbiol. 2011;6:

441–457. doi:10.2217/fmb.11.19

44. Fortwendel JR, Juvvadi PR, Perfect BZ, Rogg LE, Perfect JR, Steinbach WJ.

Transcriptional Regulation of Chitin Synthases by Calcineurin Controls Paradoxical

Growth of Aspergillus fumigatus in Response to Caspofungin. Antimicrob Agents

Chemother. 2010;54: 1555–1563. doi:10.1128/AAC.00854-09

45.   Juvvadi PR, Muñoz A, Lamoth F, Soderblom EJ, Moseley MA, Read ND, et al. Calcium-

734   Mediated Induction of Paradoxical Growth following Caspofungin Treatment Is

735   Associated with Calcineurin Activation and Phosphorylation in Aspergillus fumigatus.

736   Antimicrob Agents Chemother. 2015;59: 4946–4955. doi:10.1128/AAC.00263-15

737   46.   Iwase M, Okada S, Oguchi T, Toh-e A. Forchlorfenuron, a phenylurea cytokinin, disturbs

738   septin organization in Saccharomyces cerevisiae. Genes Genet Syst. 2004;79: 199–206.

739   doi:10.1266/ggs.79.199

740   47.   Hu Q, Nelson WJ, Spiliotis ET. Forchlorfenuron Alters Mammalian Septin Assembly,

741   Organization, and Dynamics. Journal of Biological Chemistry. 2008;283: 29563–29571.

742   doi:10.1074/jbc.M804962200

743   48.   Angelis D, Karasmanis EP, Bai X, Spiliotis ET. In Silico Docking of Forchlorfenuron (FCF)

744   to Septins Suggests that FCF Interferes with GTP Binding. PLoS One. 2014;9: e96390.

745   doi:10.1371/journal.pone.0096390

746   49.   Kim KK, Singh RK, Khazan N, Kodza A, Singh NA, Jones A, et al. Development of Potent

747   Forchlorfenuron Analogs and Their Cytotoxic Effect in Cancer Cell Lines. Sci Rep.

748   2020;10: 3241. doi:10.1038/s41598-020-59824-4

749   50.   Beauvais A, Latgé J-P. Membrane and cell wall targets in Aspergillus fumigatus. Drug

750   Resistance Updates. 2001;4: 38–49. doi:10.1054/drup.2001.0185

751   51.   Gastebois A, Mouyna I, Simenel C, Clavaud C, Coddeville B, Delepierre M, et al.

752   Characterization of a New β(1–3)-Glucan Branching Activity of Aspergillus fumigatus.

753   Journal of Biological Chemistry. 2010;285: 2386–2396. doi:10.1074/jbc.M109.077545

52. Sarthy A V., McGonigal T, Coen M, Frost DJ, Meulbroek JA, Goldman RC. Phenotype in
Candida albicans of a disruption of the BGL2 gene encoding a 1,3-β-glucosyltransferase.
Microbiology (N Y). 1997;143: 367–376. doi:10.1099/00221287-143-2-367

53. Kalebina TS, Farkaš V, Laurinavichiute DK, Gorlovoy PM, Fominov G V., Bartek P, et al.
Deletion of BGL2 results in an increased chitin level in the cell wall of Saccharomyces
cerevisiae . Antonie Van Leeuwenhoek. 2003;84: 179–184.
doi:10.1023/A:1026034123673

54. Millet N, Latgé J-P, Mouyna I. Members of Glycosyl-Hydrolase Family 17 of A. fumigatus
Differentially Affect Morphogenesis. Journal of Fungi. 2018;4: 18. doi:10.3390/jof4010018

55. Zhao W, Lu Y, Ouyang H, Zhou H, Yan J, Du T, et al. N-Glycosylation of Gel1 or Gel2 is
vital for cell wall-glucan synthesis in Aspergillus fumigatus. Glycobiology. 2013;23: 955–
968. doi:10.1093/glycob/cwt032

56. Mouyna I, Morelle W, Vai M, Monod M, Léchenne B, Fontaine T, et al. Deletion of GEL2
encoding for a β(1–3)glucanosyltransferase affects morphogenesis and virulence in
Aspergillus fumigatus. Mol Microbiol. 2005;56: 1675–1688. doi:10.1111/j.1365-
2958.2005.04654.x

57. Große V, Krappmann S. The Asexual Pathogen Aspergillus fumigatus Expresses
Functional Determinants of Aspergillus nidulans Sexual Development. Eukaryot Cell.
2008;7: 1724–1732. doi:10.1128/EC.00157-08

58. Han K-H, Han K-Y, Yu J-H, Chae K-S, Jahng K-Y, Han D-M. The nsdD gene encodes a
putative GATA-type transcription factor necessary for sexual development of Aspergillus
nidulans. Mol Microbiol. 2001;41: 299–309. doi:10.1046/j.1365-2958.2001.02472.x

59. O'Gorman CM, Fuller HT, Dyer PS. Discovery of a sexual cycle in the opportunistic

fungal pathogen Aspergillus fumigatus. Nature. 2009;457: 471–474.

doi:10.1038/nature07528

60. Szewczyk E, Krappmann S. Conserved Regulators of Mating Are Essential for

Aspergillus fumigatus Cleistothecium Formation. Eukaryot Cell. 2010;9: 774–783.

doi:10.1128/EC.00375-09

61. Li X, Chang YH. Amino-terminal protein processing in Saccharomyces cerevisiae is an

essential function that requires two distinct methionine aminopeptidases. Proceedings of

the National Academy of Sciences. 1995;92: 12357–12361.

doi:10.1073/pnas.92.26.12357

62. Brown JA, Sherlock G, Myers CL, Burrows NM, Deng C, Wu HI, et al. Global analysis of

gene function in yeast by quantitative phenotypic profiling. Mol Syst Biol. 2006;2.

doi:10.1038/msb4100043

63. Lanze CE, Gandra RM, Foderaro JE, Swenson KA, Douglas LM, Konopka JB. Plasma

Membrane MCC/Eisosome Domains Promote Stress Resistance in Fungi. Microbiology

and Molecular Biology Reviews. 2020;84. doi:10.1128/MMBR.00063-19

64. Grossmann G, Malinsky J, Stahlschmidt W, Loibl M, Weig-Meckl I, Frommer WB, et al.

Plasma membrane microdomains regulate turnover of transport proteins in yeast. J Cell

Biol. 2008;183: 1075–1088. doi:10.1083/jcb.200806035

65. Hosiner D, Sponder G, Graschopf A, Reipert S, Schweyen RJ, Schüller C, et al. Pun1p is

a metal ion-inducible, calcineurin/Crz1p-regulated plasma membrane protein required for

cell wall integrity. Biochimica et Biophysica Acta (BBA) - Biomembranes. 2011;1808:

1108–1119. doi:10.1016/j.bbamem.2011.01.002

66.   Mir SS, Fiedler D, Cashikar AG. Ssd1 Is Required for Thermotolerance and Hsp104-

800   Mediated Protein Disaggregation in Saccharomyces cerevisiae. Mol Cell Biol. 2009;29:

801   187–200. doi:10.1128/MCB.02271-07

802   67.   Xu T, Shively CA, Jin R, Eckwahl MJ, Dobry CJ, Song Q, et al. A Profile of Differentially

803   Abundant Proteins at the Yeast Cell Periphery during Pseudohyphal Growth. Journal of

804   Biological Chemistry. 2010;285: 15476–15488. doi:10.1074/jbc.M110.114926

805   68.   Alvarez FJ, Douglas LM, Rosebrock A, Konopka JB. The Sur7 Protein Regulates Plasma

806   Membrane Organization and Prevents Intracellular Cell Wall Growth in Candida albicans.

807   Mol Biol Cell. 2008;19: 5214–5225. doi:10.1091/mbc.e08-05-0479

808   69.   Ferreira ME da S, Malavazi I, Savoldi M, Brakhage AA, Goldman MHS, Kim HS, et al.

809   Transcriptome analysis of Aspergillus fumigatus exposed to voriconazole. Curr Genet.

810   2006;50: 32–44. doi:10.1007/s00294-006-0073-2

811   70.   da Silva Ferreira ME, Kress MRVZ, Savoldi M, Goldman MHS, Ha□rtl A, Heinekamp T,

812   et al. The akuBKU80 Mutant Deficient for Nonhomologous End Joining Is a Powerful Tool

813   for Analyzing Pathogenicity in Aspergillus fumigatus. Eukaryot Cell. 2006;5: 207–211.

814   doi:10.1128/EC.5.1.207-211.2006

815   71.   Fortwendel JR, Juvvadi PR, Pinchai N, Perfect BZ, Alspaugh JA, Perfect JR, et al.

816   Differential effects of inhibiting chitin and 1,3-β-D-glucan synthesis in ras and calcineurin

817   mutants of Aspergillus fumigatus. Antimicrob Agents Chemother. 2009;53: 476–482.

818   doi:10.1128/AAC.01154-08

819   72.   Bourne R. ImageJ. Fundamentals of Digital Imaging in Medicine. London: Springer

820   London; 2010. pp. 185–188. doi:10.1007/978-1-84882-087-6_9

73.   Steinbach WJ, Cramer RA, Perfect BZ, Asfaw YG, Sauer TC, Najvar LK, et al.

Calcineurin Controls Growth, Morphology, and Pathogenicity in Aspergillus fumigatus.

Eukaryot Cell. 2006;5: 1091–1103. doi:10.1128/EC.00139-06

74.   Steinbach WJ, Benjamin Jr DK, Trasi SA, Miller JL, Schell WA, Zaas AK, et al. Value of

an inhalational model of invasive aspergillosis. Med Mycol. 2004;42: 417–425.

doi:10.1080/13693780410001712034

75.   Juvvadi PR, Belina D, Soderblom EJ, Moseley MA, Steinbach WJ. Filamentous fungal-

specific septin AspE is phosphorylated in vivo and interacts with actin, tubulin and other

septins in the human pathogen Aspergillus fumigatus. Biochem Biophys Res Commun.

2013;431: 547–553. doi:10.1016/j.bbrc.2013.01.020

Figures:

Fig 1. AspA, AspB, and AspC are involved in fungal response to caspofungin. The deletion of

aspA, aspB, and aspC results in reduction of viability to echinocandins. (A) Spore dilution assay

show an increase in susceptibility to caspofungin in the Δ*aspA,* Δ*aspB,* and Δ*aspC* strains. 10^4-

10^1 conidia were plated on GMM media supplemented with caspofungin for 48 hours at 37°C.

(B, C) Deletion of *aspB* results in loss of viability when grown in caspofungin but not micafungin.

10^4 conidia were incubated for 48 hours in GMM supplemented with (B) 1µg/mL caspofungin or

(C) 1 µg/mL micafungin. Cells were incubated in CFDA for one hour then visualized. Viable and

non-viable colonies were counted. The number of viable colonies was divided by the total

number of colonies. Experiment was replicated three times. Error bar represent SEM. Student t-

test were done in Graphpad Prism, with each strain compared against the *akuB*[KU80] wild-type.

(D) E-strip plates of caspofungin and micafungin show clearer zone of effect in Δ*aspA,* Δ*aspB*,

and Δ*aspC* strains. 10^6 conidia were plated with beads and left to dry. E-strip was placed, and

plates were grown at 37°C for 48 hours. All experiments were replicated three times.

Representative images are shown in this figure.

Fig 2. Δ*aspB* strain loses cell viability when grown in caspofungin. (A) All strains had viability

when grown in GMM+UU. 10^4 conidia were grown in 4 mL GMM+UU at 37°C for 24 hours. Cells

were then treated with CFDA, which is hydrolyzed in living cells to a fluorescent ester, then

visualized. (B) Loss of viability in Δ*aspB* strain is seen after growth in the presence of

caspofungin. After 48 hours post exposure to 1 µg/mL caspofungin, cells were incubated in

CFDA for one hour then visualized. Experiments were replicated three times. Scale bar is 500

µm.

Fig 3. Δ*aspB* strain has increased hyphal damage during caspofungin exposure.

(A) No hyphal damage seen in basal conditions. 10^4 conidia were grown in GMM at 37°C for 24

hours. Cells were then washed with PIPES (pH 6.7) for 5 minutes twice. Slides were then

treated with propidium iodide (PI) solution, which stains nucleic acids. Coverslips were washed

twice with PIPES, then prepared and visualized. (B) Hyphal damage was seen in Δ*aspB* strain

treated with caspofungin. 10^4 conidia were grown in GMM at 37°C for 24 hours. Coverslips were

then incubated in caspofungin for 2 hours at 37°C. Cells were then washed with PIPES and

treated with PI solution. Experiments were replicated three times. Scale bar is 500 µm.

Fig 4. The Δ*aspB* strain treated with caspofungin has reduced inflammation and fungal burden

in immunosuppressed intranasal murine model of invasive aspergillosis.

(A-B) Mice infected with Δ*aspB* strain and treated with caspofungin had a higher chance of

survival. Ten mice per strain per condition were infected intranasally with 4×10^6 conidia. Mice

were treated with either saline (A) or caspofungin (B) on days +1 through +4 post-infection. Mice

were monitored at least twice a day for 14 days. Survival is on a Kaplan-Meier curve with log

rank pair-wise comparison (p<0.0001). (C) H&E stain of lungs after 3 days post-infection shows

that there is a decrease in inflammatory response in Δ*aspB* infected mice treated with

caspofungin. Silver stain shows less and smaller fungal lesions in Δ*aspB* infected mice treated

with caspofungin. H&E images taken on an Echo Rebel Hybrid Light microscope using a 10x

objective. Scale bar is 300 µm. Silver stain images taken on an Inverted Leica DMi8 with Leica

K5 Microscope Camera using a 40x objective. Scale bar is 100 µm.

Fig 5. Δ*nsdD* and Δ*punA* strains have a growth defect; Bgt1, Gel2, NsdD, and MapA are

involved in conidiation; and Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains have septation defects.

(A, D) Δ*nsdD* and Δ*punA* have a significant defect in radial growth after 5 days incubation. 10^4

total conidia of each strain were placed on GMM agar. Plates were incubated for 5 days and

radial growth was measured every 24 hours. Experiments were replicated three times. (B, E)

Deletion of *aspB*, *bgt1*, *gel2*, *nsdD*, or *mapA* results in reduced conidial production. Δ*punA* has

increased conidial production. Conidia were collected on day 5 of growth on GMM media using

10 mL of 0.05% Tween-80. Conidia were counted using a hemocytometer and values were

divided by the total area of growth to account for strains with growth defects. Experiments were

replicated three times. (C) Δ*aspB* and Δ*bgt1* have an increase in length of the apical

compartment. Δ*gel2*, Δ*nsdD*, and Δ*mapA* have a decrease in length of the apical compartment.

10^4 total conidia of each strain were inoculated onto coverslips immersed in GMM and

incubated for 15 hours. Coverslips were stained with aniline blue and visualized. The length of

the apical compartments (N=20) was measured using imageJ. (A, B, C) Experiment performed

at 37°C. (D, E) Experiment performed at 30°C. Error bar represent SEM. One-way ANOVA with

Tukey's multiple-comparison test were performed in Graphpad Prism and were declared

significantly different at a p-value of <0.05. Group means with different lowercase letters are

significantly different.

Fig 6. Deletion of Bgt1, Gel2, NsdD, and MapA increases susceptibility to caspofungin.

(A, C) Spore dilution assays show Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains have an increase in

susceptibility to caspofungin. Conidia (10^4-10^1) were grown on GMM media supplemented with

1 µg/mL caspofungin for 48 hours. (B, D) Deletion of *bgt1*, *gel2*, *nsdD*, and *mapA* results in

reduced, but not loss of, viability when grown in caspofungin. Deletion of *gel1* and *punA* does

not affect viability when exposed to caspofungin. 10^4 conidia were grown in GMM for 24 hours

then incubated in CFDA and visualized. Experiment was replicated three times. Error bar

represent SEM. One-way ANOVA with Tukey's multiple-comparison test were performed in

Graphpad Prism and were declared significantly different at a p-value of <0.05. Group means

with different lowercase letters are significantly different. (A, B) Experiment performed at 37°C.

(C, D) Experiment performed at 30°C.

Fig 7. Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains have reduced cell viability when grown in

caspofungin. (A, C) All strains were fully viable when grown in GMM media. Conidia (10^4) were

grown in GMM for 24 hours on coverslips then incubated in CFDA. Slides were prepared and

visualized on an Inverted Leica DMi8 with Leica K5 Microscope Camera using a 10x objective.

(B, D) Candidate deletion strains did not lose total viability, but strains Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and

Δ*mapA* had a partial reduction in viability. Conidia (10^4) were grown on coverslips immersed in

GMM supplemented with 1 µg/mL caspofungin for 48 hours. Coverslips were then incubated in

CFDA and visualized. (A, B) Experiment performed at 37°C. (C, D) Experiment performed at

30°C. All experiments were replicated three times. Scale bar is 500 µm.

Fig 8. Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains have increased hyphal damage during

caspofungin exposure. (A, C) Hyphal damage was not observed in basal conditions. Conidia

(10^4) were grown on coverslips in GMM for 24 hours. Coverslips were washed in PIPES (pH

6.7) for 5 minutes twice then treated with propidium iodide (PI) solution. Coverslips were

washed twice with PIPES, then visualized on an Inverted Leica DMi8 with Leica K5 Microscope

Camera using a 10x objective. (B, D) Strains Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* had hyphal

damage when treated with caspofungin. Conidia (10^4) were grown on coverslips in GMM for 24

hours, then incubated in caspofungin for 2 hours. Cells were then washed with PIPES and

incubated with PI. (A, B) Experiment performed at 37°C. (C, D) Experiment performed at 30°C.

Experiments were replicated three times. Scale bar is 500 μm.

Supplementary Figure 1. Caspofungin exposure alters AspB protein interactions.

(A) Principal component analysis (PCA) of the proteomics data reveals different protein

composition from AspB pulldown from GMM and caspofungin treated samples. Blue color

denotes those grown in GMM, red color denotes fungi grown in GMM+ 1 μg/mL caspofungin.

(B) 226 significantly decreased (FC < -2, p < 0.05) proteins (blue) and 106 significantly

increased (FC > 2, p < 0.05) proteins (red). Proteins under two-fold change or had a p-value

greater than 0.05 are shown in black. Red horizontal line denotes p=0.05, red vertical lines

denote ± 2 fold change. (C) Category of biological processes most enriched after exposure to

caspofungin is hyphal growth. (D) Category of biological processes most decreased after

exposure to caspofungin is protein folding. (C,D) Gene ontology enrichment for biological

processes determined by using FungiFun [32].

Supplementary Figure 2. Δ*nsdD* and Δ*punA* strains have a growth defect.

Radial growth assay shown per each day, 10^4 conidia of each strain were inoculated on GMM

agar. Plates were then incubated for 5 days, and radial growth was measured every 24 hours.

(A) Assays performed at 37°C. (B) Assays performed at 30°C. All experiments done in triplicate.

Supplementary Figure 3. Δ*bgt1*, Δ*gel2*, and Δ*nsdD* have mild sensitivity to calcofluor white.

10^4 conidia were inoculated into GMM agar supplemented with cell wall disrupting agents listed

in the figure then incubated for 3 days at 37°C, with the exception of 4 µg/mL caspofungin for

five days to ensure observation of the caspofungin paradoxical effect. Experiments were

replicated three times.

Supplementary Figure 4. Deletion of *punA* results in changed colony morphology in nikkomycin

Z.

10^4 conidia were inoculated into GMM agar supplemented with the cell wall disrupting agents

listed in the figure then incubated for 3 days at 30°C, with the exception of 4 µg/mL caspofungin

for five days to ensure observation of the caspofungin paradoxical effect. Experiments were

replicated three times and representative images are shown.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4