Deletion of core septin gene *aspB* in *Aspergillus fumigatus* results in fungicidal activity of

caspofungin

Rebecca Jean Busch[1,2], Carson Doty[3], C. Allie Mills[4], Flutur Latifi[5], Laura E. Herring[4], Vjollca

Konjufca[5], Francisco Carvallo[6], Tatiana Boluarte[1], and José M Vargas-Muñiz[1,2,7*]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, United States

[2]Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Tech, Blacksburg,

Virginia, United States

[3]School of Biological Sciences, Southern Illinois University-Carbondale, Carbondale, Illinois,

United States

[4]Michael Hooker Metabolomics and Proteomics Core Facility, Department of Pharmacology,

University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States

[5]Microbiology Program, Southern Illinois University-Carbondale, Carbondale, Illinois, United

States

[6]Department of Biomedical Sciences and Pathobiology, Virginia-Maryland College of Veterinary

Medicine, Blacksburg, Virginia, United States

[7]Fralin Life Science Institute, Virginia Tech, Blacksburg, Virginia, United States

Running title: Septin response to caspofungin

* Corresponding author

E-mail: j.vargas.muniz@vt.edu (JVM)

Abstract

Septins are a family of GTP-binding proteins found in many eukaryotic lineages. Although highly

conserved throughout many eukaryotes, their functions vary across species. In *Aspergillus*

fumigatus, the etiological agent of invasive aspergillosis, septins participate in a variety of

processes, including conidiation, septation, and responses to cell wall stress. Previous studies

determined that the Δ*aspB* strain had a greater sensitivity to anti-cell wall drugs, especially the

echinocandins, yet mechanisms behind this augmented sensitivity are unknown. We performed

cell viability staining of the deletion strains after caspofungin exposure and found that the

Δ*aspA*, Δ*aspB*, and Δ*aspC* strains had significantly lower cell viability. Concomitant with the

reduced viability, deletion strains are more susceptible to caspofungin on solid media. These

results indicate that the septin cytoskeleton is important for *A. fumigatus* survival in the

presence of caspofungin. Due to the potential of improved therapeutic outcome, we followed up

using a neutropenic murine model of invasive aspergillosis. Animals infected with the Δ*aspB*

strain and treated with caspofungin showed improved survival compared to the animals infected

with *akuB*[KU80] wild-type or complemented strains. Additionally, histological analysis showed

reduced fungal burden and inflammation in the Δ*aspB* infected, caspofungin-treated group.

Affinity purification coupled with quantitative proteomics identified proteins involved in the

septin-dependent response to caspofungin, including four candidate interactors involved in cell

wall stress response. Deletion of these candidate genes resulted in increased susceptibility to

caspofungin and moderately reduced viability post-drug exposure. Taken together, these data

suggest that septin AspB contributes to the fungistatic response to caspofungin.

Author Summary

Invasive aspergillosis is a pulmonary disease caused by the fungus *Aspergillus fumigatus* that

primarily occurs in immunocompromised patients. Invasive aspergillosis has a high mortality

rate, ranging from 50-90%. Therapy options are limited due to few available drugs with

fungicidal activity and growing global drug resistance. Treatment typically starts with triazoles,

which target the fungal cell membrane. If unsuccessful, an echinocandin, which targets the cell

wall, is given as a salvage therapy in the U.S. Echinocandins, including caspofungin, are

fungistatic against *A. fumigatus*, slowing growth of the fungus rather than killing it. Due to this,

echinocandins have a high therapeutic failure rate. Previous work suggests that deletion of the

cytoskeletal septin genes increases sensitivity to caspofungin. Here we describe our finding that

the septin genes *aspA, aspB*, and *aspC* are involved in the fungal response to caspofungin.

Additionally, the deletion of *aspB* results in fungicidal activity of this otherwise fungistatic drug.

These findings show promise for novel therapy options that block the septin-mediated response

to caspofungin.

Introduction

Aspergillus fumigatus is a ubiquitous environmental mold responsible for a wide range of

opportunistic systemic and allergic pathologies [1]. One such pathology is invasive aspergillosis

(IA), an invasive infection most common in immunocompromised patients [2]. Over 300,000

cases of IA are reported yearly, leading to a mortality rate ranging between 30-90% [1,3].

Additionally, the at-risk patient population for IA is increasing due to increasing numbers of

immunomodulating therapies, as well as emerging global diseases [4–6]. IA has the highest per-

patient cost of any invasive fungal disease, costing the United States an estimated $1.3 billion

per year [7].

IA begins when the conidia, or asexual spores, are inhaled by an individual [8,9]. Due to their

small size (~2.5 μm diameter) and hydrophobic nature, the conidia can travel down the airway

into the terminal alveoli [10]. In healthy individuals, the mucociliary escalator removes conidia

[8]. However, any remaining conidia can then germinate within the lung. Epithelial cells, alveolar

macrophages, neutrophils, and other immune cells can detect the germinating conidia, which

leads to the release of cytokines and phagocytosis of conidia by immune cells [8]. However,

immunocompromised patients lack the appropriate immune responses, which leads to the

progression of IA. Germinated conidia can then establish polarity, leading to the formation of

hyphae that invades host tissue. IA can disseminate within the host via the release of hyphal

fragments into the bloodstream. Neutropenic patients, such as those on anti-rejection

medications post-transplant, are at high risk for IA [11]. In neutropenic patients, disease often

presents with abundant hyphal growths, angioinvasion, and intra-alveolar hemorrhage [12,13].

Patients undergoing corticosteroid-induced immunosuppression are also at risk of IA, as

glucocorticoids lead to a reduction of pattern recognition receptor (PRR) signaling and inhibit

lymphocyte activation [14,15]. Pneumonia, inflammatory necrosis, and minimal hyphal growths

are seen in cases of IA in patients under corticosteroid-induced immunosuppression [12,13].

Therapeutic options to effectively treat IA are limited. The frontline treatment for IA is the triazole

class of drugs [16]. Triazole treatments are often prolonged and are now common to use as

antifungal prophylaxis in at-risk populations. Nonetheless, cases of triazole-resistant *A.*

fumigatus infections are increasing in some countries and are a major concern worldwide. This

increase in the incidence of triazole resistance led to the inclusion of *A. fumigatus* as a critical

priority in the World Health Organization 2022 Fungal Priority Pathogens List [17].

Echinocandins are a class of antifungals that inhibit the β-glucan synthase [18]. In fungi such as

C. albicans, echinocandins have a fungicidal effect [19]. In contrast, the echinocandins are

fungistatic against *A. fumigatus* [18]. For immunocompromised patients in the United States, the

echinocandins are often used as a salvage or secondary therapy in conjunction with triazoles as

they cannot reap the benefits of solely fungistatic medications [16]. Slowing the growth of the

fungi through fungistatic therapies then requires either the immune system or fungicidal drugs to

kill the fungi and clear disease [20,21]. Because of its fungistatic nature, caspofungin use as a

salvage treatment had a favorable response rate of only 45% [22]. Understanding the

mechanisms within fungi that render these medications fungistatic rather than fungicidal can aid

in developing new therapies that improve the efficacy of existing antifungal drugs.

Previous studies have shown that septins AspA, AspB, and AspC contribute to the *Aspergillus*

spp. response to the echinocandin caspofungin [23,24]. Septins are a family of highly-conserved

eukaryotic GTP-binding proteins. In *A. fumigatus,* septins are primarily involved in septation,

conidiation, and response to cell wall stress [24]. *A. fumigatus* has five septins: AspA, AspB,

AspC, AspD, and AspE [25]. AspA-D are core mitotic septins involved in polymerizing into

hexameric and octameric complexes. They are orthologs of *Saccharomyces cerevisiae*'s

Cdc11, Cdc3, Cdc12, and Cdc10, respectively [26]. AspE is a Momany Group 5 septin that is

present in filamentous fungi and other eukaryotic lineages [27]. *Aspergillus nidulans* septins

have been associated with the cell wall integrity (CWI) pathway [23]. Double deletion strains of

aspB and CWI pathway MAPK *mpkA* showed a novel phenotype when grown on caspofungin

compared to the respective single deletion strains [23]. In *Candida albicans,* deletion of septin

genes leads to aberrant localization of chitin in the cell wall, increasing susceptibility to

caspofungin [28,29]. This susceptibility is due to caspofungin inducing aberrant cytokinesis and

phosphatidylinositol (4,5)-bisphosphate relocalization [28–30]. This points out a conserved role

of septins in the fungal response to caspofungin, but the mechanism might not be conserved.

In this study, we found that septins AspA*,* AspB, and AspC contribute to fungal viability post-

caspofungin exposure in *A. fumigatus*. The deletion of *aspB* in particular produced a fungicidal

response. We also observed a higher chance of survival, reduced lung inflammation, and

reduced fungal burden were associated with Δ*aspB* infected mice treated with caspofungin than

Δ*aspB* strain treated with saline or *akuB*[KU80] and Δ*aspB::aspB* strains with either treatment in

our neutropenic murine model of IA. To gain deeper mechanistic insights, we conducted a

quantitative mass spectrometry-based proteomics analysis to identify candidate AspB-

interacting proteins during exposure to caspofungin. Based on our proteomics analysis, we

narrowed down six possible proteins involved in the fungal response to caspofungin that were

significantly increased (by at least two-fold) upon caspofungin exposure. Gene ontology (GO)

analysis indicated that these candidate protein interactors might have a role in cell wall function

and organization, suggesting that AspB may mediate cell wall responses through its interactions

with these candidates. Deletion strains for each candidate were generated and characterized to

obtain an understanding of their biology at basal conditions. Strains were then tested against

caspofungin, and four out of the six genes, *bgt1, gel2, nsdD*, and *mapA*, were implicated in

fungal viability after caspofungin exposure. Double deletion mutants lacking *aspB* and either

bgt1, *gel2*, *nsdD*, and *mapA* were generated, which indicated that their role in caspofungin

response is mediated through AspB.

Results:

***Deletion of* aspA*,* aspB*, and* aspC *Reduces Viability Post-Caspofungin Exposure*

Previous work demonstrated that the *A. fumigatus* Δ*aspB* and Δ*aspC* strain had increased

sensitivity to caspofungin on solid media using single-spot inoculation [24]. Although

informative, these assays provide a minimal understanding of the effect of caspofungin in the

septin deletion strains. In order to further characterize the role septins in *A. fumigatus* response

to echinocandins, we conducted a spore dilution assay (10^4-10^1 conidia, 1 μg/mL caspofungin),

minimum effective concentration assay (MEC), and E-strip test (10^6 conidia) to visualize the

susceptibility to caspofungin. Deletion of the core septin genes, *aspA, aspB*, and *aspC*, resulted

in increased sensitivity to caspofungin compared to the *akuB*[KU80] wild-type strain, Δ*aspB::aspB*

complemented strain, and other septin deletion strains in these assays (Fig. 1A, Supplementary

Table 4). Similarly, the Δ*aspA*, Δ*aspB*, and Δ*aspC* strains had a clearer zone of effect on the E-

strip test (Fig. 1D). To explore whether this effect was exclusive to caspofungin, we conducted

an E-strip test to visualize the susceptibility to another drug in the echinocandin class,

micafungin. The Δ*aspA and* Δ*aspC* strains showed a slight increase in susceptibility to

micafungin, as seen by a clearer zone of effect, while Δ*aspB* had the most noticeable decrease

in growth in the zone of effect (Fig. 1D). Thus, this increase in susceptibility of the Δ*aspA,*

Δ*aspB*, and Δ*aspC* strains is not limited to caspofungin, albeit it is more pronounced.

AspB is Required for Fungistatic Response to Caspofungin in vitro

Since we observed an increase in susceptibility in all three of our drug susceptibility assays, we

hypothesized that the absence of AspA, AspB, and AspC leads to a fungicidal effect of

caspofungin (Fig. 1). To determine the viability of the deletion strains during caspofungin

exposure, we utilized CFDA (5,(6)-Carboxyfluorescein Diacetate). When CFDA is cleaved by

esterases, its product is fluorescent, allowing us to use it as a way to distinguish between viable

and non-viable cells [31]. In basal conditions, 10^4 conidia were grown on coverslips in GMM+UU

(glucose minimal media + uracil and uridine) liquid media for 24 hours, followed by 1 hour

incubation in 1 µg/mL CFDA. Uracil and uridine are added to the media due to Δ*aspD* being

auxotrophic. All strains had equally viable confluent growth, indicating that the deletion of the

septin does not impact mycelial viability(Fig. 2A). Strains were then grown for 48 hours in the

presence of 1 µg/mL caspofungin, followed by incubation with CFDA. We then quantified the

number of viable and total microcolonies formed in the presence of caspofungin. The *aspA,*

aspB, and *aspC* deletion strains were significantly less viable (p<0.0001) after 48 hours of

caspofungin exposure, with only 0.92% of Δ*aspB* (p<0.0001) being viable (Fig. 1B, 2B). Δ*aspB*

also does not form many microcolonies in the presence of caspofungin. Δ*aspA* and Δ*aspC* form

microcolonies in the presence of caspofungin; however, only 16.3% and 18.5% of the colonies,

respectively, were viable at the time of microscopic examination ($p<0.0001$)(Fig. 1B, 2B). We

repeated this assay with the echinocandin micafungin in order to test if this reduction in viability

was specific to caspofungin or a general response to echinocandin exposure. In contrast to

caspofungin treatment, the majority of colonies from all strains were viable after micafungin

treatment (*akuB*^{KU80}=95.3%, Δ*aspA*=82.1%, Δ*aspB*=67.4%, Δ*aspC*=79.9%, Δ*aspD*=91.4%,

Δ*aspE*=89.8%, Δ*aspB::aspB*=92.2% viable colonies)(Fig. 1C). This indicates that the fungicidal

effect in the Δ*aspB* strain is specific to caspofungin.

AspB is Involved in Response to Caspofungin in Mature Mycelium

It is possible that most of these phenotypes that we observe are due to caspofungin acting as

the spore germinates, as exemplified by the small microcolonies in the Δ*aspB* strain. Mature

mycelia would be a more clinically relevant growth stage, and for this reason, we decided to

determine the effect of caspofungin on mature mycelia. To test this, we used propidium iodide

(PI) stain to determine hyphal damage. Strains were grown in GMM+UU for 24 hours at 37°C

before being stained with PI and visualized. No hyphal damage was seen under basal

conditions (Fig. 3A). Following the same procedure, we grew strains then incubated them with

caspofungin (1 µg/ml) for 2 hours. Coverslips were washed and stained as before, then

visualized. Similar to our viability assays, only the Δ*aspB* strain exhibits extensive hyphal

damage after 2 hours of exposure to caspofungin (Fig. 3B). Taken together, these results

suggest that AspB is needed for fungal response to caspofungin.

Deletion of* aspB *improves caspofungin treatment efficiency in a neutropenic murine

model of invasive aspergillosis

As we observed a strong fungicidal effect against the Δ*aspB,* we decided to determine if

deletion of *aspB* led to improved survival in our murine model of invasive aspergillosis. As the

Δ*aspB* strain has been previously shown to have no discernible difference in virulence

compared to the *akuB*[KU80] and Δ*aspB::aspB* strains with respect to fungal burden and lung

inflammation, any effect on survival would be attributed to the role that AspB plays in fungal

response to caspofungin [24]. To test this hypothesis, neutropenia was induced in 6-week old

male CD-1 mice using 175 mg/kg cyclophosphamide and 40 mg/kg triamcinolone acetonide.

Neutropenic mice were then intranasally infected with 4×10^6 conidia of *akuB*[KU80], Δ*aspB*, and

Δ*aspB::aspB* strains. They were subsequently treated with either 2 mg/kg caspofungin or an

equivalent volume of the saline vehicle. All groups of mice treated with saline all had the first

death on day 3 post-infection (Fig. 4A) and were not statistically significantly different. Mice

inoculated with Δ*aspB* strain and treated with caspofungin had a 70% survival rate

($p<0.0001$)(Fig. 4B). Additionally, in this group the first death did not occur until day 12 post-

infection. Compared with the next highest groups, the Δ*aspB* strain treated with saline and

akuB[KU80] treated with caspofungin had 30% survival (Fig. 4A). Lung histology was then

performed to visualize inflammation and fungal lesions. Concurrent with the survival graph,

animals infected with the Δ*aspB* strain treated with caspofungin had stunted fungal growth

compared to animals from the other groups, while inflammation around fungal cells was similar

across conditions (Fig. 4D). Previous infection models also showed no difference in

inflammation or fungal lesions between untreated *akuB*[KU80], Δ*aspB*, and Δ*aspB::aspB* strains

[24]. This suggests the caspofungin treatment reduces growth and potentially has a fungicidal

effect on the Δ*aspB* strain *in vivo.*

AspB Interactome Changes Post Caspofungin Exposure

Previous work determined that septins' localization is altered by exposure to caspofungin in *C.*

albicans and *A. fumigatus* [24,29,30]. Additionally, the protein interactome of septin AspB was

altered after exposure to caspofungin in a qualitative proteomic experiment [32]. This approach

only detected for the presence or absence of interactant proteins between basal and

caspofungin conditions, potentially missing interactions that occur in both conditions but change

in abundance [32]. To gain a more mechanistic insight into how AspB contributes to the fungal

response to caspofungin, we applied affinity purification coupled with quantitative proteomics.

An AspB-eGFP expressing strain was grown in GMM and GMM supplemented with 1 µg/mL

caspofungin for 24 hours. AspB-GFP was then purified using a GFP-Trap® affinity matrix, and

proteins bound to AspB-GFP were prepared and analyzed by LC-MS/MS to identify protein

interactors in each condition. Principal Component Analysis (PCA) indicates that the GMM- and

caspofungin-AspB interactome are distinctive from each other (Fig. S1A). A total of 226 proteins

were significantly decreased (fold change (FC) < -2, $p < 0.05$) and 106 proteins were

significantly increased (FC > 2, $p < 0.05$) upon caspofungin treatment (Fig. S1B). Among the

proteins increased upon caspofungin exposure was PpoA, a characterized fatty acid

monooxygenase. PpoA was also identified by a previous study investigating the AspB

interactome post-caspofungin exposure, indicating that our analyses were able to confirm

previously described caspofungin-specific AspB interactions [32]. Candidate protein interactors

that met the screening criteria (FC > 2 or FC<-2, $p < 0.05$) were analyzed using FungiFun3, a

gene ontology tool, to assess changes in biological processes [33]. Proteins that are known to

be involved in hyphal growth and cell wall-related processes were overrepresented among the

list of proteins increased upon caspofungin treatment (Fig. S1C). In contrast, proteins that are

known to be involved in protein folding were overrepresented in the list of proteins decreased

upon caspofungin treatment (Fig. S1D).

Δbgt1, Δgel2, ΔnsdD*, and *ΔmapA *Have a Reduction in Conidiation, While *ΔnsdD *Has a Growth Defect*

Based on our proteomics and gene ontology analyses, we narrowed down to six genes that met the following criteria for further investigation: were significantly increased after caspofungin exposure ($p < 0.05$), were at least two-fold greater in abundance ($FC > 2$), and had either a known or putative role in cell-wall related functions (Table S3). We performed the individual deletions of all six candidate genes (Table S3) to determine their role in septin-related phenotypes. We first characterized each deletion strain to understand how the gene affects growth during basal conditions. Due to the Δ*punA* not being viable at 37 °C, we decided not to pursue it further in this report. Major and biologically relevant radial growth defects are seen in Δ*nsdD* (37.8 mm average diameter compared to the 85 mm diameter of the *akuB*^{KU80} strain)(Fig. 5A). The slight reduction of radial growth in Δ*aspB* was also seen in a previous study and deemed unlikely to be biologically relevant in basal conditions, which is also the case for the Δ*gel2* strain [24]. Similarly, a previously noted defect in conidiation was seen in Δ*aspB* [24]. Δ*bgt1, *Δ*gel2, *Δ*nsdD*, and Δ*mapA* also have reduced conidiation, but not as severely as Δ*aspB* (Fig. 5B). As prior work noted that Δ*aspB* strain had delayed septation, we were interested in determining if any of the candidate gene deletion strains were also defective in septation [24]. To test this, we measured apical compartment length as an indirect method of measuring potential septation defects. Δ*bgt1* mutants have a similar increase in apical compartment length to Δ*aspB*, suggesting a delay in septation (Fig. 5C). In contrast, Δ*gel1* mutants have apical compartments similar to the *akuB*^{KU80} wild-type strain (Fig. 5C), suggesting that Gel1 is dispensable for septa formation. Δ*gel2* (70.4 μm) and Δ*mapA* (67.7 μm) mutants exhibit hyperseptation. Δ*nsdD* (37.7 μm) has a more drastic reduction in apical compartment length, but this phenotype may be due to its growth defect (Fig. 5C)[34].

Δbgt1, Δgel2, ΔnsdD*, and *ΔmapA *Have Increased Sensitivity to Caspofungin Exposure*

Since the candidate proteins showed increased interaction with AspB during caspofungin

exposure, we explored their potential role in mediating the AspB-dependent fungal response to

caspofungin. To test this, we conducted a spore dilution assay on GMM agar supplemented with

1 µg/mL caspofungin. 10^4, 10^3, 10^2, and 10^1 spores were plated and incubated at 37°C for 2

days. Deletion of the *aspB* gene displayed colony-level growth defects in the 10^3, 10^2, and 10^1

concentrations (Fig. 5D). No other deletion strains displayed as severe of a sensitivity to

caspofungin. Δ*bgt1,* Δ*gel2,* Δ*nsdD*, and Δ*mapA* lost full colony growth at 10^2 and 10^1

concentrations (Fig. 5D). In contrast, Δ*gel1* is similar to the *akuB*^KU80 wild-type and Δ*aspB::aspB*

complemented strains (Fig. 5D). The minimum effective concentration (MEC) of caspofungin for

Δ*bgt1,* Δ*gel2,* Δ*nsdD*, and Δ*mapA* was determined to be lower than that of the wild-type, and

equal to or lower than that of Δ*aspA*, Δ*aspB,* and Δ*aspC* (Table S4).

Reduced Viability of* Δbgt1, Δgel2, ΔnsdD*, and *ΔmapA *Post-Caspofungin Exposure

Since *aspB* deletion leads to a fungicidal response to caspofungin, we investigated whether any

candidate gene deletion strains also exhibited reduced viability. All strains showed equal

viability in basal conditions, confirmed qualitatively by visible fluorescence present throughout

the mycelium (Fig. 6A). We then grew all strains for 48 hours in the presence of 1 µg/mL

caspofungin and determined their viability with CFDA. The *aspB* deletion strain showed nearly

no viability at 37°C, as previously shown in Figure 1B (Fig. 5E, Fig. 6B). No other strain

demonstrated the complete loss of viability phenotype observed in the Δ*aspB*. The Δ*bgt1*,

Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains form microcolonies in the presence of caspofungin like the

akuB^KU80 strain, but they had a reduction in viability (Δ*bgt1*=37.4%, Δ*gel2*=39.5%,

Δ*nsdD*=27.8%, Δ*mapA*=26.1% compared to *akuB*^KU80 =90.4%)(Fig 5E). Δ*gel1* did not have a

293 reduction in cell viability when grown in caspofungin (Δ*gel1*=90.8%) (Fig. 5E, Fig. 6B). To

294 determine if any of the candidate gene deletion strains had a unique phenotype to other

295 echinocandins, we conducted the same assay but with micafungin, similar to Figure 1C. The

296 candidate gene deletion strains do not have a high reduction of viability when grown in

297 micafungin (Fig. 5F, Fig. S2).

Bgt1, Gel2, NsdD, and MapA are Involved in the Response to Caspofungin in Mature

Mycelium

301 Next, we determined if any of our candidate gene deletion strains were susceptible to hyphal

302 damage when exposed to caspofungin as mature mycelia. To visualize hyphal damage, we

303 utilized a propidium iodide (PI) stain. No hyphal damage was observed in basal conditions,

304 indicating that there is no defect in the cell wall at basal conditions (Fig. 6C). The experiment

305 was then repeated with a 2-hour incubation in GMM supplemented with 1 µg/mL caspofungin

306 prior to PI staining. Extensive hyphal damage was seen in Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA*

307 strains, similar to Δ*aspB,* after exposure to caspofungin (Fig. 6D). Taken together, Bgt1, Gel2,

308 NsdD, and MapA are involved in the fungal response to caspofungin in earlier stages of growth

309 and mature hyphae.

Candidate Genes are not involved in the response to other Cell Wall Disrupting Agents or

in the Caspofungin Paradoxical Effect

313 We further characterized the candidate gene deletion strains' sensitivity to other cell wall-

314 disturbing agents. To test this, we plated conidia on GMM agar and GMM supplemented with

315 either 1 µg/mL caspofungin, 100 µg/mL Congo red, 2 µg/mL nikkomycin Z, 5 µg/mL calcofluor

316 white, or 10 µg/mL calcofluor white and incubated for three days. Additionally, we plated 4

μg/mL caspofungin and incubated for five days to observe whether the deletion strains were still

capable of the caspofungin paradoxical effect. This paradoxical effect, sometimes known as the

Eagle phenomenon, is when increasing concentrations of a drug become less effective. After

incubation, Δ*aspB* shows increased susceptibility to caspofungin and Congo red, and slight

increase in susceptibility to nikkomycin Z and calcofluor white (Fig. S3). We observed Δ*gel1* had

a mild increase in susceptibility to Congo red compared to the *akuB*^KU80 wild-type (Fig. S3).

Δ*bgt1*, Δ*gel2*, and Δ*nsdD* had a mild increase in susceptibility to 10 μg/mL calcofluor white, but

not as pronounced as Δ*aspB* (Fig. S3). To avoid missing potential phenotypes, we also

conducted spot dilution assays with these drugs (Fig. S4). We again observed a slight increase

in susceptibility of Δ*bgt1*, Δ*gel2*, and Δ*nsdD* to calcofluor white, but no novel phenotypes were

seen (Fig. S4). Taken together, these candidate genes might mediate fungal response to

caspofungin while not being a part of a general fungal cell wall stress response.

Candidate genes are involved in the fungal response to caspofungin through the same

mechanisms as AspB

To further confirm that AspB is functioning in the same pathways as the candidate proteins, we

generated double deletion mutants that lack both the *aspB* and one of the genes that previously

had a phenotype to caspofungin (*bgt1*, *gel2*, *nsdD*, and *mapA*). As these strains appeared

slightly more sickly, we decided to verify that they do not have a phenotype in basal conditions

that could largely affect the interpretation of drug assays. We conducted radial growth, conidial

quantification, and spore viability assays. The Δ*aspB*Δ*nsdD* strain had a similar growth defect to

Δ*nsdD* (Fig. 7A). The other double mutants have typical radial growth, resembling either Δ*aspB*

or their other parent (Fig. 7A). All double mutant strains had reduced conidia production, similar

to their Δ*aspB* parent, or both Δ*aspB* and Δ*mapA* in the case of the Δ*aspB*Δ*mapA* strain (Fig.

7B). Spores were tested for viability by staining 1×10^7 conidia of each strain with propidium

iodide for inviability and calcofluor white as confirmation of being a spore. Heat-killed wild-type

spores were used to standardize images. All strains had viable spores, albeit all mutants had

slightly more inviable spores than the wild-type (Fig. 7C). To test for mycelial viability, a CFDA

assay was conducted, similar to that of Figures 2 and 6. All strains were viable under basal

conditions. Strains were then grown for 48 hours in the presence of 1 µg/mL caspofungin and

we determined their viability with CFDA. All double deletion strains had similar viability reduction

as the *aspB* deletion strain (Fig. 7D). To test if there is increased hyphal damage in mature

mycelium, we used propidium iodide staining on mycelium grown for 24 hours. Little to no

fluorescence was observed, suggesting no major defect in the cell wall under basal conditions

(Fig. S5A). This experiment was repeated with a 2-hour incubation in GMM supplemented with

1 µg/mL caspofungin, followed by PI staining. Similar to both Δ*aspB* and their other parent, all

double deletion strains had increased hyphal damage post-caspofungin exposure (Fig. S5B).

Taken together, these data suggest that AspB contributes to the candidate proteins role in

response to caspofungin.

Discussion

The septins are a highly conserved family of cytoskeletal proteins with a variety of cellular roles.

This includes roles in cell division, stress response, cytoskeleton organization, and scaffolding

[24,35–37]. Previous work noted the role of septins in response to cell wall stress. In *A.*

fumigatus, it was determined that the septins have a role in response to cell wall stress, notably

showing that Δ*aspB* exhibits a hypersensitive response to caspofungin exposure on solid media

[24]. Here, we aimed to further investigate the role of septins in the *A. fumigatus* response to

caspofungin. We found that deletion of *aspA, aspB*, and *aspC* reduced fungal viability after

caspofungin exposure and increases hyphal damage in mature mycelia, whereas deletion of

aspB elicited a fungicidal response. The drastic phenotype of the Δ*aspB* strain was surprising,

as *Aspergillus spp.* septins form heteropolymers consisting of hexamers (AspA-C) and octamers

(AspA-D) [38]. Thus, it would be expected that deletion of genes that abolish septin

heteropolymers would have similar phenotype. Deletion of *aspD* still allows proper formation of

hexamers but not octamers. In our work, we found that the *aspD* deletion strain was slightly

more sensitive than the *akuB*^{KU80} wild-type, Δ*aspB::aspB*, or Δ*aspE* strains, although its

increased susceptibility was not as pronounced as that of the *aspA*, *aspB*, and *aspC* deletion

strains (Fig. 1). This suggests that the presence of the hexamer is at least sufficient, if not

necessary, for proper fungal response to caspofungin. It is unknown if the sole presence of the

octamer would also be sufficient, as deletion of *aspA-C* prevents the formation of both

structures. Similarly, Δ*aspB*'s fungicidal response to clinically relevant doses of caspofungin

could be due to AspB being specifically needed to scaffold necessary proteins in response to

caspofungin, regardless of the septin complex, while the Δ*aspA* or Δ*aspC* leads to a loss of both

the hexamer and octamer, which could causes general dysfunction in AspB's role in

coordinating the fungal response to caspofungin. Proteomics analysis in *Cryptococcus*

neoformans also supports the possibility of subunit-specific protein interactions, where the

cdc3^{aspB} and *cdc10*^{aspD} septins are shown to have differences in interactomes after exposure to

cellular stressors [39]. All of the septin deletion strains retain the caspofungin paradoxical effect,

suggesting that individual septins and their heteropolymers act independently of the pathways

that contribute to the caspofungin paradoxical effect.

Septins are involved in septation, as the Δ*aspA*, Δ*aspB*, Δ*aspC*, and Δ*aspE* strains have fewer

septa and a larger apical compartment compared to the *akuB*^{KU80} strain [24]. It has been shown

that septation is a critical cellular process for maintaining the fungistatic effect of echinocandins,

as the deletion of genes involved in the septation initiation network (SIN) results in a fungicidal

effect [31]. The Δ*aspE* strain has a septation defect of similar magnitude to the Δ*aspB* strain,

but the Δ*aspE* strain does not have a drastic increase in susceptibility to the echinocandins.

Additionally, Δ*aspA* and Δ*aspC* have a septation defect, but their decrease in viability post-

caspofungin exposure is not as severe as in the Δ*aspB* strains. Furthermore, deletion of SIN

genes confers fungicidal activity to both micafungin and caspofungin. Thus, the septation defect

could partially contribute to the increased susceptibility to echinocandins in the Δ*aspA,* Δ*aspB,*

and Δ*aspC* strains; however, the septation phenotype alone is not responsible for the fungicidal

effect observed in the Δ*aspB* strain during caspofungin exposure.

The echinocandin class includes four clinically approved drugs: anidulafungin, micafungin,

caspofungin, and most recently approved, rezafungin [40,41]. These drugs all work through

noncompetitive inhibition of the β-1,3-D-glucan synthase, but alterations in their side chains

comprise the major differences between each one [18,42]. These changes in the side chains

can lead to changes in how the drug impacts fungal physiology. For instance, caspofungin is the

only echinocandin to induce the paradoxical effect [42]. Unlike micafungin, high concentrations

of caspofungin increase the levels of cytosolic calcium [43,44]. This activates calmodulin-

calcineurin signaling and leads to the caspofungin paradoxical effect [44]. Thus, our findings

add to the growing evidence that even though the different echinocandins have a shared

mechanism of action, how the fungus responds to echinocandins is drug-specific.

Animals infected with the Δ*aspB* strain and treated with caspofungin had a 70% chance of

survival, which is higher than that of animals infected with *akuB*[KU80] strain (30%). The first death

of Δ*aspB* strain infected mice treated with caspofungin was recorded on day +12 post infection,

compared to Δ*aspB* strain infected mice treated with saline or the other strains and conditions

with first deaths on day +3 or day +4. It is possible that colonies that persist under caspofungin

treatment are capable of growing and establishing infection after the drug pressure is released.

Additionally, while ΔaspB strain infected mice treated with saline still had 30% survival, this is

not statistically significant. Previous work also determined loss of AspB does not impact fungal

virulence and resulted in a similar murine survival rate [24]. Our histological analyses showed

reduced inflammation and reduced fungal lesions in the lungs of ΔaspB infected mice treated

with caspofungin. However, this assay is done on day +3 after infection when mice are still

being treated with caspofungin. Nonetheless, this increase in survival makes AspB a

prospective target for drug therapy developments, as use in conjunction with caspofungin elicits

a fungicidal response in *A. fumigatus*. Currently, only the plant cytokinin forchlorfenuron (FCF) is

known to disrupt septin organization [45]. This compound disrupts septins within both fungal and

mammalian cells by interfering with their ability to bind and hydrolyze GTP [45–47]. It is possible

that FCF could be modified and refined to target fungal-specific septins. FCF analogues have

previously been developed to better target ovarian and endometrial cancers [48]. Thus, FCF

analogues could be used in conjunction with caspofungin to create a new fungicidal therapy for

the treatment of IA. Further work is required to better understand the mechanisms of AspB's

response to caspofungin and to develop novel clinical therapies.

Our proteomic analysis identified four candidate protein interactors of septin AspB that were

significantly increased by caspofungin treatment and have a potential role in cell wall function.

One limitation of our experiment is the lack of an input control, such as GFP ectopically

expressed. Because of the lack of a GFP-only control, some hits may be nonspecific GFP-

interactants. We verified that our candidate genes were not increased in expression after

caspofungin exposure based on publicly available datasets [49]. *bgt1*, *gel2*, *nsdD*, and *mapA* all

decrease in transcript abundance after caspofungin exposure, while *gel1* and *punA* increase in

transcript abundance after caspofungin exposure [49]. This suggests that the increase in *bgt1*,

gel2, *nsdD*, and *mapA* interactions observed in our AspB pulldown experiment might not be due

to an increase in overall abundance. Additionally, to mitigate the possible false positives that

could be obtained from this process, we performed epistasis analyses and generated double

mutants of our candidate and *aspB,* in which candidate genes were epistatic to *aspB*.

Bgt1, Gel1, and Gel2 are glucanosyltransferases in *A. fumigatus* [50]. These

glucanosyltransferases work to remodel unorganized β-1,3-glucan chains in the periplasmic

space to stabilize and modify the cell wall as needed [50]. Due to their role in cell wall

homeostasis, previous work identified these three glucanosyltransferases as potential drug

targets that required further investigation [50]. Bgt1 works by hydrolyzing a β-1,3-linked

oligosaccharide and placing it on another molecule of β-1,3-glucan [51]. In *C. albicans*, deletion

of *BGL2*, the *bgt1* homologue, resulted in an increase in susceptibility to nikkomycin Z, a chitin

synthase inhibitor [52]. In contrast, *A. fumigatus* Δ*bgt1* only has a slight increase in susceptibility

to nikkomycin Z (Fig. S3). Deletion of *BGL2* in *S. cerevisiae* increased the cellular chitin content

[53]. In *A. fumigatus,* though, Δ*bgt1*Δ*bgt2* double mutants do not have a change in chitin content

[51]. A slight reduction in conidiation was observed in *A. fumigatus* Δ*bgt1* (Fig. 5A, 5B). *A.*

fumigatus Δ*bgt1* susceptibility to calcofluor white and Congo red were similar to the *akuB*[KU80]

wild-type, although compared to basal conditions there was a reduction in growth (Fig. S3) [54].

A potential mechanism for the increased susceptibility in *A. fumigatus* Δ*bgt1* and Δ*aspB* to

caspofungin is that hyphal damage may be exacerbated by the lack of interaction between

AspB and Bgt1 when either party is missing (Fig. 5E). Due to the role of septins in septation and

scaffolding, the loss of AspB in particular may be more deleterious than just the loss of Bgt1 and

thus result in the observed fungicidal effect (Fig. 1B). Gel1 and Gel2 are two GPI-anchored

glucanosyltransferase members of the Glycoside Hydrolase Family 72 (GH72) [55]. Despite

having the same enzymatic activity, deletion of *gel1* did not result in a unique phenotype, but

reduced mycelial growth and abnormal cell wall architecture was observed in Δ*gel2* during basal

conditions (Fig. 5A-B) [50,51,56]. An increase in chitin is associated with deletion of *gel2*, a

compensatory mechanism similar to deletion of *bgt1* [56]. *A. fumigatus* Gel1 is dispensable for

proper fungal response to caspofungin (Fig. 5D-E). In contrast, Gel2 does appear to be involved

in the response to caspofungin due to increased susceptibility to caspofungin and reduced

viability in caspofungin compared to *akuB*^{KU80} wild-type (Fig. 5D-E)[56]. Our proteomics analysis

noted an increased interaction between Gel2 and AspB, but cellular levels of Gel2 were not

explored. It is unknown if cellular levels of Gel2 are increased in *A. fumigatus* post-caspofungin

exposure, similar to *C. albicans*. Additionally, further work exploring the overexpression of Gel2

may uncover a new mechanism of resistance to caspofungin in *A. fumigatus*. Further work is

also needed to uncover the relationship between glucanosyltransferase and the septins in

septation and cell wall stress responses.

NsdD (**n**ever in **s**exual **d**evelopment) is a putative GATA-type transcriptional activator with roles

in *A. nidulans* for sexual development [57]. NsdD is responsible for activating sexual

development [58]. Deletion of *nsdD* results in the inability to produce cleistothecia (fruiting

bodies), while overexpression allows production of Hülle cells even in conditions which typically

block sexual development [58]. Despite being a primarily asexual fungus, *A. fumigatus* is

capable of and still holds a variety of functional sexual reproductive genes [59]. NsdD has been

implicated in cell wall remodeling and hyphal fusion [60]. Deletion of *nsdD* results in reduced

hyphal growth on minimal media (Fig. 5B) [57,60]. Drug challenges show some sensitivity of

ΔnsdD to Congo red [60] and caspofungin (Fig. 5D-E, Fig. S3). Previous work also noted

weakened hyphal tips in Δ*nsdD*, which is in accordance with the increased hyphal damage

during cell wall stress seen via PI staining (Fig. 6D)[60]. We suspect that while NsdD may have

independent roles in sexual development, its previous cell wall remodeling functions during

hyphal fusion for heterokaryon formation may have alternative roles in *A. fumigatus* cell wall

remodeling outside of sexual development. Further work is required to understand the NsdD-

mediated transcriptional response to caspofungin exposure.

MapA (gene *Afu4g06930*) is a previously uncharacterized protein in *A. fumigatus*. In *S.*

cerevisiae and *C. albicans*, cytosolic protein MAP2 is an ortholog that functions as a methionine

aminopeptidase. Deletion of MAP2 results in slightly slower growth rate and some chemical

sensitivities in *S. cerevisiae* and *C. albicans* [61,62]. It is possible that this methionine

aminopeptidase can remove the methionine to promote the maturation of proteins involved in

the response to caspofungin. However, which proteins are regulated by MapA in *A. fumigatus* is

unknown.

Although septins have been implicated in the mediation of cell wall stress, the full mechanism

how they contribute to this phenotype is not fully understood [23,24,28,29,32]. Previous

proteomic analyses and our own proteomics experiments show that MpkA, MkkA, and other cell

wall integrity components co-immunoprecipitated with AspB [32]. It is possible that AspB

interacts with the cell wall integrity pathway to facilitate fungal response to caspofungin [32]. In

A. nidulans, deletion of *aspB* partially rescues both the growth defect and susceptibility to

caspofungin of the Δ*mpkA* strain, further hinting at a possible crosstalk between the septin and

the CWI [23]. As we did not observe a difference in abundance in the CWI pathways kinases

between basal and caspofungin conditions, we did not explore the possible role of septins in

regulating the CWI. Nonetheless, further work will be needed to investigate this possibility. As

we see different pathways differentially interacting with AspB, we hypothesize that the fungicidal

response observed in the Δ*aspB* strain is not due to a singular AspB-mediated interaction that is

yet to be found, but instead by AspB having a global effect on a number of contributing proteins

and pathways. Future studies directed towards developing drugs specific to fungal septins or in

determining additional septin interactants that have key roles in the caspofungin response can

help to better understand ways in which caspofungin therapy can be improved for use against

invasive aspergillosis.

Materials and Methods

Strains, media, and conditions

A. fumigatus akuB[KU80] served as the wild-type and control strain. Δ*aspB::aspB* complemented

strain served as an additional control to determine that phenotypes were due to the deletion of

the *aspB* gene. Septin deletion strains are described in (Table S1, [24,63]). The *aspB-egfp*

strain from [24], with *aspB-egfp* expressed by the *aspB* promoter from the native loci, was used

to pulldown AspB interactants. All cultures were plated on glucose minimal media (GMM)

supplemented with 5 mM uracil and 5 mM uridine (GMM+UU) and incubated at 37°C, unless

otherwise specified.

Drug susceptibility assays

Spores were counted, diluted, and inoculated in rows onto agar plates containing caspofungin

(1 μg/mL) in 10μL quantities with a total of 10^4, 10^3, 10^2, and 10^1 conidia. Plates were incubated

for 48 hours at 37°C unless otherwise noted. For the minimum effective concentration assay, a

quantity of 2.5×10^4 spores were added to RPMI media with different concentrations of

caspofungin and incubated for 48 hours at 37°C according to CLSI guidelines [64]. We were not

able to test the *aspD* and *punA* deletion strains according to CLSI guidelines as Δ*aspD*, which is

an uracil/uridine auxotroph, could not grow in RPMI, and Δ*punA*, which has a temperature

sensitivity, could not grow at 37°C.

Fungal Viability Assay

Conidia of *akuB*[KU80], Δ*aspA-E*, and Δ*aspB::aspB* were diluted to 10^4 spores and cultured on

coverslips immersed in 4 mL of GMM+UU broth and GMM+UU+caspofungin (1 µg/ml) and

incubated at 37°C for 24 and 48 hours, respectively. To examine cell viability, coverslips were

incubated in 5-carboxyfluorescein diacetate (CFDA) (50 µg/mL 0.1 M MOPS pH 3) for 1 hour at

37°C and 70 rpm [31]. Slides were prepared for microscopy and image on an Inverted Leica

DMi8 with Leica K5 Microscope Camera using a 10x objective. Images were analyzed using

imageJ [65]. Viability in basal conditions (GMM) was determined qualitatively by noting the

presence of visible fluorescence throughout the mycelium. To quantitatively determine viability

when grown in caspofungin, viable, fluorescent colonies were manually counted and divided

over total colonies counted. A minimum of 50 colonies per strain per replicate were counted.

Spore Inviability Assay

Conidia of *akuB*[KU80], heat-killed *akuB*[KU80] (5 minutes at 95°C), Δ*aspB*, and the single and double

deletion mutants were diluted to 10^7 spores and spun down via centrifuge for 5 minutes at

maximum speed. Their media was removed and replaced with PIPES (50 mM PIPES, pH 6.7),

then vortexed to dislodge the pellet. Spores were washed a total of two times in this manner,

then vortexed and incubated with calcofluor white (1 µg/mL) and propidium iodide (12.5 µg/mL)

for 5 minutes at room temperature. Spores were spun down and washed with PIPES two more

times. Slides were prepared for microscopy and imaged on Nikon Ti-Eclipse microscope using

a 100x objective. Images were analyzed using imageJ [65]. Inviability was determined by

standardizing against the heat-killed sample, then manually counted for the number of inviable

spores and divided over total spores counted [66]. A minimum of 50 spores per strain per

replicate were counted.

Hyphal Damage Assay

Conidia (10^4) of *akuB*[KU80], Δ*aspA-E*, and Δ*aspB::aspB* were cultured on coverslips immersed in

4 mL of GMM+UU broth and incubated for 24 hours at 37°C. Coverslips were then incubated

with GMM+UU or GMM+UU+caspofungin (1 µg/ml) for 2 hours at 37°C. To examine the

damage of mature hyphae, coverslips were washed with 4 mL PIPES (pH 6.7) for 5 minutes.

PIPES was then removed and washed again with 4 mL PIPES for 5 minutes. After removing the

second PIPES wash, 500 uL of propidium iodide (PI) solution (12.5 ug/mL in 50 mM PIPES)

was added on top of the coverslip and incubated in the dark for 5 minutes. Slides were washed

in 4 mL PIPES twice as described previously. They were then prepared for microscopy and

imaged on an Inverted Leica DMi8 with Leica K5 Microscope Camera using a 10x objective.

Images were analyzed in imageJ [65].

Neutropenic Murine Model of Invasive Aspergillosis

Murine experiments followed previously established intranasal neutropenic models of IA [67,68].

Sixty 6-week old male CD1 mice (Charles River Laboratories, Raleigh, NC) were injected via

intraperitoneal route with 175 mg/kg cyclophosphamide on days −2 and +3 and 40mg/kg

triamcinolone acetonide subcutaneously on days −1 and + 6. Twenty mice per strain were

infected intranasally with 40 µL of 10^8 spores/mL conidia suspension of the *akuB*[KU80], Δ*aspB*, or

Δ*aspB::aspB* strains on day +0. On days +1 through +4, mice were injected via intraperitoneal

route with either 2 mg/kg caspofungin or saline. Mice were monitored until day +14 and

humanely euthanized via CO_2 followed by cervical dislocation if they showed severe symptoms.

Survival was plotted on a Kaplan-Meier curve with a log rank pair-wise comparison. Murine

experiments were conducted in compliance with the SIU Institutional Animal Care and Use

Committee Protocol 20-034 and Virginia Tech Protocol 23-254.

Histopathology Analysis

Mice were immunocompromised and treated as described above in [Section: *Neutropenic*

murine model of Invasive Aspergillosis]. Lungs were harvested on day +3 after infection via

isoflurane euthanasia and cervical dislocation. Tissue sections were stained using hematoxylin

and eosin (H&E) stains to visualize inflammation and Gomori's methenamine silver stain to

visualize fungal hyphae.

Protein Extraction, AspB-eGFP Fusion Protein Purification, and LC-MS/MS Analysis

The *aspB-egfp* strain from [24] was grown in GMM liquid media and GMM liquid media

supplemented with 1 µg/mL of caspofungin for 24 hours at 37°C. Protein extraction and

pulldown were completed as described by [32,69]. Fungal mycelia were homogenized to obtain

total cell lysate and AspB complexes were purified using the GFP-Trap® affinity purification

(Chromotek), as described [69]. GFP-Trap® magnetic beads were equilibrated by washing

beads three times in 500 µL ice-cold dilution buffer then resuspended in 100 µL ice cold dilution

buffer. The resin suspension is then mixed with total cell lysate (10 mg total protein) and

incubated at 4°C with gentle agitation for 2 hours. Beads were collected using a magnetic stand.

Beads were washed in 500 µL ice-cold lysis buffer and five times with 500 µL of wash buffer.

Beads were suspended in 50 µL wash buffer.

Samples were digested on-bead with trypsin followed by C18 desalting. Samples were analyzed

via LC-MS/MS on a Thermo Easy nLC 1200-QExactive HF in technical duplicate. All mass

spectra data was processed using MaxQuant (ver. 1.6.12.0) and searched against the Uniprot

Aspergillus fumigatus proteome reviewed database (Proteome ID UP000002530). MaxQuant

output was further processed via Perseus with filtering at 1% false discovery rate (FDR). Only

proteins with >1 peptide were reported.

Prioritization of Candidate Genes and Generation of Deletion Mutants

Candidate genes were chosen by prioritizing proteins with at least a two-fold increase, statistical

significance, and known or putative roles in cell wall functions as listed in FungiFun3 GO term

search for biological processes (Table S3) [33]. Deletions of *bgt1* (*Afu1g11460*; fungidb.org),

gel1 (*Afu2g01170*; fungidb.org), *gel2* (*Afu6g11390*; fungidb.org), *nsdD* (*Afu3g13870*;

fungidb.org), *mapA* (*Afu4g06930*; fungidb.org), and *punA* (*Afu6g07470*; fungidb.org) genes

were obtained by replacing the gene with the 2.4 kb *pyrG* gene from *Aspergillus parasiticus*.

Approximately 1 kb of promoter and terminator region of each gene were PCR-amplified from

AF293 genomic DNA. Deletion constructs were generated by overlap fusion PCR and

subsequently transformed into *akuB*^{KU80} *pyrG–* strain, all as previously described by [67].

Primers used in the transformation of all candidate strains are found in Table S2. Transformants

were validated via PCR screening.

Deletion of* aspB *via CRISPR/Cas9-Mediated Transformation

Double deletion and non-*akuB*^{KU80} Δ*aspB* strains were generated by deleting the *aspB* gene in

the Δ*bgt1,* Δ*gel2,* Δ*nsdD,* Δ*mapA,* AF293, and CEA10 backgrounds. CRISPR/Cas9-mediated

transformation was performed, as previously described by [70]. For the homologous repair

template, a 2890 bp hygromycin B resistance (*hph*) cassette from the pUCGH plasmid was PCR

amplified with 35 bp of *aspB* microhomology domains at each end. crRNAs were designed to

target outside the *aspB* gene. The approximately 1.8 kb *aspB* gene was replaced with the total

2960 bp *hph* plus *aspB* microhomology domains in each of the strains. The crRNA sequences

used are in Table S2. Deletion of *aspB* was verified via PCR by screening for the lack of the

aspB gene, total size of the construct, and presence of *hph*. Primers used are in Table S2.

Acknowledgments

R.J.B. was supported by the Southern Illinois University Carbondale Master's Fellowship and

Virginia Tech GRAD Fellowship. R.J.B. and J.V.M. were supported in part by 1R01AI165656-

01A1. This work was supported in part by the College of Science at Virginia Tech (Fund BR-

COS1357) and made possible by funding from the Fralin Life Sciences Institute by FLSI Award

260027. We also want to thank the members of the Vargas-Muñiz group for their critical reading

of the manuscript. We want to thank Stacey Mcgee (SIU histology core facility) and Dr. Jennifer

Harris (SIU Laboratory Animal Program Director) for providing guidance and support during

animal experiments.

References:

1. Bongomin F, Gago S, Oladele R, Denning D. Global and Multi-National Prevalence of Fungal Diseases—Estimate Precision. Journal of Fungi. 2017;3: 57. doi:10.3390/jof3040057

2. Latgé J-P, Chamilos G. Aspergillus fumigatus and Aspergillosis in 2019. Clin Microbiol Rev. 2019;33. doi:10.1128/CMR.00140-18

3. Rybak JM, Fortwendel JR, Rogers PD. Emerging threat of triazole-resistant Aspergillus fumigatus. Journal of Antimicrobial Chemotherapy. 2019;74: 835–842. doi:10.1093/jac/dky517

657   4.   Li X, Lau SK, Woo PC. Fungal infection risks associated with the use of cytokine antagonists and
658   immune checkpoint inhibitors. Exp Biol Med. 2020;245: 1104–1114.
659   doi:10.1177/1535370220939862

660   5.   Baddley JW. Clinical risk factors for invasive aspergillosis. Med Mycol. 2011;49: S7–S12.
661   doi:10.3109/13693786.2010.505204

662   6.   Gouzien L, Cocherie T, Eloy O, Legriel S, Bedos J-P, Simon C, et al. Invasive Aspergillosis associated
663   with Covid-19: A word of caution. Infect Dis Now. 2021;51: 383–386.
664   doi:10.1016/j.idnow.2020.12.008

665   7.   Benedict K, Jackson BR, Chiller T, Beer KD. Estimation of Direct Healthcare Costs of Fungal
666   Diseases in the United States. Clinical Infectious Diseases. 2019;68: 1791–1797.
667   doi:10.1093/cid/ciy776

668   8.   Dagenais TRT, Keller NP. Pathogenesis of Aspergillus fumigatus in Invasive Aspergillosis. Clin
669   Microbiol Rev. 2009;22: 447–465. doi:10.1128/CMR.00055-08

670   9.   Shlezinger N, Irmer H, Dhingra S, Beattie SR, Cramer RA, Braus GH, et al. Sterilizing immunity in
671   the lung relies on targeting fungal apoptosis-like programmed cell death. Science (1979).
672   2017;357: 1037–1041. doi:10.1126/science.aan0365

673   10.   Latgé J-P. Aspergillus fumigatus and Aspergillosis. Clin Microbiol Rev. 1999;12: 310–350.
674   doi:10.1128/CMR.12.2.310

675   11.   Morgan J, Wannemuehler KA, Marr KA, Hadley S, Kontoyiannis DP, Walsh TJ, et al. Incidence of
676   invasive aspergillosis following hematopoietic stem cell and solid organ transplantation: interim
677   results of a prospective multicenter surveillance program. Med Mycol. 2005;43: 49–58.
678   doi:10.1080/13693780400020113

679   12.   Balloy V, Huerre M, Latgé J-P, Chignard M. Differences in Patterns of Infection and Inflammation
680   for Corticosteroid Treatment and Chemotherapy in Experimental Invasive Pulmonary
681   Aspergillosis. Infect Immun. 2005;73: 494–503. doi:10.1128/IAI.73.1.494-503.2005

682   13.   Stergiopoulou T, Meletiadis J, Roilides E, Kleiner DE, Schaufele R, Roden M, et al. Host-Dependent
683   Patterns of Tissue Injury in Invasive Pulmonary Aspergillosis. Am J Clin Pathol. 2007;127: 349–
684   355. doi:10.1309/UJRV9DLC11RM3G8R

685   14.   Coutinho AE, Chapman KE. The anti-inflammatory and immunosuppressive effects of
686   glucocorticoids, recent developments and mechanistic insights. Mol Cell Endocrinol. 2011;335: 2–
687   13. doi:10.1016/j.mce.2010.04.005

688   15.   Cain DW, Cidlowski JA. Immune regulation by glucocorticoids. Nat Rev Immunol. 2017;17: 233–
689   247. doi:10.1038/nri.2017.1

690   16.   Patterson TF, Thompson GR, Denning DW, Fishman JA, Hadley S, Herbrecht R, et al. Practice
691   Guidelines for the Diagnosis and Management of Aspergillosis: 2016 Update by the Infectious
692   Diseases Society of America. Clinical Infectious Diseases. 2016;63: e1–e60.
693   doi:10.1093/cid/ciw326

694 17. World Health Organization. WHO fungal priority pathogens list to guide research, development
695   and public health action. 2022. Available:
696   https://www.who.int/publications/i/item/9789240060241

697 18. Denning DW. Echinocandin antifungal drugs. The Lancet. 2003;362: 1142–1151.
698   doi:10.1016/S0140-6736(03)14472-8

699 19. Dudiuk C, Berrio I, Leonardelli F, Morales-Lopez S, Theill L, Macedo D, et al. Antifungal activity
700   and killing kinetics of anidulafungin, caspofungin and amphotericin B against Candida auris.
701   Journal of Antimicrobial Chemotherapy. 2019;74: 2295–2302. doi:10.1093/jac/dkz178

702 20. Lewis JS, Graybill JR. Fungicidal versus Fungistatic: what's in a word? Expert Opin Pharmacother.
703   2008;9: 927–935. doi:10.1517/14656566.9.6.927

704 21. Graybill JR, Burgess DS, Hardin TC. Key issues concerning fungistatic versus fungicidal drugs.
705   European Journal of Clinical Microbiology & Infectious Diseases. 1997;16: 42–50.
706   doi:10.1007/BF01575120

707 22. Hiemenz JW, Raad II, Maertens JA, Hachem RY, Saah AJ, Sable CA, et al. Efficacy of caspofungin as
708   salvage therapy for invasive aspergillosis compared to standard therapy in a historical cohort.
709   European Journal of Clinical Microbiology & Infectious Diseases. 2010;29: 1387–1394.
710   doi:10.1007/s10096-010-1013-0

711 23. Mela A, Momany M. Septins coordinate cell wall integrity and lipid metabolism in a sphingolipid-
712   dependent process. J Cell Sci. 2022;135. doi:10.1242/jcs.258336

713 24. Vargas-Muñiz JM, Renshaw H, Richards AD, Lamoth F, Soderblom EJ, Moseley MA, et al. The
714   Aspergillus fumigatus septins play pleiotropic roles in septation, conidiation, and cell wall stress,
715   but are dispensable for virulence. Fungal Genetics and Biology. 2015;81: 41–51.
716   doi:10.1016/j.fgb.2015.05.014

717 25. Juvvadi PR, Fortwendel JR, Rogg LE, Steinbach WJ. Differential localization patterns of septins
718   during growth of the human fungal pathogen Aspergillus fumigatus reveal novel functions.
719   Biochem Biophys Res Commun. 2011;405: 238–243. doi:10.1016/j.bbrc.2011.01.017

720 26. Bertin A, McMurray MA, Grob P, Park S-S, Garcia G, Patanwala I, et al. Saccharomyces cerevisiae
721   septins: Supramolecular organization of heterooligomers and the mechanism of filament
722   assembly. Proceedings of the National Academy of Sciences. 2008;105: 8274–8279.
723   doi:10.1073/pnas.0803330105

724 27. Shuman B, Momany M. Septins From Protists to People. Front Cell Dev Biol. 2022;9.
725   doi:10.3389/fcell.2021.824850

726 28. Badrane H, Nguyen MH, Clancy CJ. Highly Dynamic and Specific Phosphatidylinositol 4,5-
727   Bisphosphate, Septin, and Cell Wall Integrity Pathway Responses Correlate with Caspofungin
728   Activity against Candida albicans. Antimicrob Agents Chemother. 2016;60: 3591–3600.
729   doi:10.1128/AAC.02711-15

29. Badrane H, Nguyen MH, Blankenship JR, Cheng S, Hao B, Mitchell AP, et al. Rapid Redistribution
of Phosphatidylinositol-(4,5)-Bisphosphate and Septins during the Candida albicans Response to
Caspofungin. Antimicrob Agents Chemother. 2012;56: 4614–4624. doi:10.1128/AAC.00112-12

30. Badrane H, Nguyen MH, Clancy CJ. Regulation of phosphatidylinositol-(4,5)-bisphosphate and
active-Rho1p levels and distribution is crucial for correct spatio-temporal cytokinesis and
echinocandin responses in Candida albicans. Antimicrob Agents Chemother. 2025;69.
doi:10.1128/aac.01900-24

31. Souza ACO, Martin-Vicente A, Nywening A V., Ge W, Lowes DJ, Peters BM, et al. Loss of Septation
Initiation Network (SIN) kinases blocks tissue invasion and unlocks echinocandin cidal activity
against Aspergillus fumigatus. PLoS Pathog. 2021;17: e1009806.
doi:10.1371/journal.ppat.1009806

32. Vargas-Muñiz JM, Renshaw H, Waitt G, Soderblom EJ, Moseley MA, Palmer JM, et al. Caspofungin
exposure alters the core septin AspB interactome of Aspergillus fumigatus. Biochem Biophys Res
Commun. 2017;485: 221–226. doi:10.1016/j.bbrc.2017.02.116

33. Garcia Lopez A, Albrecht-Eckardt D, Panagiotou G, Schäuble S. FungiFun3: systemic gene set
enrichment analysis for fungal species. Bioinformatics. 2024;40.
doi:10.1093/bioinformatics/btae620

34. Dichtl K, Samantaray S, Aimanianda V, Zhu Z, Prévost M, Latgé J, et al. Aspergillus fumigatus
devoid of cell wall β-1,3-glucan is viable, massively sheds galactomannan and is killed by septum
formation inhibitors. Mol Microbiol. 2015;95: 458–471. doi:10.1111/mmi.12877

35. HARTWELL L. Genetic control of the cell division cycle in yeast *1IV. Genes controlling bud
emergence and cytokinesis. Exp Cell Res. 1971;69: 265–276. doi:10.1016/0014-4827(71)90223-0

36. Gladfelter A. The septin cortex at the yeast mother-bud neck. Curr Opin Microbiol. 2001;4: 681–
689. doi:10.1016/S1369-5274(01)00269-7

37. McMurray MA, Thorner J. Septins: molecular partitioning and the generation of cellular
asymmetry. Cell Div. 2009;4: 18. doi:10.1186/1747-1028-4-18

38. Hernández-Rodríguez Y, Masuo S, Johnson D, Orlando R, Smith A, Couto-Rodriguez M, et al.
Distinct Septin Heteropolymers Co-Exist during Multicellular Development in the Filamentous
Fungus Aspergillus nidulans. PLoS One. 2014;9: e92819. doi:10.1371/journal.pone.0092819

39. Martinez Barrera S, Hatchell E, Byrum SD, Mackintosh SG, Kozubowski L. Quantitative analysis of
septin Cdc10 & Cdc3-associated proteome during stress response in the fungal pathogen
Cryptococcus neoformans. PLoS One. 2024;19: e0313444. doi:10.1371/journal.pone.0313444

40. Ben-Ami R. Systemic Antifungal Therapy for Invasive Pulmonary Infections. Journal of Fungi.
2023;9: 144. doi:10.3390/jof9020144

41. Syed YY. Rezafungin: First Approval. Drugs. 2023;83: 833–840. doi:10.1007/s40265-023-01891-8

42. Perlin DS. Current perspectives on echinocandin class drugs. Future Microbiol. 2011;6: 441–457.
doi:10.2217/fmb.11.19

43.   Fortwendel JR, Juvvadi PR, Perfect BZ, Rogg LE, Perfect JR, Steinbach WJ. Transcriptional
Regulation of Chitin Synthases by Calcineurin Controls Paradoxical Growth of Aspergillus
fumigatus in Response to Caspofungin. Antimicrob Agents Chemother. 2010;54: 1555–1563.
doi:10.1128/AAC.00854-09

44.   Juvvadi PR, Muñoz A, Lamoth F, Soderblom EJ, Moseley MA, Read ND, et al. Calcium-Mediated
Induction of Paradoxical Growth following Caspofungin Treatment Is Associated with Calcineurin
Activation and Phosphorylation in Aspergillus fumigatus. Antimicrob Agents Chemother. 2015;59:
4946–4955. doi:10.1128/AAC.00263-15

45.   Iwase M, Okada S, Oguchi T, Toh-e A. Forchlorfenuron, a phenylurea cytokinin, disturbs septin
organization in Saccharomyces cerevisiae. Genes Genet Syst. 2004;79: 199–206.
doi:10.1266/ggs.79.199

46.   Hu Q, Nelson WJ, Spiliotis ET. Forchlorfenuron Alters Mammalian Septin Assembly, Organization,
and Dynamics. Journal of Biological Chemistry. 2008;283: 29563–29571.
doi:10.1074/jbc.M804962200

47.   Angelis D, Karasmanis EP, Bai X, Spiliotis ET. In Silico Docking of Forchlorfenuron (FCF) to Septins
Suggests that FCF Interferes with GTP Binding. PLoS One. 2014;9: e96390.
doi:10.1371/journal.pone.0096390

48.   Kim KK, Singh RK, Khazan N, Kodza A, Singh NA, Jones A, et al. Development of Potent
Forchlorfenuron Analogs and Their Cytotoxic Effect in Cancer Cell Lines. Sci Rep. 2020;10: 3241.
doi:10.1038/s41598-020-59824-4

49.   Valero C, Colabardini AC, Chiaratto J, Pardeshi L, de Castro PA, Ferreira Filho JA, et al. Aspergillus
fumigatus Transcription Factors Involved in the Caspofungin Paradoxical Effect. mBio. 2020;11.
doi:10.1128/mBio.00816-20

50.   Beauvais A, Latgé J-P. Membrane and cell wall targets in Aspergillus fumigatus. Drug Resistance
Updates. 2001;4: 38–49. doi:10.1054/drup.2001.0185

51.   Gastebois A, Mouyna I, Simenel C, Clavaud C, Coddeville B, Delepierre M, et al. Characterization
of a New β(1–3)-Glucan Branching Activity of Aspergillus fumigatus. Journal of Biological
Chemistry. 2010;285: 2386–2396. doi:10.1074/jbc.M109.077545

52.   Sarthy A V., McGonigal T, Coen M, Frost DJ, Meulbroek JA, Goldman RC. Phenotype in Candida
albicans of a disruption of the BGL2 gene encoding a 1,3-β-glucosyltransferase. Microbiology (N
Y). 1997;143: 367–376. doi:10.1099/00221287-143-2-367

53.   Kalebina TS, Farkaš V, Laurinavichiute DK, Gorlovoy PM, Fominov G V., Bartek P, et al. Deletion of
BGL2 results in an increased chitin level in the cell wall of Saccharomyces cerevisiae . Antonie Van
Leeuwenhoek. 2003;84: 179–184. doi:10.1023/A:1026034123673

54.   Millet N, Latgé J-P, Mouyna I. Members of Glycosyl-Hydrolase Family 17 of A. fumigatus
Differentially Affect Morphogenesis. Journal of Fungi. 2018;4: 18. doi:10.3390/jof4010018

55. Zhao W, Lu Y, Ouyang H, Zhou H, Yan J, Du T, et al. N-Glycosylation of Gel1 or Gel2 is vital for cell
wall-glucan synthesis in Aspergillus fumigatus. Glycobiology. 2013;23: 955–968.
doi:10.1093/glycob/cwt032

56. Mouyna I, Morelle W, Vai M, Monod M, Léchenne B, Fontaine T, et al. Deletion of GEL2 encoding
for a β(1–3)glucanosyltransferase affects morphogenesis and virulence in Aspergillus fumigatus.
Mol Microbiol. 2005;56: 1675–1688. doi:10.1111/j.1365-2958.2005.04654.x

57. Große V, Krappmann S. The Asexual Pathogen Aspergillus fumigatus Expresses Functional
Determinants of Aspergillus nidulans Sexual Development. Eukaryot Cell. 2008;7: 1724–1732.
doi:10.1128/EC.00157-08

58. Han K-H, Han K-Y, Yu J-H, Chae K-S, Jahng K-Y, Han D-M. The nsdD gene encodes a putative GATA-
type transcription factor necessary for sexual development of Aspergillus nidulans. Mol
Microbiol. 2001;41: 299–309. doi:10.1046/j.1365-2958.2001.02472.x

59. O'Gorman CM, Fuller HT, Dyer PS. Discovery of a sexual cycle in the opportunistic fungal
pathogen Aspergillus fumigatus. Nature. 2009;457: 471–474. doi:10.1038/nature07528

60. Szewczyk E, Krappmann S. Conserved Regulators of Mating Are Essential for Aspergillus
fumigatus Cleistothecium Formation. Eukaryot Cell. 2010;9: 774–783. doi:10.1128/EC.00375-09

61. Li X, Chang YH. Amino-terminal protein processing in Saccharomyces cerevisiae is an essential
function that requires two distinct methionine aminopeptidases. Proceedings of the National
Academy of Sciences. 1995;92: 12357–12361. doi:10.1073/pnas.92.26.12357

62. Brown JA, Sherlock G, Myers CL, Burrows NM, Deng C, Wu HI, et al. Global analysis of gene
function in yeast by quantitative phenotypic profiling. Mol Syst Biol. 2006;2.
doi:10.1038/msb4100043

63. da Silva Ferreira ME, Kress MRVZ, Savoldi M, Goldman MHS, Härtl A, Heinekamp T, et al. The
akuBKU80 Mutant Deficient for Nonhomologous End Joining Is a Powerful Tool for Analyzing
Pathogenicity in Aspergillus fumigatus. Eukaryot Cell. 2006;5: 207–211. doi:10.1128/EC.5.1.207-
211.2006

64. Fortwendel JR, Juvvadi PR, Pinchai N, Perfect BZ, Alspaugh JA, Perfect JR, et al. Differential effects
of inhibiting chitin and 1,3-β-D-glucan synthesis in ras and calcineurin mutants of Aspergillus
fumigatus. Antimicrob Agents Chemother. 2009;53: 476–482. doi:10.1128/AAC.01154-08

65. Bourne R. ImageJ. Fundamentals of Digital Imaging in Medicine. London: Springer London; 2010.
pp. 185–188. doi:10.1007/978-1-84882-087-6_9

66. Renshaw H, Vargas-Muñiz JM, Richards AD, Asfaw YG, Juvvadi PR, Steinbach WJ. Distinct Roles of
Myosins in Aspergillus fumigatus Hyphal Growth and Pathogenesis. Infect Immun. 2016;84:
1556–1564. doi:10.1128/IAI.01190-15

67. Steinbach WJ, Cramer RA, Perfect BZ, Asfaw YG, Sauer TC, Najvar LK, et al. Calcineurin Controls
Growth, Morphology, and Pathogenicity in Aspergillus fumigatus. Eukaryot Cell. 2006;5: 1091–
1103. doi:10.1128/EC.00139-06

68. Steinbach WJ, Benjamin Jr DK, Trasi SA, Miller JL, Schell WA, Zaas AK, et al. Value of an
inhalational model of invasive aspergillosis. Med Mycol. 2004;42: 417–425.
doi:10.1080/13693780410001712034

69. Juvvadi PR, Belina D, Soderblom EJ, Moseley MA, Steinbach WJ. Filamentous fungal-specific
septin AspE is phosphorylated in vivo and interacts with actin, tubulin and other septins in the
human pathogen Aspergillus fumigatus. Biochem Biophys Res Commun. 2013;431: 547–553.
doi:10.1016/j.bbrc.2013.01.020

70. Al Abdallah Q, Ge W, Fortwendel JR. A Simple and Universal System for Gene Manipulation in
Aspergillus fumigatus: In Vitro-Assembled Cas9-Guide RNA Ribonucleoproteins Coupled with
Microhomology Repair Templates. mSphere. 2017;2. doi:10.1128/mSphere.00446-17

Figures:

Figure 1. AspA, AspB, and AspC are involved in fungal response to caspofungin. The

deletion of *aspA, aspB, and aspC* results in reduction of viability to echinocandins. (A) Spore

dilution assay show an increase in susceptibility to caspofungin in the Δ*aspA*, Δ*aspB,* and

Δ*aspC* strains. 10^4-10^1 conidia were plated on GMM media supplemented with caspofungin for

48 hours at 37°C. (B, C) Deletion of *aspB* results in loss of viability when grown in caspofungin

but not micafungin. 10^4 conidia were incubated for 48 hours in GMM supplemented with (B)

1μg/mL caspofungin or (C) 1 μg/mL micafungin. Cells were incubated in CFDA for one hour

then visualized. Viable and non-viable colonies were counted. The number of viable colonies

was divided by the total number of colonies. Experiment was replicated three times. Error bar

represent SEM. Student t-test were done in Graphpad Prism, with each strain compared against

the *akuB*[KU80] wild-type. Asterisks denote statistical significance levels as follows: ns p>0.05, *

p≤0.05, ** p≤0.01, *** p≤0.001, and **** p≤0.0001. (D) E-strip plates of caspofungin and

micafungin show clearer zone of effect in Δ*aspA*, Δ*aspB*, and Δ*aspC* strains. 10^6 conidia were

plated with beads and left to dry. E-strip was placed, and plates were grown at 37°C for 48

hours. All experiments were replicated three times. Representative images are shown in this

figure.

Figure 2. Δ*aspB* strain loses cell viability when grown in caspofungin. (A) All strains had

viability when grown in GMM+UU. 10^4 conidia were grown in 4 mL GMM+UU at 37°C for 24

hours. Cells were then treated with CFDA, which is hydrolyzed in living cells to a fluorescent

ester, then visualized. (B) Loss of viability in Δ*aspB* strain is seen after growth in the presence

of caspofungin. After 48 hours post exposure to 1 μg/mL caspofungin, cells were incubated in

CFDA for one hour then visualized. Experiments were replicated three times. Scale bar is 500

μm.

Figure 3. Δ*aspB* strain has increased hyphal damage during caspofungin exposure. (A)

No hyphal damage seen in basal conditions. 10^4 conidia were grown in GMM at 37°C for 24

hours. Cells were then washed with PIPES (pH 6.7) for 5 minutes twice. Slides were then

treated with propidium iodide (PI) solution, which stains nucleic acids. Coverslips were washed

twice with PIPES, then prepared and visualized. (B) Hyphal damage was seen in Δ*aspB* strain

treated with caspofungin. 10^4 conidia were grown in GMM at 37°C for 24 hours. Coverslips were

then incubated in caspofungin for 2 hours at 37°C. Cells were then washed with PIPES and

treated with PI solution. Experiments were replicated three times. Scale bar is 500 μm.

Figure 4. The Δ*aspB* strain treated with caspofungin has reduced fungal burden in an

immunosuppressed intranasal murine model of invasive aspergillosis. (A-B) Mice infected

with Δ*aspB* strain and treated with caspofungin had a higher chance of survival. Ten mice per

strain per condition were infected intranasally with 4×10^6 conidia. Mice were treated with either

saline (A) or caspofungin (B) on days +1 through +4 post-infection. Mice were monitored at least

twice a day for 14 days. Survival is on a Kaplan-Meier curve with log rank pair-wise comparison

($p<0.0001$). (C) H&E stain of lungs after 3 days post-infection shows that there is a similar

inflammatory response across conditions in the region where hyphae are present. GMS stain

shows fewer and smaller fungal lesions in Δ*aspB*-infected mice treated with caspofungin.

Images taken on an Echo Rebel Hybrid Light microscope using a 40x objective. Scale bar is 20

µm. Sections are 4 µm thick. Arrows denote examples of fungal growth.

Figure 5. Deletion of *bgt1*, *gel2*, *nsdD*, and *mapA* increases susceptibility to caspofungin.

(A) Δ*nsdD* has a significant defect in radial growth after 5 days incubation. 10^4 total conidia of

each strain were placed on GMM agar. Plates were incubated for 5 days and radial growth was

measured every 24 hours. Experiments were replicated three times. (B) Deletion of *aspB, bgt1,*

gel2, nsdD, or *mapA* results in reduced conidial production. Conidia were collected on day 5 of

growth on GMM media using 10 mL of 0.05% Tween-80. Conidia were counted using a

hemocytometer, and values were divided by the total area of growth to account for strains with

growth defects. (C) Δ*aspB* and Δ*bgt1* have an increase in the length of the apical compartment.

Δ*gel2, ΔnsdD*, and Δ*mapA* have a decrease in the length of the apical compartment. 10^4 total

conidia of each strain were inoculated onto coverslips immersed in GMM and incubated for 15

hours. Coverslips were stained with aniline blue and visualized. The length of the apical

compartments (N=20) was measured using imageJ. (D) Spore dilution assays show Δ*bgt1*,

Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains have an increase in susceptibility to caspofungin. Conidia

(10^4-10^1) were grown on GMM media supplemented with 1 µg/mL caspofungin for 48 hours. (E,

F) Deletion of *bgt1*, *gel2*, *nsdD*, and *mapA* results in reduced, but not loss of, viability when

grown in caspofungin but not micafungin. Deletion of *gel1* does not affect viability when exposed

to caspofungin or micafungin. 10^4 conidia were grown in GMM supplemented with either 1

µg/mL (E) caspofungin or (F) micafungin for 48 hours, then incubated in CFDA and visualized.

Error bars represent SEM. Experiments were replicated three times. One-way ANOVA with

Tukey's multiple-comparison test was performed in Graphpad Prism and was declared

significantly different at a p-value of <0.05. Group means with different lowercase letters are

significantly different.

Figure 6. Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* strains have reduced cell viability and

increased hyphal damage from caspofungin. (A) All strains were fully viable when grown in

GMM media. Conidia (10^4) were grown in GMM for 24 hours on coverslips then incubated in

CFDA. Slides were prepared and visualized on an Inverted Leica DMi8 with Leica K5

Microscope Camera using a 10x objective. (B) Candidate deletion strains did not lose total

viability, but strains Δ*bgt1*, Δ*gel2*, Δ*nsdD*, and Δ*mapA* had a partial reduction in viability. Conidia

(10^4) were grown on coverslips immersed in GMM supplemented with 1 µg/mL caspofungin for

48 hours. Coverslips were then incubated in CFDA and visualized. (C) Hyphal damage was not

observed in basal conditions. Conidia (10^4) were grown on coverslips in GMM for 24 hours.

Coverslips were washed in PIPES (pH 6.7) for 5 minutes twice then treated with propidium

iodide (PI) solution. Coverslips were washed twice with PIPES, then visualized on an Inverted

Leica DMi8 with Leica K5 Microscope Camera using a 10x objective. (D) Strains Δ*bgt1*, Δ*gel2*,

Δ*nsdD*, and Δ*mapA* had hyphal damage when treated with caspofungin. Conidia (10^4) were

grown on coverslips in GMM for 24 hours, then incubated in caspofungin for 2 hours. Cells were

then washed with PIPES and incubated with PI. Experiments were performed at 37°C. All

experiments were replicated three times. Scale bar is 500 µm.

Figure 7. Double-deletion mutants are similar to the Δ*aspB* single mutant under

caspofungin exposure. (A) Δ*aspB*Δ*nsdD* radial growth defect is similar to that of the Δ*nsdD*

parent. 10^4 total conidia of each strain were placed on GMM agar. Plates were incubated for 5

days and radial growth was measured every 24 hours. (B) Double deletion mutants resemble

Δ*aspB* conidiation defect. Conidia were collected on day 5 using 10 mL of 0.05% Tween-80.

Conidia were counted via hemocytometer and values were divided by the total area of growth to

account for strains with growth defects. (C) Deletion mutants' conidia are viable. 10^7 spores

were spun down, washed with PIPES twice, and incubated with propidium iodide and calcofluor

white for 5 minutes. Spores were washed twice again, then imaged and counted for viability. (D)

Δ*aspB* and double mutants have fungicidal response to caspofungin. 10^4 conidia were grown in

GMM supplemented with 1 µg/mL caspofungin for 48 hours then incubated in CFDA and

visualized. Viable colonies were counted. All experiments done in triplicate. Error bars represent

SEM. One-way ANOVA with Tukey's multiple-comparison test were performed in Graphpad

Prism and were declared significantly different at a p-value of <0.05. Group means with different

lowercase letters are significantly different.

Supplementary Figure 1. Caspofungin exposure alters AspB protein interactions. (A)

Principal component analysis (PCA) of the proteomics data reveals different protein composition

from AspB pulldown from GMM and caspofungin treated samples. Blue color denotes those

grown in GMM, red color denotes fungi grown in GMM+ 1 µg/mL caspofungin. (B) 226

significantly decreased (FC < -2, p < 0.05) proteins (blue) and 106 significantly increased (FC >

2, p < 0.05) proteins (red). Proteins under two-fold change or had a p-value greater than 0.05

are shown in black. Red horizontal line denotes p=0.05, red vertical lines denote ± 2 fold

change. (C) Category of biological processes most enriched after exposure to caspofungin is

hyphal growth. (D) Category of biological processes most decreased after exposure to

caspofungin is protein folding. (C,D) Gene ontology enrichment for biological processes

determined by using FungiFun3 [33].

Supplementary Figure 2. Candidate deletion mutants do not have reduced viability in

micafungin. (A) All strains were fully viable when grown in GMM media supplemented with

DMSO. Conidia (10^4) were grown in GMM supplemented with 1 µL/mL DMSO for 24 hours on

coverslips, then incubated in CFDA. Slides were prepared and visualized. (B) All strains have

some viability when grown in micafungin. Conidia (10^4) were grown on coverslips immersed in

GMM supplemented with 1 µg/mL micafungin for 48 hours. Coverslips were then incubated in

CFDA and visualized. Scale bar is 300 µm.

Supplementary Figure 3. Δ*bgt1*, Δ*gel2*, and Δ*nsdD* have mild sensitivity to calcofluor

white. 10^4 conidia were inoculated into GMM agar supplemented with cell wall disrupting agents

listed in the figure then incubated for 3 days at 37°C, with the exception of 4 µg/mL caspofungin

for five days to ensure observation of the caspofungin paradoxical effect. Experiments were

replicated three times.

Supplementary Figure 4. Candidate gene deletion mutants spot dilution assays confirm

Δ*bgt1*, Δ*gel2*, and Δ*nsdD* sensitivity to calcofluor white. Conidia (10^4-10^1) were grown on

GMM media supplemented with drug noted for 48 hours, then imaged. Experiments done in

triplicate.

Supplementary Figure 5. Double deletion mutants have hyphal damage after caspofungin

exposure. (A) Double deletion mutants have minimal hyphal damage in basal conditions.

Conidia (10^4) were grown on coverslips in GMM for 24 hours. Coverslips were washed in PIPES

(pH 6.7) for 5 minutes twice then treated with propidium iodide (PI) solution. Coverslips were

washed two times with PIPES and visualized after. (B) Double deletion mutants have increased

hyphal damage after caspofungin exposure. Conidia (10^4) were grown on coverslips in GMM for

24 hours, then incubated in caspofungin for 2 hours. Cells were then washed with PIPES and

incubated with PI. Scale bar is 250 μm. Experiments were replicated three times. All images

were standardized in ImageJ.

Figure 1

Figure 2

Figure 3

Figure 4

Fungal growth

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

A

B

C

Supplementary Figure 2

A

B

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5