Identification of multivariate phenotypes most influenced by mutation:

***Drosophila serrata* wings as a case study**

Cara Conradsen[1,2] and Katrina McGuigan[1,3]

1. School of the Environment

The University of Queensland;

St. Lucia, QLD, 4072 AUSTRALIA

2. European Centre for Environment and Human Health,

University of Exeter Medical School,

Environment and Sustainability Institute,

Treliever Road, Penryn Campus,

Penryn, TR10 9FE, United Kingdom

Environment and Sustainability Institute,

University of Exeter, Treliever Road,

Penryn Campus, Penryn, TR10 9FE, United Kingdom

3. k.mcguigan1@uq.edu.au

All data analysed in this paper is published: (Conradsen *et al.* 2021)

Running Title: Multivariate mutational variance

Keywords: mutational variance; subspace; size; morphology; DsGRP

ABSTRACT

The distribution of mutational effects, especially pleiotropic effects, significantly impacts phenotypic evolution and population persistence. However, our empirical understanding of these effects remains quite limited. Here, we interrogated multivariate wing shape data from 12 replicated samples of the same mutation accumulation experiment in *Drosophila serrata* to characterize both the variation in wing shape introduced by mutations, and the among-sample heterogeneity in estimates of that variation. Consistent with previous studies and expectations from statistical theory, there was considerable heterogeneity among replicate estimates of the same parameter (median ratio of largest to smallest estimate of mutational variance per trait: 7.6; estimates of the same pairwise trait correlation ranged up to a maximum of the theoretical limits: -1 to 1). However, multivariate analyses identified a trait combination associated with relatively similar magnitude of mutational variance in all population samples (ratio of largest to smallest estimate: 1.96). This major axis of mutational variance aligned closely with the major axis of among-line variance within most of the population samples, suggesting that each sample adequately estimated the predominant pattern of correlated mutational effects, despite the marked heterogeneity of individual (co)variance parameters. Mutation predominantly introduced variation along an axis of shorter, wider vs. longer, narrower wing tips. This was qualitatively the same mutational wing shape variance identified in independent studies of *Drosophila* wings, using different experimental designs and analytical approaches. Our study suggests that while estimation error can independently influence variances of individual traits and covariances among traits, multivariate analyses may be effective at revealing clear signals higher dimensional patterns of variation.

INTRODUCTION

How spontaneous mutations affect the phenotype is central to a range of evolutionary genetic phenomena, including the maintenance of quantitative genetic variance (Bürger 2000; Johnson and Barton 2005), and the nature of complex human diseases (Eyre-Walker 2010). Reflecting this broad significance, substantial effort has been devoted to characterising the input of new mutation to individual quantitative traits (Houle *et al.* 1996; Halligan and Keightley 2009; Walsh and Lynch 2018; Conradsen *et al.* 2022). However, this individual trait focus is at odds with the long-standing recognition that selection does not act on individual traits in isolation and that the shared genetic basis of traits influences the response to selection (Walsh and Blows 2009; Svensson *et al.* 2021). Mutational correlation among traits has been implicated as determining the stability of patterns of standing genetic covariance (Jones *et al.* 2003), highlighting the potential for mutation to exert long-term influence on the direction and rate of multivariate phenotypic evolution. Available data suggests mutations frequently have correlated effects across traits (e.g., Estes *et al.* 2005; McGuigan *et al.* 2014b; Mallard *et al.* 2023; Miller *et al.* 2023). However, how closely patterns of trait covariation generated by mutation match that observed in standing genetic variation ranges widely, from markedly different (e.g., Latimer *et al.* 2014) to highly concordant (e.g., Houle *et al.* 2017). Further empirical data are required to advance our understanding of the mutational correlations across sets of traits, the factors that contribute to variability in patterns of correlations, and their persistence to standing genetic variation.

High estimation error is a well-known challenge to studying genetic covariances (Klein *et al.* 1973; Klein 1974; Lynch and Walsh 1998). Random sampling error can generate strong spurious covariance among variables, particularly when the number of independent samples is small relative to the number of variables (traits) (Johnstone 2007; Blows and McGuigan 2015;

Sztepanacz and Blows 2017). Relative to genetic covariances among highly polygenic traits,

this statistical problem is likely to be exacerbated for studies of mutation: few mutations

affecting the traits are sampled, resulting in small magnitudes for covariances (i.e., small

statistical effect size). This limitation can be mitigated over hundreds of generations of

mutation-drift evolution, with accumulated mutational variance approaching the magnitude of

standing genetic variation (Lynch *et al.* 1999; Johnson and Barton 2005). However, phenotype

variation in such experiments is likely to increasingly reflect the combined effects of multiple

mutations per genotype (homozygous genetic line), which will limit insight into the distribution

of mutational effects. Furthermore, due to logistical limitations, long-term maintenance might

typically trade-off with sample size (e.g., the number of mutationally divergent genotypes

assayed), opposing the statistical gain from higher variance. Thus, mutational covariance

estimates are expected to typically have low accuracy and precision, reducing our ability to

characterize mutational effects, and to establish general patterns or causes of heterogeneity in

trait covariances.

The converse to this perspective of statistical limits on our capacity to acquire

knowledge is that multivariate analyses may substantially improve the statistical power to

estimate effects by leveraging the shared information among covarying traits (Schmitz *et al.*

1998). For example, growing availability of extensive phenotypic data in human genome-wide

association studies (Bycroft *et al.* 2018) has been exploited via multi-trait frameworks that

demonstrate greatly improved variant detection (Qi and Chatterjee 2018; Turley *et al.* 2018;

Baselmans *et al.* 2019; Wu 2020). Several studies have sought to understand mutational

covariance patterns via eigenanalyses of multi-trait mutational variance-covariance matrices

(e.g., Latimer *et al.* 2014; Houle *et al.* 2017; Dugand *et al.* 2021; Mallard *et al.* 2023). The first

eigenvector (g_{max}) of the additive genetic (co)variance matrix, **G**, reflects the dominant pattern

of shared standing genetic variation among the traits, and is of particular interest in

evolutionary genetics because it describes the phenotype that can evolve most rapidly (Lande 1979; Schluter 1996). Similarly, the leading eigenvector of the mutational (co)variance matrix, m_{max}, informs how mutation may most strongly influence **G**, and future evolutionary potential. Sampling error is well known to inflate the estimated magnitude of variance along these eigenvectors, and statistical tools have been developed to account for this (Blows and McGuigan 2015; Sztepanacz and Blows 2017). However, how well experiments might typically capture the multivariate effects of mutation, as opposed to sampling error and biological factors such as environmentally dependent mutational effects, has received little attention.

Drosophila wings have emerged as a model for studying genetics and evolution (Mezey and Houle 2005; Carreira *et al.* 2011; Houle *et al.* 2017; Dugand *et al.* 2023). Wing veins (the relative positions of which define wing shape traits) have a shared developmental basis, controlled by diffusion of patterning genes (reviewed in Blair 2007; Matamoro-Vidal *et al.* 2015), suggesting that covariance among shape traits should be pervasive. Consistent with this prediction, empirical evidence indicates substantial genetic (Mezey and Houle 2005; McGuigan and Blows 2007) and mutational (Houle *et al.* 2017; Dugand *et al.* 2021) covariance among parts of the wing. Furthermore, specific alleles causing broadscale changes across the wing (i.e., in multiple shape traits) have been identified as segregating in natural populations (e.g., Pitchers *et al.* 2019). Here, we seek to exploit the shared information expected to arise through pleiotropic mutation on wing shape to determine the statistical robustness of estimates of multivariate mutational variance in *Drosophila serrata* wing shape.

Our experimental design is based on a classical mutation accumulation experiment, in which an inbred ancestor is subdivided into daughter populations that diverge from one another via chance fixation of spontaneous mutations (Halligan and Keightley 2009). Following 20

generations of brother-sister inbreeding (and mutation accumulation), each of the 42 MA lines

were subdivided and maintained over six generations as two sublines, differing in population

size, with wing phenotypes sampled every generation (Conradsen *et al.* 2022; Figure 1). This

design, in which effects of mutations fixed during the 20 generations of mutation accumulation

are measured repeatedly, allowed us to characterize how experimental factors might influence

estimates of among-line (mutational) covariance. Specifically, the 12 experimental samples

contain the same set of fixed mutations, but potentially vary due to the influence of micro-

environment, segregating mutations, measurement and estimation errors (as detailed in

Conradsen *et al.* 2022). Through comparative analyses of the 12 mutational covariance

matrices, we seek to both determine how these factors affect estimation, and to robustly

estimate the major patterns of multivariate wing shape variation introduced by new mutation

in *D. serrata*.

METHODS

Population history and wing data collection

The experimental design was described in detail in Conradsen *et al.* (2022). Briefly, we used a

highly inbred *D. serrata* Reference Genome Panel (DsRGP) line (Reddiex *et al.* 2018),

DsGRP-226, as a progenitor to establish 200 MA lines that were maintained for 20 generations

by brother-sister inbreeding (Kannan *et al.* 2023). From these MA lines, 42 were randomly

selected and expanded to found two sublines (84 lines in total) under two population size

treatments (Small and Large: Figure 1a) (Conradsen *et al.* 2022). For each of the 84 sublines,

wings were sampled from up to six male flies in each of two replicate rearing vials for six

sequential generations: a total of 5,135 wings were available for analysis (Conradsen *et al.*

2021).

Nine wing landmark positions (Figure 1b) were recorded from images, aligned using a

generalised Procrustes least squares superimposition, and centroid size (CS, the square root of

the sum of the squared distances between each land-mark and their centroid) determined (Rohlf

2007). Inter-landmark distances (ILDs, in units of centroid size) were calculated from the

aligned coordinates. Using the model described below, we attempted to estimate the among-

line variance-covariance matrix for the 11 traits previously analysed individually by Conradsen

et al. (2022). However, this model failed to converge – a likely consequence of the relatively

few lines (42) and the small magnitude of among-line variance accumulated after 20

generations of mutation-drift evolution. We therefore report on a subset of those 11 traits, and

consider further in the Discussion how our pragmatic, but non-random, trait selection may

influence conclusions. To arrive at the subset of five ILDs, along with CS (Figure 1b), that

were retained for multivariate analysis, four of the original 11 traits (ILD1.9, ILD3.8, ILD4.8

and ILD5.6) were excluded because at least one of the 12 (two treatments measured in six

generations) among-line variance estimates was zero (see Figure 5 of Conradsen *et al.* 2022),

negatively impacting on multivariate model convergence. A further trait (ILD4.5) was

excluded as models containing this trait returned a non-positive definite asymptotic matrix

(indicating poor model convergence) in several samples.

Estimating mutational variance-covariance matrices

Prior to analysis, Mahalanobis distance was used to identify multivariate outliers (critical value

of χ^2 = 22.46, df = 6, α = 0.01), which were excluded (55 outliers across the 5135 flies, <

1.1%). To facilitate convergence in multivariate models, data for each trait was variance

standardised (mean = 0, standard deviation = 1) across all generations and treatments. We

estimated the 12 among-line (**M**) covariance matrices for our six traits, using restricted-

maximum likelihood (REML) in PROC MIXED in SAS v9.4 (SAS Institute Inc., Cary, NC.)

to fit:

$$y_{ijk} = \mu + Line_i + Vial_{j(i)} + \varepsilon_{ijk} \tag{1}$$

where y_{ijk} was the vector of six trait values for the kth wing ($k = 1\ldots6$), for the jth vial ($j = 1,2$),

within the ith line ($i = 1...42$); μ was the vector of global means per trait; the random effects of

Line, *Vial* and the residual error (ε) estimated the 6 x 6 unstructured covariance matrices

corresponding to among-line (mutational), among replicate vial and among individual flies

within a rearing vial, respectively.

To determine the number of statistically supported independent axes of among-line

variation, a factor-analytic approach (Hine and Blows 2006) was applied, using likelihood ratio

tests (LRT) to compare sequential models constraining among-line variance from zero up to

six dimensions. The LRT degrees of freedom were defined by the difference between models

in the number of parameters estimated. Factor analytic modelling is conservative but represents

the most robust approach for determining statistical support for the dimensionality of

covariance matrices (Sztepanacz and Blows 2017).

To place confidence intervals about estimates of the variance components for result

visualisation, we applied the REML-MVN method developed by Meyer and Houle (2013;

Houle and Meyer 2015). To quantify uncertainty, 10,000 random samples were drawn from

the distribution $N \sim (\hat{\boldsymbol{\theta}}, \mathbf{V})$, where $\hat{\boldsymbol{\theta}}$ was the vector of observed (REML) covariance parameter

estimates, and $\mathbf{V}$ was the asymptotic variance-covariance matrix. These samples were used to

calculate confidence intervals for both model-estimated and derived parameters (detailed

below).

Constructing null model variance-covariance matrices

The overarching null hypothesis of the current study is that we have 12 repeated measures of the same mutational (co)variance of wing traits, differing from one another only through transient segregation of recent mutations, micro-environmental effects and estimation error. We constructed a null distribution reflecting this by randomly sampling MA lines (with replacement) to create 12 experimental populations (two treatments by six generations) of 42 lines. By shuffling lines among population size treatments and generations, while retaining the within-line information (i.e., the set of replicate vials and sampled individual flies within each vial were collectively assigned their new population identity), we maintained the signal of mutation in the populations, focusing our null on the potential for sampling error to generate spurious divergence among 12 repeated measures. We generated 1000 (randomised) null datasets, and again implemented model 1 to estimate 12 **M**. Of the 12,000 null data models (1000 randomised datasets, each analysed to estimate 12 population sample **M**), nine models failed to converge, and those **M** were omitted from analyses. Interpretation of the null is thus based on all 1000 randomised datasets where nine datasets had only 11 populations for which all **M** models converged. The null matrices were taken through the same analytical pipeline (detailed below) as the observed data.

Comparing variance-covariance matrices

We used Krzanowski's subspace method to determine the extent to which the 12 **M** estimated the same multivariate axes of mutational variance. This approach is described in detail elsewhere (Krzanowski 1979; Blows *et al.* 2004; Aguirre *et al.* 2014). Briefly, R (R Core Team 2024) was used to diagonalise each **M**, and to estimate the Krzanowski subspace matrix, **H**:

$$\mathbf{H} = \sum_{t=1}^{p} \mathbf{L}_t^{\mathrm{T}} \mathbf{L}_t \qquad (2)$$

where $\mathbf{L}_t$ contains the subset of the j^{th} largest eigenvectors, as rows, of the t^{th} $\mathbf{M}$ matrix, (and

$\mathbf{L}_t^{\mathrm{T}}$ its transpose containing eigenvectors as columns) and $p = 12$ is the number of populations

to be compared. Spectral decomposition of $\mathbf{H}$ describes the degree of shared structure across

the matrices, with the eigenvalues ranging from 0 to p. Here, an eigenvalue of 12 would indicate

that all 12 $\mathbf{M}$ have variance in the direction of the associated eigenvector of $\mathbf{H}$. The

interpretation of $\mathbf{H}$'s eigenvalues depends on how j is defined (Krzanowski 1979; Blows *et al.*

2004; Aguirre *et al.* 2014). When $j > \frac{n}{2}$ (where n is the number of traits; here, $n = 6$, setting $j >$

3), the analysis is constrained to recover common subspaces of two matrices. Conversely, if j

is allowed to vary among populations, above $j = \min(j_1, j_2, \ldots j_p)$ dimensions are constrained to

be orthogonal (i.e., not contain a common subspace) for at least one population — the one with

the lowest j. Here, we followed Aguirre *et al.* (2014) in defining j for the t^{th} $\mathbf{M}$ matrix as the

number of eigenvectors required to explain at least 90% of the variance; for each of our 12

sampled matrices, this included all dimensions with statistically supported variation (Table 1,

Table S2). The 90% criterion ensured that we could interpret at least the first two eigenvalues

of $\mathbf{H}$ (Table 1). For four $\mathbf{M}$, $j = 2$, indicating that these matrices were constrained to share no

more than two common dimensions with any other matrix; for seven of the 12 $\mathbf{M}$, $j = 3$, and

these $\mathbf{M}$ may share all three dimensions. Notably, only one $\mathbf{M}$ had $j > 3$, indicating that no two

matrices were constrained to share common subspaces.

To place confidence intervals on the eigenvalues of $\mathbf{H}$, we applied equation 2 and

extracted the eigenvalues of the resulting matrix for each of the 10,000 REML samples of the

12 matrices. To determine which of the 12 matrices had variance in which parts of the subspace

captured by $\mathbf{H}$, and whether matrices differed in the magnitude of among-line variance in

shared (common) subspaces, we projected the eigenvectors of $\mathbf{H}$ through each $\mathbf{M}$ (Aguirre *et*

al. 2014). To place confidence intervals on these estimates of common-subspace variance,

eigenvectors of the observed REML $\mathbf{H}$ matrix were projected through each of the 10,000

REML-MVN samples of the matrix. The null datasets were also analysed, with equation 2

applied to estimate **H** for each of the 100 null datasets (we used $j = 3$ eigenvectors for all null

datasets).

RESULTS

Among-line variation in the 12 population samples

Consistent with previous analyses of each trait individually (Conradsen *et al.* 2022), we

observed the 12 population estimates of among-line (mutational) variance to vary in both

magnitude and statistical support for all traits (Figure 2a,b; Table S1). There were no patterns

of consistent change over time or divergence between the population size treatments (Figure

2a), suggesting the observed heterogeneity was not due to mutation-drift-selection processes.

There was statistical support for 46 (63.9%) of the 72 estimates (6 traits in 12 population

samples), with statistical support across all 12 samples for only one trait (ILD1.5, capturing

proximo-posterior wing width: Figure 1b) (Figure 2a,b). Particularly for traits with fewer

statistically supported estimates (Figure 2b) there were large differences in magnitude of

among-line variance (ratio of largest to smallest ranged from 1.93 up to 34.4, median 7.6;

Figure 2a). Nonetheless, there was little statistical support for among-line variance to differ

between the population samples - estimates typically fell well within each other's 90% CI, with

only a few exceptions for traits ILD1.2 (two largest versus five smallest estimates) and ILD2.8

(largest versus smallest) (Figure 2a).

There was considerably less statistical support for mutational covariance among traits,

with only 15% (27 of 180) of pairwise covariances statistically distinct from zero (Figure 2a,c).

Nonetheless, the average absolute correlation was 0.51 (although this value is upwardly biased

by estimates at the statistical boundary of -1 or 1: Table S1). This pattern of frequent strong

association between traits, and consistency of sign (particularly for statistically supported

estimates: Figure 2a,c) suggests that the relatively large sampling error of covariance estimates,

rather than a lack of correlated mutational effects among traits, might contribute to the limited

statistical evidence for mutational covariance. Binomial tests of the null hypothesis of an equal

number of positive and negative values rejects the null for six (of 15, 40%) trait pairs that had

consistent sign for at least 10 estimates ($P = 0.0193$) (Figure 2a,c).

The distribution of eigenvalues of each population sample's among-line covariance

matrix was consistent with mutation generating trait covariance, and with variation in

mutational effects among traits. Specifically, the first eigenvector accounted for substantially

more variance than any individual trait (Figure 3; Table 1 vs Table S1), consistent with shared

variance among some traits, while robust statistical support for two axes of mutational variance

in most population samples (Table S2) suggested that not all mutations had concordant effects

across all traits. These general conclusions are well supported by consistent evidence of at least

some statistically supported trait covariances in each population sample (median = 2 of the 15

covariances per population sample: Figure 2a), and consistent patterns of eigenvalue

distributions (Table 1; Figure 3). Notably, relative to the individual traits, there was markedly

less heterogeneity among the 12 samples in the magnitude of the total variance (ratio of largest

to smallest trace: 2.16) and eigenvalues (ratio of largest to smallest: 1.92, 2.90 and 3.08 for m_1,

m_2 and m_3, respectively: Table 1). Thus, traits might be independently affected by sampling

error in each population sample, as suggested by the absence of general patterns across traits

in the variation among the 12 population sample estimates (Figure 2a; see also Conradsen *et*

al. 2022). The multivariate analysis may be relatively effective at partitioning out independent

(uncorrelated) spurious signals affecting individual traits, resulting in more consistent

conclusions about the magnitude of variance. Nonetheless, given the heterogeneity among the

12 estimates of variance per trait and covariance per trait pair (Figure 2a), the specific

multivariate wing shapes associated with among-line variance (Table S3) may differ among

population samples. Below, we present further analyses aimed at addressing this.

Common subspace and major axis of mutational variation

There was strong support for consistent orientation of multivariate among-line variance across

the 12 population samples. Because sampling error is expected to impact each of the 12 **M**

independently, shared patterns of among-line variance provide strong evidence of the nature of

mutational input to wing shape. The major axis of the Krzanowski common subspace had an

eigenvalue of 11.68, close to the maximum value (12) attainable when the vector is present in

the sampled subspace of each of the 12 populations (Table 2). Further, the eigenvalue was

within the confidence bounds estimated from the null datasets simulating a single population

(Table 2). The eigenvalues of the other two interpretable common subspace axes (8.9 and 6.6:

Table 2) implicated further axes of mutational variance, but suggesting that other factors, such

as estimation error or micro-environmental heterogeneity, may limit the robust estimation of

this variation.

As suggested by the large eigenvalue of h_1, all 12 population samples had statistically

supported variance for this multivariate wing shape trait (Figure 4a). This common axis of wing

shape variation, representing the best estimate of the multivariate wing shape most affected by

mutation in the current experiment, reflected an axis of expansion versus contraction of the

proximal versus distal wing length (strongest, opposing, loadings on h_1 from ILD1.5 and

ILD3.7: Table 2). Notably, these two traits had the most widespread statistical support for

among-line variance, the least heterogeneity in magnitude among repeated estimates, and were

significantly negatively correlated in all but one population sample (average bounded

correlation coefficient = -0.87: Figure 2; Table S1).

The major axis of multivariate mutational input, h_1, was typically captured by the

leading eigenvector of a population sample's **M** (i.e., was aligned with m_1). This is evidenced

by: i) the angle between h_1 and eigenvectors of each population's **M** being closest to zero

(indicating identical orientation) for m_1 in 10 populations (11 – 21 degrees, median = 14,

excluding S2 and L1: Table S4); ii) the magnitude of among-line variance being greater for h_1

than for h_2 or h_3 in 11 populations (Figure 4a,b) and; iii) h_1 typically accounting for almost as

much variance as m_1 did (median = 86% of m_1: Figure 4d). In population S2, h_1 was more

closely aligned with m_2 (h_1 and m_1: 37 degrees and h_1 and m_2: 15 degrees: Table S4), reflected

in less variance along h_1 than h_2 (0.30 versus 0.36: Figure 4a,b) and h_1 accounting for only 60%

as much variance as m_1 (lower right point in Figure 4d). Nonetheless, even for this relatively

incongruent population sample, m_1 did capture some signal of the major axis of mutational

variance (h_1) (Figure 4a,d), suggesting that a single population sample might, therefore,

typically be adequate to estimate the multivariate phenotypic trait that was most strongly

influenced by mutation. Along h_1, the largest estimate of among-line variance (in population

L3) was 1.96 times greater than the smallest (L4), with estimates typically well within the CI

of all other populations (Figure 4a).

The magnitude of variance along h_1 reflected the magnitude of variance along m_1, but

was consistently lower (Figure 4d). Leading eigenvalues of genetic (and mutational)

covariance matrices are particularly vulnerable to inflation by sampling error (Hill and

Thompson 1978; McGuigan *et al.* 2014a), but the Krzanowski common subspace analysis finds

the shared pattern among population samples, rather than being strongly influenced by

independent sampling error within each population sample. Comparing variance along h_1 and

population-specific m_1 confirms the expectation that individual experiments will be prone to

overestimating multivariate mutational variance. There also remains, however, heterogeneity

among replicated estimates in the estimated magnitude of mutational variance (i.e., along h_1)

(Figure 4a). A pattern of greater variance in earlier and later generations for both population size treatments (with L3 estimate a notable exception) may suggest that heterogeneity in micro-environmental effects among generations weakly influenced the magnitude of among-line variance.

There were no clear patterns in the magnitude of variance along h_2 or h_3, although there was a weak trend for the first two generation estimates to be larger than later in time, and for the presence of among-line (mutational) variance to be more robustly supported along h_2 in the large populations and along h_3 in the small populations (Figure 4b,c). Despite this pattern in statistical support, there was little difference between the population size treatments in the magnitude of variance along h_2 (median = 0.148 vs 0.141 in small vs large population treatment, respectively; Figure 4b) or in the propensity for this axis to be most closely aligned to m_2 within each population (median angle = 0.28 vs 0.27 in small vs large population treatment, respectively; Table S4). For h_3, there was higher median variance in the small population size treatment (0.094 versus 0.059; Figure 4c) and a greater propensity for h_3 to be most closely aligned with lower eigenvectors (2x m_6, 1x m_5; Table S4) of the among-line variance matrix in the large than small population size treatment.

Notably, h_2 primarily reflects mutational variance in wing size (CS), while h_3 is most strongly influenced by ILD2.5 (Table 2): these are the only two traits for which individual trait analyses presented by Conradsen *et al.* (2022) robustly rejected the null hypothesis that among-line variance was homogenous over time. ILD1.2 and ILD2.8 also contributed to h_2 (larger wings had disproportionally longer anterior margins, while smaller wings had shorter anterior margin: Table 2, Figure 1b), and were the two traits with the greatest heterogeneity in variances and covariance among the 12 **M** (Figure 2). Importantly, despite this evidence that specific micro-environmental context may influence the observed magnitude of variance along these

355 axes, the presence of among-line variance along h_2 and h_3 in multiple independent samples of

356 the mutation accumulation lines suggests that mutation did contribute variation to these aspects

357 of wing phenotype.

DISCUSSION

Mutations contribute, on average, 0.24% to the phenotypic variance or 1.57% of the trait mean

per generation (Houle *et al.* 1996; Lynch *et al.* 1999; Conradsen *et al.* 2022). Accurately

estimating these small effects on each trait represents a clear challenge to understanding how

trait type, genetic background and environmental context might each influence mutational

characteristics, with differences in the magnitude of mutational variance reported among

studies comparable to the magnitude of differences expected from sampling error alone

(Conradsen *et al.* 2022). Here, focusing on a biased subset of traits that Conradsen *et al.* (2022)

found to have the most statistically robust estimates, we nonetheless re-derive their inference

of large sampling error effects. There was substantial heterogeneity among the 12 estimates of

among-line variance per trait (median ratio of largest to smallest estimate was 7.57) but no

apparent temporal or between-treatment divergence in magnitude (Figure 2a; Table S1),

consistent with sampling error, rather than mutation-drift-selection process driving divergence

in the magnitude of among-line variance.

The challenge posed by estimation error is greatly exacerbated for covariances (Klein

et al. 1973; Klein 1974; Lynch and Walsh 1998). Shared development (Matamoro-Vidal *et al.*

2015), known pleiotropic alleles (Pitchers *et al.* 2019) and prior estimation of mutational

covariance (Carreira *et al.* 2011; Houle *et al.* 2017; Dugand *et al.* 2021) all led us to expected

strong among-line trait covariances for wing shape traits. The magnitude of estimates (median

absolute correlation ~0.5) was consistent with this, but only one covariance (between ILD1.5

and ILD3.7; Figure 2a,c; Table S1) was robustly statistically supported. Each of our population

samples was designed to approximate the sample size of a typical mutation accumulation

experiment (Conradsen *et al.* 2022). From this perspective, for nine of the 15 trait pairs, we

would have alternatively concluded that mutations did, or did not, have correlated effects,

depending on which of the 12 replicate samples we considered. In addition to contributing wide

confidence intervals, sampling error may generate strong (significant) spurious associations

between variables. Here, in the sign of covariance estimates were consistent for some trait pairs,

suggesting that mutation generated covariance, but for other trait pairs with inconsistently

estimated direction of covariance despite some statistically supported estimate(s) (e.g., CS:

ILD2.5; Figure 2a,c) it remains unknown if the association is spurious.

Relatively few specific traits have been investigated across more than one mutational

study leaving it relatively unknown whether mutation consistently causes certain traits to

covary. Life history traits (e.g., early versus late fecundity, productivity, longevity) are the best

studied, with independent studies in several species concordantly reporting statistically

supported positive (>0.6) mutational covariances (Houle *et al.* 1994; Estes *et al.* 2005). Positive

covariance of life history traits is predicted by their joint dependence on energetic resources

(i.e., condition dependence) (Houle 1991; Rowe and Houle 1996). However, positive

covariance can also arise in mutation accumulation lines (and during mutagenesis) via linkage

(all mutations within a line are in complete linkage with one another), particularly where

mutations have directionally biased effects on the traits (Estes *et al.* 2005). When mutational

effects are not biased to predominantly increase (or decrease) trait values, different

(independent) mutation accumulation experiments are expected to report inconsistent strength

and direction of covariance of a specific trait pair due to the chance linkage of different,

independently acting, mutations. As discussed further below, replicated patterns of the same

trait covariance in independent studies thus provides support for pleiotropic mutation, rather

than linkage, as causal of observed trait covariances among wing traits. Although we were

unable to assess it in the current study, investigations in *D. melanogaster* have found no

evidence of directional bias in mutations affecting wing shape (Santiago *et al.* 1992; Houle and

Fierst 2013). Traits under stabilising selection are predicted to lack directional bias in

mutational effects (Johnson and Barton 2005), with some evidence that *Drosophila* wing shape

is under stabilizing selection (McGuigan *et al.* 2011; Houle and Fierst 2013).

Multivariate mutational variance in *Drosophila* wings

Despite the substantial variability of most individual variance and covariance parameter

estimates, the multivariate metrics (trace, eigenvalues) were notably similar among the 12

population samples (ratio of largest to smallest ~2: Table 1, Figure 4). Further, the Krzanowski

common subspace analysis suggested that each population sample was reasonably effective at

estimating the major axes of mutational variance (Figure 4). This indicates that multivariate

analyses may be effective at partitioning out estimation error independently affecting

individual traits, leveraging shared mutational information among traits to improve the

precision of estimates.

With the Krzanowski shared subspace analysis, we uncovered a single major axis, h_1,

along which there was statistically supported mutational (among-line) variance in all 12

population samples. Trait loadings on this h_1 vector (Table 2) suggest that mutation

predominantly generated variation along an axis of shorter, wider versus longer, narrower wing

tips. Intriguingly, this was qualitatively the same mutational wing shape variance identified in

independent studies of *Drosophila* wings, using different experimental designs and analytical

approaches. Dugand *et al.* (2021) estimated mutational (co)variance in an outbred, randomly

mating, *D. serrata* population founded by the DsGRP lines (including the ancestor line of the

mutation accumulation lines considered in the current study). In that outbred population

analysis, m_1 was characterized by lengthening ILD1.5 and ILD5.7 while shortening along

ILD4.6 and ILD3.9 (and vice versa: see their Table 3). Similarly, Houle *et al.* (2017) estimated

the major axis of mutational variance using inbred mutation accumulation lines of *D.*

melanogaster (see their Extended Data Figure 1), finding that mutation pushed out the

landmarks corresponding to LM5 and LM7 in the current study (increasing ILD1.5 and

ILD1.7), while pulling closer together landmarks corresponding to LM3,4 and LM9,10

(shortening distances between those landmarks, including ILD3.7, ILD3.9 and ILD4.6).

The consistency of mutational covariance patterns across these different studies of

Drosophila wings suggests bias in either the frequency or effect sizes of pleiotropic mutations,

driving persistent bias in the phenotypic variation generated by new mutation. Houle *et al.*

(2017) found that the distribution of new wing shape variation introduced by mutation in *D.*

melanogaster strongly predicted the pattern of divergence among Drosophilidae species over

40 million years (see also Houle *et al.* 2020). Rohner and Berger (2023) extended upon this,

demonstrating that developmental processes bias the frequency distribution of wing shape

variation in another family of flies, Sepsidae, in a manner that was strongly correlated with the

distribution of mutational covariance reported by Houle *et al.* (2017). The similar patterns of

mutational variance in *D. serrata* from this study and Dugand *et al.* (2021) support the

inference from Houle *et al.* (2017) and Rohner and Berger (2023) by providing evidence that

the frequency distribution of phenotypes introduced by new mutation is conserved among

genotypes. We note this evidence of conserved effects is qualitative rather than quantitative.

This reflects, in part, methodological challenges that prevent exact duplication of trait

measurement (i.e., wing imaging methods capture different landmarks). Factors influencing

model convergence (such as the ratio of the number of inbred lines to the number of traits) can

also preclude matched analyses of the same trait sets. Importantly however, variability in the

presence of statistically supported variance for each trait can also limit direct comparability.

Such inconsistency in detection of variance may reflect random sampling from the same

underlying distribution of effects, but further data is required to assess this.

The factor analytic modelling and Krzanowski shared subspace analysis revealed robust

statistical support for additional dimensions of mutational co-variance, although variability

among the 12 population samples limited insight into the exact nature of these mutational

effects. Wing size contributed strongly to the second axis of the common subspace, h_2 (Table

2), which was associated with statistically supported among-line variance in 58% (7 of 12) of

the population samples (Figure 4b). Mutational (and genetic) variance in size may vary due to

sensitivity of growth to resource availability (Cavicchi *et al.* 1985; Bitner-Mathé and Klaczko

1999). In their analysis of individual traits, Conradsen *et al.* (2022) reported significant change

in mutational effects on size among the six assay generations. Here, we found some marginal

statistical support for the direction of covariance of size with other traits (ILD2.5 and ILD3.7)

to reverse sign among some population samples (Figure 2a: non-overlapping CI among

estimates with CI either not including or marginally including zero). Kannan *et al.* (2023)

investigated the same mutation accumulation lines, maintained via brother sister mating for a

further 10 generations (30 in total). Rearing flies under different diet and temperature

conditions, they found evidence of reduced mutational variance in male wing size when

nutrition was limited (Kannan *et al.* 2023), suggesting the potential that unintended

heterogeneity in nutrition among generations may have contributed to the observed

heterogeneity in variance along this second common axis of mutational variance.

Overall, our results suggest that even relatively modest mutation accumulation

experiments may provide valuable information on the multivariate axes of traits space to which

mutation contributes the greatest new variance each generation. Such insights are valuable,

particularly for deepening our understanding of how mutation and selection interact to shape

standing genetic variance, and long-term responses to selection – for example, does mutation

predominantly introduce variance to traits with relatively high standing genetic variation (as

observed for Drosophildae wings: Houle *et al.* 2017; Dugand *et al.* 2021) or is there a mismatch

between mutational and standing genetic variance implicating selection acting against new

mutations (for example, as observed for thermal performance traits: Latimer *et al.* 2014). The

current study, prior studies of *Drosophila* wings (Houle and Fierst 2013; Houle *et al.* 2017;

Dugand *et al.* 2021), and studies of other trait types and species (e.g., Estes *et al.* 2005; Latimer

et al. 2014; Hine *et al.* 2018; Mallard *et al.* 2023; Miller *et al.* 2023) have all highlighted the

challenge of detecting mutational variance for much of the trait space, with (near) zero

eigenvalues (so-called nearly null spaces: Gomulkiewicz and Houle 2009) observed in all

studies. While it will remain challenging to characterize the properties of such nearly null

phenotypes, determining whether they reflect chance sampling of mutations in a single

experiment, or persistently low input of mutation, is an important first step to improving our

understanding of how mutation might limit evovable variation.

In conclusion, we have shown substantial variability in the estimated magnitude (and

confidence range) of mutational (co)variances of traits, but also that this heterogeneity, driven

by uncontrollable micro-environmental variation and estimation error, can be partitioned out

in multivariate analysis. The variability we observe is likely at the lower end of the range given

our selection of mutationally variable traits (i.e., exclusion of any trait with a zero estimate of

among-line variance in any of the 12 samples) coupled with our focus on morphological traits,

which might typically have lower non-genetic contributions to observed variance than other

types of traits (Houle *et al.* 1996). The heterogeneity among estimates suggests that we will

have low confidence in the magnitude of individual mutational variances and pairwise

mutational covariances. Yet, uncorrelated non-mutational contributions to trait variability

might allow multivariate analyses to generate relatively robust insight into general

characteristics of mutational covariances. Qualitatively similar patterns of multivariate

mutational effects on *Drosophila* wing shape have been reported in different species, consistent

with relatively conserved pleiotropic effects of mutations. Drosophiladae wings, whose

development is programmed by highly conserved developmental patterning genes (Neto-Silva

et al. 2009; Matamoro-Vidal *et al.* 2015) and which evolve slowly (Houle *et al.* 2017) may be

atypical in having such conserved patterns of mutational covariance. Nonetheless, the

experimental approaches of replicated estimation and leveraging multivariate analyses have

promise for improving confidence in estimates of mutational covariance, and of detecting any

true differences in patterns among genetic backgrounds.

DATA AVAILABILITY

All data analysed in this paper are available at doi 10.6084/m9.figshare.14913051.v3

ACKNOWLEDGMENTS

The authors thank Stephen F. Chenoweth for providing us with the D. serrata MA lines, and

Adam Reddiex, Nicholas Appleton, Jack Price and Derek Sun for their help with maintaining

the flies and collecting the data. The authors also thank Emma Hine for helpful discussions

about analyses.

CONFLICTS OF INTEREST

None declared.

LITERATURE CITED

Aguirre, J. D., E. Hine, K. McGuigan and M. W. Blows, 2014 Comparing G: multivariate analysis of genetic variation in multiple populations. Heredity 112**:** 21-29, 10.1038/hdy.2013.12.

Baselmans, B. M., R. Jansen, H. F. Ip, J. van Dongen, A. Abdellaoui, M. P. van de Weijer, . . . G. Willemsen, 2019 Multivariate genome-wide analyses of the well-being spectrum. Nature Genetics 51**:** 445-451.

Bitner-Mathé, B. C., and L. B. Klaczko, 1999 Plasticity of *Drosophila melanogaster* wing morphology: effects of sex, temperature and density. Genetica 105**:** 203-210.

Blair, S. S., 2007 Wing vein Patterning in *Drosophila* and the analysis of intercellular signaling. Annual Review of Cell and Developmental Biology 23**:** 293-319, 10.1146/annurev.cellbio.23.090506.123606.

Blows, M. W., S. F. Chenoweth and E. Hine, 2004 Orientation of the genetic variance-covariance matrix and the fitness surface for multiple male sexually selected traits. Am Nat 163**:** 329-340, 10.1086/381941.

Blows, M. W., and K. McGuigan, 2015 The distribution of genetic variance across phenotypic space and the response to selection. Molecular Ecology 24**:** 2056-2072, 10.1111/mec.13023.

Bürger, R., 2000 *The Mathematical Theory of Selection, Recombination, and Mutation*. Wiley, Chichester.

Bycroft, C., C. Freeman, D. Petkova, G. Band, L. T. Elliott, K. Sharp, . . . J. O'Connell, 2018 The UK Biobank resource with deep phenotyping and genomic data. Nature 562**:** 203-209.

Carreira, V. P., I. M. Soto, J. Mensch and J. J. Fanara, 2011 Genetic basis of wing morphogenesis in *Drosophila*: Sexual dimorphism and non-allometric effects of shape variation. BMC Developmental Biology 11**:** 32, 10.1186/1471-213x-11-32.

Cavicchi, S., D. Guerra, G. Giorgi and C. Pezzoli, 1985 Temperature-related divergence in experimental populations of *Drosophila melanogaster*. I. Genetic and developmental basis of wing size and shape variation. Genetics 109**:** 665-689.

Conradsen, C., M. W. Blows and K. McGuigan, 2021 Variability of Mutational Variance Estimates: MA_dserrata_wings. Figshare https://doi.org/10.6084/m9.figshare.14913051.v3, https://doi.org/10.6084/m9.figshare.14913051.v3.

Conradsen, C., M. W. Blows and K. McGuigan, 2022 Causes of variability in estimates of mutational variance from mutation accumulation experiments. Genetics 221**:** iyac060, 10.1093/genetics/iyac060.

Dugand, R. J., J. D. Aguirre, E. Hine, M. W. Blows and K. McGuigan, 2021 The contribution of mutation and selection to multivariate quantitative genetic variance in an outbred population of *Drosophila serrata*. Proceedings of the National Academy of Sciences 118**:** e2026217118.

Dugand, R. J., M. W. Blows and K. McGuigan, 2023 Using inbreeding to test the contribution of non-additive genetic effects to additive genetic variance: A case study in *Drosophila*

serrata. Proceedings of the Royal Society B-Biological Sciences 290,
10.1098/rspb.2022.2111.

Estes, S., B. C. Ajie, M. Lynch and P. C. Phillips, 2005 Spontaneous mutational correlations
for life-history, morphological and behavioral characters in *Caenorhabditis elegans*.
Genetics 170: 645-653.

Eyre-Walker, A., 2010 Genetic architecture of a complex trait and its implications for fitness
and genome-wide association studies. Proceedings of the National Academy of
Sciences of the United States of America 107: 1752-1756, 10.1073/pnas.0906182107.

Gomulkiewicz, R., and D. Houle, 2009 Demographic and genetic constraints on evolution. The
American Naturalist 174: E218-E229, 10.1086/645086.

Halligan, D. L., and P. D. Keightley, 2009 Spontaneous mutation accumulation studies in
evolutionary genetics. Annual Review of Ecology, Evolution and Systematics 40: 151-
172, DOI 10.1146/annurev.ecolsys.39.110707.173437.

Hill, W. G., and R. Thompson, 1978 Probabilities of non-positive definite between-group or
genetic covariance matrices. Biometrics 34: 429-439.

Hine, E., and M. W. Blows, 2006 Determining the effective dimensionality of the genetic
variance–covariance matrix. Genetics 173: 1135-1144.

Hine, E., D. E. Runcie, K. McGuigan and M. W. Blows, 2018 Uneven distribution of
mutational variance across the transcriptome of *Drosophila serrata* revealed by high-
dimensional analysis of gene expression. Genetics 209: 1319-1328,
10.1534/genetics.118.300757.

Houle, D., 1991 Genetic covariance of fitness correlates: What genetic correlations are made
of and why it matters. Evolution 45: 630-648.

Houle, D., G. H. Bolstad and T. F. Hansen, 2020 Fly wing evolutionary rate is a near-isometric
function of mutational variation. bioRxiv: 2020.2008.2027.268938,
10.1101/2020.08.27.268938.

Houle, D., G. H. Bolstad, K. van der Linde and T. F. Hansen, 2017 Mutation predicts 40 million
years of fly wing evolution. Nature 548: 447-450.

Houle, D., and J. Fierst, 2013 Properties of spontaneous mutational variance and covariance
for wing size and shape in *Drosophila melanogaster*. Evolution 67: 1116-1130,
10.1111/j.1558-5646.2012.01838.x.

Houle, D., K. A. Hughes, D. K. Hoffmaster, J. Ihara, S. Assimacopoulos and B. Charlesworth,
1994 The effects of spontaneous mutation on quantitative traits. I. Variances and
covariances of life history traits. Genetics 138: 773-785.

Houle, D., and K. Meyer, 2015 Estimating sampling error of evolutionary statistics based on
genetic covariance matrices using maximum likelihood. Journal of Evolutionary
Biology 28: 1542-1549.

Houle, D., B. Morikawa and M. Lynch, 1996 Comparing mutational variabilities. Genetics
143: 1467-1483.

Johnson, T., and N. Barton, 2005 Theoretical models of selection and mutation on quantitative
traits. Philosophical Transactions of the Royal Society B: Biological Sciences 360:
1411-1425, DOI 10.1098/rstb.2005.1667.

Johnstone, I. M., 2007 High dimensional statistical inference and random matrices.
Proceedings of the International Congress of Mathematics**: 307-333.

Jones, A. G., S. J. Arnold, R. Bürger and D. Houle, 2003 Stability of the G-matrix in a
population experiencing pleiotropic mutation, stabilizing selection, and genetic drift.
Evolution 57**: 1747-1760.

Kannan, A., R. J. Dugand, N. C. Appleton, S. F. Chenoweth, C. M. Sgro and K. McGuigan,
2023 Environmental dependence of mutational (co)variances of adaptive traits.
Evolution 77**: 2341–2351, 10.1093/evolut/qpad154.

Klein, T. W., 1974 Heritability and genetic correlation: statistical power, population
comparisons, and sample size. Behavior Genetics 4**: 171-189.

Klein, T. W., J. DeFries and C. T. Finkbeiner, 1973 Heritability and genetic correlation:
standard errors of estimates and sample size. Behavior Genetics 3**: 355-364.

Krzanowski, W., 1979 Between-groups comparison of principal components. Journal of the
American Statistical Association 74**: 703-707.

Lande, R., 1979 Quantitative genetic analysis of multivariate evolution, applied to brain:body
size allometry. Evolution 33**: 402-416.

Latimer, C. A. L., K. McGuigan, R. S. Wilson, M. W. Blows and S. F. Chenoweth, 2014 The
contribution of spontaneous mutations to thermal sensitivity curve variation in
Drosophila serrata. Evolution 68**: 1824-1837, Doi 10.1111/Evo.12392.

Lynch, M., J. Blanchard, D. Houle, T. Kibota, S. Schultz, L. Vassilieva and J. Willis, 1999
Spontaneous deleterious mutation. Evolution 53**: 645-663.

Lynch, M., and B. Walsh, 1998 Correlations between characters, pp. 629-656 in *Genetics and*
Analysis of Quantitative Traits. Sinauer, Sunderland, MA.

Mallard, F., L. Noble, C. F. Baer and H. Teotónio, 2023 Variation in mutational (co) variances.
G3: Genes, Genomes, Genetics 13**: jkac335.

Matamoro-Vidal, A., I. Salazar-Ciudad and D. Houle, 2015 Making quantitative morphological
variation from basic developmental processes: Where are we? The case of the
Drosophila wing. Developmental Dynamics 244**: 1058-1073, 10.1002/dvdy.24255.

McGuigan, K., and M. W. Blows, 2007 The phenotypic and genetic covariance structure of
Drosophilid wings. Evolution 61**: 902-911.

McGuigan, K., J. M. Collet, S. L. Allen, S. F. Chenoweth and M. W. Blows, 2014a Pleiotropic
mutations are subject to strong stabilizing selection. Genetics 197**: 1051-1062.

McGuigan, K., J. M. Collet, E. A. McGraw, Y. X. H. Ye, S. L. Allen, S. F. Chenoweth and M.
W. Blows, 2014b The nature and extent of mutational pleiotropy in gene expression of
male *Drosophila serrata*. Genetics 196**: 911-921, 10.1534/genetics.114.161232.

McGuigan, K., L. Rowe and M. W. Blows, 2011 Pleiotropy, apparent stabilizing selection and
uncovering fitness optima. Trends in Ecology & Evolution 26**: 22-29, DOI
10.1016/j.tree.2010.10.008.

Meyer, K., and D. Houle, 2013 Sampling based approximation of confidence intervals for
functions of genetic covariance matrices. Proceedings of the Association for the
Advancement of Animal Breeding and Genetics 20**: 523-526.

Mezey, J. G., and D. Houle, 2005 The dimensionality of genetic variation for wing shape in
Drosophila melanogaster Evolution 59**: 1027-1038, 10.1554/04-491.

Miller, C. L., D. R. Sun, L. H. Thornton and K. McGuigan, 2023 The contribution of mutation to variation in temperature-dependent sprint speed in zebrafish, *Danio rerio*. American Naturalist 202: 519-533, 10.1086/726011.

Neto-Silva, R. M., B. S. Wells and L. A. Johnston, 2009 Mechanisms of growth and homeostasis in the *Drosophila* wing. Annual Review of Cell and Developmental Biology 25: 197-220.

Pitchers, W., J. Nye, E. J. Márquez, A. Kowalski, I. Dworkin and D. Houle, 2019 A multivariate Genome-Wide Association Study of wing shape in *Drosophila melanogaster*. Genetics 211: 1429-1447, 10.1534/genetics.118.301342.

Qi, G., and N. Chatterjee, 2018 Heritability informed power optimization (HIPO) leads to enhanced detection of genetic associations across multiple traits. PLoS Genetics 14: e1007549.

R Core Team, 2024 R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Reddiex, A. J., S. L. Allen and S. F. Chenoweth, 2018 A genomic reference panel for *Drosophila serrata*. G3: Genes, Genomes, Genetics 8: 1335-1346.

Rohlf, F., 2007 tpsRelw version 1.45. Department of Ecology and Evolution, State University of New York, Stony Brook.

Rohner, P. T., and D. Berger, 2023 Developmental bias predicts 60 million years of wing shape evolution. Proceedings of the National Academy of Sciences of the United States of America 120: e2211210120, 10.1073/pnas.2211210120.

Rowe, L., and D. Houle, 1996 The lek paradox and the capture of genetic variance by condition dependent traits. Proceedings of the Royal Society B-Biological Sciences 263: 1415-1421.

Santiago, E., J. Albornoz, A. Dominguez, M. A. Toro and C. Lopez-Fanjul, 1992 The distribution of spontaneous mutations on quantitative traits and fitness in *Drosophila melanogaster*. Genetics 132: 771-781.

Schluter, D., 1996 Adaptive radiation along genetic lines of least resistance. Evolution 50: 1766-1774.

Schmitz, S., S. S. Cherny and D. W. Fulker, 1998 Increase in power through multivariate analyses. Behavior Genetics 28: 357-363, 10.1023/a:1021669602220.

Svensson, E. I., S. J. Arnold, R. Burger, K. Csillery, J. Draghi, J. M. Henshaw, . . . A. Runemark, 2021 Correlational selection in the age of genomics. Nature Ecology & Evolution 5: 562-573, 10.1038/s41559-021-01413-3.

Sztepanacz, J. L., and M. W. Blows, 2017 Accounting for sampling error in genetic eigenvalues using random matrix theory. Genetics 206: 1271-1284.

Turley, P., R. K. Walters, O. Maghzian, A. Okbay, J. J. Lee, M. A. Fontana, . . . N. A. Furlotte, 2018 Multi-trait analysis of genome-wide association summary statistics using MTAG. Nature Genetics 50: 229-237.

Walsh, B., and M. W. Blows, 2009 Abundant genetic variation + strong selection = multivariate genetic constraints: A geometric view of adaptation. Annual Review of Ecology Evolution and Systematics 40: 41-59, 10.1146/annurev.ecolsys.110308.120232.

691   Walsh, B., and M. Lynch, 2018 Maintenance of quantitative genetic variation, pp. 1015–1078
692   in *Evolution and Selection of Quantitative Traits*. Oxford University Press.

693   Wu, C., 2020 Multi-trait genome-wide analyses of the brain imaging phenotypes in UK
694   Biobank. Genetics 215**:** 947-958.

TABLES AND FIGURES

Table 1. The variance associated with independent axes of trait mutational variance. Eigenanalyses were performed separately for **M** in each population size treatment (Treat: Small or Large) in each generation (Gen: 1 through 6). The eigenvalues (i.e., λ) and their 90% CIs* are reported, along with the number of eigenvectors (*j*) accounting for at least 90% of the variation. These *j* eigenvectors were included in the estimation of the Krzanowski subspace. The estimated **M** and the eigenvector loadings are reported in Tables S1 and S3, respectively.

Treat	Gen	m_1	m_2	m_3	m_4	m_5	m_6	*j*
Small	1	0.495	0.102	0.084	0.027	0.001	-0.024	3
		0.193; 0.798	0.014; 0.191	0.004; 0.165	-0.036; 0.092	-0.025; 0.027	-0.073; 0.026	
	2	0.506	0.270	0.066	0.015	-0.014	-0.050	2
		0.206; 0.804	0.133; 0.402	-0.005; 0.136	-0.013; 0.043	-0.040; 0.012	-0.091; -0.009	
	3	0.276	0.116	0.037	0.016	-0.011	-0.056	2
		0.079; 0.472	0.032; 0.197	-0.028; 0.102	-0.039; 0.071	-0.042; 0.020	-0.138; 0.028	
	4	0.279	0.205	0.051	0.019	0.000	-0.013	3
		0.087; 0.471	0.029; 0.375	-0.021; 0.121	-0.008; 0.045	-0.025; 0.025	-0.044; 0.017	
	5	0.429	0.159	0.114	0.029	0.010	-0.07	3
		0.213; 0.648	0.023; 0.297	0.008; 0.219	-0.001; 0.060	-0.015; 0.034	-0.147; 0.007	
	6	0.439	0.093	0.047	0.021	-0.004	-0.037	2
		0.238; 0.641	0.020; 0.165	-0.001; 0.094	-0.006; 0.049	-0.044; 0.034	-0.099; 0.025	
Large	1	0.446	0.267	0.070	0.033	0.008	-0.007	3
		0.211; 0.676	0.131; 0.406	0.007; 0.136	-0.017; 0.082	-0.013; 0.030	-0.040; 0.026	
	2	0.313	0.215	0.077	0.052	0.025	0.000	4
		0.164; 0.460	0.073; 0.354	0.015; 0.138	-0.006; 0.110	-0.000; 0.050	-0.020; 0.019	
	3	0.529	0.144	0.070	0.020	-0.020	-0.038	2
		0.272; 0.792	-0.014; 0.300	0.004; 0.137	-0.009; 0.051	-0.069; 0.029	-0.115; 0.039	
	4	0.303	0.160	0.101	0.020	-0.006	-0.029	3

	0.144; 0.462	0.046; 0.273	0.030; 0.170	-0.016; 0.058	-0.025; 0.014	-0.069; 0.010
5	0.304	0.163	0.059	0.023	0.001	0.000	3
	0.151; 0.461	0.067; 0.257	-0.019; 0.136	-0.015; 0.060	-0.054; 0.057	-0.021; 0.020
6	0.398	0.129	0.112	0.022	0.004	-0.055	3
	0.194; 0.602	0.050; 0.209	0.010; 0.214	0.004; 0.039	-0.035; 0.042	-0.111; 0.002

* CIs were determined by projecting each eigenvector (e_1: e_6 as reported in Table S3) through the 10,000 REML-MVN samples of the respective population's **M**. Although a positive lower CI suggests statistical support for non-zero variance, REML-MVN (and Bayesian MCMC posterior) CI for eigenvalues are known to be biased, leading to false positive inference for leading eigenvalues (Sztepanacz and Blows 2017). We rely instead on the robust Factor Analytic modelling (Table S2) to infer statistical support.

Table 2. Eigenvalues and eigenvectors of the Krzanowski subspace matrix, H. The 95%

CI (below estimate) for the eigenvalues (λ) of the observed data **H** were determined from

Krzanowski subspace analyses applied to each of the 10,000 REML-MVN samples of each of

the 12 **M***. For the null data **H**, the reported eigenvalue is the mean (and 95% CI the 5%

quantiles) of the Krzanowski subspace analyses applied to the 12 **M** estimated from the 1000

randomised datasets.

	h_1	h_2	h_3
λ	11.62	8.86	6.62
	10.35; 11.49	6.87; 9.27	5.16; 7.14
Null λ	11.38	10.93	8.56
	10.61; 11.79	9.35; 11.66	5.77; 10.88
CS	-0.37	0.76	-0.32
ILD1.2	0.27	0.41	0.36
ILD1.5	0.65	-0.05	-0.46
ILD2.5	0.20	-0.03	0.64
ILD2.8	0.16	0.42	0.34
ILD3.7	-0.55	-0.27	0.19

* Eigenvalues of **H** are bounded statistics (ranging from 0 to p – here, 0 to 12). In conjunction

with the nature of sampling error, which independently affects each estimated **M** and thus

contributes dissimilarity, this has resulted in the eigenvalue of h_1 falling above its upper REML-

MVN CI. Importantly, the observed λ_1 is within the range observed for the null λ_1, which

estimates the error in h_1 when the 12 samples of **M** are all drawn from the same population.

Figure 1. Schematic of (a) experimental design and (b) *Drosophila* wing traits. (a) 42 MA

lines each founded two sublines: Small (S; 12 virgin males and 12 virgin females) and Large

(L; 144 virgin males and 144 virgin females, distributed evenly among 12 vials). These 84 lines

(S and L subline per 42 MA lines) were maintained at these census population sizes for six

generations (first two shown here). Each generation, all emergent flies from the 12 vials per L

subline were pooled prior to virgin collection. For S sublines, the focal vial contributed

offspring to the next generation, while a second, replicate, vial (grey shaded) was used for wing

collection only. Each generation, six randomly chosen males were sampled from each of two

vials per line (focal and replicate vials for S; randomly chosen two for L). (b) Positions of nine

landmarks were recorded and aligned. Six traits were analysed, five inter-landmark distances

(named by their end landmarks: ILD1.2, ILD1.5, ILD2.5, ILD2.8, ILD3.7) and wing size,

characterised as centroid size (CS).

Figure 2. The variability among the 12 population sample estimates of among-line variance and covariance. (a) The REML estimate (point) and REML-MVN bootstrap confidence intervals of among-line variance (on diagonal, grey boxes; 90% CI) and covariances (below the diagonal; 95% CI) in each population size treatment (Small or Large, closed and open symbols, respectively) in each of six generations. The horizontal dashed line indicates zero; statistical support was concluded when CIs did not include zero (a one-tailed test for variances and two-tailed for covariances). Red stars highlight estimates where CI did not overlap with at least one other estimate of the same parameter. Inset plots are the counts of the 12 estimates of (b) variance for each trait or (c) pairwise covariances between traits that were (black) or were not (grey) statistically supported as different from zero. For the covariances,

39   the counts of estimates are further plotted by sign (negative or positive). See Figure 1b for trait

40   definitions and Table S1 for the parameter estimates. NB: two estimates of covariance between

41   ILD2.5 and ILD3.7 were exactly 0; these were classified as positive.

42
43

Figure 3. The distribution of among-line variation of six wing traits in 12 population

samples. Open boxplots represent distribution across the 12 **M** of the ranked eigenvalues

(largest to smallest, x-axis 1st to 6th), plotted as the proportion of the variance in their respective

M. For comparison, among-line variances of individual traits (diagonals of Table S1) were

ranked from largest to smallest within each population sample, and their distribution also

plotted (grey boxplots). Note, the combination of the six traits (i.e., eigenvector) associated

with a specific ordered eigenvalue (1st through 6th), and the specific trait ranked 1st through 6th

may differ among the 12 matrices. For each boxplot, the median of the distribution is shown

as a black band, boxes represent the interquartile range, whiskers represent 1.5 times the

interquartile range, and estimates that fell outside this range are designated with crosses.

Figure 4. The among-line variance associated with each of the three eigenvectors of the Krzanowski's subspace. Left-hand panels: The magnitude of the among-line variance (90% REML-MVN CI) is plotted for (a) h_1, (b) h_2 and (c) h_3. Within each plot, the populations are ordered by generation (x-axis), then by population size treatments (Small or Large, closed and open symbols, respectively). The dashed lines indicate zero. Note that the y-axis range differs among panels, reflecting decreasing total variance associated with each successive common subspace vector (a to c). Right-hand panels: the magnitude of variance along eigenvectors of

62   **H** are plotted against the same-rank eigenvalue of each population sample's respective **M**. (a)

63   h_1 vs m_1, (b) h_2 vs m_2 and (c) h_3 vs m_3. Population size treatment is again indicated by symbol

64   (Small or Large, closed and open symbols, respectively). The dashed blue line indicates the

65   regression slope; solid grey line is the 1:1 line indicating equal magnitude along x and y axes.