Not just mutations: Inbreeding depression persists without genetic variation

Nicolás Bonel [1,2]*, Christoph Grunau [3], Patrice David [2]

[1] Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS—CCT—CONICET Bahía Blanca), Camino de la Carrindanga km 7, Bahía Blanca 8000, Argentina.

[2] Centre d'Écologie Fonctionnelle et Évolutive, UMR 5175, CNRS—Université de Montpellier, Université Paul-Valéry Montpellier—École Pratique des Hautes Études—IRD, 34293 Montpellier Cedex 05, France.

[3] IHPE, Univ Perpignan Via Domitia, CNRS, IFREMER, Univ Montpellier, Perpignan, France

* To whom correspondence should be addressed: nicobonel@gmail.com (N. Bonel)

Running head: Persistence of inbreeding depression in genetically uniform strains

Key words: Inbreeding depression, *Physa acuta*, epigenetics, inbred lines, self-fertilization, fitness traits, evolutionary dynamics

ABSTRACT

Inbreeding depression (ID), the decline in fitness upon inbreeding, is thought to result

from a decrease in genetic heterozygosity enhancing phenotypic effects of recessive

deleterious mutations. However, emerging evidence suggests that mutations may not

explain ID completely. In this study, we test whether ID can emerge even in contexts

where genetic heterozygosity does not vary. To that end, highly inbred lines

(F=0.99999997) of the freshwater snail *Physa acuta* were used to produce individuals

with varying levels of parental relatedness (self-fertilization, sibling crosses, and cousin

crosses), though with identical genomic heterozygosity. Several fitness traits declined

significantly with increasing parental relatedness, a pattern characteristic of ID, and

quantitatively representing a non-negligible fraction of the ID usually observed in

natural, genetically diverse populations of *Physa acuta*. Individual-based simulations

showed that mutation rates compatible with values of ID found in natural populations

are way too low to generate as much ID as observed in our experiment. These findings

are consistent with the hypothesis that epigenetic changes, in addition to mutations,

could contribute to a rapid regeneration of ID and explain the persistence of detectable

ID in sets of genetically identical individuals.

INTRODUCTION

Inbreeding depression (ID) is the loss in fitness that occurs in offspring of consanguineous matings and is one of the most general observations in evolutionary biology[1,2]. Understanding the mechanisms of ID is important because it strongly constrains the evolution of mating systems (self-fertilization rates, self-incompatibility) and is a major concern in plant breeding and animal conservation[3]. ID is classically explained by the presence of recessive deleterious mutations in the genome, whose effects are exposed by homozygosity[2]. Most of the mutations that occur spontaneously at every generation are indeed deleterious and partly recessive[2,4–6]. In large diploid organisms, however, genomic mutation rates are typically small, implying that the standing inbreeding load is accumulated through many generations until a steady state of mutation, selection, and drift is reached. This view inspired the idea that a substantial fraction of standing ID could be removed in conditions that drastically reduce heterozygosity by increasing the elimination or fixation rates of alleles. The so-called genetic purging is expected to occur by selection when unusual amounts of systematic inbreeding (such as high selfing rates) are applied to large populations, exposing deleterious alleles to selection[7,8]. ID can also be lost by drift when population size is severely reduced[8,9], as alleles (deleterious or not) get fixed and heterozygosity is lost.

Inbreeding depression is, however, surprisingly difficult to eliminate completely[10]. First, while some evidence of genetic purging has come from comparative studies in natural populations as well as a few experiments[11–14], some residual ID frequently persists even in populations subject to recurrent inbreeding[12,13,15,16] Second, ID is also not systematically eliminated by drift, even within lines with extremely reduced effective size and highly homozygous, such as those propagated by continuous sib-mating or continuous self-fertilization (e.g.[17]). A classical argument explaining the difficulty of purging refers to the genetic architecture of the mutation load. If ID relies on the joint effect of many loci with very small effects, allele frequency changes under selection are expected to be very slow[18,19]. This argument, however, does not apply to fixation by drift. Purging and fixation may also be impeded by linkage and selective interference between loci[19]. Indeed, linkage disequilibrium among deleterious mutations may preserve heterozygosity (and ID) in some regions of the genome, when two deleterious alleles with strong effects (such as lethals) are in repulsion, or when stretches of weak-effect deleterious mutations collectively form pseudo-overdominant

blocks[10]. Thus, linkage can delay the loss of heterozygosity in small populations and

inbred lines. However, this loss eventually takes place given sufficient time, as

recombination ends up breaking the associations that prevent homozygosity. Once

homozygosity is attained in a genomic region, the sole mechanism that may reintroduce

heterozygosity, and regenerate ID, is the occurrence of spontaneous mutations.

Mutations, however, are rare and, within a selfing or sib-mating line, they are quickly

lost or fixed, so they are not expected to yield substantial levels of ID. Overall, an

extended period of very low effective population sizes, such as in an anciently

established inbred line, makes it unlikely to observe significant levels of ID. In other

words, aside from a few recent mutations, two individuals from the same inbred line are

essentially genetic clones. As a result, crossing them is genetically equivalent to self-

fertilization, and the fitness of offspring from crossbreeding and selfing should not

differ.

Recent studies, however, suggest another process that could regenerate

inbreeding depression in genetically uniform lines faster than spontaneous mutation:

epigenetic change (reviewed by[20]). Support for a role of epigenetics in ID comes from

several lines of evidence[21–24]. First, ID can be greatly reduced by applying a

demethylating enzyme[24]. Second, in maize, the continued decline in performance within

selfing lines from generations 7 to 11 is accompanied by a global increase in DNA

methylation while residual heterozygosity is extremely low[17]. This suggests that

epigenetic change can co-occur with sustained phenotypic deterioration even when

genetic homozygosity itself does not increase anymore. Similarly, in *Arabidopsis,*

heterosis (the reciprocal of ID) is observed when crossing isogenic epi-RILs (i.e. selfing

lines differing only in methylation patterns;[25]).

Epigenetic modifications can be independent of genetic variation ("pure" or

autonomous epigenetic variants, *sensu* Richards[26]) thus being a possible source of

heritable phenotypic variation in the absence of DNA sequence change[27]. A growing

body of evidence suggests that incomplete reprogramming and erasure of epigenetic

marks during gametogenesis and zygote development could promote stable

transgenerational transmission of some epigenetic variants[28]. Epimutations are

generated at higher rates than genetic mutations and tend to revert to the wild-type in a

few generations[29]. A fraction of the standing ID could therefore reflect fast-generated

epigenetic changes transmitted over one or a few generations. Phenotypic consequences

of autonomous epigenetic inheritance have been classically studied using systems with a

known lack of genetic variation (e.g.[21]). In a similar way, if autonomous epigenetic

changes play a role in ID, differences in offspring fitness should persist among different

types of crosses (self-fertilized, sib-mating, cousin-mating) even when all parents used

are genetically identical (i.e. clonemates). Yet, this approach has not been undertaken

yet.

Here we tested whether significant ID can be detected even when differences in

genomic heterozygosity are extremely low. To that end, we created "outbred clones" by

intercrossing inbred lines. We produced two highly inbred lines of the hermaphrodite

freshwater snail *Physa acuta* (a naturally outcrossing species) that have undergone more

than 28 generations of enforced self-fertilization, ensuring minimal residual

heterozygosity. We chose this species because it is easy to raise in the laboratory, ID

has been repeatedly observed in its natural (outbred) populations, and it can be forced to

self-fertilize (the most efficient way to eliminate heterozygosity). Individuals within

each line are genetically homozygous and identical, except for neo-mutations in the last

few generations. We crossed individuals from two different inbred lines and obtained a

set of offspring that were heterozygous and identical to one another ("outbred clone").

We then performed different types of matings with individuals from this outbred clone:

self-fertilization, sibling pair-crossing, and cousin pair-crossings. These treatments

yielded second-generation offspring with the same expected genomic heterozygosity

(ignoring recent mutations), but that differed in pedigree inbreeding, i.e. the number of

generations tracing back to the common ancestor of the two uniting gametes. We chose

to start with outbred parents (though genetically belonging to the same "clone") rather

than directly from inbred ones, because highly inbred individuals are rather weak, as

expected, a condition we deemed "unnatural". In contrast, the results obtained with

normally heterozygous, outbred individuals, are representative of processes that could

contribute to ID in natural contexts.

To evaluate whether parental relatedness (mother and father being either the

same individual, or siblings, or cousins) affected progeny fitness, we measured juvenile

survival, adult body size, probability of laying eggs (in isolation) at the age of 50 days,

and female fecundity after insemination. Classical genetic theory predicts the same

average performances in all three types of parental relatedness as offspring all have the

same genomic heterozygosity.

RESULTS

Figure 1 recapitulates the pedigree of the parents (F_2) we used to produce F_3 snails by either self-fertilizaton, sib-mating or cousin-mating. Table 1 and Figure 2 summarize our main results, while linear models are reported in detail in Tables S1-S3.

Juvenile survival significantly differed across treatments. The highest survival was in the offspring of cousin-cross (average 0.76) whereas survival in the selfing and sib-cross treatments was reduced by 18% and 9%, respectively, compared to cousin-cross (Table 1, Fig. 2A). Body weight showed a similar pattern, the cousin-cross treatment had the highest mean, with a reduction of 20% and 14% in the self-fertilization and sib-cross treatment, respectively (Table 1, Fig. 2B). Despite this difference in body weight, there was no significant effect of parental relatedness on female fecundity at 53 days, after insemination (Table 1, Fig. 2C). In contrast, we found a strong treatment effect on the probability of having produced eggs by self-fertilization at the age of 50 days. This probability was the highest in cousin-cross treatment and it decreased by 77% and 35% in the selfing and sib-cross treatments, respectively (Table 1, Fig. 2D).

We genotyped F_2 individuals, which represent inter-line crosses in our design (their offspring are F_3), using microsatellites fixed for alternative alleles in the two inbred lines. As expected, almost all F_2 were fully heterozygous, with rare exceptions where one or two loci appeared homozygous in a few individuals, likely due to genotyping errors (Fig. 3). In the F_3 generation, we found no significant differences in heterozygosity among mating treatments ($X^2_2 = 0.475$, $P = 0.789$, binomial model on the proportion of heterozygous loci). The multilocus heterozygosity distribution within each treatment was centered around 0.5, consistent with expectations for $F_2 \times F_2$ progeny: selfing ($H_O = 0.51 \pm 0.05$), sib-cross ($H_O = 0.49 \pm 0.16$), and cousin-cross ($H_O = 0.52 \pm 0.16$; Fig. 3).

We conducted simulations replicating our experimental design to determine whether the observed differences between treatments could result from incomplete fixation, linkage, or spontaneous mutation. The mutational parameters were selected to align with previous estimates of ID in natural, outbred populations of *Physa acuta*[30]. The simulations showed that offspring fitness from full-sib and cousin crosses was only slightly higher than that from the selfing treatment, with differences of 0.8% and 2.0%, respectively—well below the observed differences in juvenile survival, body weight,

and the probability to lay eggs in isolation (Fig. 4). We tested various parameter sets

compatible with naturally observed levels of ID, adjusting linkage, contributions of

lethal mutations, and selection or dominance coefficients, but these had minimal effects

on the results. The largest predicted difference in fitness between cousin-cross, sib-

cross, and self-fertilization occurred when the source of ID was driven quasi-exclusively

by mutations to semi-lethals (d_1=0.49, $d2$=0.01) and when these mutations were nearly

fully lethal (s_1=0.99). However, even under these extreme conditions, average fitness

differences never exceeded 5% (see Fig. S7). While individual replicate runs varied

considerably within each parameter set, the mean predictions consistently fell well

below the observed ratios in all situations. Simulations could match the experimental

results only when the mutation rate was artificially increased by approximately 10- to

12 fold beyond the levels of ID estimated for natural populations of *P. acuta* (Fig. S8).

DISCUSSION

Our study reveals substantial inbreeding depression (ID) in the progeny of

experimental lines, characterized by distinct levels of parental relatedness. Three fitness

traits —juvenile survival, body mass, and self-fertility—declined as parental relatedness

increased (self<sibs<cousins), while female fecundity showed no significant

differences. The amount of ID in juvenile survival (0.18) and body mass (0.20) in the

selfing treatment, relative to the cousin cross, is lower than typical values reported for

outbred and self-fertilized offspring of wild-caught *Physa* acuta, where ID estimates are

approximately 0.55 for juvenile survival[30,31] and 0.38 for body mass[31]. Comparable data

for self-fertility are not available. Despite these lower ID values, the levels observed in

our experiment—amounting to one-third to one-half of those in natural outbred

populations for key traits—are striking and somewhat paradoxical given the unique

design of our study: all three types of offspring have identical genomic homozygosity.

During the establishment of the inbred lines, deleterious mutations have been either

eliminated or fixed, resulting in identical levels of heterozygosity across all F_3

categories (selfed, sib-cross, and cousin-cross)—all expected to retain half the

heterozygosity of the genetically uniform F_2 . Under mutation-based ID theory, we

would predict similar average fitness across treatments, yet our results clearly indicate

otherwise.

One potential explanation for this result is that some F_2 or F_3 individuals may

have failed to successfully cross-fertilize, possibly due to a loss of cross-fertilization

ability after 28 generations of enforced selfing. However, two key arguments dispel this

possibility. First, additional crosses confirmed that these individuals could both fertilize

and be fertilized by their partners (see Supporting Information: *Outcrossing ability of*

inbred lines). Second, microsatellite genotyping supported the expected patterns, with

F_2 individuals showing full heterozygosity for the diagnostic markers. Occasional

apparent homozygosity at one or two loci is likely due to reading errors and does not

indicate selfed offspring, which would be fully homozygous like their parents.

Moreover, observed microsatellite heterozygosity did not differ among the three classes

of F_3 offspring, and was consistently at 0.5, as expected. Note that in classical ID

experiments, self-fertilized broods typically exhibit half the heterozygosity of

outcrossed ones.

In the absence of differences in genomic heterozygosity, what explains the

observed ID? A first hypothesis is that the reduced fitness in the selfing treatment may

stem from maternal effects[32]. However, such maternal effects in this case would not be

linked to variation in maternal quality, as mothers were randomly assigned to all

treatments. Instead, they might reflect a parental component of ID, where less energy is

allocated to self-fertilized eggs compared to outcrossed ones (e.g.[20]). While plausible,

this argument fails to account for the fitness differences between the sib- and cousin-

cross treatments. Moreover, fitness traits in *Physa acuta*, including juvenile survival, are

not known to be strongly influenced by maternal effects. For instance, outcrossed

offspring of inbred mothers do not exhibit lower fitness than offspring of outbred

mothers (*cf.*[14,31]), even though the inbred mothers themselves are less fit than outbred

ones.

Alternatively, the observed variation in ID among mating treatments could result

from the long-term persistence of localized heterozygosity in specific genomic regions

within our inbred lines. This may occur due to linkage disequilibrium among deleterious

mutations, constituting pseudo-overdominant blocks[10]. Such blocks would not be

detected by the seven microsatellites used in our study, which represent only a small

fraction of the genome. However, our simulations do not support the hypothesis that

pseudo-overdominance, driven by tightly linked deleterious alleles, could persist or

accumulate over the 25+ generations of extreme inbreeding experienced by our lines.

Specifically, our simulations show that mutations generate negligible fitness differences

among offspring from selfing, sib-, and cousin-crosses. Even when we artificially

increased linkage by reducing the genome size to the unrealistic value of 1 Morgan

(despite *Physa acuta* having 19 pairs of chromosomes), the resulting fitness ratios

remained virtually unchanged. This suggests that linkage disequilibria among mutations

are insufficient to maintain heterozygosity in highly inbred lines such as those used in

our study. In contrast, simulations suggest that fitness contrasts are maximized when

inbreeding depression arises from semi-lethal mutations. While these mutations may not

persist for long, they can occasionally be present in a heterozygous state in the parents

at the time of experimentation. It is important to note that lethal mutations are lost at a

rate of one-third per generation in selfing lines. Indeed, homozygous lethal individuals

are unviable, so that among viable offspring produced by a heterozygote, one-third are

homozygous wild-type, and two-thirds heterozygous. As a result, the half-life of lethal

mutations in selfing lines is approximately 2-3 generations. Heterozygosity resulting

from mutations with milder effects has an even shorter half-life, as homozygous

mutants are not completely eliminated from the population. Thus, in long-established

selfing lines, the heterozygous load is likely to reflect only a few generations (2-3) of

accumulated spontaneous mutations.

Even in the most favorable scenario, which assumes an unusually high

proportion of lethal mutations, our findings remain highly unlikely. In our reference

scenario, we modeled an average per-haploid genome rate of spontaneous semi-lethal

mutations of $U_1 = 0.028$. Although direct estimates of U_1 are unavailable for *Physa*

acuta, this value is close to those reported for *Drosophila melanogaster*: 0.003 to 0.017

for the large chromosomes 2 and 3, and 0.006 to 0.034 for the full genome[33–35]. Across

all simulations, cousin-crossed or sib-crossed offspring consistently showed, on

average, a small but positive viability advantage (<5%) over self-fertilized offspring.

This aligns with the persistence of spontaneous mutations over two or three generations,

as discussed earlier. However, this predicted advantage is far smaller than the observed

differences in juvenile survival. Notably, fitness ratios across simulations exhibit

significant stochastic variation (Fig. 4), particularly when lethal mutations are frequent.

This variability is expected, given the stochastic nature of recent mutations in the lines

just a few generations before the experiment. While the observed survival estimates

remain improbable, a small fraction of simulations occasionally yield survival ratios

that are similar to those observed: 3.8 % for Sibling/Selfing, 0.8% for Cousin/Selfing,

and 6% for Cousin/Sibling (Fig. 4). To definitively rule out a purely mutational origin

for these survival differences, replication with independent line pairs would be

necessary.

Body size and the proportion of self-fertile individuals at the age of 50 days also

showed clear signs of ID, with ratios even higher than those observed for juvenile

survival. These traits expressed later in life, typically contribute less to overall ID than

juvenile survival in wild-caught *Physa acuta* and are less susceptible to purging [14,30,31].

This suggests that the genetic load affecting these traits is more strongly influenced by

small-effect alleles than by semi-lethal mutations. Consequently, simulating these traits

may require parameter adjustments, such as lower per-generation genomic mutation

rates and a higher proportion of small-effect mutations. However, such adjustments

would inevitably reduce residual heterozygosity within the lines, further decreasing all

ratios and pushing simulated values even farther from the observed results.

Taken together, the unexpectedly high ID observed across three independent

traits in our experiment is highly improbable under a mutational genetic model. To

reconcile our findings with such a model, one would need to assume a strong (> 10-

fold) increase in mutation rates in the inbred lines compared to the original populations

(see Fig. S8). While there is evidence that mutation rates can increase in individuals in

poor condition, as may be the case in our inbred lines, the magnitude required remains

implausible. For instance, studies in *Drosophila* have shown that individuals carrying a

load of fixed deleterious mutations—those in poor condition—tend to accumulate

spontaneous mutations at higher rates than healthier individuals[35,36]. However the

observed increases in mutation rates were modest (up to 2-fold) and did not extend to

lethal mutations[35].

Our results therefore suggest either an increase in mutation rates much higher

than previous measures, or a source of fast-renewed ID unrelated to mutation.

Differences in fitness-related traits could instead arise from the accumulation of

deleterious epimutations, such as DNA methylations, small RNAs, and histone

modifications. For instance, in maize, inbred lines exhibit a persistent decrease in fitness

alongside a consistent rise in genomic methylation over generations of inbreeding[17],

likely due to disruptions in methylation resetting mechanisms. However, this is unlikely

in our study since the founders (F_2) of the experimental lines were all highly outbred

individuals, a condition that would typically restore the overall methylation level back

to normal[17]. Instead, our findings support the hypothesis that the observed ID may be

driven by epigenetic similarity between uniting gametes.

Epigenetic marks can be transmitted across generations with incomplete fidelity,

suggesting that this information is stored within gametes through mechanisms that

remain poorly understood (but see[37], and references therein for the role of small RNAs

in epigenetic inheritance). Two gametes produced by the same individual are expected

to be more similar in their epigenetic information than two gametes produced by full-

sibs, while the latter are more similar than those of cousins. This gradient of similarity

arises from the increasing number of meiotic cycles, along with subsequent zygote

development and epigenetic resetting events, separating paternal and maternal gametes

from selfing to sib-cross to cousin-cross (Fig. 1).

Although no direct evidence currently supports this hypothesis, some studies

suggest that increased fitness (heterosis) can result from epigenetic dissimilarity

between parents, as observed in the crossing of isogenic epi-RILs in *Arabidopsis*[25]. In

Physa acuta, as well as in other mollusks, parental environmental history can influence

offspring's traits through trans-generational plasticity (e.g.,[38–40]). In *Biomphalaria*

glabrata, a freshwater pulmonate gastropod closely related to *Physa*, DNA

methyltransferase inhibitors have been shown to disrupt methylation patterns, leading to

changes in growth and gene expression that persist in the untreated progeny of exposed

individuals[41]. Such environmental effects may contribute to epigenetic differences in

gametes produced by different individuals. More broadly, considering that epimutations

are generated at higher rates than genetic mutations[29], it is plausible that a fraction of the

standing ID originates from rapidly generated epigenetic changes. These changes,

transmitted over one or a few generations, could lead to a decline in performance as

parental relatedness increases.

In summary, our study demonstrates that inbreeding depression (ID), typically

characterized by reduced offspring fitness with increasing parental relatedness, can

appear even in genetically uniform populations. This finding challenges traditional

genetic models, as in such contexts individuals cannot differ in genomic heterozygosity.

The levels of ID we observed, impacting both juvenile and adult traits, are equivalent to

a substantial fraction of the ID typically reported in natural polymorphic populations for

the same traits. Although spontaneous mutations continuously arise and create minor

genetic differences among individuals, these mutations alone seem insufficient to

account for the magnitude of effects we observed across three distinct traits. By

definition effects that are not due to genetic differences, should be called epigenetic,

irrespective of their material basis. While genetic mutations undoubtedly contribute to

ID in populations, it would be premature to assume they account for the entirety of its

effects, particularly given the notorious difficulty of fully eliminating ID through purge

or fixation. Our findings suggest that epigenetic mechanisms may play a more

prominent role in ID than previously recognized – in line with emerging evidence that

epimutations and epigenetic inheritance contribute to heritable phenotypic variation in

general. Future studies, encompassing a broader range of lines and species, are essential

to fully unravel the role and impact of epigenetics to inbreeding depression.

METHODS

Study organism and experimental conditions

Physa acuta is a widespread simultaneously hermaphroditic freshwater snail that

reproduces mostly through cross-fertilization. Snails that have no access to mates can

resort to self-fertilization[42,43]. When a potential mate is present, however, individuals

nearly always cross-fertilize[44,45]. These snails mate frequently, with unilateral

copulations (*i. e*., they adopt either the male or the female sex role during a given

copulation[46]), and can store sperm for long periods. Sexual maturity is reached in

around 6 weeks under laboratory conditions at 25 °C.

Throughout the experiments—and also during the establishment of experimental

lines—snails were maintained under standard laboratory conditions (25 °C, a 12:12

photoperiod, water renewed twice a week, and *ad-libitum* food in the form of boiled

ground lettuce). Each individual was kept in a 75 mL plastic box. In addition, albino

lines of *P.acuta* are maintained in our laboratory. Albinism (a single-locus recessive

trait), can be used to assess paternity success of wild-type focal individuals because

wild-type offspring produced by albino mothers must have a wild-type father.

Field sampling and breeding design

The first step was to collect snails from the field and generate two inbred lines through

continuous self-fertilization over many generations. Self-fertilization was enforced by

keeping mature individuals isolated until they lay eggs, usually two or three weeks after

the age at which they normally start laying eggs when they have mates available[47]. The

two ancestral wild-caught snails were part of a larger sample taken from a wild

population at the "Pont Romain" in the Lirou river (43°43'47.0"N, 3°49'50.4"E), located

close to the village of Les Matelles, 15 km north of Montpellier (France) on October 23,

2013. The initiation of 44 inbred lines is described in Janicke et al.[48], who used them for

an experiment after the third generation of self-fertilization. A subset of the lines were

propagated by continuous self-fertilization until May 2019, a time at which they had all

reached at least 25 generations of selfing. The inbreeding coefficient at these stage is

therefore expected to be, at least, $1-1/2^{25} = 0.99999997$. To evaluate whether inbred

individuals were still capable of cross fertilization, we performed a preliminary analysis

described in Supporting Information. Once confirmed that individuals from inbred lines

were fully capable of performing cross-fertilization, we conducted the main assay.

Estimating inbreeding depression in outbred clones

We selected two inbred lines named 120 and 110. At this stage, these lines had

undergone 28 and 29 generations of self-fertilization, respectively. The residual

heterozygosity is expected to be $< 4.10^{-9}$ times that of the wild-caught individuals,

ignoring recent mutations. As a first approximation, we therefore consider inbred lines

as fully-homozygous clones. The pedigree and the mating protocol described below are

depicted in figure 1 and in figures S1 to S6 from the appendix.

First step: obtaining of F$_2$ snails

One individual from each inbred line, 120 and 110, was isolated and forced to self-

fertilize to produce the F_1. F_1 offspring were isolated at day 28 after egg-laying (still

immature) and then raised in isolation until maturity for three more weeks. We assigned

a code letter to the F_1 siblings for identification (120_a, 120_b, 120_c, and so on; and, 110_a,

110_b, 110_c, and so on), constituted inter-line pairs of F_1 individuals (e.g. 120_a x 110_a,

120_b x 110_b), and let them mate for one week. To differentiate the 120 snail from its 110

partner in each pair, we added small dots of harmless car-body paint of different colors

on their shell[49]. After mating, F_1 individuals were re-isolated to lay F_2 eggs. We

successfully collected and raised the resulting F_2 hatchlings for 10 families *a* to *j* (Fig.

1).

F_2 snails belonged to two different types, hereafter type S and type T. The S-type

F_2 offspring were laid by a mother from the 120 line, and sired by fathers from the 110

line, while T-type offspring came from the reciprocal combination (mother 110 – father

120). As these two types are present within all 10 families *a* to *j*, we end up with 20

categories (S_{a-j} and T_{a-j}), each represented by several offspring. Note that irrespective of

the category they belonged to, all F_2 were expected to be heterozygous and genetically

identical to one another (i.e. they form a heterozygous clone; Fig. 1 and S1). We

proceeded until we obtained at least 40 F_2 descendants of type S and 10 of type T for

each family. This procedure sometimes required taking several successive clutches or

even re-mating the same pair to stimulate egg-laying. These descendants were raised

and isolated before maturity, as previously mentioned, and kept until fully mature and

ready for mating. Then, we used them to generate different types of F_3 snails.

Second step: self-fertilization, sib-, and cousin pair-crosses

We submitted each F_2 snail to one of three mating treatments: (i) selfing, (ii) sib-cross,

and (iii) cousin-cross. Sib and cousin crosses were themselves divided in two subtypes

depending on direction of exchange of genetic material (one- and two-ways), giving

five treatments in total. We explain this in more detail in the following.

The first treatment included S-type individuals that were forced to self-fertilize

(Selfing, Figs. 1 and S2). Thus, ten F_2 S-type individuals per family (i.e., $S_{a,1\text{-}10}$, $S_{b,1\text{-}10}$,

..., $S_{j,1\text{-}10}$) were isolated and eggs were collected three days after we detected the first

clutch. The second treatment consisted in pair-crossings between two S-type snails from

the same family (called "one-way sib cross"). So, for instance, in family *a*, we crossed

individuals $S_{a,11}$ … $S_{a,20}$, according to the following pairs: $S_{a,11}/S_{a,16}$, $S_{a,12}/S_{a,17}$, and so on

(and similarly for S_b to S_j) (Figs. 2 and S3). Note that in these crosses the two parents

are full sibs and they have the same mother and the same father. The third treatment

consisted in pair-crossings within the same family (e.g. family *a*) but between S and T

individuals, which we defined as "two-way sib cross". For instance, we paired snails

from S*a* (numbered from 21 to 25) with snails from T*a* (numbered from 1 to 5), which

gave us the following combinations: $S_{a,21}$ / $T_{a,1}$, $S_{a,22}$ / $T_{a,2}$, and $S_{a,23}$ / $T_{a,3}$, and so on.

The two parents in these crosses are full-sibs but not obtained in the same way (the

father of one is the mother of the other and reciprocally; Figs. 2 and S4). The fourth

treatment involved pair-crossings between two F_2 S-type snails (numbered from 26 to

35) from different families, which we called the one-way cousin cross (the two parents

are cousins, belonging to the same type). Because of insufficient individuals in families

h and *j* we were able to do this for four pairs of families. The resulting combinations

were: $S_{a,26\text{-}35}$ / $S_{b,26\text{-}35}$, $S_{c,26\text{-}35}$ / $S_{d,26\text{-}35}$, $S_{e,26\text{-}35}$ / $S_{f,26\text{-}35}$ and $S_{g,26\text{-}35}$ / $S_{i,26\text{-}35}$ (Figs. 2 and

S5). Finally, the fifth treatment consisted in pair-crossings between snails of the S-type

(36 to 40) and of the T-type (6 to 10) from different families, which we defined as the

two-way cousin cross. The combinations were: $S_{a,36\text{-}40}$ / $T_{b,6\text{-}10}$, $S_{c,36\text{-}40}$ / $T_{e,6\text{-}10}$ or $S_{d,36\text{-}40}$ /

$T_{f,6-10}$, and so on (Figs. 2 and S6). When necessary, we added different color dots of car-

body paint on each individual's shell to differentiate snails from each family and origin.

After pairing for three days, mated snails were re-isolated to lay eggs for three

days. We collected two successive F_3 clutches for each F_2 parent, by transferring the

parent to a new box after three days (Fig. S9). These F_3 clutches constitute the

generation on which we evaluated fitness traits.

Third step: assessing and comparing fitness components in F₃ snails

Eggs were counted in all F_3 clutches right after collection and live juveniles were

counted in the boxes 14 days later. We estimated juvenile survival as the ratio of

juveniles over eggs. Next, we randomly selected one of the surviving F_3 individuals per

box and raised it in isolation until age 50 days. At this age we weighed each snail and

determined whether it had initiated egg laying or not). Given that each individual was

kept in isolation, this egg-laying represents self-fertilization. Then, we paired it with a

same-age albino partner for 3 days. Once inseminated, we isolated the F_3 individual

once more, weighed it a second time, and allowed it to lay eggs for 3 days. Finally, we

counted these eggs that represent female fecundity of F_3 under cross-fertilization

conditions. We therefore have four traits on the F_3: juvenile survival, body mass at 50-

53 days (the mean of the two measures), self-fertilization ability at 50 days (yes/no),

female fecundity after insemination at 54-57 days (number of eggs).

Heterozygosity

We genotyped a subset of the F_2 and F_3 individuals of different categories using 7

microsatellite loci chosen because preliminary data indicated that the inbred lines 110

and 120 were fixed for different alleles. The loci were AF108762, AF108764[50], Pac1,

Pac2,[51] Pasu1-2, Pasu1-9, Pasu1-11 (*cf.*[30]) grouped into two multiplexes. We expected

that the F_2 would be fully heterozygous (except for a portion of misreadings), whereas

in the F_3 generation, half of the loci would, on average, exhibit heterozygosity. We

assessed the number of heterozygous and homozygous loci for each individual.

However, it was inevitable that a small fraction of the genotypes remained unamplified,

leading to missing data. Moreover, in microsatellite readings heterozygous genotypes

are frequently misconstrued as homozygous, i.e. one allele behaves as partly

"dominant"[52]. To avoid any bias, we kept the readings blind to the sample's origin.

Also, we refrained from making corrections, even in instances where the reading error

was evident (e.g. one homozygous locus and all other loci heterozygous in a F_2).

Statistical analyses

All analyses were performed with *R* 4.0.2 packages *lme4*[53] and *nlme*[54] by fitting

generalized linear mixed models (GLMMs) for binomial variables (juvenile survival,

ability to self-fertilize, proportion of heterozygous loci) or linear mixed models (LMMs)

for Gaussian continuous variables (body weight, female fecundity). Our linear models

included as fixed effects: (i) mating treatment (selfing, sib-cross, and cousin-cross), (ii)

the cross type (one vs. two ways) when appropriate (i.e. for sib- and cousin-cross), and

(iii) the maternal type (S vs. T) within two-way crosses (recall that in the selfing and

one-way crosses all mothers are S). We proceeded by first running separate analyses

within the sib-mating and cousin-mating treatments in order to test the mating treatment

and maternal type effects. As these effects were not significant, we pooled all

subcategories within the sib-cross and cousin-cross treatments in order to test the

mating treatment effect using the whole dataset. In each model we added an appropriate

structure of random effects to account for various levels of replication within each level

of fixed factors (e.g. identity of the maternal family: *a*, *b*, *c*; pair of parental families:

a/b, *c/d*; identity of the mother: 1 to 35; and/or of the parental pair: 21/1, 22/2 described

for each model in the results). Observation number was also added as a random factor in

the binomial models on juvenile survival in order to account for overdispersion[55,56].

Tests of fixed effects were made using LRT (Likelihood-ratio tests, which compare the

likelihood of models with and without the factor of interest) keeping all random effects

present in the model. Random effects were tested using LRT with the corrections

indicated by Zuur et al.[57], chap. 5, pp. 123–125). Post hoc tests were performed when

the treatment effect was significant using Holm-Bonferroni correction for multiple-

testing to test effect sizes among different treatments.

Simulations

We made a set of individual-based simulations to evaluate whether the observed

changes in fitness between the three type of mating treatments (selfing, sib-cross,

cousin-cross) could be compatible with a purely genetic model of inbreeding depression

(*i.e.* deleterious-mutation based). Each individual genome was modelled as a single

chromosome on which deleterious mutations were located. The deleterious mutations

were of two types: small-effect and large-effect (quasi-lethal), each type with its own

rate, degree of recessivity, and effect. We modelled both the process of extraction from

a large population and the propagation of two inbred lines over 28 generations with the

following events at each generation: mutation, meiosis, recombination, selection, and

drift within each line. Then, we simulated the creation of F_1, F_2, and the three types of

crosses giving the F_3 as in our experiment. The output of each simulation was a set of

three average fitness values for each of the three types of F_3. For each set of parameter,

1000 independent replicates were made to produce expected distributions of the relative

fitnesses of the three categories. The details on the simulations are explained in Support

Information in *Simulation Details*.

ACKNOWLEDGMENTS

We thank Tim Janicke for starting the inbred lines. This work was funded by CeMEB ("Projets de Recherche Exploratoires du CeMEB" grant to project "Epigenetics of inbreeding depression (EPID)" by P. David and C. Grunau) and ANR (FRIDA project, grant ANR-23-CE02-0033-01 to P. David). N. Bonel was partially supported by the "Programa de Financiamiento Parcial de Estadías en el Exterior para Investigadores Asistentes", National Scientific and Technical Research Council CONICET (Res. Nº 558 1236/08; 4118/16).

AUTHORS' CONTRIBUTIONS

Conceptualization, CG and PD; methodology, NB and PD; formal analysis, NB and PD; data curation, NB and PD; writing—original draft preparation, NB and PD; writing—review and editing, NB and PD; supervision, PD; project administration, PD; funding acquisition, PD. All authors have read and agreed to the published version of the manuscript.

REFERENCES

1. Charlesworth, B. & Charlesworth, D. The genetic basis of inbreeding depression. *Genet. Res.* **74**, 329–340 (1999).

2. Charlesworth, D. & Willis, J. H. The genetics of inbreeding depression. *Nat Rev Genet* **10**, 783–796 (2009).

3. Charlesworth, B. & Charlesworth, D. Some evolutionary consequences of deleterious mutations. *Genetica* **102/103**, 3–19 (1998).

4. Simmons, M. J. & Crow, J. F. Mutations affecting fitness in drosophila populations. *Annu. Rev. Genet.* **11**, 49–78 (1977).

5. Agrawal, A. F. & Whitlock, M. C. Inferences about the distribution of dominance drawn from yeast gene knockout data. *Genetics* **187**, 553–566 (2011).

6. Huber, C. D., Durvasula, A., Hancock, A. M. & Lohmueller, K. E. Gene expression drives the evolution of dominance. *Nat Commun* **9**, 2750 (2018).

7. Lande, R. & Schemske, D. W. The evolution of self-fertilization and inbreeding depression in plants. I. Genetic models. *Evolution* **39**, 24–40 (1985).

8. Glémin, S. How are deleterious mutations purged? Drift versus nonrandom mating. *Evolution* **57**, 2678–2687 (2003).

9. Kirkpatrick, M. & Jarne, P. The effects of a bottleneck on inbreeding depression and the genetic load. *The American Naturalist* **155**, 154–167 (2000).

10.   Waller, D. M. Addressing Darwin's dilemma: Can pseudo-overdominance explain

persistent inbreeding depression and load? *Evolution* **75**, 779–793 (2021).

11.   Barrett, S. C. H. & Charlesworth, D. Effects of a change in the level of inbreeding on

the genetic load. *Nature* **352**, 522–524 (1991).

12.   Carr, D. E. & Dudash, M. R. The effects of five generations of enforced selfing on

potential male and female function in *Mimulus guttatus*. *Evolution* **51**, 1797–1807 (1997).

13.   Chelo, I. M., Carvalho, S., Roque, M., Proulx, S. R. & Teotónio, H. The genetic basis

and experimental evolution of inbreeding depression in Caenorhabditis elegans. *Heredity* **112**,

248–254 (2014).

14.   Noël, E. *et al.* Reduced mate availability leads to evolution of self-fertilization and

purging of inbreeding depression in a hermaphrodite. *Evolution* **70**, 625–640 (2016).

15.   Byers, D. L. & Waller, D. M. Do plant populations purge their genetic load? Effects of

population size and mating history on inbreeding depression. *Annu. Rev. Ecol. Syst.* **30**, 479–

513 (1999).

16.   Crnokrak, P. & Barrett, S. C. H. Perspective: purging the genetic load: a review of the

experimental evidence. *Evolution* **56**, 2347–2358 (2002).

17.   Han, T. *et al.* An epigenetic basis of inbreeding depression in maize. *Sci. Adv.* **7**,

eabg5442 (2021).

18.   Charlesworth, D. & Charlesworth, B. Inbreeding depression with heterozygote

advantage and its effect on selection for modifiers changing the outcrossing rate. *Evolution* **44**,

870–888 (1990).

19.   Lande, R., Schemske, D. W. & Schultz, S. T. High inbreeding depression, selective

interference among loci, and the threshold selfing rate for purging recessive lethal mutations.

Evolution **48**, 965–978 (1994).

20.   Cheptou, P.-O. The evolutionary ecology of inbreeding depression in wild plant

populations and its impact on plant mating systems. *Front. Plant Sci.* **15**, 1359037 (2024).

21.   Bossdorf, O., Richards, C. L. & Pigliucci, M. Epigenetics for ecologists. *Ecology*

Letters **11**, 106–115 (2008).

22.   Cheptou, P.-O. & Donohue, K. Epigenetics as a new avenue for the role of inbreeding

depression in evolutionary ecology. *Heredity* **110**, 205–206 (2013).

23.   Biémont, C. Inbreeding effects in the epigenetic era. *Nat Rev Genet* **11**, 234–234 (2010).

24.   Vergeer, P., Wagemaker, N. (C. A. M.) & Ouborg, N. J. Evidence for an epigenetic

role in inbreeding depression. *Biol. Lett.* **8**, 798–801 (2012).

25.   Dapp, M. *et al.* Heterosis and inbreeding depression of epigenetic Arabidopsis hybrids.

Nature Plants **1**, 15092 (2015).

26.   Richards, E. J. Inherited epigenetic variation — revisiting soft inheritance. *Nat Rev*

Genet **7**, 395–401 (2006).

27.   Johannes, F. *et al.* Assessing the Impact of Transgenerational Epigenetic Variation on

Complex Traits. *PLoS Genet* **5**, e1000530 (2009).

28.   Preite, V. *et al.* Increased transgenerational epigenetic variation, but not predictable

epigenetic variants, after environmental exposure in two apomictic dandelion lineages. *Ecology*
and Evolution **8**, 3047–3059 (2018).

29.   Van Der Graaf, A. *et al.* Rate, spectrum, and evolutionary dynamics of spontaneous
epimutations. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6676–6681 (2015).

30.   Escobar, J. S., Nicot, A. & David, P. The different sources of variation in inbreeding
depression, heterosis and outbreeding depression in a metapopulation of *Physa acuta*. *Genetics*
180, 1593–1608 (2008).

31.   Janicke, T., Vellnow, N., Sarda, V. & David, P. Sex-specific inbreeding depression
depends on the strength of male-male competition: inbreeding depression in a hermaphrodite.
Evolution n/a-n/a (2013) doi:10.1111/evo.12167.

32.   McCarthy, T. M. & Sih, A. Relatedness of mates influences mating behaviour and
reproductive success of the hermaphroditic freshwater snail <i>Physa gyrina. *Evolutionary*
Ecology Research **10**, 77–94 (2008).

33.   Fry, J. D., Keightley, P. D., Heinsohn, S. L. & Nuzhdin, S. V. New estimates of the
rates and effects of mildly deleterious mutation in *Drosophila melanogaster*. *Proc. Natl. Acad.*
Sci. U.S.A. **96**, 574–579 (1999).

34.   Charlesworth, B., Borthwick, H., Bartolomé, C. & Pignatelli, P. Estimates of the
genomic mutation rate for detrimental alleles in *Drosophila melanogaster*. *Genetics* **167**, 815–
826 (2004).

35.   Sharp, N. P. & Agrawal, A. F. Evidence for elevated mutation rates in low-quality
genotypes. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 6142–6146 (2012).

36.   Agrawal, A. F. & Wang, A. D. Increased transmission of mutations by low-condition
females: evidence for condition-dependent DNA repair. *PLoS Biol* **6**, e30 (2008).

37.   Cecere, G. Small RNAs in epigenetic inheritance: from mechanisms to trait
transmission. *FEBS Letters* **595**, 2953–2977 (2021).

38.   Luquet, E. & Tariel, J. Offspring reaction norms shaped by parental environment:
interaction between within- and trans-generational plasticity of inducible defenses. *BMC Evol*
Biol **16**, 209 (2016).

39.   Fallet, M., Luquet, E., David, P. & Cosseau, C. Epigenetic inheritance and
intergenerational effects in mollusks. *Gene* **729**, 144166 (2020).

40.   Tariel, J., Plénet, S. & Luquet, É. Transgenerational plasticity of inducible defences:
Combined effects of grand-parental, parental and current environments. *Ecology and Evolution*
10, 2367–2376 (2020).

41.   Luviano, N. *et al.* The methylome of Biomphalaria glabrata and other mollusks:
enduring modification of epigenetic landscape and phenotypic traits by a new DNA methylation
inhibitor. *Epigenetics & Chromatin* **14**, 48 (2021).

42.   Henry, P.-Y., Bousset, L., Sourrouille, P. & Jarne, P. Partial selfing, ecological
disturbance and reproductive assurance in an invasive freshwater snail. *Heredity* **95**, 428–436
(2005).

43.   Jarne, P., Pointier, J.-P., David, P. & Koene, J. M. Basommatophoran gastropods. in
The evolution of primary sexual characters in animals 173–196 (Oxford University Press,
2010).

44.   Pélissié, B., Jarne, P. & David, P. Sexual selection without sexual dimorphism: bateman
gradients in a simultaneous hermaphrodite: sexual selection in a hermaphrodite. *Evolution* **66**,
66–81 (2012).

45.   Noël, E. *et al.* Reduced mate availability leads to evolution of self-fertilization and
purging of inbreeding depression in a hermaphrodite: selfing evolution in a hermaphroditic
snail. *Evolution* **70**, 625–640 (2016).

46.   Wethington, A. R. & Dillon, Jr, R. T. Gender choice and gender conflict in a non-
reciprocally mating simultaneous hermaphrodite, the freshwater snail,Physa. *Animal Behaviour*
51, 1107–1118 (1996).

47.   Tsitrone, A., Duperron, S. & David, P. Delayed Selfing as an Optimal Mating Strategy
in Preferentially Outcrossing Species: Theoretical Analysis of the Optimal Age at First
Reproduction in Relation to Mate Availability. *The American Naturalist* **162**, 318–331 (2003).

48.   Janicke, T. *et al.* Environmental effects on the genetic architecture of fitness
components in a simultaneous hermaphrodite. *Journal of Animal Ecology* **91**, 124–137 (2022).

49.   Henry, P. & Jarne, P. Marking hard-shelled gastropods: tag loss, impact on life-history
traits, and perspectives in biology. *Invertebrate Biology* **126**, 138–153 (2007).

50.   Monsutti, A. & Perrin, N. Dinucleotide microsatellite loci reveal a high selfing rate in
the freshwater snail *Physa acuta*. *Molecular Ecology* **8**, 1076–1078 (1999).

51.   Sourrouille, P., Debain, C. & Jarne, P. Microsatellite variation in the freshwater snail
Physa acuta. *Molecular Ecology Notes* **3**, 21–23 (2003).

52.   David, P., Pujol, B., Viard, F., Castella, V. & Goudet, J. Reliable selfing rate estimates
from imperfect population genetic data. *Molecular Ecology* **16**, 2474–2487 (2007).

53.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models
using lme4. *arXiv:1406.5823 [stat]* (2014).

54.   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Core Team. nlme: linear and
nonlinear mixed effects models. R package version 3.1-131. (2017).

55.   Elston, D. A., Moss, R., Boulinier, T., Arrowsmith, C. & Lambin, X. Analysis of
aggregation, a worked example: numbers of ticks on red grouse chicks. *Parasitology* **122**, 563–
569 (2001).

56.   Browne, W. J., Subramanian, S. V., Jones, K. & Goldstein, H. Variance partitioning in
multilevel logistics models with over-dispersion. *Journal of the Royal Statistical Society* **168**,
599–613 (2005).

57.   Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A. & Smith, G. M. *Mixed Effects*
Models and Extensions in Ecology with R. (Springer New York, New York, NY, 2009).
doi:10.1007/978-0-387-87458-6.

TABLES

Table 1. Summary of means (±SE) and statistical significance of the mating treatment type effects on fitness traits measured on F_3 individuals of the hermaphrodite snail *Physa acuta*. Number of observations are indicated in parentheses. In this table, as in fig.2, we did not distinguish, within ecah mating treatment, the different possible types of crosses (one-way versus two-way, see fig.1) because they were not statistically different. Detailed results are reported in supplementary tables S1-S3.

Variable			Mating treatment	Treatment effect	Treatment comparison	Effect size
	Selfing (SF)	Sib-cross (SC)	Cousin-cross (CC)			
Juvenile survival	0.62 ± 0.03 (93)	0.69 ± 0.03 (193)	0.76 ± 0.02 (153)	$X^2_2 = 15.40$; P **<0.001**	SF vs. CC	-4.04***
					SF vs. SC	-2.74*
					SC vs. CC	-2.33*
Body weight	41.6 ± 5.3 (82)	44.9 ± 2.6 (152)	52.3 ± 1.8 (134)	$X^2_2 = 10.06$; P **= 0.007**	SF vs. CC	-2.98**
					SF vs. SC	-1.94 .
					SC vs. CC	-2.23 .
Fecundity rate	12.0 ± 0.4 (79)	11.5 ± 0.5 (146)	11.3 ± 0.4 (127)	$X^2_2 = 1.53$; P = 0.466	—	—
					—	—
					—	—
Probability of producing selfed eggs at the age of 50 days	0.10 ± 0.06 (82)	0.28 ± 0.05 (152)	0.43 ± 0.07 (134)	$X^2_2 = 19.69$; P **<0.001**	SF vs. CC	-4.25***
					SF vs. SC	-3.09**
					SC vs. CC	-2.0 *

Treatment means (±SE, with) are reported in natural scale for the four traits (proportion of eggs surviving, milligrams, eggs laid per day, and proportion of laying individuals respectively); however statistical models are generalized linear mixed models on the appropriate scale (Binomial for juvenile survival and probability of producing selfed eggs, with logit link; Gaussian for body weight and fecundity, identity link), and include appropriate random effects to represent the structure of our design (not shown here, see methods and detailed results in Supplementary materials and tables S1-S3). Mating treatment effect sizes (estimated mean difference on transformed scale divided by its SD) were tested using post hoc tests with the Holm-Bonferroni correction for multiple testing. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$ after correction.

FIGURES

Figure 1. Pedigree of the F₂ snails used in this study, and how they were crossed to produce the F₃ on which fitness traits were measured. Parents were two individual snails extracted from independent inbred lines (named 120 and 110) derived from wild-caught ancestors throughr 28 or 29 successive generations of self-fertilization in the laboratory, respectively. F₁ individuals (inbred sublines) resulted from self-fertilization of parents, and were also homozygous. The F₁ pair crossing yielded a set of individuals that constituted the experimental strains (F₂ snails), which are all heterozygous and clonal to one another but with different type of origins, S and T. F₂ snails from the S type had their mothers from the 120 line and their fathers from the 110 line, and *vice versa* for those from type T. The experimental strains were constituted by ten families (*a* to *j*) per type of origin.

Figure 2. Fitness traits measured on F₃ individuals of the hermaphrodite snail *Physa acuta*. **A.** Juvenile survival (proportion of eggs that become juveniles alive after 14 days). **B.** Body weight at 50-53 days (the mean of the two measures, before and after pairing the focal snail with a virgin albino partner). **C.** Female fecundity rate after insemination (number of eggs per day produced by the focal snail). **D.** Probability of having produced self-fertilized eggs at 50 days just before pair mating. Mating treatment effect sizes (difference between two types divided by its SD) were tested using post hoc tests with the Holm-Bonferroni correction for multiple-testing (*$P <$ 0.05; **$P < 0.01$; ***$P < 0.001$). SF, selfing; SC, sib-cross; CC, cousin-cross. Black dots are mean values ± 1SE. We did not separate one-way and two-way crosses in the SC and CC category as differences between them were small and not significant (see supplementary tables S1-S3).

Figure 3. Distributions of multilocus heterozygosity for a subset of F_2 and F_3 individuals of different categories using 7 microsatellite loci (AF108762, AF108764, Pac1, Pac2, Pasu1-2, Pasu1-9, Pasu1-11) chosen because preliminary data indicated that the inbred lines 110 and 120 were fixed for different alleles. The microsatellites were grouped into two multiplexes. Overall, F_2 individuals were fully heterozygous (with a few exceptions) whereas in the F_3 heterozygosity was spread around 0.5 within each mating treatment. Red dots are mean values ± 1SE.

Figure 4. Violin plots of the distribution of fitness ratios from 1,000 simulations of our experiment, using the reference parameter set, compatible with available data on ID in *Physa acuta* (see Methods and *Simulations* in Supporting Information). Parameters include the length of the genome ($L = 10$ Morgans), the dominance coefficient and effect size of semi-lethals ($h_1 = 0.02$, $s_1 = 0.9$), and those of small-effect mutations ($h_2 = 0.2$, $s_2 = 0.05$), with both mutation types contributing to a total inbreeding depression of $d=0.5$ in natural populations ($d_1 = 0.3$ for semi-lethals and $d_2 = 0.2$ for small-effect mutations). Black dots represent the mean fitness ratio, with 95% confidence intervals (CI) shown. The observed ratios for juvenile survival (red line), body mass (blue line), and the proportion of fertile individuals at 50 days (green line) were for Sib/Self: 1.11, 1.08, and 2.80 (not shown, out of range); for Cous/Self: 1.23, 1.26, and 4.30 (not shown); and for Cous/Sib: 1.10, 1.16, and 1.54, respectively. Simulations with different parameter sets are in Figure S7 from Supporting Information.