In sync with sound: Interpersonal synchronization as an objective measure of listening engagement

Lotte Lambrechts[1], Bernd Accou[1], Jonas Vanthornhout[1], Bart Boets[2], Tom Francart[1]

Affiliation:

1. Department of Neurosciences, ExpORL, KU Leuven - University of Leuven, Leuven (3000), Belgium
2. Department of Neurosciences, CDP, KU Leuven - University of Leuven, Leuven (3000), Belgium

Corresponding author: Lotte Lambrechts

Email: lotte.lambrechts@kuleuven.be

Author Contributions: L.L., B.A., J.V., B.B. and T.F. conceptualized the study. L.L. and B.A. collected the data. L.L. and B.A. analyzed the data. L.L., B.A., J.V., B.B. and T.F. interpreted the data. L.L. wrote the first draft of the manuscript. L.L., B.A., J.V., B.B. and T.F. reviewed and edited the manuscript.

Abstract

Listening engagement plays a crucial role in effective communication and knowledge transfer, reflecting a state of deep absorption in an auditory stimulus. The available behavioral assessments of listening engagement are limited in capturing its multidimensional nature. Measuring listening engagement objectively using brain and body signals is a compelling alternative. In particular, interpersonal synchronization—observed through synchronized physiological responses among listeners—may offer a promising objective measure. In this study, we examined synchronization of brain activity, heart rate, and electrodermal activity between participants while they listened to engaging and non-engaging stories. Subjective engagement ratings were collected as a ground truth reference. Results showed higher interpersonal synchronization across all modalities when participants listened to engaging stories compared to non-engaging ones, with significant correlations between synchronization measures and ratings of subjective engagement. These results confirm that synchronization may be a reliable, objective marker of listening engagement. We anticipate that these results will provide a powerful framework for future research in relevant domains, including education, audiology, and communication science, and enable impactful real-world applications, such as automatically tracking audience engagement in education, entertainment, and broader societal contexts.

Significance Statement

We provide the first direct empirical evidence that interpersonal physiological synchronization serves as a robust and reliable marker of listening engagement. Using a novel experimental paradigm with ecologically valid auditory stimuli, we demonstrate that heightened engagement leads to increased neural and physiological synchronization among listeners, while disengagement reduces it. This objective measure significantly correlates with established behavioral assessments, validating its relevance and reliability. Our innovative approach offers a powerful framework for future research in the relevant domains (education, audiology and communication

40 science) and enables impactful real-world applications, such as automatic tracking of audience
41 engagement in education, entertainment, and broader societal domains.
42
43 **Main Text**
44
45 **Introduction**
46
When listening to an auditory stimulus, the brain processes and interprets sound, tracking its
complex acoustic dynamics(1–3). Neural tracking of speech has been established as a reliable and
valid measure of natural speech processing and perception, allowing for an objective assessment
of speech understanding (e.g.,(3, 4)). Moreover, neural tracking is modulated by attention to
speech (e.g.,(5, 6)). Recent evidence suggests that whether people are engaged with speech is
represented not only in the brain but also in physiological signals throughout the body(7). This
paper further explores listening engagement and aims to delineate objective markers.

Listening engagement is a complex, multifaceted construct, defined as a state of deep listening
where a person pays attention to a stimulus and is completely absorbed by it (8). It involves a
cognitive and emotional connection to a sound(9) and may also involve narrative presence (8, 10),
a state where listeners lose awareness of their surroundings and feel immersed in the narrative
world. Listening engagement yields several beneficial outcomes: It is linked to enhanced speech
processing and communication, enhanced memory retention (11, 12), listening enjoyment (9, 13),
and the experience of a state of flow (8, 14).

Thus far, listening engagement has mostly been evaluated through behavioral self-report rating
scales (8, 10, 15), such as the 'Narrative absorption scale' and the 'Narrative engagement scale'.
These well-validated measures can reliably capture variability in listening engagement (16).
Nevertheless, behavioral self-report measures hold inherent limitations, as they may disturb the
immediate experience and induce biases (e.g., socially desirable responses when students have
to rate their engagement with a lecture). Additionally, they provide only limited temporal resolution,
failing to capture moment-to-moment fluctuations in engagement. To address these limitations, an
objective measure of listening engagement could provide valuable insights by minimizing bias and
allowing continuous assessment. Specifically, we will explore how brain and body responses may
serve as reliable indicators of listening engagement.

Previous research has shown that when people consciously process engaging stories, their
physiological responses tend to synchronize (7, 17, 18). Interpersonal synchronization occurs
across multiple physiological modalities, including brain activity, heart rate (HR), and electrodermal
activity (EDA). However, there remains considerable uncertainty regarding the drivers of
interpersonal synchronization. While it has been observed in direct conversational interactions(19),
it also occurs when individuals (jointly) passively listen to the same auditory stimulus (7, 17, 18).
Notably, interpersonal synchronization has been found even when participants listen separately to
the same stimulus at different moments (7, 17), suggesting that synchronization is not uniquely
driven by direct social interaction (20, 21), but rather by shared cognitive and emotional
engagement with the content. However, of course, social interactions may amplify these effects.

Various mechanisms have been proposed to explain interpersonal synchrony. Bevilacqua et al.
(13) and Madsen & Parra (7, 17) suggest that it arises from the conscious and attentive processing
of an engaging stimulus, driven by shared neural oscillations and cognitive processes in response
to the content. For example, attentive listening to an engaging stimulus modulates brain activity in
a way that aligns across individuals, leading to synchronization. Pérez et al. (18) support this view,
proposing that HR synchronizes across participants in response to cognitive processes when
individuals listen to the same stimulus. However, other studies suggest that HR may also be
influenced by emotional responses elicited by the stimulus, suggesting a broader interplay among
cognitive and affective factors (22). Given that bodily responses such as HR and EDA are closely

linked to emotions and arousal (23–25), emotional processing likely plays a role in listening
engagement alongside cognitive mechanisms.

To further understand the relationship between listening and interpersonal synchronization,
Madsen and Parra (7, 17) showed that synchronization decreased when participants were
distracted from listening through a cognitively demanding task (distracted condition). This suggests
that synchronization may be influenced by attention. Despite these interesting findings, no
conclusions regarding the relationship between synchronization and listening engagement can be
validated, as they only manipulated attention and not listening engagement. In addition, previous
studies have either lacked a behavioral ground truth (7, 16, 17) or used behavioural measures not
directly relevant for engagement (26), making it challenging to ascertain whether the auditory
stimulation was genuinely engaging.

The present study aims to address these gaps by employing an objective synchronization measure
to distinguish between engaging and non-engaging conditions and to examine the relationship
between synchronization and listening engagement. To overcome limitations in prior studies, we
meticulously selected engaging and dull stimuli for the participant group and incorporated a
behavioral measure of engagement. We hypothesize that interpersonal synchronization will be
higher when individuals are engaged with the stimulus and that synchronization levels will predict
self-rated listening engagement.

Results
109
We built a test setup and designed an experimental paradigm to investigate the relationship
between listening engagement and interpersonal synchronization. 40 participants were exposed to
three auditory conditions: engaging (listening to an engaging podcast), nonengaging (listening to a
nonengaging recording), and distracted (listening to a nonengaging recording while solving riddles).
Brain activity, EDA, and HR were recorded. Physiological synchronization was assessed via
intersubject correlations (ISC) across modalities. Behavioral engagement was measured through
a newly developed Listening Engagement Scale, based on validated tools, along with content-
based multiple-choice questions. Statistical analyses included linear mixed-effects modeling to
assess condition effects and Spearman correlations between physiological synchronization and
behavioral ratings.
120

<u>Participants rate engaging stimuli higher than non-engaging stimuli on the behavioral Listening</u>
<u>Engagement Scale</u>

Figure 1 (bottom, right panel) contains the averaged (across all questions) subjective engagement
ratings for the different conditions. Due to system failure, we lost responses for the first engaging
condition of one participant, the first non-engaging condition of three participants, and the second
non-engaging condition of 4 participants.
The mean listening engagement score for the engaging condition was 4.8. The mean score for the
non-engaging stimuli was 2.1. The linear mixed effects model showed highly significant differences
in subjective engagement between both conditions (β = -2.70, SE = 0.098, t = -27.73, p < .001).
130
<u>Participants display higher interpersonal synchronization when listening to an engaging stimulus</u>

The top three panels in *Figure 1* show synchronization of physiological responses between
individuals while listening to auditory stimuli with varying engagement levels. Statistical significance
of each data point was calculated using circular shifting.

A highly significant effect of condition could be identified for all physiological responses, based on
ANOVA comparisons with and without condition: brain synchronization ($\chi^2(2)$ = 158.24, p < .0001),
HR synchronization ($\chi^2(2)$ = 119.44, p < .0001), EDA synchronization ($\chi^2(2)$ = 73.392, p < .0001).

Post-hoc comparisons with Holm correction (**Table 1**) revealed significantly higher synchronization
in the engaging condition (E) than in the non-engaging (N) and distracted condition (D) for all
physiological responses. Additionally, brain synchronization was higher in the non-engaging
condition than in the distracted condition, but this effect was not observed for HR and EDA.

We also looked at neural tracking results. Significant neural tracking could be found for each
condition, but not for every participant, especially in the distracted condition. Using a similar
analysis approach as for the synchronization measures, we identified a significant effect of
condition for neural tracking ($\chi^2(2) = 45.033$, $p < .0001$). Post-hoc comparisons showed significant
neural tracking differences between the three conditions (**Table 1**).

<u>Relation between physiological synchrony measures and behavioral engagement ratings</u>

We analyzed the relationship between the physiological and behavioral data. *Figure 2* presents
scatter plots depicting the association between the different interpersonal synchronization values
and the scores on the Listening Engagement Scale. Each correlation plot includes data for all
participants and conditions involving attentive listening (excluding the distracted condition, as we
did not assess any subjective ratings). The first panel contains a scatter plot for brain
synchronization, the second for HR synchronization, and the third panel for EDA synchronization.
All correlations were significant, indicating a clear relationship between objective synchronization
measures and subjective engagement ratings across all modalities. For the different body
responses, the scatter plots indicate that higher engagement ratings correspond to higher
physiological synchronization between participants' physiological responses.

Additionally, we conducted a PLS regression using physiological responses as predictors and
behavioral data as the outcome variable. Based on three predictor variables, the regression model
explained 88% of the variance in behavioral rating outcomes. To assess how well the behavioral
ratings could be reconstructed from these physiological responses, we used Spearman's
correlation to compare the reconstructed data with the Listening Engagement Scale results. This
analysis revealed a strong correlation of 0.68 between the PLS measure and the engagement
scores, indicating a meaningful relationship between physiological responses and self-reported
engagement (see *Figure 2 - last panel*). The observed correlation was higher than any of the
individual synchronization results; however, no statements can be made regarding significance.

Discussion
168
This paper aims to evaluate interpersonal synchronization as a measure of listening engagement.
Twenty duos were invited to our lab and listened to auditory stimuli varying in engagement level
(ranging from engaging/interesting to non-engaging/uninteresting. Meanwhile, we recorded
physiological brain and body responses. As an extra control condition, participants also completed
a distracting task while listening to the stimuli. Research indicates that physiological
synchronization between individuals may be a valuable indicator of listening engagement.

For each of the physiological modalities and across all conditions, significant interpersonal
synchronization was observed when people were presented with an auditory stimulus. This finding
supports past results that link interpersonal synchronization to conscious speech processing (7,
16, 17). Our results show that synchronization was higher when participants listened in an engaged
manner and significantly lower when the stimulus was non-engaging. This pattern was consistent
across all the brain and body signals (EDA and HR), suggesting that engagement activates
additional mental processes reflected in physiological responses (27, 28). These processes might
drive interpersonal synchronization across the physiological responses. However, identifying the
exact cause of this effect is challenging and requires further research. Nevertheless, our findings
demonstrate that synchrony does not merely result from attentive listening, as participants were
encouraged to listen attentively in both the engaging and non-engaging conditions. Even under

these conditions, synchronization decreased significantly in the non-engaging condition,
suggesting that engagement is driven by an involuntary process, beyond mere attention.

We also saw that synchronization declined when participants were distracted compared to when
they listened attentively, aligning with previous studies (7, 17, 18). Through brain responses,
including neural tracking and brain-to-brain synchronization, we were able to distinguish between
non-engaged listening and distracted listening, indicating that engagement extends beyond
attention and conscious processing, which can be captured in measures of synchronization (7, 16,
17). These findings support the idea that synchronization can be used as a proxy for listening
engagement. However, synchronization did not significantly differ between non-engaged and
distracted listening conditions for HR and EDA.

Additionally, we examined the relationship between self-rated listening engagement and objective
interpersonal synchronization data. Results revealed strong and consistent correlations between
subjective ratings and objective synchronization measures across all three physiological
responses. This supports the hypothesis that synchronization can predict listening engagement at
the individual subject level.

PLS analysis revealed a strong relationship between a combined physiological measure and
behavioral engagement scores. Although listening engagement is a highly individualized,
subjective, and hard-to-capture cognitive-affective construct (8, 9), interpersonal synchronization
across these very basic objective physiological measures seems to relate strongly to it. Further
analysis could determine whether a single measure, a subset, or the full combination is most
effective. At present, the combined measure exhibits the highest correlation, outperforming
individual measures. Future research with larger datasets and more advanced models could further
refine these findings. Ultimately, these findings could inform the development of sensors and
systems capable of automatically monitoring and tracking audience focus and engagement, with
promising applications in society, education and entertainment.

While encouraging, these results stem from a deliberately sharp contrast in stimulus types. To
maximize the contrast between engaging and non-engaging stimuli, we selected content that was
either highly interesting or very dull. However, in everyday life, auditory input often falls along a
more nuanced spectrum. Future studies should incorporate a broader range of engagement levels
to determine whether a continuous gradient in synchronization can be observed.

Listening engagement is a multidimensional construct influenced by various factors, making it
challenging to control all potential confounds. For example, different speakers were used across
the conditions, and individual preferences for voice timbre could have affected engagement. To
partly control for this, we ensured all speakers were female, eliminating potential effects of the
speaker's sex. However, factors such as fatigue, anxiety, and personal interest (8, 9, 29) may also
play a role. Future research should also further investigate how speaker engageability
characteristics (e.g., speech rate, prosody, vocal quality) influence listening engagement.

Another key challenge is differentiating between listening engagement and auditory attention. In all
active listening conditions, participants were instructed to listen and focus attentively to the
presented stimulus. To ensure compliance, participants were informed beforehand that they would
be asked to answer multiple-choice questions. While this design theoretically ensures comparable
attention levels across both listening activities, engaged and attentive listening may be more
intricately linked in practice. As a result, it remains unclear whether the observed synchronization
differences between conditions are driven purely by engagement or variation in attention. Future
studies should further disentangle these overlapping constructs.

Finally, through future analyses, we will examine dyadic effects within these data to gain deeper
insights into interpersonal synchronization dynamics and explore whether synchronization is
enhanced by physical proximity and real-life presence.

Conclusion

In this study, participants listened to auditory stimuli while we measured their brain activity, HR, and EDA. The stimuli were designed to be either engaging or non-engaging. Our analysis revealed that interpersonal synchronization was significantly higher in the engaging condition across all physiological responses. Additionally, individual differences in interpersonal synchronization closely correlated with subjective listening engagement scores. These findings strongly suggest that interpersonal synchronization can serve as an objective marker of listening engagement.

Materials and Methods

Participants

Participants subscribed to the experiment in dyads. They came to the lab and participated with someone they knew well (e.g., friends, romantic partners, or family members). We included 40 participants, with a mean age of 21 years (range: 18 to 25 years). Of these, 4 were male. All participants had normal hearing, confirmed by a digit triplet screening test (SRT = -10 dB SNR) (30) and/ or a tonal audiogram (thresholds < 25 dB HL at all audiometric frequencies), with a mean Fletcher index of 13 ± 6 dB HL. Exclusion criteria were attention or concentration problems, neuropathological disorders, communication disorders and language disorders. The Medical Ethics Committee UZ Leuven/Research (KU Leuven) approved the study with reference S68492.

Setup

Testing took place in dyads in a soundproof room (Faraday cage). Participants sat next to each other, approximately 1.35 meters apart, facing each other at a 90° angle (chairs turned 45° towards the loudspeaker). A loudspeaker and monitor were positioned 1.5 meters in front of the participants. The setup was calibrated, and stimuli were presented at 60 dB A, equivalent to an everyday conversation level. See *figure 3* for a full overview.

Procedure

This study measured brain and physiological measures and subjective listening engagement in participants exposed to various auditory stimuli.

Experimental conditions. The experiment consisted of **three different conditions**, each involving the presentation of an auditory stimulus:

1. Engaging condition: Participants listened attentively to a highly engaging stimulus.
2. Non-engaging condition: Participants listened attentively to a stimulus designed to be less engaging.
3. Distracted condition: Participants completed cognitively demanding riddles while an auditory stimulus played in the background, discouraging attention and engagement.
Participants were instructed to listen attentively to the speech stimulus in the engaging and non-engaging conditions. Engagement levels were manipulated by presenting stimuli with varying degrees of intrinsic interest. The **engaging stimulus** was a 46:29-minute episode of a popular Flemish podcast. This stimulus was expected to be inherently interesting, easy to follow, and able to capture individuals' attention ('De Volksjury' episode 19 'Maddie McCann', ranked second place on "Spotify Wrapped" of 2024). The **non-engaging stimulus** comprised self-recorded excerpts from Dutch debates in the European parliament (25:19 minutes duration). This content was chosen for its neutral/dull and less captivating nature. The engaging and non-engaging stimuli were split into two parts, so we had two engaging and two non-engaging conditions. After both the engaging and non-engaging conditions, we assessed self-perceived listening engagement, as will be discussed under 'Behavioral measure - p. 6'. For the **distracted condition,** the participants listened

to another non-engaging stimulus (a 12:42-minute self-recorded excerpt of the European
Parliament) while solving riddles. This condition was designed to minimize both attention and
engagement, serving as a control to differentiate synchronization as a measure of engagement
from mere attentional effects.

The full experimental procedure and order of the protocol can be found in *Figure 3*. Each dyad
followed the same protocol, with the order of conditions (engaging, non-engaging, and distracted)
randomized across dyads to mitigate potential training or fatigue effects.

Brain and physiological measures

We recorded three objective measures: brain activity via electroencephalography (EEG), EDA, and
HR via electrocardiography (ECG). Data were collected using a BioSemi ActiveTwo system
(BioSemi - The Netherlands). For EEG, a 64-electrode headcap was applied according to the
conventional 10-20 system. For HR, ECG was measured using two flat electrodes applied at the
left lumbar region and below the left clavicle, following established protocols (7, 17, 31). EDA was
measured using 2 passive Nihon Kohden electrodes, applied on the pointer and middle finger of
the left hand. All signals were recorded at a sampling rate of 2048 Hz, and the recording systems
of both participants were synchronized.

Signal processing

All signal processing (both preprocessing and signal processing) was done in Python (version
3.10), using Numpy (32), Scipy (33) and Pandas (DOI 10.5281/zenodo.3509134).

<u>Interpersonal synchronization</u>

Preprocessing was different for each included sensor. All signals were trimmed to match the
shortest stimulus length (12:37 minutes) for fair comparison.

EEG

EEG was first high-pass filtered using a 4th-order Butterworth filter with a 0.5 Hz cutoff to remove
sensor drift. Artifacts exceeding 500 µV were removed, and linear interpolation was performed
across the affected intervals. Eyeblink artifacts were removed using a multichannel Wiener Filter
(34). All channels were re-referenced to a common average reference. Lastly, the signal was
downsampled to 64 Hz. All filtering steps were done using zero-phase filtering. Additionally, to find
shared information within the EEG from different subjects (for the synchronization analysis), we
applied correlated component analysis (corrCA)(35) to calculate components from the EEG that
are maximally correlated between participants. Input for corrCA consists of 64-channel EEG data,
from which ISCs must be calculated.

ECG

From the raw ECG signal, we extracted instantaneous HR. The two ECG leads were re-referenced
to each other (ECG1-ECG2) to suppress common-mode noise and enhance the signal-to-noise
ratio. We applied the algorithm by Pan & Tompkins et al. (36) to find the R-peaks in the ECG signal.
Instantaneous HR was determined by calculating the time intervals between the peaks using the
ECG-detectors package in Python, based on the ECG processing code provided by Perez et
al.(18).

EDA

For EDA, we smoothed the raw signal using a 1.5-second Bartlett window. We computed the
difference between consecutive samples to remove low-frequency trends and emphasize rapid

changes in the signal (37), and we applied a sigmoid transformation (i.e., $y = \frac{1}{1+e^{-x}}$, with x = EDA
amplitude) to compress large peaks. This scaled the data between 0 and 1, where 0 indicates no
change or a decrease, and values above 0 indicate an increase.

To assess interpersonal synchronization, we computed ISCs, which quantify the co-fluctuation of
each body response (EEG, HR, EDA) among multiple participants. This involved calculating the
average correlations of one participant's responses with those of all other participants. This resulted
in a single average correlation value for each participant for each modality for each condition, which
could be interpreted as the listening engagement performance of that specific participant in relation
to the whole group. A high synchronization value would indicate a high degree of listening
engagement, whereas a low value would indicate that the participant was not very involved with
the auditory stimulus material. To assess significance, we applied circular shifting. We shifted one
of the two signals and then recalculated the correlation. Each value was then Fisher transformed.
This procedure was repeated 1000 times to generate a null distribution, from which we derived a
p-value for each synchronization value.

<u>Neural tracking of the auditory stimulus</u>

All stimuli were sampled at 48 kHz and normalized for RMS. We extracted the envelope using a
gammatone   filterbank   (Brian2   and   Brian2hears   toolkits;   https://github.com/brian-
team/brian2hears), where the sound wave was filtered in 28 subbands between 500 and 5000 Hz.
Next, the absolute value of each signal was raised by a power of 0.6 to extract the slow modulations
(38). The envelope was downsampled to 64 Hz and used as the sound feature in neural tracking.
For consistency, all stimuli were trimmed to match the shortest length (12:37 minutes).

No CorrCA was applied for neural tracking analysis. Before analysis, both the EEG data and the
speech envelope were bandpass filtered between 0.5 and 4 Hz with a 4th-order Butterworth filter.
To quantify neural tracking, we reconstructed the envelope from EEG data using a linear decoder
trained via ridge regression. The decoder had an integration window of 250 ms (or 16 samples),
with the regularization parameter (lambda) set to max(abs(EEG covariance matrix)) to prevent
overfitting. Linear weights for each channel at different time delays were calculated to minimize the
mean squared error between the reconstructed and actual speech envelope. We applied leave-
one-fold-out cross-validation, where each recording was split into N non-overlapping folds of 120
seconds. The decoder was trained on all but one fold and then tested on the held-out fold. After
training the decoder for a specific stimulus, the envelope was reconstructed from the EEG data and
correlated with the speech envelope to obtain a correlation reconstruction accuracy value. This
value indicated how well the stimulus was represented in the brain over time.

Statistical significance was calculated for each stimulus separately using circ-shifting (2), which is
similar to the procedure described earlier for the synchronization analysis. In this case, the pre-
processed speech envelope was randomly shifted in time to generate new noisy correlations. We
applied the same Fisher transformation before using all the new correlation values to generate the
null distribution.

<u>Behavioral measure</u>

We evaluated participants' experienced listening engagement and enjoyment after each engaging
and non-engaging active listening condition. We created a new scale by combining and integrating
two validated scales, tapping into self-perceived listening engagement(10, 15). Specifically, we
combined the 'attention' subscale from the Narrative Absorption Scale and adapted the
Engagement Questionnaire by Hannum & Simon(15) to focus more on listening experiences. This
new Listening Engagement Scale consisted of 15 statements about their listening experience,
which participants rated on a 1-7 Likert scale. For further analysis, we calculated the mean of these
15 statements for each condition separately per participant. We also administered a short multiple-

choice content questionnaire to assess content processing quality and motivate participants to listen actively.

Data analysis

All data visualization and analysis were conducted in R Statistical Software (version 4.3.1, R Core Team, 2023). Statistical significance of behavioral ratings and synchronization measures across different conditions was evaluated using a linear mixed-effects model, with synchronization as the dependent variable, condition as a fixed effect, and participants as a random effect. To determine whether the different conditions had a distinct effect on the synchronization measure, we compared two linear mixed-effects models—one including condition as a predictor and one excluding it—using an ANOVA Likelihood Ratio Test to determine significant differences. We used the following models:

Full model: Synchronization value ~ condition + (1|participants)

Reduced model: Synchronization value ~ (1|participants)

Post-hoc comparisons were corrected for multiple comparisons using a Holm correction.
In addition, we calculated Spearman correlations between synchronization values and behavioral ratings. We calculated one correlation per modality across all participants and conditions, each containing 4 data points per participant (two engaging and two non-engaging conditions). Since this violated the assumption of independence, we generated a new null distribution by shuffling the synchronization data by matching the values of each participant to the values of another participant and then again calculating the synchronization-behavior correlation each time. This reshuffling process was repeated 1000 times. This procedure was applied consistently across all included physiological signals.

To enhance our understanding of the combined physiological responses in relation to the behavioral rating, we computed a summarized index using partial least squares (PLS) regression (39). Given the high correlation among the different physiological responses, this approach gave us a comprehensive measure while addressing multicollinearity. We applied PLS regression using a leave-one-out cross-validation approach, where the model was trained with three components on data from 39 out of 40 participants, leaving one out for validation. We selected three components as this was the maximum, based on the number of predictor variables (39). This process was repeated for each participant. We reconstructed the behavioral data using the estimated values from the three PLS components of the regression model. Then we evaluated the alignment between the reconstructed and actual behavioral measures using Spearman's rank correlation. This final step allowed us to determine how well the PLS regression reconstructed measure aligned with the ground truth behavioral data.

Acknowledgments

The authors thank Phebe Dekeyzer and Jade Willaert for their help during data collection.

References

1.   C. Brodbeck, J. Z. Simon, Continuous speech processing. *Curr Opin Physiol* [Preprint] (2020).

2.   M. J. Crosse, *et al.*, Linear Modeling of Neurophysiological Responses to Speech and
Other Continuous Stimuli: Methodological Considerations for Applied Research. *Front*
Neurosci [Preprint] (2021).

3.   J. Vanthornhout, L. Decruy, J. Wouters, J. Z. Simon, T. Francart, Speech Intelligibility
Predicted from Neural Entrainment of the Speech Envelope. *JARO - Journal of the*
Association for Research in Otolaryngology **19**, 181–191 (2018).

4.   I. Iotzov, L. C. Parra, EEG can predict speech intelligibility. *J Neural Eng* **16** (2019).

5.   J. Vanthornhout, L. Decruy, T. Francart, Effect of Task and Attention on Neural Tracking of
Speech. *Front Neurosci* **13** (2019).

6.   N. Ding, J. Z. Simon, Emergence of neural encoding of auditory objects while listening to
competing speakers. *Proc Natl Acad Sci U S A* **109**, 11854–11859 (2012).

7.   J. Madsen, L. C. Parra, Bidirectional brain-body interactions during natural story listening.
Cell Rep **43** (2024).

8.   R. Busselle, H. Bilandzic, Measuring Narrative Engagement. *Media Psychol* **12**, 321–347
(2009).

9.   B. Herrmann, I. S. Johnsrude, A model of listening engagement (MoLE). *Hear Res* **397**
(2020).

10. M. M. Kuijpers, F. Hakemulder, E. S. Tan, M. M. Doicaru, Exploring absorbing reading
experiences. *Sci Study Lit* **4**, 89–122 (2014).

11. S. S. Cohen, L. C. Parra, Memorable audiovisual narratives synchronize sensory and
supramodal neural responses. *eNeuro* **3** (2016).

12. H. Song, E. S. Finn, M. D. Rosenberg, Neural signatures of attentional engagement during
narratives and its consequences for event memory. *PSYCHOLOGICAL AND COGNITIVE*
SCIENCES **118** (2021).

13. D. Bevilacqua, *et al.*, Brain-to-brain synchrony and learning outcomes vary by student–
teacher dynamics: Evidence from a real-world classroom electroencephalography study. *J*
Cogn Neurosci **31**, 401–411 (2018).

14. M. Cziksentmihalyi, Flow: The Psychology of Optimal Experience. *Harper & Row* (1990).

15. M. E. Hannum, C. T. Simons, Development of the engagement questionnaire (EQ): A tool
to measure panelist engagement during sensory and consumer evaluations. *Food Qual*
Prefer **81** (2020).

16. V. C. Irsik, I. S. Johnsrude, B. Herrmann, Neural Activity during Story Listening Is
Synchronized across Individuals Despite Acoustic Masking. *J Cogn Neurosci* **34**, 933–950
(2022).

17. J. Madsen, L. C. Parra, Cognitive processing of a common stimulus synchronizes brains,
hearts, and eyes. *PNAS Nexus* **1** (2022).

18. P. Pérez, *et al.*, Conscious processing of narrative stimuli synchronizes heart rate
between individuals. *Cell Rep* **36** (2021).

19. S. Kinreich, A. Djalovski, L. Kraus, Y. Louzoun, R. Feldman, Brain-to-Brain Synchrony
during Naturalistic Social Interactions. *Sci Rep* **7** (2017).

20. A. Pérez, M. Carreiras, J. A. Duñabeitia, Brain-To-brain entrainment: EEG interbrain
synchronization while speaking and listening. *Sci Rep* **7** (2017).

21. R. V. Palumbo, *et al.*, Interpersonal Autonomic Physiology: A Systematic Review of the
Literature. *Personality and Social Psychology Review* **21**, 99–141 (2017).

22. X. Deng, *et al.*, Parental involvement affects parent-adolescents brain-to-brain synchrony
when experiencing different emotions together: An EEG-based hyperscanning study.
Behavioural Brain Research **458** (2024).

23. A. Anthiyur Aravindan, S. Kalyan Chappidi, A. Thumma, R. Palanisamy, Prediction of
Arousal and Valence State from Electrodermal Activity using Wavelet based ResNet50
Model in *Current Directions in Biomedical Engineering*, (Walter de Gruyter GmbH, 2023),
pp. 555–558.

24. M. Kołodziej, P. Tarnowski, A. Majkowski, R. J. Rak, Electrodermal activity measurements
for detection of emotional arousal. *Bulletin of the Polish Academy of Sciences: Technical
Sciences* **67**, 813–826 (2019).

25. R. Sánchez-Reolid, F. López de la Rosa, D. Sánchez-Reolid, M. T. López, A. Fernández-
Caballero, Machine Learning Techniques for Arousal Classification from Electrodermal
Activity: A Systematic Review. *Sensors* [Preprint] (2022).

26. I. Chatterjee, M. Goršič, J. D. Clapp, D. Novak, Automatic Estimation of Interpersonal
Engagement During Naturalistic Conversation Using Dyadic Physiological Measurements.
Front Neurosci **15** (2021).

27. J. F. Thayer, A. L. Hansen, E. Saus-Rose, B. H. Johnsen, Heart rate variability, prefrontal
neural function, and cognitive performance: The neurovisceral integration perspective on
self-regulation, adaptation, and health. *Annals of Behavioral Medicine* [Preprint] (2009).

28. A. Marois, D. Mouratille, Y. Pratviel, C. Chamberland, S. Tremblay, Using Cardiac and
Electrodermal Activity as Cognitive Markers for Interruptions and Distraction in a
Surveillance Simulation in *Neuroergonomics and Cognitive Engineering*, (AHFE
International, 2024).

29. M. K. Pichora-Fuller, *et al.*, Hearing impairment and cognitive energy: The framework for
understanding effortful listening (FUEL). *Ear Hear* **37**, 5S-27S (2016).

30.   S. Denys, M. Hofmann, A. van Wieringen, J. Wouters, Improving the efficiency of the digit
triplet test using digit scoring with variable adaptive step sizes. *Int J Audiol* **58**, 670–677
(2019).

31.   M. Metshein, A. Krivošei, A. Abdullayev, P. Annus, O. Märtens, Non-Standard Electrode
Placement Strategies for ECG Signal Acquisition. *Sensors* **22** (2022).

32.   C. R. Harris, *et al.*, Array programming with NumPy. *Nature* [Preprint] (2020).

33.   P. Virtanen, *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nat Methods **17**, 261–272 (2020).

34.   B. Somers, T. Francart, A. Bertrand, A generic EEG artifact removal algorithm based on
the multi-channel Wiener filter. *J Neural Eng* **15** (2018).

35.   L. C. Parra, S. Haufe, J. P. Dmochowski, Correlated Components Analysis - Extracting
Reliable Dimensions in Multivariate Data. (2018). https://doi.org/10.51628/001c.7125.

36.   J. Pan, W. J. Tompkins, A Real-Time QRS Detection Algorithm. *IEEE Trans Biomed Eng* **32**
(1985).

37.   C. D. Marci, J. Ham, E. Moran, S. P. Orr, Physiologic correlates of perceived therapist
empathy and social-emotional process during psychotherapy. *Journal of Nervous and*
Mental Disease **195**, 103–111 (2007).

38.   W. Biesmans, N. Das, T. Francart, A. Bertrand, Auditory-inspired speech envelope
extraction methods for improved EEG-based auditory attention detection in a cocktail
party scenario. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* **25**,
402–412 (2017).

39.   J. A. Wegelin, "A Survey of Partial Least Squares (PLS) Methods, with Emphasis on the
Two-Block Case" (2000).

504
505
506

507   **Figures and Tables**
508
509   **Figure 1.**

510
Figure 1: Each data point represents the mean correlation between one participant's response and
all other participants' responses for a given stimulus and physiological measure. Colored points
indicate significance. All measures of listening engagement—both behavioral and physiological
(through synchronization)—showed higher engagement in the engaging condition compared to the
non-engaging condition. Engaging stimuli were rated significantly higher than non-engaging ones
(p < 0.0001). In the physiological data, higher levels of synchronization and neural tracking were
observed in the engaging condition relative to the non-engaging condition. Notably, brain
synchronization and neural tracking also showed significant differences between the non-engaging
and distracted conditions.

520
521

  Figure 2.

Figure 2: Physiological synchronization (brain, HR and EDA) correlates significantly with behavioral listening engagement ratings. Additionally, a combined physiological measure (predicted through partial least squares regression) also significantly correlated with behavioral listening engagement.

529 **Figure 3.**

Figure 3: Overview of the experimental setup, the implemented physiological and behavioral
measures, and the full experimental procedure. The conditions were presented in a random order
but were always administered in their entirety (Engaging – Non-Engaging – Distracted).

536 **Table 1.**

538 *Table 1: Contrasting condition effects for each of the neural and physiological measures.*

	Contrast	β (estimate)	SE	t	Adjusted p-value
Brain	**E - N**	0.00644	0.0005	12.878	**<0.0001**
	E - D	-0.00883	0.000608	-14.529	**<0.0001**
	N - D	-0.00239	0.000611	-3.913	**0.0001**
Heart rate	**E - N**	0.04332	0.00356	12.174	**<0.0001**
	E - D	-0.04136	0.00433	-9.557	**<0.0001**
	N - D	0.00196	0.00435	0.451	0.6525
Electrodermal activity	**E - N**	0.0145	0.00156	9.261	**<0.0001**
	E - D	-0.0115	0.00190	-6.036	**<0.0001**
	N - D	0.0030	0.00191	1.571	0.1181
Neural tracking	**E - N**	0.0228	0.00471	4.839	**<0.0001**
	E - D	-0.0388	0.00574	-6.766	**<0.0001**
	N - D	0.0228	0.00471	4.839	**<0.0001**

539 *Note. Holm-corrected post hoc comparisons; bold indicates significant contrasts.(E = Engaging*
540 *condition, N = non-engaging condition, D = Distracted condition)*