Title: In Sync With Sound: Interpersonal synchronization as an
objective measure of listening engagement

Authors: Lotte Lambrechts[1], Bernd Accou[1], Jonas Vanthornhout[1], Bart Boets[2], Tom Francart[1]

Corresponding author: Lotte Lambrechts (lotte.lambrechts@kuleuven.be)

Affiliation:

1. Department of Neurosciences, ExpORL, KU Leuven - University of Leuven, Leuven (3000),
Belgium
2. Department of Neurosciences, CDP, KU Leuven - University of Leuven, Leuven (3000),
Belgium

Adress : Herestraat 49 – 721, 3000 Leuven

Phone: +32 16 32 82 60

Email: lotte.lambrechts@kuleuven.be

Author Contributions: L.L., B.A., J.V., B.B. and T.F. conceptualized the study. L.L. and B.A. collected
the data. L.L. and B.A. analyzed the data. L.L., B.A., J.V., B.B. and T.F. interpreted the data. L.L. wrote
the first draft of the manuscript. L.L., B.A., J.V., B.B. and T.F. reviewed and edited the manuscript.

Competing Interest Statement: No competing interests

Keywords: Story listening, Listening engagement, Physiological measurements, Interpersonal
synchronization

Word count: 4652

26   29

Abstract

Listening engagement plays a crucial role in
effective communication and knowledge
transfer, reflecting a state of deep absorption in
an auditory stimulus. The available behavioral
assessments of listening engagement are
limited in capturing its multidimensional nature.
Measuring listening engagement objectively
using brain and body signals is a compelling
alternative. In particular, interpersonal
synchronization—observed through
synchronized neural and physiological
responses among listeners—may offer a
promising objective measure. In this study, we
examined synchronization of brain activity,
heart rate, and electrodermal activity between
participants while they listened to engaging and
non-engaging stories. Subjective engagement
ratings were collected as a ground truth
reference. Results showed higher interpersonal
synchronization across all modalities when
participants listened to engaging stories
compared to non-engaging ones, with
significant correlations between
synchronization measures and ratings of
subjective engagement. These results confirm
that synchronization may be a reliable,
objective marker of listening engagement. We
anticipate that these results will provide a
powerful framework for future research in
relevant domains, including auditory
neuroscience, education, and communication
science, and enable impactful real-world
applications, such as automatically tracking
audience engagement in education,
entertainment, and broader societal contexts.

Introduction

Every second of the day, we encounter various sounds, with speech one of the most pervasive. Where some of these seem negligible (train station announcements), carry little meaning (listening to an automated voice message) or don't capture you're interest (listening to a boring lecture), others, like a conversation with a dear friend, will encourage you to be fully present and absorb every detail. Despite these vastly different listening experiences, in all cases our brains will process and interpret these speech sounds, tracking their complex acoustic dynamics [1], [2], [3]. Neural tracking of speech has been established as a reliable and valid measure of natural speech processing and perception, allowing for an objective assessment of speech understanding [1], [4]. Research has shown that this neural representation of speech is modulated by auditory attention to speech [5], [6]. Auditory attention has previously been understood as the conscious decision to direct one's focus to an external or internal presence [7], [8]. While this has been shown to significantly affect speech processing, it does not completely account for the complexities of everyday listening experiences. Research suggests that when you are genuinely interested and motivated to listen to a speech source, you can enter a state of listening engagement in which you fully absorb everything that is being said. Based on previous research, we have defined listening engagement as deep **immersion in an auditory stimulus, where individuals feel connected to the narrative and its characters** [9]. Listening engagement is different from attention, in that it also involves an unconscious and affective response related to the content or the speaker [9], [10]. It may involve narrative presence [9], [11], a state where listeners lose awareness of their surroundings and feel immersed in the narrative world. Listening engagement yields several beneficial outcomes: It is linked to enhanced speech processing and communication, enhanced memory retention [12], [13], higher listening enjoyment [10], [14], and the experience of a state of flow [9], [15]. Consequently, improving listening engagement could positively impact daily life by for instance enhancing education and media entertainment.

Thus far, listening engagement has mostly been evaluated through behavioral self-report rating scales [9], [11], [16], such as the 'Narrative absorption scale' and the 'Narrative engagement scale'. These well-validated measures can reliably capture variability in listening engagement [17]. Nevertheless, behavioral self-report measures hold inherent limitations, as they may disturb the immediate experience and induce biases (e.g., socially desirable responses when students have to rate their engagement with a lecture). Additionally, they provide only limited temporal resolution, failing to capture moment-to-moment fluctuations in engagement. To address these limitations, an objective measure of listening engagement could provide valuable insights by minimizing bias and allowing continuous assessment. With this research, we will explore how brain and body responses may serve as reliable indicators of listening engagement. Given the strong affective response we expect it to elicit, the combination of brain and body responses may better reflect

listening engagement than just brain response
alone [18], [19].

In the field of socio-affective neurosciences, it
has already been shown that when people
share an experience, their brain and body
responses will synchronize [20], [21], [22], [23],
[24]. Recently, researchers have started to
explore the possibility of interpreting listening to
the same sound, also as a shared experience.
Following this, there is some initial evidence
that when people consciously process
engaging stories, their physiological responses
tend to synchronize [25], [26], [27].
Interpersonal synchronization occurs across
multiple physiological modalities, including
brain activity, heart rate (HR), and
electrodermal activity (EDA). However,
considerable uncertainty remains regarding the
factors that drive interpersonal synchronization.
While it has been observed in direct
conversational interactions [28], it also occurs
when individuals (jointly) passively listen to the
same auditory stimulus [25], [26], [27].
Remarkably, interpersonal synchronization has
been found even when participants listen
separately to the same stimulus at different
moments [26], [27], suggesting that
synchronization is not uniquely driven by direct
social interaction [25], [29], but rather by shared
cognitive and emotional engagement with the
content. However, of course, social interactions
may amplify these effects.

Various mechanisms have been proposed to
explain interpersonal synchrony. Bevilacqua et
al. [14] and Madsen & Parra [26], [27] suggest
that it arises from the conscious and attentive
processing of an engaging stimulus, driven by
shared neural oscillations and cognitive
processes in response to the content. For

example, attentive listening to an engaging
stimulus modulates brain activity in a way
that aligns across individuals, leading to
synchronization. Pérez et al. [25] support
this view, proposing that HR synchronizes
across participants in response to cognitive
processes when individuals listen to the
same stimulus. However, other studies
suggest that HR may also be influenced by
emotional responses elicited by the
stimulus, suggesting a broader interplay
among cognitive and affective factors [30].
Given that bodily responses such as HR
and EDA are closely linked to emotions and
arousal [18], [31], [32], emotional
processing likely plays a role in listening
engagement alongside cognitive
mechanisms.

To further understand the relationship
between listening and interpersonal
synchronization, Madsen and Parra [26],
[27] showed that synchronization
decreased when participants were
distracted from listening through a
cognitively demanding task (distracted
condition). This suggests that
synchronization may be influenced by
attention, which is known to be a precursor
of engagement. Despite these interesting
findings, no conclusions regarding the
relationship between synchronization and
listening engagement can be validated, as
they only manipulated attention and not
listening engagement. In addition, previous
studies have either lacked a behavioral
ground truth [17], [26], [27] or used
behavioral measures not directly relevant
for engagement [33], making it challenging

to ascertain whether the auditory stimulation was genuinely engaging.

The present study aims to address these gaps by employing an objective synchronization measure to distinguish between engaging and non-engaging conditions and to examine the relationship between synchronization and listening engagement. To overcome limitations in prior studies, we meticulously selected engaging and dull stimuli for the participant group and incorporated a behavioral measure of engagement. We hypothesize that interpersonal synchronization will be higher when individuals are engaged with the stimulus and that synchronization levels will predict self-rated listening engagement.

Materials and Methods

Participants

Participants subscribed to the experiment in dyads. They came to the lab and participated with someone they knew well (e.g., friends, romantic partners, or family members). After evaluating the inclusion criteria, we included all 40 participants. They had a mean age of 21 years (range: 18 to 25 years). Of these, 4 were male. All participants had normal hearing, confirmed by a digit triplet screening test (SRT = -10 dB SNR) [34] and/ or a tonal audiogram (thresholds < 25 dB HL at all audiometric frequencies), with a mean Fletcher index of 13 ± 6 dB HL. Other exclusion criteria were attention or concentration problems, neuropathological disorders, communication disorders and language disorders. These criteria were evaluated at the beginning of each session through a screening questionnaire. The Medical Ethics Committee UZ Leuven/Research (KU Leuven) approved the study with reference S68492 and each participant provided informed consent before the start of the session.

Setup

Testing took place in dyads in a soundproof room (Faraday cage). Participants sat next to each other, approximately 1.35 meters apart, facing each other at a 90° angle (chairs turned 45° towards the loudspeaker). A loudspeaker and monitor were positioned 1.5 meters in front of the participants. The setup was calibrated, and stimuli were presented at 60 dB A, equivalent to an everyday conversation level. See *figure 1* for a full overview.

Figure 1: Overview of the experimental setup, the implemented physiological and behavioral measures, and the full experimental procedure. The conditions were presented in a random order but were always administered in their entirety (Engaging – Non-Engaging – Distracted).

Experimental Design

This study measured brain and physiological responses, as well as subjective listening engagement, in participants exposed to various auditory stimuli.

The experiment consisted of **three different experimental conditions**, each involving the presentation of an auditory stimulus:

1. Engaging condition: Participants listened attentively to a highly engaging stimulus.
2. Non-engaging condition: Participants listened attentively to a stimulus designed to be less engaging.
3. Distracted condition: Participants completed cognitively demanding riddles while an auditory stimulus played in the background, discouraging attention and engagement.

Participants were instructed to listen attentively to the speech stimulus in the engaging and non-engaging conditions. Engagement levels were manipulated by presenting stimuli with varying degrees of intrinsic interest. The **engaging stimulus** was a 46:29-minute episode of a popular Flemish podcast. This stimulus was expected to be inherently interesting, easy to follow, and able to capture individuals' attention ('De Volksjury' episode 19 'Maddie McCann', ranked second place on "Spotify Wrapped" of

2024). The **non-engaging stimulus** comprised excerpts from Dutch debates in the European parliament (25:19 minutes duration), which were self-recorded by one of the researchers. This content was chosen for its neutral/dull and less captivating nature. The engaging and non-engaging stimuli were split into two parts, so we had two engaging and two non-engaging conditions. To minimize the impact of differences between the stimuli on the subject's responses, we selected stimuli featuring only female voices. After both the engaging and non-engaging conditions, we assessed self-perceived listening engagement, as will be discussed under 'Behavioral measure - p. 9'. For the **distracted condition,** the participants listened to another non-engaging stimulus (a 12:42-minute self-recorded excerpt of the European Parliament) while solving riddles. Participants were given a new riddle every 2 minutes and were instructed to remember their solutions. After this condition, the solutions were orally discussed. This condition was designed to minimize not only engagement but also attention, serving as a control to differentiate synchronization as a measure of engagement from mere attentional effects.

The full experimental procedure and order of the protocol can be found in *Figure 1*. Each dyad followed the same protocol and same stimuli, with the order of conditions (engaging, non-engaging, and distracted) randomized across dyads to mitigate potential training or fatigue effects.

Brain and physiological measures

We recorded three objective measures: brain activity via electroencephalography (EEG), EDA, and HR via electrocardiography (ECG). Data were collected using a BioSemi ActiveTwo system (BioSemi - The Netherlands). For EEG, a 64-electrode headcap was applied according to the 10-20 system. We aimed to keep electrode DC offsets (relative to CMS) between -20 and 20 mV. For HR, ECG was measured using two flat electrodes applied at the left lumbar region and below the left clavicle, following established methods [26], [27], [35]. EDA was measured using 2 passive Nihon Kohden electrodes, applied on the pointer and middle finger of the left hand. All signals were recorded at a sampling rate of 2048 Hz, and the recording systems of both participants were synchronized.

Signal processing

All signal processing (both preprocessing and signal processing) was done in Python (version 3.10), using Numpy [36], Scipy [37] and Pandas (DOI 10.5281/zenodo.3509134).

Interpersonal synchronization

Preprocessing was different for each included sensor. All signals were trimmed to match the shortest stimulus length (12:37 minutes) for fair comparison.

EEG

EEG was first high-pass filtered using a 4th-order Butterworth filter with a 0.5 Hz cutoff to remove sensor drift. Artifacts exceeding 500 µV were removed, and linear interpolation was performed across the affected intervals. Eyeblink artifacts were removed using a multichannel Wiener Filter [38]. All channels were re-referenced to a common average reference. Lastly, the signal was downsampled to 64 Hz. All filtering steps were done using zero-phase filtering. Additionally, to find the relevant shared information within the EEG

from different subjects (for the synchronization
analysis), we applied correlated component
analysis (corrCA) [39]. CorrCA optimizes
weights of a linear combination of EEG
channels to maximize correlation between
subjects' EEG and transforms the information
to components that are ordered from highest to
lowest correlation. These resulting EEG
components, calculated from the original 64-
channel EEG data are used as input to perform
the ISC calculation. To do this, we selected only
the first component, as these will be maximally
correlated.

ECG

From the raw ECG signal, we extracted
instantaneous HR. The two ECG leads were re-
referenced to each other (ECG1-ECG2) to
suppress common-mode noise and enhance
the signal-to-noise ratio. We applied the
algorithm by Pan & Tompkins et al. [40] to find
the R-peaks in the ECG signal. Instantaneous
HR was determined as the inverse of the time
intervals and interpolated to obtain a sampling
rate of 64Hz, similar to the method of Pérez et
al., [25].

EDA

For EDA, we smoothed the raw signal using a
1.5-second Bartlett window. We computed the
difference between consecutive samples to
remove low-frequency trends [41],
downsampled the signal to 64 Hz and applied a
sigmoid transformation (i.e., $y = \frac{1}{1+e^{-x}}$, with x =
EDA amplitude) to compress large peaks. This
scaled the data between 0 and 1, where 0
indicates no change or a decrease, and values
above 0 indicate an increase.

To assess interpersonal synchronization, we
computed ISCs, which quantify the co-
fluctuation of each body response (EEG, HR,
EDA) among multiple participants. This
involved calculating the average of all the
correlations of one participant's responses with
those of all other participants. This resulted in a
single averaged correlation value for each
participant for each modality for each condition,
which could be interpreted as the listening
engagement performance of that specific
participant in relation to the whole group. A high
synchronization value would indicate a high
degree of listening engagement, whereas a low
value would indicate that the participant was not
very involved with the auditory stimulus
material. To assess significance, we applied
circular shifting for each correlation that was
calculated (before averaging). We shifted one
of the two signals and then recalculated the
correlation. Each value was then Fisher
transformed. This procedure was repeated a
1000 times to generate a null distribution, from
which we derived a p-value for each
synchronization value.

<u>Neural tracking of the auditory stimulus</u>

All stimuli were sampled at 48 kHz and
normalized for RMS. We extracted the
envelope using a gammatone filterbank (Brian2
and Brian2hears toolkits;
https://github.com/brian-team/brian2hears),
where the sound wave was filtered in 28
subbands between 500 and 5000 Hz. Next, the
absolute value of each signal was raised by a
power of 0.6 to extract the slow modulations
[42]. The envelope was calculated as the
average of these 28 subband envelopes. The
envelope was downsampled to 64 Hz and used
as the stimulus feature in neural tracking. For

consistency, all stimuli were trimmed to match
the shortest length (12:37 minutes).

No CorrCA was applied for neural tracking
analysis. Before analysis, both the EEG data
and the speech envelope were bandpass
filtered between 0.5 and 4 Hz with a zero phase
4th-order Butterworth filter. To quantify neural
tracking, we reconstructed the envelope from
EEG data using a linear decoder trained via
ridge regression. The decoder had an
integration window of 250 ms (or 16 samples),
with the regularization parameter (lambda) set
to max(abs(EEG covariance matrix)) to prevent
overfitting. Linear weights for each channel at
different time delays were calculated to
minimize the mean squared error between the
reconstructed and actual speech envelope. We
applied leave-one-fold-out cross-validation,
where each recording was split into N non-
overlapping folds of 120 seconds. The decoder
was trained on all but one fold and then tested
on the held-out fold. After training the decoder
for a specific stimulus, the envelope was
reconstructed from the EEG data and
correlated with the speech envelope to obtain a
correlation reconstruction accuracy value. This
value indicated how well the stimulus was
represented in the brain over time.

Statistical significance was calculated for each
stimulus separately using circ-shifting [2], which
is similar to the procedure described earlier for
the synchronization analysis. In this case, the
pre-processed speech envelope was randomly
shifted in time to generate new noisy
correlations. We applied the same Fisher
transformation before using all the new
correlation values to generate the null
distribution.

Behavioral measure

We evaluated participants' subjective listening
engagement and enjoyment after each
engaging and non-engaging active listening
condition. We integrated two validated scales,
tapping into self-perceived listening
engagement. Specifically, we combined the
'attention' subscale from the Narrative
Absorption Scale [11] and adapted the
Engagement Questionnaire by Hannum &
Simon [16] so it was more applicable for
evaluating listening experiences. This new
Listening Engagement Scale consisted of 15
statements about their listening experience,
which participants rated on a 1-7 Likert scale.
We encouraged participants to be as accurate
and honest as possible in their responses. For
further analysis, we calculated the mean of
these 15 statements for each condition
separately per participant. We also
administered a short multiple-choice content
questionnaire to assess content processing
quality and motivate participants to listen
actively.

Statistical analysis

All data visualization and analysis were
conducted in R Statistical Software (version
4.3.1, R Core Team, 2023). Statistical
significance of behavioral ratings and
synchronization measures across different
conditions was evaluated using a linear mixed-
effects model, with synchronization as the
dependent variable, condition as a fixed effect,
and participants as a random effect. To
determine whether the different conditions had
a distinct effect on the synchronization
measure, we compared two linear mixed-
effects models—one including condition as a
predictor and one excluding it—using an

ANOVA Likelihood Ratio Test to determine
significant differences. We used the following
models:

<u>Full model</u>:

Synchronization value ~ condition +
(1|participants)

<u>Reduced model</u>:

Synchronization value ~ (1|participants)

Post-hoc comparisons were corrected for
multiple comparisons using a Holm correction.
In addition, we calculated Spearman
correlations between synchronization values
and behavioral ratings. We calculated one
correlation per modality across all participants
and conditions, each containing 4 data points
per participant (two engaging and two non-
engaging conditions). Since this violated the
assumption of independence, we generated a
new null distribution by shuffling the
synchronization data by matching the values of
each participant to the values of another
participant and then again calculating the
synchronization-behavior correlation each time.
This reshuffling process was repeated a 1000
times. This procedure was applied consistently
across all included physiological signals.

To enhance our understanding of the combined
physiological responses in relation to the
behavioral rating, we computed a summarized
index using partial least squares (PLS)
regression [43]. Given the high correlation
among the different physiological responses,
this approach gave us a comprehensive
measure while addressing multicollinearity. We
applied PLS regression using a leave-one-out
cross-validation approach, where the model
was trained with three components on data

from 39 out of 40 participants, leaving one out
for validation. We selected three components
as this was the maximum, based on the number
of predictor variables [43]. This process was
repeated for each participant. We
reconstructed the behavioral data using the
estimated values from the three PLS
components of the regression model. Then we
evaluated the alignment between the
reconstructed and actual behavioral measures
using Spearman's rank correlation. This final
step allowed us to determine how well the PLS
regression reconstructed measure aligned with
the ground truth behavioral data.

Results
<u>Participants rate engaging stimuli higher than</u>
<u>non-engaging stimuli on the behavioral</u>
<u>Listening Engagement Scale</u>

Figure 2 (bottom, right panel) contains the
averaged (across all questions) subjective
engagement ratings for the different conditions.
Due to system failure, we lost responses for the
first engaging condition of one participant, the
first non-engaging condition of three
participants, and the second non-engaging
condition of 4 participants. The mean listening
engagement score for the engaging condition
was 4.8. The mean score for the non-engaging
stimuli was 2.1. The linear mixed effects model
showed highly significant differences in
subjective engagement between both
conditions (β = -2.70, SE = 0.098, t = -27.73, p
< .001).

<u>Participants display higher interpersonal</u>
<u>synchronization when listening to an engaging</u>
<u>stimulus</u>

The top three panels in *Figure 2* show synchronization of physiological responses between individuals while listening to auditory stimuli with varying engagement levels. Statistical significance of each data point was calculated using circular shifting.

A highly significant effect of condition could be identified for all physiological responses, based on ANOVA comparisons with and without condition: brain synchronization ($\chi^2(2)$ = 158.24, p < .0001), HR synchronization ($\chi^2(2)$ = 119.44, p < .0001), EDA synchronization ($\chi^2(2)$ = 73.392, p < .0001). Post-hoc comparisons with Holm correction (**Table 1**) revealed significantly higher synchronization in the engaging condition (E) than in the non-engaging (N) and distracted condition (D) for all physiological responses. Additionally, brain synchronization was higher in the non-engaging condition than in the distracted condition, but this effect was not observed for HR and EDA.

Figure 2: Each data point represents the mean correlation between one participant's response and all other participants' responses for a given stimulus and physiological measure. Colored points indicate significance. All measures of listening engagement—both behavioral and physiological (through synchronization)—showed higher engagement in the engaging condition compared to the non-engaging condition. Engaging stimuli were rated significantly higher than non-engaging ones (p < 0.0001). In the physiological data, higher levels of synchronization and neural tracking were observed in the engaging condition relative to the non-engaging condition. Notably, brain synchronization and

671 *neural tracking also showed significant differences between the non-engaging and distracted*
672 *conditions.*

673 *Table 1: Contrasting condition effects for each of the neural and physiological measures.*

	Contrast	β (estimate)	SE	t	Adjusted p-value
Brain	**E - N**	0.00644	0.0005	12.878	**<0.0001**
	E - D	-0.00883	0.000608	-14.529	**<0.0001**
	N - D	-0.00239	0.000611	-3.913	**0.0001**
Heart rate	**E - N**	0.04332	0.00356	12.174	**<0.0001**
	E - D	-0.04136	0.00433	-9.557	**<0.0001**
	N - D	0.00196	0.00435	0.451	0.6525
Electrodermal activity	**E - N**	0.0145	0.00156	9.261	**<0.0001**
	E - D	-0.0115	0.00190	-6.036	**<0.0001**
	N - D	0.0030	0.00191	1.571	0.1181
Neural tracking	**E - N**	0.0228	0.00471	4.839	**<0.0001**
	E - D	-0.0388	0.00574	-6.766	**<0.0001**
	N - D	0.0228	0.00471	4.839	**<0.0001**

Note. Holm-corrected post hoc comparisons; bold indicates significant contrasts.(E = Engaging
condition, N = non-engaging condition, D = Distracted condition)

We also looked at neural tracking results.
Significant neural tracking could be found for
each condition, but not for every participant,
especially in the distracted condition. Using a
similar analysis approach as for the
synchronization measures, we identified a
significant effect of condition for neural tracking
($\chi^2(2)$ = 45.033, p < .0001). Post-hoc

comparisons showed significant neural tracking differences between the three conditions (**Table 1**).

Relation between physiological synchrony measures and behavioral engagement ratings

We analyzed the relationship between the physiological and behavioral data. *Figure 3* presents scatter plots depicting the association between the different interpersonal synchronization values and the scores on the Listening Engagement Scale. Each correlation plot includes data for all participants and conditions involving attentive listening (excluding the distracted condition, as we did not assess any subjective ratings). The first panel contains a scatter plot for brain synchronization, the second for HR synchronization, and the third panel for EDA synchronization. All correlations were significant, indicating a clear relationship between objective synchronization measures and subjective engagement ratings across all modalities. For the different body responses, the scatter plots indicate that higher engagement ratings correspond to higher physiological synchronization between participants' physiological responses.

Figure 3: Physiological synchronization (brain, HR and EDA) correlates significantly with behavioral listening engagement ratings. Additionally, a combined physiological measure (predicted through partial least squares regression) also significantly correlated with behavioral listening engagement.

Additionally, we conducted a PLS regression using physiological responses as predictors and behavioral data as the outcome variable. Based on three predictor variables, the regression model explained 88% of the variance in behavioral rating outcomes. To assess how well the behavioral ratings could be reconstructed from these physiological responses, we used Spearman's correlation to compare the reconstructed data with the Listening Engagement Scale results. This analysis revealed a strong correlation of 0.68 between the PLS measure and the engagement scores, indicating a meaningful relationship between physiological responses and self-reported engagement (see *Figure 3 - last panel*). The observed correlation was higher than any of the individual synchronization results; however, no statements can be made regarding significance.

Discussion

This paper aims to evaluate interpersonal synchronization as a measure of listening engagement. Twenty duos were invited to our

lab and listened to auditory stimuli varying in
engagement level (ranging from
engaging/interesting to non-
engaging/uninteresting). Meanwhile, we
recorded physiological brain and body
responses. As an extra control condition,
participants also completed a distracting task
while listening to the stimuli. Our research
indicates that physiological synchronization
between individuals may be a valuable
indicator of listening engagement.

For each of the physiological modalities and
across all conditions, significant interpersonal
synchronization was observed when people
were presented with an auditory stimulus. This
finding supports past results that link
interpersonal synchronization to conscious
speech processing [17], [26], [27]. Extending
previous research, our results show that
synchronization was higher when participants
listened in an engaged manner and significantly
lower when the stimulus was non-engaging.
This pattern was consistent across all the brain
and body signals (EDA and HR), suggesting
that engagement activates additional mental
processes reflected in physiological responses
[44], [45]. These processes might drive
interpersonal synchronization across the
physiological responses. However, identifying
the exact cause of this effect is challenging and
requires further research. Nevertheless, our
findings expand significantly on previous
research, demonstrating that synchrony does
not merely result from attentive listening, as
participants were encouraged to listen
attentively in both the engaging and non-
engaging conditions.

We also saw that synchronization declined
when participants were distracted compared to

when they listened attentively, aligning with
previous studies [25], [26], [27]. Through brain
responses, including neural tracking and brain-
to-brain synchronization, we were able to
distinguish between non-engaged listening and
distracted listening, indicating that engagement
extends beyond attention and conscious
processing [17], [26], [27]. These findings
support the idea that synchronization can be
used as a proxy for listening engagement.
However, synchronization did not significantly
differ between non-engaged and distracted
listening conditions for HR and EDA. From
these results, at this point, we cannot make any
conclusions as to why this is not the case. It is
plausible that attention effects are uniquely
cognitively driven and therefore only
represented in the brain and that the body
responses are more reflective of affective
experiences, related to engagement [18], [32].
To the best of our knowledge, this hypothesis
has not been tested yet. Further research
should be carried out to tease this more apart.

Additionally, we examined the relationship
between self-rated listening engagement and
objective interpersonal synchronization data.
Results revealed strong and consistent
correlations between subjective ratings and
objective synchronization measures across all
three physiological responses. While these
correlations may be driven by condition effects,
their strength and consistency support the
hypothesis that synchronization can predict
listening engagement at the individual subject
level. PLS analysis revealed a strong
relationship between a combined physiological
measure and behavioral engagement scores.
Although listening engagement is a highly
individualized, subjective, and hard-to-capture
cognitive-affective construct [9], [10],

interpersonal synchronization across these very basic objective physiological measures seems to relate strongly to it. Further analysis could determine whether a single measure, a subset, or the full combination is most effective. At present, the combined measure exhibits the highest correlation, outperforming individual measures. Future research with larger datasets and more advanced models could further refine these findings. Ultimately, these findings could inform the development of sensors and systems capable of automatically monitoring and tracking audience focus and engagement, with promising applications in society, education and entertainment.

While encouraging, these results stem from a deliberately sharp contrast in stimulus types. To maximize the contrast between engaging and non-engaging stimuli, we selected content that was either highly interesting or very dull. However, in everyday life, auditory input often falls along a more nuanced spectrum. Future studies should incorporate a broader range of engagement levels to determine whether a continuous gradient in synchronization can be observed.

Listening engagement is a multidimensional construct influenced by various factors, making it challenging to control all potential confounds [9], [10]. For example, different speakers were used across the conditions, and individual preferences for voice timbre could have affected engagement. To partly control for this, we ensured all speakers were female, eliminating potential effects of the speaker's sex. However, factors such as fatigue, anxiety, and personal interest [7], [9], [10] may also play a role. Future research should also further investigate how speaker engageability characteristics (e.g., speech rate, prosody, vocal quality) influence listening engagement.

Another key challenge is differentiating between listening engagement and auditory attention. In all active listening conditions, participants were instructed to listen and focus attentively to the presented stimulus. To ensure compliance, participants were informed beforehand that they would be asked to answer multiple-choice questions. While this design theoretically ensures comparable attention levels across both listening activities, engaged and attentive listening may be more intricately linked in practice. As a result, it remains unclear how much of the observed synchronization differences between conditions were driven by engagement or variation in attention. Future studies should further disentangle these overlapping constructs.

Finally, through future analyses, we will examine dyadic effects within these data to gain deeper insights into interpersonal synchronization dynamics and explore whether synchronization is enhanced by physical proximity and real-life presence.

Conclusion

In this study, participants listened to auditory stimuli while we measured their brain activity, HR, and EDA. The stimuli were designed to be either engaging or non-engaging. Our analysis revealed that interpersonal synchronization was significantly higher in the engaging condition across all physiological responses. Additionally, individual differences in interpersonal synchronization closely correlated with subjective listening engagement scores. These findings strongly suggest that

898 interpersonal synchronization can serve as an
899 objective marker of listening engagement.

References

[1] J. Vanthornhout, L. Decruy, J. Wouters, J. Z. Simon, and T. Francart, "Speech Intelligibility Predicted from Neural Entrainment of the Speech Envelope," *JARO - Journal of the Association for Research in Otolaryngology*, vol. 19, no. 2, pp. 181–191, Apr. 2018, doi: 10.1007/s10162-018-0654-z.

[2] M. J. Crosse, N. J. Zuk, G. M. Di Liberto, A. R. Nidiffer, S. Molholm, and E. C. Lalor, "Linear Modeling of Neurophysiological Responses to Speech and Other Continuous Stimuli: Methodological Considerations for Applied Research," Nov. 22, 2021, *Frontiers Media S.A.* doi: 10.3389/fnins.2021.705621.

[3] C. Brodbeck and J. Z. Simon, "Continuous speech processing," Dec. 01, 2020, *Elsevier Ltd*. doi: 10.1016/j.cophys.2020.07.014.

[4] I. Iotzov and L. C. Parra, "EEG can predict speech intelligibility," *J Neural Eng*, vol. 16, no. 3, 2019, doi: 10.1088/1741-2552/ab07fe.

[5] J. Vanthornhout, "Towards an objective and automatic measure of speech understanding," 2019.

[6] N. Ding and J. Z. Simon, "Emergence of neural encoding of auditory objects while listening to competing speakers," *Proc Natl Acad Sci U S A*, vol. 109, no. 29, pp. 11854–11859, Jul. 2012, doi: 10.1073/pnas.1205381109.

[7] M. K. Pichora-Fuller *et al.*, "Hearing impairment and cognitive energy: The framework for understanding effortful listening (FUEL)," *Ear Hear*, vol. 37, pp. 5S-27S, 2016, doi: 10.1097/AUD.0000000000000312.

[8] N. A. Phillips, "The implications of cognitive aging for listening and the framework for understanding effortful listening (FUEL)," in *Ear and Hearing*, Lippincott Williams and Wilkins, 2016, pp. 44S-51S. doi: 10.1097/AUD.0000000000000309.

[9] R. Busselle and H. Bilandzic, "Measuring Narrative Engagement," *Media Psychol*, vol. 12, no. 4, pp. 321–347, Nov. 2009, doi: 10.1080/15213260903287259.

[10] B. Herrmann and I. S. Johnsrude, "A model of listening engagement (MoLE)," *Hear Res*, vol. 397, Nov. 2020, doi: 10.1016/j.heares.2020.108016.

[11] M. M. Kuijpers, F. Hakemulder, E. S. Tan, and M. M. Doicaru, "Exploring

absorbing reading experiences," *Sci Study Lit*, vol. 4, no. 1, pp. 89–122, Oct. 2014, doi: 10.1075/ssol.4.1.05kui.

[12] S. S. Cohen and L. C. Parra, "Memorable audiovisual narratives synchronize sensory and supramodal neural responses," *eNeuro*, vol. 3, no. 6, Jan. 2016, doi: 10.1523/ENEURO.0203-16.2016.

[13] H. Song, E. S. Finn, and M. D. Rosenberg, "Neural signatures of attentional engagement during narratives and its consequences for event memory," *PSYCHOLOGICAL AND COGNITIVE SCIENCES*, vol. 118, no. 33, 2021.

[14] D. Bevilacqua *et al.*, "Brain-to-brain synchrony and learning outcomes vary by student–teacher dynamics: Evidence from a real-world classroom electroencephalography study," *J Cogn Neurosci*, vol. 31, no. 3, pp. 401–411, 2018, doi: 10.1162/jocn_a_01274.

[15] M. Cziksentmihalyi, "Flow: The Psychology of Optimal Experience," *Harper & Row*, 1990.

[16] M. E. Hannum and C. T. Simons, "Development of the engagement questionnaire (EQ): A tool to measure panelist engagement during sensory and consumer evaluations," *Food Qual Prefer*, vol. 81, Apr. 2020, doi: 10.1016/j.foodqual.2019.103840.

[17] V. C. Irsik, I. S. Johnsrude, and B. Herrmann, "Neural Activity during Story Listening Is Synchronized across Individuals Despite Acoustic Masking," *J Cogn Neurosci*, vol. 34, no. 6, pp. 933–950, May 2022, doi: 10.1162/jocn_a_01842.

[18] M. Kołodziej, P. Tarnowski, A. Majkowski, and R. J. Rak, "Electrodermal activity measurements for detection of emotional arousal," *Bulletin of the Polish Academy of Sciences: Technical Sciences*, vol. 67, no. 4, pp. 813–826, 2019, doi: 10.24425/bpasts.2019.130190.

[19] C. A. F. Wascher, "Heart rate as a measure of emotional arousal in evolutionary biology," Aug. 16, 2021, *Royal Society Publishing*. doi: 10.1098/rstb.2020.0479.

[20] V. Leong, E. Byrne, K. Clackson, S. Georgieva, S. Lam, and S. Wass, "Speaker gaze increases information coupling between infant and adult brains," *Proc Natl Acad Sci U S A*, vol. 114, no. 50, pp. 13290–13295,

Dec. 2017, doi: 10.1073/pnas.1702493114.

[21] S. F. Waters, T. V. West, H. R. Karnilowicz, and W. B. Mendes, "Affect contagion between Mothers and Infants: Examining valence and touch," *J Exp Psychol Gen*, vol. 146, no. 7, pp. 1043–1051, Jul. 2017, doi: 10.1037/xge0000322.

[22] S. V. Wass *et al.*, "Parental neural responsivity to infants' visual attention: How mature brains influence immature brains during social interaction," *PLoS Biol*, vol. 16, no. 12, Dec. 2018, doi: 10.1371/journal.pbio.2006328.

[23] S. V. Wass, M. Whitehorn, I. Marriott Haresign, E. Phillips, and V. Leong, "Interpersonal Neural Entrainment during Early Social Interaction," Apr. 01, 2020, *Elsevier Ltd*. doi: 10.1016/j.tics.2020.01.006.

[24] C. G. Smith, E. J. H. Jones, T. Charman, K. Clackson, F. U. Mirza, and S. V. Wass, "Vocalization and physiological hyperarousal in infant-caregiver dyads where the caregiver has elevated anxiety," *Dev Psychopathol*, 2022, doi: 10.1017/S095457942100153X.

[25] P. Pérez *et al.*, "Conscious processing of narrative stimuli synchronizes heart rate between individuals," *Cell Rep*, vol. 36, no. 11, Sep. 2021, doi: 10.1016/j.celrep.2021.109692.

[26] J. Madsen and L. C. Parra, "Bidirectional brain-body interactions during natural story listening," *Cell Rep*, vol. 43, no. 4, Apr. 2024, doi: 10.1016/j.celrep.2024.114081.

[27] J. Madsen and L. C. Parra, "Cognitive processing of a common stimulus synchronizes brains, hearts, and eyes," *PNAS Nexus*, vol. 1, no. 1, Apr. 2022, doi: 10.1093/pnasnexus/pgac020.

[28] S. Kinreich, A. Djalovski, L. Kraus, Y. Louzoun, and R. Feldman, "Brain-to-Brain Synchrony during Naturalistic Social Interactions," *Sci Rep*, vol. 7, no. 1, Dec. 2017, doi: 10.1038/s41598-017-17339-5.

[29] R. V. Palumbo *et al.*, "Interpersonal Autonomic Physiology: A Systematic Review of the Literature," *Personality and Social Psychology Review*, vol. 21, no. 2, pp. 99–141, May 2017, doi: 10.1177/1088868316628405.

[30] X. Deng *et al.*, "Parental involvement affects parent-adolescents brain-to-brain synchrony when experiencing different emotions together: An EEG-

based hyperscanning study," *Behavioural Brain Research*, vol. 458, Feb. 2024, doi: 10.1016/j.bbr.2023.114734.

[31] A. Anthiyur Aravindan, S. Kalyan Chappidi, A. Thumma, and R. Palanisamy, "Prediction of Arousal and Valence State from Electrodermal Activity using Wavelet based ResNet50 Model," in *Current Directions in Biomedical Engineering*, Walter de Gruyter GmbH, Sep. 2023, pp. 555–558. doi: 10.1515/cdbme-2023-1139.

[32] R. Sánchez-Reolid, F. López de la Rosa, D. Sánchez-Reolid, M. T. López, and A. Fernández-Caballero, "Machine Learning Techniques for Arousal Classification from Electrodermal Activity: A Systematic Review," Nov. 01, 2022, *MDPI*. doi: 10.3390/s22228886.

[33] I. Chatterjee, M. Goršič, J. D. Clapp, and D. Novak, "Automatic Estimation of Interpersonal Engagement During Naturalistic Conversation Using Dyadic Physiological Measurements," *Front Neurosci*, vol. 15, Oct. 2021, doi: 10.3389/fnins.2021.757381.

[34] S. Denys, M. Hofmann, A. van Wieringen, and J. Wouters, "Improving the efficiency of the digit triplet test using digit scoring with variable adaptive step sizes," *Int J Audiol*, vol. 58, no. 10, pp. 670–677, Oct. 2019, doi: 10.1080/14992027.2019.1622042.

[35] M. Metshein, A. Krivošei, A. Abdullayev, P. Annus, and O. Märtens, "Non-Standard Electrode Placement Strategies for ECG Signal Acquisition," *Sensors*, vol. 22, no. 23, Dec. 2022, doi: 10.3390/s22239351.

[36] C. R. Harris *et al.*, "Array programming with NumPy," Sep. 17, 2020, *Nature Research*. doi: 10.1038/s41586-020-2649-2.

[37] P. Virtanen *et al.*, "SciPy 1.0: fundamental algorithms for scientific computing in Python," *Nat Methods*, vol. 17, no. 3, pp. 261–272, Mar. 2020, doi: 10.1038/s41592-019-0686-2.

[38] B. Somers, T. Francart, and A. Bertrand, "A generic EEG artifact removal algorithm based on the multi-channel Wiener filter," *J Neural Eng*, vol. 15, no. 3, Feb. 2018, doi: 10.1088/1741-2552/aaac92.

[39] L. C. Parra, S. Haufe, and J. P. Dmochowski, "Correlated Components Analysis - Extracting

Reliable Dimensions in Multivariate Data," Jan. 2018, doi: 10.51628/001c.7125.

[40] J. Pan and W. J. Tompkins, "A Real-Time QRS Detection Algorithm," *IEEE Trans Biomed Eng*, vol. 32, no. 3, 1985.

[41] C. D. Marci, J. Ham, E. Moran, and S. P. Orr, "Physiologic correlates of perceived therapist empathy and social-emotional process during psychotherapy," *Journal of Nervous and Mental Disease*, vol. 195, no. 2, pp. 103–111, Feb. 2007, doi: 10.1097/01.nmd.0000253731.71025.fc.

[42] W. Biesmans, N. Das, T. Francart, and A. Bertrand, "Auditory-inspired speech envelope extraction methods for improved EEG-based auditory attention detection in a cocktail party scenario," *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, vol. 25, no. 5, pp. 402–412, May 2017, doi: 10.1109/TNSRE.2016.2571900.

[43] J. A. Wegelin, "A Survey of Partial Least Squares (PLS) Methods, with Emphasis on the Two-Block Case," 2000.

[44] J. F. Thayer, A. L. Hansen, E. Saus-Rose, and B. H. Johnsen, "Heart rate variability, prefrontal neural function, and cognitive performance: The neurovisceral integration perspective on self-regulation, adaptation, and health," Apr. 2009. doi: 10.1007/s12160-009-9101-z.

[45] A. Marois, D. Mouratille, Y. Pratviel, C. Chamberland, and S. Tremblay, "Using Cardiac and Electrodermal Activity as Cognitive Markers for Interruptions and Distraction in a Surveillance Simulation," in *Neuroergonomics and Cognitive Engineering*, AHFE International, 2024. doi: 10.54941/ahfe1004733.