Macrogenetic atlas of prokaryotes

Chao Yang[1,*], Hao Huang[1], Naike Wang[1], Xavier Didelot[2], Ruifu Yang[3], Yujun Cui[3], Daniel
Falush[1,*]

[1] The Center for Microbes, Development and Health, CAS Key Laboratory of Molecular Virology
and Immunology, Shanghai Institute of Immunity and Infection, Chinese Academy of Sciences,
Shanghai, China;
[2] School of Life Sciences and Department of Statistics, University of Warwick, UK
[3] State Key Laboratory of Pathogen and Biosecurity, Academy of Military Medical Sciences,
Beijing, China.

* Corresponding authors:
C.Y.: cyang@siii.cas.cn; D.F.: daniel.falush@siii.cas.cn

Abstract

Macrogenetics investigates the patterns and predictors of intraspecific genetic variation across
diverse taxa, offering a framework to address longstanding evolutionary hypotheses. Here, we
present a comprehensive macrogenetic atlas of prokaryotes (MAP), integrating genomic data
(summarized by 30 parameters in 12 categories) from 15,235 prokaryotic species and population
genetic data (35 parameters in 7 categories) from 786 species with phylogenetic, phenotypic, and
ecological data. We illustrate the power and potential of the MAP by investigating the
determinants of fundamental genomic and population genetic parameters. For example, we
provide evidence that genome size in prokaryotes depends on a balance between deletion bias and
the beneficial effects of additional genes. We also investigate how the diversity within prokaryotic
species is structured. We show that long-range and short-range genetic linkage represent quite
different features of evolutionary dynamics and contribute independently to genetic diversity,
contradicting the expectation of neutral theory. Instead, we show that as within-species diversity
increases, selection becomes an increasingly strong force structuring diversity, for example
creating convergent ecospecies structures in *Streptococcus mitis* and *S. oralis*. Overall, our atlas
represents a widely applicable resource and offers novel insights into the drivers of
macroevolution and the life-cycle of prokaryotic species.

Introduction

Over the past 50 years, population genetics has greatly advanced our understanding of fundamental questions regarding the nature of genetic variation and evolution, while providing essential underpinnings for fields such as evolutionary biology and ecology[1]. More recently, the rapid advent of high-throughput sequencing technologies has dramatically propelled this field forward by enabling the generation of large-scale genomic data with unprecedented scope[2,3]. Classical population genetics focuses primarily on genetic diversity and evolutionary forces within individual species. Now, with the ongoing accumulation of data across many taxa, the time is ripe for a novel view—macrogenetics—which embraces broad taxonomic, spatial or temporal scales across dozens to thousands of species[4,5].

Macrogenetics holds significant promise for addressing longstanding evolutionary hypotheses. For instance, recent macrogenetic studies have uncovered a positive correlation between species diversity within a community and intrapopulation genetic diversity in fish and mammals[6,7], providing evidence for the species-genetic diversity correlation (SGDC) concept[8]. Additionally, macrogenetic studies also offer novel insights into what determines genetic diversity[9], revealing the species traits (fecundity and propagule size) underlying genetic diversity[10] and latitudinal gradients in genetic diversity[11].

Prokaryotes make up more biomass than all organisms other than plants combined[12]. Furthermore, a single prokaryotic species can have lineages within it that specialize to multiple distinct niches, for example differing in host range, pathogenicity and nutrient sources, the classic example being *Escherichia coli*[13]. This feature contrasts with eukaryotes where there is typically limited genetic differentiation and specialization within species. This difference raises a number of specific questions such as how are ecological niches partitioned between prokaryotic species, and how does phenotypic and niche diversity affect the genomic and population genetic characteristics of each species. These differences also mean that the process of speciation is likely to be driven by a different set of driving forces, which are collectively less well understood than those in eukaryotes[14].

Here we present a comprehensive macrogenetic atlas of prokaryotes, integrating large-scale genomic and population genetic data with phylogenetic, phenotypic, and ecological metadata. This atlas is a widely applicable resource, and we demonstrate its utility in addressing long-standing evolutionary hypotheses and offering novel insights into the process of speciation.

Results

Overview of macrogenetic atlas of prokaryotes

We analyzed 317,542 assembled genomes across 15,235 species (503 archaeal and 14,732 bacterial species; NCBI taxonomy). Following deduplication and quality control, a representative (*Rep.*) dataset for each species was constructed to minimize biases from repeat sampling (see Methods). To incorporate microevolution data, we also constructed one or more clonal group (CG) datasets for each species, consisting of sets of closely related genomes.

For genomic parameters (GP), we used the median value of *Rep.* genomes as the species estimate.

To estimate population genetic parameters (PGP), we performed random sampling of 10 *Rep.*

genomes for each species across 10 independent replications, accounting for variable sample sizes.

The median PGP value from these replications was used as the species estimate. The CG dataset

was exclusively used to estimate mutation rates and recombination levels. In total, we obtained 30

parameters from 12 categories of GP for all 15,235 species and 35 parameters from 7 categories of

PGP for 786 species (**Fig. 1a,** Supplementary Tables 1 and 2). Moreover, we correlated these

parameters with phylogenetic trees, as well as phenotypic (10 categories) and ecological (4

categories) data retrieved from public databases[15–17].

We refer to the integrated GP and PGP data as the macrogenetic atlas of prokaryotes (MAP). The

parameter values were transformed into rank percentages for each species to make them

distribution-agnostic and to provide a comprehensive overview (**Fig. 1b,c**). We quantified the

correlations among various categories (using r for quantitative data or η for categorical data), as

well as their relationships with the phylogenetic tree (Pagel's λ), demonstrating prevalent

correlations and strong phylogenetic signals for most parameters (**Fig. 1d**). The distribution of

different parameters was assessed (**Fig. 1e**), showing similar GP distributions for archaea and

bacteria, with the exception that archaeal species tend to have shorter gene lengths. Most

parameters exhibited normal-like, left-skewed, or right-skewed distributions, while bimodal

distributions were rarely identified, with the notable exceptions of GC content and codon bias

index (CBI).

Interestingly, we observed that some "well-studied" species (with >100 *Rep.* genomes) exhibit

"extreme" parameter values. For instance, the human pathogen *Helicobacter pylori* displays an

exceptionally high proportion of restriction-modification (RM) system genes, comprising 5.1% of

its genome, ranking first among 15,235 species. Additionally, we identified "extreme" genera. For

example, the cell-wall-deficient helical *Spiroplasma* bacterial genus has an exceptionally low

average gene length of 278 bp, with 7 of the lowest 10 species in gene length belonging to this

genus.

Determinants of genomic characteristics

The MAP enabled a comprehensive investigation of fundamental questions, such as what

determines genome size. To explore this, we extracted several genomic parameter (genome size,

gene length and GC content) related network (r or $\eta \geq 0.3$, **Fig. 2a**) and adjusted for phylogenetic

signals in the correlations (quantified by *r-adj*), due to the significant phylogenetic signals

observed across most parameters. Following adjustment, the correlation coefficients for most

parameter pairs decreased, with a notable reduction in the correlation between genome size and

GC content ($r = 0.5$, *r-adj* $= 0.1$).

Genome size

Genome replication is costly, so natural selection will remove unnecessary sequence. In support of

this prediction, we have found a negative correlation between genome size and growth rate (*r-adj*

$= -0.13$, **Fig. 2b**). Genome streamlining theories, which have found good empirical support in

eukaryotes, therefore predict that species where selection is more effective should have shorter,

more compact genomes[18,19]. In prokaryotes, a convenient way to measure the effectiveness of

selection is using *dN/dS* (the ratio of non-synonymous to synonymous substitution rates), with

lower *dN/dS* or high *dS/dN* values found in species where purifying selection is more effective.

Previous work in bacteria has found that *dN/dS* is negatively correlated with genome size, which

goes against the expectation under the "streamlining" theory, since weaker selection should cause

both a higher *dN/dS* and a higher genome size[19]. It has therefore been proposed that in prokaryotes

stronger selection allows a larger number of beneficial genes to remain in the genome, in the face

of ubiquitous deletion bias[19]. Other authors have found a negative correlation between mutation

rate (*u*) and genome size[20]. We validated the previous observations of negative correlations

between *dN/dS* and genome size (*r-adj* = -0.22), as well as between *u* and genome size (*r-adj* =

-0.26). Furthermore, we showed that the compound parameter *dS/dN×u* (*r-adj* = -0.59) was most

predictive of genome size (**Fig. 2b**). This observation provides additional support for a "balance"

model, with the two factors acting in opposition to each other to determine the overall genome

size.

We have also found several other factors influencing genome size including the habitat,

temperature and oxygen tolerance. For example, aerobic species living in soil and in low

temperatures tend to have larger genomes. Some of these factors, such as gene number,

core-/pan-genes, RM-genes, tRNA counts, and cell size are likely outcome factors of genome size.

Other factors correlated with genome size are metabolic pathways, cell shape and sporulation (**Fig.**

2b).

Gene length

Similar reasoning to the determinants of genome size should also predict that species with more

effective selection should also have longer genes. Gene length is more strongly correlated with

dN/dS (*r-adj* = -0.31) than *u* (*r-adj* = -0.14) or the compound parameter *dS/dN×u* (*r-adj* = -0.29,

Fig 2c), implying that selection rather than deletion bias is a driver of gene length. Moreover,

there is a strong positive correlation between gene length and coding proportion (*r-adj* = 0.48, **Fig.**

2c), which could be explained by non-coding regulatory regions being of similar length for long

and short genes.

GC content

GC content is highly variable in bacteria despite empirical evidence of a universal mutation bias

towards AT[21]. We identified a significant negative correlation between mutation rate and GC

content (*r-adj* = -0.19, **Fig. 2d**), suggesting that a high input of A and Ts through mutation acts to

lower the GC content, while the factors counteracting this mutation bias need time to act. Habitat

emerged as another potential causal factor, with soil-isolated species exhibiting higher GC content.

In contrast, GC content had a significant positive effect on codon bias (*r-adj* = 0.84). Furthermore,

we observed that aerobic species with rod-shaped cells and monotrichous or polar flagella tend to

have an elevated GC content (**Fig. 2d**).

Determinants of population genetic characteristics

Genetic diversity

Classical population genetic theory predicts that in a well-mixed population, synonymous
nucleotide diversity π_s should be proportional to the product on the mutation rate u and the
effective population size N_e[9]. In practice, we do not have a direct measure of N_e for any species,
however in eukaryotes, the negative correlation between propagule size or fecundity and π_s[10] is
likely to be driven by higher N_e of fecund organisms and those with small propagules. In our atlas,
there is a negative correlation (*r-adj* = -0.14) between cell size and within-species π_s (**Fig. 3a,b**),
consistent with the pattern seen in eukaryotes. *dN/dS* can be used as a proxy for N_e[14], and we find
a strong negative correlation between *dN/dS* and π_s (*r-adj* = -0.76), consistent with theory. There is
also a weak correlation between the mutation rate u and π_s (*r-adj* = 0.13). However, the
relationships between other parameters is more complex. Specifically, there is an unexpected
negative correlation between the compound parameter *dS/dN*×u and π_s (*r-adj* = -0.73).
177
In terms of potential ecological factors (**Fig. 3b**), the habitat with the highest within-species π_s are
plant roots, which might reflect the stability of these habitats and the large bacterial population
sizes associated with them. By contrast, fermented food has the lowest diversity, which might
reflect artificial selection but also the bottlenecks associated with the fermentation cycle.
Concordant with this, species involved in fermentation have lower diversity than those that fix
nitrogen. There is also a positive correlation with temperature (*r-adj* = 0.24), similar to the higher
genetic diversity observed in eukaryotic species living in tropical regions[11], which might reflect
warmer habitats supporting larger populations of both prokaryotes and eukaryotes, as well as
seasonal bottlecks in colder climbs.
187
According to population genetic theory, genetic linkage, as measured by linkage disequilibrium
(LD) values within a species are determined by N_e and the recombination rate[22]. In species which
maintain large well-mixed populations for long periods, recombination will break down LD,
meaning that there should be low LD at all genetic distances and high π_s. We have examined
correlations between π_s and LD at different genetic distances. Concordant with theory, there is a
strong negative correlation between short-range LD (10 bp) and π_s (*r-adj* = -0.58). However, while
long-range LD (10 kb) and short-range LD are highly correlated (*r-adj* = 0.68), the correlation
between long-range LD and π_s is not significant. Furthermore, for species with similar levels of
short-range LD, there is a pronounced positive correlation between long-range LD and π_s (**Fig. 4a**),
contradicting the null expectation based on neutral population genetic theory. Thus long-range and
short-range LD reflect evolutionary dynamics that contribute independently in opposite directions
to genetic diversity.
200
Recombination is expected to positively influence genetic diversity, for example by reducing the
fraction of the genome influenced by selective sweeps. We identified significant positive
correlations between π_s and several recombination-related parameters (**Fig. 3b**): the proportion of
genome sites affected by recombination (*Rec. c*[23], *r-adj* = 0.72), the ratio of effects of
recombination and mutation (*r/m*[24], *r-adj* = 0.32) and the ratio of homoplasic and non-homoplasic
alleles (*h/m*[25], *r-adj* = 0.16). The correlations between *c*, *h/m* and π_s are more likely mediated by
LD, given the stronger correlation of these parameters with LD than with π_s (*c* vs r^2 10 bp: *r-adj* =
0.89, *h/m* vs r^2 10 kb: *r-adj* = 0.73). However, the correlation between *r/m* and π_s is stronger than

with LD (r^2 10 bp: *r-adj* = 0.2, r^2 10 kb: *r-adj* = 0.19), suggesting that LD statistics do not capture

the full effect of recombination on diversity.

Mutation rate

The determinants of mutation rate remain poorly understood. One potentially revealing

explanation, the drift-barrier hypothesis, suggests that selection minimizes mutation rates to

enhance replication fidelity, with selection efficiency constrained by genetic drift[26]. However, we

did not observe a significant correlation between *dN/dS* and *u* (*p > 0.05*). Instead, temperature

emerged as a potential causal factor (*r-adj* = 0.23), with species in warmer environments showing

higher mutation rates (**Fig. 3c**).

Gene fluidity

Lifestyle, effective population size, and genome size have been reported as major determinants of

prokaryotic pangenomes[27–29]. Consistent with previous studies, we identified significant

correlations between habitat (η = 0.53), *dN/dS* (*r-adj* = 0.53)*,* genome size (*r-adj* = 0.24) and gene

fluidity (**Fig. 3d**), measured by the ratio of unique gene families to the total gene families in

genome pairs[28]. Gene fluidity positively affects pan-gene number (*r-adj* = 0.63), but has a weak

effect on core-gene number (*r* = -0.1).

We found a strong positive correlation between the number of RM genes (*r-adj* = 0.57) and gene

fluidity. Given the role of RM genes in regulating gene transfer[30], we propose RM genes as a

putative causal factor for gene fluidity. Moreover, the number of mobile genetic elements, such as

phages (*r-adj* = 0.29) and insertion sequence elements (ISE, *r-adj* = 0.2), are also putative causal

factors due to their role in promoting gene transfer.

Other factors, including cell shape, growth rate, metabolic pathways and oxygen tolerance, were

also significantly correlated with gene fluidity, though the causal direction remain unclear. Notably,

a strong connection was found between cell shape and gene fluidity, with rod-shaped species

exhibiting more than a two-fold higher gene fluidity than spherical-shaped species. Furthermore,

the correlations between *u*, *r/m*, and gene fluidity suggest shared drivers and molecular

mechanisms that influence both core and accessory genomic variations, as well as homologous

and non-homologous recombination.

Absence of panmictic species

Many bacterial species disperse effectively around the globe and have enormous population sizes.

Accordingly, we have observed several species with very high genetic diversity, comparable to

that in most diverse eukaryotes[31], such as *Pseudomonas fluorescens* (**Fig. 4a,b**). Classical

population genetic theory suggests that these species should be panmictic, characterized by low

LD at all genetic distances. However, while the most diverse species do have low short-range LD,

consistent with high recombination rates, the diverse species are all highly structured (**Fig. 4a,b**),

reflected by their moderate long-range LD. Consequently, there is a paradoxical absence of

panmictic species.

An example of a well-mixed species with low long-range LD is *Vibrio parahaemolyticus*, which

has moderate diversity. Previous analysis has suggested that diversity of *V. parhaemolyticus* is

moderate because its population size has expanded only recently[32] and that if population size were

maintained, equilibrium diversity would be much higher, and short-range LD would become lower,

which makes it a prime candidate to evolve into a panmictic species. However, *V.*

parahaemolyticus is also in the initial stages of functional diversification into ecospecies, which

are highly differentiated from each other, but only in a small fraction of the genome[33,34]. If

long-range LD increases over time due to an expansion of functional diversification across the

genome, then *V. parahaemolyticus* would not evolve into a panmictic species but instead come to

resemble the structured high diversity species that we have observed.

Selection as a driver of population structure

Our hypothesis is that panmictic bacteria are absent because in species that have maintained large

population sizes, selection will progressively create long-range LD, due to inevitable functional

diversification. To test this hypothesis, we examined the genetic basis of population structure

within bacteria. Specifically, we used principal component analysis (PCA) to identify the major

population structure in each species. We hypothesized that this structure should show stronger

signatures of selection in more diverse species. As a simple measure of selection, we looked for an

enrichment of non-synonymous sites amongst the SNPs most strongly associated with particular

PC components, as measured by SNP loadings. We expect selection to be evident when species

have maintained their current population size for long time periods, which should result in similar

values of short- and long-range LD. Accordingly, we introduce an LD saturation statistic, defined

by the ratio of long-range to short-range LD.

As predicted by the hypothesis, we observed a strong enrichment of such non-synonymous sites

amongst species with high saturation levels (**Fig. 4c**), which we tested statistically by computing

the correlation between enrichment and saturation when the latter is >0.5 (zones 3 and 4 in **Fig.**

4a,c). The correlation is strongest for species with lower levels of short-range LD (zone 4),

indicating selection is more effective for highly recombining species (**Fig. 4c**), supporting our

hypothesis. The absence of correlation for species with low saturation (zones 1 and 2 in **Fig. 4a,c**)

shows that selection only becomes evident in patterns of genome wide diversity for species that

have maintained their current population size for long periods.

Convergent ecospecies

Given the evidence that selection has molded genetic structure, we asked whether we could

identify species further along the road to functional differentiation than *V. parahaemolyticus*. To

this end, we examined species with higher diversity and many genomes (>100 *Rep.* genomes).

Two such species are *Streptococcus mitis* and *S. oralis*. The PCA loading distribution for both

resembled that seen for previous ecospecies[33,34], implying high differentiation in specific regions

and high gene flow elsewhere (**Fig. 4d,e**). Furthermore, there is strong enrichment of

non-synonymous SNPs and genes of particular categories within SNPs with extreme PC1 loadings.

For both species, COG categories D (cell cycle control, cell division, chromosome partitioning)

and M (cell wall/membrane/envelope biogenesis), and KEGG category ko03036 (chromosome

and associated proteins) were invoved, suggesting functional divergence in cell division and

biogenesis. The congruence of the differentiation signal is further supported by a correlation of
0.41 between PC1 gene loadings in the two species and an overlap of the majority of the extreme
PC1 loadings genes in the D, M and ko03036 categories (**Fig. 4f**).
299
To characterize these putative ecospecies, we constructed phylogenetic trees based on SNPs with
extreme loadings (**Fig. 4d,e**). We found two putative ecospecies for *S. mitis* and three for *S. oralis*,
which also correspond to distinct clusters in strain PCA loadings. To determine whether they had
evolved independently, we constructed a joint tree for the shared high loading genes, which imply
independent origin of ecospecies in the two species (**Fig. 4f**). Finally, we constructed phylogenetic
trees ased on remaining SNPs to examine the pattern of differentiation in the rest of the genome.
Distinct ecospecies clusters are visible implying modest genome-wide differentiation, which
contrasts with the absence of differentiation in most of the genome seen in *V. parahaemolyticus*
and *H. pylori*[33,34]. This result is consistent with the *Streptococcus* ecospecies being older and
futher along the path of functional differentiation and helps to explain the higher long-range LD
seen in these two species.
311
312 **Discussion**
313 The MAP resource should be useful to scientists interested in a particular microbe, by providing a
314 simple summary of its genomic, population genetic and other parameters and allowing easy
315 comparison between all available prokaryotic species. It also represents a resource for
316 macrogenomic and population genetic analysis, including raw data and non-redundant datasets.
317
We have illustrated the power and potential of MAP by addressing a series of fundamental
questions about genomic and population genetic parameters, and by comparing our results to those
obtained from eukaryotes, which remain better studied. For example, we provide evidence that
genome size depends on a balance between deletion bias and the beneficial effects of additional
genes[19]. In addition to validating previous observations with a larger dataset, we show that the
compound parameter $dS/dN \times u$, which reflects the combined effects of N_e and mutation rate, is
more than twice as correlated with genome size as previously tested causal factors. Moreover, we
have uncovered a surprisingly strong relationship between cell shape and genome fluidity, with
rod-shaped species having more than two-fold higher fluidity than spherical ones.
327
Our analyses and resources are available based on two taxonomic classifications, namely NCBI
and Genome Taxonomy Database (GTDB)[35,36]. The problem of bacterial species definitions
remains unsolved and large databases inevitably contain errors[37,38]. NCBI is based on user
inputted classifications, while GTDB uses an automatic algorithm, based on a universal average
nucleotide identity cutoff of 0.95. While GTDB has advantages of consistency and quality control
over NCBI, it often splits taxonomically diverse named species into multiple putative species
(Supplementary Fig. 1). In many cases, these species do not correspond to coherent evolutionary
units. In particular, the most diverse species, such as *P. flourescens*, have very low short-range LD,
indicating that gene flow is effective in distributing diversity within the species, and accordingly
they are amongst the best examples of biological species, despite being split into >10 species by
GTDB (Supplementary Fig. 1). Therefore GTDB is not especially well-suited for macrogenomic
analysis, especially for highly diverse species, and we presented resulted based on NCBI

classification. We nevertheless provide an alternative version of parameter calculation and

correlation analysis results based on GTDB (Supplementary Tables 3-4), and show a strong

correlation between parameter relationships derived from two taxonomies ($r = 0.97$,

Supplementary Fig. 2), indicating that the great majority of our results are robust to species

definitions.

In investigating the determinants of genetic diversity, we found that long-range and short-range

LD contribute independently, contradicting the expectation of netural population genetic theory.

Specifically, there is a pronounced positive correlation between long-range LD and π_s for species

exhibiting similar levels of short-range LD. This correlation seems to underly a paradoxical

feature, namely that despite there being several highly diverse species in the data, panmictic

species with low LD at all distances are absent, suggesting they are also rare or absent in nature[32].

Our resolution of the paradox is that as within-species diversity increases, selection becomes an

increasingly strong force structuring diversity.

An important implication of our finding is that we can get insight into the ecological pressures

acting on genetically diverse prokaryotic species, simply by characterizing how their diversity is

structured. One example of structure generated by selection is bacterial ecospecies, which are

characterized by fixed differences in a small fraction of the genome, but high recombination

elsewhere. We have previously described the first two bacterial ecospecies in *V. parahaemolyticus*

and *H. pylori.* In *V. parahaemolyticus* the structure reflects diversification at motility related genes,

while in *H. pylori* differentiation has occurred in genes related to immune escape and nutrient

uptake [33,34]. Both species have moderate diversity and ecospecies structure is restricted to a small

fraction of the genome.

Here we have identified two additional species with ecospecies-like structure, namely *S. mitis* and

S. oralis. These two sister taxa both inhabit the mouths of humans and other animals and there is a

substantial convergence in the genetic basis of the ecospecies structure, with the same cell division

and biogenesis associated genome categories and genes involved, suggesting similar ecological

driving forces and/or adaptive potentials in both species. The ecospecies appear to be older than

those identified previously and show evidence for some level of differentiation genome-wide,

which is likely to be a precursor to full speciation.

Extrapolating from early to late stage ecospecies divergence and beyond, we can begin to

reconstruct the life-cycle of many prokaryotic species. However, ecospecies is only one type of

structure caused by selection. We have previously hypothesized the existence of bacterial

ecoclines, which exhibit continuous variation driven by a quantitative trait[39] and the genetic

structure in the most diverse species such as *P. flourescens* resists any immediate classification.

Characterizing the full range of prokaryotic genetic structures should provide a unique new

window into the forces shaping the most diverse and adaptable organisms on our planet.

Methods
Genome datasets
Deduplication and quality control
We downloaded all assembled genomes (n=317,542) included in the Genome Taxonomy Database
(GTDB; https://gtdb.ecogenomic.org, release 207)[35,36]. We performed deduplications based on the
Biosample accessions, retaining only the genome with highest N50 for each Biosample. Genomes
with completeness <90% or contamination >5% (estimated by CheckM[40]) were excluded.
388
The remaining genomes (n=268,463) were grouped into species based on National Center for
Biotechnology Information (NCBI) taxonomy (taxid) and GTDB classifications. For each species,
the earliest sequenced complete genomes or the genome with highest N50 (if no complete genome
was available) were selected as the reference genomes. All genomes of a species were aligned
against the reference using MUMmer v3.23[41] to generate whole-genome alignments. Genomes
with <60% coverage against the reference (n=4,280) were excluded.
395
Representative datasets and clonal groups
While reliable estimation of population genetic parameters relies on random representative
datasets, biased sampling (*e.g.*, strains from disease outbreaks) is common for public datasets,
especially for pathogenic species. To minimize the effect of repeat sampling, we generated
representative genome datasets. For each species, core-genome (regions present in >95% strains)
single-nucleotide polymorphisms (SNPs) were identified using SNP-sites v2.5.1[42], as previously
described[43]. SNPs located in repetitive regions identified by TRF v4.07b[44] or BLASTN self-search
were excluded. Pairwise SNP-distance between strains were calculated using snp-dists v0.8.2
(https://github.com/tseemann/snp-dists). Genomes of a species were assigned to complete-linkage
clusters (CLCs) based on SNP-distance. For representative datasets of each species, we used a
"relative value" - 1% median pairwise SNP-distance as the cutoff, and selected one genome with
highest N50 from a CLC or these singletons as representative genomes.
408
409 Mutation rate and recombination parameter estimations rely on clonal groups (CGs) consisting of
410 closely related genomes. We used an "absolute value" – 5,000 SNPs (total SNP threshold per
411 branch for reliable recombination detection[45]) as the cutoff to identify CLCs representing CGs.
412
Genomic parameters
We calculated 12 categories of genomic parameters for all the representative genomes, using the
median value of representative genomes to represent each species. Genome size and GC content
were calculated using Quast v5.0.2[46]. We re-annotated representative genomes using Prokka
v1.13[47] and extracted or calculated the gene number and coding proportion based on annotation
results. Codon bias parameters (CBI, Fop, ENc) were estimated using codonW v1.3
(https://codonw.sourceforge.net/) based on coding sequences. The number of tRNA and rRNA
genes was extracted from Prokka annotation results.
421
The standard protein sequences of restriction-modification (R-M) system genes were retrieved
from REBASE[48] and used for searching against annotated protein sequences to identify R-M
genes using BLASTp, as previously described[49]. Repetitive regions of genomes were identified

using TRF v4.07b[44] and BLASTN self-search. CRISPR arrays were identified using

CRISPRidentify v1.1.0[50]. Phage sequences were identified using PhiSpy v3.7.8[51]. Insertion

sequence (IS) elements were identified using ISEScan v1.7.2.3[52]. Default parameters were used

for these analysis.

Population genetics parameters

We calculated 7 categories of population genetics parameters for species with ≥10 representative

genomes and for CGs with ≥10 genomes. To account for the varying number of representative

genomes per species, which can introduce bias in parameter estimation, we performed random

sampling of 10 genomes from all representative genomes. The random sampling was

independently replicated 10 times, and the median value of the parameters from the replications

with ≥5 successful outcomes was used to represent each species.

π_s and dN/dS

For each species, the coding sequence alignment of representative genomes were extracted from

whole-genome alignment. Pairwise non-synonymous divergence (dN), synonymous divergence

(dS), and dN/dS were calculated with PAML v4.10.0[53] using the yn00 algorithm. The mean

pairwise value of dS was used to represent the synonymous nucleotide diversity (π_s).

Recombination

Recombination parameters were calculated for each species using three different methods. For

each species, we calculated five parameters using mcorr[23] based on all the representative genomes:

mutational divergence (θ_{pool}), recombinational divergence (φ_{pool}), mean fragment size (δ), fraction

of the sample diversity resulting from recombination (c), and the relative effect of recombination

to mutation ($r/m = \varphi_{pool}/\theta_{pool}$).

We calculated the ratio of homoplasic and non-homoplasic alleles (h/m) using ConSpeciFix

v1.3.0[25] based on randomly sampled representative genomes, as well as based on all the genomes

for each CG. The median h/m value obtained from random sampling was used to represent each

species.

Using all the genomes for each CG, we calculated the parameters of relative rate of recombination

to mutation (R/θ), mean divergence between donor and recipient (v), δ and r/m using

ClonalFrameML v1.12[24].

Linkage disequilibrium

Linkage disequilibrium (LD) correlation coefficient between two loci (r^2) at distances of 1 bp, 10

bp, 100 bp, 1 kb, and 10 kb was calculated using Haploview v4.2[54] based on randomly sampled

representative genomes. The r^2 values were retained only when the number of loci pairs within the

given distance exceeded 1,000. We also calculated the LD satuation level, defined as the ratio of r^2

between different distances (e.g. r^2 at 10 kb / r^2 at 10 bp).

Core-genome, pan-genome and gene fluidity

For each species, the number of core- and pan-genes was calculated using Panaroo v1.2.8[55] based

on re-annotated results of randomly sampled representative genomes. Gene fluidity was measured

by the mean pairwise ratio between the number of genes not shared and the total number of genes.

Mutation rate

For each CG of a species, we reconstructed maximum-likelihood (ML) phylogenetic trees using

FastTree v 2.1.10[56] based on non-repetitive and non-recombinational SNPs, as previously

described[43]. The isolation year of strains were extracted from the corresponding BioSample

information. We used the ML tree and isolation year information to calculate the mutation rate

with BactDating v1.1.0[57]. Only mutation rates with effective sample sizes (ESS) >100 were

retained. The median mutation rate of CGs was used to represent each species.

Phenotype and ecology data

The phenotype and ecology data were extracted from three sources: BacDive database[15], Madin *et*

al[16] and Secaira-Morocho *et al*[17]. The median value of quantitative data and the major status of

category data of multiple strains was used to represent each species.

Correlation analysis

We used the Python packages pingouin and scipy.stats to calculate correlations between different

parameter types. Specifically, we generated all pairwise combinations of parameters and

performed correlation analyses on parameter pairs containing more than 100 observations. For

quantitative data, we applied three methods to compute the correlation coefficient (r): Pearson,

Spearman, and Kendall. For categorical data, we performed logistic regression and quantified

associations with categorical variables using the eta coefficient (η). When significant correlations

were detected ($p < 0.05$ and coefficient r or $\eta > 0.1$), the highest correlation coefficient was

retained to represent the relationship between the parameters.

We conducted phylogenetic signal analysis on quantitative data parameters using phylolm v2.6.2[58].

The phylogenetic trees based on universally distributed genes (archaea: 53 genes, bacteria: 120

genes) used by GTDB[35,36] were employed in the analysis. Pagel's λ value was used to quantify the

strength of the phylogenetic signal.

Principal component analysis (PCA)

PCA was performed using PLINK v1.9[59] based on core-SNPs with a minor allele frequency (MAF)

threshold of 2%. The SNP loading for each principal component (PC) was calculated as described

in Liu *et al.*[39], reflecting the correlation between SNPs and PCs. Briefly, strain loadings were

derived by multiplying the eigenvectors by the square root of the eigenvalues corresponding to

each PC. SNP loadings were then computed by matrix multiplication between strain loadings and

the SNP matrix.

SNP annotation and gene enrichment analysis

Synonymous and non-synonymous SNPs were annotated using SnpEff v4.3t[60], based on the

re-annotated reference genome. Genes associated with SNPs in the top 1% of SNP loadings on

PC1 were extracted for subsequent analysis. Functional enrichment was performed using the R

package ClusterProfiler v4.2.2[61].

Between-species homologous gene analysis

Homologous genes between *Streptococcus mitis* and *S. oralis* were identified using BLASTp

based on annotated protein sequences, with a minimum identity and coverage of 50%. Protein

sequences of homologous genes associated with SNPs in the top 1% of SNP loadings on PC1 were

used for phylogenetic tree construction using FastTree v2.1.10[56].

Data availability

All genome sequences are available from the Genome Taxonomy Database (GTDB;

https://gtdb.ecogenomic.org, release 207). Essential data files for replication, including

whole-genome alignments, the SNP matrix, sorted metadata, and calculated results, are available

on Zenodo (https://zenodo.org/records/14880463) and Supplementary Tables. Source data are

provided with this paper.

References

1.   Charlesworth, B. & Charlesworth, D. Population genetics from 1966 to 2016. *Heredity* **118**
(2017).

2.   Goodwin, S., McPherson, J. D. & McCombie, W. R. Coming of age: Ten years of
next-generation sequencing technologies. *Nature Reviews Genetics* **17** (2016).

3.   De Coster, W., Weissensteiner, M. H. & Sedlazeck, F. J. Towards population-scale
long-read sequencing. *Nature Reviews Genetics* **22** (2021).

4.   Blanchet, S., Prunier, J. G. & De Kort, H. Time to Go Bigger: Emerging Patterns in
Macrogenetics. *Trends in Genetics* **33**, 579–580 (2017).

5.   Leigh, D. M. *et al.* Opportunities and challenges of macrogenetic studies. *Nature Reviews*
Genetics **22**, 791–807 (2021).

6.   Manel, S. *et al.* Global determinants of freshwater and marine fish genetic diversity. *Nat*
Commun **11** (2020).

7.   Theodoridis, S. *et al.* Evolutionary history and past climate change shape the distribution
of genetic diversity in terrestrial mammals. *Nat Commun* **11** (2020).

8.   Vellend, M. & Geber, M. A. Connections between species diversity and genetic diversity.
Ecology Letters **8** (2005).

9.   Ellegren, H. & Galtier, N. Determinants of genetic diversity. *Nature Reviews Genetics* **17**,
422–433 (2016).

10.   Romiguier, J. *et al.* Comparative population genomics in animals uncovers the
determinants of genetic diversity. *Nature* **515**, 261–263 (2014).

11.   Miraldo, A. *et al.* An Anthropocene map of genetic diversity. *Science (1979)* **353** (2016).

12.   Bar-On, Y. M., Phillips, R. & Milo, R. The biomass distribution on Earth. *Proceedings of*
the National Academy of Sciences **115**, 6506–6511 (2018).

13.   Denamur, E., Clermont, O., Bonacorsi, S. & Gordon, D. The population genetics of
pathogenic Escherichia coli. *Nat Rev Microbiol* **19**, 37–54 (2021).

14.   Charlesworth, B. Effective population size and patterns of molecular evolution and
variation. *Nat Rev Genet* **10** (2009).

15.   Reimer, L. C. *et al.* BacDive in 2022: The knowledge base for standardized bacterial and
archaeal data. *Nucleic Acids Res* **50**, (2022).

16.   Madin, J. S. *et al.* A synthesis of bacterial and archaeal phenotypic trait data. *Sci Data* **7**,
(2020).

17.   Secaira-Morocho, H., Chede, A., Gonzalez-de-Salceda, L., Garcia-Pichel, F. & Zhu, Q. An
evolutionary optimum amid moderate heritability in prokaryotic cell size. *Cell Rep* **43**,
114268 (2024).

18.   Lynch, M. & Conery, J. S. The Origins of Genome Complexity. *Science (1979)* **302**,
1401–1404 (2003).

19.   Sela, I., Wolf, Y. I. & Koonin, E. V. Theory of prokaryotic genome evolution. *Proc Natl*
Acad Sci U S A **113**, 11399–11407 (2016).

20.   Bourguignon, T. *et al.* Increased Mutation Rate Is Linked to Genome Reduction in
Prokaryotes. *Current Biology* **30**, 3848-3855.e4 (2020).

21.   Hershberg, R. & Petrov, D. A. Evidence that mutation is universally biased towards AT in
bacteria. *PLoS Genet* **6** (2010).

22.   Slatkin, M. Linkage disequilibrium - Understanding the evolutionary past and mapping the
medical future. *Nature Reviews Genetics* **9** (2008).

23.   Lin, M. & Kussell, E. Inferring bacterial recombination rates from large-scale sequencing
datasets. *Nat Methods* **16**, 199–204 (2019).

24.   Didelot, X. & Wilson, D. J. ClonalFrameML: Efficient Inference of Recombination in
Whole Bacterial Genomes. *PLoS Comput Biol* **11**, 1–18 (2015).

25.   Bobay, L. M., Ellis, B. S. H. & Ochman, H. ConSpeciFix: Classifying prokaryotic species
based on gene flow. *Bioinformatics* **34**, (2018).

26.   Lynch, M. *et al.* Genetic drift, selection and the evolution of the mutation rate. *Nature*
Reviews Genetics **17** (2016).

27.   Dewar, A. E., Hao, C., Belcher, L. J., Ghoul, M. & West, S. A. Bacterial lifestyle shapes
pangenomes. *Proceedings of the National Academy of Sciences* **121** (2024).

28.   Andreani, N. A., Hesse, E. & Vos, M. Prokaryote genome fluidity is dependent on effective
population size. *ISME Journal* **11**, 1719–1721 (2017).

29.   Bobay, L.-M. & Ochman, H. Factors driving effective population size and pan-genome
evolution in bacteria. *BMC Evol Biol* **18**, 153 (2018).

30.   Oliveira, P. H., Touchon, M. & Rocha, E. P. C. Regulation of genetic flux between bacteria
by restriction-modification systems. *Proc Natl Acad Sci U S A* **113**, 5658–5663 (2016).

31.   Dey, A., Chan, C. K. W., Thomas, C. G. & Cutter, A. D. Molecular hyperdiversity defines
populations of the nematode Caenorhabditis brenneri. *Proc Natl Acad Sci U S A* **110**,
(2013).

32.   Yang, C., Cui, Y., Didelot, X., Yang, R. & Falush, D. Why panmictic bacteria are rare.
bioRxiv (2018).

33.   Cui, Y. *et al.* The landscape of coadaptation in Vibrio parahaemolyticus. *Elife* **9**, e54136
(2020).

34.   Tourrette, E. *et al.* An ancient ecospecies of Helicobacter pylori. *Nature* **635**, 178–185
(2024).

35.   Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat Biotechnol* **36**, 996 (2018).

36.   Parks, D. H. *et al.* GTDB: An ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy.
Nucleic Acids Res **50**, D785–D794 (2022).

37.   Rosselló-Mora, R. & Amann, R. The species concept for prokaryotes. *FEMS Microbiol*
Rev **25**, (2001).

38.   VanInsberghe, D., Arevalo, P., Chien, D. & Polz, M. F. How can microbial population
genomics inform community ecology? *Philosophical Transactions of the Royal Society B:*
Biological Sciences **375**, 20190253 (2020).

39.   Liu, S., Svensson, S. L. & Falush, D. A putative ecocline in Klebsiella pneumoniae.
bioRxiv (2025).

40.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
Assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043–1055 (2015).

41.   Delcher, A. L., Salzberg, S. L. & Phillippy, A. M. Using MUMmer to Identify Similar
Regions in Large Sequence Sets . *Curr Protoc Bioinformatics* **00**, (2003).

42.   Page, A. J. *et al.* SNP-sites: rapid efficient extraction of SNPs from multi-FASTA
alignments. *Microb Genom* **2**, e000056 (2016).

43.   Yang, C. *et al.* Outbreak dynamics of foodborne pathogen Vibrio parahaemolyticus over a
seventeen year period implies hidden reservoirs. *Nat Microbiol* **7**, 1221–1229 (2022).

44.   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*
Res **27**, 573–580 (1999).

45.   Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant
bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* **43**, e15 (2015).

46.   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

47.   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069
(2014).

48.   Roberts, R. J., Vincze, T., Posfai, J. & Macelis, D. REBASE: a database for DNA
restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res* **51**, (2023).

49.   Yang, C. *et al.* Recent mixing of Vibrio parahaemolyticus populations. *ISME J* **13**,
2578–2588 (2019).

50.   Mitrofanov, A. *et al.* CRISPRidentify: Identification of CRISPR arrays using machine
learning approach. *Nucleic Acids Res* **49**, (2021).

51.   Akhter, S., Aziz, R. K. & Edwards, R. A. PhiSpy: A novel algorithm for finding prophages
in bacterial genomes that combines similarity-and composition-based strategies. *Nucleic*
Acids Res **40**, (2012).

52.   Xie, Z. & Tang, H. ISEScan: automated identification of insertion sequence elements in
prokaryotic genomes. *Bioinformatics* **33**, (2017).

53.   Yang, Z. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**,
(2007).

54.   Barrett, J. C. Haploview: Visualization and analysis of SNP genotype data. *Cold Spring*
Harb Protoc **4**, (2009).

55.   Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo
pipeline. *Genome Biol* **21**, 180 (2020).

56.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately
644   Maximum-Likelihood Trees for Large Alignments. *PLoS One* **5**, e9490 (2010).
645   57.   Didelot, X., Croucher, N. J., Bentley, S. D., Harris, S. R. & Wilson, D. J. Bayesian
646   inference of ancestral dates on bacterial phylogenetic trees. *Nucleic Acids Res* **46**, (2018).
647   58.   Tung Ho, L. S. & Ané, C. A linear-time algorithm for gaussian and non-gaussian trait
648   evolution models. *Syst Biol* **63**, (2014).
649   59.   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-based
650   linkage analyses. *Am J Hum Genet* **81**, 559–575 (2007).
651   60.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
652   polymorphisms, SnpEff. *Fly (Austin)* **6**, (2012).
653   61.   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
654   *The Innovation* **2**, 100141 (2021).
655

Acknowledgements
This work was supported by National Key Research and Development Program of China (No.
2022YFC2304700 to F.D.), National Natural Science Foundation of China (No. 32270003 and No.
32000008 to C.Y., No. 32170640 and No.32211550014 to D.F.), Youth Innovation Promotion
Association, Chinese Academy of Sciences (No. 2022278 to C.Y.) and Shanghai Rising-Star
Program (No. 23QA1410500 to C.Y.).
662

663   **Author contributions**
664   C.Y., D.F., R.Y., and Y.C. initiated and coordinated the study. C.Y., H.H. and N.W. contributed to
665   data collectionand and performed bioinformatics analysis. All authors contributed to interpretation
666   of the data. C.Y. and D.F. wrote the first draft of the paper and D.X., R.Y., and Y.C. reviewed and
667   revised the paper. All authors read and approved the final manuscript.
668

669   **Competing interests**
670   The authors declare no competing interests.
671

Figure legends

Figure 1. Overview of the macrogenetic atlas of prokaryotes (MAP). a) Study flowchart. **b,c)**

Heatmaps displaying the rank percentages of genomic parameters (GP) and population genetic

parameters (PGP) for each species. The bars at the top indicate the kingdom and phylum of each

species, while the heatmap colors represent rank percentages. **d)** Correlation heatmap among

various parameters. Phylogenetic signals were quantified using Pagel's λ. **e)** Distribution of

different GP and PGP. Red and blue colors represent archaea and bacteria, respectively. PGP is

shown only for bacteria, as few archaeal species met the criteria for PGP estimation (≥10 *Rep.*

genomes).

Figure 2. Determinants of genomic characteristics. a) Correlation network of three GPs

(genome size, gene length and GC content, r or $\eta \geq 0.3$) and related parameters. Node color

represents parameter categories, while line color indicates the type or direction of the correlation.

Line width and transparency indicate the strength of the correlation coefficient. **b-d)** Three GPs

related parameters before and after phylogenetic signal adjustment ($r\text{-}adj \geq 0.1$). Bar color

indicates the direction or type of correlation. Green and orange stars indicate putative causal and

consequential factors, respectively. Violin and box plots display the distribution of GPs across

different categories, with only categories containing ≥100 species shown. Point density plots show

detailed correlations between the GPs and their putative causal/outcome factors, with color

representing point density.

Figure 3. Determinants of population genetics characteristics. a) Correlation network of three

PGs (genetic diversity, mutation rate, and gene fluidity, r or $\eta \geq 0.2$) and their related parameters.

Node and line color, width and transparency are as in Fig. 2a. **b-d)** Three PGs related parameters

before and after phylogenetic signal adjustment ($r\text{-}adj \geq 0.1$). Bar and point colors are as in Fig.

2b-d. Only categories containing ≥10 species are shown.

Figure 4. Determinants of population structure and ecospecies. a) Relationships between

short-range LD (10 bp), long-range LD (10 kb), and genetic diversity. Point colors represent

genetic diversity levels. Species with ≥100 representative genomes are outlined in black. Blue

dashed lines indicate LD saturation levels of 0.5 (y=0.5x) and 1 (y=x). **b)** Phylogenetic trees of

seven selected species. **c)** Correlations between LD saturation and N/S (top 1% vs other SNPs),

the ratio of non-synonymous to synonymous sites (N/S) for SNPs in the top 1% of SNP loadings

on PC1 (SNP-top1%-PC1) relative to the ratio of N/S for SNPs outside the top 1%

(SNP-other-PC1). **d,e)** Top: Principal components analysis (left, point colors indicate inferred

ecospecies) and functional enrichment of genes related to SNP-top1%-PC1 (right) in

Streptococcus mitis and *S. oralis.* Middle: Distribution of squared SNP loadings on PC1 across the

genome, with point colors indicating COG or KEGG categories. Dashed lines represent the top 1%

SNP loading cutoff. Bottom: Phylogenetic trees based on SNPs with top and bottom 1% SNP

loadings on PC1 and other SNPs. Branch colors indicate inferred ecospecies. **f)** Top: Shared

homologous genes associated with SNP-top1%-PC1 between two species. Middle: Correlation of

gene loading (average SNP loading per gene) between species. Point colors indicate COG or

KEGG categories. Bottom: Phylogenetic trees based on protein sequences of shared genes

associated with SNP-top1%-PC1 between two species.