Freshwater snail faecal metagenomes reveal environmental reservoirs of antimicrobial resistance genes across two continents

Angus M. O'Ferrall [a], Alexandra Juhász [a, b], Sam Jones [a], Peter Makaula [c], Gladys Namacha [c], Shaali Ame [d], David Oguttu [e], Aidan Foo [a, f], Sekeleghe A. Kayuni [a, c], James E. LaCourse [a], Janelisa Musaya [c, g], J. Russell Stothard [a], Adam P. Roberts [a] *

[a] Department of Tropical Disease Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

[b] Institute of Medical Microbiology, Semmelweis University, Budapest, Hungary

[c] Malawi Liverpool Wellcome Programme, Queen Elizabeth Central Hospital, Blantyre, Malawi

[d] Neglected Tropical Diseases Program, Ministry of Health, Stone Town, Zanzibar, Tanzania

[e] Vector Control Division, Ministry of Health, Kampala, Uganda

[f] Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

[g] Department of Pathology, School of Medicine and Oral Health, Kamuzu University of Health Sciences, Blantyre, Malawi

* Corresponding author – E-mail: adam.roberts@lstmed.ac.uk

Abstract

The transfer of antimicrobial resistance genes (ARGs) from environmental microbes to pathogens is a critical but underexplored One Health driver of antimicrobial resistance (AMR). Here, we evaluate freshwater snails, which are geographically widespread aquatic invertebrates, as environmental reservoirs of ARGs. We collected faeces from eight gastropod genera at 15 freshwater locations across Malawi, Uganda, Zanzibar, and the United Kingdom, and conducted the first freshwater snail faecal metagenomic sequencing study. Proteobacteria dominated the faecal microbiomes, within which Aeromonadales were strongly associated with the total ARG load. We detected putative ARGs conferring resistance to 13 distinct antibiotic classes, including carbapenem resistance genes in all countries. In Malawi, we linked $bla_{OXA-181}$ (bla_{OXA-48}-like), a putatively mobile carbapenemase gene, to *Shewanella xiamenensis*—the gene's known progenitor—in a metagenome-assembled genome. We detected another bla_{OXA-48}-like variant by read-mapping from a sample in the United Kingdom and identified mobile colistin resistance (*mcr*)-like genes at 11 of 15 locations. Our findings highlight freshwater snail microbiomes as One Health-relevant, environmental reservoirs of clinically important ARGs.

Keywords

AMR, Carbapenemase, CphA, Gastropoda, MCR, One Health, OXA-181

 MAIN TEXT

Antimicrobial resistance (AMR) is a growing global health emergency driven by industrial scale

use of antimicrobials in medicine, veterinary practice, and agriculture since their discovery in

the twentieth century (1). The impact of AMR on human health is largely driven by the

acquisition of antimicrobial resistance genes (ARGs) in pathogenic bacteria (2). Consequently,

most data used to investigate AMR originate from disease-causing organisms, yet there is

increasing recognition that One Health frameworks accounting for the interdependence of

human, animal, and environmental health systems need consideration to better understand

the origins, drivers, and spread of AMR (3). This approach is essential for assessing how AMR

may be linked to crossing planetary boundaries, such as those related to freshwater use,

biodiversity loss, and climate change (4).

Outside the context of AMR, freshwater gastropod snails are already of interest in a One

Health context. Snails are globally distributed in aquatic environments, often playing critical

roles as obligate intermediate hosts in the transmission of neglected tropical diseases when

freshwater sources are contaminated with human and animal excreta. For instance, snails of

the *Bulinus*, *Biomphalaria*, and *Oncomelania* genera are key in the transmission of

schistosomiasis (5), while *Lymnaea* snails transmit fascioliasis (6). Recent efforts to

characterize the microbiomes of medically important freshwater snails are generally motivated

by the growing recognition that host-associated microorganisms significantly influence host-

parasite interactions (7). The microbiomes of *Biomphalaria* snails have been shown by 16S

taxonomic profiling to be diverse and distinct from those of their freshwater environments (8),

yet bacteriome transplant experiments have demonstrated the potential for these microbiomes

to change depending on the surrounding environmental bacteria (9). Freshwater snail

populations are highly mobile (10), capable of translocating due to changes in ecosystems

caused by flooding or the formation of artificial water bodies (11,12), and are distributed across

a range of human-influenced areas (13,14). Considering that the environments where

freshwater snails graze—typically on detritus—can be contaminated with AMR enteric bacteria

at sites of interaction with humans and animals, we wanted to explore the potential role of

snails as widespread and persistent environmental reservoirs of ARGs (15). This may be

particularly relevant in Low- and Middle-Income Countries (LMICs), where AMR is

disproportionately linked to increased mortality (16), and where poor environmental health

infrastructure is linked to AMR transmission across various One Health compartments (17,18).

The analysis of metagenomic sequence data is increasingly being used in the study of

complex microbial samples and AMR, made possible by advances in next-generation

sequencing and computational science technologies in recent years (19). Metagenomic

studies are not reliant on culture steps that alter microbial communities and are therefore able
to yield non-selective data on sample composition (20). Metagenomic analyses generally
utilise short-read sequence data, which when coupled with the complexity of microbiome
samples poses challenges in the assembly of metagenome-assembled genomes (MAGs) for
low-abundance organisms without ultra-deep sequencing (21,22). Approaches that map reads
directly to reference databases are therefore useful for taxonomic profiling and the detection
of low-abundance target genes as they do not rely on metagenome assembly, and have
proved to be more sensitive than assembly-based approaches for the detection of ARGs in
complex samples such as faeces (23). When combining read-based profiling with assembly-
based approaches that uncover the genomic context of AMR—linking ARGs to specific hosts
and mobile genetic elements (MGEs)—metagenomics can enable a comprehensive
understanding of the resistome and its potential impacts on microbial ecology and global
health.

Here, we present a metagenomic dataset characterising the first freshwater snail faecal
metagenomes from 15 locations within Malawi, Uganda, Zanzibar (a self-governing region
within the Republic of Tanzania) and the United Kingdom (*Supplementary Table 1*).
Acknowledging that freshwater snail microbiomes vary by organ (24), and to mirror the natural
mechanisms of faecal bacterial transfer from gastropods to the environment, we sampled
faeces excreted by snails immediately after they were collected from their habitats, rather than
extracting DNA directly from snail organs or rearing snails under laboratory conditions prior to
DNA extraction. We use whole-genome metagenomics to investigate the role of freshwater
gastropods as reservoirs, and potentially vectors, of AMR. Our goal is to identify clinically
relevant ARGs within these common aquatic invertebrates and to examine their genomic
context, providing insights and targets for future studies on ARG transfer to human and animal
pathogens. We utilize direct read mapping for taxonomic profiling and ARG identification,
followed by less sensitive but highly insightful assembly-based methods to explore the
genomic context of ARGs.

Results

Sample characteristics

In 2023 and 2024, we collected freshwater snails at 15 freshwater locations in four
administrative regions across two continents, targeting snails of a single genus at each
location. In Africa, collections were made from six locations in Malawi (five *Bulinus* and one
Cleopatra), three locations in Uganda (one each of *Biomphalaria*, *Bulinus*, and *Physella*), and
four locations in Zanzibar (one each of *Bulinus*, *Cleopatra*, *Lanistes*, and *Pila*). In Europe, two
locations in the United Kingdom were sampled (one *Anisus* and one *Lymnaea*). At each

location, groups of 50 snails from the identified genus were gathered ahead of faecal harvesting. For downstream analysis, we pooled the faeces of each group of snails to obtain enough biomass for DNA extraction and Illumina short-read sequencing, resulting in 15 metagenomes, each representing a specific snail genus at a specific location (*Supplementary Table 1*). The number of 150 bp Illumina paired end reads per sample ranged from 9,340,457 to 35,991,622.

The freshwater snail faecal microbiome is diverse, varied, yet consistently rich in Proteobacteria

After read processing and quality control, reads classified as bacterial by Kraken2 underwent species abundance estimation with Bracken (25). Proteobacteria was the most abundant phylum observed, with a mean relative abundance of 66.27% (*Figure 1a*). Enterobacterales, an order within the Proteobacteria containing multiple species of clinical concern due to their ability to acquire and exchange ARGs, were well represented, with a mean relative abundance of 4.79%. The Uk1 sample was a notable outlier in which the relative abundance of Enterobacterales was estimated to be 41.70%. When excluding Uk1, the mean relative abundance of Enterobacterales dropped to 2.15%. Four orders were more abundant than Enterobacterales (Burkholderiales, Rhizobiales, Aeromonadales, and Pseudomonadales) all of which had a mean relative abundance of greater than 5% (*Figure 1b*). All samples exhibited high community (alpha) diversity. Shannon's diversity index ranged from 3.881 (M5) – 7.372 (Z2) (mean = 6.474, median = 6.888). Simpson's diversity index, in which scores from 0 – 1 are possible (1 = maximum possible diversity), ranged from 0.793 (M5) – 0.999 (Z2) (mean = 0.972, median = 0.994). We observed that alpha diversity (Shannon's diversity index) was negatively corelated with the relative abundance of Proteobacteria (Pearson correlation coefficient = -0.600; *p* = 0.0184).

Relative abundance data were aggregated at the order level, subjected to centred log-ratio (CLR) transformation, and subsequently standardized (CLR-z) to facilitate visual comparisons between taxa with varying absolute abundances (*Figure 1c*). From the 160 bacterial orders identified across all samples, the 50 most abundant orders, accounting for 97.92% of the total microbiome, were selected for hierarchical clustering and visualization. Samples from the same administrative region clustered closely together, with the exception of Malawian samples, which were split into two distinct groups. This split corresponded to the two districts where the samples were collected: samples M1, M2, M3, and M4, collected in Mangochi District on the southern shorelines of Lake Malawi, clustered together, while samples M5 and M6, collected in Nsanje District, over 275 km from Mangochi District sampling sites, formed a separate cluster. Although they clustered together, northern hemisphere samples (United

Kingdom) did not form a highly divergent cluster despite likely differences in climate-related and environmental conditions. No clear trend in taxonomic profiles was observed by snail genus. Microbiomes from *Bulinus* snails (the most-represented genus, obtained from seven locations) showed notable divergence: the sample collected in Zanzibar clustered at one extreme of the dendrogram alongside samples from other snail genera, distinct from *Bulinus* samples collected in Uganda and the two Malawian districts.

Figure 1. Exploration and comparison of bacterial taxa within freshwater snail faeces: a) Relative abundance of the top phyla; b) Relative abundance of the top orders; c) Clustered

heatmap of standardised order abundances (CLR-z) across snail hosts and geographic locations. Column and row trees cluster samples and taxa, respectively, based on a hierarchical clustering approach.

When investigating at species level, a Bray-Curtis dissimilarity (beta diversity) matrix (*Supplementary Table 2*) showed a range of divergence between Malawian *Bulinus* snails. We observed moderate-high divergence (0.649) between two samples (M2, M4) collected a little over five kilometres from each other, both from stagnant water bodies close to the Lake Malawi shoreline, suggesting that local environmental factors play a key role in influencing the microbiome.

Proteobacteria, specifically Aeromonadales, drive the ARG load

Using a read-based k-mer alignment (KMA) method to screen for ARGs against the ResFinder database (26,27), reads mapped to 54 unique ARGs in total (minimum 90% query identity and 60% template coverage), belonging to 41 ARG Cluster90s (in which reference genes share minimum 90% nucleotide identity) (23). ARG diversity (the number of unique ARGs identified by read mapping in any given sample) ranged from 1 – 16 (mean = 7.07, median = 6). We detected fewer ARGs by screening metagenome assemblies against the ResFinder database (mean = 2.53, median = 2) (*Supplementary Figure 1*). Across all samples, 10 of the 38 ARGs (26.3%) assembled on contiguous sequences (contigs) could be binned to high-quality (>90% completeness, <5% contamination) or medium-quality (>50% completeness, <10% contamination) metagenome-assembled genomes (MAGs). These ARGs were all binned to MAGs classified as Proteobacteria (genera: *Aeromonas*, n = 7; *Shewanella*, n = 1; *Stenotrophomonas*, n = 2) (*Figure 5*).

We calculated the normalised abundance of each ARG from read-mapping data, expressed as RPKM (Reads Per Kilobase per Million mapped bacterial reads), then summed the RPKM of all ARGs to calculate total ARG load per sample. ARG load ranged from 3.16 – 357.04 RPKM (mean = 87.99, median = 59.62). The relationship observed between relative abundance of Proteobacteria (the predominant bacterial order) and ARG load appeared to be non-linear and was linearised with a log transformation of ARG load (*Supplementary Figure 2*). Proteobacteria relative abundance was positively correlated with log-transformed ARG load in faecal microbiomes (Pearson correlation coefficient = 0.684; *p* = 0.0049) (*Figure 2a*). A significant association between Proteobacteria relative abundance and ARG diversity was not detected (Pearson correlation coefficient = 0.122; *p* = 0.6654). Among the 20 orders of bacteria with a mean relative abundance above 1%, Aeromonadales (belonging to the Proteobacteria) were strongly and positively correlated with ARG load (Pearson correlation

coefficient = 0.918; *p* < 0.0001) (*Figure 2b*), while no other orders were associated with either positive or negative correlation (*Supplementary Table 3*).

Figure 2. Associations between Proteobacteria and ARG load: a) Relative abundance of Proteobacteria versus log(ARG load) with a linear regression line and associated Pearson correlation coefficients indicating improved fit of data following log-transformation of ARG load; b) Relative abundance of the five most abundant orders within Proteobacteria versus ARG load with associated Pearson correlation coefficients indicating that Aeromonadales contribute significantly to ARG load within the Proteobacteria. Within-plot Labels in part (a) are snail genera. Relationships between relative abundance and ARG load in part (b) either appeared linear (Aeromonadales) or non-correlated (others), hence transformations were not applied.

ARGs conferring resistance to first- and last-line antibiotics detected

When we aggregated ARG relative abundance (RPKM) by class (i.e., antibiotic that the ARGs confer resistance to), beta-lactamase genes (excluding carbapenemases) were the most abundant, contributing to 53.58% of the total ARG load across samples, followed by genes conferring resistance to colistin (14.38%), carbapenems (10.27%), sulfonamides (5.32%), tetracyclines (5.12%), aminoglycosides (2.77%), fosfomycin (2.15%), macrolides (2.14%), trimethoprim (1.90%), lincosamides (0.51%) chloramphenicol (0.47%), fusidic acid (0.19%),

and quinolones (0.03%) (*Supplementary Figure 3*). Efflux pump genes conferring multidrug

resistance contributed 1.16% of the total ARG load.

Beta-lactamase genes from the OXA-12 Cluster90 were the most prevalent in our samples

and were detected in fourteen of the fifteen metagenomes (*Figure 3*). We also detected ARGs

from the CphA Cluster90, a family of carbapenem-hydrolyzing metallo-beta-lactamase genes,

in ten of the fifteen samples. Three OXA-carbapenemase genes were identified, each from a

different country. These included two ARGs from the OXA-48 Cluster90 ($bla_{OXA-181}$ and $bla_{OXA-547}$,

in Malawi and the United Kingdom, respectively) of concern globally due to plasmid-borne

dissemination amongst Enterobacterales (28), and $bla_{OXA-212}$ (Uganda) from the OXA-211

Cluster90 of carbapenemase genes that naturally occur in *Acinetobacter johnsonii* (29). We

also detected putative colistin resistance genes, predominantly variants of the *mcr-7.1* gene,

in eleven metagenomes.

Figure 3. Heatmap of normalized ARG Cluster90 abundances across samples, expressed as

log(RPKM). Cluster90s are colour-coded by the antibiotic class that they are considered to

confer resistance against.

MGEs in snail faecal microbiomes are associated with ARGs

To investigate the relationship between MGEs and ARGs in metagenome assemblies, we screened all contigs for insertion sequence (IS) elements (ISfinder database (30)) in addition to ARGs (ResFinder database (27)). We identified two contigs containing IS elements and ARGs, neither of which could be binned to a putative MAG. A 4042 bp contig (Ug2 NODE_1302: *Bulinus* spp. faeces, Lake Victoria, Uganda) contained *sul1* (sulfonamide resistant dihydropteroate synthase) and *bla*OXA-10 (beta-lactamase) genes, on which both ARGs were located upstream of IS*6100*. The gene cluster also contained *qacEdelta* (a gene conferring resistance to antiseptics) and *yhbS* (putative N-acetyltransferase). The contig displays 100% query cover with one complete genome in the GenBank Nucleotide database— *Aeromonas hydrophila* (accession no. AP025277)—with 99.97% identity, but also maintains high query cover (> 95%) and identity (> 99%) with chromosomal and plasmid sequences from various other bacteria (*Pseudomonas aeruginosa*, *Vibrio cholerae*, *Morganella morganii*, *Salmonella enterica*, *Klebsiella pneumoniae*). A second, shorter, 3505 bp contig (M2 NODE_687: *Bulinus* spp. faeces, Mangochi District, Malawi), aligned with 99.94% nucleotide identity to Ug NODE_1302 over the region containing *qacEdelta, sul1* and *yhbS* upstream of IS*6100* (*Figure 4*), with a single nucleotide polymorphism (SNP) in the *sul1* gene. The similarity observed between the two contigs from Ug2 and M2, and with sequences from a range of bacterial species in GenBank, demonstrates the widespread nature of certain ARGs associated with MGEs, which we show can be found in freshwater snail faeces.

*Figure 4: Gene cluster comparison between Ug2 NODE_1302 and M2 NODE_687 containing bla*OXA-10 *(Ug2 only), qacEdelta1, sul1 and yhbS upstream of IS6100. Regions coding for hypothetical proteins are not displayed.*

We detected between 8 and 97 putative IS elements per sample (mean = 37.8, median = 29). However, we could only link these directly to ARGs on contigs in the above two examples. Searches for ARGs associated with IS elements in transposons should ideally capture 24.34 kbp of DNA (based on the length of composite and unit transposons in The Transposon

Registry) (31). Due to the limitations in assembling large contigs in diverse metagenomic

samples (22), and the associated inability to capture all ARGs and MGEs on contigs (the

median length of contigs containing ARGs in our dataset is 2.10 kbp), we constructed a

generalized linear model (*Supplementary Information 1*) with a negative binomial distribution

to model the relationship and show associations where MGEs and ARGs could not be directly

linked by assembly. The choice of negative binomial distribution was made to account for

overdispersion in the ARG count data, which is common in datasets involving count outcomes

such as this where the variance in the number of ARGs assembled per sample (4.12) exceeds

the mean (2.53).

Table 1: Outputs of a negative binomial generalized linear model exploring associations

between counts of ARGs and IS elements in metagenome assemblies.

Predictor	Estimate	Standard Error	95% confidence interval	*p*-value
Intercept (β_0)	-17.3585	0.3934	-18.1757, -16.6292	<0.0001
IS elements assembled (β_1)	0.01393	0.0067	0.0007, 0.0271	0.0383

In tangible terms, the model outputs (*Table 1*) indicate that the addition of 10 IS elements to a

snail faecal microbiome is associated, on average, with a 14.9% increase in the number of

ARGs (assuming that the assembly length remains constant), while ARGs are expected to be

identified at a rate of 0.030 per Mb assembly (or 30 per Gb), in the absence of IS elements

(solutions available in *Supplementary Information 1*).

Progenitors of mobile carbapenemase and colistin resistance genes identified

One of the two *bla*$_{OXA-48}$-like carbapenemase genes identified by read-mapping was also

detected in metagenome assemblies. NODE_173 from the M5 metagenome (*Cleopatra* spp.

faeces, Nsanje District, Malawi) contained *bla*$_{OXA-181}$ on a 20996 bp contig with 17 open-

reading frames (ORFs). The contig was binned to a high-quality MAG classified as *Shewanella*

xiamenensis (*Figure 5*). When the *bla*$_{OXA-181}$-containing contig was queried against the

GenBank Nucleotide database, the closest alignment was to chromosomal DNA from an *S.*

xiamenensis isolate (accession no. AP026732) obtained from urban drain water from Vietnam

(100% query cover, 98.51% identity) (32). With only one SNP in a non-coding region, the

contig also aligns with 99.92% identity over a 1309 bp region (8073 – 9381) to various

plasmids from the Enterobacterales containing the internationally disseminated *bla*$_{OXA-181}$ gene

and a truncated *lysR* (Δ*lysR*) gene, including the KP3-A plasmid from which *bla*$_{OXA-181}$ was first

characterised in a clinical *K. pneumoniae* isolate (accession no. JN205800) (33). The region

containing *bla*$_{OXA-181}$ and Δ*lysR* within the KP3-A (and other) plasmid(s) can be traced back to

its origin, *S. xiamenensis*, where *bla*$_{OXA-181}$ is found adjacent to a complete *lysR* gene, as

previously shown (34) and also visible in our annotated sequences (*Supplementary Figure 5*). The ability to detect, assemble, and bin this critical priority ARG is aided by the high relative abundance of *Shewanella* in the M5 sample (5.07%). We also observed a strong *Shewanella* signal in the Uk2 sample (relative abundance = 1.33%), from which another $bla_{\text{OXA-48}}$-like variant ($bla_{\text{OXA-547}}$) was detected by read-mapping, when compared to the remaining 13 samples in which no $bla_{\text{OXA-48}}$-like genes were detectable (mean *Shewanella* relative abundance = 0.11%).

Figure 5. Summary of medium- and high-quality MAGs – a) Genera; b) AMR determinants (annotated as '-like' if nucleotide identity is <100% when compared to reference); c) IS families.

Two *mcr-7*-like lipid A phosphoethanolamine transferase genes were detected in a high-quality *Aeromonas jandaei* MAG recovered from the Ug3 metagenome (*Physella* spp. faeces, Kimi Island, Uganda) on separate contigs (NODE_28 and NODE_32: 86.88% and 81.93% nucleotide identity to *mcr-7.1*, respectively). Lipid A phosphoethanolamine transferase enzymes cause a reduction in overall net-negative charge of the outer membrane, conferring resistance to colistin among some gram-negative bacteria, and are a major public health concern due to their propensity to be mobilised from progenitor species chromosomes to plasmids, limiting last-resort antibiotic treatment options for infections caused by multi-drug-resistant pathogens (35). *Aeromonas* spp. are the known progenitors of *mcr-3* and *mcr-7* (36). In our 15 samples, reads mapped to the MCR-3 Cluster90 in one and the MCR-7 Cluster90 in ten, underscoring the widespread prevalence of putative *Aeromonas*-associated colistin resistance genes in the sampled environment.

Discussion

Our findings provide novel insights into the role that freshwater snails play as AMR reservoirs, revealing a potential widespread environmental source of clinically relevant ARGs. Through analysis of snail faecal metagenomes from diverse geographic locations, we demonstrate that

their microbiomes harbour a wide array of bacterial species and ARGs, including those conferring resistance to both first- and last-line antibiotics. Identification of environmental progenitors of potentially mobile ARGs conferring resistance to carbapenems and colistin is particularly noteworthy in sub-Saharan Africa, where such antibiotics are not commonly available or prescribed for patients with severe infections caused by Enterobacterales, but where resistance among Enterobacterales to other treatment options such as penicillins, third-generation cephalosporins, and gentamicin has increased (37). A likely increase in carbapenem prescribing as a result of this may introduce additional selective pressures to aquatic habitats, driving selection and transfer of ARGs from environmental reservoirs to clinically relevant bacteria (38). Our results emphasize the potential for snails to contribute to this dissemination in freshwater ecosystems, particularly in those modified by anthropogenic change with deficits in local WASH (water, sanitation, and hygiene) infrastructures (3,18).

Proteobacteria dominated the snail faecal microbiomes, reinforcing results from previous studies that found Proteobacteria to be highly abundant in other freshwater snail microbiomes such as the gut and haemolymph (8,39–41). The significant correlations observed between Proteobacteria abundance and both alpha diversity and ARG load suggest that an abundance of Proteobacteria reduces overall microbiome diversity while increasing the relative abundance of ARGs within the microbial community. Specifically, we show that this effect is associated with Aeromonadales, within which we identified geographically widespread reservoirs of three ARG Cluster90s (CphA, MCR-7, OXA-12). The Proteobacteria order and *Aeromonas* genus have both recently been shown to be core taxa in gut samples extracted from laboratory-reared *Biomphalaria* snails (41); as well as signalling that snail faecal microbiomes are a reflection the intestinal tract, our results indicate that these core features are preserved across various snail genera and within their natural habitats. Meanwhile, our visualisation of hierarchical clusters at the order level demonstrates the variation in numerous other bacterial taxa observed between countries (and between districts in the case of Malawi). The differences observed between proximal samples from Malawian *Bulinus* snails at the species level highlight the potential role of local environmental conditions, such as water pollution, climate, and antibiotic exposure, in addition to host-specific characteristics, in shaping faecal microbiomes beyond the core taxa.

We recovered eight high-quality MAGs and 21 medium-quality MAGs from 20 bacterial genera in total, yet 4773 unique species from 1246 genera were detected in total by Kraken read profiling, demonstrating the limitations of assembly-based approaches in complex samples such as these. The inability to recover medium- to high-quality MAGs (with >50% completeness) in 1226 of the 1246 genera detected by read profiling (98.39%) suggests that some ARGs from low-abundance species are likely to have been undetectable even by our

read-based approach in which 60% template coverage was needed to call a hit, and hence

the ARG profiles that we present here are likely biased towards those found in high-abundance

genera such as *Aeromonas*. our The detection of ARGs from the CphA, MCR-7 and OXA-12

Cluster90s in 66.7%, 66.7% and 93.3% of samples, respectively, all of which originate from

Aeromonas species (42,43), is a reflection of this. The *mcr-7.1* colistin resistance gene was

first identified on plasmids in *K. pneumoniae* isolates from chickens in China (44), but

chromosome-borne variants of the genes have more recently been identified in *Aeromonas*

isolated from poultry in the United States (45), and in global environmental sources (water)

(46). The suggestion that *Aeromonas* species are the natural reservoirs of *mcr-3* and *mcr*-7

was since proven by robust evolutionary models (36), and we were able to assemble and bin

mcr-7-like variants to a putative *Aeromonas jandaei* MAG. Our results signpost to the potential

for emergence of new variants of these ARGs from freshwater gastropod faeces, and

specifically, the *Aeromonas* reservoirs within.

Similar reservoirs of ARGs in low-abundance species may be more widespread than our data

indicate. We demonstrate the presence of $bla_{OXA-181}$ and $bla_{OXA-547}$ in samples from Malawi and

the United Kingdom, respectively, both of which are bla_{OXA-48}-like genes. In the case of $bla_{OXA-181}$,

we could bin the >20 kbp contig containing the gene to a high-quality MAG of *S.*

xiamenensis, the species known to be the gene's progenitor (34). *Shewanella* species have

been identified as the progenitors of numerous bla_{OXA-48} variants (47), and the enzymes coded

for by these genes hydrolyze carbapenems and confer resistance to most beta-lactam, beta-

lactamase inhibitor combinations. They emerged and proliferated over the last two decades

to become the most prevalent carbapenemases amongst Enterobacterales in many parts of

Europe, Northern Africa, and the Middle East (28). Consequently, they are a major public

health concern, with carbapenemase-producing Enterobacterales (CPE) being listed by the

World Health Organisation as critical priority pathogens for which new antimicrobial agents

are urgently needed (48).

To our knowledge, this is the first report of the $bla_{OXA-181}$ variant in Malawi, a country where

CPE are uncommon in the clinical setting, but extended-spectrum beta-lactamase (ESBL)-

producing Enterobacterales resistant to a range of other beta-lactam antibiotics are, and

contribute to increased mortality in hospital patients with bloodstream infections (49,50).

ESBL-producing Enterobacterales have also been widely detected in water systems as well

as human and animal faeces in Malawi (50,51). Efforts to characterize the selective pressures

behind the spread of ESBL-producing Enterobacterales in Malawian wastewater and

freshwater are ongoing, but low environmental antimicrobial concentrations are known to drive

the transfer and persistence of ARGs in water systems (38). This selective pressure may also

facilitate the emergence of bla_{OXA-48}-like genes from environmental reservoirs where

environmental and human health intersect. Meanwhile, such reservoirs of carbapenem resistance genes could allow bacteria to react rapidly if carbapenems are introduced more routinely into clinical practice, limiting their efficacy. Events like this are often driven by horizontal gene transfer (HGT) associated with MGEs such as ISs (3), which we show are abundant within snail faeces. We were able to demonstrate in general via a statistical model that IS elements are associated with an increase in the number of different ARGs assembled in freshwater snail faecal metagenomes. An example of this is shown by the identification of a *sul1*-containing gene cluster upstream of IS*6100* in samples from Malawi and Uganda.

Our study does have several limitations. For example, the use of pooled snail faeces for metagenomic analysis, while necessary to obtain sufficient biomass for DNA extraction and sequencing, limits the ability to assess individual snail variability and the potential heterogeneity in microbiome composition and ARG content in snails from the same location. Meanwhile, the reliance on short-read sequencing, although effective for taxonomic profiling and ARG identification, poses challenges in assembling complete genes and genomes, particularly for low-abundance species, potentially leading to an underestimation of the full diversity of ARGs. This limitation is compounded by the inherent bias toward detecting ARGs in more abundant taxa which may skew the representation of ARGs in less abundant microbial communities. Inherently, short-read metagenome assemblies from diverse samples are highly fragmented. This limited our ability to uncover various examples of MGEs linked to ARGs on composite transposons, despite our ability to show a statistical association with regression modelling. Furthermore, while this study highlights the potential role of freshwater snails as AMR reservoirs, the cross-sectional nature of the sampling limits our ability to infer causality or the dynamics of ARG transmission over time. Longitudinal studies are needed to better understand temporal changes in ARG prevalence and the environmental factors influencing these patterns. Finally, while the study identifies ARGs of clinical relevance, it does not explore the functional impact of these ARGs, nor does it confirm their potential for HGT to human pathogens, a critical future step to quantify the risks associated with our findings.

Our study represents the first metagenomic analysis of AMR in freshwater gastropod faeces, providing important insights into the potential role of these invertebrates as reservoirs of clinically relevant ARGs. These findings have significant implications for AMR surveillance within One Health frameworks, particularly in areas where agricultural animals, wild animals and humans utilise the same open water sources. This reservoir of ARGs poses a threat to public health in an era of industrial-scale antibiotic use, particularly where access to clean water and sanitation is limited. Future research should build on these results through longitudinal sampling and by investigating how environmental factors, such as antibiotic contamination, influence ARG diversity and abundance in snail microbiomes. Additional

sampling in nearby human and animal populations to map potential transmission of ARGs from snail-associated bacteria to human pathogens will be essential for designing targeted interventions to curb AMR spread from and into shared aquatic environments.

Methods

Sampling collection, processing, and sequencing

In 2023 and 2024, freshwater snails were collected from 15 freshwater locations in four administrative regions: Malawi, n = 6; Uganda, n = 3, Zanzibar (Republic of Tanzania), n = 4; United Kingdom, n = 2. From each location, 50 morphologically identical snails of the same genus were identified by a competent medical malacologist (JRS), rinsed and then pooled in sterile water. The faecal material from each snail group was collected after 24 hours. DNA was extracted from PBS-washed faecal samples with the QIAamp PowerFecal Pro DNA Kit (QIAGEN, Germany), as per manufacturer's instructions. Concentration, purity and integrity of extracted DNA was assessed using a Qubit 4 fluorometer (Thermo Fisher Scientific, USA), NanoDrop One spectrophotometry (Thermo Fisher Scientific, USA), and the TapeStation System (Agilent Technologies, USA). Purified DNA extracts were shipped to Genewiz/AZENTA (UK) for library preparation, validation, and shotgun metagenomic sequencing. Sequencing was carried out with the Illumina NovaSeq instrument using a 2 x 150 bp paired-end configuration.

Bioinformatics

Read processing

Raw FASTQ sequence files were quality assessed using FASTQC (https://github.com/s-andrews/FastQC) Version 0.11.9. Trimming with Trimmomatic (52) Version 0.39 was performed to remove adaptor sequences and low-quality bases with a sliding window quality cutoff of Q20 and a minimum read length of 50 bp. Usual practice in metagenomic studies is to filter out host DNA before progressing to analysis of microbial reads by aligning all reads to host reference genomes. As our study involves analysis of pooled DNA samples from hosts whose genomes are not available in databases, we adapted the protocol outlined by Lu et al. (25) to remove contaminant and host DNA. Trimmed reads were classified using Kraken 2 (53) Version 2.1.3. Kraken 2 works by matching read k-mers against a database of k-mers from existing genomes. We aligned reads to the standard Kraken 2 database, which includes RefSeq archaea, bacteria, viruses, plasmid complete genomes, UniVec Core and the human reference genome, GRCh38. In a two-step approach using the KrakenTools "extract_kraken_reads.py" script, contaminant reads identified as eukaryotic were first filtered out, before reads identified as bacterial in origin were extracted. The larger read set obtained

after the first filtering step was used for read-based ARG profiling, as including only the reads

identified as bacterial by Kraken 2 may potentially have removed some ARG variants. The

twice filtered bacterial-only reads were used for taxonomic profiling and assembly.

<u>Taxonomic profiling</u>

Kraken 2 reports containing bacterial read taxonomic assignments were used to run species

abundance estimation with Bracken (54), with a minimum number of reads required for a

classification at the specified rank set at 10 to reduce noise from potential low-abundance

contaminant species, as per default recommendations. A Bray-Curtis dissimilarity matrix

containing pairwise dissimilarities among microbiome samples was generated by running the

KrakenTools "beta_diversity.py" script with Bracken outputs, to assess bacterial beta diversity.

<u>ARG database and clustering</u>

Sequences in the ResFinder database (27) Version 2.3.2 were downloaded, then clustered

using CD-HIT (55) Version 4.8.1 to identify ARG clusters containing sequences with a

minimum nucleotide identity of 90%, referred to as Cluster90s.

<u>Read-based ARG detection</u>

To identify ARGs within each sample, reads were aligned to reference sequences in the

ResFinder database using KMA (26) Version 1.4.14, a tool designed to map raw reads directly

against redundant databases such as ARG databases. Alignments were then filtered to retain

only those with a minimum 90% query identity and 60% template coverage. For all reference

sequences with alignment hits after filtering, the corresponding Cluster90 was identified and

annotated manually.

<u>Metagenome assembly, binning, and classification</u>

Reads identified as being bacterial by Kraken 2 were assembled with metaSPAdes (56)

Version 3.11.1. QUAST (57) Version 5.0.2 was used to assess the quality of metagenome

assemblies. Contigs were indexed using the Burrows Wheeler Aligner (BWA) Version 0.7.17

before bacterial reads were aligned to the indexed contigs with "bwa-mem" (58). The contigs

and resulting sorted BAM files were parsed to the "jgi_summarize_bam_contig_depth" script

from MetaBAT2 (59) Version 2.17, before the resulting depth files were used by MetaBAT2 to

bin assembled contigs of ≥ 2,500 bp to putative genomes (MAGs). CheckM (60) Version 1.1.2

was used to assess the completeness and contamination of each MAG. CheckM outputs were

compared to standards set by the genome standards consortium to classify MAGs as high-

quality (>90% completeness, <5% contamination), medium-quality MAGs (>50%

completeness, <10% contamination) or low-quality (<50% completeness and/or >10%

contamination) (61). MAGs deemed as medium- or high-quality proceeded to further analysis.
MAGs underwent taxonomic classification using GTDB-Tk (62) Version 2.1.1 against the
Genome Taxonomy Database Version 09-RS220 (24th April 2024).

<u>Assembly-based profiling of ARGs and MGEs</u>

Metagenome assemblies were screened against the ResFinder (27) and ISfinder (30)
databases using ABRicate (https://github.com/tseemann/abricate) Version 1.0.1 at default
settings (minimum 80% identity and 80% query cover) to identify contigs containing ARGs and
ISs. Compared to our read-mapping approach, in which we used a higher identity threshold to
identify hits (90%) in combination with a lower minimum query cover (60%) due to the more
fragmented nature of using raw sequences, the assembly-based approach allowed for a
broader detection of putative AMR determinants and MGEs, which could then be
characterised further within their genomic context following annotation. Annotations were
performed using Bakta (63) Version 1.9.3 (full database version 5.1). To elucidate the potential
origins and dissemination pathways of ARGs, selected contigs were subjected to a sequence
similarity search using the Nucleotide Basic Local Alignment Search Tool (BLAST) webtool
(megablast algorithm) against the GenBank Nucleotide database. This step aimed to identify
existing homologous sequences and/or gene clusters, thereby pinpointing potential source
organisms. The Clinker webtool (https://cagecat.bioinformatics.nl/tools/clinker) (64) was used
to visualise gene cluster comparisons.

Statistical analysis and visualisation

Species abundance and read-based ARG assay data were combined with
hierarchical/classification data and sample metadata to generate a
TreeSummarizedExperiment data container (65). Data associated with MAGs were handled
separately. Within the R environment, downstream data wrangling, analysis, and visualisation
took place predominantly using the R/Bioconductor *mia* and *miaViz* (Version 1.12.0)
packages, and the R/CRAN *ggplot2* (Version 3.5.1) and *pheatmap* (Version 1.0.12) packages.
CLR transformation was applied to microbial composition data (66) to normalize for
compositional effects, with a pseudocount added to manage zero values using the *mia*
package. Hierarchical clustering of both taxa and samples was applied to standardised CLR-
transformed data (CLR-z) using Euclidean distance and complete linkage, as implemented by
default in the *pheatmap* package. Pearson correlation was used to explore the relationships
between numerical variables and ARG load, assuming linearity in the data. When exploratory
analysis revealed non-linear trends in the data, an appropriate transformation (e.g. log
transformation) was applied. Generalized linear regression with a negative binomial

distribution was carried out with the R/CRAN *MASS* package (Version 7.3.60.2) to assess the
association between IS element counts and ARG counts in metagenome assemblies.

<u>SEQUENCE AND DATA AVAILABILITY</u>

Short reads from all samples sequenced in this study have been deposited in the Sequence
Read Archive (SRA) under BioProject PRJNA1211045. Data and code used to carry out
analyses in R are available at https://github.com/amoreo71/freshwater_snail_faecal.

<u>FUNDING</u>

AMO'F is funded by the Medical Research Council via the Liverpool School of Tropical
Medicine / Lancaster University doctoral training partnership (grant no.
MR/W007037/1). Fieldwork was supported by the Wellcome Trust within a Joint Investigator
Award (grant no. 220818/Z/20/Z) and by MÁEÖ (grant no. 2023-2024/183544). APR
acknowledges funding from the Medical Research Council, Biotechnology and Biological
Sciences Research Council and Natural Environmental Research Council which are all
Councils of UK Research and Innovation (grant no. MR/W030578/1) under the umbrella of the
JPIAMR (Joint Programming Initiative on Antimicrobial Resistance), and UKRI through the
Strength in Places Fund (grant no. SIPF 36348). The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.

<u>AUTHOR CONTRIBUTIONS</u>

The study was conceived and designed by AMO'F, JM, JRS and APR. Sampling was carried
out by AMO'F, AJ, SJ, PM, GM, SA, DO, SAK, JEL and JRS. Formal analysis was performed
by AMO'F, AF, JRS and APR. The original draft was prepared by AMO'F, JRS and APR, then
reviewed and edited by all authors. All authors approved the final version of the manuscript.

<u>ACKNOWLEDGEMENTS</u>

The authors thank Dr Simon Wagstaff and Mr Andrew Bennet for provision and maintenance
of the scientific computing environment at LSTM used to carry out bioinformatic analysis. The
authors acknowledge staff and local people in each of the sampling areas who aided the
sampling teams.

COMPETING INTERESTS

None to declare.

REFERENCES

1.   Hutchings MI, Truman AW, Wilkinson B. Antibiotics: past, present and future. Curr Opin Microbiol. 2019 Oct 1;51:72–80.

2.   Tenover FC. Mechanisms of antimicrobial resistance in bacteria. Am J Med. 2006 Jun;119(6 Suppl 1):S3-10; discussion S62-70.

3.   Despotovic M, de Nies L, Busi SB, Wilmes P. Reservoirs of antimicrobial resistance in the context of One Health. Curr Opin Microbiol. 2023 Jun;73:102291.

4.   Rockström J, Steffen W, Noone K, Persson Å, Chapin FS, Lambin EF, et al. A safe operating space for humanity. Nature. 2009 Sep;461(7263):472–5.

5.   Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. Lancet. 2014 Jun 28;383(9936):2253–64.

6.   Caravedo MA, Cabada MM. Human Fascioliasis: Current epidemiological status and strategies for diagnosis, treatment, and control. Res Rep Trop Med. 2020 Nov 26;11:149–58.

7.   Liu H, Macdonald CA, Cook J, Anderson IC, Singh BK. An ecological loop: host microbiomes across multitrophic interactions. Trends Ecol Evol. 2019 Dec;34(12):1118–30.

8.   Chevalier FD, Diaz R, McDew-White M, Anderson TJC, Le Clec'h W. The hemolymph of *Biomphalaria* snail vectors of schistosomiasis supports a diverse microbiome. Environ Microbiol. 2020 Dec;22(12):5450–66.

9.   Schols R, Vanoverberghe I, Huyse T, Decaestecker E. Host-bacteriome transplants of the schistosome snail host *Biomphalaria glabrata* reflect species-specific associations. FEMS Microbiol Ecol. 2023 Sep 1;99(9):fiad101.

10.   Madsen H, Frandsen F. The spread of freshwater snails including those of medical and veterinary importance. Acta Trop. 1989 May;46(3):139–46.

11.   Lv SB, He TT, Hu F, Li YF, Yuan M, Xie JZ, et al. The impact of flooding on snail spread: the case of endemic schistosomiasis areas in Jiangxi Province, China. Trop Med Infect Dis. 2023 Apr 30;8(5):259.

12.   Alharbi MH, Condemine C, Christiansen R, LaCourse EJ, Makaula P, Stanton MC, et
al. *Biomphalaria pfeifferi* snails and intestinal schistosomiasis, Lake Malawi, Africa, 2017–
2018. Emerg Infect Dis. 2019 Mar;25(3):613–5.

13.   Pedersen UB, Stendel M, Midzi N, Mduluza T, Soko W, Stensgaard AS, et al.
Modelling climate change impact on the spatial distribution of fresh water snails hosting
trematodes in Zimbabwe. Parasit Vectors. 2014 Dec;7(1):1–12.

14.   Kock KD, Wolmarans CT, Bornman M. Distribution and habitats of *Biomphalaria*
pfeifferi, snail intermediate host of Schistosoma mansoni, in South Africa. Water SA. 2004
Aug 3;30(1):29–36.

15.   O'Ferrall AM, Musaya J, Stothard JR, Roberts AP. Aligning antimicrobial resistance
surveillance with schistosomiasis research: an interlinked One Health approach. Trans R
Soc Trop Med Hyg. 2024 Aug 1;118(8):498–504.

16.   Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Aguilar GR, Gray A, et al. Global
burden of bacterial antimicrobial resistance in 2019: a systematic analysis. Lancet. 2022 Feb
12;399(10325):629–55.

17.   Olaru ID, Walther B, Schaumburg F. Zoonotic sources and the spread of
antimicrobial resistance from the perspective of low and middle-income countries. Infect Dis
Poverty. 2023 Jun 14;12(1):59.

18.   Musicha P, Morse T, Cocker D, Mugisha L, Jewell CP, Feasey NA. Time to define
One Health approaches to tackling antimicrobial resistance. Nat Commun. 2024 Oct
10;15(1):8782.

19.   Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. Shotgun metagenomics,
from sampling to analysis. Nat Biotechnol. 2017 Sep;35(9):833–44.

20.   Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, et al. A new
genomic blueprint of the human gut microbiota. Nature. 2019 Apr;568(7753):499–504.

21.   Kim CY, Lee M, Yang S, Kim K, Yong D, Kim HR, et al. Human reference gut
microbiome catalog including newly assembled genomes from under-represented Asian
metagenomes. Genome Med. 2021 Aug 27;13(1):134.

22.   Howe A, Chain PSG. Challenges and opportunities in understanding microbial
communities with metagenome assembly (accompanied by IPython Notebook tutorial). Front
Microbiol. 2015 Jul 9;6:678.

23.   Lee K, Raguideau S, Sirén K, Asnicar F, Cumbo F, Hildebrand F, et al. Population-
level impacts of antibiotic usage on the human gut microbiome. Nat Commun. 2023 Mar
2;14(1):1191.

24.   Carruthers LV, Nordmeyer SC, Anderson TJc, Chevalier FD, Le Clec'h W. Organ-
specific microbiomes of *Biomphalaria snails* [Preprint]. 2024. Available from:
http://biorxiv.org/lookup/doi/10.1101/2024.06.11.598555

25.   Lu J, Rincon N, Wood DE, Breitwieser FP, Pockrandt C, Langmead B, et al.
Metagenome analysis using the Kraken software suite. Nat Protoc. 2022 Dec;17(12):2815–
39.

26.   Clausen PTLC, Aarestrup FM, Lund O. Rapid and precise alignment of raw reads
against redundant databases with KMA. BMC Bioinformatics. 2018 Aug 29;19(1):307.

27.   Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder
4.0 for predictions of phenotypes from genotypes. J Antimicrob Chemother. 2020 Dec
1;75(12):3491–500.

28.   Boyd SE, Holmes A, Peck R, Livermore DM, Hope W. OXA-48-Like β-Lactamases:
global epidemiology, treatment options, and development pipeline. Antimicrob Agents
Chemother. 2022 Jul 20;66(8):e00216-22.

29.   Figueiredo S, Bonnin RA, Poirel L, Duranteau J, Nordmann P. Identification of the
naturally occurring genes encoding carbapenem-hydrolysing oxacillinases from
Acinetobacter haemolyticus, *Acinetobacter johnsonii*, and *Acinetobacter calcoaceticus*. Clin
Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis. 2012 Sep;18(9):907–13.

30.   Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference
centre for bacterial insertion sequences. Nucleic Acids Res. 2006 Jan 1;34(suppl_1):D32–6.

31.   Nielsen TK, Browne PD, Hansen LH. Antibiotic resistance genes are differentially
mobilized according to resistance mechanism. GigaScience. 2022 Jan 1;11:giac072.

32.   Dao TD, Takemura T, Kasuga I, Hirabayashi A, Nga NT, Anh PHQ, et al. Emergence
of plasmid-mediated RND-type efflux pump gene cluster *tmexCD-toprJ* in *Shewanella*
xiamenensis in a water environment [Preprint]. 2022. Available from:
http://biorxiv.org/lookup/doi/10.1101/2022.09.15.508154

33.   Potron A, Nordmann P, Lafeuille E, Maskari Z, Rashdi F, Poirel L. Characterization of
OXA-181, a carbapenem-hydrolyzing class D-lactamase from *Klebsiella pneumoniae*.
Antimicrob Agents Chemother. 2011 Sep 15;55:4896–9.

34.   Potron A, Poirel L, Nordmann P. Origin of OXA-181, an emerging carbapenem-
hydrolyzing oxacillinase, as a chromosomal gene in *Shewanella xiamenensis*. Antimicrob
Agents Chemother. 2011 Sep;55(9):4405–7.

35.   Samantha A, Vrielink A. Lipid A phosphoethanolamine transferase: regulation,
structure and immune response. J Mol Biol. 2020 Aug 21;432(18):5184–96.

36.   Guo Y, Zou G, Kerdsin A, Schultsz C, Hu C, Bei W, et al. Characterization of NMCR-
3, NMCR-4 and NMCR-5, three novel non-mobile colistin resistance determinants:
Implications for MCR-3, MCR-7, and MCR-5 progenitors, respectively. Drug Resist Updat.
2024 Jul 1;75:101088.

37.   Kowalski M, Obama BM, Catho G, Dewez JE, Merglen A, Ruef M, et al. Antimicrobial
resistance in Enterobacterales infections among children in sub-Saharan Africa: a systematic
review and meta-analysis. eClinicalMedicine. 2024 Apr; 70:102512.

38.   Murray AK, Stanton IC, Tipper HJ, Wilkinson H, Schmidt W, Hart A, et al. A critical
meta-analysis of predicted no effect concentrations for antimicrobial resistance selection in
the environment. Water Res. 2024 Nov 15;266:122310.

39.   Silva TM, Melo ES, Lopes ACS, Veras DL, Duarte CR, Alves LC, et al.
Characterization of the bacterial microbiota of *Biomphalaria glabrata* (Say, 1818) (Mollusca:
Gastropoda) from Brazil. Lett Appl Microbiol. 2013 Jul;57(1):19–25.

40.   Hu Z, Tong Q, Chang J, Yu J, Li S, Niu H, et al. Gut bacterial communities in the
freshwater snail *Planorbella trivolvis* and their modification by a non-herbivorous diet. PeerJ.
2021 Feb 12;9:e10716.

41.   Li P, Hong J, Wu M, Yuan Z, Li D, Wu Z, et al. Metagenomic analysis reveals
variations in gut microbiomes of the *Schistosoma mansoni*-transmitting snails *Biomphalaria
straminea* and *Biomphalaria glabrata*. Microorganisms. 2023 Sep 28;11(10):2419.

42.   Chen PL, Ko WC, Wu CJ. Complexity of β-lactamases among clinical *Aeromonas*
isolates and its clinical implications. J Microbiol Immunol Infect. 2012 Dec 1;45(6):398–403.

43.   Alksne LE, Rasmussen BA. Expression of the AsbA1, OXA-12, and AsbM1 beta-
lactamases in *Aeromonas jandaei* AER 14 is coordinated by a two-component regulon. J
Bacteriol. 1997 Mar;179(6):2006–13.

44.   Yang YQ, Li YX, Lei CW, Zhang AY, Wang HN. Novel plasmid-mediated colistin
resistance gene mcr-7.1 in *Klebsiella pneumoniae*. J Antimicrob Chemother. 2018 Jul
1;73(7):1791–5.

45.   Wang Y, Hou N, Rasooly R, Gu Y, He X. Prevalence and genetic analysis of

chromosomal mcr-3/7 in *Aeromonas* from U.S. animal-derived samples. Front Microbiol.

2021 Apr 30;12:667406.

46.   Khedher MB, Baron SA, Riziki T, Ruimy R, Raoult D, Diene SM, et al. Massive

analysis of 64,628 bacterial genomes to decipher water reservoir and origin of mobile colistin

resistance genes: is there another role for these enzymes? Sci Rep. 2020 Apr 6;10(1):5970.

47.   Tacão M, Araújo S, Vendas M, Alves A, Henriques I. *Shewanella* species as the origin

of *bla*OXA-48 genes: insights into gene diversity, associated phenotypes and possible

transfer mechanisms. Int J Antimicrob Agents. 2018 Mar 1;51(3):340–8.

48.   World Health Organisation. WHO publishes list of bacteria for which new antibiotics

are urgently needed [Internet]. 2017 [cited 2023 Jun 15]. Available from:

https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-

antibiotics-are-urgently-needed

49.   Lewis JM, Mphasa M, Banda R, Beale MA, Heinz E, Mallewa J, et al. Colonization

dynamics of extended-spectrum beta-lactamase-producing Enterobacterales in the gut of

Malawian adults. Nat Microbiol. 2022 Oct;7(10):1593–604.

50.   Lester R, Musicha P, Kawaza K, Langton J, Mango J, Mangochi H, et al. Effect of

resistance to third-generation cephalosporins on morbidity and mortality from bloodstream

infections in Blantyre, Malawi: a prospective cohort study. Lancet Microbe. 2022 Dec

1;3(12):e922–30.

51.   Cocker D, Chidziwisano K, Mphasa M, Mwapasa T, Lewis JM, Rowlingson B, et al.

Investigating One Health risks for human colonisation with extended spectrum β-lactamase-

producing *Escherichia coli* and *Klebsiella pneumoniae* in Malawian households: a

longitudinal cohort study. Lancet Microbe. 2023 Jul 1;4(7):e534–43.

52.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014 Aug 1;30(15):2114–20.

53.   Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2.

Genome Biol. 2019 Nov 28;20(1):257.

54.   Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: estimating species abundance

in metagenomics data. PeerJ Comput Sci. 2017 Jan 2;3:e104.

55.   Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-

generation sequencing data. Bioinforma Oxf Engl. 2012 Dec 1;28(23):3150–2.

56.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile

metagenomic assembler. Genome Res. 2017 May;27(5):824–34.

57.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for

genome assemblies. Bioinforma Oxf Engl. 2013 Apr 15;29(8):1072–5.

58.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM [PREPRINT]. 2013. Available from: https://arxiv.org/abs/1303.3997

59.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive

binning algorithm for robust and efficient genome reconstruction from metagenome

assemblies. PeerJ. 2019 Jul 26;7:e7359.

60.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing

the quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res. 2015 Jul;25(7):1043.

61.   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK,

et al. Minimum information about a single amplified genome (MISAG) and a metagenome-

assembled genome (MIMAG) of bacteria and archaea. Nat Biotechnol. 2017 Aug

8;35(8):725–31.

62.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics. 2020 Mar 15;36(6):1925–7.

63.   Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. Bakta:

rapid and standardized annotation of bacterial genomes via alignment-free sequence

identification. Microb Genomics. 2021;7(11):000685.

64.   Gilchrist CLM, Chooi YH. clinker & clustermap.js: automatic generation of gene

cluster comparison figures. Bioinforma Oxf Engl. 2021 Aug 25;37(16):2473–5.

65.   Huang R, Soneson C, Ernst FGM, Rue-Albrecht KC, Yu G, Hicks SC, et al.

TreeSummarizedExperiment: a S4 class for data with hierarchical structure.

F1000Research. 2021 Mar 2;9:1246.

66.   Calle ML. Statistical analysis of metagenomics data. Genomics Inform. 2019 Mar

31;17(1):e6.