Yes-associated Protein Induces Age-dependent Inflammatory Signaling in the Pulmonary

Endothelium

Memet T. Emin,[1] Alexandra M. Dubuisson,[1,2] Prisha Sujin Kumar,[1] Carsten Knutsen,[3] Cristina M. Alvira,[3]

and Rebecca F. Hough*,[1]

[1]Pediatric Critical Care, Hospitalist, and Palliative Medicine, Department of Pediatrics, Columbia

University Irving Medical Center, [2] Louisiana State University Health Sciences Center Shreveport,

[3]Department of Pediatrics, University of California, San Francisco School of Medicine

*Address correspondence to:

Rebecca F. Hough

Department of Pediatrics

3959 Broadway, CHN10-24

New York, New York 10032, USA.

Phone: 212.305.2370

Email: rft2106@cumc.columbia.edu

ORCiD: 0000-0002-0203-4146

Conflict of interest statement: The authors have declared that no conflict of interest exists.

Running Title: Age-dependent YAP signaling in the lung endothelium

Abstract

Acute Lung Injury (ALI) causes the highly lethal Acute Respiratory Distress Syndrome (ARDS) in

children and adults, for which therapy is lacking. Children with Pediatric ARDS (PARDS) have a

mortality rate that is about half of adults with ARDS. Improved ALI measures can be reproduced in

rodent models with juvenile animals, suggesting that physiologic differences may underlie these different

outcomes. Here, we show that pneumonia-induced ALI caused inflammatory signaling in the endothelium

of adult mice which depended on Yes-associated protein (YAP). This signaling was not present in 21-day-

old weanling mice. Transcriptomic analysis of lung endothelial responses revealed nuclear factor kappa-B

(NF-κB) as significantly increased with ALI in adult versus weanling mice. Blockade of YAP signaling

protected against ALI and NF-κB nuclear translocation in adult mice. Our results demonstrate an

important signaling cascade in the lung endothelium of adult mice that is not present in weanlings. We

suggest other pathways may also exhibit age-dependent inflammatory signaling, which would have

important implications for therapeutics in the adult and pediatric age groups.

Introduction

Acute lung injury (ALI) causes the Acute Respiratory Distress Syndrome (ARDS) and Pediatric ARDS (PARDS). Mortality rates in adult patients with ARDS are double those of PARDS patients (1–3). Preclinical studies in rodents show decreased lung injury measures in juvenile mice (4–6), suggesting that age-specific physiological differences underlie the different outcomes in ARDS and PARDS.

The Hippo pathway and its main effectors Yes-associated Protein (YAP) and PDZ-binding motif (TAZ) are vital for growth and differentiation in lung epithelium and in non-pulmonary organs (7–9). YAP is essential for the development of the endothelial barrier in the retina and the brain (10) and interacts with tight junction components in the lung epithelium (11). Recently, YAP was shown to protect against lung injury (12, 13). We hypothesized that YAP responses could underlie the differences in lung injury responses in juvenile and adult rodents.

Here, we instilled *Pseudomonas aeruginosa* into adult and weanling mice, and showed that lung injury measures are improved in weanlings. We show robust YAP protein expression in response to *P. aeruginosa* infection in adult mice, but not in weanlings. YAP knockdown prevented lung injury in adult mice, suggesting a causal role of YAP in our injury model. Knockdown of YAP decreased nuclear translocation of NF-κB in adult mice, suggesting that YAP contributes to pro-inflammatory signaling in the pulmonary endothelium of adult, but not juvenile mice.

Methods

Materials. For immunoblot, YAP mouse antibody (Cat#12395, 1:1000), YAP rabbit antibody (cat#14074, 1:1000), GAPDH mouse antibody (Cat#97166, 1:1000), and NF-κB p65 rabbit antibody (Cat#8242, 1:1000) were purchased from Cell Signaling Technology; HDAC1 mouse monoclonal antibody (Cat# 66085, 1:1000) was purchased from Proteintech; and actin rabbit antibody (cat# A2066, 1:1000) was purchased from Sigma-Aldrich. For lung cell isolation, CD31 rat monoclonal antibody clone MEC 13.3 (1:500, catalog 557355) was purchased from BD Biosciences. CD45 rat monoclonal antibody (1:600, catalog 14-0451-85), CD41 rat monoclonal antibody (1:500, catalog 14-0411-85), and CD326 rat

monoclonal antibody (1:500, catalog 14-5791-85) were purchased from Invitrogen. siRNA constructs

targeting Yap1 (siRNA ID #s76160) and Taz (siRNA ID #s211845) were purchased from ThermoFisher

Scientific. Male and female C57BL/6J mice were purchased from Jackson Laboratory

(RRID:IMSR_JAX:000664). *P. aeruginosa* strain K was obtained from Alice Prince (Columbia

University).

Pseudomonas-induced lung injury. P. aeruginosa was prepared as previously described (14). We

anesthetized mice and intranasally (i.n.) instilled 2.5 x 10^5 colony forming units (CFUs) *P. aeruginosa* at

exponential growth phase in 50µl sterile PBS (adult) or 20µl (weanling). 24 hours later, we anesthetized

the mice, performed a tracheotomy, and collected BAL as described (14) for cell count, protein

quantification, cytokine measurement, and determination of CFUs. We retained lungs after BAL for

downstream experiments.

SpO_2 measurement. For non-invasive O_2 saturation (SpO_2) measurements, we anesthetized mice and

removed the fur from right thigh. We placed a MouseOx® Pulse Oximeter sensor (Starr Life Sciences

Corp, Oakmont, PA) on the right femoral artery region and recorded SpO_2 after a 15-min stabilization

period.

YAP RT-PCR. We isolated RNA from whole lung using miRNeasy kit (Qiagen). We performed YAP RT-

PCR by using TaqMan Gene Expression Master Mix with QuantStudio™ 5 Real-Time PCR Instrument

(Thermo Fisher Scientific Applied Biosystems).

YAP knockdown. Anesthetized mice were in vivo transfected with YAP and TAZ siRNA (50µg in 150µL)

as in (14, 15) by tail vein injection 24 hours before *P. aeruginosa* instillation.

Immunoblot. Without freezing, we enriched lungs for cytoplasmic and nuclear fractions using NE-PER

Nuclear and Cytoplasmic Extraction Reagents (Thermo Fisher Scientific, cat #78833). Equal amounts of

protein were separated by SDS-PAGE, transferred onto nitrocellulose membrane, blocked in

StartingBlock Blocking Buffer (Thermo Scientific / Pierce, catalog 37543), and immunoblotting was

performed.

Endothelial cell isolation. Endothelial cells from C57BL/6J mice were isolated as described (16, 17) with

the following modifications. Lungs were digested for 15 min at 37°C in 0.38mg/ml Liberase (Roche) in

D-PBS + 0.1% albumin. We used a DynaMag-2 Magnet and Dynabeads Sheep Anti-Rat IgG (Invitrogen)

to deplete CD45+ and CD41+ cells, then positively enrich three times for CD31+ cells. These procedures

yielded ~10^5 cells per mouse lung. Bead isolation by this method yielded endothelial cells that were ~93%

pure and had <5% leukocyte contamination.

Bulk RNA-seq. RNA was isolated from freshly isolated endothelial cells bound to magnetic beads

(miRNeasy). RNA-seq was performed by Azenta. We aligned sequences to the GRCm39 mouse genome

using STAR aligner. We analyzed differential gene expression (DE) with EdgeR using a generalized linear

model with thresholds for significance set at <0.05 FDR (Benjamani-Hochenberg) and 1>|Log2(FC)|.

Results

Differential lung injury responses in weanling and adult mice.

To determine lung injury responses, we intranasally instilled weanling (P21-24) and adult (8-10 week old)

mice with *P. aeruginosa*. Twenty-four hours later, we assessed mice for lung injury by measurement of

bronchoalveolar lavage (BAL) protein, BAL cell count, BAL neutrophils, extra-vascular lung water, BAL

cytokines, and non-invasive measurement of O_2 saturation (SpO$_2$). Following *P. aeruginosa* instillation,

adult mice had an increase in BAL protein, BAL cell count, BAL neutrophils, EVLW, BAL Il-6 and BAL

TNF-α (**Fig 1A-E**), and a decrease in SpO$_2$ (**Fig 1F**) compared to weanling mice. Saline treatment did not

produce different responses in any lung injury measurements.

Given the difference in size of the mice (~10g vs. ~25g), we considered whether delivery of *P. aeruginosa*

could differ between the two age groups. In addition, decreased clearance of *P. aeruginosa* has been noted

in mice >20 days old (18). To assess delivery, we instilled the same number of CFUs in the same volume

to both age groups, and again in two different volumes. The lung injury results did not differ when the

volume was changed. To assess bacterial clearance, we measured BAL CFUs 24h after instillation and

noted no difference between the two age groups (**Supplemental Fig 1**). These findings suggest that a

similar number of CFUs was delivered in both ages and that bacterial clearance did not differ significantly

between the age groups.

P. aeruginosa-induced changes in YAP protein expression differ with age

We determined changes in expression of YAP in response to *P. aeruginosa*-induced lung injury. Following

BAL and a vascular wash, we enriched whole lungs from *P. aeruginosa* and saline treated mice into

nuclear and cytoplasmic portions. We found that adult mice had robust increases in YAP protein

expression in response to *P. aeruginosa* instillation in both cytoplasmic and nuclear fractions (**Fig 2A, B**).

In contrast, weanling mice had decreased cytoplasm expression and no change in nuclear expression of

YAP. Of note, there was no difference in YAP protein expression between age groups in saline-treated

controls (**Supplemental Fig 2**), suggesting that baseline YAP expression does not differ between the two

age groups. We assessed YAP mRNA expression with RT-PCR (**Supplemental Fig 3**) and noted that YAP

mRNA expression in both weanling and adult mice decreased upon *P. aeruginosa*-induced injury,

suggesting a post-transcriptional mechanism of increased YAP protein expression in adult mice.

P. aeruginosa-induced YAP protein increases occur in lung endothelial cells

Given the substantial increase in inflammatory cells in the BAL of adult versus weanling mice (**Fig 1B-**

C), we considered whether leukocyte expression of YAP (19) contributed to YAP increases in adult mice.

To determine the cell type in which YAP expression increased, we enriched for three fractions: lung

endothelial cells, epithelial cells, and leukocytes plus platelets. We performed immunoblot for YAP in

each of the enriched cell populations. We noted a substantial increase in YAP protein expression in

endothelial cells (**Fig 2C**). There was no significant increase in YAP expression in epithelial cells (**Fig**

2D), and no YAP protein expression was detected in the leukocyte- and platelet-enriched fraction in PBS

or *P. aeruginosa*-treated mice.

Vascular YAP mediates P. aeruginosa-induced lung injury in adult mice

Since YAP expression increases substantially in adult mice, but not in weanling mice, we hypothesized

that adult mice exhibit YAP-dependent lung injury responses that weanling mice lack. Thus, we targeted

endothelial YAP by tail vein injection of siRNA-targeted YAP and its paralog TAZ. Injection of YAP- and

TAZ-targeted siRNA decreased YAP expression in *P. aeruginosa* infected mice by ~80% (**Fig 3A**). We

assessed lung injury measures as in **Fig 1** and found that YAP knockdown prevented *P. aeruginosa*-

induced increases in BAL cell count, IL-6, and TNFα, as well as *P. aeruginosa*-induced decreases in SpO$_2$

(**Fig 3B-D**). This suggests that endothelial YAP expression is important for inflammatory responses in *P.*

aeruginosa-induced lung injury. Notably, YAP knockdown had no effect on BAL protein and EVLW,

suggesting that endothelial YAP may not be essential for barrier failure.

YAP expression drives endothelial NF-κB responses in P. aeruginosa-infected adult mice

To determine differences in *P. aeruginosa*-induced lung injury endothelial responses between weanling

and adult mice, we performed bulk RNA sequencing on freshly isolated endothelial cells from mice

treated with i.n. *P. aeruginosa* and control mice treated with i.n. PBS. In weanling mice, 2256 genes

showed increased expression in response to *P. aeruginosa*-induced injury and 2651 showed decreased

expression. In adult mice, 2732 genes increased expression and 3476 decreased. Of note, gene expression

differences between both age groups did not differ greatly at baseline (**Fig 4A**). However, transcriptomic

responses to lung injury differ vastly between age groups. This suggests substantial differences in lung

endothelial responses to inflammatory injury of weanling and adult mice.

We noted that there was no significant alteration in *yap1* gene expression with *P. aeruginosa* instillation in

either weanling or adult endothelia. We identified pathways that exhibited increased expression in adult

mice in response to *P. aeruginosa*-induced lung injury relative to weanling mice. The most significant

increases were found in TNFα signaling via NF-κB (**Fig 4B**). Although both weanling and adult

endothelial cells increased their expression of genes involved in this pathway, the adult endothelial

responses were increased relative to responses in weanling mice.

We considered whether YAP drives NF-κB nuclear translocation, leading to a more robust inflammatory

response in adult mice. First, we determined whether *P. aeruginosa* instillation drove NF-κB translocation

to the nucleus in adult mice. 24h after *P. aeruginosa* instillation, we enriched whole lung into nuclear and

cytoplasmic fractions. We noted a significant increase in NF-κB in the nuclear fraction and a decrease in

the cytoplasmic fraction (**Fig 4C**). In weanlings, there was an increase in NF-κB in the nuclear fraction,

but no significant difference in NF-κB quantity in the cytoplasmic fraction upon instillation of *P.*

aeruginosa (**Supplemental Fig 4**), suggesting that NF-κB nuclear translocation in response to *P.*

aeruginosa in weanlings may be decreased in comparison to adult mice. To determine whether the

increase in YAP expression in adult mice drove NF-κB translocation, we knocked down YAP expression

by vascular injection of siRNA as in **Fig 3A** and instilled *P. aeruginosa* into controls and siRNA-treated

mice 24h later. 24 hours after *P. aeruginosa* instillation, we measured NF-κB in the nuclear and

cytoplasmic fractions. We noted that YAP knockdown increased NF-κB protein expression in the

cytoplasm and decreased YAP expression in the nucleus (**Fig 4D**), suggesting that YAP mediated NF-κB

nuclear translocation.

Discussion

Our results differ from previous reports that endothelial YAP protects against lung injury (12, 13). YAP

appeared to inhibit NF-κB nuclear translocation by promoting degradation of tumor necrosis factor

receptor-associated factor 6 (TRAF-6) (13). In the epithelium, nuclear YAP may increase *ikba* gene

expression, inhibiting NF-κB nuclear translocation (20). Of note, in our transcriptomics dataset we noted

that expression of the *ikba* mouse homologue *nfkbia* was induced in both weanling and adult mice. This

suggests YAP-independent drivers of transcription that are active in weanling mice. In the above studies,

cell-specific YAP knockouts were used. It is possible that knockout of YAP leads to up-regulation of

alternative pathways (21). We suggest that our use of knockdown of YAP may more accurately reflect the

results of potential pharmacologic inhibition. The role of YAP in lung injury and repair is complex and

requires further study.

In neonatal mice in the early alveolar stage of lung development, NF-κB is known to play a protective

role—ultimately promoting angiogenesis (17, 22–24). To our knowledge, this role has not been described

in weanling mice in the late alveolar stage of lung development. However, it is possible that NF-κB

nuclear translocation in weanling mice results in signaling that differs from canonical pro-inflammatory

signaling in adults.

Our study has limitations. First, the signaling that leads to differential YAP expression between the

weanling and adult age groups is unclear. Our RT-PCR (**Supplemental Fig 3**) and transcriptomics data

suggest that the regulation is post-transcriptional, yet the drivers of these differences remain unknown.

Second, our transcriptomics do not reveal the specific endothelial cell types that drive the inflammatory

responses. We and others have identified the post-capillary venule as a primary location of inflammatory

signaling (25–29, 14). However, a gene expression signature of this cell type to our knowledge remains

elusive. Finally, the mechanism(s) by which YAP drives NF-κB nuclear translocation in adult mice

remains unclear. This deserves further investigation.

Here, we show evidence for novel age-dependent inflammatory signaling in the pulmonary endothelium.

Although pneumonia is a common cause of ARDS and PARDS, ALI signaling that leads to hypoxemia is

orchestrated by the endothelium (15, 30). Our transcriptomics dataset suggests that there are multiple

differences in inflammatory responses in the pulmonary endothelium between weanling and adult age

groups, of which YAP-driven NF-κB is one example. Further investigation into age-dependent

mechanisms of lung injury could aid in the tailoring of therapies across the lifespan.

Grants

This work was supported by NIH K08HL148403; a Stony Wold-Herbert Fund Grant-in-aid; and a

Columbia University Department of Pediatrics Innovation Nucleation Fund Award (R.F.H.).

Acknowledgements

The authors are grateful to Jahar Bhattacharya, Wellington Cardoso, Mohammad N. Islam, and Galina

Gusarova for contributive discussions.

Figure Legends

Figure 1. Responses to intranasal *Pseudomonas aeruginosa* in weanling and adult mice. Analyses

were performed 24h after intranasal instillation of 2.5 x 10^5 CFU of *P. aeruginosa* or PBS. (**A-C, E**) BAL

was obtained by instillation of ice cold PBS intratracheally twice with ~80% recovery. (**A**) BAL protein

was determined by BCA assay. (**B-C**) BAL cell and neutrophil counts were determined with Vetscan

hematology analyzer. (**D**) ELVW was determined in separate experiments. (**E**) BAL cytokines were

measured by multiplex array (Eve Technologies). (**F**) SpO_2 was determined in anesthetized mice on a

warming blanket using a MouseOx® Pulse Oximeter. Data are shown as mean±SE. Number of replicates

are shown by dots (≥4 for all groups). *BAL*, bronchoalveolar lavage; *3wo*, 3 week-old mice;

Pseudomonas, *Pseudomonas aeruginosa; PMNs*, neutrophils; *EVLW*, extravascular lung water; *IL-6*,

interleukin-6; *TNF-α*, tumor necrosis factor α; *SpO₂*, non-invasive O_2 saturation. *$p<0.05$, ***$p<0.001$,

****$p<0.0001$. Multiple comparisons were analyzed by two-way ANOVA; paired differences were

compared by unpaired 2-tailed t test.

Figure 2. YAP expression responses to *P. aeruginosa*-induced lung injury in weanling and adult

mice. Representative gels and scatter plots show immunoblotting and densitometry of enriched

cytoplasmic (**A**) and nuclear (**B**) fractions of whole lung, and enriched lung endothelial cells (**C**) and

epithelial cells (**D**). All animals were treated with i.n. PBS or 2.5 x 10^5 CFU of *P. aeruginosa* 24 hours

before lung removal. Antibodies used were YAP rabbit monoclonal, GAPDH mouse monoclonal, HDAC

mouse monoclonal, and actin rabbit polyclonal. Lanes were run on the same gel. *3wo*, 3 week-old mice;

n≥4 as indicated by dots. *n.s.*, not significant. *$p < 0.05$, ****$p<0.0001$. Differences between groups were

compared by t test.

Figure 3. YAP mediates *P. aeruginosa*-induced lung injury in adult mice. Mice were injected

intravenously with YAP- and TAZ-targeted siRNA or scRNA control 24h before *P. aeruginosa* intranasal

instillation. All mice were treated with *P. aeruginosa*. (**A**) Representative gel and scatter plots show

immunoblotting and densitometry of whole lung cytoplasmic fraction. Antibodies used were YAP rabbit

monoclonal and GAPDH mouse monoclonal. Note that there was high variability of lung YAP expression,

but statistical significance remained when the highest outlier from the scRNA-treated group was removed

from the analysis. (**B**) BAL cell counts were determined with Vetscan hematology analyzer. (**C**) BAL

cytokines were measured by multiplex array. (**D**) SpO_2 was determined using a MouseOx® Pulse

Oximeter. Number of replicates are shown by triangles (≥ 4 for all groups). *BAL*, bronchoalveolar lavage;

3wo, 3 week-old mice; *Pseudomonas*, *Pseudomonas aeruginosa; IL-6*, interleukin-6; *TNF-α*, tumor

necrosis factor α; *SpO_2*, non-invasive O_2 saturation. *$p < 0.05$, **< 0.01. Paired differences were compared

by t test.

Figure 4. YAP expression drives endothelial NF-κB responses in *P. aeruginosa*-infected adult mice.

(**A**) Principal component analysis of four treatment groups. PC1, treatment; PC2, age, orange, saline-

treated; blue, *P. aeruginosa*-treated. •, adult; **x**, weanling. (**B**) Top pathways with up-regulated genes in

adult mice relative to weanling mice in response to *P. aeruginosa* instillation. (**C**) Representative gel and

scatter plots show immunoblotting and densitometry of whole lung cytoplasmic and nuclear enriched

fractions (**C**) 24h after PBS or *P. aeruginosa* instillation and (**D**) following pretreatment with YAP siRNA

by tail vein injection and *P. aeruginosa* instillation. Antibodies used were NF-κB p65 rabbit monoclonal,

YAP rabbit monoclonal, HDAC mouse monoclonal, and actin rabbit polyclonal. Number of replicates are

shown by squares or triangles (≥3 for all groups). *NF-κB, nuclear factor-kappa B; scRNA, scrambled*

siRNA; siRNA, YAP- and TAZ-directed siRNA. *$p<0.05$, **$p<0.01$, ****$p<0.0001$. Paired differences

were compared by t test.

References

1. **Dowell JC**, **Parvathaneni K**, **Thomas NJ**, **Khemani RG**, **Yehya N**. Epidemiology of Cause of Death in Pediatric Acute Respiratory Distress Syndrome. *Crit Care Med* 46: 1811–1819, 2018. doi: 10.1097/CCM.0000000000003371.

2. **Matthay MA**, **Zemans RL**, **Zimmerman GA**, **Arabi YM**, **Beitler JR**, **Mercat A**, **Herridge M**, **Randolph AG**, **Calfee CS**. Acute respiratory distress syndrome. *Nat Rev Primer* 5: 18, 2019. doi: 10.1038/s41572-019-0069-0.

3. **Schouten LR**, **van Kaam AH**, **Kohse F**, **Veltkamp F**, **Bos LD**, **de Beer FM**, **van Hooijdonk RT**, **Horn J**, **Straat M**, **Witteveen E**, **Glas GJ**, **Wieske L**, **van Vught LA**, **Wiewel MA**, **Ingelse SA**, **Cortjens B**, **van Woensel JB**, **Bos AP**, **Walther T**, **Schultz MJ**, **Wösten-van Asperen RM**. Age-dependent differences in pulmonary host responses in ARDS: a prospective observational cohort study. *Ann Intensive Care* 9: 55, 2019. doi: 10.1186/s13613-019-0529-4.

4. **Smith LS**, **Gharib SA**, **Frevert CW**, **Martin TR**. Effects of age on the synergistic interactions between lipopolysaccharide and mechanical ventilation in mice. *Am J Respir Cell Mol Biol* 43: 475–486, 2010. doi: 10.1165/rcmb.2009-0039OC.

5. **Schouten LR**, **Helmerhorst HJ**, **Wagenaar GT**, **Haltenhof T**, **Lutter R**, **Roelofs JJ**, **van Woensel JB**, **van Kaam AH**, **Bos AP**, **Schultz MJ**, **Walther T**, **Wosten-van Asperen RM**. Age-Dependent Changes in the Pulmonary Renin-Angiotensin System Are Associated With Severity of Lung Injury in a Model of Acute Lung Injury in Rats. *Crit Care Med* 44: e1226–e1235, 2016. doi: 10.1097/CCM.0000000000002008.

6. **Ingelse SA**, **Juschten J**, **Maas MAW**, **Matute-Bello G**, **Juffermans NP**, **van Woensel JBM**, **Bem RA**. Fluid restriction reduces pulmonary edema in a model of acute lung injury in mechanically ventilated rats. *PLoS One* 14: e0210172, 2019. doi: 10.1371/journal.pone.0210172.

7. **Totaro A**, **Panciera T**, **Piccolo S**. YAP/TAZ upstream signals and downstream responses. *Nat Cell Biol* 20: 888–899, 2018. doi: 10.1038/s41556-018-0142-z.

8. **Zou R**, **Xu Y**, **Feng Y**, **Shen M**, **Yuan F**, **Yuan Y**. YAP nuclear-cytoplasmic translocation is regulated by mechanical signaling, protein modification, and metabolism. *Cell Biol Int* 44: 1416–1425, 2020. doi: 10.1002/cbin.11345.

9. **van Soldt BJ**, **Cardoso WV**. Hippo-Yap/Taz signaling: Complex network interactions and impact in epithelial cell behavior. *Wiley Interdiscip Rev Dev Biol* 9: e371, 2020. doi: 10.1002/wdev.371.

10. **Kim J**, **Kim YH**, **Kim J**, **Park DY**, **Bae H**, **Lee DH**, **Kim KH**, **Hong SP**, **Jang SP**, **Kubota Y**, **Kwon YG**, **Lim DS**, **Koh GY**. YAP/TAZ regulates sprouting angiogenesis and vascular barrier maturation. *J Clin Invest* 127: 3441–3461, 2017. doi: 10.1172/JCI93825.

11. **Zhou B**, **Flodby P**, **Luo J**, **Castillo DR**, **Liu Y**, **Yu F-X**, **McConnell A**, **Varghese B**, **Li G**, **Chimge N-O**, **Sunohara M**, **Koss MN**, **Elatre W**, **Conti P**, **Liebler JM**, **Yang C**, **Marconett CN**, **Laird-Offringa IA**, **Minoo P**, **Guan K**, **Stripp BR**, **Crandall ED**, **Borok Z**. Claudin-18-mediated YAP activity regulates lung stem and progenitor cell homeostasis and tumorigenesis. *J Clin Invest* 128: 970–984, 2018. doi: 10.1172/JCI90429.

12. **Su K**, **Wang J**, Lv Y, **Tian M**, **Zhao Y-Y**, **Minshall RD**, **Hu G**. YAP Expression in Endothelial
Cells Prevents Ventilator-induced Lung Injury. .

13. **Lv Y**, **Kim K**, **Sheng Y**, **Cho J**, **Qian Z**, **Zhao Y-Y**, **Hu G**, **Pan D**, **Malik AB**, **Hu G**. YAP Controls
Endothelial Activation and Vascular Inflammation Through TRAF6. *Circ Res* 123: 43–56, 2018.
doi: 10.1161/CIRCRESAHA.118.313143.

14. **Emin MT**, **Lee MJ**, **Bhattacharya J**, **Hough RF**. Mitochondria of lung venular capillaries mediate
lung-liver crosstalk in pneumonia. .

15. **Hough RF**, **Islam MN**, **Gusarova GA**, **Jin G**, **Das S**, **Bhattacharya J**. Endothelial mitochondria
determine rapid barrier failure in chemical lung injury. *JCI Insight* 4, 2019. doi:
10.1172/jci.insight.124329.

16. **Schmidt EP**, **Yang Y**, **Janssen WJ**, **Gandjeva A**, **Perez MJ**, **Barthel L**, **Zemans RL**, **Bowman JC**,
Koyanagi DE, **Yunt ZX**, **Smith LP**, **Cheng SS**, **Overdier KH**, **Thompson KR**, **Geraci MW**,
Douglas IS, **Pearse DB**, **Tuder RM**. The pulmonary endothelial glycocalyx regulates neutrophil
adhesion and lung injury during experimental sepsis. *Nat Med* 18: 1217–23, 2012. doi:
10.1038/nm.2843.

17. **Liu M**, **Iosef C**, **Rao S**, **Domingo-Gonzalez R**, **Fu S**, **Snider P**, **Conway SJ**, **Umbach GS**,
Heilshorn SC, **Dewi RE**, **Dahl MJ**, **Null DM**, **Albertine KH**, **Alvira CM**. Transforming Growth
Factor-induced Protein Promotes NF-κB-mediated Angiogenesis during Postnatal Lung
Development. *Am J Respir Cell Mol Biol* 64: 318–330, 2021. doi: 10.1165/rcmb.2020-0153OC.

18. **Sordelli DO**, **Djafari M**, **García VE**, **Fontán PA**, **Döring G**. Age-dependent pulmonary clearance of
Pseudomonas aeruginosa in a mouse model: diminished migration of polymorphonuclear leukocytes
to N-formyl-methionyl-leucyl-phenylalanine. *Infect Immun* 60: 1724–1727, 1992. doi:
10.1128/iai.60.4.1724-1727.1992.

19. **Meli VS**, **Atcha H**, **Veerasubramanian PK**, **Nagalla RR**, **Luu TU**, **Chen EY**, **Guerrero-Juarez**
CF, **Yamaga K**, **Pandori W**, **Hsieh JY**, **Downing TL**, **Fruman DA**, **Lodoen MB**, **Plikus MV**,
Wang W, **Liu WF**. YAP-mediated mechanotransduction tunes the macrophage inflammatory
response. *Sci Adv* 6: eabb8471, 2020. doi: 10.1126/sciadv.abb8471.

20. **LaCanna R**, **Liccardo D**, **Zhang P**, **Tragesser L**, **Wang Y**, **Cao T**, **Chapman HA**, **Morrisey EE**,
Shen H, **Koch WJ**, **Kosmider B**, **Wolfson MR**, **Tian Y**. Yap/Taz regulate alveolar regeneration and
resolution of lung inflammation. *J Clin Invest* 129: 2107–2122, 2019. doi: 10.1172/JCI125014.

21. **Moroishi T**, **Park HW**, **Qin B**, **Chen Q**, **Meng Z**, **Plouffe SW**, **Taniguchi K**, **Yu F-X**, **Karin M**,
Pan D, **Guan K-L**. A YAP/TAZ-induced feedback mechanism regulates Hippo pathway
homeostasis. *Genes Dev* 29: 1271–1284, 2015. doi: 10.1101/gad.262816.115.

22. **Alvira CM**, **Abate A**, **Yang G**, **Dennery PA**, **Rabinovitch M**. Nuclear Factor-κB Activation in
Neonatal Mouse Lung Protects against Lipopolysaccharide-induced Inflammation. *Am J Respir Crit*
Care Med 175: 805–815, 2007. doi: 10.1164/rccm.200608-1162OC.

23. **Iosef C**, **Alastalo T-P**, **Hou Y**, **Chen C**, **Adams ES**, **Lyu S-C**, **Cornfield DN**, **Alvira CM**. Inhibiting
NF-κB in the developing lung disrupts angiogenesis and alveolarization. *Am J Physiol - Lung Cell*
Mol Physiol 302: L1023–L1036, 2012. doi: 10.1152/ajplung.00230.2011.

24. **Alvira CM**. Nuclear factor-kappa-B signaling in lung development and disease: one pathway,
numerous functions. *Birt Defects Res A Clin Mol Teratol* 100: 202–216, 2014. doi:
10.1002/bdra.23233.

25. **Schneeberger EE**. Segmental differentiation of endothelial intercellular junctions in intra-acinar
arteries and veins of the rat lung. *Circ Res* 49: 1102–11, 1981.

26. **Qiao RL**, **Bhattacharya J**. Segmental barrier properties of the pulmonary microvascular bed. *J Appl*
Physiol 1985 71: 2152–9, 1991. doi: 10.1152/jappl.1991.71.6.2152.

27. **Ying X**, **Minamiya Y**, **Fu C**, **Bhattacharya J**. Ca2+ waves in lung capillary endothelium. *Circ Res*
79: 898–908, 1996. doi: 10.1161/01.res.79.4.898.

28. **Kuebler WM**, **Ying X**, **Singh B**, **Issekutz AC**, **Bhattacharya J**. Pressure is proinflammatory in lung
venular capillaries. *J Clin Invest* 104: 495–502, 1999. doi: 10.1172/JCI6872.

29. **Kuebler WM**, **Ying X**, **Bhattacharya J**. Pressure-induced endothelial Ca(2+) oscillations in lung
capillaries. *Am J Physiol Lung Cell Mol Physiol* 282: L917-23, 2002. doi:
10.1152/ajplung.00275.2001.

30. **Hough RF**, **Bhattacharya S**, **Bhattacharya J**. Crosstalk signaling between alveoli and capillaries.
Pulm Circ 8: 2045894018783735, 2018. doi: 10.1177/2045894018783735.

Figure 1: Responses to intranasal *Pseudomonas aeruginosa* in weanling and adult mice

Figure 2: YAP expression responses to *P. aeruginosa*-induced lung injury in weanling and adult mice

A Cytoplasmic:

3wo: 9wo:

YAP

GAPDH

YAP/GAPDH relative intensity

*   ****

PBS *Pseudomonas*   PBS *Pseudomonas*

B Nuclear:

3wo: 9wo:

YAP

HDAC

YAP/HDAC1 relative intensity

ns   ****

PBS *Pseudomonas*   PBS *Pseudomonas*

C   **D**

CD31+:   CD326+:

YAP   YAP

actin   actin

YAP/actin relative intensity

*   ns

PBS *Pseudomonas*   PBS *Pseudomonas*

Figure 3: YAP mediates *P. aeruginosa*-induced lung injury in adult mice

Figure 4: YAP expression drives endothelial NF-κB responses in *P. aeruginosa*-infected adult mice