Arabidopsis thaliana Iron Superoxide Dismutase FeSOD1 Protects ARGONAUTE 1 in a Copper-Dependent Manner

Ariel H. Tomassi[1,2], Ana Perea-García[1,3], Guillermo Rodrigo[4], Javier Sánchez-Vicente[2], Adriana E. Cisneros[2], Marta Olmos[3], Amparo Andrés-Bordería[3], Lucio López-Dolz[2], José-Antonio Daròs[2], Lola Peñarrubia[3,*] and Alberto Carbonell[2,*]

[1]Contributed equally

[2]*Instituto de Biología Molecular y Celular de Plantas (Consejo Superior de Investigaciones Científicas–Universitat Politècnica de València), Valencia, Spain.*

[3]*Departament de Bioquímica i Biología Molecular, Universitat de València, Burjassot, Valencia, Spain*

[4]*Institute for Integrative Systems Biology (I2SysBio), CSIC – University of Valencia, Paterna 46980, Spain*

*Corresponding authors

Author's e-mail addresses:

Ariel H. Tomassi: ahtomas1@ibmcp.upv.es

Ana Perea-García: ana.perea@uv.es

Guillermo Rodrigo: guillermo.rodrigo@csic.es

Javier Sánchez-Vicente: jsanchez@madeinplant.com

Adriana E. Cisneros: adgoncis@ibmcp.upv.es

Marta Olmos: marta.olmos@uv.es

Amparo Andrés-Bordería: Amparo.Andres-Borderia@uv.es

Lucio López-Dolz: lulodolz@gmail.com

José-Antonio Daròs: jadaros@ibmcp.upv.es

Lola Peñarrubia: Lola.Penarrubia@uv.es

Alberto Carbonell: acarbonell@ibmcp.upv.es

Submission date: February 28th 2025

Number of Figures: 6

Number of Tables: 0

Supplementary data: 3 supplemental datasets, 1 supplemental figure, 1 supplemental table, 1 supplemental text

Word count: 4074

Running title: FeSOD1 protects AGO1 in response to copper deficiency

HIGHLIGHT

AGO1 is essential for Cu-deficiency responses but is sensitive to oxidative stress. FeSOD1

interacts with AGO1 and protects it from superoxide radical-induced degradation, thereby

preserving miRNA-mediated gene silencing pathways.

ABSTRACT

Copper (Cu) deficiency compromises plant growth and limits crop productivity. Plants respond

to Cu scarcity by activating the expression of several microRNAs, known as Cu-miRNAs,

which degrade mRNAs from various cuproteins to conserve Cu. Cu-miRNAs, like most

plant miRNAs, associate with ARGONAUTE 1 (AGO1), the primary effector protein of

miRNA-mediated gene silencing pathways, whose function is typically modulated by

interacting proteins acting as cofactors. However, how AGO1 is regulated and functions under

Cu deficiency remains unknown. Here, we searched for AGO1 interactors in *Arabidopsis*

thaliana plants expressing a functional AGO1 protein tagged with the Twin-Strep-tag (TST)

polypeptide, grown under Cu-sufficient or Cu-deficient conditions. TST-AGO1 complexes

were affinity-purified, and proteins were identified using tandem mass spectrometry.

Interestingly, the iron superoxide dismutase 1 (FeSOD1) encoded by *FSD1*, was enriched in

TST-AGO1 complexes purified from plants grown under Cu deficiency. Moreover, *fsd1-2*

mutant plants showed reduced levels of AGO1 compared to wild-type plants under Cu

sufficiency, while both Cu-miRNA-specific and general AGO1 target mRNAs accumulated to

higher levels in *fsd1-2* plants under both Cu-deficient and Cu-sufficient conditions compared

to wild-type plants. These findings suggest that FeSOD1 is essential for proper AGO1 function,

and that its superoxide dismutase activity, which mitigates oxidative stress, enhances AGO1

stability, particularly under Cu deficiency.

Keywords: AGO1, AP-MS, *Arabidopsis thaliana*, copper, FeSOD1, miRNA, TST

INTRODUCTION

Plants require rapid and flexible responses to changing environmental conditions throughout their developmental stages, particularly under stress. MicroRNAs (miRNAs) are a class of small RNAs (sRNAs) that regulate gene expression post-transcriptionally and function as key players in the interplay between development and stress responses (Borges and Martienssen, 2015; Song *et al.*, 2019). miRNAs regulate a wide range of developmental processes including hypocotyl elongation, root development and the transition from vegetative to reproductive phases (Schwab *et al.*, 2005; Meng *et al.*, 2010). Conversely, certain miRNAs are integral to stress responses, modulating plant adaptation to a multitude of environmental conditions such as nutrient availability, low temperatures and drought (Patel *et al.*, 2017; Martin *et al.*, 2010; Buhtz *et al.*, 2010). For instance, miRNAs involved in nutrient stress, including transition metal deficiencies, play significant roles in plant adaptation (Patel *et al.*, 2017) by regulating processes like metal uptake, chelation, antioxidant response and hormone signaling (Ding *et al.*, 2020).

The availability of essential transition metals, such as copper (Cu) and iron (Fe), often limit agricultural productivity (Broadley *et al.*, 2012). These metals, essentials as cofactors in metalloproteins central to life, can also cause toxicity when in excess, leading to increased reactive oxidative species (ROS) production (Farooq *et al.*, 2019). Due to their dual nature as essential and toxic elements, plants have developed complex homeostatic networks to assure their uptake and distribution while preventing toxicity (Puig and Peñarrubia, 2009; Pilon, 2011; Kobayashi *et al.*, 2019; Kumar *et al.*, 2021). ROS are also produced under metal scarcity, as the malfunctioning of key redox metalloproteins in respiratory and photosynthetic electron transport chains causes oxidative stress (Shahbaz and Pilon, 2019). Among ROS, the superoxide radical is one of the most harmful species, capable of damaging macromolecules. ROS detoxification involves superoxide dismutases (SODs), which in plants use different metals as cofactors, functioning in various cellular compartments under diverse conditions. Plant SODs include FeSODs, Cu and zinc (Zn)SODs (Cu/ZnSODs) and manganese (Mn) MnSODs (Ravet and Pilon, 2013).

In *Arabidopsis thaliana* (Arabidopsis), the denoted as Cu-miRNAs play crucial roles in plant responses under Cu deficiency, orchestrating metal redistribution and substituting Cu/ZnSODs with FeSODs (Yamasaki *et al.*, 2007; Pilon, 2017). Cu-miRNAs are proposed to modulate metalloprotein mRNAs post-transcriptionally, establishing a metal prioritization hierarchy and preventing interference with metal-sensitive processes (Perea-García *et al.*, 2022; Shahbaz and Pilon, 2019). Cu-miRNAs, including miR397, miR398, miR408 and

miR857, are induced under Cu deficiency and primarily target mRNAs encoding Cu/ZnSODs
(*CSD1* and *CSD2*), plantacyanin and laccases (Pilon, 2017). Cu-miRNAs containing *cis*-
regulatory domains in their promoters, recognized by a SQUAMOSA BINDING PROTEIN-
LIKE (SPL) family member, SPL7, the master transcription factor driving Cu-deficiency
responses (Yamasaki *et al.*, 2009; Bernal *et al.*, 2012). In addition to Cu-miRNAs, plasma
membrane Cu transporters genes (*COPT1*, *COPT2* and *COPT6*) as well as *FSD1* (encoding
FeSOD1) are regulated similarly (Yamasaki *et al.*, 2007; Bernal *et al.*, 2012; Garcia-Molina *et*
al., 2013). These processes enhance high affinity Cu uptake and metalloprotein substitution as
primary responses to nutrient scarcity.

Importantly, SPL7 belongs to a large family of transcription factors mainly involved in the
control of developmental phase transitions and reproduction (Huijser and Schmid, 2011).
Based on their sizes and similarities, the Arabidopsis SPL family can be grouped in two
subfamilies: large SPLs (including SPL1, SPL7, SPL12, SPL14 and SPL16) and small SPLs
(the remaining 11 SPLs) (Guo *et al.*, 2008; Xing *et al.*, 2010). The small SPLs, with the
exception of SPL8, share a miRNA response element that is complementary to miR156
responsive in the timing of the vegetative and reproductive phase transitions (Rhoades *et al.*,
2002; Schwab *et al.*, 2005; Wu and Poethig, 2006; Gandikota *et al.*, 2007). *SPL3*
overexpressing plants, resistant to miR156 regulation, display a strong reduction in SPL7-
mediated Cu-deficiency responses, probably due to competition between SPL7 and miR156-
regulated SPL3 for *cis* element binding in target promoters (Perea-García *et al.*, 2021).
Therefore, the balance of miR156 SPL targets and non-targets is essential for SPL7-mediated
Cu-deficiency adaptation at different levels, underscoring that the effect of miRNA on its
targets could serve to reinforce the miRNA non-target function. Thus, while Cu-miRNAs
function as post-transcriptional modulators of metalloproteins, allowing flexible Cu
redistribution, other miRNAs, such as miR156, operate upstream, balancing the relative
abundance of SPL transcription factors to integrate Cu-deficiency responses with
developmental processes.

ARGONAUTE (AGO) proteins act as the effector proteins in sRNA-mediated RNA
silencing pathways in eukaryotes (Meister, 2013). In plants, AGOs regulate essential biological
processes such as development, stress responses, genome stability and defense against
pathogens, particularly to viruses (Carbonell, 2017; Fang and Qi, 2016; Carbonell and
Carrington, 2015). They associate with sRNAs such as miRNAs and siRNAs to form the RNA-
induced silencing complex (RISC), which directs the endonucleolytic cleavage or translational
repression of sequence-complementary target RNAs (Huntzinger and Izaurralde, 2011).

Arabidopsis has 10 AGO members that have functionally diverged during the specialization of

sRNA-based RNA silencing pathways (Chapman and Carrington, 2007). AGO1 is the principal

miRNA-associating AGO (Vaucheret *et al.*, 2004) and, consequently, *ago1* mutants show

severe growth defects (Morel *et al.*, 2002). AGO1 protein levels, are regulated at different

transcriptional and posttranscriptional levels (Molinier, 2020). For instance, AGO1 can be

ubiquitinated and degraded through the ubiquitin-proteasome pathway that tags AGOs with

ubiquitin molecules for their degradation by the 26S proteasome (Earley *et al.*, 2010), or

through autophagy by sequestration of AGO1 proteins in autophagosomes (Derrien *et al.*,

2012). In contrast, AGO1 protein is stabilized by its binding to sRNAs (Vaucheret *et al.*, 2006;

Hacquard *et al.*, 2022), as unloaded AGO1 is more prone to degradation, by its sequestration

into stress granules (Leung *et al.*, 2006) or through its association with co-factors. AGO1

physically interacts with several proteins, including chaperones involved in RISC loading such

as HEAT SHOCK PROTEIN 70 (HSP70), HSP90 and CYCLOPHILLIN 40/SQUINT

(CYP40/SQN), which facilitate the conformational opening of AGO1 during sRNA loading

(Iki *et al.*, 2010; Iwasaki *et al.*, 2010; Iki *et al.*, 2012). Moreover, TRANSPORTIN 1 (TRN1),

ENHANCED MiRNA ACTIVITY 1 (EMA1) β and CONSTITUTIVE ALTERATIONS IN

THE SMALL RNAS PATHWAYS 9 (CARP9) proteins interact with AGO1 to modulate

miRNA loading (Wang *et al.*, 2011; Cui *et al.*, 2016; Tomassi *et al.*, 2020), while PROTEIN

ARGININE METHYLTRANSFERASE 5 (PRMT5) associate with AGO1 to promote its

methylation (Martín-Merchán *et al.*, 2024). Finally, proteins known to interact with AGO1 in

response to stress are limited to DAMAGE-SPECIFIC DNA BINDING PROTEIN 2 (DDB2),

TUDOR DOMAN-CONTAINING PROTEIN 1 (TSN1) and TSN2 and SWI/SNF complexes

in response to UV, general stress and hormone or cold stress, respectively (Liu *et al.*, 2018;

Gutierrez-Beltran *et al.*, 2021; Schalk *et al.*, 2017). However, proteins specifically interacting

with AGO1 during the metal stress response have not been identified.

To better understand AGO1's role in response to Cu deficiency, we searched for AGO1

interactors in Arabidopsis transgenic lines expressing a functional AGO1 protein tagged with

the bidentate polypeptide Twin-Strep-tag (TST), grown under Cu-sufficient or Cu-deficient

conditions. AGO1 complexes were affinity-purified, and proteins were identified by tandem

mass spectrometry (AP-MS or affinity purification mass spectrometry). Interestingly, *FSD1-*

encoded FeSOD1, recently proposed as an AGO1 interactor as a result of a yeast two-hybrid

screening (Bressendorff *et al.*, 2023), was enriched in AGO1 complexes purified from plants

grown under Cu deficiency. *fsd1-2* mutant plants accumulated lower levels of AGO1 compared

to Col-0 plants under Cu sufficiency, and both Cu-miRNA-specific and general AGO1 target

mRNAs accumulated to higher levels in *fsd1-2* plants grown under Cu deficiency and sufficiency, respectively, compared to Col-0. These results underscore the role of FeSOD1 in protecting AGO1 stability, especially under Cu deficiency, where its function in miRNA-mediated gene silencing pathways becomes particularly important.

RESULTS

Generation and functional characterization of Arabidopsis transgenic lines expressing TST-tagged AGO1

TST is an advanced affinity tag used for the purification and detection of recombinant proteins under non-denaturing, mild conditions (Schmidt *et al.*, 2013). It consists of a tandem arrangement of two Strep-tag II sequences, designed to enhance binding affinity to Strep-Tactin, a derivative of streptavidin. The TST coding sequence was inserted after AGO1 start codon to tag the N-terminus of AGO1, a previously done with hemagglutinin (HA) or FLAG peptides (Baumberger and Baulcombe, 2005; Carbonell *et al.*, 2012), in the *pAGO1:TST-AGO1* construct, which includes authentic AGO1 regulatory sequences (Figure 1A). As a control, the *pAGO1:TST-GFP* construct, designed to express TST-tagged GFP under the same AGO1 regulatory sequences, was also generated (Figure 1A). To test the functionality of TST-AGO1, *pAGO1:TST-AGO1* and *pAGO1:TST-GFP* constructs were transformed in both Col-0 and *ago1-25* genetic backgrounds. At least, 24 independent T1 transgenic lines were recovered for each construct in both backgrounds. Interestingly, *ago1-25* phenotypic defects, such as slow growth or delayed flowering, were suppressed in plants expressing *pAGO1:TST-AGO1*, but not in plants expressing *pAGO1:TST-GFP* (Figure 1B and 1C). In contrast, plants expressing either construct in the Col-0 background were phenotypically similar to non-transformed Col-0 plants (Figure 1B and 1C). Importantly, plants expressing *pAGO1:TST-AGO1* or *pAGO1-TST-GFP* constructs accumulated high levels of TST-AGO1 and TST-GFP proteins, respectively, as observed in the western blot analysis (Figure 1D). To further investigate TST-AGO1 functionality, AGO1-dependent *TAS1c* tasiRNA 255 (tasiR255) formation was analyzed in T1 transgenic plants. TasiR255 accumulation levels were restored to Col-0 levels in *ago1-25* plants transformed with *pAGO1:TST-AGO1* but not in *ago1-25* plants transformed with *pAGO1:TST-GFP* (Figure 1D). Finally, tasiR255 levels in transgenic lines expressing either construct in the Col-0 background were similar than those of non-transformed Col-0 plants. Overall, these results indicate that the addition of a TST tag at the 5' terminus of Arabidopsis AGO1 does not affect AGO1 functionality. They also suggest that the expression of TST-AGO1 in a Col-0 background does not impair plant growth or AGO1 functionality.

AP-MS analysis of AGO1 protein complexes in response to Cu deficiency stress

Given the previously described key role of miRNA-mediate gene regulation in response to Cu deficiency (Pilon, 2017; Perea-García *et al.*, 2022), we analyzed the AGO1 protein interactome under Cu deficiency compared to Cu sufficiency, using three independent T4 pAGO1:TST-AGO1 transgenic lines to study AGO1 function in both conditions: lines 1, 2 and 3. As a negative control, one T4 *pAGO1:TST-GFP* transgenic line was also grown in parallel under the same conditions. Seedlings of each transgenic line were grown for 10 days under Cu sufficiency (1 µM $CuSO_4$) or under Cu deficiency (100 µM bathocuproine disulphonate, BCS) conditions. AGO1 and GFP protein complexes were purified by affinity chromatography in native conditions. Proteins from the purified complexes were resolved using denaturing polyacrylamide gel electrophoresis (PAGE) and subsequently subjected to in-gel digestion with trypsin. The resulting peptides were extracted from the gel and analyzed by liquid chromatography coupled with tandem mass spectrometry to identify the proteins in the purified complexes. Data S1 shows all hits resulting from proteomic analysis of complexes purified from seedlings of each transgenic line grown under Cu sufficiency and Cu deficiency.

Next, we examined the dynamics of the AGO1 protein interactome in response to Cu deficiency. Proteins from purified samples were processed using the ProteinPilot (Shilov *et al.*, 2007) and Mascot (Perkins *et al.*, 1999) software tools, and the identified proteins were matched against the *Viridiplantae* protein database to filter for plant-specific proteins (Figure 2A). These plant proteins were further analyzed using the BLAST tool (Altschul *et al.*, 1990) against the NCBI Genomes to identify Arabidopsis-specific proteins. The total number of proteins identified in the three *pAGO1:TST-AGO1* lines under Cu deficiency were similar (2185, 2099 and 2002 for lines 1, 2 and 3, respectively), although higher than the number observed in all lines under Cu sufficiency (1565, 1680 and 1749 for lines 1, 2 and 3, respectively) (Figure 2B). The total number of proteins identified in the *pAGO1:TST-GFP* line under Cu deficiency and sufficiency was rather similar (2080 and 1975, respectively) (Figure 2B). We then quantified the change in the abundance of AP-MS interactors under these two conditions (Data S2), discarding those proteins present in the *pAGO1:TST-GFP* control purifications. A $\log_2$ fold change value was assigned for each protein comparison, with $\log_2$ fold > 0 indicating that the protein is more abundant in Cu deficiency and $\log_2$ fold < 0 indicating higher abundance in Cu sufficiency, respectively. A Venn diagram showed that 89 proteins were more abundant in Cu deficiency than in Cu sufficiency in the three lines (Figure 2C, Data S3). Among these proteins, we considered the Fe superoxide dismutase 1 protein

(FeSOD1) for our subsequent analyses, as it was recently identified as an AGO1 interactor in

yeast two hybrid assays (Bressendorff *et al.*, 2023) and was also known to participate in

oxidative stress responses in a Cu-dependent manner (Melicher *et al.*, 2022).

AGO1 and FeSOD1 fine-tune Cu-responsive SOD gene expression

To assess the impact of Cu deficiency on plant growth, Arabidopsis Col-0, *fsd1-2* (loss-of-

function mutant for *FSD1*), and *ago1-25* (mutant for *AGO1*) seedlings were grown under Cu-

sufficient and Cu-deficient conditions (Figure 3A), and their root lengths were quantified

(Figure 3B). Under Cu sufficiency, all genotypes displayed healthy growth, with *fsd1-2*

seedlings showing a slight but statistically significant increase in root length compared to Col-

0 ($P < 0.05$), while *ago1-25* seedlings exhibiting a pronounced decrease as observed before in

other *ago1* mutants (Trolet *et al.*, 2019). Under Cu deficiency, root growth was significantly

inhibited across all genotypes (Figure 3A, 3B).

Expression of SOD genes is an excellent marker for Cu status in plant cells (Yamasaki *et*

al., 2009; Andrés-Colás *et al.*, 2013). To examine SOD expression under our experimental

conditions, mRNA accumulation of *CSD1*, *CSD2*, *FSD1* and *MSD1* was measured by RT-

qPCR (Figure 3C) in Col-0, *fsd1-2* and *ago1-25* seedlings. As previously reported (Yamasaki

et al., 2009), expression of Cu/Zn SODs encoded by *CSD1* and *CSD2* in Col-0 seedlings was

high under Cu sufficiency but remained low under Cu deficiency (Figure 3A). Although the

expression pattern in *ago1-25* was similar, *CSD* expression was significantly higher in most

cases (Figure 3C). Since *CSD* expression is regulated by *miR398*, the impaired AGO1 function

in *ago1-25* could explain this difference. On the other hand, expression of FeSOD1, encoded

by *FSD1,* remained low under Cu sufficiency and was higher under Cu deficiency in both

genotypes, although it was significantly lower in *ago1-25* under Cu deficiency (Figure 3C).

This suggests an indirect effect of AGO1 function on *FSD1* transcription under Cu deficiency,

potentially due to miR156 influencing the ratio of different SPL members and thereby affecting

SPL7's occupancy of Cu deficiency target promoters (Perea-García *et al.*, 2021). Finally,

expression of MnSOD, encoded by MSD1, remained approximately constant across both

genotypes and conditions (Figure 3C).

FeSOD1 is essential for AGO1 stability and functionality, particularly under Cu

deficiency

Next, we investigated whether FeSOD1 function was required for proper AGO1 expression.

To do so, AGO1 protein levels were measured by western blot in 10-day-old Col-0, *fsd1-2* and

ago1-25 seedlings grown under Cu sufficiency and Cu deficiency conditions (Figure 4A).

Surprisingly, AGO1 protein levels in Col-0 plants drastically dropped (around 90%) in Cu

deficiency compared to Cu sufficiency. This decrease was similar to that observed in *ago1-25*

plants grown under both Cu conditions. Additionally, AGO1 protein levels also decreased by

about 80% in *fsd1-2* seedlings compared to Col-0 plants under Cu sufficiency (Figure 4A).

These results indicate an important decrease in AGO1 protein in Col-0 plants grown content

under Cu deficiency, and in *fsd1-2* seedlings grown under both Cu conditions.

To further investigate whether the reduction in AGO1 protein levels was due to

transcriptional regulation, we analyzed *AGO1* mRNA levels by RT-qPCR (Figure 4B) in the

same samples used for AGO1 protein analysis (Figure 4A). Under Cu sufficiency, *AGO1*

transcript levels were significantly higher in *fsd1-2* compared to Col-0, suggesting that

FeSOD1 function may influence *AGO1* expression. Additionally, *ago1-25* mutants exhibited a

significant increase in *AGO1* mRNA levels relative to Col-0, likely due to impaired miR168-

mediated AGO1 mRNA regulation caused by reduced AGO1 activity (Vaucheret *et al.*, 2006).

Under Cu deficiency, *AGO1* transcript accumulation markedly increased in Col-0, *fsd1-2* and

ago1-25 seedlings compared to Cu sufficiency, with *ago1-25* displaying the highest expression

levels (Figure 4B). These results indicate that the reduction in AGO1 protein levels under Cu

deficiency is not due to decreased AGO1 transcript accumulation.

To further assess the relationship between *AGO1* mRNA and protein levels, we calculated

the relative ratio of *AGO1* mRNA to AGO1 protein accumulation in the same samples (Figure

4C). Under Cu sufficiency, this ratio (R) was much higher in *fsd1-2* (R=4.9) and *ago1-25*

(R=40.6) mutants compared to Col-0 (R=1), indicating that despite increased *AGO1* transcript

levels, AGO1 protein accumulation remained low in these mutants. Under Cu deficiency, the

AGO1 mRNA/protein ratio increased more drastically across all genotypes, particularly in

ago1-25, which exhibited the highest ratio (R=50.2) among the conditions tested.

To determine whether the decrease in AGO1 protein content also affected its functionality

under Cu deficiency, the expression of several Cu-miRNA targets (*ARPN,* miR160 target; *CCS,*

CSD1, CSD2, miR398 targets*;* GST25, miR396 target; *LAC3*, miR397 target) was analyzed by

RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings (Figure 5A). As expected, the

expression of all Cu-miRNA targets (except for *GSTU25*) significantly increased in *ago1-25*

compared to Col-0. Similarly, all miRNA targets were induced to varying degrees in *fsd1-2*

compared to Col-0, with the increase in *CSD1* being statistically significant (Figure 5A). These

findings confirm that AGO1 functionality requires the presence of FeSOD1 under Cu

deficiency.

Next, we wondered whether FeSOD1 activity was also required for general AGO1 function

under Cu sufficiency and for regulating other non-Cu-miRNA targets. The expression of

different well-known miRNA targets (*ARF8*, miR167 target; *NF-YA9*, miR169 target; *SPL3*

and *SPL9*, miR156 targets; *TCP4*, miR159 target) was analyzed by RT-qPCR in 10-day-old

Col-0, *fsd1-2* and *ago1-25* seedlings grown under Cu sufficiency (Figure 5B). As expected, the

expression of all miRNA targets was significantly higher in *ago1-25* compared to Col-0.

Interestingly, the expression of all miRNA targets also increased in *fsd1-2* compared to Col-0,

with a statistically significant increase observed for four out of five miRNA targets (*ARF8*,

SPL3, *SPL9* and *TCP4*). Notably, the increased expression of miR156 targets (*SPL3* and *SPL9*)

is relevant, as these targets compete with SPL7 in Cu-deficiency responses (Perea-García *et*

al., 2021). Finally, the expression of the miRNA targets mentioned above was also analyzed in

seedlings of the different genotypes grown under Cu deficiency (Figure S1). The expression of

all miRNA targets was higher in both *fsd1-2* and *ago1-25* compared to Col-0, although this

increase was significant only for *NF-YA9* and *SPL3* in *ago1-25* (Figure S1).

In conclusion, our results indicate that FeSOD1 is necessary for proper AGO1 function in

miRNA-mediated gene silencing. We propose a model (Figure 6) in which FeSOD1's

superoxide dismutase activity protects against oxidative stress, contributing to the increased

stability of the AGO1 protein. This protective role of FeSOD1 is especially important under

Cu deficiency due to the elevated oxidative stress and the critical role played by Cu-miRNAs

in the plant's response to this nutrient deficit. Absence of FeSOD1 in *fsd1-2* mutants results in

AGO1 degradation independently of the Cu levels (Figure 6).

DISCUSSION

This study investigates plant responses under Cu deficiency conditions, where Cu-miRNAs

regulate the mRNAs encoding cuproteins. Different studies have reported that numerous

miRNAs are induced by a variety of environmental stresses, suggesting the existence of shared

signaling pathways (Martin *et al.*, 2010). Plants face a conflictive situation when responding

to adverse conditions, give the extensive role of miRNAs in stress responses and the instability

of the key protein AGO1 within RISC under suboptimal conditions (Fang and Qi, 2016). To

solve this conflict, providing a protective environment to preserve AGO1 function could be

crucial for an effective stress response. Here, we explored how Cu deficiency influences the

regulation of miRNA targets mediated by AGO1. We analyzed the AGO1 protein interactome

in *pAGO1:TST-AGO1* Arabidopsis transgenic plants grown under both normal and Cu-

deficient conditions. Due to the adverse effects of Cu deficiency on AGO1 function, we

searched for interactor proteins with a protective effect on AGO1. Notably, a well-documented

response to Cu deficiency is the substitution of Cu/ZnSODs with FeSODs to support

antioxidative defense (Yamasaki *et al.*, 2009). In this context, FeSOD1 was enriched in TST-

AGO1 complexes purified from all three *pAGO1:TST-AGO1* lines grown in Cu deficiency.

Since FeSOD1 was recently identified as an AGO1 interactor in yeast two-hybrid assays

(Bressendorff *et al.*, 2023), the role of FeSOD1-AGO1 interaction became the main focus of

our study.

Particularly striking is the opposite pattern observed between *AGO1* mRNA and protein

levels, suggesting that either AGO1 translation or stability is severely compromised under Cu

deficiency. This is evidenced by the significant decrease in endogenous AGO1 protein levels

in Col-0 plants grown under Cu-deficient conditions compared to Cu-sufficient conditions. A

similar opposite pattern is also observed in *fsd1-2* and, more drastically, in the *ago1-25* mutant,

possibly indicating a compensatory increase in *AGO1* mRNA synthesis in response to low

AGO1 protein levels (Figure 4).

The elevated ROS levels induced by Cu deficiency appear to be a key factor explaining

AGO1 degradation, as the presence of FeSOD1 detoxifies the cellular oxidative state and

safeguards AGO1 function in regulating Cu-miRNA genes or other known targets (Figure 5A,

Figure 6, Figure S1). Consistently, as peroxide is the reaction product of SODs, the peroxidase

PRX32 is also identified as antioxidant molecules interacting with AGO1 under Cu deficiency

(Data S3). Whether ROS-induced AGO1 degradation occurs via the ubiquitin-proteasome

system, the autophagy pathway or simultaneously through both remains unknown. It is possible

that specific proteins target AGO1 for degradation, as reported for the F-box protein FBW2

that assembles an SCF complex that selectively targets for proteolysis AGO1 when it is

unloaded (Hacquard *et al.*, 2022).

Alongside Cu-miRNAs, SPL7 can also induce *FSD1* expression (Yamasaki *et al.*, 2009).

Our results indicate that defects in AGO1 function impact Cu-miRNAs and *FSD1* expression

(Figure 3C). The reduced expression of *FSD1* in *ago1-25* likely results from defective miR156

targeting. Indirectly, *FSD1* expression is affected by miR156, which main function is to inhibit

the gene expression of some members of the SPL protein family (especially SPL3), while it

does not affect other members such as SPL7. The purpose seems to be reducing competition

between SPL factors for target promoter binding (Perea-García *et al.*, 2021). Consequently, the

defective targeting of miR156 in *ago1-25* explains the increased expression of non-SPL7

members (SPL3 and SPL9), potentially disrupting SPL7 responses, including *FSD1* expression

(Figure 5). Additionally, the expression of miR398 targets, *CSDs* and *CCS*, is also reduced in

ago1-25 (Figure 3C).

To test whether the FeSOD1-AGO1 interaction functions reciprocally, we examined the role of FeSOD1 in AGO1 function. FeSOD1 is essential for ROS response, specifically in eliminating superoxide radicals (Ravet and Pilon, 2013). In this sense, the interaction between AGO1 and FeSOD1 suggests that antioxidant protection is necessary for AGO1 function. To check this, we studied the AGO1 protein content in *fsd1-2* mutants. Homozygous *fsd1-2* seedlings grown under Cu-deficient and Cu-excess conditions exhibit reduced ROS neutralization, affecting phenotypes like increased chlorosis (Melicher *et al.*, 2022). Moreover, this capacity depends on Cu availability, as *FSD1* expression is regulated by Cu status. AGO1 levels depend on Cu status and the presence of FeSOD1, supporting the hypothesis that Cu deficiency and FeSOD1 have a protective role against oxidative damage to AGO1 (Figure 3B). Furthermore, the effects on different Cu-miRNA targets (*ARPN*, *CCS*, *CSD2*, *GSTU25*, *LAC3*) (Figure 5A) highlights the defect in AGO1 function in the absence of this protective barrier. Taken together, our findings reveal the challenges plants face under Cu deficiency in maintaining RNA interference efficiency, given the essential role of Cu-miRNAs and the stability of AGO1 protein under these conditions. The substitution of Cu/ZnSOD with FeSOD1 serves as a solution to this compromise (Figure 6). Increased Fe demand for FeSOD1 highlights the interaction between Cu and Fe homeostasis (Perea-García *et al.*, 2021; Waters *et al.*, 2012). Notably, IRT1, the main Fe transporter, was enriched in TST-AGO1 complexes purified under Cu deficiency. Further studies are needed to clarify the role of the AGO1-IRT1 interaction, although an indirect association via IRT1-RNA binding proteins with the RISC may be plausible.

While stress conditions often reveal molecular imbalances, such as RNA interference functioning with unstable AGO1, these balances could also exist under normal conditions, as observed here for AGO1 under Cu sufficiency. Despite low *FSD1* levels under Cu sufficiency, the expression of non-Cu-miRNAs, such as miR167 (*ARF8*) (Wu *et al.*, 2006), miR169 (*NF-YA9*) (Sorin *et al.*, 2014), miR156 (*SPL3* and *SPL9*) (Kim *et al.*, 2012) and miR159 (*TCP4*) (Palatnik *et al.*, 2007) was also affected in *fsd1-2* (Figure 5B). These findings suggest a broader scenario (Figure 6) where maintaining an antioxidant environment around AGO1 is crucial for proper RNA interference function, particularly under Cu deficiency.

MATERIALS AND METHODS

Plant materials, growth conditions and plant phenotyping

ago1-25 is a hypomorphic mutant obtained in an EMS mutagenesis and described before
(Morel *et al.*, 2002). *fsd1-2* (GABI_740E11) is a mutant line derived from a T-DNA insertion
and described before (Melicher *et al.*, 2022). Wild-type Arabidopsis cv Columbia 0 (Col-0),
ago1-25 and *fsd1-2* seeds were grown on plates where Cu concentration was 1 mM, considered
as Cu sufficiency, as previously described (Perea-García *et al.*, 2020). For severe Cu deficiency
condition, 100 μM of (BCS) was added to the growth medium. Seedlings were grown for 10
days at neutral photoperiodic conditions (12 h light, 23°C/12 h darkness, 16°C). Root length
of 10-day old seedlings was measured using ImageJ (Abramoff *et al.*, 2004).
409

Transgene constructs
pENTR-pAGO1-3xHA-AGO1 (Carbonell *et al.*, 2012) was digested with *Hind*III and *Kpn*I, and
the two fragments (7812 bp and 1425 bp) were gel-purified. The 1425 bp fragment was ligated
into *Hind*III/*Kpn*I-digested *pBSIIKS+* to generate *pBS-pAGO1-3xHA*. A 4266 bp linear *pBS-*
pAGO1 fragment lacking the 3xHA tag was amplified using *pBS-pAGO1-3xHA* as template
and oligonucleotide pair D2475/D2476, gel purified and assembled with a 136 bp *TST-AGO1*
fragment using a plasmid containing a *TST-GFP* cassette and oligonucleotide pair D2477/2478
in the presence of GeneArt Gibson Assembly HiFi Master Mix (Invitrogen) to generate *pBS-*
pAGO1-TST. *pBS-pAGO1-TST* was digested with *Hind*III-*Kpn*I, and the resulting 1437 bp
fragment was ligated to the 7812 bp mentioned before, to generate *pENTR-pAGO1-TST-AGO1*.
Finally, the *pAGO1-TST-AGO1* cassette was cloned by LR recombination into *pMDC99*, a
Gateway-compatible plant transformation vector (Curtis and Grossniklaus, 2003), to generate
pMDC99-pAGO1-TST-AGO1. Nucleotide and amino acid sequences of TST-AGO1 and TST-
GFP are listed in Text S1.
A 6003 bp and an 863 bp fragment was amplified from *pENTR-pAGO1-3xHA-AGO1*
(Carbonell *et al.*, 2012) and *TST-GFP* plasmid with oligonucleotide pair AC-22/AC-23 and
AC-20/AC-21, respectively. Both fragments were assembled in the presence of GeneArt
Gibson Assembly HiFi Master Mix (Invitrogen) to generate *pENTR-pAGO1-TST-GFP*. The
pAGO1-TST-GFP was transferred by LR recombination into *pMDC99* to generate *pMDC99-*
pAGO1-TST-GFP.
430

RNA and protein preparation
Total RNA form Arabidopsis seedlings or inflorescences was isolated as before (Cisneros *et*
al., 2023). Triplicate samples from pools of Arabidopsis seedlings or inflorescences were
analyzed. Total protein extracts were prepared in 2X PDB buffer (0.0625 M Tris pH 6.8, 2%

SDS, 10% glycerol, 10% mercaptoethanol, 0.02% bromophenol blue) in a 1:5 tissue:buffer

ratio.

Real-time RT-qPCR

cDNA synthesis and qPCR analysis were performed essentially as before (Cisneros *et al.*,

2025). Briefly, five hundred ng of DNaseI-treated total RNA was used to produce cDNA from

10-day-old Arabidopsis seedlings with the PrimeScript RT Reagent Kit (Perfect Real Time,

Takara) according to manufacturer's instructions. RT-qPCR was done on optical 96-well plates

in a QuantStudio 3 Real-Time PCR system (Thermo Fisher Scientific, Waltham, MA, USA)

following these conditions: 20 s at 95ºC, followed by 40 cycles of 95ºC for 3 s and 60ºC for 30

s, with an additional melt curve stage consisting of 15 s at 95ºC, 1 min at 60ºC and 15 s at 95ºC.

The 20-μl reaction included 10 μl of 2× TB Green Premix Ex Taq (Takara), 2 μl of diluted

complementary DNA (1:5), 0.4 μl of ROX II Reference Dye (50X) and 300 nM of each gene-

specific primer. Oligonucleotides used for RT-qPCR are listed in Table S1. Target mRNA

expression levels were calculated relative to reference genes *ACT2* and *UBQ10*, using the delta

delta cycle threshold comparative method of QuantStudio Design and Analysis software,

version 1.5.1 (Thermo Fisher Scientific). Three independent biological replicates, and two

technical replicates for each biological replicate were analyzed.

Small RNA blot assays

Small RNA blot assays and band quantification from radioactive membranes were done as

described (Cisneros *et al.*, 2022). Oligonucleotides used as probes for sRNA blots are listed in

Table S1.

Protein blot assays

Proteins were separated in NuPAGE Novex 4-12% Bis-Tris gels (Invitrogen), transferred to

Protran nitrocellulose membranes (Amersham) and detected by chemiluminescence using

specific antibodies and SuperSignal West Pico PLUS chemiluminescent substrate

(ThermoFisher Scientific) as before (Cisneros *et al.*, 2025). Here, for detection of TST-tagged

proteins, StrepMAB-Classic HRP Twin-Strep-Tag antibody (IBA) at a 1:10000 dilution was

used. For detection of endogenous AGO1 and HISTONE3 (H3), anti-AGO1 (Agrisera) and

anti-Histone 3 (Abcam) were used at 1:10000 and 1:4000 dilutions, respectively, and

conjugated with 1:20000 of goat anti-rabbit IgG horseradish peroxidase secondary antibody

(ThermoFisher Scientific). Images were acquired with an ImageQuant 800 CCD imager

(Cytiva) and analyzed with ImageQuantTL v10.2 (Cytiva). Ponceau red S solution (Thermo

Fisher Scientific) staining of membranes was used to verify the global protein content of the

samples.

Statistical analysis

Statistical tests are described in the figure legends. Significant differences were determined

with two-tailed Student's *t*-test ($P < 0.05$).

Affinity purification of protein complexes

Protein complexes containing TST-tagged protein products were purified from 1.5 g of 12-

day-old seedlings by affinity chromatography in native conditions using a 1-ml Strep-Tactin

XT spin column (IBA) as previously described (Martinez *et al.*, 2016).

Protein identification by liquid chromatography coupled with tandem mass spectrometry

Protein preparations were separated by SDS-PAGE using 4% and 10% acrylamide for stacking

and resolving gel, respectively. Electrophoresis was stopped when the front entered 1 cm into

the resolving gel, and the gel was stained with Coomassie. The bands were cut for subsequent

in-gel digestion with trypsin (12.5 ng/ul). The digest was acidified with 0.5% TFA to stop

digestion. The peptides were desalted using C18 columns according to the manufacturer's

protocol (Pierce C18 Spin Columns, Thermo Scientific). The peptides were dried in a speedvac

and resuspended in 15 µL of mobile phase (2% ACN, 0.05% TFA) for injection into HPLC.

The mixture of tryptic peptides was analyzed by high-resolution mass spectrometry using an

Orbitrap Fusion (Thermo Scientific) in "Data Dependent Acquisition" mode, equipped with a

nanoESi ion source. The peptide extract, diluted in 2% acetonitrile (ACN) and 0.05%

trifluoroacetic acid (TFA), was first separated in an Ultimate 3000 Dionex UHPLC (Thermo

Scientific) under the following chromatographic conditions: preconcentration on a C18 pre-

column (PepMap100, 5 µm, 300 µm x 5 mm, Thermo Scientific) at a flow rate of 5 µL/min for

3 minutes in 98% acetonitrile and 0.1% TFA. The chromatographic gradient of 4–40%

acetonitrile was applied at a flow rate of 300 nL/min on a C18 nano-column (Acclaim PepMap

RSLC 75 µm x 50 cm, Thermo Scientific) over 60 minutes. The total chromatography time

was 85 minutes. The Orbitrap Fusion operated in positive mode with a voltage of 2 kV. The

mass range in "full scan" mode was 400–1500 m/z in the Orbitrap at a resolution of 120,000,

with an AGC target of $4x10^5$ and a dynamic "injection time." Peptide analysis was performed

in "data dependent acquisition" mode, operating in MS/MS Top 50 mode, fragmenting as many

high-intensity ions as possible per "full scan" within a cycle time of 3 seconds. Fragmentation

spectra were acquired in the linear trap by CID fragmentation at a collision energy of 35%,

with a dynamic injection time and an AGC target of $1x10^2$. A dynamic exclusion time of 15

seconds was included for each ion to avoid repetitive ion fragmentation. Protein identification

was done using Proteome Discoverer 2.1 (PD2.1, Thermo Fisher Scientific). Proteins were

identified with the following conditions: database, Uniprot_Arabidopsis-thaliana + AGO-

sequences; tolerance, 10 ppm for Orbitrap and 0.02 Da for Ion Trap; maximum of 1 missed

cleavages; *b-ions weight 1 (CID) and 0.3 (HCD); *y-ions weigh: 1 (CID-HCD); dynamic

modifications, oxidation (Met); fixed modifications, carbamidomethylation (Cys) (*depends

on the fragmentation type: CID or HCD); acceptance criteria, Percolator (Delta CN<0.05);

FDR of 1% against decoy database. Resulting Excel sheets contain the identifications at the

peptide and protein levels (Data S1). The data included in the report are filtered by a 1% FDR

based on the probability values obtained by Percolator. The minimum number of peptides per

protein is 1.

Computational analysis

A previously developed bioinformatic pipeline (Ambrós *et al.*, 2021)was used to process all

information obtained via proteomics. From the raw proteomic lists, all plant proteins were

identified and mapped to Arabidopsis genes (proteins), keeping the normalized MS peak area

as a quantitative measure. Specifically, we identified proteins from green plants by GenInfo

Identifier (GI) through Mascot analysis and conducted sequence alignments using BLAST to

identify individual genes in *A. thaliana*, using data from The Arabidopsis Information Resource

(TAIR) version 10 (Lamesch *et al.*, 2012). For each hit, the associated amino acid sequence

was retrieved from the NCBI database in FASTA format using the Biopython Entrez module.

Only alignments with e-values < 0.0001 were considered, and any redundant elements were

excluded. The protein list obtained from the Cu deficiency condition was compared against the

list obtained from the Cu sufficiency condition in a quantitative manner (*i.e.*, by subtracting

the MS peak areas) for each transgenic line, discarding those proteins present in control

pAGO1-TST-GFP lists for each condition. After the control filtering, if a protein was in one

list but not in the other, we considered $|log_2 \text{ fold}| = 100$ (arbitrary value, denoted by Inf in the

figures); positive for proteins detected in Cu deficiency but not in sufficiency and negative for

proteins detected in Cu sufficiency but not in deficiency. All genes (proteins) in the lists were

functionally annotated according to the information from TAIR10.

Gene and virus identifiers

Arabidopsis gene identifiers are: *ACT2* (AT3G18780), *AGO1* (AT1G48410), *ARF8* (AT5G37020), *ARPN* (AT2G02850), *CCS* (AT1G12520), *CSD1* (AT1G08830), *CSD2* (AT2G28190), *FSD1* (AT4G25100), *GSTU25* (AT1G17180), *LAC3* (AT2G30210), *MSD1* (AT3G10920) *NF-YA9* (AT3G20910), *SPL3* (AT2G33810), *SPL9* (AT2G42200), *TCP4* (AT3G15030), *UBQ10* (AT4G05320).

ACKOWLEDGEMENTS

This work was supported by grants or fellowships from MCIN/AEI/10.13039/501100011033 (co-financed by the European Regional Development Fund) [RTI2018-095118-A-I00, PID2021-122186OB-I00 and RYC-2017-21648 to A.C.; PID2021-127671NB-I00 to G.R.; PID2020-116940RB-I00 to L.P. and A.P.G.; BIO2017-83184-R and PID2020-114691RB-I00 to J.A.D.; PRE2019-088439 to A.E.C.], and from the Generalitat Valenciana (CIPROM/2022/21 to G.R. and J.A.D.) We thank Dr. Tomás Takác for sharing *fsd1-2* seeds and Carlos Fuentes from the Proteomics Unit of University of Córdoba for the assistance in preparing and analyzing purified proteins.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

A.C. and L.P. planned and designed the research. A.H.T., A.P.G., A.C., A.E.C., L.L.D., A.A.B. and M.O. performed most of the experiments. J.S.V. and J.A.D. performed AGO1 affinity purifications. G.R. did the computational analysis of AP-MS samples. A.H.T., A.P.G, L.P and A.C. analyzed data. A.C. and L.P. wrote the manuscript with input from rest of the authors.

DATA AVAILABILITY STATEMENT

All data relating to this manuscript can be found within the manuscript and its supplementary files. Data that support the findings of this study are available from the corresponding authors upon reasonable request. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al.*, 2022) partner repository with the dataset identifier PXD057108.

SUPPLEMENTAL DATA

Data S1. Proteins identified by AP-MS in *pAGO1:TST-AGO1* and *pAGO1:TST-GFP*

transgenic lines grown under Cu deficiency and Cu sufficiency.

Data S2. Proteins identified by AP-MS in each of the three *pAGO1:TST-AGO1* transgenic lines

grown under Cu deficiency versus Cu sufficiency, and not present in *pAGO1:TST-GFP* AP-

MS samples for each condition.

Data S3. Proteins identified by AP-MS in all three *pAGO1:TST-AGO1* transgenic lines grown

under Cu deficiency versus Cu sufficiency, and not present in *pAGO1:TST-GFP* AP-MS

samples for each condition.

Figure S1. Accumulation of miRNA target mRNAs in different Arabidopsis genotypes under

Cu deficiency.

Table S1. Name, sequence and use of oligonucleotides used in the present study.

Text S1. Nucleotide and amino acid sequences of TST-based inserts.

REFERENCES

Abramoff, M., Magalhaes, P. and Ram, S. (2004) Image processing with ImageJ. *Biophotonics International*, **11**, 36–42.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990) Basic local alignment search tool. *J Mol Biol*, **215**, 403–10.

Ambrós, S., Gómez-Muñoz, N., Giménez-Santamarina, S., Sánchez-Vicente, J., Navarro-López, J., Martínez, F., Daròs, J.-A. and Rodrigo, G. (2021) Molecular signatures of silencing suppression degeneracy from a complex RNA virus. *PLOS Computational Biology*, **17**, e1009166.

Andrés-Colás, N., Perea-García, A., Mayo de Andrés, S., Garcia-Molina, A., Dorcey, E., Rodríguez-Navarro, S., Pérez-Amador, M.A., Puig, S. and Peñarrubia, L. (2013) Comparison of global responses to mild deficiency and excess copper levels in Arabidopsis seedlings†. *Metallomics*, **5**, 1234–1246.

Baumberger, N. and Baulcombe, D.C. (2005) Arabidopsis ARGONAUTE1 is an RNA slicer that selectively recruits microRNAs and short interfering RNAs. *Proc Natl Acad Sci U S A*, **102**, 11928–33.

Bernal, M., Casero, D., Singh, V., et al. (2012) Transcriptome Sequencing Identifies SPL7-Regulated Copper Acquisition Genes FRO4/FRO5 and the Copper Dependence of Iron Homeostasis in Arabidopsis. *The Plant Cell*, **24**, 738–761.

Borges, F. and Martienssen, R.A. (2015) The expanding world of small RNAs in plants. *Nat Rev Mol Cell Biol*, **16**, 727–41.

Bressendorff, S., Kausika, S., Zobbe Sjøgaard, I.M., Oksbjerg, E., Michels, A., Poulsen, C. and Brodersen, P. (2023) The N-coil and the globular N-terminal domain of plant ARGONAUTE1 are interaction hubs for regulatory factors. *Biochemical Journal*, BCJ20230025.

Broadley, M., Brown, P., Buerkert, A., et al. (2012) Contributing Authors. In P. Marschner, ed. *Marschner's Mineral Nutrition of Higher Plants (Third Edition)*. San Diego: Academic Press, p. xi. Available at: https://www.sciencedirect.com/science/article/pii/B978012384905200025X [Accessed October 24, 2024].

Buhtz, A., Pieritz, J., Springer, F. and Kehr, J. (2010) Phloem small RNAs, nutrient stress responses, and systemic mobility. *BMC Plant Biology*, **10**, 64.

Carbonell, A. (2017) Plant ARGONAUTEs: Features, Functions, and Unknowns. *Methods Mol Biol*, **1640**, 1–21.

Carbonell, A. and Carrington, J.C. (2015) Antiviral roles of plant ARGONAUTES. *Curr Opin Plant Biol*, **27**, 111–7.

Carbonell, A., Fahlgren, N., Garcia-Ruiz, H., Gilbert, K.B., Montgomery, T.A., Nguyen, T., Cuperus, J.T. and Carrington, J.C. (2012) Functional analysis of three Arabidopsis ARGONAUTES using slicer-defective mutants. *Plant Cell*, **24**, 3613–29.

Carrió-Seguí, A., Garcia-Molina, A., Sanz, A. and Peñarrubia, L. (2015) Defective Copper Transport in the copt5 Mutant Affects Cadmium Tolerance. *Plant and Cell Physiology*, **56**, 442–454.

Chapman, E.J. and Carrington, J.C. (2007) Specialization and evolution of endogenous small RNA pathways. *Nat Rev Genet*, **8**, 884–96.

Cisneros, A.E., Alarcia, A., Llorens-Gámez, J.J., Puertes, A., Juárez-Molina, M., Primc, A. and Carbonell, A. (2025) Syn-tasiR-VIGS: Virus-Based Targeted RNAi in Plants By Synthetic Trans Acting Small Interfering RNAs Derived from Minimal Precursors. *Nucleic Acids Research (in press)*. https://doi.org/10.1093/nar/gkaf183

Cisneros, A.E., Martín-García, T., Primc, A., Kuziuta, W., Sánchez-Vicente, J., Aragonés, V., Daròs, J.-A. and Carbonell, A. (2023) Transgene-free, virus-based gene silencing in plants by artificial microRNAs derived from minimal precursors. *Nucleic Acids Research*, **51**, 10719–10736.

Cisneros, A.E., Torre-Montaña, A. de la and Carbonell, A. (2022) Systemic silencing of an endogenous plant gene by two classes of mobile 21-nucleotide artificial small RNAs. *Plant J*, **110**, 1166–1181.

Cui, Y., Fang, X. and Qi, Y. (2016) TRANSPORTIN1 promotes the association of microRNA with ARGONAUTE1 in Arabidopsis. *Plant Cell*, **28**, 2576–2585.

Curtis, M.D. and Grossniklaus, U. (2003) A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiol*, **133**, 462–9.

Derrien, B., Baumberger, N., Schepetilnikov, M., Viotti, C., De Cillia, J., Ziegler-Graff, V., Isono, E., Schumacher, K. and Genschik, P. (2012) Degradation of the antiviral component ARGONAUTE1 by the autophagy pathway. *Proc Natl Acad Sci U S A*, **109**, 15942–6.

Ding, Y., Ding, L., Xia, Y., Wang, F. and Zhu, C. (2020) Emerging Roles of microRNAs in Plant Heavy Metal Tolerance and Homeostasis. *J. Agric. Food Chem.*, **68**, 1958–1965.

Earley, K., Smith, M., Weber, R., Gregory, B. and Poethig, R. (2010) An endogenous F-box protein regulates ARGONAUTE1 in Arabidopsis thaliana. *Silence*, **1**, 15.

Fang, X. and Qi, Y. (2016) RNAi in Plants: An Argonaute-Centered View. *Plant Cell*, **28**, 272–85.

Farooq, M.A., Niazi, A.K., Akhtar, J., Saifullah, Farooq, M., Souri, Z., Karimi, N. and Rengel, Z. (2019) Acquiring control: The evolution of ROS-Induced oxidative stress and redox signaling pathways in plant stress responses. *Plant Physiology and Biochemistry*, **141**, 353–369.

Gandikota, M., Birkenbihl, R.P., Hohmann, S., Cardon, G.H., Saedler, H. and Huijser, P. (2007) The miRNA156/157 recognition element in the 3' UTR of the Arabidopsis SBP box gene SPL3 prevents early flowering by translational inhibition in seedlings. *Plant J*, **49**, 683–93.

Garcia-Molina, A., Andrés-Colás, N., Perea-García, A., Neumann, U., Dodani, S.C., Huijser, P., Peñarrubia, L. and Puig, S. (2013) The Arabidopsis COPT6 Transport Protein Functions in Copper Distribution Under Copper-Deficient Conditions. *Plant and Cell Physiology*, **54**, 1378–1390.

Guo, A.-Y., Zhu, Q.-H., Gu, X., Ge, S., Yang, J. and Luo, J. (2008) Genome-wide identification and evolutionary analysis of the plant specific SBP-box transcription factor family. *Gene*, **418**, 1–8.

Gutierrez-Beltran, E., Elander, P.H., Dalman, K., Dayhoff, G.W., Moschou, P.N., Uversky, V.N., Crespo, J.L. and Bozhkov, P.V. (2021) Tudor staphylococcal nuclease is a docking platform for stress granule components and is essential for SnRK1 activation in Arabidopsis. *The EMBO Journal*, **40**, e105043.

Hacquard, T., Clavel, M., Baldrich, P., et al. (2022) The Arabidopsis F-box protein FBW2 targets AGO1 for degradation to prevent spurious loading of illegitimate small RNA. *Cell Reports*, **39**, 110671.

Huijser, P. and Schmid, M. (2011) The control of developmental phase transitions in plants. *Development*, **138**, 4117–4129.

Huntzinger, E. and Izaurralde, E. (2011) Gene silencing by microRNAs: contributions of translational repression and mRNA decay. *Nat Rev Genet*, **12**, 99–110.

Iki, T., Yoshikawa, M., Meshi, T. and Ishikawa, M. (2012) Cyclophilin 40 facilitates HSP90-mediated RISC assembly in plants. *EMBO J*, **31**, 267–78.

Iki, T., Yoshikawa, M., Nishikiori, M., Jaudal, M.C., Matsumoto-Yokoyama, E., Mitsuhara, I., Meshi, T. and Ishikawa, M. (2010) In vitro assembly of plant RNA-induced silencing complexes facilitated by molecular chaperone HSP90. *Mol Cell*, **39**, 282–91.

Iwasaki, S., Kobayashi, M., Yoda, M., Sakaguchi, Y., Katsuma, S., Suzuki, T. and Tomari, Y. (2010) Hsc70/Hsp90 chaperone machinery mediates ATP-dependent RISC loading of small RNA duplexes. *Mol Cell*, **39**, 292–9.

Kim, J.J., Lee, J.H., Kim, W., Jung, H.S., Huijser, P. and Ahn, J.H. (2012) The microRNA156-SQUAMOSA PROMOTER BINDING PROTEIN-LIKE3 Module Regulates Ambient Temperature-Responsive Flowering via FLOWERING LOCUS T in Arabidopsis. *Plant Physiology*, **159**, 461–478.

Kobayashi, T., Nozoye, T. and Nishizawa, N.K. (2019) Iron transport and its regulation in plants. *Free Radical Biology and Medicine*, **133**, 11–20.

Kumar, V., Pandita, S., Singh Sidhu, G.P., Sharma, A., Khanna, K., Kaur, P., Bali, A.S. and Setia, R. (2021) Copper bioavailability, uptake, toxicity and tolerance in plants: A comprehensive review. *Chemosphere*, **262**, 127810.

Lamesch, P., Berardini, T.Z., Li, D., et al. (2012) The Arabidopsis Information Resource (TAIR): improved gene annotation and new tools. *Nucleic Acids Research*, **40**, D1202–D1210.

Leung, A.K.L., Calabrese, J.M. and Sharp, P.A. (2006) Quantitative analysis of Argonaute protein reveals microRNA-dependent localization to stress granules. *Proceedings of the National Academy of Sciences*, **103**, 18125–18130.

Liu, C., Xin, Y., Xu, L., Cai, Z., Xue, Y., Liu, Yong, Xie, D., Liu, Yule and Qi, Y. (2018) *Arabidopsis* ARGONAUTE 1 Binds Chromatin to Promote Gene Transcription in Response to Hormones and Stresses. *Developmental Cell*, **44**, 348-361.e7.

Martin, R.C., Liu, P.P., Goloviznina, N.A. and Nonogaki, H. (2010) microRNA, seeds, and Darwin?: diverse function of miRNA in seed biology and plant responses to stress. *J Exp Bot*, **61**, 2229–34.

Martinez, F., Rodrigo, G., Aragones, V., Ruiz, M., Lodewijk, I., Fernandez, U., Elena, S.F. and Daros, J.A. (2016) Interaction network of tobacco etch potyvirus NIa protein with the host proteome during infection. *BMC Genomics*, **17**, 87.

Martín-Merchán, A., Lavatelli, A., Engler, C., González-Miguel, V.M., Moro, B., Rosano, G.L. and Bologna, N.G. (2024) Arabidopsis AGO1 N-terminal extension acts as an essential hub for PRMT5 interaction and post-translational modifications. *Nucleic Acids Research*, gkae387.

Meister, G. (2013) Argonaute proteins: functional insights and emerging roles. *Nat Rev Genet*, **14**, 447–59.

Melicher, P., Dvořák, P., Krasylenko, Y., Shapiguzov, A., Kangasjärvi, J., Šamaj, J. and Takáč, T. (2022) Arabidopsis Iron Superoxide Dismutase FSD1 Protects Against Methyl Viologen-Induced Oxidative Stress in a Copper-Dependent Manner. *Front. Plant Sci.*, **13**. Available at: https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2022.823561/full [Accessed October 4, 2024].

Meng, Y., Ma, X., Chen, D., Wu, P. and Chen, M. (2010) MicroRNA-mediated signaling involved in plant root development. *Biochemical and Biophysical Research Communications*, **393**, 345–349.

Mira, H., Martínez-García, F. and Peñarrubia, L. (2001) Evidence for the plant-specific intercellular transport of the Arabidopsis copper chaperone CCH. *The Plant Journal*, **25**, 521–528.

Molinier, J. (2020) The Multilayer's Control of ARGONAUTE 1 Contents. *Molecular Plant*, **13**, 1–3.

Morel, J.B., Godon, C., Mourrain, P., Beclin, C., Boutet, S., Feuerbach, F., Proux, F. and Vaucheret, H. (2002) Fertile hypomorphic ARGONAUTE (ago1) mutants impaired in post-transcriptional gene silencing and virus resistance. *Plant Cell*, **14**, 629–39.

Palatnik, J.F., Wollmann, H., Schommer, C., et al. (2007) Sequence and Expression Differences Underlie Functional Specialization of *Arabidopsis* MicroRNAs miR159 and miR319. *Developmental Cell*, **13**, 115–125.

Patel, P., Yadav, K. and Ganapathi, T.R. (2017) Small and Hungry: MicroRNAs in Micronutrient Homeostasis of Plants. *MicroRNA*, **6**, 22–41.

Perea-García, A., Andrés-Bordería, A., Huijser, P. and Peñarrubia, L. (2021) The Copper-microRNA Pathway Is Integrated with Developmental and Environmental Stress Responses in Arabidopsis thaliana. *Int J Mol Sci*, **22**, 9547.

Perea-García, A., Andrés-Bordería, A., Vera-Sirera, F., Pérez-Amador, M.A., Puig, S. and Peñarrubia, L. (2020) Deregulated High Affinity Copper Transport Alters Iron Homeostasis in Arabidopsis. *Front Plant Sci*, **11**, 1106.

Perea-García, A., Puig, S. and Peñarrubia, L. (2022) The role of post-transcriptional modulators of metalloproteins in response to metal deficiencies. *Journal of Experimental Botany*, **73**, 1735–1750.

Perez-Riverol, Y., Bai, J., Bandla, C., et al. (2022) The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Research*, **50**, D543–D552.

Perkins, D.N., Pappin, D.J.C., Creasy, D.M. and Cottrell, J.S. (1999) Probability-based protein identification by searching sequence databases using mass spectrometry data. *ELECTROPHORESIS*, **20**, 3551–3567.

Pilon, M. (2011) Moving copper in plants. *New Phytologist*, **192**, 305–307.

Pilon, M. (2017) The copper microRNAs. *New Phytologist*, **213**, 1030–1035.

Puig, S. and Peñarrubia, L. (2009) Placing metal micronutrients in context: transport and distribution in plants. *Current Opinion in Plant Biology*, **12**, 299–306.

Ravet, K. and Pilon, M. (2013) Copper and Iron Homeostasis in Plants: The Challenges of Oxidative Stress. *Antioxidants & Redox Signaling*, **19**, 919–932.

Rhoades, M.W., Reinhart, B.J., Lim, L.P., Burge, C.B., Bartel, B. and Bartel, D.P. (2002) Prediction of plant microRNA targets. *Cell*, **110**, 513–20.

Schalk, C., Cognat, V., Graindorge, S., Vincent, T., Voinnet, O. and Molinier, J. (2017) Small RNA-mediated repair of UV-induced DNA lesions by the DNA DAMAGE-BINDING PROTEIN 2 and ARGONAUTE 1. *Proc Natl Acad Sci U S A*. Available at: https://www.ncbi.nlm.nih.gov/pubmed/28325872 http://www.pnas.org/content/early/2017/03/20/1618834114.full.pdf.

Schmidt, T.G., Batz, L., Bonet, L., et al. (2013) Development of the Twin-Strep-tag(R) and its application for purification of recombinant proteins from cell culture supernatants. *Protein expression and purification*, **92**, 54–61.

Schwab, R., Palatnik, J.F., Riester, M., Schommer, C., Schmid, M. and Weigel, D. (2005) Specific effects of microRNAs on the plant transcriptome. *Developmental cell*, **8**, 517–27.

Shahbaz, M. and Pilon, M. (2019) Conserved Cu-MicroRNAs in Arabidopsis thaliana Function in Copper Economy under Deficiency. *Plants*, **8**, 141.

Shilov, I.V., Seymour, S.L., Patel, A.A., Loboda, A., Tang, W.H., Keating, S.P., Hunter, C.L., Nuwaysir, L.M. and Schaeffer, D.A. (2007) The Paragon Algorithm, a Next

Generation Search Engine That Uses Sequence Temperature Values and Feature Probabilities to Identify Peptides from Tandem Mass Spectra*. *Molecular & Cellular Proteomics*, **6**, 1638–1655.

Song, X., Li, Y., Cao, X. and Qi, Y. (2019) MicroRNAs and Their Regulatory Roles in Plant-Environment Interactions. *Annu Rev Plant Biol*. Available at: https://www.ncbi.nlm.nih.gov/pubmed/30848930.

Sorin, C., Declerck, M., Christ, A., et al. (2014) A miR169 isoform regulates specific NF-YA targets and root architecture in Arabidopsis. *New Phytologist*, **202**, 1197–1211.

Tomassi, A.H., Re, D.A., Romani, F., Cambiagno, D.A., Gonzalo, L., Moreno, J.E., Arce, A.L. and Manavella, P.A. (2020) The Intrinsically Disordered Protein CARP9 Bridges HYL1 to AGO1 in the Nucleus to Promote MicroRNA Activity. *Plant Physiology*, **184**, 316–329.

Trolet, A., Baldrich, P., Criqui, M.-C., Dubois, M., Clavel, M., Meyers, B.C. and Genschik, P. (2019) Cell Cycle–Dependent Regulation and Function of ARGONAUTE1 in Plants. *The Plant Cell*, **31**, 1734–1750.

Vaucheret, H., Mallory, A.C. and Bartel, D.P. (2006) AGO1 homeostasis entails coexpression of MIR168 and AGO1 and preferential stabilization of miR168 by AGO1. *Mol Cell*, **22**, 129–36.

Vaucheret, H., Vazquez, F., Crete, P. and Bartel, D.P. (2004) The action of ARGONAUTE1 in the miRNA pathway and its regulation by the miRNA pathway are crucial for plant development. *Genes Dev*, **18**, 1187–97.

Wang, W., Ye, R., Xin, Y., Fang, X., Li, C., Shi, H., Zhou, X. and Qi, Y. (2011) An importin beta protein negatively regulates MicroRNA activity in Arabidopsis. *Plant Cell*, **23**, 3565–76.

Waters, B.M., McInturf, S.A. and Stein, R.J. (2012) Rosette iron deficiency transcript and microRNA profiling reveals links between copper and iron homeostasis in Arabidopsis thaliana. *Journal of Experimental Botany*, **63**, 5903–5918.

Wu, G. and Poethig, R.S. (2006) Temporal regulation of shoot development in Arabidopsis thalianaby miR156 and its target SPL3. *Development*, **133**, 3539–3547.

Wu, M.-F., Tian, Q. and Reed, J.W. (2006) Arabidopsis microRNA167 controls patterns of ARF6 and ARF8 expression, and regulates both female and male reproduction. *Development*, **133**, 4211–4218.

Xing, S., Salinas, M., Höhmann, S., Berndtgen, R. and Huijser, P. (2010) miR156-Targeted and Nontargeted SBP-Box Transcription Factors Act in Concert to Secure Male Fertility in Arabidopsis. *The Plant Cell*, **22**, 3935–3950.

Yamasaki, H., Abdel-Ghany, S.E., Cohu, C.M., Kobayashi, Y., Shikanai, T. and Pilon, M. (2007) Regulation of Copper Homeostasis by Micro-RNA in *Arabidopsis*. *Journal of Biological Chemistry*, **282**, 16369–16378.

Yamasaki, H., Hayashi, M., Fukazawa, M., Kobayashi, Y. and Shikanai, T. (2009) SQUAMOSA Promoter Binding Protein–Like7 Is a Central Regulator for Copper Homeostasis in Arabidopsis. *The Plant Cell*, **21**, 347–361.

FIGURE LEGENDS

Figure 1. Phenotypic and molecular analyses of Arabidopsis Col-0 and *ago1-25* transgenic plants expressing TST-tagged AGO1 or GFP. A, Diagram of the constructs. TST, GFP and AGO1 sequences are shown in orange, green and dark blue boxes, respectively. AGO1 endogenous promoter and 3' UTR regulatory sequences are shown in a light blue arrow and box, respectively. **B**, Pictures of 21-day-old Col-0 (top panel) and *ago1-25* (bottom panel) T1 transgenic plants, homozygous for the transgene. Col-0 and *ago1-25* non-transgenic controls (NTCs) are included. **C**, Box plot representing the mean flowering time of Col-0 and *ago1-25* NTC and T1 transgenic plants (*n* = 24). Pairwise Student's *t*-test comparisons are represented with an * if significantly different (*P* < 0.05) compared to Col-0 NTC. **D**, Accumulation of *TAS1c*-dependent tasiRNA (tasiR255) and TST-AGO1 in T1 transgenic lines. The graph at the top shows the mean (*n* = 3) relative to tasiR255 levels + standard deviation (Col-0 = 1.0). One blot from three biological replicates is shown. Each biological replicate is a pool of at least eight independent lines that were randomly selected. U6 and L-Rubisco (ribulose-1,5-bisphosphate carboxylase/oxygenase) blots are shown as loading controls. Other details are as in C.

Figure 2. Protein analyses in TST-AGO1 complexes purified by AP-MS. A, Flowchart of the process for plant protein identification. Proteins from purified samples are processed through ProteinPilot and Mascot software tools, and identified proteins are matched against the *Viridiplantae* protein database to filter for plant proteins. The plant proteins are further analyzed using the Blast tool against the NCBI Genomes database to identify Arabidopsis-specific proteins. **B**, Bar chart showing the total number of proteins identified by AP-MS in the three different *pAGO1-TST-AGO1* transgenic lines grown under copper (Cu) deficiency (light grey bars) or Cu sufficiency (dark grey bars). **C**, Venn diagram comparing the total number of proteins in Cu deficiency versus Cu sufficiency identified in the three *pAGO1:TST-AGO1* transgenic lines. The cutoff value for significant fold change is set at $\log_2$ fold = 1, meaning that only proteins enriched in Cu deficiency versus Cu sufficiency are represented. Overlapping section indicates the proteins identified in purified samples from independent transgenic lines.

Figure 3. Analysis of Arabidopsis superoxide dismutase genes under copper sufficiency and deficiency conditions in Col-0, *fsd1-2* and *ago1-25* seedlings. A, Photographs of 10-day-old of Col-0, *fsd1-2* and *ago1-25* seedlings grown on plates with MS medium in Cu sufficiency (1 µM Cu) or deficiency (100 µM BCS). **B**, Quantification of root length of Col-0,

fsd1-2 and *ago1-25* seedlings grown in the same conditions than in A. Mean relative level (*n* = 40-45) + standard error of root length. Pairwise Student's *t*-test comparisons are represented with an * if significantly different (*P* < 0.05) compared to Col-0 in each condition. **C**, Relative expression of *CSD1, CSD2, FSD1* and *MSD1* mRNAs determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control (Cu sufficiency, 1 µM CuSO$_4$) and Cu deficiency (100 µM BCS). Mean relative level (*n* = 3) + standard error of mRNAs after normalization to *ACTIN2*. Bars with an * are significantly different from that of the Col-0 control sample (*P* < 0.05 in all pairwise Student's *t*-test comparisons). Other *t*-test comparisons are also shown. Each biological replicate is a pool of at least twenty independent lines that were randomly selected.

Figure 4. Accumulation of AGO1 protein and mRNA in Arabidopsis Col-0, *fsd1-2* and *ago1-25* seedlings. A, Accumulation of endogenous AGO1 protein in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control (Cu sufficiency) and Cu deficiency. The graph at the top shows the mean (*n* = 3) relative to endogenous AGO1 levels + standard deviation (Col-0 = 1.0). One blot from three biological replicates is shown. Each biological replicate is a pool of at least twenty independent lines that were randomly selected. HISTONE H3 (H3) blot is shown as loading control. **B**, Relative expression of *AGO1* determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control and Cu deficiency. Mean relative level (*n* = 3) + standard error of mRNA after normalization to *ACTIN2* (Col-0 = 1.0). Bars with an * are significantly different from that of the Col-0 control sample (*P* < 0.05 in all pairwise Student's *t*-test comparisons). Each biological replicate is a pool of at least twenty independent lines that were randomly selected.

Figure 5. Accumulation of miRNA target genes in Arabidopsis Col-0, *fsd1-2* and *ago1-25* seedlings. A, Relative expression of Cu-miRNA targets *ARPN*, *CCS*, *LAC3* and *GSTU25* determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under Cu deficiency. Mean relative level (*n* = 3) + standard error of mRNAs after normalization to *ACTIN2* (Col-0 = 1.0). Bars with an * are significantly different from that of the Col-0 control sample (*P* < 0.05 in all pairwise Student's *t*-test comparisons). Each biological replicate is a pool of at least twenty independent lines that were randomly selected. **B,** Relative expression of miRNA targets *ARF8*, *NF-YA9, SPL3*, *SPL9* and *TCP4* determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under Cu sufficiency (1 µM CuSO$_4$). Other details are as in A.

Figure 6. Model for the anti-superoxide protection of the AGO1 function by the FeSOD1 dome under Cu deficiency. Top, under Cu sufficiency Cu-miRNA and *FSD1* expression are essentially blocked. A basal level of iron superoxide dismutase FeSOD1 allows the protection of AGO1 from reactive oxygen species (ROS) derived from normal cellular metabolism. Middle, under Cu deficiency, the increase presence of ROS, including superoxide anion, induces a certain level of AGO1 degradation. However, in these conditions SPL7 outcompetes SPL3 and SPL9 for binding to cis-regulatory transcriptional elements (GTACs) present in promoters of Cu-miRNA genes and FeSOD1-encoding *FSD1,* inducing their expression. Increased levels of FeSOD1 play a protective role of the effector complex of RNAi by detoxifying superoxide anion. By directly interacting with AGO1 and/or associated RNA Binding Proteins (RBP), FeSOD1 promotes AGO1 stability contributing to Cu-miRNA and general AGO1 functions. Bottom, in *fsd1* mutants, the lack of FeSOD impedes proper detoxification of ROS and leads to AGO1 degradation in both Cu sufficiency (left) and Cu deficiency (right) conditions.

Figure 1. Phenotypic and molecular analyses of Arabidopsis Col-0 and *ago1-25* transgenic plants expressing TST-tagged AGO1 or GFP. **A**, Diagram of the constructs. TST, GFP and AGO1 sequences are shown in orange, green and dark blue boxes, respectively. AGO1 endogenous promoter and 3' UTR regulatory sequences are shown in a light blue arrow and box, respectively. **B**, Pictures of 21-day-old Col-0 (top panel) and *ago1-25* (bottom panel) T1 transgenic plants, homozygous for the transgene. Col-0 and *ago1-25* non-transgenic controls (NTCs) are included. **C**, Box plot representing the mean flowering time of Col-0 and *ago1-25* NTC and T1 transgenic plants ($n = 24$). All data points are shown on the plots. Pairwise Student's *t*-test comparisons are represented with an * if significantly different ($P < 0.05$) compared to Col-0 NTC. **D**, Accumulation of *TAS1c*-dependent tasiRNA (tasiR255) and TST-AGO1 in T1 transgenic lines. The graph at the top shows the mean ($n = 3$) relative to tasiR255 levels + standard deviation (Col-0 = 1.0). One blot from three biological replicates is shown. Each biological replicate is a pool of at least eight independent lines that were randomly selected. U6 and L-Rubisco (ribulose-1,5-bisphosphate carboxylase/oxygenase) blots are shown as loading controls. Other details are as in C.

Figure 2. Protein analyses in TST-AGO1 complexes purified by AP-MS. **A**, Flowchart of the process for plant protein identification. Proteins from purified samples are processed through ProteinPilot and Mascot software tools, and identified proteins are matched against the *Viridiplantae* protein database to filter for plant proteins. The plant proteins are further analyzed using the Blast tool against the NCBI Genomes database to identify Arabidopsis-specific proteins. **B**, Bar chart showing the total number of proteins identified by AP-MS in the three different *pAGO1-TST-AGO1* transgenic lines grown under copper (Cu) deficiency (light grey bars) or Cu sufficiency (dark grey bars). **C**, Venn diagram comparing the total number of proteins in Cu deficiency versus Cu sufficiency identified in the three *pAGO1:TST-AGO1* transgenic lines and not in *pAGO1:TST-GFP*. The cutoff value for significant fold change is set at $\log_2$ fold = 1, meaning that only proteins enriched in Cu deficiency versus Cu sufficiency are represented. Overlapping section indicates the proteins identified in purified samples from independent transgenic lines.

A **Plant phenotypes**

Col-0   *fsd1-2*   *ago1-25*

Cu-sufficiency

Cu-deficiency

B **Root length quantification**

C **Expression of Arabidopsis superoxide dismutase genes**

Figure 3. Analysis of Arabidopsis superoxide dismutase genes under copper sufficiency and deficiency conditions in Col-0, *fsd1-2* and *ago1-25* seedlings. A, Photographs of 10-day-old of Col-0, *fsd1-2* and *ago1-25* seedlings grown on plates with MS medium in Cu sufficiency (1 µM Cu) or deficiency (100 µM BCS). **B,** Quantification of root length of Col-0, *fsd1-2* and *ago1-25* seedlings grown in the same conditions than in A. Mean relative level (n = 40-45) + standard error of root length. Pairwise Student's t-test comparisons are represented with an * if significantly different (P < 0.05) compared to Col-0 in each condition. Other t-test comparisons are also shown. **C,** Relative expression of *CSD1, CSD2, FSD1* and *MSD1* mRNAs was determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control (Cu sufficiency, 1 µM CuSO$_4$) and Cu deficiency (100 µM BCS). Mean relative level (n = 3) + standard error of mRNAs after normalization to *ACTIN2*. Bars with an * are significantly different from that of the Col-0 control sample (P < 0.05 in all pairwise Student's t-test comparisons). Other t-test comparisons are also shown. Each biological replicate is a pool of at least twenty independent lines that were randomly selected.

Figure 4. Accumulation of AGO1 protein and mRNA in Arabidopsis Col-0, *fsd1-2* and *ago1-25* seedlings. A, Accumulation of endogenous AGO1 protein in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control (Cu sufficiency) and Cu deficiency. The graph at the top shows the mean (n = 3) relative to endogenous AGO1 levels + standard deviation (Col-0 = 1.0). One blot from three biological replicates is shown. Each biological replicate is a pool of at least twenty independent lines that were randomly selected. HISTONE H3 (H3) blot is shown as loading control. **B**, Relative expression of *AGO1* determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control and Cu deficiency. Mean relative level (n = 3) + standard error of mRNA after normalization to *ACTIN2* (Col-0 = 1.0). Bars with an * are significantly different from that of the Col-0 control sample (P < 0.05 in all pairwise Student's *t*-test comparisons). Each biological replicate is a pool of at least twenty independent lines that were randomly selected. **C**, Relative AGO1 mRNA/AGO1 protein accumulation in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under control and Cu deficiency.

Figure 5. Accumulation of miRNA target genes in Arabidopsis Col-0, *fsd1-2* and *ago1-25* seedlings. A, Relative expression of Cu-miRNA targets *ARPN*, *CCS*, *CSD1*, *CSD2*, *LAC3* and *GSTU25* determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under Cu deficiency. Mean relative level ($n = 3$) + standard error of mRNAs after normalization to *ACTIN2* (Col-0 = 1.0). Bars with an * are significantly different from that of the Col-0 control sample ($P < 0.05$ in all pairwise Student's *t*-test comparisons). Each biological replicate is a pool of at least twenty independent lines that were randomly selected. **B**, Relative expression of miRNA targets *ARF8*, *NF-YA9*, *SPL3*, *SPL9* and *TCP4* determined by RT-qPCR in 10-day-old Col-0, *fsd1-2* and *ago1-25* seedlings, grown under Cu sufficiency (1 μM $CuSO_4$). Other details are as in panel A.

Figure 6. Model for the anti-superoxide protection of the AGO1 function by the FeSOD dome under Cu deficiency. Top, under Cu sufficiency Cu-miRNA and FSD1 expression are essentially blocked. A basal level of iron superoxide dismutase FeSOD allows the protection of AGO1 from reactive oxygen species derived from normal cellular metabolism. Middle, under Cu deficiency, the increase presence of ROS, including superoxide anion, induces a certain level of AGO1 degradation. However, in these conditions SPL7 outcompetes SPL3 and SPL9 for binding to cis-regulatory transcriptional elements (GTACs) present in promoters of Cu-miRNA genes and FeSOD-encoding *FSD1,* inducing their expression. Increased levels of FeSOD play a protective role of the effector complex of RNAi by detoxifying superoxide anion. By directly interacting with AGO1 and/or associated RNA Binding Proteins (RBP), FeSOD promotes AGO1 stability contributing to Cu-miRNA and general AGO1 functions. Bottom, in *fsd1* mutants, the lack of FeSOD impedes proper detoxification of ROS and leads to AGO1 degradation in both Cu sufficiency (left) and Cu deficiency (right) conditions.