Diversity and single-cell activity of endolithic microbes in sediment-hosted carbonate nodules within and below the sulfate-methane transition zone

Sergio A. Parra[a]#, Magdalena J. Mayr[a,b], James Mullahoo[a], Laura K. Quinn[c&], Rebecca L. Wipfler[a], Victoria J. Orphan[a,b]#

[a]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California, USA

[b]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, USA

[c]Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California, USA

Running Head: **Endolithic microbes in carbonate nodules**

#Address correspondence to Sergio A. Parra, serg.a.parra@gmail.com; Victoria J. Orphan, vorphan@caltech.edu.

&Present Address: Laura K. Quinn, Saint-Gobain Ceramics & Plastics, Inc., Northborough, MA, USA

Abstract

Authigenic carbonate concretions ('nodules') precipitate in marine seep sediments as a result of anaerobic oxidation of methane (AOM). These rocks host active endolithic microbial communities and persist as important methane sinks. Still, how these communities and their activity differ from those in adjacent seep sediments, particularly as a function of proximity to the sulfate-methane transition zone (SMTZ), remains understudied. Here, we sampled sediments and nodules within and below the SMTZ (0-57 cm deep) at four active deep-sea seep areas in Santa Monica Basin, CA. Measurements of high nodule porosities (43-51%) coupled to strong similarities between sediment and nodule 16S rRNA-based community profiles, including below the SMTZ, suggest continued perfusion and exchange between buried nodules and surrounding sediment. Shared, depth-dependent transitions in methanotrophic taxa (ANME-1, ANME-2, ANME-3) and methanogenic taxa (*Methanofastidiosales*) below the SMTZ were also consistent with trends in porewater methane and sulfate concentrations, porewater DIC, and nodule $\delta^{13}C$ values — underscoring the impact of different geochemical conditions on community structure and suggestive of under-characterized physiological plasticity in ANME-1. Laboratory-based BONCAT incubations of nodules within the SMTZ over ~14-weeks revealed active sulfide production and translationally active endolithic microorganisms. However, cells from parallel nodule incubations recovered below the SMTZ showed weak-to-negligible BONCAT-based activity despite similar cell abundances — suggestive of low activity on shorter timescales or dormancy. Together, these data challenge the interpretation of passively recorded microbiological signatures in seep sediment-hosted carbonate nodules and expand our understanding of how these endolithic communities may be actively shaped by past and present conditions.

Importance

This study advances earlier investigations of microbial communities in buried seep carbonate nodules by integrating microbiological profiles of nodules and sediments, sediment geochemistry, single-cell activity measurements, and nodule mineral, geochemical, and physical characteristics within and below the sulfate-methane transition zone in deep-sea methane seeps. This approach allows us to view how nodule-hosted, endolithic microbial communities change relative to their surrounding sediments across multiple geochemical contexts and better understand how formation history and environmental conditions might affect community identity and metabolic function. Results indicate that the seep nodule and surrounding sediment communities are closely linked across diverse geochemical conditions. This connectivity between sediments and carbonate nodules is distinct from that observed in exhumed seep carbonates, with implications regarding how microbial community composition within these nodules are interpreted, suggesting that instead of a passive recorder of the communities at the time of formation, these nodules appear to retain diverse, metabolically viable communities.

Introduction

Authigenic carbonates at marine methane seeps form as the result of microbially-mediated anaerobic oxidation of methane (AOM) in sediments, which draws down a substantial amount of the subsurface methane produced at these seeps (1, 2). Because these deposits persist over geologic timescales, these carbonates have been interpreted as a record of past AOM activity in ancient marine environments (3–7). However, several lab-based measurements of elevated methane oxidation rates in recovered seep carbonates demonstrate that these rocks continue to

host viable microorganisms and AOM activity, representing a persistent methane sink whose

activity and ecology remains under-studied compared to seep sediments (8–11).

In particular, vuggy carbonate concretions, or 'nodules' observed in seep sediments (12–15) have

not been investigated as thoroughly as denser, seafloor-exposed seep carbonates — whose

metabolically diverse, endolithic communities have been observed to differ significantly from

sediments and sediment-hosted nodules collected from the same sites, e.g., (16). As such, seep

nodules represent a unique opportunity to study seep carbonate microbial community structure

and activity in comparison to their host sediments in geochemical context.

For example, AOM in marine sediments primarily occurs within a distinct redox regime known

as the sulfate-methane transition zone (SMTZ), where syntrophic activity between anaerobic,

methanotrophic (ANME) archaea and sulfate-reducing bacteria (SRB) act to consume porewater

methane and sulfate (17–20). This zone of AOM activity varies in depth and thickness largely

controlled by methane flux, e.g., (21), transitioning from methane-poor, sulfate-rich overlying

sediment horizons to deeper sulfate-poor, methane-rich sediments. The impact of relative

proximity to the SMTZ on the endolithic community structure and activity within sediment-

hosted carbonate nodules remains poorly characterized, with limited sampling from existing

studies focusing exclusively within this zone (14–16, 22). In sediments, this geochemical

stratification often results in regimes of distinct ANME/SRB lineages indicative of niche

differentiation (22–29), where the observed persistence of certain ANME clades in sulfate-

depleted seep sediments remains an area of active investigation (30–32). As such, the SMTZ

represents a useful comparative framework in which to study the similarities and differences

with co-occurring carbonate nodules across this globally important geochemical transition zone.

Previous work examining nodules recovered from seep sediments reported high similarities in

the 16S rRNA diversity and community structure with host sediments (14–16). However, the

extent to which these DNA-based community profiles represent extant, metabolically viable

communities actively exchanged with host sediment or preserved, ancient sediment communities

remains unclear (14, 15), with implications how different traces of AOM are ultimately

interpreted and/or preserved in sediment-hosted nodules.

In this study, we expanded the characterization of sediment-hosted authigenic carbonate nodules

from four methane seep areas at Santa Monica Mounds 800 and 863 within the Santa Monica

Basin (33). Specifically, we integrated the biological and porewater geochemical profiles of 1 m

long, ROV-deployed sediment cores within active deep-sea seeps with the structural and

microbiological profiles of seep carbonate nodules, effectively capturing changes in both

sediment and associated nodules within and well below the SMTZ. Laboratory microcosm

experiments were also used to characterize differences in bulk respiration and cellular anabolic

activity by the endolithic community across this transition zone using geochemical assays and

single cell bioorthogonal non-canonical amino acid tagging (BONCAT) incubations. This

combined approach allows us to probe the historical and current capacity of seep carbonate

nodules as hosts for continued AOM activity in geochemical context, compared to their

surrounding sediments.

Results

Sediment and nodule sampling within Santa Monica Basin methane seeps

Nodules and sediment push cores in this study were collected in May 2021 during the WF05-21

Southern California oceanographic expedition on the *R/V Western Flyer* using the *ROV Doc*

Ricketts, both owned and operated by the Monterey Bay Aquarium Research Institute (MBARI).

93 samples (7 cores, 82 sediment horizons, 11 nodule horizons) were recovered in total from two

active methane seep sites within the Santa Monica Basin: Santa Monica Mound Site 800 (SMM

800: 33.799438N /118.64672E; 805m depth; also known as the NE mound) and Santa Monica

Mound Site 863 (SMM 863: 33.7888N /118.6683W; 863 m; also known as the SW mound); (33,

34). At these sites, we investigated 4 distinct seep areas: SMM 800-I, SMM 800-II, SMM 800-

III, and SMM 863 (Fig. 1). Within each seep area, 1-2 sediment cores were collected, including

3, 1.22 m long cores (LC) and 4, standard, 30 cm long push cores (PC), with quasi-duplicate

('paired') cores taken in close spatial proximity (~30-50 cm apart) to capture comparable

environments, where possible. One sediment core (PC64, SMM 800-II) was collected during a

previous cruise and sampled from the same microbial mat within 5 m from the 2021 sampling

location (Supplemental Material).

124
Fig. 1 – A) General location of the Santa Monica Basin seep sites (NE Mound, 800m, and SW mound, 863m)
sampled for this study; B) Multi-beam bathymetry map of the sampling seep areas at SMM 800 and SMM 863; C)
higher-resolution bathymetry map of the seep areas sampled at SMM 800: SMM 800-I, SMM 800-II, and SMM
800-III. Bathymetry colors in B) and C) correspond to mapped depth. Bathymetry maps were collected by MBARI
in 2018.

Morphological diversity in seep sediment-hosted nodules

Across both short and long sediment push cores from all four seep areas, we observed distinct

depth intervals of hardened, grey-brown calcitic material ranging in size from <cm rubble to >cm

concretions, or nodules. Representative nodules from the four seep areas are shown in Fig. 2,

with a complete list of the nodule-bearing sediment horizons included in this study summarized

in Table S1. We observed strong morphological differences between the nodules recovered from

cores at SMM 800 and SMM 863. Nodules from SMM 800 (Fig. 2A-C) were rougher, more

irregularly shaped, and featured a greater number of mm to cm-sized voids. These nodules were

also less friable, with lobate, grey-brown, coarse-grained growths often covering black, thin

bivalve shell fragments. These features were consistent between the three seep areas sampled at

SMM 800, as well as down core (7 nodule-bearing horizons total). In contrast, nodules from

SMM 863 (Fig. 2D) were less irregularly shaped, with a less porous, finer grained, brown-

colored material that resembled packed sediment hosting smaller lithics within. This morphology

was also consistently observed between cores and across the five nodule-bearing horizons

sampled at SMM 863.

145
Fig. 2 – Characteristic nodules recovered from each of the four seep areas in this study; Nodule morphology is
remarkably consistent with depth (expressed as centimeters below seafloor, cmbsf); ruler markings are in cm; A)
Nodules recovered from LC62's 36-42-cmbsf horizons at SMM 800-I; B) Nodules from PC44's 6-9-cmbsf horizon
at SMM 800-II; C) Nodules recovered from PC69 (3-12-cmbsf) at SMM 800-III; D) Nodules recovered from the 39-
42-cmbsf horizon of LC66 at SMM 863.

Nodule porosity differences and pore size distributions

Open porosity is defined as the volume of connected pores relative to the volume of bulk solid,

effectively representing the pore space that can freely exchange with the surrounding sediment

environment. Water intrusion analyses of a subset of nodules (Table 1) revealed that the open

porosities of two nodules sampled from SMM 800 (800-I, 800-II) were substantially higher than

that of the nodule from SMM 863.

Table 1

Water intrusion-derived open porosities for select nodules

Seep Area	SMM 800-I	SMM 800-II	SMM 863
Core ID	LC62	PC44	LC66
Core Horizon (cmbsf)	36-42	9-12	39-42
Open Porosity (%)	51.36±0.84	49.52±1.33	43.03±0.78

Hg-porosimetry enabled the measurement of pore size distributions between 6 and 350 μm,

revealing several pore-size regimes within a SMM 863 nodule collected at depth (39-42 cmbsf),

including peaks corresponding to 11, 30, and 300 μm, and a slightly higher proportion of pores

with sizes >100 μm (Fig. 3). A similarly deep-sourced nodule from SMM 800-I had fewer,

although more evenly distributed pore sizes in the observed range. We also saw a smaller peak of

pore diameters at 90 μm and fewer pore diameters above 100 μm compared to the nodule from

SMM 863. A shallower nodule from SMM 800-II (9-12 cmbsf) also had a relatively low, but

even pore-size distribution below 100 μm, in addition to the occurrence of pore sizes between

100 and 250 μm, with minor peaks at 300 and 350 μm.

169
Fig. 3 – Hg-porosimetry curves detailing pore diameter versus incremental intrusion for three nodules sampled from
SMM 800-I, SMM 800-II, and SMM 863. Pore sizes observable with porosimetry range from 6- to 350-μm. The
three nodules shed information on the three distinct nodule-featuring regions identified in this data set: 1) SMM 800
deep sediment (LC62), 2) SMM 800 shallow sediment (PC44), and 3) SMM 863 deep sediment (LC66)

Aragonite dominates nodule carbonate mineralogy, with increased Mg-calcite fractions in

deeper nodules

Bulk powder XRD-derived mineralogies showed SMM 800 nodules were largely composed of

aragonite (>78.2%), with minor contributions from Mg-calcite, calcite, and dolomite (Table 2).

Conversely, nodules from SMM 863 were more aragonite-poor (<43.1%) and featured greater

contributions from Mg-calcite (45.1%, 65.1%) than the nodules from SMM 800. Collectively,

however, we did observe a greater proportion of Mg-calcite in the nodules sourced from deeper

sediment horizons.

Table 2
Key mineral phases for select nodules, as determined by XRD

Seep Area	Core ID	Depth (cmbsf)	Aragonite	Mg-Calcite[a]	Calcite	Dolomite	Quartz
SMM 800-I	LC62	24-27	85.2%	4.5%	0.6%	3.3%	6.4%
		48-51	78.2%	11.3%	3.8%	0.03%	6.7%
SMM 800-II	PC44	9-12	90.2%	2.1%	2%	1.6%	3.3%
SMM 800-III	PC69	3-12	97%	0.1%	1.3%	0.8%	0.9%
SMM 863	LC74	33-39	43.1%	45.1%	0.5%	4.8%	0.5%

	LC66	39-42	26.1%	65.1%	7.1%	1.6%	0.03%

a: Mg-Calcite corresponds to $(Ca_{0.9}Mg_{0.1})O_3$, as measured by (35)

Geochemical localization of the SMTZ and $\delta^{13}C$ trends in porewater DIC and nodules

To characterize the geochemical environments at SMM 800 and SMM 863, we focused on the

vertical trends of three characteristic porewater species that inform the sulfate-methane transition

zone (SMTZ): methane (CH_4), sulfate (SO_4^{2-}), and sulfide (H_2S), shown in Fig. 4. At SMM 800-I

(LC62) collected within a sulfur-oxidizing microbial mat (Fig. S1), porewater extracts revealed

high AOM activity, with a near-complete drop in sulfate concentration by 6-9 cmbsf, which

remained low down core (Fig. 4A). This drop was matched by a concurrent rise in porewater

sulfide and methane, although we observed more dynamic behavior in the sulfide and methane

concentrations with depth, including a notable decrease in both species below their peaks at 6-9

cmbsf. SMM 800-II (PC64) showed a similar sulfate profile, reaching its minimum by 3-4

cmbsf, with values measured at ~3 mM to the base of the short core at 9 cmbsf (Fig. 4B).

Similarly, this drop was accompanied by a steep rise in porewater methane and sulfide, although

as with SMM 800-I, we observed a subsequent drop in concentration below the 3-4 cmbsf

horizon. At SMM 800-III (PC43), also collected in a sulfur-oxidizing microbial mat (Fig. S3),

the peak in AOM activity also occurred at shallower depths (4-5 cmbsf), and, distinct from SMM

800-II, sulfate concentration dropped below detection below this depth (Fig. 4C). At SMM 863

(LC66), only porewater sulfate concentrations were measured. However, compared to the seep

areas at SMM 800, we observed a much deeper penetration of sulfate into the sediment, dropping

below detection by 21-24 cmbsf in this microbial mat habitat, suggestive of a lower methane flux

(Fig. 4D).

Fig. 4 – Sulfate, methane, and sulfide concentration profiles collected for sediment porewaters at the seep areas (A-D) investigated in this study. We plot the median depth in centimeters below seafloor (cmbsf), where concentrations represent 3-cm depth horizons. The sulfate-methane transition zones (SMTZ), as determined by near-complete(<<1mM) sulfate drawdown in each core, are also demonstrated with colored arrows. Colored bars represent the horizons where nodules were recovered for further analysis. Nodules recovered from a different, co-located core at the seep area are also specified. PC64 from SMM 800-II was collected a year previously. We note that there are greater uncertainties with measured sulfide concentrations above the detection limit (>25mM, e.g., in PC43 from SMM 800-III). Only sulfate porewater concentration was collected for the core at SMM 863.

Porewater concentrations and the $\delta^{13}C$ values of dissolved inorganic carbon (DIC, $\delta^{13}C_{pw}$) in addition to the inorganic carbon isotopic concentration of the nodules ($\delta^{13}C_{nod}$) were also measured across the four areas in this study (Fig. 5). These data provide additional information into active AOM horizons where highly [13]C-depleted bicarbonate is produced during methane oxidation and subsequently serves as the source of precipitation of authigenic carbonates (36). At SMM 800-I (LC62), porewater DIC concentrations steeply increased from <10mM to as high as ~42mM in the shallow subsurface (0-6 cmbsf) before stabilizing down core. Interestingly, this occurred in two distinct stages at SMM 800-I, which experienced a second rise in DIC to ~36mM at 15-18 cmbsf to the base of the core at 54 cmbsf (Fig. 5A). At both SMM 800-I and

SMM 800-III (PC43), this shallow spike in DIC concentration was paired with the peak in

negative $\delta^{13}C_{pw}$, reaching values as low as -59.20‰, followed by a steady increase in $\delta^{13}C$ values

down core. This indicated a peak in AOM in the shallow sediment horizons, consistent with the

depth of the sulfate minima. SMM 800-II (PC64) exhibited a different trend, with comparatively

less $\delta^{13}C_{pw}$ (-46.24‰, Fig. 5B). Notably, both DIC concentration and $\delta^{13}C_{pw}$ increased with

depth, perhaps reflective of a mixture of both heterotrophic sulfate reduction and AOM at this

site (37, 38). In comparison, $\delta^{13}C_{nod}$ exhibited a wide range of values across all four seep areas (-

30.09‰ to -53.79‰). Notably, nodules from shallower depths at both SMM 800-I and SMM863

had greater ^{13}C enrichment than the nodules recovered deeper in the cores, where the $\delta^{13}C_{nod}$ at

SMM 800-I was slightly more ^{13}C-depleted (-46.03‰) than its surrounding porewater (-

41.26‰).

235
Fig. 5 – Sediment porewater DIC, $\delta^{13}C_{porewater}$, and $\delta^{13}C_{nodule}$ profiles collected at the seep areas investigated in this
study. We plot the median depth in centimeters below seafloor (cmbsf), where values represent multiple depth
horizons. Because of the <3‰ spread between $\delta^{13}C_{nodule}$ replicates, values are simply averaged for comparison
(individual values are included in Table S2). Colored arrows indicate an estimate for the location of the SMTZ, as in

Fig. 4. PC64 from SMM 800-II was collected a year previously. Only $\delta^{13}C_{nodule}$ was able to be collected for the cores
at SMM 863.

16S rRNA gene comparison of phylum-level diversity in nodules and host sediments

Dominant microbial phyla across all sampled sediment and nodules (Fig. S6-S12) broadly

resembled reports from other carbonate-bearing methane seep environments from Hydrate

Ridge, OR and Eel River Basin, CA, which were often dominated by members of the

Halobacteriota (putative methane-oxidizers largely affiliated with ANME archaea lineages),

Euryarchaeota, and *Desulfobacterota* phyla (14, 16, 22). Additionally, members of the archaeal

Thermoplasmatota, *Asgardarchaeota*, and *Crenarchaeota* were consistently recovered above 1%

relative abundance. Of the bacterial phyla, *Desulfobacterota* were consistently present, with

additional representation by members of the *Chloroflexi* and *Caldatribacteriota* (the latter

exclusively associated with JS1, Table S11), and occasional detection (>1%) of taxa belonging to

the *Planctomycetota*, *Acidobacteriota*, and *Fermentibacterota*. Notably, members from the

Fermentibacterota, first characterized in anaerobic digesters by (39), were only substantially

recovered in three nodules (1.05-2.28% relative abundance) at both the SMM 800-II and SMM

800-III seep areas and rarely recovered (<<1%) within the associated host sediments (Fig. S9,

S11).

Similar 16S rRNA community profiles between sediments and nodules from shallow

sediment horizons

In the shorter push cores from seep areas SMM 800-II and 800-III, *Halobacteriota* lineages

consistently present in seep sediments were ANME-1a, ANME-2b, and ANME-2c (Fig. 6).

Interestingly, these lineages persisted past the shallow zone of sulfate depletion in these cores.

ANME-2c relative abundances were largely consistent with depth through 9-12 cmbsf at both

SMM 800-II and 800-III. However, we observed contrasting patterns in ANME-2b and ANME-

1a abundances. Deeper horizons in SMM 800-II (PC44) showed a drop in the relative abundance

of ANME-2b, concurrent with an increase in ANME-1a below 6 cmbsf. In the cores from SMM

800-III (PC43 and PC69) however, ANME-2b relative abundance and comparatively less

ANME-1a were recovered from the deeper horizons, although this trend was much weaker in

PC69 (Fig 6C). Within the three nodules recovered from these push cores, the dominant

Halobacteriota lineages largely resembled those of their parent sediment horizons, where notable

exceptions include an enrichment of ANME-1a (>20%) in the nodules from PC44 at SMM 800-

II. Still, Shannon-Wiener and Inverse Simpson indices based on archaeal 16S ASV proportional

abundances (Table S3) showed that archaeal diversity was greater in the sediment communities

from SMM 800-II and SMM 800-III than in the nodules.

275
Fig. 6 – Bubble plots summarizing the depth-resolved, 16S relative abundances for groups within *Halobacteriota*
(featuring ANME), *Euryarchaeota (*featuring methanogens, e.g., *Methanofastidiosales*), and *Desulfobacterota*
(featuring sulfate-reducing partners to ANME) in PC64 and PC44 at SMM 800-II (A, B), as well as PC43 and PC69
at SMM 800-III (C, D) . Where possible, ASVs within these phyla were grouped at the genus-level, or at the next
highest taxonomic level possible, and then cut off below a minimum 1% relative abundance. Communities extracted
from nodules are highlighted in grey and often represent larger depth horizons. The location of the sulfate-methane
transition zone (SMTZ) according to porewater sulfate concentrations from the same or co-located sediment core is
indicated with the colored arrows.

Dominant *Desulfobacterota* lineages within these shallow sediment horizons at SMM 800-II and

800-III included two main sulfate-reducing bacterial lineages, Seep-SRB1 and Seep-SRB2, that

have been previously documented to form syntrophic associations with ANME (40–43), where

their respective abundances appeared to covary with those of their ANME partners (Fig. 6BD).

In the sulfate-rich, near-seafloor sediments (0-3 cmbsf) at both seep areas, a large diversity of

other putative sulfate-reducing bacterial groups was also detected above 1%. These SRB lineages

were largely affiliated with groups not known to directly participate in syntrophic AOM. By

comparison, *Desulfatiglans* (not currently known to be involved in AOM) was recovered from

both seep areas at low abundances as deep as 12 cmbsf, including within the nodule from PC69

at SMM 800-III. While uncommon in seep ecosystems, this group has been observed in cold

seep sediments in the Okinawa Trough that also harbor ANME-1 (29). Shannon-Wiener and

Inverse Simpson indices based on bacterial 16S ASV proportional abundances (Table S3)

indicated that the sediments surrounding the nodules from these seep areas had higher diversity,

as found with the Archaea.

Deeper sediment and nodule 16S rRNA surveys show persistence of *Halobacteriota* and

***Desulfobacterota* and putative methanogenic lineages**

Similarly to SMM 800-II and 800-III, SMM 800-I (LC62) had a shallow SMTZ (sulfate depleted

by 5-6 cmbsf) and both sediments and nodules were enriched in *Halobacteriota*, *Euryarchaeota*,

and *Desulfobacterota* lineages within and well below the sulfate depletion depth. Archaeal

diversity was largely dominated by ANME-1a (up to 66.5% relative abundance) which was

recovered throughout the core at depths below 3 cmbsf. In the deeper horizons, this group was

accompanied by ANME-1b at lower relative abundance (Fig. 7). Notably, within this long core,

we also documented two distinct transitions in both archaeal and bacterial community

composition, which occurred between 4-27 cmbsf and between 33-57 cmbsf, representing the

base of the core.

310
Fig. 7 – Bubble plots summarizing the depth-resolved, 16S relative abundances for groups within *Halobacteriota*,
Euryarchaeota, and *Desulfobacterota* in LC62 at SMM 800-I. Where possible, ASVs within these phyla were
grouped at the genus-level, or at the next highest taxonomic level possible, and then cut off below a minimum 1%
relative abundance. Communities extracted from nodules are highlighted in grey and often represent larger depth
horizons. The location of the sulfate-methane transition zone (SMTZ) according to porewater sulfate concentrations
from the same or co-located sediment core is indicated with the colored arrows. A) Relative abundances within
prominent phyla *Halobacteriota* (featuring ANME) and *Euryarchaeota* (featuring methanogens such as
Methanofastidiosales). B) Relative abundances within *Desulfobacterota* – the phylum with sulfate-reducing partners
to ANME.

The first transition showed a shift from the consistent presence of ANME-2c (1-10% relative

abundance), an unclassified ANME-1 lineage, and the putative syntrophic sulfate-reducing

bacterial lineages Seep-SRB1 and Seep-SRB2 in the shallow sediment down to 27 cmbsf to an

underlying zone ~24 cm thick where each of these groups, with the exception of sporadic

occurrence of ANME-2c, were below detection (Fig. 7). The second transition, occurring

between 33-57 cmbsf (well below the SMTZ), was marked by the emergence of ANME-2b and a

putative methylotrophic methanogen lineage *Methanofastidiosales* (formerly the WSA2 group,

<20% relative abundance) largely dominated by a singular ASV (Table S11). The

Methanofastidiosales are a poorly characterized group originally described from anaerobic

sludge digesters (44) and observed in hydrothermal systems, gas hydrates, methane seeps, and

methane-rich coastal sediments (29, 32, 45–48). This ASV was also detected above 1%

abundance in the shallower push core sediments from SMM 800-II (PC464, PC44) and 800-III

(PC69), including a nodule in PC69 (Fig. 6AC). A similar transition was observed for

Desulfobacterota lineages, where the dominance of ANME partners Seep-SRB1 and Seep-SRB2

was succeeded by the persistent, though minor (<10%) presence of *Desulfatiglans* (Fig. 7B).

Additional transitions in the archaeal and bacterial communities occurred in the uppermost 3 cm

and deepest horizons of the core below 51 cmbsf, with the near seafloor sediments largely devoid

of ANME lineages, but supporting a high diversity of *Desulfobacterota* not recovered in deeper

horizons, including Seep-SRB4 and Desulfobulbus lineages, while the deepest horizons (51-54

cmbsf) showed an increase in *Halobacteriota* diversity, with the reappearance of members of the

ANME-2c and detection of ANME-3, alongside the reoccurrence of Seep-SRB1. Compared to

ANME-2c ASVs, the dominant Seep-SRB1 ASVs in the deepest core interval were different

from the dominant ASVs in the intermediate depth horizons (3-27 cmbsf), although we observed

a few shared Seep-SRB1 ASVs between these regions (Table S11).

Direct comparisons of the carbonate nodules (n=4) with paired sediments across these transitions

revealed strong similarities between representative archaeal and bacterial taxa, including major

members within *Halobacteriota* and *Desulfobacterota*, down to the ASV level (Fig. 7, Table

S11). Beyond these taxa, both Shannon-Wiener and Inverse Simpson indices suggested the

sampled nodule bacterial communities were consistently more diverse than their surrounding

sediments, though this trend did not extend to the archaeal communities (Table S3).

The long cores recovered from SMM 863 revealed a slightly different community structure from

SMM 800-I. Across both SMM 863 cores (LC74, LC66), we observed a clear transition in the

dominant ANME and putative sulfate-reducing bacterial groups around 21-24 cmbsf (Fig. 8),

aligned with the approximate depth of the SMTZ at this site. Here, members of the

Halobacteriota above this transition zone were dominated by ANME-2c and known syntrophic

partners belonging to the Seep-SRB1 as well as other uncharacterized lineages within the

Desulfosarcinaceae. We also identified ANME-3 at varying abundances in the upper (<12

cmbsf) sediment in both cores. Below the transition at 21-24 cmbsf, ANME-2c was succeeded

by lineages within ANME-1, primarily ANME-1b (up to 75% relative abundance). This

transition corresponded with the appearance of the Seep-SRB2 lineage, previously observed to

partner with ANME-1 (43, 49, 50). This depth distribution pattern was distinct from the other

main ANME syntrophic partner, Seep-SRB1, which was consistently detected above 1%

throughout both cores (Fig. 8B). Interestingly, we observed another transition in both cores,

beginning at 36-39 cmbsf in LC74 and at 42-45 cmbsf in LC66, where ANME-2c reappeared

alongside ANME-3. Here, the lower abundance of ANME-2c at depth was represented by the

same dominant ANME-2c ASVs recovered in the upper sediment, while the ANME-3 ASVs at

depth largely differed from those recovered near the top of the core (Table S11). Nodules

sampled from this transition (n=3) revealed similar ANME-1b and ANME-1a lineages as in the

surrounding sediments, down to the ASV level (Fig. 8A, Table S11), but lacked consistent

representation from the other minor ANME lineages, including the uncharacterized ANME-1

group (in LC66) and ANME-2c (in both LC66 and LC74). Similarities were notably less

apparent among the *Desulfobacterota*, where lineages present in the host sediment were often

absent from the nodules (i.e., Seep-SRB1 and Seep-SRB2) and lineages not observed in the

immediate sediment horizon were observed in the hosted nodule (e.g., *Desulfatiglans* in LC66).

374
Fig. 8 – Bubble plots summarizing the depth-resolved, 16S relative abundances for groups within *Halobacteriota*,
Euryarchaeota, and *Desulfobacterota* in LC74 and LC66 at SMM 863. Where possible, ASVs within these phyla
were grouped at the genus-level, or at the next highest taxonomic level possible, and then cut off below a minimum
1% relative abundance. Communities extracted from nodules are highlighted in grey and often represent larger depth
horizons. The location of the sulfate-methane transition zone (SMTZ) according to porewater sulfate concentrations
from the same or co-located sediment core is indicated with the colored arrows. A) Relative abundances within
prominent phyla *Halobacteriota* (featuring ANME) and *Euryarchaeota* (featuring methanogens such as
Methanofastidiosales). B) Relative abundances within *Desulfobacterota* – the phylum with sulfate-reducing partners
to ANME.

Cell abundance and fluorescence *in situ* hybridization (FISH) in nodules and sediments

Quantification of SYBR Gold®-stained cells and cell aggregates (cell associations > 5 μm)

extracts were consistently higher (often 1-2 orders of magnitude) in sediment compared with

nodules across the sites, with typical single cell abundances ranging between 10^6-10^7 cells mg^{-1}

in sediments compared with 10^5-10^6 cells mg^{-1} in nodules (Table 3). While differences in cell

abundance between sediments and nodules were observed, a substantial decrease in cell

abundance between the deeper sediment horizons (e.g., 2.7×10^7 at 54-57 cmbsf) compared with

shallower depths (e.g., 3.59×10^7 at 9-12 cmbsf) was not detected.

Table 3
Aggregate/single-cell/cell clump SYBR Gold counts in nodules and paired sediment horizons

Seep Area	Core ID	Core Horizon (cmbsf)	Avg Single Cells per mg Sample	SE	Avg Aggregates per mg Sample	SE
SMM 800-I	LC62	36-39	9.53×10^6	3.69×10^5	8.00×10^3	1.58×10^3
		39-42	1.30×10^7	4.05×10^5	1.02×10^4	1.60×10^3
		(Nodule) 36-42	3.89×10^5	1.86×10^4	5.65×10^2	1.73×10^2
		51-54	2.98×10^7	2.36×10^6	4.63×10^4	6.63×10^3
		54-57	2.70×10^7	2.18×10^6	3.11×10^4	1.89×10^3
		(Nodule) 51-57	5.11×10^5	1.73×10^4	2.56×10^3	2.78×10^2
SMM 800-II	PC44	6-9	4.05×10^7	2.10×10^6	8.28×10^4	2.17×10^4
		(Nodule) 6-9	4.21×10^5	1.54×10^4	4.63×10^3	7.19×10^2
		9-12	3.59×10^7	2.01×10^6	2.88×10^4	8.87×10^3
		(Nodule) 9-12	1.15×10^6	7.56×10^4	2.89×10^3	4.11×10^2
SMM 800-III	PC69	3-6	5.50×10^7	3.78×10^6	3.62×10^4	9.54×10^3
		6-9	4.42×10^7	2.64×10^6	4.14×10^4	1.61×10^4
		9-12	3.97×10^7	1.84×10^6	3.63×10^4	1.24×10^4
		(Nodule) 3-12	1.09×10^6	6.52×10^4	1.77×10^3	2.38×10^2
SMM863	LC66	42-45	3.01×10^7	9.51×105	2.64×10^4	3.02×10^3
		(Nodule) 42-45	1.70×10^6	8.68×10^4	4.01×103	9.20×10^2
	LC74	42-45	7.45×10^6	4.70×10^5	3.16×10^4	4.11×10^3
		(Nodule) 42-45	1.60×10^6	1.54×10^5	1.08×10^4	1.51×10^3

The spatial associations in aggregates and morphology of Bacteria, Archaea, and ANME-1 cells

were also examined within the recovered nodules by fluorescence *in situ* hybridization (FISH);

(Fig. 9). The persistence of highly auto-fluorescent material within the long cores from seep

areas SMM 800-I and SMM 863 made it difficult to clearly discern a positive fluorescence signal

from FISH, however we were able to observe positive hybridization with the ANME-1 probe,

supporting 16S rRNA results. Notably, we observed large aggregates with morphology

consistent with ANME-SRB consortia reported from seep sediments at a higher frequency within

nodules recovered from the shallower sediment horizons (< 9-12 cmbsf) at seep areas SMM

800-II and 800-III, compared to the nodules recovered from deeper sediment intervals at SMM

800-I and SMM 863, where the FISH signal from cells was dimmer and cell aggregates were

more loosely bound in an exopolymeric matrix (Fig. 9FG, Fig. S14).

406
407   **Fig. 9** – Representative examples of carbonate nodule associated cells and aggregates recovered from SMM 800-II,
408   800-III, and 863, visualized by fluorescence in situ hybridization (FISH) using probes targeting *Bacteria* (EUB338
409   I-III, in green), ANME-1 (ANME1-350, ANME1-728; in red), and *Archaea* (ARCH915, in pink). Cell extracts for
410   nodules from SMM 800-I were not shown due to strong interference from auto-fluorescent exopolymeric material.

Single-cell translational activity measurements (BONCAT) show large shift in nodule endolithic activity between shallow sediments and deeper horizons below the SMTZ

A subset of nodules (n=7) from 5 sediment cores were separated from their host sediment and incubated with the methionine analog HPG under sulfate- and methane-rich conditions to examine translational activity by endolithic microorganisms using biorthogonal noncanonical amino acid tagging (BONCAT), e.g., (49, 51, 52). Nodules were recovered after a 14-week incubation at 4ºC, and BONCAT analysis of carbonate-associated cells showed high heterogeneity in activity for both single cells and cell aggregates between nodules, ranging from less than 1% up to 39% of the recovered cells. The relative fraction of extracted cells and cell aggregates (>5 µm) with a confirmed BONCAT signal compared to those without is provided in Table 4. Representative examples of BONCAT stained cells and aggregates from each of the seep areas are shown in Fig. 10.

Table 4
Proportion of BONCAT-active cells to non-BONCAT-active cells from incubated nodules

Seep Area	Core ID	Nodule Horizons (cmbsf)	BONCAT-Active Single Cells	BONCAT-Active Aggregates
SMM 800-I	LC62	36-42	<1% (n = 765)	*no aggregates observed*
		51-57	<1% (n = 767)	0% (n=1)
SMM 800-II	PC44	6-9	39% (n = 792)	60% (n = 10)
		9-12	14% (n = 1011)	40% (n = 5)
SMM 800-III	PC69	3-12	4% (n = 1041)	62% (n = 21)
SMM863	LC66	42-45	<1% (n = 1062)	*no aggregates observed*
	LC74	42-45	7% (n = 857)	100% (n = 2)

Fig. 10 – Cell extracts from recovered carbonate nodules at all four seep areas, displaying the presence/absence of newly synthesized proteins with BONCAT (in red) after a 14-week incubation with methionine analog, L-Homopropargylglycine (HPG). DAPI staining of nucleic acids indicating cells is in cyan. Positive BONCAT signal was consistently observed in all extracts from the shallower nodules at SMM 800-II and 800-III, compared to the consistent lack of signal at SMM 800-I and 863.

Nodule-hosted cells recovered from SMM 800-I (LC62, 36-42 and 51-57 cmbsf, both well below the SMTZ), did not exhibit evidence of HPG assimilation among the >700 cells examined per sample. Additionally, these nodules lacked cell aggregates commonly found in shallower sediment horizons. The absence of a BONCAT signal aligns with the minimal sulfide production observed prior to the HPG incubation period (Fig. S15). We observed similar results with a deep nodule recovered from SMM863 below the SMTZ (LC66, 42-45 cmbsf). However, the nodule recovered from the parallel core (LC74) from the same depth horizon did show a weak BONCAT signal (7% of total cells screened) in single cells and aggregates, despite lack of observed sulfide production prior to HPG incubation (Fig. S15). This signal indicated that these endolithic habitats below the SMTZ continued to support viable microorganisms (Table 4).

The low microbial activity and lack of cell aggregates observed in the majority of nodule

incubations from the deeper sediment intervals contrasted with the nodules recovered from the

shallow horizons within the SMTZ (between 3-12 cm) at SMM 800-II and 800-III. Here, we

observed active HPG assimilation for both single cells and cell aggregates, with up to 39% of the

single cells (n=792) and 62% of aggregates (n=21) showing a positive BONCAT signal (Table

4). FISH-based identification of BONCAT positive cells indicated a large fraction of these active

cells were Archaea (ARCH915 probe, Fig. S16). These samples actively produced sulfide prior

to the incubation, reaching values as high as 8 mM after 30 days (Fig. S15), supporting their

likely involvement in sulfate-coupled methane oxidation.

Discussion

In this study, increased depth sampling directly within active seeps using ROV-deployed long

cores provided a window into microbial community transitions within and below the SMTZ. We

sampled seven sediment cores containing carbonate nodules directly within microbial mat-

associated, active seep habitats from four seep areas associated with two carbonate mounds in

the Santa Monica Basin. The three seep areas at SMM 800 showed consistent geochemical

profiles with high, sulfate-coupled AOM activity characterized by a shallow (<9 cmbsf), sharply

defined SMTZ, whereas the two long cores from SMM 863 instead revealed a much deeper (~21

cmbsf) sulfate depletion zone suggestive of lower methane flux and AOM activity (21).

Carbonate nodules recovered from all four seep areas across these distinct geochemical zones

enabled us to investigate the physical properties, along with the diversity and metabolic activity

of nodule-hosted microbial communities relative to their host sediments, with implications for

the degrees to which these incipient carbonate nodules preserve records of past AOM and

continue to serve as habitats for *in situ* endolithic communities.

Closely-linked sediment and nodule-hosted microbial communities indicate ongoing

exchange throughout and below the SMTZ

16S rRNA-based community profiles from recovered nodules at all the characterized seep areas

reinforced a close similarity to those of their host sediments, regardless of depth and

geochemical regime. Non-metric multidimensional scaling (NMDS) ordination of all sediment

and nodule-derived ASVs (Fig. S13) also showed a consistent, area-specific clustering between

nodules and sediments. This coupling generally agrees with previous studies comparing various

carbonate substrates, including nodules, from seep sites at Hydrate Ridge, OR; Eel River Basin,

CA; and the Sea of Japan (14–16, 22).

Collectively, these shared microbial community profiles across major geochemical transition

zones are strong evidence of continuous colonization and response to *in situ* conditions which

collectively influence nodules and sediments, rather than a time-integrated, microbial

'thumbprint' of the surrounding sediment.

This is further supported by the high open porosities measured for the nodules in this study,

regardless of depth and pore size distributions (< 350 µm), which likely facilitate continuous

colonization by surrounding seep sediments. Indeed, 13-month seafloor colonization experiments

of sterilized carbonate substrates within and on the periphery of deep-sea methane seeps

previously demonstrated active colonization by seep sediment microorganisms, including

ANME-1 (16). Although the mechanisms of potential exchange between nodules and sediments

remain unknown — especially since motility has not been definitively observed in ANME —

previous studies have hypothesized the involvement of fluid advection and/or methane ebullition

in facilitating active carbonate colonization (14, 16, 53, 54).

Further, the high nodule open porosities support sufficient connectivity for methane and sulfate

perfusion, along with the diffusion of small organic molecules (e.g., HPG, used in BONCAT

experiments). As such, these nodules likely experience very similar, current geochemical

pressures as the surrounding sediments, which supports the shared microbial community

transitions observed in both sediments and nodules from SMM 800 and SMM 863. Indeed,

subsurface methane fluxes can vary over periods that can last from days to centuries (55–57),

where changes in geochemical stratification exert an active role in structuring the sediment

microbial communities performing AOM and facilitating authigenic carbonate precipitation (1,

14, 36, 58–60).

Isotopic and mineral signatures from the nodules recovered below the SMTZ also record

carbonate precipitation in different geochemical regimes from the present environments

observed. Despite being a sulfate-depleted environment, the consistently depleted $\delta^{13}C$ values for

these nodules — below that of marine organic carbon, -25‰ (61) — support at least a partly

AOM-driven origin, and thus preclude exclusive precipitation under the current geochemical

conditions. Indeed, all the nodules at SMM 800-I were aragonite-rich (78.2-85.2%), suggestive

of a history largely informed by shallower, sulfate-bearing sediments (62, 63). By contrast, the

greater Mg-calcite content (45.1-65.1%) in the nodules from SMM 863 support a longer-lived

interval in a deeper, more sulfate-limited AOM environment (6, 59, 64, 65). Together, these data

suggest that the nodules recovered at depth are likely to have experienced microbial community

shifts to reflect communities better adapted to *in situ* conditions, which agrees with a model of

continued colonization from the surrounding sediments.

Persistence of ANME lineages suggest potential adaptation or survival under sulfate-poor conditions below SMTZ

Long cores collected from both SMM 800-I and SMM 863 revealed the consistent dominance of ANME-1 lineages over ANME-2 in the deeper sulfate-depleted sediments and nodules, consistent with several previous studies that have noted this general pattern in seep ecosystems worldwide (24, 26, 28, 30–32, 66, 67). The occurrence of ANME-1 lineages well below the SMTZ, frequently representing the same ASVs as those found within zones of high AOM activity, may reflect thus greater physiological plasticity among this archaeal lineage.

Interestingly, the difference in community profiles below the SMTZ between SMM 800-I (LC62) and SMM 863 (LC74, LC66) attests to additional environmental factors driving niche differentiation in seep sediments and nodules in this geochemical zone. While the dominance of ANME-1b below the SMTZ at SMM 863 may be linked in part to AOM with known ANME partners Seep-SRB1 and Seep-SRB2, the low abundance (<1%) of *Desulfobacterota* sequences below the SMTZ at SMM 800-I suggest that the ANME-1a cells at depth may persist without a sulfate-reducing partner. FISH data from several environments have documented the occurrence of ANME-1 as single cells without an attached bacterial partner (23, 66, 68–71), further supportive of ANME-1's metabolic flexibility beyond sulfate-coupled AOM.

Notably, the occurrence of putative methylotrophic methanogens within the *Methanofastidiosales* at depth supports the transition to a methanogenic regime below the SMTZ, where ANME-1a and *Caldatribacteriota* (exclusively JS1, Table S11) were found to co-occur. JS1 has been previously identified in anaerobic and/or methanogenic seep sediments worldwide (16, 26, 72–74), and hypothesized to participate in hydrocarbon degradation that may supply key substrates for methanogenesis (44, 73–75). Indeed, it has also been hypothesized that

at least some members of the ANME-1 may be capable of methanogenesis (31, 70, 76–80).

Although porewater methane concentrations do not conclusively support methane production,

previous methane measurements at SMM 800 (33) do support a biological origin for methane,

based on the purity and isotopic signature (-70.8‰ δ^{13}C). Similarly, the observed rise in

porewater DIC and associated δ^{13}C in the deeper sediment horizons at SMM 800-I (LC62) may

be explained by ^{13}C-enriched DIC produced during methanogenesis (81, 82). Enriched ^{13}C

content associated with an increased dolomite fraction in a deeper nodule from SMM 800-I

(LC62, 24-27 cmbsf) also agrees with carbonate formation in historically methanogenic sediment

regimes below the SMTZ (6, 64) — a phenomenon additionally observed for a sub-SMTZ

nodule recovered from SMM 863 (LC74, 33-39 cmbsf).

Beyond methanogenesis, the observed co-occurrences of ANME-1 lineages,

Methanofastidiosales, and JS1 may also attest to Fe-coupled AOM below the SMTZ at SMM

800-I. Fe-coupled AOM has been observed in sulfate-depleted marine sediments (29, 83–91),

where explicit associations between ANME-1a/1b, JS1, and *Methanofastidiosales* have been

statistically supported within seep sediments likely hosting Fe-AOM (29). Further, the detection

of ANME-3 in the lower (51-57 cmbsf) horizons of core LC62 at SMM 800-I aligns with

observations of ANME-3 in sulfate-depleted marine sediments and microcosms supporting Fe-

AOM (83, 85). Ultimately, lack of dissolved iron concentrations for the sediment or discernible

iron mineral content in the nodules precludes geochemical support for Fe-AOM at SMM 800-I.

Still, past observations of this metabolic regime and its connection to frequently observed

microbial taxa in globally-distributed seep sediments below the SMTZ — including sediments

and nodules at the seep areas in this study — suggest that the extent of AOM below the SMTZ at

active seeps may be under-constrained.

More broadly, the co-occurrence of both ANME/SRB and methanogenic lineages at SMM 800-I

may also suggest *in situ* mixing of AOM and methanogenic regimes. These regimes have been

hypothesized to co-occur in marine sediments (92) and have been reported for sediments in the

Bothnian Sea (87). Significant correlations between ANME clades and methanogen groups from

sulfate-depleted sediments at the Okinawa Trough have also suggested a degree of cryptic

methane cycling (29). Collectively, these studies underscore the complexity of resolving distinct

metabolic regimes in seep sediments.

Interestingly, the weaker-to-negligible BONCAT signal in incubated nodules from below the

SMTZ at SMM 800-I and SMM 863 under sulfate and methane-replete conditions indicates that

substrate concentrations alone may be insufficient to stimulate measurable AOM activity in

nodules at depth over the time period of the incubation (3.5 months). This was despite recovering

high cell abundances that included both single cells and aggregate morphologies consistent with

ANME (including FISH-detectable ANME-1) in the deeper nodules. Together, these data

suggest that measuring metabolic activity for incubated nodules recovered at depth may require

longer observation periods, or more sensitive measurements.

In this regard, the minimal activity observed for carbonate nodules below the SMTZ may be

compared to that of seafloor carbonates in low-activity seep areas. Indeed, *in situ* experiments

transplanting seafloor-exposed carbonates from low to high activity seepage areas by (16)

reported shifts towards microbial community structures enriched in lineages involved in AOM

and sulfur-based metabolisms after 13 months. Further, sulfide measurements from incubated

low activity seafloor carbonates in (11) showed that reactivation of AOM in hosted endolithic

communities could take months to years. By contrast, previous $^{14}CH_4$ radiotracer measurements

of methane oxidation rates from low-activity seep carbonates, incubated under similar

conditions, showed detectable, if low methane oxidation (<200 nmol after 7 days) after the

introduction of AOM-favorable conditions (8).

Collectively, these data may also point to the understudied nature of metabolic dormancy in seep

microbial communities. Previous studies have diverged on interpretation of gene-based profiles

of seep sediments and buried seep carbonates, where relict, or 'fossil' DNA has been proposed to

persist in marine sediments (93–95) and cited in buried seep carbonate crusts (58, 59, 90), such

that community profiles are not representative of metabolically viable cells. However, recent

studies have suggested that 'fossil' DNA negligibly affects taxonomic surveys of environmental

samples, especially in marine sediments below the bioturbation zone and even across systems

where physical protection of DNA (e.g., seep carbonates) may generate bias (96, 97). As such,

we consider 16S rRNA-based nodule community profiles to largely reflect the intact cell

assemblage observed in nodules from all depths, with clear variation in metabolic activity with

increasing depth below the SMTZ, representing a mixture of dormant and active microorganisms

within this endolithic habitat.

Future investigations at methane seeps should therefore seek to better characterize the role of

AOM in both the sediment and nodules below the SMTZ, which remains poorly understood.

Longer (>1 year), lab-based studies examining the dynamics and range of metabolisms leveraged

in these methane-rich, sulfate-poor conditions may shed additional information on factors driving

metabolic activity in low-energy seep environments and how sensitive these communities are to

the types of geochemical changes characteristic of transient methane seep activity.

Mechanisms, timescales for sediment community entrapment or exchange in nodules

remain underexplored

Although continued exposure and exchange between seep nodules and sediments is strongly

supported by the observations presented in this study, the processes or conditions that may act to

decouple the nodule environment from its surroundings remain to be more clearly defined.

Minor offsets observed between sediment and nodule communities in NMDS ordination of all

sediment and nodule-derived ASVs (Fig. S13) may reflect subtle substrate-level differences in

community structure. Further, we observed higher community richness in shallow sediments

compared to nodules at SMM 800-II and 800-III (Table S3). This trend has also been observed

previously in methane seeps at Hydrate Ride, OR (14, 22), and may suggest a local environment

more favorable to certain microbial taxa. Indeed, lower cell and cell aggregate abundances in the

nodules from SMM 800 and SMM 863 relative to the surrounding sediments may result from

niche-specific differences and/or likelihood of some barriers to free exchange between sediments

and porous carbonate nodules.

Minor discrepancies between sediment-nodule pairs have also been hypothesized to originate

from translocation events that disrupt the original emplacement of incipient seep carbonates.

These translocation events act on a local basis and may be driven by sudden gas hydrate

decomposition or ebullition events (14). Still, the prevalence of this phenomenon is not known,

as the formation and depositional history of seep nodules remains poorly constrained. Sediment-

hosted seep carbonates are thought to be continuously subject to burial by continuous sediment

transport from nearby coasts, uplift from tectonic movements or gas hydrate accumulation (6, 33,

64), and dissolution from nearby sulfide oxidation and/or aerobic methane oxidation (98–103). In

addition to precipitation dynamics, these forces are all expected to shape the duration and nature

of exchange between sediments and hosted carbonate nodules in ways that outline future paths of

investigation. Although the ages of the nodules in this study remain unconstrained (though

loosely estimated at 10^2-10^3 years, (33)), the data presented in this study suggest that over the

interval since their formation, diagenetic forces have not acted to significantly decouple sediment

and nodule microbial communities.

Additional sampling of seep sediment-hosted nodules should seek to more clearly define the

time-constrained formation and diagenetic history captured by these understudied carbonates

relative to their surrounding sediments, especially in instances where nodule-hosted microbial

communities and surrounding sediment communities differ. Uranium-thorium (U-Th) dating,

combined with $\delta^{13}C$ and X-ray diffraction (XRD) characterizations of these sediment-hosted

carbonates can better address their age, provenance, and subsequent diagenetic history. In turn,

these studies can advance our understanding of controls informing continued microbial

habitation in seep sediment-hosted nodules.

637   Conclusions

In this study, we integrated microbiological and geochemical analyses of paired seep carbonate

nodules and host sediments spanning the SMTZ and below at two previously described seep sites

in the Santa Monica Basin, CA. We extend prior characterizations of seep sediment-hosted

nodules with this integrated analysis, mapping changes in nodule-hosted microbial community

structure and activity of these endolithic microorganisms across steep geochemical and

microbiological transitions. By sampling within and below the SMTZ, we captured similar

community transitions in both nodules and sediments, suggestive of continued perfusion and

exchange between nodules and surrounding sediments, regardless of geochemical conditions.

Porosity and mineralogy measurements also indicated sediment-nodule connectivity was feasible

and had likely persisted throughout the nodule's formation, regardless of formation environment.

Unique community structures beneath the SMTZ also attested to different metabolic regimes

captured in both sediments and nodules, with the potential for co-occurrence of methanogenesis

and AOM. Still, BONCAT activity measurements under sulfate/methane-rich conditions showed

that the deeper nodules collected below the SMTZ were slow or no longer capable of producing

measurable evidence of AOM within the 14-week incubation period compared to nodules from

near-seabed horizons supporting active, sulfate-coupled AOM. Future investigations should

explore the range of AOM and non-AOM metabolisms feasible in seep sediment ecosystems in

wider detail and seek to more accurately constrain the range of diagenetic histories captured by

seep nodules to better constrain the past and current degree of AOM activity recorded in these

endolithic habitats.

Materials and Methods

Sample collection

Sediment cores were collected with the manipulator arm using the *ROV Doc Ricketts* during

dives DR1329 and DR1333 during the WF05-21 Southern California cruise on the *R/V Western*

Flyer. Additionally, we included data from one core (PC64) collected on dive DR1248 by ROV

Doc Ricketts at SMM 800 in February 2020 during the WF02-20 Southern California cruise on

the *R/V Western Flyer*. Additional information on site contexts and additional sample recovery

details may be found in the Supplemental Material. A further detailed summary of all the

sediment and nodules in the study is also included in Table S10.

Upon recovery, all cores were kept in a 4°C walk-in cooler until processing. Within hours of

collection, sediment cores were documented and sectioned shipboard into 1- or 3-cm horizons by

upward extrusion. Nodules were manually identified during core sectioning and kept within the

associated sediment horizons. These horizons were then bagged in sterile Whirl-Pak bags, sealed

in Ar-flushed Mylar bags (IMPAK Corporation), and maintained in cold storage at 4°C for, and

during, transport to the laboratory.

In the laboratory (<2 weeks after sample collection), nodules >1cm in diameter were manually

extracted from the bagged, anoxic sediment horizons under N_2 and rinsed three times in 0.22 μm-

filtered, 3X phosphate-buffered saline (PBS, as in (104)) to remove any adhered sediment. After

rinsing, nodules from each horizon were then sub-sampled for various physical, geochemical,

microbiological analyses (further detailed by section titled in the Supplemental Material). Due to

limited sample available for multiple analyses, some nodules were combined across adjacent

sediment horizons and treated as the representation of a larger segment of the sediment core

collected (Table S10). We conducted a second cleaning step for the rinsed nodules used for

DNA-based community analysis and visualization by fluorescence microscopy to minimize

sediment contamination following the optimized protocol for DNA extraction from carbonate

nodules described in (14). In (14), rigorous testing for sufficient removal of surface

contamination from the sediment hosted nodules was tested using *E. coli* as a tracer, with the

following protocol demonstrating complete removal of DNA from the exogenously introduced

microbes. Briefly, this surface treatment included additional rinses with cold, 0.22 μm-filtered

3X PBS, sonication, and centrifugation (additional details are provided in the Supplemental

Material). Rinsed nodules designated for activity measurements with bioorthogonal non-

canonical amino acid tagging (BONCAT) (51) were immediately transferred to separate, sterile,

acid-washed media bottles, submerged in cold, 0.22 μm-filtered, N_2-sparged artificial seawater

(media composition available in Supplemental Material Table 1.10), and stoppered with an acid-

treated butyl rubber stopper (Th. Geyer GmbH & Co. KG). After immersion, bottle headspaces

were flushed with pure CH_4 for 1 minute with a needle and then pressurized to 2 bar. Samples

were incubated in the dark at 4°C.

Methane measurements

On both the WF05-21 and WF0-20 cruises, sediment methane samples were collected by 1 mL

cut off syringe from each sediment core horizon and immediately transferred into 3 mL of 5M

NaOH in a pre-weighed gas-tight 10 mL Restek vial, shaken to mix, and then stored upside down

at room temperature until analysis. In the laboratory, methane concentrations in the headspace of

the vials were measured with a Hewlett Packard (HP) 5890 Series II Plus Gas Chromatography

system with a flame ionization detector, connected to a HP 7694 Headspace Sampler with a 1

mL sample loop. The GC column was a 30 m HP-Plot/Q column with inner diameter (I.D.) of

0.32 mm. The software used for analysis was Agilent GC ChemStation (version A.09.03 [1417]).

A calibration series ranging from 250 ppm (v/v) to 5000 ppm (v/v) was run immediately prior to

samples using lab grade CH_4 gas (100%) injected into sealed 10 mL Restek vials using a gas-

tight syringe. Methane volumetric ppm values were converted to μmol assuming ideal gas

behavior at standard temperature and pressure. These values were normalized relative to the

mass of sediment added to the Restek vials during the cruise, where masses were corrected for

porosity by weighing sediment from each horizon before and after drying in an oven.

Sulfate measurements

On both cruises, sediment porewater was collected in pressure-driven 'squeezers' (105), using

argon to compress the sediment section against a 47 mm, 0.22 μm polycarbonate filter (Thermo

Fisher) to recover sediment-free porewater in gas-tight, 60 mL syringes while minimizing

exposure to air. Porewater from each sediment horizon was then filtered through a 0.22 μm

polyethersulfone (PES) syringe filter (Tisch) into 2 mL Eppendorf tubes and stored at -20°C for

later analysis. In the laboratory, the concentration of major ion species (e.g. sulfate) was

determined by ion chromatography at the Resnick Sustainability Institute's Water and

Environment Lab at the California Institute of Technology. Specifically, samples were run on a

250 mm Dionex IonPac instrument (Thermo Fisher) with AS19-4μm and CS16-4μm columns for

anions and cations, respectively — each with 50 mm versions as guard columns. 100 μL of each

porewater sample were diluted 50x in milliQ ultrapure DI water (final volume 5 mL) before

running. Standard curves were generated using calibration standards diluted similarly to samples,

with 100 μL of 500 mM NaCl added to mimic typical seawater sample peak behavior. Analyte

peaks were automatically integrated and major ion species, including sulfate, were automatically

calculated using the prepared standards with the Chromeleon Dionex IC (Thermo Fisher, v7.2.9)

software, with manual inspection to ensure regular peak shapes.

Sulfide measurements

Samples for porewater sulfide concentrations were prepared by the addition of 0.5mL of 0.22 μm

filtered porewater from each sediment horizon to 0.5 mL of 0.5M zinc acetate in a 1.5 mL

Eppendorf tube to precipitate zinc sulfide. Due to its lability, sulfide samples were processed first

after porewater collection from the squeezer apparatus. In the laboratory, the precipitated sulfide

was quantified spectrophotometrically according to the Cline sulfide assay protocol (106).

Briefly, this involved the acidification of the sample and reaction with N, N-dimethyl-p-

phenylenediamine dihydrochloride in presence of ferric chloride. The resulting methylene blue

solution's absorbance was measured in duplicate at 670-nm wavelength with a microplate reader

(Sunrise, Tecan). Concentrations were calculated after subtracting absorbance values from DI

H_2O blanks from concurrently measured zinc sulfide standards prepared from sodium sulfide

solutions (0.12 mM to 25 mM). We additionally measured 10x and 50x dilutions to allow for

quantification of highly sulfidic samples within the linear color range.

DIC and $\delta^{13}C$ analyses

We measured dissolved inorganic carbon (DIC) concentrations and its $\delta^{13}C$ in porewater (C_{pw}),

as well as the $\delta^{13}C$ of nodule (C_{nod}) where sufficient porewater or nodule material allowed. After

extraction and rinsing, nodule samples were air dried and ground to a fine powder with an agate

mortar and pestle, which was cleaned with DI water and ethanol between samples. Replicate

subsamples of powdered nodules (0.45-0.65-mg) were added to 12 mL, acid-washed and

combusted Exetainer vials (Labco). The vials were then flushed with helium for 2 minutes and

acidified with 200 μL 43.5% phosphoric acid. To measure porewater DIC and $\delta^{13}C_{pw}$, 1 mL of

extracted porewater was immediately added via syringe to 12 mL, acid-washed and combusted

Exetainer vials (Labco) pre-flushed with helium and pre-loaded with 100 μL 43.5% phosphoric

acid during the WF05-21 and WF02-20 cruises. In the laboratory, the released CO_2 gas was

measured on a GasBench II coupled to a Delta V Plus isotope ratio mass spectrometer (IRMS) at

the Stable Isotope Facility at the California Institute of Technology.

Concentrations of DIC were determined based on comparison of the average total peak area of

sample injections to replicate sodium bicarbonate standard curves. The 6-point dilution standard

curves for the porewater DIC used ultrapure DI H_2O as a background standard and a solution

series (1, 2.5, 5, 10, 20, and 30mM) of sodium bicarbonate (-2.9‰).

$\delta^{13}C_{nod}$ and $\delta^{13}C_{pw}$ values were corrected for instrument dependency on CO_2 concentration and

then normalized to the VPDB scale with a two-point calibration (107) using the Merck ($\delta^{13}C$ = -

49.2‰) and Carrara Marble (CM 2013, $\delta^{13}C$ = +2‰) carbonate standards. Here, we use solid

carbonate powder isotope standards for both $\delta^{13}C_{nod}$ and $\delta^{13}C_{pw}$ on the basis of a quantitative

conversion (i.e., total carbonate consumption) and since O isotopes are not considered for this

study.

Nodule porosity & porosimetry

During nodule sub-sampling, select nodules (where sufficient material allowed) were

additionally rinsed in deionized water to remove any salts and dried at room temperature for

structural and mineralogical analysis. To characterize the internal structure of the recovered

carbonate nodules, we determined their open porosity using a water intrusion (or Archimedes)

method. We followed Method A of ASTM D792 (108), which collects the dry mass, the

immersed mass (i.e., mass of fluid displaced), and the saturated mass (i.e., mass of fluid-

saturated object) of a nodule sample with DI water, according to Eq. 1:

777
$$\mu_a = \frac{(m_3 - m_1)}{(m_3 - m_2)} \times 100 \tag{1}$$

where μ_a is the apparent, or open, porosity (expressed as a percentage of the sample bulk

volume), m_1 is the dry mass, m_2 is the immersed mass, and m_3 is the saturated mass. This was

achieved with a density apparatus attached to a fine scale (Mettler Toledo). Additionally, we

measured the pore size distribution of paired samples with Hg-porosimetry. This method uses a

pressurized, non-wetting liquid (mercury) to intrude the nodule pore spaces, where the

cumulative volume of intruded mercury is recorded at each pressure interval, corresponding to a

certain pore size (109). We used a Micromeritics AutoPore IV Mercury Intrusion Porosimeter to

measure open pore size distribution between 6 and 350 µm by recording cumulative mercury

intrusion volume at 90 intervals within a pressure range of 3.51 and 206 kPa.

XRD

Nodule subsamples for mineralogical characterization with X-ray diffraction (XRD) were air-

dried and ground to fine powder with an agate mortar and pestle, cleaned with ethanol between

each use. Resulting powders were analyzed with a Rigaku SmartLab powder diffractometer at

the X-Ray Crystallography Facility in the Beckman Institute at the California Institute of

Technology. We used a Cu Kα source at 50 kV and 40 mA. Scans were run from 5° to 70° 2θ at

a scanning speed of 0.02° 2θ/s. Rigaku SmartLab Studio II was used to fit and identify peaks and

the 2020 Crystallography Open Database was used for phase identification. The relative semi-

quantification of clearly identifiable phases was calculated according to the reference intensity

ratio (RIR) based on a corundum standard and documented in weight percent. For the phases, the

RIR method used the (104) peak heights of calcite, Mg-calcite and dolomite, the (111) peak

height of aragonite, and the (101) peak height for quartz. Here, Mg-calcite corresponds to

$(Ca_{0.9}Mg_{0.1})O_3$, as measured by (35).

16S rRNA gene sequencing

To characterize and compare the diversity and composition of the microbial communities across

the seep areas represented in the dataset, we sequenced the V4 and V5 regions of the 16S rRNA

gene. Here, 2mL of sediment from each core horizon was sampled by sterile 1 mL cutoff

disposable syringe into a 2 mL screwcap Eppendorf tube and immediately stored at -80°C for

DNA extraction. In the laboratory, doubly rinsed nodules intended for sequencing were also

finely ground and frozen at -80°C after extraction from sediment and cleaning as described

above. DNA was extracted from 0.15-0.3 g (wet weight) of thawed sediment or finely ground

nodule powder using the DNeasy PowerSoil Pro Kit (Qiagen). Target sequences were then

amplified with a 2-step Illumina sequencing strategy (110). Briefly, this involved an initial

amplification with 515f (5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-

GTGYCAGCMGCCGCGGTAA-3') and 926r (5'-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-CCGYCAATTYMTTTRAGTTT-3')

primers, followed by a second amplification to include the necessary Illumina barcodes prior to

being sequenced.

Adapters from paired-end sequences were removed using 'Cutadapt' (v2.9), (111). Trimmed

reads were then input into QIIME2 (v2020.11), (112), where we used a DADA2 pipeline

(v1.14.1), (113) to filter and denoise the reads as well as remove chimeric sequences. ASVs with

assigned taxonomy were generated from the merged, non-chimeric sequences in QIIME2 using

the Silva V138.1 SSURef database feature classifier (114). Classified, non-zero ASVs for all the

sediment and nodules in the study are included in Table S11. Additional details on modified

DNA extraction protocols, PCR amplification, gene sequencing, and read processing are also

included in the Supplemental Material.

During subsequent analysis, we calculated Shannon-Wiener and Inverse Simpson diversity

indices for archaeal and bacterial ASVs from the carbonate nodules and their host sediments.

Further information on how these indices were calculated are detailed in the Supplemental

Material. Additionally, we performed a non-metric multidimensional scaling (NMDS) ordination

of all carbonate nodule and sediment-derived ASVs. Further details on how this ordination was

performed is also detailed in the Supplemental Material.

Fluorescence in situ hybridization (FISH)

During sediment core sectioning, 0.5-0.7 mL sediment per horizon was immediately added to 0.5

mL of a 0.22 μm filtered 4% paraformaldehyde solution in 3X PBS and fixed overnight at 4°C.

Afterwards, the sample was washed twice with 0.22 μm filtered 1X PBS and then re-suspended

in a 50% ethanol: 1X PBS medium and stored at -20°C. During nodule extraction in the

laboratory, doubly rinsed nodules were finely ground with a sterile agate mortar and pestle.

Immediately after grinding, the ground nodules were also fixed and washed as was done for the

sediments, but re-suspended in a 70% ethanol: 1X PBS medium prior to storage at -20°C. To

visualize the cells with minimal interference from sediment and mineral particles, we then

performed a density separation using percoll, based on a modified protocol outlined in (8, 23,

115). Briefly, 100-200 μL fixed sediment or carbonate was added to 400 μL of a chilled, 0.22 μm

filtered 'combination' buffer made of 0.5% v/v Tween 20, 3 mM sodium pyrophosphate, and

0.35% wt/v PVP (116). Samples were then sonicated on ice over 3 cycles at amplitude 20 on a

Q55 sonicator (Qsonica), with 10 seconds of active sonication and 10 seconds of rest in between

to avoid overheating. Next, sample tubes were shaken on a vortex mixer (Vortex-Genie 2,

Scientific Industries) for 30 minutes at room temperature. We then added 400 μL of chilled, 0.22

μm filtered 1X PBS before mixing and overlaying the mixed sample on 950 μL of a chilled

percoll density gradient. The gradient tubes were then centrifuged at 18,000 x g for 25 minutes at

4ºC (Microfuge 18, Beckman Coulter). After centrifugation, the density gradient supernatant was

transferred to a new set of sample tubes and centrifuged again at 16,000 x g for 3 minutes at 4ºC

to form cell pellets. The pellets were washed twice with 0.22 μm filtered 1X PBS.

The washed cells were then stained with a SYBR® Gold (Thermo Fisher) solution at a 25X final

concentration in ultrapure DI H_2O in the dark for 15 minutes at room temperature. After staining,

the cells were centrifuged at 16,000 x g for 3 minutes at room temperature, re-suspended in

ultrapure DI H_2O, and stored overnight at 4ºC. The next day, the cells were mounted via vacuum

filtration onto a 0.2 µm black polycarbonate filter and a 5µm PVDF backing filter (Millipore

Sigma). The filters were washed twice in ultrapure DI H_2O, air dried, and then stored in the dark

at 4ºC until counting. Prior to coverslip placement, each filter received ~ 50 µL

Vectashield® Antifade Mounting Medium (VectorLabs). For counts, we used an Olympus BX51

fluorescence microscope using a 100x objective lens (Olympus). Thirty fields of view per

filtered sample were surveyed to determine the number of cell aggregates and single cells. For

these counts, any association of cells larger than 5 µm were considered aggregates. Cell and

aggregate counts were normalized per unit dry mass for all sample types.

To observe the presence and morphologies associated with particular taxa, we used fluorescence

in situ hybridization (FISH). For FISH, the washed, percoll-extracted cells were re-suspended in

ultrapure DI H_2O and stored overnight at 4ºC. The following day, cell suspensions were dried at

room temperature onto separate 6 mm wells on PTFE- and L-lysine-coated glass slides (Tekdon).

The hybridization probes EUB338 [5'-GCTGCCTCCCGTAGGAGT-3' (117), 3' conjugated

Alexa Fluor™ 488 dye], EUB338 II [5'- GCAGCCACCCGTAGGTGT-3' (118), dual-labeled

Alexa Fluor™ 488 dye], EUB338 III [5'- GCTGCCACCCGTAGGTGT-3' (118), dual-labeled

Alexa Fluor™ 488 dye], ANME1-350 [5'-AGTTTTCGCGCCTGATGC-3' (19), dual-labeled

Alexa Fluor™ 546 dye], ANME1-728 [5'- GGTCTGGTCAGACGCCTT-3', designed using

ARB (119) to hit the ANME1 Clade of the Silva 138.1 database), 5' conjugated Alexa Fluor™

546 dye], and ARCH-915 [5'-GTGCTCCCCCGCCAATTCCT-3' (120), dual-labeled Alexa

Fluor™ 647 dye] were used for the mounted cell extracts. The hybridization probe mix was

prepared in sterile, 40% formamide concentration, with final concentrations of 5-ng/µL for each

probe (Integrated DNA Technologies). Each well on the slide then received 10 µL of probe mix

and was hybridized in the dark, overnight at 46°C. The next day, the glass slides with hybridized

cells were immersed and incubated in a pre-warmed hybridization wash in the dark for 10

minutes at 48°C. The slides were then rinsed in ultrapure DI H_2O and allowed to air dry (in the

dark) completely. FISH hybridization probe and wash recipes are provided in Tables S4-S5.

Prior to coverslip placement, each well also received 10 µL of a 90% Citifluor™ mounting

medium (Electron Microscopy Sciences) with 4.5 ng/µL 4',6-diamidino-2-phenylindole (DAPI)

for nucleic acid visualization. Hybridized cells were then visualized with an ELYRA S.1 / Axio

Observer Z1 super resolution microscope using a 100x objective lens (Zeiss). For control

experiments, each glass slide had a well with mounted cells subjected to the entire FISH protocol

but used ultrapure DI H_2O instead of probes. For these wells, negligible signal was observed.

Single cell translational activity measurements with BONCAT

Translational activity was observed by amending a subset of the bottled nodules with an anoxic

L-Homopropargylglycine (HPG) solution. HPG works as a methionine analog and is

incorporated during protein synthesis without significantly altering community composition or

metabolic activity (49, 51, 52). After incorporation, proteins containing HPG can be

fluorescently labeled through copper-catalyzed azide-alkyne 'click' chemistry, which enables

detection via fluorescence microscopy (49, 52, 121, 122).

In this study, the nodules previously maintained in N_2-sparged artificial seawater with a 2 bar

CH_4 headspace were separated into replicate experimental and control bottles. Where possible,

the nodules were selected to minimize mass differences between experimental and control

groups. Each new sterile, acid-washed bottle received 50 mL of 0.22 μm-filtered, N_2-sparged

artificial seawater of the same composition (Table S6, sufficient to immerse the nodules).

Throughout transfer and immersion, all bottles were constantly maintained under active N_2 flow

to minimize oxygen exposure. After immersion, all bottle headspaces were flushed with pure

CH_4 for 1 minute and pressurized to 2 bar. Samples were then incubated in the dark at 4°C for a

period of 5 weeks, during which the media was periodically sampled to monitor sulfide

concentrations in control and experimental bottles. After determining comparable sulfide

production between each control and experimental group (Fig. S15), the artificial seawater media

was replaced with freshly prepared 0.22 μm-filtered, N_2-sparged artificial seawater. Additionally,

each experimental bottle received aqueous 0.22 μm-filtered, N_2-sparged HPG to a final

concentration of 200 μM. Throughout media replacement and HPG addition, all bottles were

constantly maintained under active N_2 flow to minimize oxygen exposure. After immersion, all

bottle headspaces were again flushed with pure CH_4 for 1 minute and pressurized to 2 bar.

Samples were then incubated in the dark at 4°C for a period of 14 weeks to ensure sufficient

HPG assimilation.

At the end of the incubation period, nodules from both control and experimental bottles were

ground, fixed, and washed as previously described for FISH hybridization. Cells from each

fixed, ground nodule were then extracted and mounted onto separate 6 mm wells on PTFE- and

L-lysine-coated glass slides as previously described. For the BONCAT dye click reaction, the

glass-mounted cell extracts were then dehydrated by sequential immersion in 50% (1 minute),

80% (3 minutes), and 90% (3 minutes) ethanol: DI H_2O baths and dried room temperature.

We used BONCAT azide dye Oregon Green 488 (VectorLabs) at a final concentration of 20 µM.

10-µL of the dye solution was then added to each well and incubated in the dark for 30 minutes.

For the incubation, we flushed a lidded sample container with Ar, in which we placed the

mounted cells to minimize oxygen exposure, which can reduce the fluorescent signal. After

incubation, the slides were washed in 0.22 µm-filtered 1X PBS for 10 minutes (in the dark),

quickly rinsed in ultrapure DI H_2O, and allowed to air dry in the dark prior to FISH

hybridization. Hybridization probes EUB338 [5' conjugated cyanine 3 dye], EUB338 II [3'

conjugated cyanine 3 dye], EUB338 III [5' conjugated cyanine 3 dye], and ARCH-915 [dual-

labeled Alexa Fluor™ 647 dye] were used. Hybridization was done at 35% formamide. Washing

and addition of DAPI-Citifluor® mounting medium was performed as previously described.

BONCAT click reaction and FISH hybridization recipes are listed in Tables S7-S9.

Click-dyed, hybridized cells were then visualized with an ELYRA S.1 / Axio Observer Z1 super

resolution microscope using a 100x objective lens (Zeiss). Fraction of cells with HPG signal

were evaluated by counting HPG-incubated, extracted cells with observable signal above a mean

background threshold calculated from the control cells' auto-fluorescence in the same channel.

For FISH and BONCAT control experiments, each glass slide had a well with mounted, non-

HPG labeled cells subjected to the entire FISH protocol but used ultrapure DI H_2O instead of

probes. For these cells in these negative control wells, a negligible signal was observed.

Data availability

The Santa Monica Mounds Low Altitude Survey System (LASS) survey was done in the Spring

of 2018 using *ROV Ventana* on *R/V Rachel Carson*. The Santa Monica Mounds Mapping AUV

surveys were done multiple times from *R/V Zephyr* and *R/V Carson*. Featured grids and

photomosaics will be available at MGDS https://www.marine-geo.org/.

The raw DNA sequence reads generated from the SMM 800 sediment cores in this study were

initially submitted to NCBI under project accession number PRJNA1196099 (sample accession

numbers: SAMN45814414-SAMN45814434; SAMN45814441- SAMN45814448) and are

accessible at https://www.ncbi.nlm.nih.gov/bioproject/ PRJNA1196099. The rest of the raw

DNA sequence reads generated for this study (SMM 800 nodules, SMM 863 samples,

experimental controls) were submitted to NCBI under accession numbers PRJNA1207588 and

accessible at https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1207588.

Acknowledgements

We are indebted to the captain, crew, and pilots of the *ROV Doc Ricketts* from the WF05-21

cruise, who made this work possible. We also thank David Caress, Charles Paull, Eric Martin,

and the Autonomous Systems Operations group at the Monterey Bay Aquarium Research

Institute (MBARI), as well as the crews of *R/V Zephyr* and *R/V Carson* for collection,

processing, and analysis of bathymetry maps of the Santa Monica Basin. We thank S. Lim

(University of Nevada, Las Vegas), J.S. Magyar (Caltech), and S. Goffredi (Occidental College)

for their help in sample collection and processing during the WF05-21 Southern California cruise

of the *R/V Western Flyer*, S. Connon (Caltech) for technical support and barcoding of 16S

amplicons for sequencing, D. Utter (Caltech) and H. Sapers (Northern Arizona University) for

support during data processing.

S.A. Parra and V.J. Orphan conceptualized the study and acquired financial support for the

study. J. Mullahoo analyzed dissolved porewater methane. L.K. Quinn analyzed nodule open

porosity and Hg-porosimetry. R.L. Wipfler analyzed porewater sulfate concentrations. S.A. Parra

processed and analyzed all other data presented in the study, including nodule selection,

porewater sulfide measurements, DIC/δ^{13}C measurements, x-ray diffraction, DNA extraction,

16S rRNA sequence analyses, fluorescence microscopy, and BONCAT incubations. M.J. Mayr

and V.J. Orphan supervised the study. S.A. Parra wrote the original draft and incorporated edits

before and during the publication stage. M.J. Mayr, J. Mullahoo, L.K. Quinn, R.L. Wipfler, and

V.J. Orphan all participated in critical review and commentary on the paper draft before and

during the publication stage.

This work was supported in part by NASA's Interdisciplinary Consortia for Astrobiology

Research (ICAR) program under Award Number AWD-005316-G4 (to V.J. Orphan), the

National Science Foundation under Award Number 2048666 (to V.J. Orphan), and NASA's

Future Investigators in NASA Earth and Space Science and Technology (FINESST) program

under Award Number 80NSSC22K1336 (to S.A. Parra). This material is based upon work

supported by the U.S. Department of Energy, Office of Science Biological and Environmental

Research Program under Award Number DE-SC0022991 (to V.J. Orphan).

Disclaimer: This report was prepared as an account of work sponsored by an agency of the

United States Government. Neither the United States Government nor any agency thereof, nor

any of their employees, makes any warranty, express or implied, or assumes any legal liability or

responsibility for the accuracy, completeness, or usefulness of any information, apparatus,

product, or process disclosed, or represents that its use would not infringe privately owned rights.

Reference herein to any specific commercial product, process, or service by trade name,

trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement,

recommendation, or favoring by the United States Government or any agency thereof. The views

and opinions of authors expressed herein do not necessarily state or reflect those of the United

States Government or any agency thereof.

References

1. Reeburgh WS. 2007. Oceanic Methane Biogeochemistry. Chem Rev 107:486–513.

2. Boetius A, Wenzhöfer F. 2013. Seafloor oxygen consumption fuelled by methane from
cold seeps. 9. Nature Geosci 6:725–734.

3. Aharon P, Schwarcz HP, Roberts HH. 1997. Radiometric dating of submarine
hydrocarbon seeps in the Gulf of Mexico. GSA Bulletin 109:568–579.

4. Peckmann J, Thiel V. 2004. Carbon cycling at ancient methane–seeps. Chemical Geology
205:443–467.

5. Teichert BMA, Bohrmann G, Suess E. 2005. Chemoherms on Hydrate Ridge — Unique
microbially-mediated carbonate build-ups growing into the water column.
Palaeogeography, Palaeoclimatology, Palaeoecology 227:67–85.

6. Naehr TH, Eichhubl P, Orphan VJ, Hovland M, Paull CK, Ussler W, Lorenson TD,
Greene HG. 2007. Authigenic carbonate formation at hydrocarbon seeps in continental
margin sediments: A comparative study. Deep Sea Research Part II: Topical Studies in
Oceanography 54:1268–1291.

7.   Haas A, Peckmann J, Elvert M, Sahling H, Bohrmann G. 2010. Patterns of carbonate

1009   authigenesis at the Kouilou pockmarks on the Congo deep-sea fan. Marine Geology

1010   268:129–136.

1011   8.   Marlow JJ, Steele JA, Ziebis W, Thurber AR, Levin LA, Orphan VJ. 2014. Carbonate-

1012   hosted methanotrophy represents an unrecognized methane sink in the deep sea. 1. Nature

1013   Communications 5:5094.

1014   9.   Marlow J, Peckmann J, Orphan V. 2015. Autoendoliths: a distinct type of rock-hosted

1015   microbial life. Geobiology 13:303–307.

1016   10.   Marlow JJ, Hoer D, Jungbluth SP, Reynard LM, Gartman A, Chavez MS, El-Naggar MY,

1017   Tuross N, Orphan VJ, Girguis PR. 2021. Carbonate-hosted microbial communities are

1018   prolific and pervasive methane oxidizers at geologically diverse marine methane seep

1019   sites. PNAS 118.

1020   11.   Mayr MJ, Parra SA, Connon SA, Narayanan AK, Murali R, Cremiere A, Orphan VJ.

1021   2025. Distinct Microbial Communities Within and On Seep Carbonates Support Long-

1022   term Anaerobic Oxidation of Methane and Novel pMMO Diversity. bioRxiv

1023   https://doi.org/10.1101/2025.02.04.636526.

1024   12.   Lein AYu. 2004. Authigenic Carbonate Formation in the Ocean. Lithology and Mineral

1025   Resources 39:1–30.

1026   13.   Watanabe Y, Nakai S, Hiruta A, Matsumoto R, Yoshida K. 2008. U–Th dating of

1027   carbonate nodules from methane seeps off Joetsu, Eastern Margin of Japan Sea. Earth and

1028   Planetary Science Letters 272:89–96.

1029   14.   Mason OU, Case DH, Naehr TH, Lee RW, Thomas RB, Bailey JV, Orphan VJ. 2015.

1030   Comparison of Archaeal and Bacterial Diversity in Methane Seep Carbonate Nodules and

1031   Host Sediments, Eel River Basin and Hydrate Ridge, USA. Microb Ecol 70:766–784.

1032   15.   Yanagawa K, Shiraishi F, Tanigawa Y, Maeda T, Mustapha NA, Owari S, Tomaru H,

1033   Matsumoto R, Kano A. 2019. Endolithic Microbial Habitats Hosted in Carbonate Nodules

1034   Currently Forming within Sediment at a High Methane Flux Site in the Sea of Japan. 11.

1035   Geosciences 9:463.

1036   16.   Case DH, Pasulka AL, Marlow JJ, Grupe BM, Levin LA, Orphan VJ. 2015. Methane Seep

1037   Carbonates Host Distinct, Diverse, and Dynamic Microbial Assemblages. mBio 6.

1038   17.   Hoehler TM, Alperin MJ, Albert DB, Martens CS. 1994. Field and laboratory studies of

1039   methane oxidation in an anoxic marine sediment: Evidence for a methanogen-sulfate

1040   reducer consortium. Global Biogeochemical Cycles 8:451–463.

1041   18.   Hinrichs K-U, Hayes JM, Sylva SP, Brewer PG, DeLong EF. 1999. Methane-consuming

1042   archaebacteria in marine sediments. Nature 398:802–805.

1043   19.   Boetius A, Ravenschlag K, Schubert CJ, Rickert D, Widdel F, Gieseke A, Amann R,

1044   Jørgensen BB, Witte U, Pfannkuche O. 2000. A marine microbial consortium apparently

1045   mediating anaerobic oxidation of methane. Nature 407:623–626.

1046   20.   Orphan VJ, House CH, Hinrichs K-U, McKeegan KD, DeLong EF. 2001. Methane-

1047   Consuming Archaea Revealed by Directly Coupled Isotopic and Phylogenetic Analysis.

1048   Science 293:484–487.

21.   Borowski W, Paull C, Ussler W. 1996. Marine Pore-water Sulfate Profiles Indicate in situ

Methane Flux from Underlying Gas Hydrate. Geology 655–658.

22.   Marlow JJ, Steele JA, Case DH, Connon SA, Levin LA, Orphan VJ. 2014. Microbial

abundance and diversity patterns associated with sediments and carbonates from the

methane seep environments of Hydrate Ridge, OR. Front Mar Sci 1.

23.   Orphan VJ, House CH, Hinrichs K-U, McKeegan KD, DeLong EF. 2002. Multiple

archaeal groups mediate methane oxidation in anoxic cold seep sediments. Proceedings of

the National Academy of Sciences 99:7663–7668.

24.   Orphan VJ, Ussler W, Naehr TH, House CH, Hinrichs K-U, Paull CK. 2004. Geological,

geochemical, and microbiological heterogeneity of the seafloor around methane vents in

the Eel River Basin, offshore California. Chemical Geology 205:265–289.

25.   Blumenberg M, Seifert R, Reitner J, Pape T, Michaelis W. 2004. Membrane lipid patterns

typify distinct anaerobic methanotrophic consortia. Proceedings of the National Academy

of Sciences 101:11111–11116.

26.   Harrison BK, Zhang H, Berelson W, Orphan VJ. 2009. Variations in Archaeal and

Bacterial Diversity Associated with the Sulfate-Methane Transition Zone in Continental

Margin Sediments (Santa Barbara Basin, California). Applied and Environmental

Microbiology 75:1487–1499.

27.   Rossel PE, Elvert M, Ramette A, Boetius A, Hinrichs K-U. 2011. Factors controlling the

distribution of anaerobic methanotrophic communities in marine environments: Evidence

from intact polar membrane lipids. Geochimica et Cosmochimica Acta 75:164–184.

28.   Yanagawa K, Sunamura M, Lever MA, Morono Y, Hiruta A, Ishizaki O, Matsumoto R,

Urabe T, Inagaki F. 2011. Niche Separation of Methanotrophic Archaea (ANME-1 and -2)

in Methane-Seep Sediments of the Eastern Japan Sea Offshore Joetsu. Geomicrobiology

Journal 28:118–129.

29.   Chen Y, Xu C, Wu N, Sun Z, Liu C, Zhen Y, Xin Y, Zhang X, Geng W, Cao H, Zhai B,

Li J, Qin S, Zhou Y. 2022. Diversity of Anaerobic Methane Oxidizers in the Cold Seep

Sediments of the Okinawa Trough. Front Microbiol 13.

30.   Cui H, Su X, Chen F, Holland M, Yang S, Liang J, Su P, Dong H, Hou W. 2019.

Microbial diversity of two cold seep systems in gas hydrate-bearing sediments in the

South China Sea. Marine Environmental Research 144:230–239.

31.   Chen J, Li Y, Zhong C, Xu Z, Lu G, Jing H, Liu H. 2023. Genomic Insights into Niche

Partitioning across Sediment Depth among Anaerobic Methane-Oxidizing Archaea in

Global Methane Seeps. mSystems 8:e01179-22.

32.   Zhang C, Fang Y-X, Yin X, Lai H, Kuang Z, Zhang T, Xu X-P, Wegener G, Wang J-H,

Dong X. 2023. The majority of microorganisms in gas hydrate-bearing subseafloor

sediments ferment macromolecules. Microbiome 11:37.

33.   Paull CK, Normark WR, Ussler W, Caress DW, Keaten R. 2008. Association among

active seafloor deformation, mound formation, and gas hydrate growth and accumulation

within the seafloor of the Santa Monica Basin, offshore California. Marine Geology

250:258–275.

34. Normark WR, Piper DJW, Sliter R. 2006. Sea-level and tectonic control of middle to late

Pleistocene turbidite systems in Santa Monica Basin, offshore California. Sedimentology

53:867–897.

35. Althoff PL. 1977. Structural refinements of dolomite and a magnesian calcite and

implications for dolomite formation in the marine environment. American Mineralogist

62:772–783.

36. Ussler W, Paull CK. 2008. Rates of anaerobic oxidation of methane and authigenic

carbonate mineralization in methane-rich deep-sea sediments inferred from models and

geochemical profiles. Earth and Planetary Science Letters 266:271–287.

37. Komada T, Burdige DJ, Li H-L, Magen C, Chanton JP, Cada AK. 2016. Organic matter

cycling across the sulfate-methane transition zone of the Santa Barbara Basin, California

Borderland. Geochimica et Cosmochimica Acta 176:259–278.

38. Tarnovetskii IY, Merkel AY, Kanapatskiy TA, Ivanova EA, Gulin MB, Toshchakov S,

Pimenov NV. 2018. Decoupling between sulfate reduction and the anaerobic oxidation of

methane in the shallow methane seep of the Black sea. FEMS Microbiology Letters

365:fny235.

39. Kirkegaard RH, Dueholm MS, McIlroy SJ, Nierychlo M, Karst SM, Albertsen M, Nielsen

PH. 2016. Genomic insights into members of the candidate phylum Hyd24-12 common in

mesophilic anaerobic digesters. The ISME Journal 10:2352–2364.

40.   Schreiber L, Holler T, Knittel K, Meyerdierks A, Amann R. 2010. Identification of the

1110   dominant sulfate-reducing bacterial partner of anaerobic methanotrophs of the ANME-2

1111   clade. Environmental Microbiology 12:2327–2340.

1112   41.   McGlynn SE, Chadwick GL, Kempes CP, Orphan VJ. 2015. Single cell activity reveals

1113   direct electron transfer in methanotrophic consortia. Nature 526:531–535.

1114   42.   Metcalfe KS, Murali R, Mullin SW, Connon SA, Orphan VJ. 2021. Experimentally-

1115   validated correlation analysis reveals new anaerobic methane oxidation partnerships with

1116   consortium-level heterogeneity in diazotrophy. ISME J 15:377–396.

1117   43.   Murali R, Yu H, Speth DR, Wu F, Metcalfe KS, Crémière A, Laso-Pèrez R, Malmstrom

1118   RR, Goudeau D, Woyke T, Hatzenpichler R, Chadwick GL, Connon SA, Orphan VJ.

1119   2023. Physiological potential and evolutionary trajectories of syntrophic sulfate-reducing

1120   bacterial partners of anaerobic methanotrophic archaea. PLOS Biology 21:e3002292.

1121   44.   Nobu MK, Narihiro T, Kuroda K, Mei R, Liu W-T. 2016. Chasing the elusive

1122   Euryarchaeota class WSA2: genomes reveal a uniquely fastidious methyl-reducing

1123   methanogen. 10. ISME J 10:2478–2487.

1124   45.   Dhillon A, Lever M, Lloyd KG, Albert DB, Sogin ML, Teske A. 2005. Methanogen

1125   Diversity Evidenced by Molecular Characterization of Methyl Coenzyme M Reductase A

1126   (mcrA) Genes in Hydrothermal Sediments of the Guaymas Basin. Applied and

1127   Environmental Microbiology 71:4592–4601.

1128   46.   Teske A, Wegener G, Chanton JP, White D, MacGregor B, Hoer D, de Beer D, Zhuang G,

1129   Saxton MA, Joye SB, Lizarralde D, Soule SA, Ruff SE. 2021. Microbial Communities

Under Distinct Thermal and Geochemical Regimes in Axial and Off-Axis Sediments of

Guaymas Basin. Frontiers in Microbiology 12.

47. Lai H, Deng Y, Yang L, Liang J, Dai L, Li L, Fang Y, Liu L, Kuang Z. 2023. Origin of

natural gas within the deep-sea uncompacted sediments of the Shenhu area, northern

South China Sea: Geochemical and methanogenic cultivation results. Marine and

Petroleum Geology 147:106015.

48. Coon GR, Duesing PD, Paul R, Baily JA, Lloyd KG. 2023. Biological methane production

and accumulation under sulfate-rich conditions at Cape Lookout Bight, NC. Front

Microbiol 14.

49. Hatzenpichler R, Connon SA, Goudeau D, Malmstrom RR, Woyke T, Orphan VJ. 2016.

Visualizing in situ translational activity for identifying and sorting slow-growing

archaeal–bacterial consortia. PNAS 113:E4069–E4078.

50. Krukenberg V, Riedel D, Gruber-Vodicka HR, Buttigieg PL, Tegetmeyer HE, Boetius A,

Wegener G. 2018. Gene expression and ultrastructure of meso- and thermophilic

methanotrophic consortia. Environmental Microbiology 20:1651–1666.

51. Hatzenpichler R, Scheller S, Tavormina PL, Babin BM, Tirrell DA, Orphan VJ. 2014. In

situ visualization of newly synthesized proteins in environmental microbes using amino

acid tagging and click chemistry. Environmental Microbiology 16:2568–2590.

52. Hatzenpichler R, Orphan VJ. 2016. Detection of Protein-Synthesizing Microorganisms in

the Environment via Bioorthogonal Noncanonical Amino Acid Tagging (BONCAT), p.

145–157. *In* McGenity, TJ, Timmis, KN, Nogales, B (eds.), Hydrocarbon and Lipid

1151   Microbiology Protocols: Single-Cell and Single-Molecule Methods. Springer, Berlin,

1152   Heidelberg.

1153 53. Schmale O, Leifer I, Deimling JS v., Stolle C, Krause S, Kießlich K, Frahm A, Treude T.

1154   2015. Bubble Transport Mechanism: Indications for a gas bubble-mediated inoculation of

1155   benthic methanotrophs into the water column. Continental Shelf Research 103:70–78.

1156 54. Crémière A, Lepland A, Chand S, Sahy D, Kirsimäe K, Bau M, Whitehouse MJ, Noble

1157   SR, Martma T, Thorsnes T, Brunstad H. 2016. Fluid source and methane-related

1158   diagenetic processes recorded in cold seep carbonates from the Alvheim channel, central

1159   North Sea. Chemical Geology 432:16–33.

1160 55. Tryon MD, Brown KM, Torres ME. 2002. Fluid and chemical flux in and out of sediments

1161   hosting methane hydrate deposits on Hydrate Ridge, OR, II: Hydrological processes. Earth

1162   and Planetary Science Letters 201:541–557.

1163 56. Treude T, Boetius A, Knittel K, Wallmann K, Barker Jørgensen B. 2003. Anaerobic

1164   oxidation of methane above gas hydrates at Hydrate Ridge, NE Pacific Ocean. Mar Ecol

1165   Prog Ser 264:1–14.

1166 57. Heeschen KU, Collier RW, de Angelis MA, Suess E, Rehder G, Linke P, Klinkhammer

1167   GP. 2005. Methane sources, distributions, and fluxes from cold vent sites at Hydrate

1168   Ridge, Cascadia Margin. Global Biogeochemical Cycles 19.

1169 58. Stadnitskaia A, Muyzer G, Abbas B, Coolen MJL, Hopmans EC, Baas M, van Weering

1170   TCE, Ivanov MK, Poludetkina E, Sinninghe Damsté JS. 2005. Biomarker and 16S rDNA

1171   evidence for anaerobic oxidation of methane and related carbonate precipitation in deep-

1172   sea mud volcanoes of the Sorokin Trough, Black Sea. Marine Geology 217:67–96.

1173   59.   Stadnitskaia A, Nadezhkin D, Abbas B, Blinova V, Ivanov MK, Sinninghe Damsté JS.

1174   2008. Carbonate formation by anaerobic oxidation of methane: Evidence from lipid

1175   biomarker and fossil 16S rDNA. Geochimica et Cosmochimica Acta 72:1824–1836.

1176   60.   Lloyd KG, Albert DB, Biddle JF, Chanton JP, Pizarro O, Teske A. 2010. Spatial Structure

1177   and Activity of Sedimentary Microbial Communities Underlying a Beggiatoa spp. Mat in

1178   a Gulf of Mexico Hydrocarbon Seep. PLOS ONE 5:e8738.

1179   61.   Meister P, Liu B, Khalili A, Böttcher ME, Jørgensen BB. 2019. Factors controlling the

1180   carbon isotope composition of dissolved inorganic carbon and methane in marine

1181   porewater: An evaluation by reaction-transport modelling. Journal of Marine Systems

1182   200:103227.

1183   62.   Peckmann J, Reimer A, Luth U, Luth C, Hansen BT, Heinicke C, Hoefs J, Reitner J. 2001.

1184   Methane-derived carbonates and authigenic pyrite from the northwestern Black Sea.

1185   Marine Geology 177:129–150.

1186   63.   Magalhães VH, Pinheiro LM, Ivanov MK, Kozlova E, Blinova V, Kolganova J,

1187   Vasconcelos C, McKenzie JA, Bernasconi SM, Kopf AJ, Díaz-del-Río V, González FJ,

1188   Somoza L. 2012. Formation processes of methane-derived authigenic carbonates from the

1189   Gulf of Cadiz. Sedimentary Geology 243–244:155–168.

1190   64.   Greinert J, Bohrmann G, Suess E. 2001. Gas Hydrate-Associated Carbonates and

1191   Methane-Venting at Hydrate Ridge: Classification, Distribution, and Origin of Authigenic

Lithologies, p. 99–113. *In* Natural Gas Hydrates: Occurrence, Distribution, and Detection.

American Geophysical Union (AGU).

65.   Aloisi G, Bouloubassi I, Heijs SK, Pancost RD, Pierre C, Sinninghe Damsté JS, Gottschal

JC, Forney LJ, Rouchy J-M. 2002. CH4-consuming microorganisms and the formation of

carbonate crusts at cold seeps. Earth and Planetary Science Letters 203:195–203.

66.   Knittel K, Lösekann T, Boetius A, Kort R, Amann R. 2005. Diversity and Distribution of

Methanotrophic Archaea at Cold Seeps. Applied and Environmental Microbiology

71:467–479.

67.   Orcutt B, Boetius A, Elvert M, Samarkin V, Joye SB. 2005. Molecular biogeochemistry of

sulfate reduction, methanogenesis and the anaerobic oxidation of methane at Gulf of

Mexico cold seeps. Geochimica et Cosmochimica Acta 69:4267–4281.

68.   Treude T, Knittel K, Blumenberg M, Seifert R, Boetius A. 2005. Subsurface Microbial

Methanotrophic Mats in the Black Sea. Applied and Environmental Microbiology

71:6375–6378.

69.   Treude T, Orphan V, Knittel K, Gieseke A, House CH, Boetius A. 2007. Consumption of

Methane and CO2 by Methanotrophic Microbial Mats from Gas Seeps of the Anoxic

Black Sea. Applied and Environmental Microbiology 73:2271–2283.

70.   House CH, Orphan VJ, Turk KA, Thomas B, Pernthaler A, Vrentas JM, Joye SB. 2009.

Extensive carbon isotopic heterogeneity among methane seep microbiota. Environmental

Microbiology 11:2207–2215.

71.   Ruff SE, Kuhfuss H, Wegener G, Lott C, Ramette A, Wiedling J, Knittel K, Weber M.

1213   2016. Methane Seep in Shallow-Water Permeable Sediment Harbors High Diversity of

1214   Anaerobic Methanotrophic Communities, Elba, Italy. Frontiers in Microbiology 7.

1215   72.   Webster G, Parkes RJ, Fry JC, Weightman AJ. 2004. Widespread Occurrence of a Novel

1216   Division of Bacteria Identified by 16S rRNA Gene Sequences Originally Found in Deep

1217   Marine Sediments. Applied and Environmental Microbiology 70:5708–5713.

1218   73.   Lee YM, Hwang K, Lee JI, Kim M, Hwang CY, Noh H-J, Choi H, Lee HK, Chun J, Hong

1219   SG, Shin SC. 2018. Genomic Insight Into the Predominance of Candidate Phylum

1220   Atribacteria JS1 Lineage in Marine Sediments. Front Microbiol 9.

1221   74.   Chakraborty A, Ruff SE, Dong X, Ellefson ED, Li C, Brooks JM, McBee J, Bernard BB,

1222   Hubert CRJ. 2020. Hydrocarbon seepage in the deep seabed links subsurface and seafloor

1223   biospheres. PNAS 117:11029–11037.

1224   75.   Liu Y-F, Qi Z-Z, Shou L-B, Liu J-F, Yang S-Z, Gu J-D, Mu B-Z. 2019. Anaerobic

1225   hydrocarbon degradation in candidate phylum 'Atribacteria' (JS1) inferred from

1226   genomics. The ISME Journal 13:2377–2390.

1227   76.   Biddle JF, Cardman Z, Mendlovitz H, Albert DB, Lloyd KG, Boetius A, Teske A. 2012.

1228   Anaerobic oxidation of methane at different temperature regimes in Guaymas Basin

1229   hydrothermal sediments. ISME J 6:1018–1031.

1230   77.   Bertram S, Blumenberg M, Michaelis W, Siegert M, Krüger M, Seifert R. 2013.

1231   Methanogenic capabilities of ANME-archaea deduced from -labelling approaches.

1232   Environmental Microbiology 15:2384–2393.

1233   78.   Beulig F, Røy H, McGlynn SE, Jørgensen BB. 2019. Cryptic CH4 cycling in the sulfate–

1234   methane transition of marine sediments apparently mediated by ANME-1 archaea. The

1235   ISME Journal 13:250–262.

1236   79.   Kevorkian RT, Callahan S, Winstead R, Lloyd KG. 2021. ANME-1 archaea may drive

1237   methane accumulation and removal in estuarine sediments. Environ Microbiol Rep

1238   13:185–194.

1239   80.   Chadwick GL, Skennerton CT, Laso-Pérez R, Leu AO, Speth DR, Yu H, Morgan-Lang C,

1240   Hatzenpichler R, Goudeau D, Malmstrom R, Brazelton WJ, Woyke T, Hallam SJ, Tyson

1241   GW, Wegener G, Boetius A, Orphan VJ. 2022. Comparative genomics reveals electron

1242   transfer and syntrophic mechanisms differentiating methanotrophic and methanogenic

1243   archaea. PLOS Biology 20:e3001508.

1244   81.   Conrad R. 2005. Quantification of methanogenic pathways using stable carbon isotopic

1245   signatures: a review and a proposal. Organic Geochemistry 36:739–752.

1246   82.   Chatterjee S, Dickens GR, Bhatnagar G, Chapman WG, Dugan B, Snyder GT, Hirasaki

1247   GJ. 2011. Pore water sulfate, alkalinity, and carbon isotope profiles in shallow sediment

1248   above marine gas hydrate systems: A numerical modeling perspective. Journal of

1249   Geophysical Research: Solid Earth 116.

1250   83.   Beal EJ, House CH, Orphan VJ. 2009. Manganese- and Iron-Dependent Marine Methane

1251   Oxidation. Science 325:184–187.

84. Riedinger N, Formolo MJ, Lyons TW, Henkel S, Beck A, Kasten S. 2014. An inorganic

geochemical argument for coupled anaerobic oxidation of methane and iron reduction in

marine sediments. Geobiology 12:172–181.

85. Oni OE, Miyatake T, Kasten S, Richter-Heitmann T, Fischer D, Wagenknecht L,

Ksenofontov V, Kulkarni A, Blumers M, Shylin S, Costa B, Klingelhöfer G, Friedrich

MW. 2015. Distinct microbial populations are tightly linked to the profile of dissolved

iron in the methanic sediments of the Helgoland mud area, North Sea. Front Microbiol 6.

86. Sun Z, Wei H, Zhang X, Shang L, Yin X, Sun Y, Xu L, Huang W, Zhang X. 2015. A

unique Fe-rich carbonate chimney associated with cold seeps in the Northern Okinawa

Trough, East China Sea. Deep Sea Research Part I: Oceanographic Research Papers

95:37–53.

87. Egger M, Rasigraf O, Sapart CJ, Jilbert T, Jetten MSM, Röckmann T, van der Veen C,

Bândă N, Kartal B, Ettwig KF, Slomp CP. 2015. Iron-Mediated Anaerobic Oxidation of

Methane in Brackish Coastal Sediments. Environ Sci Technol 49:277–283.

88. Egger M, Hagens M, Sapart CJ, Dijkstra N, van Helmond NAGM, Mogollón JM,

Risgaard-Petersen N, van der Veen C, Kasten S, Riedinger N, Böttcher ME, Röckmann T,

Jørgensen BB, Slomp CP. 2017. Iron oxide reduction in methane-rich deep Baltic Sea

sediments. Geochimica et Cosmochimica Acta 207:256–276.

89. Rooze J, Egger M, Tsandev I, Slomp CP. 2016. Iron-dependent anaerobic oxidation of

methane in coastal surface sediments: Potential controls and impact. Limnology and

Oceanography 61:S267–S282.

90.   Li J, Peng X, Bai S, Chen Z, Van Nostrand JD. 2018. Biogeochemical processes

1274   controlling authigenic carbonate formation within the sediment column from the Okinawa

1275   Trough. Geochimica et Cosmochimica Acta 222:363–382.

1276   91.   Aromokeye DA, Kulkarni AC, Elvert M, Wegener G, Henkel S, Coffinet S, Eickhorst T,

1277   Oni OE, Richter-Heitmann T, Schnakenberg A, Taubner H, Wunder L, Yin X, Zhu Q,

1278   Hinrichs K-U, Kasten S, Friedrich MW. 2020. Rates and Microbial Players of Iron-Driven

1279   Anaerobic Oxidation of Methane in Methanic Marine Sediments. Front Microbiol 10.

1280   92.   Mitterer RM. 2010. Methanogenesis and sulfate reduction in marine sediments: A new

1281   model. Earth and Planetary Science Letters 295:358–366.

1282   93.   Dell'Anno A, Stefano B, Danovaro R. 2002. Quantification, base composition, and fate of

1283   extracellular DNA in marine sediments. Limnology and Oceanography 47:899–905.

1284   94.   Dell'Anno A, Danovaro R. 2005. Extracellular DNA Plays a Key Role in Deep-Sea

1285   Ecosystem Functioning. Science 309:2179–2179.

1286   95.   Torti A, Lever MA, Jørgensen BB. 2015. Origin, dynamics, and implications of

1287   extracellular DNA pools in marine sediments. Marine Genomics 24:185–196.

1288   96.   Lennon JT, Muscarella ME, Placella SA, Lehmkuhl BK. 2018. How, When, and Where

1289   Relic DNA Affects Microbial Diversity. mBio 9:10.1128/mbio.00637-18.

1290   97.   Torti A, Jørgensen BB, Lever MA. 2018. Preservation of microbial DNA in marine

1291   sediments: insights from extracellular DNA pools. Environmental Microbiology 20:4526–

1292   4542.

98.   Cai W-J, Chen F, Powell EN, Walker SE, Parsons-Hubbard KM, Staff GM, Wang Y,

Ashton-Alcox KA, Callender WR, Brett CE. 2006. Preferential dissolution of carbonate

shells driven by petroleum seep activity in the Gulf of Mexico. Earth and Planetary

Science Letters 248:227–243.

99.   Himmler T, Brinkmann F, Bohrmann G, Peckmann J. 2011. Corrosion patterns of seep-

carbonates from the eastern Mediterranean Sea. Terra Nova 23:206–212.

100. Matsumoto R. 1990. Vuggy carbonate crust formed by hydrocarbon seepage on the

continental shelf of Baffin Island, northeast Canada. Geochemical Journal 24:143–158.

101. Carson B, Kastner M, Bartlett D, Jaeger J, Jannasch H, Weinstein Y. 2003. Implications of

carbon flux from the Cascadia accretionary prism: results from long-term, in situ

measurements at ODP Site 892B. Marine Geology 198:159–180.

102. Krause S, Aloisi G, Engel A, Liebetrau V, Treude T. 2014. Enhanced Calcite Dissolution

in the Presence of the Aerobic Methanotroph Methylosinus trichosporium.

Geomicrobiology Journal 31:325–337.

103. Leprich DJ, Flood BE, Schroedl PR, Ricci E, Marlow JJ, Girguis PR, Bailey JV. 2021.

Sulfur bacteria promote dissolution of authigenic carbonates at marine methane seeps.

ISME J 15:2043–2056.

104. Dulbecco R, Vogt M. 1954. PLAQUE FORMATION AND ISOLATION OF PURE

LINES WITH POLIOMYELITIS VIRUSES. Journal of Experimental Medicine 99:167–

182.

105. Reeburgh WS. 1967. An Improved Interstitial Water Sampler. Limnology and

Oceanography 12:163–165.

106. Cline JD. 1969. Spectrophotometric Determination of Hydrogen Sulfide in Natural

Waters. Limnology and Oceanography 14:454–458.

107. Coplen TB, Brand WA, Gehre M, Gröning M, Meijer HAJ, Toman B, Verkouteren RM.

2006. New Guidelines for δ13C Measurements. Anal Chem 78:2439–2441.

108.   Standard Test Methods for Density and Specific Gravity (Relative Density) of Plastics by

Displacement. https://www.astm.org/d0792-20.html. Retrieved 19 September 2024.

109. Giesche H. 2006. Mercury Porosimetry: A General (Practical) Overview. Particle &

Particle Systems Characterization 23:9–19.

110. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a

Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence

Data on the MiSeq Illumina Sequencing Platform. Appl Environ Microbiol 79:5112–5120.

111. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. 1. EMBnet.journal 17:10–12.

112. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H,

Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn

CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R,

Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M,

Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K,

Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S,

Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D,

Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E,

Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV,

Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF,

Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB,

Rivers A, Robeson MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ,

Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-

Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M,

Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ,

Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG. 2019. Reproducible, interactive,

scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 37:852–

857.

113. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods

13:581–583.

114. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Research 41:D590–D596.

115. Mayr MJ, Orphan VJ. 2024. Extraction of microbial cells from methane-seep carbonate

rocks for single-cell analyses (e.g. BONCAT-FISH) using "combination buffer" and

percoll density centrifugation https://doi.org/10.17504/protocols.io.rm7vzjk55lx1/v1.

116.   Eichorst SA, Strasser F, Woyke T, Schintlmeister A, Wagner M, Woebken D. 2015.

1356   Advancements in the application of NanoSIMS and Raman microspectroscopy to

1357   investigate the activity of microbial cells in soils. FEMS Microbiol Ecol 91:fiv106.

1358   117.   Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. 1990.

1359   Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for

1360   analyzing mixed microbial populations. Applied and Environmental Microbiology

1361   56:1919–1925.

1362   118.   Daims H, Brühl A, Amann R, Schleifer K-H, Wagner M. 1999. The Domain-specific

1363   Probe EUB338 is Insufficient for the Detection of all *Bacteria*: Development and

1364   Evaluation of a more Comprehensive Probe Set. Systematic and Applied Microbiology

1365   22:434–444.

1366   119.   Westram R, Bader K, Prüsse E, Kumar Y, Meier H, Glöckner FO, Ludwig W. 2011. ARB:

1367   A Software Environment for Sequence Data, p. 399–406. *In* Handbook of Molecular

1368   Microbial Ecology I. John Wiley & Sons, Ltd.

1369   120.   Stahl DA, Amann R. 1991. Development and application of nucleic acid probes in bacterial

1370   systematics, p.205-248. *In* Stackebrandt, E. and Goodfellow, M., Eds., Nucleic Acid

1371   Techniques in Bacterial Systematics, John Wiley & Sons Ltd.

1372   121.   Dieterich DC, Link AJ, Graumann J, Tirrell DA, Schuman EM. 2006. Selective

1373   identification of newly synthesized proteins in mammalian cells using bioorthogonal

1374   noncanonical amino acid tagging (BONCAT). Proceedings of the National Academy of

1375   Sciences 103:9482–9487.

1376 122. Sletten EM, Bertozzi CR. 2009. Bioorthogonal Chemistry: Fishing for Selectivity in a Sea

1377   of Functionality. Angewandte Chemie International Edition 48:6974–6998.