Title: Elusive tropical forest canopy diversity revealed through environmental DNA contained
in rainwater

Short title: Rain eDNA to monitor rainforest biodiversity

Author list:

Lucie Zinger[1,†]*, Anne-Sophie Benoiston[1], Yves Cuenot[1], Céline Leroy[1,3-4], Eliane Louisanna[4],
Lucie Moreau[1], Frédéric Petitclerc[4], Finn Piatscheck[4], Jérôme Orivel[1,4], Cécile Richard-
Hansen[4,5], Lou Hansen-Chaffard[5], Uxue Suescun[1], Valérie Troispoux[4], Frédéric Boyer[6],
Jérôme Chave[1], Thibaud Decaëns[7], Antoine Fouquet[1], Johan Pansu[8], Jérémy Raynaud[1],
Rodolphe Rougerie[9], Lucas Sire[9], Heidy Schimann[10], Pierre Taberlet[6,11], Amaia Iribar[1]*

Affiliations

[1] Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), UMR 5300 CNRS,
IRD, INP, UT3, Toulouse, France.

[3] AMAP, Univ Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France.

[4] UMR Ecologie des Forêts de Guyane (EcoFoG), INRAE, CNRS, AgroParisTech, CIRAD,
Université de Guyane, Université des Antilles, Campus agronomique, Kourou, France.

[5] OFB-Office Français de la Biodiversité/DRAS Campus agronomique de Kourou

[6] Laboratoire d'Ecologie Alpine (LECA), UGA, UMR 5553 CNRS, University of Savoie Mont
Blanc, Grenoble, France.

[7] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

[8] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA,
Villeurbanne, France.

[9] Institut de Systématique, Évolution, Biodiversité (ISYEB), UMR 7205 Museum National
d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE-PSL, Université des Antilles, Paris,
France

[10] Biodiversité, Gènes et Communautés (BIOGECO), INRAE, Université de Bordeaux,
Cestas, France

[11] PT: UiT – The Arctic University of Norway, Tromsø Museum, Tromsø, Norway.

[†] former address: Instituto Tecnológico Vale, Rua Boaventura da Silva 955, 66055-090,
Belém, Pará, Brazil.

*Correspondence: lucie@zinger.fr, amaya.iribar-canseco@cnrs.fr

ORCID:

Lucie Zinger: 0000-0002-3400-5825

Anne-Sophie Benoiston: 0000-0001-9446-5703

Yves Cuenot: 0000-0002-9755-811X

Céline Leroy: 0000-0003-4859-8040

Eliane Louisanna: 0000-0002-2082-5726

Lucie Moreau: 0000-0002-3859-0435

Frédéric Petitclerc: 0000-0002-2015-1258

Finn Piatscheck: 0000-0002-1796-6288

Jérôme Orivel: 0000-0002-5636-3228

Cécile Richard-Hansen: 0000-0002-9353-791X

Valérie Troispoux: 0000-0002-1118-1367

Frédéric Boyer: 0000-0003-0021-9590

Jerome Chave: 0000-0002-7766-1347

Thibaud Decaëns: 0000-0003-0915-3882

Antoine Fouquet: 0000-0003-4060-0281

Johan Pansu: 0000-0003-0256-0258

Rodolphe Rougerie: 0000-0003-0937-2815

Lucas Sire: 0000-0002-2911-6109

Heidy Schimann: 0000-0002-9139-920X

Pierre Taberlet: 0000-0002-3554-5954

Amaia Iribar: 0000-0002-4709-648X

Abstract

Tropical rainforest canopies shelter an under-explored reservoir of biodiversity. In recent years, the amplification and sequencing of taxonomically informative DNA fragments from environmental samples (i.e. eDNA) has revolutionized biomonitoring. Here, we explore the potential of DNA contained in canopy throughfall water to sample the biological diversity of rainforest canopies. By sampling rainwash eDNA in two 1ha-plots, one mature Amazonian forest and a nearby tree plantation, we were able to detect 170 plant taxa, 72 vertebrate taxa including mammals, birds, and amphibians, and 313 insect taxa including e.g. mosquitoes, ants, beetles. The taxonomic composition retrieved in these two plots reflected their different disturbance status, and the plantation exhibited a much lower diversity. We also found that rainwash eDNA persists over ten days in passive eDNA collectors while providing a local picture of the diversity. This suggests that this method can be scaled up for a cost-effective environmental management of rainforest', and more generally of all forests canopies.

Teaser

Rainwash DNA reveals the identity of plants, animals, and insects diversity, offering a practical tool for monitoring canopy biodiversity.

Keywords

Rainwater, environmental DNA, metabarcoding, tropical forests, conservation biology, terrestrial ecology, canopy biodiversity

Introduction

Tropical rainforests are indispensable components of Earth's systems, playing critical roles
in maintaining global biogeochemical cycles and supporting human well-being through
essential services such as food provisioning, water quality regulation, and global climate
regulation (*1*). Tropical rainforests also harbour a tremendous and unique reservoir of
biodiversity, a vast part being largely unexplored (*2, 3*). Within these ecosystems, the forest
canopy is a particularly dynamic and significant zone, representing the active interface
between the terrestrial biosphere and the atmosphere, and a recognized hotspot of
biodiversity (*4*). However, canopies' inaccessibility has historically limited their scientific
exploration, leaving substantial knowledge gaps. With the increasing pressures tropical
rainforests face, including deforestation, biological invasions, and climate change (*5–7*),
improving methods to monitor rainforests biodiversity, and in particular that of their canopies
is now a pressing societal demand.

In recent years, the amplification and sequencing of taxonomically-informative DNA
fragments from environmental samples (*8–10*), i.e. eDNA metabarcoding, has revolutionised
biomonitoring across all domains of life. Environmental matrices such as soil (*9, 11*),
invertebrates bulk (*12, 13*), marine and freshwater (*14–16*), or even air (*17–19*), are being
commonly sampled environmental matrices for such purposes. However, these
environmental matrices present several caveats for the sampling of terrestrial aboveground
biodiversity (*20*). Specifically, the DNA found in soil samples is dominated by that of
belowground organisms and overall transport poorly, making the probability of detecting
aboveground organisms very low (*9, 20*). The ingested DNA (i.e. iDNA) contained in
hematophagous and/or phytophagous arthropod bulks can effectively sample vertebrates or
plant communities (*12, 13*) but this approach is destructive and hence not desirable. Filtering
of air (*17–19*) or of water from local streams or lakes (*14–16*) is less invasive, but present
several uncertainties related to the spatio-temporal window of the signal retrieved (*21, 22*).

Organisms inhabiting the canopy naturally release DNA through tissue renewal and
excretion, releasing cellular debris, urine, feces, and other materials that can adsorb onto the
vegetation surfaces, creating a valuable eDNA source for biomonitoring. As such, analysis of
eDNA from leaf swabs has been shown as a promising approach to detect terrestrial
vertebrates (*23*). However, translating this approach to large-scale biodiversity assessments
in tropical rainforests necessitates a considerable sampling effort, demanding alternative
sampling protocols of this material. The sampling of canopy throughfall water represents
such an alternative, as rainwash water transports and concentrates the DNA adsorbed on
canopy leaves and branches. Accordingly, both experimental and *in situ* studies have
demonstrated that it was possible to detect canopy arthropod species directly from rainwash
water collected directly below the canopy (*24, 25*). Rainwash waters therefore appear as
extremely relevant for implementing large-scale eDNA-based monitoring of tropical forest
canopies, these ecosystems being further characterised by an equatorial rainfall regime, with
abundant rainfall throughout the year, albeit seasonal drier periods.

Here, we assessed the potential of the DNA contained in rainwash water collected under the
canopy for the monitoring of plants, vertebrates and insects diversity in tropical rainforests
(Figure 1). To this end, we conducted two sampling campaigns in two contrasted amazonian
forests: one old growth forest, and a rubber (*Hevea brasiliensis*, Euphorbiaceae) and
rosewood (*Aniba rosaeodora*, Lauraceae) tree plantation (Figure 1a). To collect canopy

rainwash water, we repurposed umbrellas (Figure S1) and equipped them with four types of
passive eDNA samplers (Figure 1b). The first campaign, hereafter referred to as the
"temporal experiment", aimed at assessing the temporal dynamics of rainwash eDNA in
terms of (i) diversity/composition in Molecular Operational Taxonomic Units (here after
referred to as MOTUs) through DNA metabarcoding (Table S1), and of (ii) eDNA persistence
through quantification of an exogenous DNA pulverised on canopy leaves at the start of the
experiment, using both digital PCR (dPCR) and DNA metabarcoding. This exogenous DNA
was obtained from carrot, this species, *Daucus carota*, belonging to Apiaceae, a family
absent from the studied area. The second campaign, hereafter referred to as the "spatial
experiment", aimed at assessing the spatial heterogeneity of the signal and sampling
coverage needed to maximise the representativeness of the plot-scale diversity retrieved by
rainwash eDNA. To this end, we collected rainwash eDNA every 20 meters in each 1ha plot
using the most effective eDNA passive sampler tested during the first sampling campaign.

134 Results and Discussion

135 Rainwash eDNA captures canopy diversity

We first asked what was the overall taxonomic composition of the signal yielded by rainwash
eDNA. Considering all samples from the temporal experiment, we were able to retrieve a
total of 562 taxa (i.e. identified taxa - to any rank from phylum to species - regardless of the
sequence similarity against its closer match in Genbank). Of these, the plant and vertebrate
taxa identified to a fine taxonomic level are known to be present in South America (Figure 2,
Table S2). For plants, we detected a total of 170 taxa, encompassing 50 orders and 56
species. These figures are low for a tropical forest, and can be due to the overall low
resolution of the plant DNA marker used here, a portion of the *rbc*L gene. Thirteen of these
taxa corresponded to mosses and liverworts, 14 to ferns including the arborescent ferns
family, 3 to gymnosperms, and 16 to green algae. The remaining taxa corresponded to
flowering plant families known to occur in the study area, such as Lecythidaceae,
Melastomataceae, Fabaceae, Chrysobalanaceae, Vochysiaceae*,* Euphorbiaceae*,*
Annonacae and, in much lower numbers, angiosperm epiphytes such as Bromeliaceae or
Orchidaceae.

Regarding vertebrates, after careful curation of common contaminants (human, beef and
pig) and remaining tag-jumps (*26, 27*) originating from positive controls, we retrieved a total
of 72 taxa (spanning 21 vertebrate orders). These included 11 amphibian taxa, including the
frog *Leptodactylus nesiotus*, known to occur in the area (*28*), 24 bird taxa (including e.g.
parrots, toucans), four squamate taxa, and 32 mammal taxa. Most mammals corresponded
to opossums and rodents, but also comprised two genera of neotropical monkeys (i.e.
Saguinus and *Pithecia*), one genera of sloth (*Choloepus*) and of anteater (*Cyclopes*), two
genera of bats (*Carollia* and *Rhinophylla*), and two genera of procyonids (i.e. *Potos* and
Nasua). All taxa corresponded to species strictly or occasionally arboreal (*29*). We also
detected two fish taxa. Fishes have also been detected on leaf swabs (*23*) but since it is
unclear whether these corresponded to sporadic contamination or to the digestion product of
another vertebrate, we excluded them from the analysis. Similarly, we detected sporadically
the domestic cat. While it may correspond to true presences - the two plots being close from
populated areas, we adopted a conservative filtration procedure and excluded these
sequences from the analysis (see Methods).

For insects, we retrieved a total of 313 taxa, representing 20 orders. We also found 5 non-
insect arthropod taxa (i.e. 2 spiders, 2 springtails, and one isopod). The most represented
insect orders (in terms of diversity of taxa) were Coleoptera, Diptera (including several
species of fruit- and sandflies, mosquitoes), Hymenoptera and Lepidoptera, which are also
the most diverse orders in these ecosystems. Other insects such as cockroaches, termites,
true-bugs, grasshoppers or dragonflies were also identified. Some of the taxa identified at a
fine taxonomic are not usually found in the neotropical realm. However, the sequence
identity scores with the closest matches in Genbank were overall quite low (i.e. below 90%)
for insect taxa identified at either fine or coarse taxonomic resolution. This finding
exemplifies the large incompleteness of the reference database and the resulting difficulty in
achieving accurate taxonomic assignments, particularly for understudied tropical insect
groups, and could explain at least partly why the number of retrieved arthropod taxa here
was low compared to expectations for such environments (*30*).

We also investigated the relevance of the approach from a conservation point of view. Few
taxa were in the vulnerable to critically endangered status, but a careful consideration of
their identity and of their distribution range reveals that they do not occur in French Guiana,
and most likely correspond to the mis-assignment of sister taxa occurring locally but that are
not referenced in DNA databases, leading to an erroneous taxonomic assignment at the
species level (i.e. the local species belonging to the genera *Campephilus, Dipteryx,* and
Swartzia species were wrongly assigned to *Campephilus imperialis, Dipteryx alata* and
Swartzia bahiensis, respectively). This illustrates how important it is to carefully verify the
taxon list retrieved from eDNA surveys, as well as the need for pursuing efforts for
populating DNA reference databases, particularly for tropical diversity, which remains largely
under-represented (*20, 31*). The rest were either found without IUCN status or were explicitly
not evaluated, thus exemplifying the usefulness of rainwash eDNA to document the
distribution of data deficient taxa, and its relevance for conservation efforts since over half of
data deficient species are projected to face extinction (*5*).

Finally, we assessed the ecological relevance of the signal retrieved. We found the
taxonomic diversity and composition in the two types of forest to be consistent with our
expectations. First, the taxonomic diversity was 1.3 to 1.9 times higher in the old-growth
forest plot depending of the taxonomic group, with 155 plant taxa (vs. 111 in the plantation),
61 vertebrate taxa (vs. 32) and 276 (vs. 153) insect taxa. The retrieved taxonomic
composition also mirrored the contrasted disturbance status of the two forest plots (Figure
2). The rubber plantation plot yielded a higher proportion of reads assigned to family
Euphorbiaceae (to which the rubber tree belongs), with its most representative MOTU
corresponding to the rubber tree itself (*Hevea,* Table S2). We also detected greater
abundance of termites belonging to the genus *Coptotermes*, to which the rubber termite
belongs. In the plantation plot, we also detected a higher relative abundance of Lauraceae
(to which rosewood belongs). We also found a higher proportion of reads assigned to
Fabaceae, Melastomataceae, Hypericaceae*,* and Piperaceae, as well as to the frog
Leptodactylus nesiotus. All these taxa are often associated with disturbed or more open
habitats (*28, 32*). For insects, we also found a higher relative abundance of mosquitoes, ants
and cockroaches in the rubber plantation. Conversely, the old-growth forest plot harboured
higher relative abundance of Lecythidaceae, Chrysobalanaceae, Vochysiaceae*,* and
Caryocaraceae, as well as the frog *Trachycephalus hadroceps*. These taxa are typically

found in old-growth forests (*32, 33*). The old-growth forest also harboured more MOTUs and
sequences assigned to phytophagous taxa such as Lepidoptera and Coleoptera.

Optimising the sampling of rainwash eDNA

Making rainwash eDNA-based biomonitoring operational for routine analyses by
environmental stakeholders requires developing sampling devices and strategies (i) that are
cost effective and - if possible - low-tech for higher transferability, (ii) that limit the number of
sampling campaigns while (iii) being able to generate biodiversity inventories that are
spatially and temporally representative of the study area. This assessment is even more
important for an effective implementation in tropical countries, where challenges of
accessibility to sites are more important.

Passive collection of eDNA

We developed a low-cost eDNA collection method where rainwash eDNA accumulates
passively. This can be done either by filtration, as commonly done in waterbodies (*14–16*)
but here filtration was not performed actively by pumping, but passively by gravity. Passive
eDNA collection can also be done through passive fixation of eDNA molecules on different
types of matrices (*34, 35*) (Figure 1b). We assessed the efficiency of five passive eDNA
samplers in fixing rainwash eDNA after 1 day of immersion in rainwash water. First, we
compared the MOTU diversity retrieved by the different types of passive eDNA samplers,
here estimated as Hill numbers set at $q = 1$, i.e. a number of equivalent species
downweighting rare MOTUs. This index tends to the exponential of the Shannon index. We
chose it because it is more robust for DNA metabarcoding data, due to its lower sensitivity to
sequencing depth (Figure S2) or sampling effort, but also to other molecular low-abundance
artefacts (*36–39*). Second, we compared the retrieved amount of carrot spike-in DNA across
eDNA sampler types (Figure 1a), either in terms of copy number of a single-copy gene, *Dau*
c1, quantified through digital PCR (dPCR), or based on the average number of Apiaceae
sequencing reads retrieved in metabarcoding data. We considered these two measures here
because while they correlated well (*40*) (Figure S3), they also have their own *pros* and *cons*
(see Methods).

Filtration by gravity with Waterra filters captured higher MOTU diversity for plants and to a
lesser extent for vertebrates compared to the other passive captors (Figure S4a, Table S3).
A similar trend was observed for invertebrates, but was not significant. This collection
method was also more effective in fixing the carrot spike-in DNA, followed by cotton strips
after 1 day of immersion in rainwash water (Figure S4b, Table S3), but also for longer
immersion periods (Figure S5). Hence, we recommend to use Waterra filters with filtration by
gravity to maximise the sampling of rainwash eDNA. Fixation on cotton strips, which is more
affordable, could be considered as an effective alternative collection method, but will require
greater fixation surface and/or more passive sampler replicates per rainwash water samplers
to reach a level of detection similar to passive filtration with Waterra filters.

Temporal coverage of rainwash eDNA

Sampling tropical diversity effectively with rainwash eDNA also requires identifying the
temporal sampling window that maximises the diversity retrieved due to an accumulation of
rain events (Figure 1a), and of eDNA molecules coming from the canopy, but at the same
time without losing too much information due to eDNA decay. Here, we evaluated this

property on the Waterra filters obtained from the temporal experimental setup (Figure 1a) by
considering three criteria: MOTU diversity dynamics, MOTU compositional temporal
turnover, and spike-in DNA decay.

We analysed how MOTUs diversity of each target clade varied with the duration of the eDNA
collection. For this specific analysis, we measured MOTU diversity at $q = 0$ (i.e. MOTU
richness) on rarefied data because of the higher sensitivity of this index to subtle changes in
species diversity and composition that would be caused by rare species (*38*), in our case of
MOTUs becoming rarer due to eDNA decay. We fitted several functions modelling
population dynamics (*41*) on the temporal trends of MOTU diversity and found that the
Ricker function, which exhibits an optimum, was the best fit (Figure 3a). The fact that the
temporal dynamics of MOTU diversity exhibits an optimum can be attributed to the reduced
influxes of new eDNA molecules through time resulting from fewer and less intense rain
events (Figure 1a), coupled with a stable eDNA degradation. Based on this model, we
estimated the initial MOTU accumulation rate, which here ranged between 5 and 10 MOTUs
per day after installation, as well as the time at which the diversity is maximal (Figure 3b,
Table S4), here between 8 to 20 days after the installation of rainwash eDNA samplers.

We also identified a time-decay pattern, i.e. a decrease of MOTUs compositional similarity
as the duration between temporal sampling points increases (Figure S6). This supports the
idea that the observed diversity dynamics of MOTUs is a balance between eDNA
persistence in the matrix and influx of new molecules. However, this relationship was weaker
for animals, and statistically non-significant for insects. We explain this by the fact that (i)
several samples collected at close dates already exhibited very few to no shared animal
MOTUs and that this phenomenon would be exacerbated using rarefied data, and (ii) all
target clades exhibited a lower MOTUs diversity for the largest temporal intervals between
sampling points (Figure 3). These properties may artificially inflate the weight of the MOTUs
shared between samples in the similarity computation and of the similarity value itself, thus
flattering the distance-decay pattern.

Finally, we estimated the decay rate of the carrot spike-in DNA using a negative exponential
function (*42*), from which we estimated the time of decay to 5% of its initial quantity to about
29 days (see Methods, Figure 3a-b, Table S4).

We conclude that our passive sampling of rainwash eDNA can integrate relevant signal from
8 to 40 days depending on the initial amount of eDNA at the site, making it less punctual
than one-shot samplings, as it is the case for the sampling of soil, water, and air or leaves
swabs, which needs to be repeated. This property also makes the implementation of the
protocol more feasible for routine analyses, either by reducing the need to frequent onsite
visits, and/or by implementing monthly sampling for real-time monitoring, hence providing
valuable insights into temporal trends and potential shifts in the studied area. These results
are a first assessment of rainwash eDNA dynamics in real-life conditions, even though they
miss processes involved in eDNA dynamics in general (*43*), such as the type from which the
DNA detected originates (e.g. tissues, faeces, pollen). The frequency and intensity of rain
events is an obvious important driver of these dynamics (see an example below), limiting the
relevance of deploying the approach in areas that receive lower precipitation annually or
during certain times of the year, such as seasonally dry tropical forests. We therefore believe
that our estimate of rainwash eDNA decay should be assessed more systematically using a
similar, simple approach with an exogenous DNA easy to obtain and monitor as we did here.

Spatial distribution of rainwash eDNA

Finally, we assessed the spatial sampling effort required to sample diversity within one 1 ha
plot. In our spatial experiment (Figure 1), rainwash eDNA was sampled every 20 metres in
each plot during 40 days. Precipitations were strongly reduced during this experiment,
resulting in an overall reduced detection power, especially for vertebrates. While we were
able to detect several vertebrates taxa known to occur locally (e.g. *Saguinus*, *Oecomys*, or
Ramphastos), the level of contamination was quite high. This illustrates the fact that
infrequent or single rain events (artificial or not) capture insufficient amounts of eDNA for
robust downstream analyses. Here, we adopted a conservative approach by restricting our
spatial analysis to plant and invertebrate data, for which we built rarefaction curves of MOTU
diversity (*sensu* Gotelli & Colwell [44], i.e. the average diversity observed when increasing
the number of samples), this time with $q=1$ (i.e. exponential of the Shannon index) to obtain
a more robust estimate of the plot-scale diversity [36–39] (Methods, Figure 4).

These rarefaction curves confirmed our previous observation (Figure 2) that the diversity
recovered from rainwash eDNA is significantly higher in the old-growth forest for the targeted
clades. This illustrates again the relevance of rainwash eDNA to detect differences of
diversity across ecosystem types.

We then fitted species rarefaction models to the rarefaction curves [45] (Methods, Figure 4),
from which we estimated the plot-scale diversity, the diversity covered by our current
sampling effort (i.e. n=16 sampling points), as well as the number of sampling points
required to cover 70% and 90% of the plot-scale diversity (Figure 4, Table S6). We found
that these figures varied greatly across target clades, the plant diversity being relatively well
covered (i.e. > 70%) with our current sampling effort. By contrast, we covered at best 46% of
the insect plot-scale diversity detectable with our approach. While we do not exclude that
these differences are partly due to the lower resolution of our markers [9], this result remain
consistent with the tremendous diversity of insects usually described in tropical forests and
at comparable spatial scales in comparison to that of plants [30]. Reaching 90% of the plot-
scale diversity would require at least 40 samples for plants, and much more for invertebrates
(Figure 4, Table S6).

This suggests that the signal retrieved from rainwash eDNA, while acceptable to describe
the diversity of non rare MOTUs at the plot scale , remains local, similarly to what is reported
for soil eDNA [20] or from temperate rainwash eDNA [25]. This conclusion is confirmed by
the overall low to null MOTU compositional similarity between sampling points as well as the
absence of a clear distance-decay pattern (Figure S7).

332   Advantages of rainwash eDNA to monitor tropical rainforests

Our findings demonstrate that rainwash eDNA is a promising material for the monitoring of
biodiversity in the tropical forest canopy by overcoming the challenges of working in a
spatially heterogeneous environment both horizontally and vertically [46–48], and where
eDNA is not transported as easily as in aquatic ecosystems. Our results show that to sample
aboveground biodiversity, rainwash eDNA provides an effective non-destructive alternative
to iDNA retrieved from arthropod bulks that has been also used for the same purpose[12,
13]. They also show that it offers a better representation of terrestrial above-ground
biodiversity than soil eDNA, the latter being dominated by the eDNA of plants and soil

organisms (*9, 20*). Sampling the two matrices could hence provide a holistic view of both
above- and belowground diversity.

Water eDNA from streams or lakes have been shown to provide useful information on
terrestrial diversity (*14–16*). However, this material can only be collected in the presence of
water bodies. In addition, water eDNA is transported across highly variable distances
depending on a multitude of factors (e.g. particle size distribution, water body
hydrogeomorphological properties, eDNA decay rate, etc.) (*22*). The water eDNA signal is
also temporally variable, not only due to the demographic changes in the target biological
community *per se*, but also because the fluctuation in frequency and intensity of rainfall
events determines terrestrial material transport into water bodies (*49*). While also subjected
to fluctuations in rainfall events, rainwash eDNA exhibits a localised spatial distribution
because it is sampled closer to its source community. In addition, our study demonstrates
that the community composition is detectable for at least two weeks. This suggests that the
spatial dynamics of rainwash eDNA is easier to interpret than stream/lake water eDNA, and
could offer a more localised and reliable approach for terrestrial biodiversity monitoring.

Air eDNA has also shown promise to describe terrestrial biodiversity, in particular that of
vertebrates. However, current proof of concepts have been mostly conducted in zoos or
similar systems with unvegetated soils - leading to high amounts of particles in the air,
and/or harbouring a low diversity, and yet a high density of animals(*18, 19, 23, 50*). These
conditions facilitate animal detection and are not typical of tropical forests, where both
animal and air particle density are expected to be much lower. In contrast, our assessment
of rainwash eDNA has been done in natural conditions. In addition, air eDNA is likely to
depend on the same spatio-temporal uncertainties as of stream/lake water eDNA (e.g.
winds, rainfall events, etc) (*21*). Further experiments in natural conditions will be necessary
to assess the advantages and/or complementarity of the two environmental matrices.

Apart from the limitations linked to molecular biases and contaminations that are common to
all eDNA-based assessments (*9, 51*), an important limit of our approach is the dependance
to rainfall events. Given the persistence of rainwash eDNA found in moist tropical forests,
this problem appears negligible for its appropriate sampling. However, it makes the
approach less relevant for forests receiving less and irregular precipitation, such as tropical
dry forests, cerrado forests or caatinga forests, where our approach would be relevant only
during the wet season. On the other hand, we expect this approach to be very promising in
temperate forests, where diversity is lower and where DNA will likely persist longer through
time.

In conclusion, our study demonstrates the potential of rainwash eDNA as a powerful tool for
monitoring the diversity of aboveground tropical forests' plants and animals, which remains
difficult to study with traditional and other eDNA-based methods. Our work establishes a
robust methodological framework for an effective sampling of rainwash eDNA, and opens
the door for further refinements for both community-level census and detection of patrimonial
or human/environmental health importance. By enabling large-scale and non-invasive
monitoring, this method can inform or assess tropical forests' conservation strategies, and
contribute to the protection of these irreplaceable ecosystems.

Methods

Study area

The study was conducted in the municipality of Sinnamary, which features two types of forest: an old-growth forest and a tree plantation. The equatorial climate in the studied area is characterised by two climatic periods: a long rainy period from mid-December to mid-August often interrupted by a short drier period between March and April and a longer dry season for the rest of the year. In this area, the annual rainfall is 3,041 mm with mean air temperature of 26°C (±1-1.5°C across the year). The 1 ha plot (plot P16) of old-growth forest is located in the Paracou Research Station (*52*) (lat: 5°18′N, lon: 52°53′W, https://paracou.cirad.fr/website). The Paracou research station managed by the French Agricultural Research Centre for International Development (CIRAD) comprises 5,000 ha of tropical rainforest in a hilly *terra firme* region. This site is a long-term research station where plant communities are well characterised and precipitation is monitored daily (Figure 1a). The 1 ha plot of tree plantation is located in a former rubber tree (*Hevea brasiliensis*, Euphorbiaceae) plantation also managed by CIRAD, where rosewood (*Aniba rosaeodora*, Lauraceae) was more recently introduced and of which the exact current floristic composition is unknown. It is located 5 km away from the Paracou research station (lat: 5°17'N, lon: 52°54'W).

eDNA sampling

Temporal experiment

To collect rainwash eDNA, we repurposed conventional umbrellas and plastic bottles into a low-cost rainwash water sampler with integrated passive eDNA sampler holders (instructions described in Figure S1). We equipped these samplers with four types of eDNA passive captors. First, we used Waterra eDNA filter (600 cm^2 polyethersulfone filter, pore size of ⌀ 0.45μm, Waterra USA Inc.), which are commonly used for eDNA sampling in aquatic ecosystems (*53*) and that we here consider as a reference. We plugged the Waterra filters at the bottom of the water tank of the rainwash water sampler with silicon tubes to passively collect eDNA through gravity filtration (*54*) (Figure 1b). We also tested 3 alternative, cost-effective passive captors based on eDNA fixation. First, we considered Mixed Cellulose Ester membrane filters (⌀ 47 mm, pore size of 0.45μm, Merck KGaA, Darmstadt, Germany), which have recently proved effective for studying aquatic biodiversity with eDNA (*34*). We also considered ceramic mosaics of 25x25 mm, as ceramic has DNA-binding properties, even over long time periods (*55, 56*). Finally, we considered cotton strips of 25x25mm, which also have the potential to passively sample aquatic eDNA (*57*). Ceramics and cotton strips have also long been used to measure biofilm colonisation/productivity rates (*58*) and decomposition (*59, 60*), respectively. As biofilms also have eDNA-fixing properties (*35*), this property may increase the efficiency of these passive eDNA samplers for our purpose. All passive eDNA samplers were placed into the rainwash collector water tank.

A total of 5 rainwash water samplers were installed in each 1 ha forest plot at randomly selected locations away from the plot edges. We assessed the efficiency of the passive captors and quantified eDNA persistence through time by introducing an exogenous DNA just before the installation of the rainwash water samplers. We chose carrot (*Daucus carota*, *Apiaceae*) as a spike-in due to its widespread availability and because this plant family is

absent from the study area and more generally from the Guiana Shield (except the aromatic
plant *Eryngium foetidum*, which is not cultivated near the study sites (*61*)). To do so, we
pulverised about 10 mL of carrot juice on canopy leaves at each sampling point with a
conventional streamer. About 20 minutes after pulverisation, we installed below pulverised
canopy leaves our rainwash water samplers, and then simulated an initial rain by pulverising
around 40 mL of water (made of rainwater collected on site a few days before the start of the
experiment (2:5) complemented with bottled springwater (3:5) to obtain a sufficient volume).
eDNA passive captors were sampled after 1, 5, 10, 20 and 40 days for subsequent
molecular analyses and placed in 50 mL sterile falcon tubes (Waterra filter excepted). We
then added 50 mL of Longmire buffer (for 1L of buffer: 100mL 1M Tris, 100ml 1M EDTA,
2mL 5M NaCl and 798 mL of molecular biology grade H20) in each falcon tube and Waterra
filter to store and preserve eDNA passive captors at room temperature (*62*) before molecular
analyses. We thus obtained a total of 200 samples (i.e. 2 plots, 5 sampling points, 5
sampling dates, 4 passive eDNA samplers) for the temporal experiment.

Spatial experiment

In the same plots, we conducted another sampling campaign in mid February 2024 where
we placed one rainwash water sampler every 20 metres, resulting in 16 points per plot.
Following the results obtained in the temporal experiment, the water samplers were
equipped with one Waterra filter each. The sampling devices were left at the field for 40 days
(Figure 1). We thus obtained a total of 32 samples for the spatial experiment.

Molecular analyses

DNA extractions

Each falcon tube containing a passive eDNA sampler immersed in Longmire buffer was
shaken with a horizontal shaker at 40 Hz for 10 min solubilize the DNA, after what we
removed the captor matrix. We also shaked the Waterra filters' samples at 40 Hz for 10 min,
after what we transferred the buffer into 50 mL falcon tubes. We then added 33 mL of
absolute ethanol to all tubes and subsequently stored them at -20°C overnight. The following
day, the tubes were removed from the freezer, left for 30 min at room temperature, and
centrifuged at 8500 for 15 minutes at 15°C. The supernatant was then discarded, and 720µL
of ATL buffer (QIAGEN; Hilden, Germany) was added to each pellet. The mixture was then
transferred to 2 mL tubes, to which we added 20 µL of proteinase K (QIAGEN; Hilden,
Germany), followed by an incubation at 56°C for 2 hours for digestion. NucleospinSoil®
Macherey-Nagel kit was used, following the protocol from step 6 (adjust binding conditions)
onward. The resulting DNA extracts were finally analysed using a Nanodrop ND-1000
(Thermo Fisher Scientific, Inc.) to determine the yield and purity of the DNA.

DNA metabarcoding

We characterised the community composition of plants, arthropods and vertebrates through
eDNA metabarcoding, using the Sper03 marker (targeting a subregion of the rbcL gene, fwd
5'-CACCACAAACAGARACTAAAARC-3', rev 5'-TCCAYACRGTTGTCCATGTACC-3') for
plants, the Inse01 marker (*9*) (fwd 5'-RGACGAGAAGACCCTATARA-3', rev: 5'-
ACGCTGTTATCCCTAARGTA-3') for insects, and the Vert01 marker (fwd 5'-
TTAGATACCCCACTATGC-3', rev 5'-TAGAACAGGCTCCTCTAG-3') for vertebrates,
together with a human-blocking primer:

CTATGCTTAGCCCTAAACCTCAACAGTTAAATCAACAAAACTGCT-Spc3 (*63*). PCR
reactions were performed in triplicate for each sample in 20 µL. Each reaction consisted of
10 µL of Amplitaq Gold 360 Master Mix (Applied Biosystems), 2 µL of a mix of fwd and rev
primers (0.5 µM each final), 6 µL of Nuclease-free water, and 2 µL of eDNA diluted 10 times
beforehand. For each PCR reaction, primers were labelled with unique combinations of 8-nt-
long tags at the 5′ end of the forward and reverse primers (*64*) to distinguish PCR replicates
after sequencing. PCR thermocycling conditions are indicated in Table S1. The final
molecular experimental design thus included for each marker 600 PCR reactions
corresponding to samples, 33 corresponding to extraction blank controls, and 30 to PCR
blank controls for the temporal experiment. For the spatial experiment, it contained 96 PCR
reactions corresponding to samples, 8 to extraction blank controls, and 3 to PCR blank
controls. All PCRs of the same marker were pooled in equal volumes without standardising
the PCR products concentration to allow a more quantitative comparison of the abundance
of taxa sequencing reads across samples (*40*) (see also Figure S3). This pool was then used
to build sequencing libraries using the Illumina Truseq Nano HT (Illumina), following the
manufacturer protocol but without PCR enrichment of ligation products to minimise the
formation of tag-jumps (*65*). Sequencing was conducted at the GetPlage sequencing
platform (Toulouse, France) for the temporal experiment, and at Novogene (Cambridge, UK)
for the spatial experiment, with the characteristics indicated in Table S1.

Bioinformatics and data preparation

Primary bioinformatics/curation procedures, i.e. reads pairing, demultiplexing, dereplication,
exclusion of obvious artefacts and singletons, and clustering of sequences to form Molecular
Operational Taxonomic Units (MOTUs) were made with the `OBITools v1.2.11`
(https://anaconda.org/bioconda/obitools, (*66*)) and the `sumaclust v1.0.31` clustering
algorithm (https://anaconda.org/bioconda/sumaclust, (*67*)), using a snakemake workflow
(https://github.com/AnneSoBen/obitools_workflow). The taxonomic assignment was done
with the `ecotag` program (*66*) with the reference databases indicated in Table S1. These
reference databases were built with the `ecoPCR` program (*68*) from Genbank. Next, we
conducted a secondary curation procedure on sequencing data with the `metabaR R`
package (https://github.com/metabaRfactory/metabaR, (*69*)), in order to further curate the
data from contaminants, tag-jumps or other degraded sequences. Parameters used in this
process are summarised in Table S1. Associated codes and diagnostic plots/statistics are
available (Key resource Table). Finally, for vertebrates and plant data, we also verified that
the taxon names retrieved had occurrence records in South America using the R package
`rgbif v3.8.0` (https://CRAN.R-project.org/package=rgbif, (*70*)) of the Global Biodiversity
Information Facility (GBIF, www.gbif.org). We also manually verified the distribution of each
MOTUs in the PCR plate design to ensure they did not result from tag-jumps from positive
controls. In some cases, we further compared their sequences against Genbank (assessed
in Aug and Oct 2024) using BLAST (*71*). An overview of the dataset size at the main steps of
the data curation process is shown in Table S6.

At the end of the data cleaning procedure, we checked how MOTUs diversity varied with
sequencing depth for all markers and samples (Figure S1). While rarefaction curves did not
saturate when considering MOTU richness, they did so when weighting MOTUs by their
relative abundances using the Hill number framework (*72*, *73*). The Hill numbers, which
correspond to an equivalent number of taxa, is more appropriate to analyse diversity with

DNA metabarcoding data when its *q* order is > 0 (*36*, *37*). We thus present here alpha and
beta diversity analyses for Hill's *q* = 1 in most cases, corresponding to the exponential of the
Shannon index for alpha diversity, and the Horn dissimilarity index for beta diversity (*73*). We
did so for all analyses, except for those related to temporal MOTU diversity and
compositional dynamics, where accounting for rare MOTUs is important due to eDNA decay
processes. For this specific analysis, we computed diversity indices at q = 0, corresponding
to MOTUs richness for alpha diversity, and the Sorensen index for beta diversity, but only
after data rarefaction at 1000 reads / sample. All Hill numbers were calculated with the
`hillR` R package `v0.5.2` (https://CRAN.R-project.org/package=hillR, (*74*)). Finally, we
retrieved the IUCN status - when available - for all the taxa found using the R package
`gbif.range` (https://github.com/8Ginette8/gbif.range, (*75*)). Note that we obtained a
reduced diversity per sample overall for the spatial experiment ((Figure S1, Table S6), most
likely due to the rarity of rain events during the spatial experiment (Figure 1). We further
found that the vertebrate data was excessively contaminated. Because the contaminations
pertained after rerunning the PCRs from scratch while most PCR blank controls were
negative, we concluded that our DNA extracts were compromised for analysing vertebrates,
either due to much rarer traces of local DNA than in the temporal experiment, or due to
higher amounts of contaminations, or both. Consequently, we excluded this dataset from
downstream analyses.

dPCR assays

We also quantified the amount of carrot DNA for each passive eDNA sampler obtained at
each sampled spatial and temporal points through digital PCR (dPCR) on a gene specific to
carrots. For the dPCR-based approach, we quantified the copy number of the carrot *Dau c1*
gene using the QIAcuity Digital PCR System (Qiagen; Hilden, Germany). The single copy
gene targeted encodes for an allergen protein and is now routinely traced through PCR. In
this study, we used the primers Dauc1F (5'-CCAGAGCCATTCACTCGAGATC-3') and
Dauc1R (5'-ACTGTATCAACATCAAGGACAATGC-3') (*76*), together with the the QIAcuity
EG PCR kit 25mL and QIAcuity Nanoplate 26k 24-well kits (QIAGEN; Hilden, Germany).
Each dPCR reaction was performed in a final volume of 40 µL, consisting of 13.3µL of
EvaGreen, 3.2 µL of the two primers mixed at 0.4 µM final concentration each, 13.5µL of
water and 10 µL of template DNA. We also performed one positive control per plate,
adjusting the above conditions for 1 µL of carrot juice DNA extract as template DNA.
Thermocycling conditions were as follow: an initial denaturation step at 95°C for 2 min,
followed by 50 cycles of denaturation at 95°C for 15 sec, primer annealing at 58°C for 15
sec, and elongation at 72°C for 15 sec, and a final elongation step at 72°C for 5 min. The
results were obtained from the machine with default parameters. For a selection of samples
(i.e. Waterra filters obtained at t1 and t20 only), we also conducted dPCRs with the plant
Sper03 marker (see Key resource table) using the reaction mix as above except that we
adjusted the annealing temperature at 50°C (Table S1)

Statistical analyses

All analyses performed with R were done with the R version 4.4.1 (*77*) and the `tidyverse`
package (*78*).

Taxonomic composition overview

To gain a first overview of the relevance of the taxonomic composition obtained from
rainwash eDNA, we summed all MOTUs counts across sampling temporal and spatial points
for each forest type of experiment 1 data. We aggregated the data at the plot scale rather
than considering each spatial point individually due to the punctual information retrieved in
each sample (see below for a more formal analysis on this aspect). We also aggregated the
counts of all MOTUs assigned to the same taxonomic name.

Identifying the best passive eDNA sampler

We identified the best type of passive eDNA sampler based on capacity to quickly fix a high
diversity of taxa and high amounts of carrot spike-in DNA (i.e. one day after passive eDNA
samplers immersion in rainwash water).

For the analysis of diversity detected by each passive sampler type, we considered the
diversity of MOTUs at q=1 as explained above. For the analysis of carrot spike-in DNA, we
used both dPCR assay data (expressed as a number of copies per µL of DNA extract) and a
subset of DNA metabarcoding MOTUs classified as *Apiaceae* (expressed as a number of
Apiaceae sequencing reads per PCR reaction). Both were not normally distributed, which is
why the downstream analysis mostly consisted of non-parametric tests. We found the two
types of data to be well correlated (Spearman's ρ = 0.56, p < 0.001, Figure S3a). A similar
trend was also observed when comparing the gene copy number of our plant marker and the
corresponding total Sper03 sequencing reads per PCR reaction for a subset of samples
(Spearman's ρ = 0.7, p < 0.001, Figure S3b). This adds to previous evidence that DNA
metabarcoding number of reads can be quantitative to a certain extent (*40*) when PCR
products are sequenced together on the same lane and were not subjected to concentration
standardisation prior sequencing. Nevertheless, we present here the two data types as they
do not have the same detection levels, the single-copy nuclear gene Dau c1 being less
detectable than the multi-copy chloroplastic gene used for DNA metabarcoding of plants, but
the latter being more likely to be distorted by different molecular biases.

We then assessed the difference of the diversity, which also did not comply the assumption
of normality, or the amount of carrot spike-in DNA detected across passive sampler types
with Kruskall-Wallis tests, followed by *post-hoc* Wilcoxon tests with FDR-adjusted p-values
when the former identified significant differences. Based on the results obtained, we
considered only the best passive eDNA passive sampler for downstream analyses. These
statistical analyses were run with the `R` package `rstatix` `v0.7.2` ([https://cran.r-](https://cran.r-project.org/package=rstatix)
[project.org/package=rstatix](https://cran.r-project.org/package=rstatix), (*79*)).

Rainwash eDNA temporal dynamics

To identify the optimal temporal window for passive sampling of rainwash eDNA, we
assessed the dynamics (i) of both MOTUs diversity and composition as well as (ii) of the
carrot spike-in DNA. We did not expect the two plots to differ for these properties and
preliminary analyses using mixed models did not retain this random effect. We thus analysed
the data without accounting for this factor.

For this specific analysis, we considered MOTU diversity at q=0 after data rarefaction (see
bioinformatics and data preparation section) rather than at q=1. We did so because it gives

more weight to rare species (*38*), and is thus more sensitive to detect MOTU accumulation
through time by better accounting for MOTUs that would become rarer due to eDNA decay.
We expected MOTU diversity to accumulate rapidly at the start of the experiment and then
either (i) to accumulate more slowly if the carrying capacity of the filter is not reached at t_n
and/or if the decay rate of eDNA is lower than that of the influx of new ones, (ii) to reach a
plateau if the carrying capacity of the filter is reached and/or if an equilibrium between the
decay rate of eDNA and the influx of new ones, and finally (iii) a decrease of MOTU diversity
when the decay rate of rainwash eDNA exceeds that of the influx of new eDNA molecules
(e.g. due to reduced rain events). To this end, we fitted several non-linear functions (*41*), i.e.
power ($D = at^b$), monomolecular ($D = a(1-e^{-bt})$), or Ricker ($D = ate^{-bt}$) on plants, vertebrates and
insects MOTU diversity (D) patterns through time (t). We selected the models exhibiting the
lowest AIC, here built with the Ricker function, from which we estimated the initial diversity
accumulation rate (a) and time at which the diversity was maximal ($T_{Dmax} = 1/b$). Models
predicted values as well as their associated standard deviations and confidence intervals
were obtained with the `R` package `propagate v1.0.6`
(https://CRAN.R-project.org/package=propagate, (*80*)).

We also analysed the MOTU compositional turnover at *q*=0 (see above) using a distance-
decay relationship approach, as we expected samples collected at close sampling date to
exhibit higher similarity than those collected at distant sampling dates, due to both eDNA
persistence/decay and arrival of new eDNA molecules through time. The temporal distance-
decay pattern was studied by considering only the pairwise comparisons involving the same
spatial sampling point. It was assessed with a semi-log linear model, which is often used to
model distance-decay patterns (*81*) and permutation tests to assess the slope significance
with the `R` package `lmPerm` (https://CRAN.R-project.org/package=lmPerm, (*82*)). To
circumvent the log transformation problem on null similarity values, we added the smallest
non-null similarity value to all similarity values.

Finally we determined the persistence of rainwash eDNA by estimating the decay of the
carrot spike-in DNA using a first-order decay model (i.e. a negative exponential function) of
the form $C_t = C_0 e^{-\lambda t}$ where C_0 is the initial eDNA concentration, t is the time in days and λ the
DNA decay rate constant (per day), from which then derived the time of rainwash eDNA
detectability as $t_{0.05} = log(0.05)/\lambda$, corresponding to the time it would take for the carrot spike-
in DNA to fall below 5% of its initial concentration. While more complex decay models have
been found to outperform the one considered here to estimate eDNA decay rate constants
(*42*), our approach allows for cross condition comparisons more easily and appears to be
biased only for very slow decay rates (e.g. at low temperature and in absence of light), which
we do not expect to be the case here. We fitted the decay model only on *Apiaceae* average
reads abundance per PCR reaction. We did not consider here the number of Dau c1 copies /
µL DNA extract, as the detection was too limited to obtain reliable fits and because
quantities obtained are expressed in a different unit than DNA metabarcoding data.

Rainwash eDNA spatial coverage

To assess the optimal spatial sampling effort, we built rarefaction curves for each target
clade from data obtained in the spatial experiment (Figure 1a), by averaging all diversity
values obtained from all possible sample combinations at each sampling intensity (in number
of sampling points per plot) and for each plot separately, as we expected to two plot to
exhibit different diversity and spatial accumulation rates (*44*). Here, MOTU diversity was

measured at q=1 instead of q=0 on rarefied data. The main reason for this choice lies in the robustness of the diversity measure at q=1 to differences in sequencing efforts and potential remaining errors (*36, 37*), as indicated above. In addition, the rate of increase of the uncertainty associated with species addition at q=1 is reduced compared to q=0 (*38, 39*). Finally, unlike the MOTU temporal dynamics analysed above, we did not seek to detect subtle changes in species abundances with increasing sampling effort.

We then fitted on MOTU accumulation curves two simple functions classically used to describe species-area/accumulation curves and exhibiting an upper asymptote in order to estimate the plot-scale diversity and from this, of our sampling coverage. These functions corresponded to the Monod ($DN = D_{max}N/(k_N+N)$) and Negative exponential ($D = D_{max}(1-e^{-kN})$) models (*45*), where D is the diversity, D_{max} the upper asymptote, corresponding here to a theoretical plot-scale diversity detectable with our approach over the period of time considered in our study, N the number of samples, k_N the "half velocity constant", in our case the number of samples required to reach 50% of the plot-scale diversity, and k a MOTU accumulation rate. As for above, we selected the function with the lowest AIC, which corresponded to the Monod models in all cases. We used these models to predict the number of spatial sampling points required to sample 70% and 90% of the plot-scale diversity. Models' predicted values and associated confidence intervals were obtained with the `R` package `propagate` (*80*) as above. Finally, we also evaluated the pairwise MOTU compositional similarities at q=1 between samples across or within each forest plot. We also assessed the relationship between MOTU compositional similarity and geographic distances graphically.

Data and code availability

Raw and pre-processed amplicon and dPCR data as well as data quality reports and analyses scripts are all stored on data.InDoRES (will be provided upon manuscript acceptance). Any additional information required to reanalyse the data reported in this paper is available from the lead contact upon request

References

1. FAO, *The State of the World's Biodiversity for Food and Agriculture* (FAO Commission on Genetic Resources for Food and Agriculture Assessments, Rome, Italy, 2019).

2. A. Antonelli, A. Zizka, F. A. Carvalho, R. Scharn, C. D. Bacon, D. Silvestro, F. L. Condamine, Amazonia is the primary source of Neotropical biodiversity. *Proceedings of the National Academy of Sciences* **115**, 6034–6039 (2018).

3. G. Kier, H. Kreft, T. M. Lee, W. Jetz, P. L. Ibisch, C. Nowicki, J. Mutke, W. Barthlott, A global assessment of endemism and species richness across island and mainland regions. *Proceedings of the National Academy of Sciences* **106**, 9322–9327 (2009).

4. A. Nakamura, R. L. Kitching, M. Cao, T. J. Creedy, T. M. Fayle, M. Freiberg, C. N. Hewitt, T. Itioka, L. P. Koh, K. Ma, Y. Malhi, A. Mitchell, V. Novotny, C. M. P. Ozanne, L. Song, H. Wang, L. A. Ashton, Forests and Their Canopies: Achievements and Horizons in Canopy Science. *Trends in Ecology & Evolution* **32**, 438–451 (2017).

5. IPBES, *The IPBES Regional Assessment Report on Biodiversity and Ecosystem Services for the Americas* (Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany, 2018).

6. D. M. Lapola, P. Pinho, J. Barlow, L. E. O. C. Aragão, E. Berenguer, R. Carmenta, H. M. Liddy, H. Seixas, C. V. J. Silva, C. H. L. Silva-Junior, A. A. C. Alencar, L. O. Anderson, D. Armenteras, V. Brovkin, K. Calders, J. Chambers, L. Chini, M. H. Costa, B. L. Faria, P. M. Fearnside, J. Ferreira, L. Gatti, V. H. Gutierrez-Velez, Z. Han, K. Hibbard, C. Koven, P. Lawrence, J. Pongratz, B. T. T. Portela, M. Rounsevell, A. C. Ruane, R. Schaldach, S. S. da Silva, C. von Randow, W. S. Walker, The drivers and impacts of Amazon forest degradation. *Science* **379**, eabp8622 (2023).

7. F. Li, W. Li, F. Li, Y. Long, S. Guo, X. Li, C. Lin, J. Li, Global projections of future wilderness decline under multiple IPCC Special Report on Emissions Scenarios. *Resources, Conservation and Recycling* **177**, 105983 (2022).

8. P. Taberlet, E. Coissac, F. Pompanon, C. Brochmann, E. Willerslev, Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* **21**, 2045–2050 (2012).

9. P. Taberlet, A. Bonin, L. Zinger, E. Coissac, *Environmental DNA: For Biodiversity Research and Monitoring* (Oxford University Press, 2018).

10. P. Taberlet, E. Coissac, M. Hajibabaei, L. H. Rieseberg, Environmental DNA. *Molecular Ecology* **21**, 1789–1793 (2012).

11. L. Zinger, P. Taberlet, H. Schimann, A. Bonin, F. Boyer, M. De Barba, P. Gaucher, L. Gielly, C. Giguet-Covex, A. Iribar, others, Body size determines soil community assembly in a tropical forest. *Molecular ecology* **28**, 528–543 (2019).

12. A. Kocher, J.-C. Gantier, P. Gaborit, L. Zinger, H. Holota, S. Valiere, I. Dusfour, R. Girod, A.-L. Bañuls, J. Murienne, Vector soup: high-throughput identification of Neotropical phlebotomine sand flies using metabarcoding. *Molecular ecology resources* (2016).

13. S. Calvignac-Spencer, K. Merkel, N. Kutzner, H. Kühl, C. Boesch, P. M. Kappeler, S. Metzger, G. Schubert, F. H. Leendertz, Carrion fly-derived DNA as a tool for comprehensive and cost-effective assessment of mammalian biodiversity. *Molecular Ecology* **22**, 915–924 (2013).

14. H. A. Broadhurst, L. M. Gregory, E. K. Bleakley, J. C. Perkins, J. V. Lavin, P. Bolton, S. S. Browett, C. V. Howe, N. Singleton, D. Tansley, N. G. Sales, A. D. McDevitt, Mapping differences in mammalian distributions and diversity using environmental DNA from rivers. *Science of The Total Environment* **801**, 149724 (2021).

15. I. Cantera, O. Coutant, C. Jézéquel, J.-B. Decotte, T. Dejean, A. Iribar, R. Vigouroux, A. Valentini, J. Murienne, S. Brosse, Low level of anthropization linked to harsh vertebrate biodiversity declines in Amazonia. *Nat Commun* **13**, 3290 (2022).

16. A. Polanco F., M. Mutis Martinezguerra, V. Marques, F. Villa-Navarro, G. H. Borrero Pérez, M.-C. Cheutin, T. Dejean, R. Hocdé, J.-B. Juhel, E. Maire, S. Manel, M. Spescha, A.

Valentini, D. Mouillot, C. Albouy, L. Pellissier, Detecting aquatic and terrestrial biodiversity in a tropical estuary using environmental DNA. *Biotropica* **53**, 1606–1619 (2021).

17. C. Lynggaard, M. F. Bertelsen, C. V. Jensen, M. S. Johnson, T. G. Frøslev, M. T. Olsen, K. Bohmann, Airborne environmental DNA for terrestrial vertebrate community monitoring. *Current Biology* **32**, 701-707.e5 (2022).

18. E. L. Clare, C. K. Economou, F. J. Bennett, C. E. Dyer, K. Adams, B. McRobie, R. Drinkwater, J. E. Littlefair, Measuring biodiversity from DNA in the air. *Current Biology* **32**, 693-700.e5 (2022).

19. M. D. Johnson, M. Fokar, R. D. Cox, M. A. Barnes, Airborne environmental DNA metabarcoding detects more diversity, with less sampling effort, than a traditional plant community survey. *BMC Ecol Evol* **21**, 218 (2021).

20. L. Zinger, J. Donald, S. Brosse, M. A. Gonzalez, A. Iribar, C. Leroy, J. Murienne, J. Orivel, H. Schimann, P. Taberlet, others, Advances and prospects of environmental DNA in neotropical rainforests. *Advances in ecological research* **62**, 331–373 (2020).

21. .R. L. Tulloch, C. I. M. Adams, M. A. Barnes, E. L. Clare, H. C. van de Ven, A. Cridge, F. Encinas-Viso, K. Fernandes, D. M. Gleeson, E. Hill, A. J. M. Hopkins, A. M. Kearns, G. C. Kroos, A. J. MacDonald, F. Martoni, A. McGaughran, T. G. McLay, L. E. Neaves, P. Nevill, A. Pugh, K. J. Robinson, F. Roger, T. V. Steinrucken, M. van der Heyde, C. Villacorta-Rath, J. Vivian, E. Hahn, Winds of change: Charting a pathway to ecosystem monitoring using airborne environmental DNA. (2025).

22. D. Pont, Predicting downstream transport distance of fish eDNA in lotic environments. *Molecular Ecology Resources* **24**, e13934 (2024).

23. C. Lynggaard, S. Calvignac-Spencer, C. A. Chapman, U. Kalbitzer, F. H. Leendertz, P. A. Omeja, E. A. Opito, D. Sarkar, K. Bohmann, J. F. Gogarten, Vertebrate environmental DNA from leaf swabs. *Current Biology* **33**, R853–R854 (2023).

24. K. Yoneya, M. Ushio, T. Miki, Non-destructive collection and metabarcoding of arthropod environmental DNA remained on a terrestrial plant. *Sci Rep* **13**, 7125 (2023).

25. T.-H. Macher, R. Schütz, T. Hörren, A. J. Beermann, F. Leese, It's raining species: Rainwash eDNA metabarcoding as a minimally invasive method to assess tree canopy invertebrate diversity. *Environmental DNA* **5**, 3–11 (2023).

26. I. B. Schnell, K. Bohmann, M. T. P. Gilbert, Tag jumps illuminated - reducing sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources*, n/a-n/a (2015).

27. J. A. Leonard, O. Shanks, M. Hofreiter, E. Kreuz, L. Hodges, W. Ream, R. K. Wayne, R. C. Fleischer, Animal DNA in PCR reagents plagues ancient DNA research. *Journal of Archaeological Science* **34**, 1361–1366 (2007).

28. R. Jairam, A. Fouquet, First records of Leptodactylus nesiotus (Anura: Leptodactylidae) for Guyana, Suriname and French Guiana. *Herpetology Notes* (2018).

29. L. H. Emmons, F. Feer, *Neotropical Rainforest Mammals: A Field Guide* (University of Chicago Press, Chicago, IL, 1997; https://press.uchicago.edu/ucp/books/book/chicago/N/bo3628148.html).

30. Y. Basset, L. Cizek, P. Cuénoud, R. K. Didham, F. Guilhaumon, O. Missa, V. Novotny, F. Ødegaard, T. Roslin, J. Schmidl, A. K. Tishechkin, N. N. Winchester, D. W. Roubik, H.-P. Aberlenc, J. Bail, H. Barrios, J. R. Bridle, G. Castaño-Meneses, B. Corbara, G. Curletti, W. Duarte da Rocha, D. De Bakker, J. H. C. Delabie, A. Dejean, L. L. Fagan, A. Floren, R. L. Kitching, E. Medianero, S. E. Miller, E. Gama de Oliveira, J. Orivel, M. Pollet, M. Rapp, S. P. Ribeiro, Y. Roisin, J. B. Schmidt, L. Sørensen, M. Leponce, Arthropod diversity in a tropical forest. *Science* **338**, 1481–1484 (2012).

31. R. A. F. de Lima, A. A. de Oliveira, G. D. Colletta, T. B. Flores, R. L. G. Coelho, P. Dias, G. P. Frey, A. Iribar, R. R. Rodrigues, V. C. Souza, J. Chave, Can plant DNA barcoding be implemented in species-rich tropical regions? A perspective from São Paulo State, Brazil. *Genet. Mol. Biol.* **41**, 661–670 (2018).

32. H. Richard, J. Ateni, *Guide des arbres de Guyane* (2022; https://www.onf.fr/vivre-la-foret/+/1699::le-guide-des-arbres-de-guyane.html).

33. A. Fouquet, N. Vidal, M. Dewynter, The Amphibians of the Mitaraka massif, French Guiana. *Zoosystema* **41**, 359 (2019).

34. C. Bessey, S. Neil Jarman, T. Simpson, H. Miller, T. Stewart, J. Kenneth Keesing, O. Berry, Passive eDNA collection enhances aquatic biodiversity analysis. *Commun Biol* **4**, 1–12 (2021).

35. S. F. Rivera, V. Vasselon, A. Bouchez, F. Rimet, eDNA metabarcoding from aquatic biofilms allows studying spatial and temporal fluctuations of fish communities from Lake Geneva. *Environmental DNA* **5**, 570–581 (2023).

36. I. Calderón-Sanou, T. Münkemüller, F. Boyer, L. Zinger, W. Thuiller, From environmental DNA sequences to ecological conclusions: How strong is the influence of methodological choices? *Journal of Biogeography* **47**, 193–206 (2020).

37. A. Alberdi, M. T. P. Gilbert, A guide to the application of Hill numbers to DNA-based diversity analyses. *Mol Ecol Resour* **19**, 804–817 (2019).

38. C. Ricotta, E. Feoli, Hill numbers everywhere. Does it make ecological sense? *Ecological Indicators* **161**, 111971 (2024).

39. B. Haegeman, J. Hamelin, J. Moriarty, P. Neal, J. Dushoff, J. S. Weitz, Robust estimation of microbial diversity in theory and in practice. *The ISME Journal* **7**, 1092–1101 (2013).

40. D. Wang, K. B. Trimbos, S. I. F. Gomes, H. Jacquemyn, V. S. F. T. Merckx, Metabarcoding read abundances of orchid mycorrhizal fungi are correlated to copy numbers estimated using ddPCR. *New Phytologist* **242**, 1825–1834 (2024).

41. B. Bolker, *Ecological Models and Data in R* (Princeton University Press, 2008).

42. E. Andruszkiewicz Allan, W. G. Zhang, A. C. Lavery, A. F. Govindarajan, Environmental DNA shedding and decay rates from diverse animal forms and thermal regimes. *Environmental DNA* **3**, 492–514 (2021).

43. MatthewA. Barnes, CameronR. Turner, The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics*, 1–17 (2015).

44. N. J. Gotelli, R. K. Colwell, Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* **4**, 379–391 (2001).

45. J. Dengler, Which function describes the species–area relationship best? A review and empirical evaluation. *Journal of Biogeography* **36**, 728–744 (2009).

46. E. W. Basham, J. A. Baecher, D. H. Klinges, B. R. Scheffers, Vertical stratification patterns of tropical forest vertebrates: a meta-analysis. *Biological Reviews* **98**, 99–114 (2023).

47. J. Terborgh, The Vertical Component of Plant Species Diversity in Temperate and Tropical Forests. *The American Naturalist* **126**, 760–776 (1985).

48. D. de Souza Amorim, B. V. Brown, D. Boscolo, R. Ale-Rocha, D. M. Alvarez-Garcia, M. I. P. A. Balbi, A. de Marco Barbosa, R. S. Capellari, C. J. B. de Carvalho, M. S. Couri, R. de Vilhena Perez Dios, D. A. Fachin, G. B. Ferro, H. F. Flores, L. M. Frare, F. M. Gudin, M. Hauser, C. J. E. Lamas, K. G. Lindsay, M. A. T. Marinho, D. W. A. Marques, S. A. Marshall, C. Mello-Patiu, M. A. Menezes, M. N. Morales, S. S. Nihei, S. S. Oliveira, G. Pirani, G. C. Ribeiro, P. R. Riccardi, M. D. de Santis, D. Santos, J. R. dos Santos, V. C. Silva, E. M. Wood, J. A. Rafael, Vertical stratification of insect abundance and species richness in an Amazonian tropical forest. *Sci Rep* **12**, 1734 (2022).

49. Z. R. Staley, J. D. Chuong, S. J. Hill, J. Grabuski, S. Shokralla, M. Hajibabaei, T. A. Edge, Fecal source tracking and eDNA profiling in an urban creek following an extreme rain event. *Sci Rep* **8**, 14390 (2018).

50. N. R. Garrett, J. Watkins, N. B. Simmons, B. Fenton, A. Maeda-Obregon, D. E. Sanchez, E. M. Froehlich, F. M. Walker, J. E. Littlefair, E. L. Clare, Airborne eDNA documents a diverse and ecologically complex tropical bat and other mammal community. *Environmental DNA* **5**, 350–362 (2023).

51. L. Zinger, A. Bonin, I. G. Alsos, M. Bálint, H. Bik, F. Boyer, A. A. Chariton, S. Creer, E. Coissac, B. E. Deagle, others, DNA metabarcoding—Need for robust experimental designs to draw sound ecological conclusions. *Molecular ecology* **28**, 1857–1862 (2019).

52. S. Gourlet-Fleury (ed), J.-M. Guehl (ed), O. Laroussinie (ed), *Ecology and Management of a Neotropical Rainforest : Lessons Drawn from Paracou, a Long-Term Experimental Research Site in French Guiana* (Elsevier, 2004; https://agritrop.cirad.fr/522004/).

53. S. Peixoto, C. Chaves, G. Velo-Antón, P. Beja, B. Egeter, Species detection from aquatic eDNA: Assessing the importance of capture methods. *Environmental DNA* **3**, 435–448 (2021).

54. S. Oka, M. Miya, T. Sado, Gravity filtration of environmental DNA: A simple, fast, and power-free method. *Methods X* **9** (2022).

55. E. Piccinini, N. Sadr, I. Martin, Ceramic materials lead to underestimated DNA quantifications: a method for reliable measurements. *Eur Cell Mater* **20**, 38–44 (2010).

56. E. J. Green, C. F. Speller, Novel Substrates as Sources of Ancient DNA: Prospects and Hurdles. *Genes (Basel)* **8**, 180 (2017).

57. C. Bessey, Y. Gao, Y. B. Truong, H. Miller, S. N. Jarman, O. Berry, Comparison of materials for rapid passive collection of environmental DNA. *Molecular Ecology Resources* **22**, 2559–2572 (2022).

58. A. Cattaneo, M. C. Amireault, How Artificial Are Artificial Substrata for Periphyton? *Journal of the North American Benthological Society* **11**, 244–256 (1992).

59. A. F. Harrison, P. M. Latter, D. W. H. Walton, Institute of Terrestrial Ecology, Eds., *Cotton Strip Assay: An Index of Decomposition in Soils* (Institute of Terrestrial Ecology, Grange-over-Sands, Cumbria, 1988)*ITE symposium*.

60. S. Tiegs, S. Langhans, K. Tockner, M. O. Gessner, Cotton strips as a leaf surrogate to measure decomposition in river floodplain habitats. *Journal of the North American Benthological Society* **26**, 70–77 (2007).

61. P. E. Berry, A. L. Weitzman, Checklist of the Plants of the Guiana Shield (Venezuela: Amazonas, Bolivar, Delta Amacuro; Guyana, Surinam, French Guiana). (2007).

62. M. A. Renshaw, B. P. Olds, C. L. Jerde, M. M. McVeigh, D. M. Lodge, The room temperature preservation of filtered environmental DNA samples and assimilation into a phenol–chloroform–isoamyl alcohol DNA extraction. *Molecular Ecology Resources* **15**, 168–176 (2015).

63. T. Riaz, W. Shehzad, A. Viari, F. Pompanon, P. Taberlet, E. Coissac, ecoPrimers: inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Research* **39**, e145 (2011).

64. E. Coissac, OligoTag: a program for designing sets of tags for next-generation sequencing of multiplexed samples. *Methods Mol Biol* **888**, 13–31 (2012).

65. K. Bohmann, V. Elbrecht, C. Carøe, I. Bista, F. Leese, M. Bunce, D. W. Yu, M. Seymour, A. J. Dumbrell, S. Creer, Strategies for sample labelling and library preparation in DNA metabarcoding studies. *Molecular Ecology Resources* **22**, 1231–1246 (2022).

66. F. Boyer, C. Mercier, A. Bonin, Y. Le Bras, P. Taberlet, E. Coissac, OBITools : a Unix-inspired software package for DNA metabarcoding. *Molecular Ecology Resources* **16**, 176–182 (2016).

67. C. Mercier, F. Boyer, E. Kopylova, P. Taberlet, A. Bonin, E. Coissac, SUMATRA and SUMACLUST: fast and exact comparison and clustering of sequences. *Programs and Abstracts of the SeqBio 2013 Workshop*, 27–29 (2013).

68. G. Ficetola, E. Coissac, S. Zundel, T. Riaz, W. Shehzad, J. Bessiere, P. Taberlet, F. Pompanon, An In silico approach for the evaluation of DNA barcodes. *BMC Genomics* **11**, 434 (2010).

69. L. Zinger, C. Lionnet, A.-S. Benoiston, J. Donald, C. Mercier, F. Boyer, metabaR: An r package for the evaluation and improvement of DNA metabarcoding data quality. *Methods in Ecology and Evolution* **12**, 586–592 (2021).

70. S. Chamberlain, D. Oldoni, V. Barve, P. Desmet, L. Geffert, D. Mcglinn, K. Ram, rOpenSci (https://ropensci.org/), J. Waller [aut, cre, rgbif: Interface to the Global Biodiversity Information Facility API, version 3.8.0 (2024); https://cran.r-project.org/web/packages/rgbif/index.html.

71. S. F. Altschul, T. L. Madden, A. A. Schäffer, J. Zhang, Z. Zhang, W. Miller, D. J. Lipman, Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* **25**, 3389–3402 (1997).

72. M. O. Hill, Diversity and evenness: a unifying notation and its consequences. *Ecology* **54**, 427–432 (1973).

73. A. Chao, C.-H. Chiu, L. Jost, Unifying species diversity, phylogenetic diversity, functional diversity, and related similarity and differentiation measures through Hill numbers. *Annual Review of Ecology, Evolution, and Systematics* **45**, 297–324 (2014).

74. D. Li, hillR: taxonomic, functional, and phylogenetic diversity and similarity through Hill Numbers. *Journal of Open Source Software* **3**, 1041 (2018).

75. Y. Chauvier, O. Hagen, S. Pinkert, gbif.range - An R package to generate species range maps based on ecoregions and a user-friendly GBIF wrapper, version v1.1 (2022); https://www.doi.org/10.16904/envidat.352.

76. S. Pafundo, M. Gullì, N. Marmiroli, Comparison of DNA Extraction Methods and Development of Duplex PCR and Real-Time PCR To Detect Tomato, Carrot, and Celery in Food. *J. Agric. Food Chem.* **59**, 10414–10424 (2011).

77. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria, 2024; https://www.R-project.org/).

78. H. Wickham, M. Averick, J. Bryan, W. Chang, L. D. McGowan, R. François, G. Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. L. Pedersen, E. Miller, S. M. Bache, K. Müller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, H. Yutani, Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).

79. A. Kassambara, rstatix: Pipe-Friendly Framework for Basic Statistical Tests, version 0.7.2 (2023); https://cran.r-project.org/web/packages/rstatix/index.html.

80. A.-N. Spiess, propagate: Propagation of Uncertainty, version 1.0-6 (2018); https://cran.r-project.org/web/packages/propagate/index.html.

81. J. C. Nekola, P. S. White, The distance decay of similarity in biogeography and ecology. *Journal of Biogeography* **26**, 867–878 (1999).

82. B. Wheeler, M. Torchiano, lmPerm: Permutation Tests for Linear Models, version 2.1.0 (2016); https://cran.r-project.org/web/packages/lmPerm/index.html.

Acknowledgements

We thank the Paracou forest research station in French Guiana, managed and supported by CIRAD, UMR EcoFoG (https://paracou.cirad.fr), and financially supported by a French Investissement d'Avenir programme (Labex CEBA ANR-10-LABX-25-01). We also thank the Genotoul bioinformatics platform of Toulouse Midi-Pyrénées (Bioinfo Genotoul) for providing computing and storage resources. LZ is deeply indebted to Markus Gastauer and to the Fundação de Ciência, Tecnologia, Inovação e Desenvolvimento Sustentável GUAMÁ, from whom she benefited from the grant Bolsa Estímulo à Inovação (BEI) no 094-2023. The work was funded by the OFB (Office Français de la Biodiversité) and has benefitted from 'Investissement d'Avenir' grants managed by Agence Nationale de la Recherche (CEBA: ANR-10-LABX-25-01; TULIP: ANR-10-LABX-0041).

Author contributions

All authors contributed to the conception of the research.

Fr.P. developed the rainwash water collectors.

A.-S.B., A.I., C.L., E.L., Fi.P., Fr.P., V.T, J.O and C.R.-H. performed the fieldwork.

Y.C., A.I., E.L., L.M., V.T., and U.S. performed the molecular work.

A.-S.B. performed the bioinformatics analyses with the help of L.Z. and J.R.

L.Z. performed the statistical analyses with the help of J. P.

A.I. and L.Z. obtained the funding and managed the project.

L.Z. wrote the first draft of the paper and all authors contributed to revisions.

Declaration of interests

The authors declare no competing interests.

Supplemental information titles and legends

Figure S1: Construction diagram of rainwash water collectors from below (topleft), above (topright), a front view from different sides (bottom). The fastener is made of galvanized steel wire (1.5mm diameter), the strings are in 1.5mm nylon cord, the bottle corresponds to a classical 1.5L plastic bottle (here « Dilo » mineral water), and the filter corresponds to the Waterra filter.

Figure S2: Rarefaction curves for plants, vertebrates, and insects metabarcoding data depicting the per-sample accumulation of MOTUs Hill's diversity with increasing sequencing depth. D0 = observed MOTU richness (i.e. Hill's q=0) and D1 ~ exponential of the Shannon index (i.e. Hill's q=1). **a)** Data from the temporal experiment, **b)** Data from the spatial experiment. "Monoculture" corresponds to the tree plantation.

Figure S3: **Correlation between gene copy numbers and sequencing yields per sample** for **a)** the *Dau c1* gene, as assessed through dPCR, and the average number of Apiaceae reads/pcr reaction (n=3 per sample) in DNA metabarcoding data for all samples analysed, and **b)** the Sper03 gene, as assessed through dPCR, and the average total number of reads/pcr reaction (n=3 per sample) in DNA metabarcoding data for Waterra samples. The Spearman correlation test statistics are also indicated.

Figure S4: Efficiency of different passive eDNA samplers after 1 day of immersion in rainwash water in terms of fixation of: **(a)** MOTUs diversity (Hill's q=1) and **(b)** carrot DNA expressed either as a number of *Dau c1* copies (i.e. dPCR data), or as a number of sequencing reads of *Apiaceae* (i.e. DNA metabarcoding data). Stars indicate the significant differences between passive eDNA sampler types identified in the Wilcoxon *post-hoc* tests (FDR-corrected p-values). Values from the tree plantation and the old-growth forest are shown with colours.

Figure S5: Overview of the amount of carrot spike-in DNA across types of passive eDNA samplers (columns) throughout the whole temporal experiment in terms of (rows) (i) number of copies of the nuclear single-copy gene Dau c1 per µL of DNA extract, (ii) average number of *Apiaceae* reads per PCR in DNA metabarcoding data, and (iii) average relative abundance of *Apiaceae* over the average total number of reads per PCR.

Figure S6: Temporal distance - similarity decay pattern of MOTU composition (as measured with the Hill number framework at *q=0* on rarefied data) across sampling dates for each spatial sampling point and for each target clade.

Figure S7: Spatial patterns of MOTU composition (as measured with the Hill number framework at *q*=1) across sampling plots and points for each target clade. **a)** MOTU compositional similarity between vs. within forest plots. **b)** Distance-decay patterns.

Table S1: DNA markers and associated molecular and bioinformatics procedures and parameters used in this study. See Key resources Table for primers and software references.

Table S2: Full list of taxa retrieved in experiment 1 after data curation

Table S3: Kruskall and Wilcoxon test statistics comparing the amount of carrot spike-in DNA
and MOTU diversity (q=1) for each target clade across passive eDNA sampler types.
Abbreviations are as in Figure 1. Significant FDR-corrected p-values (p.adj) are in bold.

Table S4: Coefficient estimates of MOTU diversity (q = 0 on rarefied data) accumulation and
spike-in DNA decay (assessed here only with metabarcoding data) through time with their
associated FDR-adjusted p-values. a : initial MOTU diversity accumulation rate, TD_{max} : time
at which MOTU diversity is maximal, λ : eDNA decay rate, $t_{0.05}$: time of detectability at 5%,
i.e. the time it would take for eDNA to fall below 5% of its initial quantity. n.a.: not applicable.

Table S5: Coefficient estimates of the spatial accumulation of MOTU diversity (at q=1) with
their associated FDR-adjusted p-values. D_{max} : plot-scale estimated MOTU diversity, N_{D50} (=
k_N), $N_{D70,}$ N_{D90}: number of samples required to reach 50, 70 and 90% of the plot-scale diversity.
D_{N16}: Coverage (in %) of the plot-scale diversity with current sampling effort (n=16).

Table S6: Summary statistics on raw and post-filtered data. Positive and blank controls are
excluded in post-curation statistics.

Figures

Figure 1: Overview of the study experimental designs. **a)** Description of the temporal and spatial experiments in terms of precipitation received throughout the sampling periods, initial setting (i.e. addition of carrot spike-in or not), as well as the spatial and temporal distribution of the sampling points. Orange plots: tree plantation, green plots: old-growth forest. **b)** Pictures of a canopy rainwash sampler (see also Figure S1) and of passive eDNA samplers (Cm: ceramic mosaics, CEm: Cellulose Ester membrane Cs: cotton strips, Wf: Waterra filters.

Figure 2: Taxonomic composition of canopy rainwash eDNA in the tree plantation and
the old-growth forest plots for **a**) plants, **b**) vertebrates and **c**) insects in the temporal
experiment. Only taxa with a relative abundance (in terms of % reads) > 0.5% in the total
dataset are shown (summarized at the family levels for plants and insects. The percentage
of reads reported here corresponds to sequencing reads summed across all samples of the
same condition (n = 25 per forest type, i.e. 5 spatial and 5 temporal sampling points). Grey
cells correspond to null values, black ones to very small and yellow ones to high values.

Figure 3: Dynamics and optimal temporal sampling window of rainwash eDNA.
Dynamics of **a**) MOTU diversity (q=0, rarefied data) of plants, vertebrates or insects, **b**)
amount of carrot spike-in DNA. Grey lines and grey transparent ribbons in (**a-b**) correspond
to the models' fits (see also Table S4) and 95% confidence interval. **c**) Temporal window of
detection in terms of maximum diversity for each target clade or of 5% of spike-in initial
concentration, with confidence intervals at 95%.

Figure 4: Accumulation of rainwash eDNA diversity with increasing sampling effort for
plants and insects. Points correspond to observed diversity values. The different line types
represent parameters or curves derived from the Monod models' fits (see also Table S5).
Vertical lines correspond to the number of samples required to sample 70 and 90% of the
plot-scale diversity. For insects, 90% of the diversity requires exceeds 100 samples (see
Table S5). The transparent ribbons represent the confidence intervals at 95% of
extrapolated diversity.

Supplementary Material

Supplementary Figures

Figure S1: Construction diagram of rainwash water collectors from below (topleft), above (topright), a front view from different sides (bottom). The fastener is made of galvanized steel wire (1.5mm diameter), the strings are in 1.5mm nylon cord, the bottle corresponds to a classical 1.5L plastic bottle (here « Dilo » mineral water), and the filter corresponds to the Waterra filter.

Figure S2: Rarefaction curves for plants, vertebrates, and insects metabarcoding data
depicting the per-sample accumulation of MOTUs Hill's diversity with increasing sequencing
depth. D0 = observed MOTU richness (i.e. Hill's q=0) and D1 ~ exponential of the Shannon
index (i.e. Hill's q=1). **a)** Data from the temporal experiment, **b)** Data from the spatial
experiment. "Monoculture" corresponds to the tree plantation.

Figure S3: **Correlation between gene copy numbers and sequencing yields per sample**
for **a)** the *Dau c1* gene, as assessed through dPCR, and the average number of Apiaceae
reads/pcr reaction (n=3 per sample) in DNA metabarcoding data for all samples analysed,
and **b)** the Sper03 gene, as assessed through dPCR, and the average total number of
reads/pcr reaction (n=3 per sample) in DNA metabarcoding data for Waterra samples. The
Spearman correlation test statistics are also indicated.

Figure S4: Efficiency of different passive eDNA samplers after 1 day of immersion in
rainwash water in terms of fixation of: **(a)** MOTUs diversity (Hill's *q*=1) and **(b)** carrot DNA
expressed either as a number of *Dau c1* copies (i.e. dPCR data), or as a number of
sequencing reads of *Apiaceae* (i.e. DNA metabarcoding data). Stars indicate the significant
differences between passive eDNA sampler types identified in the Wilcoxon *post-hoc* tests
(FDR-corrected p-values). Values from the tree plantation and the old-growth forest are
shown with colours.

Figure S5: Overview of the amount of carrot spike-in DNA across types of passive eDNA samplers (columns) throughout the whole temporal experiment in terms of (rows) (i) number of copies of the nuclear single-copy gene Dau c1 per µL of DNA extract, (ii) average number of *Apiaceae* reads per PCR in DNA metabarcoding data, and (iii) average relative abundance of *Apiaceae* over the average total number of reads per PCR.

831 **Figure S6:** Temporal distance - similarity decay pattern of MOTU composition (as measured
832 with the Hill number framework at *q=0* on rarefied data) across sampling dates for each
833 spatial sampling point and for each target clade.

Figure S7: Spatial patterns of MOTU composition (as measured with the Hill number framework at *q*=1) across sampling plots and points for each target clade. **a)** MOTU compositional similarity between vs. within forest plots. **b)** Distance-decay patterns.

843 Supplementary Tables

844 **Table S1:** DNA markers and associated molecular and bioinformatics procedures and
845 parameters used in this study. See Key resources Table for primers and software
846 references.

		Sper03	Vert01	Inse01
Marker characteristics	Target clades	Viridiplantae	Vertebrata	Arthropoda/Insecta
	Target gene	rbcL	mt 12S	mt 16S
	Fwd sequence (5'-3')	CACCACAAACAGARACT AAARC	TTAGATACCCCACTATGC	RGACGAGAAGACCCT ATARA
	human blocking primer (5'-3')	n.a.	CTATGCTTAGCCCTAAAC CTCAACAGTTAAATCAAC AAAACTGCT-Spc3	n.a.
	rev sequence (5'-3')	TCCAYACRGTTGTCCAT GTACC	TAGAACAGGCTCCTCTAG	ACGCTGTTATCCCTAA RGTA
	avg length (nt)	185	97	155
PCR thermocycling conditions	initial denaturation	10 min at 95°C		
	nb. cycles denaturation, annealing, elongation,	40 cycles: 95°C (1 min) 52°C (1 min) 72°C (1 min 30 s)	50 cycles: 95°C (30 s) 50°C (30 s) 72°C (30 s)	40 cycles: 95°C (30 s) 52°C (30 s) 72°C (30 s)
	final elongation	72°C (7 min)	72°C (7 min)	72°C (7 min)
Sequencing conditions	sequencing platform	NovaSeq 6000 - Illumina		
	chemistry	Flowcell SP - 2x250 PE		
Bioinformatics parameters: I[ary] curation (obitools)	read pairing	exclusion of paired-reads with alignments with scores < 40		
	demultiplexing	2 mismatches allowed on primers, no mismatches on tags		
	spurious sequences	containing Ns, singletons		
		length < 50	length < 50	length < 55
	clustering	97% sequence similarity		
	databases building characteristics	- from EMBL 2021-12-21 - only Viridiplantae references - references with a family name defined - references of length between 30 to 300 nt.	- from Genbank 259 - references with a family name defined - references of length between 40 to 145 nt.	- from Genbank 259 - references with a family name defined - references of length between 125 to 285 nt.
Bioinformatics parameters: II[ary] curation (metabaR)	contaminant detection	maximum abundance in controls		
	minimum similarity against	85%	80%	75%

	reference databases			
	minimum relative abundance allowed (tag-jump control)	0.05	0.05	0.05
	semi-manual checks for species distribution	done	done	too large databases incompleteness

850 **Table S2:** Full list of taxa retrieved in experiment 1 after data curation (=>
851 table_metabar_taxalist_all.tsv)

853 **Table S3:** Kruskall and Wilcoxon test statistics comparing the amount of carrot spike-in DNA
854 and MOTU diversity (q=1) for each target clade across passive eDNA sampler types.
855 Abbreviations are as in Figure 1. Significant FDR-corrected p-values (p.adj) are in bold.

Data type	Statistics				
	Kruskall test n = 30	statistic	p-value		
		13.2	**7.53e-4**		
	Wilcoxon test n = from 5 to 9 each	Passive sampler 1 (n)	Passive sampler 2 (n)	statistic W	p.adj
		CEm	Cm	47.5	0.870
Plant MOTU diversity		CEm	Cs	30.5	0.304
		CEm	Wf	9	**0.004**
		Cm	Cs	32	0.285
		Cm	Wf	4	**7.8e-4**
		Cs	Wf	10	**0.004**
	Kruskall test n = 35	statistic	p-value		
		8.54	**0.0361**		
	Wilcoxon test n = from 8 to 9 each	Passive sampler 1 (n)	Passive sampler 2 (n)	statistic W	p.adj
		CEm	Cm	30.5	0.479
		CEm	Cs	42.5	0.56
Vertebrate MOTU diversity		CEm	Wf	16	0.101 (=0.034 non adjusted)
		Cm	Cs	52	0.198
		Cm	Wf	22	0.198
		Cs	Wf	11	0.101 (=0.018 non adjusted)
Insect MOTU diversity	Kruskall test n = 40	statistic	p-value		
		3.46	0.327		
Concentration (copies Dau c1/µL DNA extract)	Kruskall test n = 40	statistic	p-value		
		20.9	**0.000109**		
	Wilcoxon test n = 10 each	Passive sampler 1 (n)	Passive sampler 2 (n)	statistic W	p.adj

		CEm	Cm	75	0.076
		CEm	Cs	40	0.481
		CEm	Wf	5	**0.000618**
		Cm	Cs	25	0.076
		Cm	Wf	0	**0.0000648**
		Cs	Wf	12	**0.006**
Apiaceae sequencing yield (avg reads/pcr)	Kruskall test n = 40	statistic	p-value		
		10.4	**0.0151**		
	Wilcoxon test n = 10 each	Passive sampler 1 (n)	Passive sampler 2 (n)	statistic W	p.adj
		CEm	Cm	44	0.880
		CEm	Cs	45	0.880
		CEm	Wf	11	**0.013**
		Cm	Cs	49	0.970
		Cm	Wf	16	**0.027**
		Cs	Wf	21	0.062

Table S4: Coefficient estimates of MOTU diversity (q = 0 on rarefied data) accumulation and spike-in DNA decay (assessed here only with metabarcoding data) through time with their associated FDR-adjusted p-values. a : initial MOTU diversity accumulation rate, TD_{max} : time at which MOTU diversity is maximal, λ : eDNA decay rate, $t_{0.05}$: time of detectability at 5%, i.e. the time it would take for eDNA to fall below 5% of its initial quantity. n.a.: not applicable.

Clade	Model type	Coefficient	Coefficient value	Confidence interval (95%)	p.adj
Plants	Ricker	a	9.86	7.50 - 12.76	**3.86e-10**
		TD_{max}	14.33	11.56 - 17.86	**6.89e-13**
Vertebrates	Ricker	a	2.31	1.51 - 3.40	**5.24e-06**
		TD_{max}	11.74	8.60 - 16.19	**4.21e-08**
Insects	Ricker	a	8.59	5.19 - 13.57	**4.54e-05**
		TD_{max}	13.14	8.94 - 19.46	**1.08e-06**
Apiaceae sequencing yield (avg number of reads/pcr)	Negative exponential	λ	0.10	0.07 - 0.14	**8.29e-07**
		$t_{0.05}$	29.13	21.18 - 40.95	n.a.

864 **Table S5:** Coefficient estimates of the spatial accumulation of MOTU diversity (at q=1) with
865 their associated FDR-adjusted p-values. D_{max} : plot-scale estimated MOTU diversity, N_{D50} (=
866 k_N), N_{D70}, N_{D90}: number of samples required to reach 50, 70 and 90% of the plot-scale diversity.
867 D_{N16}: Coverage (in %) of the plot-scale diversity with current sampling effort (n=16).

	Forest type	Tree plantation			Olg-growth		
Clade	Coefficients	Estimate	CI (95%)	p.adj	Estimate	CI (95%)	p.adj
Plant	D_{max}	23.820	23.598 - 24.056	**0**	31.764	31.534 - 31.996	**0**
	$N_{D50}(=k_N)$	4.602	4.482 - 4.723	**0**	5.771	5.671 - 5.872	**0**
	D_{N16}	77.662	77.357 - 77.966	n.a.	73.492	73.288 - 73.696	n.a.
	N_{D70}	10.5	10 - 11	n.a.	13.5	13 - 14	n.a.
	N_{D90}	39	35 - 43	n.a.	50	45 - 55	n.a.
Insects	D_{max}	34.675	33.699 - 35.711	**0**	74.399	71.729 - 77.282	**0**
	$N_{D50}(=k_N)$	18.539	17.779 - 19.347	**0**	23.199	22.061 - 24.430	**0**
	D_{N16}	46.324	46.022 - 46.624	n.a.	40.817	40.52 - 41.109	n.a.
	N_{D70}	41	39 - 43	n.a.	50.5	48 - 53	n.a.
	N_{D90}	136	124 - 148	n.a.	161.5	148 - 175	n.a.

869 **Table S6:** Summary statistics on raw and post-filtered data. Positive and blank controls are
870 excluded in post-curation statistics.

			Sper03	Vert01	Inse01
Temporal experiment	raw data[1]	total # reads total # samples	25,059,048 200	52,612,351 200	34,535,786 200
	Primary curation	total # reads total # motus total # samples	23,492,156 65,717 200	50,058,332 55,001 200	31,891,191 51,237 200
		# reads (±sd) / sample # motus (±sd) / sample	117,460 (±91,260) 146 (± 139)	250,291 (± 230,054) 230 (± 129)	159,455 (± 192,441) 165 (± 177)
	Secondary curation	total # reads total # motus total # samples	17,905,314 12,723 200	10,854,793 4,252 200	18,622,452 12,187
		# reads (±sd) / sample # motus (±sd) / sample	89,526 (± 78,408) 63 (± 95)	54,273 (± 96,010) 21 (± 39)	93,112 (± 163,141) 60 (± 107)
Spatial experiment	raw data[1]	total # reads total # samples	5,828,776 32	12,846,173 32	2,197,546 32
	Primary curation	total # reads total # motus total # samples	5,605,882 21,811 32	12,580,901 6,561 32	1,971,798 6,175 32
		# reads (±sd) / sample # motus (±sd) / sample	175,183 (± 110,215) 519 (± 240)	393,153 (± 358,917) 188 (± 138)	61,618 (± 31,796) 124 (± 86)
	Secondary curation	total # reads total # motus total # samples	4,399,092 6,071 32	3,472,721 200 32	437,406 1318 32
		# reads (±sd) / sample # motus (±sd) / sample	137,471 (± 93,061) 189 (± 140)	108,522 (± 282,061) 6 (± 8)	13,668 (± 14,896) 41 (± 32)

871 [1] Provided only after read pairing and demultiplexing, the sequencing lane containing
872 samples from another project.