Unraveling new characteristics of γδ T cells using scRNA-seq in TCR KO chicken

Theresa von Heyl[1], Milena Brunner[1], Antonia Schmauser[1], Christine Wurmser[2], Simon
Früh[3,4], Benjamin Schusser[1*]

[1] Reproductive Biotechnology, TUM School of Life Science, Technical University of Munich, 85354
Freising, Germany
[2] Chair of Animal Physiology and Immunology, TUM School of Life Sciences, Technical University
of Munich, 85354 Freising, Germany
[3] Department of Veterinary Sciences, Ludwig-Maximilians-Universität München, 80539 Munich,
Germany
[4] Virale Integration, Tumorigenesis and Virus evolution, Institute of Virology, Freie Universität
Berlin, 14163 Berlin, Germany
11
* Correspondence:
Benjamin Schusser
benjamin.schusser@tum.de

Abstract

The characterization of T cells in chickens has proven to be challenging, primarily due to the lack of
specific markers to differentiate classical Th1, Th2, and Th17 subsets. Furthermore, chickens possess
a notably high proportion of γδ T cells, making them a unique model for investigating the poorly
understood role of these cells not only in chickens but also in mammals. To gain deeper insights into
the functions and characteristics of the αβ and γδ T cell subsets in chickens, whole transcriptome
analysis (WTA) on $CD3^+$ single cells isolated from wild type (WT), TCR Cβ knockout (KO) and
TCR Cγ KO chickens at embryonic day (ED) 18, day (d) 3, and d14 post-hatch was conducted. The
results indicate that γδ T cells exhibit cytolytic activity in both TCR Cβ KO and WT chickens. A
distinct cluster of γδ T cells expressing genes associated with the interferon pathway was revealed in
TCR Cβ KO chickens, which may contribute to the severe phenotype in these chickens, which is
characterized by severe inflammation of spleen, gut and stomach. Additionally, novel gene markers
were identified that precisely define the αβ and γδ T cell subsets. These findings provide new insights
into chicken T cell biology and contribute to a better understanding of the mechanisms underlying
the severe phenotype observed in TCR Cβ KO chickens.
30

31 **Keywords: T cells, chicken, gamma/delta T cells, genetically modified animals, scRNA-seq**

1   Background

Chickens are an indispensable livestock species, as not only the chicken meat but also the eggs are a principal protein source worldwide. Compared to other livestock species, chicken meat is a healthier and more resource-efficient source of protein (Farrell 2011, Salter 2018). Livestock diseases often pose significant threats, leading to great economic losses. Moreover, diseases like avian influenza pose a considerable zoonotic risk to humans (Bodewes, Bestebroer et al. 2015, Rahman, Sobur et al. 2020, Abdelwhab and Mettenleiter 2023). Therefore, a detailed understanding of the chicken immune system is essential to ensure the health of chickens and humans.

T cells are key players in the adaptive immunity of chickens (Cooper and Alder 2006). They are categorized into $\alpha\beta$ and $\gamma\delta$ T cells based on their T cell receptor chains (Chen, Sowder et al. 1989). The role of $\alpha\beta$ T cells is defined through co-expression of CD4 and CD8 (Arstila, Vainio et al. 1994, Luhtala 1998). The $CD4^+$ $\alpha\beta$ T cells are activated through antigen peptides presented on MHC class II molecules, leading to production of cytokines, targeted immune responses and the stimulation of B cells to secrete immunoglobulins (Arstila, Vainio et al. 1994). The $CD8^+$ T cells can directly lyse pathogens by secreting cytotoxic cytokines upon activation through peptides on MHC class I molecules (Luhtala 1998). The balance between protective inflammation and tissue damage is stabilized by regulatory T cells (T_{reg}), described by their transcription factor *FoxP3* (Burkhardt, Elleder et al. 2022). Compared to mammals, there is no clear classification into Th1, Th2 and Th17 cells of $\alpha\beta$ T cells in chickens, as the classification is based on the secretion of various interleukins, some of which have not yet been described (Mousset, Hobo et al. 2019).

The other major T cell subset, $\gamma\delta$ T cells, remains poorly understood in both chickens and mammals. Chickens present a compelling model for studying the role of $\gamma\delta$ T cells in greater detail, as they are a $\gamma\delta$-high species, with comprising up to 50% $\gamma\delta$ T cells of circulating lymphocytes throughout their lifespan (Sowder, Chen et al. 1988). In contrast, mice and humans typically harbor only 1–10% $\gamma\delta$ T cells in the peripheral blood(Sowder, Chen et al. 1988, Holderness, Hedges et al. 2013). To date, chicken $\gamma\delta$ T cells are known to exhibit cytotoxic activity when stimulated *in vitro* (Fenzl, Gobel et al. 2017). They are abundantly present in peripheral epithelial tissues (Hayday, Roberts et al. 2000), playing a crucial role in the immune response to infections such as Salmonella (Berndt and Methner 2001). $\gamma\delta$ T cell subsets have been previously characterized by Berndt et al. into three groups: $CD8^{high+}$, $CD8^{dim+}$, and $CD8^{neg-}$. These $\gamma\delta$ T cell subsets respond differently to infections like salmonellosis (Berndt, Pieper et al. 2006).

To characterize T cell subsets in more detail, we generated knockout chickens that lack $\alpha\beta$ or $\gamma\delta$ T cells, respectively. The phenotypical analysis of the TCR Cγ KO chicken showed a comparable phenotype to wild type (WT) chickens (von Heyl, Klinger et al. 2023). Additionally, infection of TCR Cγ KO chickens with MDV showed increased disease incidence and tumor formation compared to their WT siblings, leading to the conclusion that $\gamma\delta$ T cells play an important role in the pathogenesis and tumor formation (Sabsabi, Kheimar et al. 2024). Strikingly, TCR Cβ KO chickens exhibit a severe phenotype two weeks after hatch, with granuloma formation at the comb, beak, and legs. In addition, inflammation of the stomach and spleen, as well as an impaired development of the lymphatic organs was observed. The cause of this phenotype and a potential contribution of the remaining $\gamma\delta$ T cells remain elusive (von Heyl, Klinger et al. 2023).

Using scRNA-seq analysis, Maxwell et al. identified leukocyte populations such as B cells, eosinophils, basophils, monocytes, red blood cells, and thrombocytes in adult 24 week old laying hens. The description of T cell subtypes proved to be challenging in chickens, but it led to new insights regarding cytotoxic characteristics of particular T cells (Maxwell, Soderlund et al. 2024).

Here, we used scRNA-seq to perform whole transcriptome analysis (WTA) of CD3$^+$ T cells sorted
from the thymus and peripheral blood mononuclear cells (PBMCs) of WT, TCR Cβ KO and TCR Cγ
KO animals on embryonal day 18 (ED18), day 3 (d3) and day 14 (d14) post hatch. Our data revealed
two different cytotoxic γδ T cell clusters in the TCR Cβ KO at 14 days post-hatch, which were absent
in TCR Cγ KO and WT chickens as well as potential gene markers to identify γδ and αβ T cell
clusters.

2   Methods:

2.1   Animals:

Two genetically modified chicken lines, TCR Cβ KO and TCR Cγ KO, derived from the White
Leghorn genetic background (Lohmann Selected Leghorn [LSL]; Lohmann-Tierzucht GmbH,
Cuxhaven, Germany), were used in this study. The generation and phenotypic characterization of
these lines have been described previously (von Heyl, Klinger et al. 2023). Animal experiments were
approved by the government of Upper Bavaria, Germany (ROB-55.2-2532.Vet_02-17-101 & 55.2-1-
54-2532-104-2015). Experiments were performed according to the German Welfare Act and
European Union Normative for Care and Use of Experimental Animals. All animals received a
commercial standard diet and water *ad libitum*.

2.2   Organ / Tissue collection:

For the sampling of the thymus of ED18 chicks, TCR Cγ KO and TCR Cβ KO eggs were incubated
for 18 days in egg incubators (HEKA-Brutgeräte, Rietberg, Germany) at standard conditions (37,8°C.
55% relative humidity, six rotations/day). Infertile eggs or dead embryos were removed by candling
the eggs on day 7 of incubation. *In ovo* genotyping was performed on ED10. On ED18, eggs were
opened, and embryos were decapitated and the thymus was collected from the embryos. For the other
time points, TCR Cβ KO, TCR Cγ KO, and their WT sibling were hatched. Blood was drawn for
genotyping on d1 post hatch. On d3 and d14 post hatch, chickens were euthanized by terminal
bleeding, and the thymus was removed. From all genotypes (WT, TCR Cβ KO and TCR Cγ KO) n=3
animals irrespective of sex were used at each time point (ED18, d3, d14).

2.3   Genotyping assays:

For *in ovo* genotyping, allantois samples were collected on ED10. Briefly, a candling light was used
to determine fertility of the egg, embryo viability, and to mark the areas for creating a hole with a
lancet where no blood vessels were visible. The allantoic fluid was collected using an insulin syringe
with a 30G cannula (BD Micro Fine^{TM}, Heidelberg, Germany). Allantoic fluid was directly used for
genotyping without a gDNA isolation step.
For the hatched chicks, a drop of blood was collected one day after hatch from the *vena jugularis*
using a syringe with a 27G cannula (B. Braun, Meisungen, Germany). gDNA was isolated from
blood, by adding 1μL blood into 200μL STM buffer and centrifuging for 5min at 100g. After
removing the supernatant, 400μL TEN buffer containing 100μg/mL Pronase E was added, and the
mix was incubated for 1h at 37°C (250rpm). The Pronase E was subsequently inactivated at 65°C for
20min (Bailes, Devers et al. 2007).
Chickens were genotyped as described before (von Heyl et al.). Briefly, primers were used to detect
TCR Cβ KO chickens: Forward: 5' GGTTCGAAATGACCGACCAAGC 3'; Reverse: 5'
GGCTTGCACACTCAGCTCTATAG 3'. A second primer pair was used to detect the TCR Cβ WT
allele: Forward: 5' GGTTCGAAATGACCGACCAAGC 3'; Reverse: 5'

CACACCATTCACCTTCCAGAC 3'. FIREPol Multiplex DNA Polymerase Mastermix (Solis
Biodyne, Tartu, Estonia) was used according to manufactures instructions with an annealing
temperature (Tm) of 58°C. For the TCR Cγ KO chickens' primers were used to detect the TCR Cγ
KO: Forward: 5' GCCATTCCTATTCCCATCCTAAGT 3'; Reverse: 5'
GGTTCGAAATGACCGACCAAGC 3'. A second primer pair was used to detect the WT constant
region of the TCR γ chain: Forward: 5' GAGCTCCACGCCATGAAACCATAG 3'; Reverse: 5'
GTTGTCACTGTCACTGGCTG 3'. FIREPol Multiplex DNA Polymerase Mastermix (Solis
Biodyne, Tartu, Estonia) was used according to manufactures instructions with an annealing Tm of
60°C.

2.4   Cell sorting:

The removed thymus was strained through a 100µm cell strainer into a falcon holding 5mL ice-cold
PBS. Lymphocytes were isolated from blood and the strained thymus single-cell solution using
histopaque density gradient centrifugation (Sigma, Taufkirchen, Germany). Cells were washed with
1% BSA in PBS and stained with CD3-PE (clone CT-3; 0,5µg/mL) (Biozol, Eching, Germany) for
20 min in the dark on ice. Subsequently, cells were again washed, and cell sorting was performed
using a CytoFLEX SRT Cell sorter (BD, Heidelberg, Germany). Data were analyzed with FlowJo
10.8.1 software (FlowJo, Ashland, USA).

2.5   Multiplexing:

In order to multiplex all samples from each timepoint the BD Flex Single-Cell Multiplexing Kit A, B,
and C (BD, Heidelberg, Germany) were used according to the manufacturer's protocol. Briefly, cells
were resuspended in 180µL PBS+ 1%FBS. 20µL of each sample tag was added and incubated.
Samples were washed with PBS+ 1%FBS and pooled together.

2.6   Single-cell capture and cDNA Synthesis

The BD Rhapsody™ cDNA Kit and BD Rhapsody Enhanced Cartridge Reagent Kit (BD,
Heidelberg, Germany) were used according to the manufacturer's protocol. The 8-Lane Cartridge
was loaded with 70.000 cells.

2.7   mRNA Whole Transcriptome Analysis and Sample Tag

The BD Rhapsody™ WTA Reagent Kit (BD, Heidelberg, Germany) was used according to the
manufacturers protocol (23-24119(02)). Briefly, random priming and extension (RPE) reactions of
the Exonuclease I-treated beads containing the whole cellular transcriptome and cDNA was
performed. After this step two parallel workflows were performed, one for the WTA libraries, using
the RPE product from the supernatant and the other one for the Sample Tag libraries from the post-
RPE beads. The RPE supernatant underwent WTA library preparation following the manufacturers'
instructions from the BD Rhapsody™ System mRNA Whole Transcriptome Analysis (WTA) Library
Preparation Protocol that included the steps of RPE cleanup, performing RPE PCR, RPE PCR
cleanup, performing WTA Index PCR and cleanup. The Sample Tag products were amplified and
purified through PCR. A Sample Tag Index PCR was performed to add full-length Illumina
Sequencing adapters and indices.

WTA libraries and Sample Tag libraries were pooled and sequenced on a NovaSeqX 10B Sequencer
by Novogene (Cambridge, UK). Libraries were sequenced as paired-end 2x150bp reads with an
estimated depth of 1,250 M reads per cartridge lane.

2.8   Single-cell data analysis

2.8.1 Preprocessing of scRNA-seq raw data:

Fastq files were processed using the BD Rhapsody™ WTA Analysis Pipeline, v1.9.1 in the Seven
Bridges Genomics cloud platform. WTA reads were aligned to the *Gallus gallus* (chicken) GRCg7b
reference genome (Accession Number GCF_016699485.2) and demultiplexed. The pipeline
generates molecular counts per cell, reads per cell, metrics and an alignment file.

2.8.2 Integration, Clustering and Annotation

Bioinformatic analyses were conducted in RStudio (Version 2024.04.2+764) using Seurat (Version
5.1.0). Cluster analyses were performed following the Seurat integration workflow (Version
5.2.0; https://satijalab.org/seurat/articles/integration_introduction). Differentially expressed genes
(DEGs) were identified from the integrated clusters according to the Seurat pipeline
(https://satijalab.org/seurat/articles/pbmc3k_tutorial.html). Functional classification of gene markers
was carried out using PANTHER (Version 19.0) for gene ontology (GO) analysis. Cluster annotation
was performed manually based on the inspection of DEGs.
176

3   Results:

3.1   Integrated Transcriptomic Landscapes of CD3$^+$ T cells in Thymus and PBMCs

In this study, whole transcriptome analysis was conducted on sorted CD3$^+$ T cells isolated from WT,
TCR Cβ KO, and TCR Cγ KO animals. Thymus samples were collected from ED18, d3, and d14
chickens, while PBMC samples were obtained from d3 and d14 chickens (Figure 1A). Due to the
limited blood volume in ED18 embryos, it was not feasible to sort the required 20,000 cells per
embryo. Generally, the CD3$^+$ population in the thymus ranged from 15% to 49,4% at all timepoints.
The CD3$^+$ population in blood increased from d3 to d14. At d3 the population ranged between 0,73%
to 8,98% of all PBMCs with no differences between the genotypes. At d14 TCR Cβ KO chickens
showed a smaller population of CD3$^+$ PBMCs (mean 2,57%), compared to TCR Cγ KO (mean
8,88%) and WT (mean 10,17%) animals (Figure 1B and C and Supplement Figure 1-3). From each
sample 20,000 cells were sorted, except for one TCR Cβ KO thymus sample at d14 where only 8,000
sorted cells were achieved. The quality of the sorting captured using BD Rhapsody^{TM} Scanner was
summarized for each timepoint in Table 1 and 2. The Sequencing quality was overall good since over
96% of the reads passed the quality filter and over 90% of the reads were assigned to a putative cell
at all three timepoints (Table 1 and 2). At all three timepoints over 90% of the reads were aligned to a
putative cell. At d3 the highest mean with 14,650.78 reads per cell was found. On ED18 39,080, on
d3 45,884 and on d14 80,075 viable cells were loaded. The cell multiplet rate ranged from 7,2% to
15,8% (Table 1). The single cell captures overall showed a good viability of the cells with a bead
efficiency of 94,6%-98,2% (Table 1). Clusters from all timepoints, organs and genotypes were
integrated for comparison, resulting in a total of 17 identified Clusters. In general, the clusters of
TCR Cβ KO and TCR Cγ KO showed only few overlaps in the thymus samples (Figure 3 A-C).

3.2   Annotation of lineage defining markers revealed functional states of T cell subclusters

All three timepoints were integrated into a single data set. Clusters of CD3□ T cells were manually
annotated based on the expression of lineage-defining marker genes within the integrated dataset.
The markers used for cluster identification are listed in Table 3 (see also Supplementary Figure 4).

The two largest clusters contained a similar number of cells: n = 13,848 effector αβ T cells (16.96%),
defined by the expression of *CD28, ICOS,* and *CD8A*, and n = 13,830 naïve αβ T cells (16.95%),
characterized by *FOXO1, KLF2, IL7R, P2PRY8, CD38,* and *RORA*. The largest γδ T cell cluster
comprised n = 7,330 mature γδ T cells (8.98%). Annotation of the T cell subclusters revealed distinct
functional states across the cell cycle, including differentiating, proliferating, and activated T cells,
which collectively contributed to the circular arrangement of clusters in the UMAP representation.
Despite prior enrichment for CD3□ T cells, small contaminating populations were consistently
detected across most samples, including thrombocytes (n = 1,401 cells; 1.7%), erythrocytes (n = 290
cells; 0.3%), and myeloid cells (n = 138 cells; 0.2%). Thrombocytes were identified by expression of
GP1BB, GP9, and *F13A1*; erythrocytes by *HBE1, SLC4A1, EPB42, SPTB,* and *DMTN*; and myeloid
cells by *ANXA1, CD74, PDCD1LG2, BLB1,* and *BLB2* (Figure 2 A and B; Tables 3–4).

3.3   Subclustering of T cells on ED18 reveals division active precursor T cells

Subclustering the ED18 dataset, TCR Cβ KO chickens exhibited a major subcluster of mature γδ T
cells (34,8%, n= 3,313) in the thymus, along with two minor clusters of activated γδ T cells (Cluster
8 =14,4%, n= 1369; Cluster 15= 0,34%, n=32 cells) and a minor cluster of proliferating γδ T cells
(10,4%, n = 992 cells). Additionally, division-active precursor (3,4%, n=320 cells) and differentiating
(4,7%, n= 447 cells) αβ T cells were observed. In TCR Cγ KO chickens, effector (31,7%, n= 3483)
and differentiating (45,4%, n= 4979) αβ T cells constituted major subpopulations together with the
division-active precursor cells (8,5%, n= 938 cells). Proliferating αβ (5,1%, n= 561) T cells were also
observed. In the wild type siblings, the largest three clusters were also effector T cells (23,5%, n=
1959), differentiating αβ T cells (19,5%, n= 1626) and mature γδ T cells (16,9%, n= 1412), which is
comparable to the largest clusters in the TCR Cβ KO and TCR Cγ KO chickens (Figure 3A).

3.4   Subclustering on d3 shows activated γδ T cells in PBMCs of TCR Cβ KO chickens

Subclustering data from d3 revealed in the thymus of WT chickens, the major clusters contained
effector αβ (33,4%, n= 1459) and differentiating αβ T cells (17,0%%, n= 744), naïve T cells (7,8%,
n= 341) and mature γδ T cells (8,0%, n= 352). Minor clusters were proliferating αβ (3,6% n= 158)
and proliferating γδ (5,7%, n= 249) T cells. In the thymus of TCR Cβ KO the main clusters were also
mature γδ T cells (25,1% n=737) and proliferating γδ T cells (12,8%, n=375). But other than in WT
also activated γδ T cells (17,3%, n=507) were a dominant population in the thymus of TCR Cβ KO
chickens. In the TCR Cγ KO chickens the principal cell type were as in WT chickens effector αβ T
cells (40,7%, n=1764), naïve αβ T cells (19,8%, n= 859) and differentiating αβ T cells (9,9%,
n=429).
While in PBMCs WT chickens exhibited major clusters of mature αβ T cells (43,8%, n= 1784), naïve
T cells (13,0% n= 531) and differentiating γδ T cells (12,9%, n= 525). In PBMCs of TCR Cβ KO
birds a principal group of differentiating γδ T cells (PBMCs: 22,9%, n= 482) and naive T cells
(26,2%, n= 552), with a minor cluster of cytolytic T cells (6,7%, n=142) were designated. By
comparing the PBMCs of TCR Cγ KO and WT chickens, TCR Cγ KO birds displayed an increased
population of naïve T cells (57,7%, n= 2159) (Figure 3B).

3.5   Cluster annotation reveals different activated γδ T cell clusters in TCR Cβ KO chickens
**   on d14**

Next, the data from d14 was subclustered. Comparing TCR Cγ KO and WT T cell clusters in the
chicken's thymus on d14 the clusters of effector αβ T cells (WT: 26,9%, n= 1676; TCR Cγ KO:
30,2%, n= 1544), differentiating αβ T cells (WT: 12,4%, n= 773; TCR Cγ KO: 11,0%, n= 560), and

naïve T cells (WT: 12,2%, n= 762; TCR Cγ KO: 22,1%, n= 1130) were mostly similar. In the thymus of TCR Cβ KO chickens mature (18,0%, n=780), differentiating (10,6%, n= 461) and interferon activated γδ T cells (10,2%, n=442) formed major clusters. The PBMCs of TCR Cγ KO chickens showed one dominant cell population: naïve αβ T cells (54,9%, n=2542). Both, TCR Cβ KO and WT chickens, have equal clusters of differentiating γδ T cells (TCR Cβ KO: 23,0%, n= 1175; WT: 19,9%, n= 1018) and naïve T cells (TCR Cβ KO: 19,9%, n= 921; WT: 31,5%, n= 1611) on d14 in the PBMCs. Surprisingly, in TCR Cβ KO chickens we identified two large clusters of activated T cells. The first cluster expressed genes associated with the interferon pathway, including *MX1, STAT1,* and *IFNGR1*, and was therefore designated as interferon-activated γδ T cells (12.0%, n= 520). The second group, which represented the major population in TCR Cβ KO chickens at day 14, consisted of cytolytic T cells characterized by the expression of *GZMM* and *GNLY* (31.0%, n= 1,345) (Figure 3C).

3.6   TARP as a useful marker to identify γδ T cell clusters

To describe γδ T cells in more detail, *TARP* was identified for γδ T cells as a potential marker to annotate γδ T cell clusters since it showed high expression (>1) in over 75% of the cells from TCR Cβ KO chickens at every timepoint in both PBMCs and thymus. In wild type *TARP* was intermediately expressed (=0) in around 50% of the cells and had a negative expression in TCR Cγ KO chickens (< -1) at all three timepoints (Figure 4A). *MAF, SOX13, GATA3, KK34, INPP5A, PDE1A, CD82* and *IL20RA* showed equal high expression in TCR Cβ KO chickens and intermediate expression in WT chickens. However, the relative expression of *SOX13, GATA3, PDE1A* and *CD82* were higher compared to *MAF, KK34, INPP3A* and *IL20RA*. Leading to the conclusion that these genes were found in specific γδ T cells clusters only. Feature Plot indicated the location of cells expressing genes describing γδ T cells on the integrated UMAP. While *TARP, GATA3* and *CD82* seem to be expressed in all γδ T cell clusters, *PDE1A, KK34, INPP5A* and *IL20RA* are only expressed in specific γδ T cell clusters (Figure 3B). *MAF* was only expressed in activated and naïve T cells and was not exclusively expressed in γδ T cells but also αβ T cells. *KK34* was only expressed in naïve γδ T cells. Alongside, the markers *PDE1A* and *CD82* were highly expressed in the γδ T cells. However, only *PDE1A* seemed to be completely absent in TCR Cγ KO chickens (Figure 4B). To identify a large population of αβ T cells *CD4, CD8A, CD5 and CCR7* were used. All four genes were absent in TCR Cβ KO chickens in the thymus. In PBMCs we found high expression in around 20% of the cells from TCR Cβ KO chickens at d3 and d14, indicating differentiated CD8$^+$ γδ T cells. Moreover, high expression (>1) of *CD4* and *CD8A* in 40-60% of the cells in the thymus of TCR Cγ KO chickens at all timepoints was found and intermediate expression in WT chickens at all timepoints in the thymus (Figure 5A). While *CD4* and *CD8A* are present in both naïve and activated αβ T cells, *CD5* was present in effector αβ T cells and *CCR7* was used to identify naïve αβ T cells. (Figure 5B).

3.7   Interferon pathway activated in TCR Cβ KO chickens

Comparing genes which are significantly upregulated in TCR Cβ KO compared to WT chickens at d14 after hatch, revealed genes which are involved in the interferon pathway (Figure 6A). The string networks showed the close relation of the genes involved into interferon pathway, such as *STAT1, MX1* and *OASL* (Figure 6B). A heatmap analysis of genes that are involved in the interferon pathway (*STAT1, IFI6, MX1, USP41, IFNGR1, OASL, RSAD2, IFIT5)* showed strong expression (>2) of these genes in the thymus and PBMCs of TCR Cβ KO chickens on d14. While there are less cell clusters in WT and TCR Cγ KO chickens on d14 that express genes involved in the interferon pathway.

Interestingly, in the Thymus and PBMCs on d3 and ED18 less expression of interferon-stimulated
genes (Figure 6C) was detected.

3.8   Cytolytic CD8+ γδ T cells expressing Granulysin and Granzyme A

The analysis of cytolytic genes on d14 in the T cells showed, that TCR Cβ KO chicken express
cytokines that are related to lysis and apoptosis. High expression (n= 2) of Granulysin (*GNLY*),
Granzyme A (*GZMA*), Granzyme M (*GZMM*) and *IL2RB* in the PBMCs of TCR Cβ KO chickens
was detected on d14, while this was not observed in the thymus of these chickens. Also, on ED18
and d3 there is no such upregulation of cytotoxic γδ T cells visible in any genotype (Figure 7A). The
string networks show the close relation of the cytokines and chemokines involved in cytotoxicity
(Figure 7B).

4   Discussion:

Single cell RNA-sequencing is a novel tool to understand gene expression patterns in cells to define
clusters (Tang, Barbacioru et al. 2009, Eschke, Moore et al. 2023). In this study, we used WTA to
analyze T cell clusters in TCR KO and WT chickens from embryonic day 18 (ED18) until two weeks
post hatch (d14).
Data integration was performed across all genotypes (TCR Cβ KO, TCR Cγ KO, and wild type),
developmental stages (ED18, d3, and d14), and organs (thymus and PBMCs). This approach enables
a more comprehensive comparison across the different layers of the analysis. However, integrating
all datasets also introduces the risk of misclassification of a subset of single cells. Although T cells
can be subdivided into αβ and γδ lineages, both share common transcriptional programs related to
processes such as proliferation and differentiation. As a result, cells from different T-cell subgroups
may cluster together despite belonging to distinct lineages (Hao, Hao et al. 2021)
The first T cells that move into the thymus are TCR1[+] cells which occur first on ED12 and are first
detectable in the spleen by ED15. The second wave of T cells colonize the thymus at ED15 and move
towards spleen on ED19 (Chen, Sowder et al. 1989, Coltey, Bucy et al. 1989, Chen, Gobel et al.
1994). Even though there should be some CD3[+] T cells in the blood of ED18, we were not able to
sort enough cells for scRNA-seq.
Through the use of TCR KO chicken, it was possible to detect different genes that are specific
markers for γδ or αβ T cells. Before, *TARP, MAF, GATA3, KK34* and *INPP5A* were described as γδ
T cell markers (Maxwell, Soderlund et al. 2024). Likewise, in this study we also found that *TARP*
and *GATA3* are highly+ expressed in the γδ T cells. While *MAF* is only expressed in activated and
mature T cells and is not exclusively expressed in γδ T cells but also αβ T cells. The genes *KK34* and
INPP5A show a low expression but are exclusively expressed in γδ T cells. Alongside we found that
the markers *PDE1A* and *CD82* are highly expressed in the γδ T cells. However, only *PDE1A* seems
to be absent in TCR Cγ KO chickens, while *TARP* and *CD82* is also expressed in some cells of the
TCR Cγ KO chickens. *IL20RA* showed low expression but is a selective marker for the detailed
description of γδ T cells. *IL20RA* is known to be a subunit of the IL20 receptor and is involved in the
regulation of inflammatory mediators by regulating the *JAK-STAT* pathway (Lamichhane, Mo et al.
2021). *CD82* (also known as *KAI1* or *R2*) is an inducible membrane marker that was initially
identified from a T cell activation study (Gaugitsch, Hofer et al. 1991). Here, we highlight for the
first time, that *CD82* is not exclusively expressed in γδ T cells, but comparing TCR Cβ KO with TCR
Cγ KO and WT chickens, we see that most of γδ T cells express *CD82* while only few αβ T cells
express *CD82*. Until now there is no literature on γδ T cells expressing the calmodulin-dependent
phosphodiesterase 1A (*PDE1A*) exclusively. This study reveals for the first time the absence of
PDE1A in TCR Cγ KO chickens and shows that only a specific γδ T cell cluster express *PDE1A*.

Before, scRNA-seq in chicken was performed in peripheral blood leukocytes (Maxwell, Soderlund
et al. 2024). Here, the T cell clusters were subdivided in CD4, T_{regs}, CD8, γδ T cells and cytolytic
cells. However, in our study we subdivided the T cells not only by the markers from Maxwell et al
(2024) but also further due to their differentiation status into proliferation, differentiation, naïve T
cells and mature T cells (Figure 2,3). Likewise, in comparison to Maxwell et al. we found cytolytic T
cells, that show high expression of Granulysin and Granzyme A and M in TCR Cβ KO animals 14
days after hatch (Maxwell, Soderlund et al. 2024). Other than shown before, here we can show that
this cytolytic T cells are γδ T cells in TCR Cβ KO birds. Interestingly, these cytolytic γδ T cells do
neither express perforin nor Granzyme B.
Depending on the expression of specific cytokines, helper T cells are subdivided in Th1, Th2, Th17
and T_{regs} in humans and mammals (Raphael, Nalawade et al. 2015). Still, it was not possible to define
these T cell subsets for the helper T cells in chickens due to missing annotations of some cytokines
and transcription factors. In this analysis, the T cells were subdivided into different clusters
depending on their activation status. However, we did not find individual clusters for different helper
T cells subclass, since in this study the sorted CD3[+] T cells in the wildtype showed no expression of
IFN-γ (Th1), IL-4 (Th2), IL-17 (Th17) or IL-10 (T_{reg}). Even though these markers are already
annotated in the chicken genome.
In addition to T cell populations, we identified minor clusters of thrombocytes (expressing *GP1BB*,
GP9, and *F13A1*), erythrocytes (expressing *HBE1, SLC4A1*, and *EPB42*), and myeloid cells
(expressing *BLB1* and *BLB2*) (see Table 4; Figures 2 and 3), despite the fact that samples were sorted
for CD3□ T cells. These clusters are clearly distinct from the T cell populations, represent less than
2% of all cells, and therefore do not interfere with downstream analyses. Maxwell et al. (2024) used
an additional immunomagnetic separation step to reduce thrombocytes based on the expression of
CD41/61. However, even using a cleanup step still a population of 5-7% Thrombocyte was measured
in their samples (Maxwell, Soderlund et al. 2024). These irregularities do not interfere with the T cell
analysis, as these cell clusters can be clearly annotated using specific markers.
In the thymus at both ED18 and d3, we identified two clusters (clusters 8 and 15) that we annotated
as activated γδ T cells. These clusters shared core γδ T cell markers such as *TARP, ZAP70*, and
CD28, supporting their classification. Despite this common identity, they exhibited distinct activation
marker profiles. Cluster 8 expressed *CHAC1, ASNS, MTHFD2, SLC1A4, PSAT1,* and *CD44*, while
cluster 15 was characterized by the expression of *IL2RA (CD25), IL2RB, IL1R1,* and *IL18R1* (see
Tables 4, Figures 2 and 3). Although the activation markers differed between the two clusters, both
were classified broadly as activated γδ T cells, as a more refined subclassification was not feasible
based on the available data.
In this study, we identified for the first time a subset of γδ T cells that exhibit the expression of
genes associated with the interferon pathway, including *MX1, STAT1*, and *OASL* (Mountford, Gheyas
et al. 2022). TCR Cβ KO chickens display a severe phenotype as early as 14 days post hatch,
characterized by inflammation of the gut, spleen, and skin. This condition is driven by an imbalance
between humoral and cytotoxic responses within the adaptive immune system (von Heyl, Klinger et
al. 2023). The interferon-activated γδ T cells described here might play a defining role in the severity
of the phenotype observed in TCR Cβ KO chickens. Further studies are needed to determine whether
these T cells are also present in other inflammatory diseases in chickens and to assess if their
differentiation is triggered by the activation of dysregulated monocytes.
377

5   Conclusion

This study revealed that cytolytic T cells, characterized by the expression of Granulysin and
Granzymes, are predominantly γδ T cells in TCR Cβ KO chickens. Both the cytolytic γδ T cells and

the interferon-activated γδ T cells are likely contributors to the severe phenotype observed in TCR
Cβ KO chickens. However, the precise mechanisms underlying this phenotype remain unclear,
highlighting the need for future sequencing analyses of monocytes and B cells to further elucidate the
drivers of this response.

Overall, this study provides new insights into chicken T cell biology, including T cell markers for
specific subsets such as αβ T cells and γδ T cells, as well as their functional states, such as
proliferation and differentiation. Notably, the use of TCR Cβ KO chickens has uncovered novel
characteristics of γδ T cells, offering valuable perspectives for further research.

6   List of abbreviations

αβ	alpha beta
γδ	gamma delta
B cell	B Lymphocyte
CD	Cluster of differentiation
d	day
DEG	Differentially expressed gene
ED	Embryonal day
FBS	Fetal bovine serum
G	Gauge
GO	Gene ontology
KO	Knockout
PBMC	Peripheral blood mononuclear cells
PBS	Phosphate-buffered saline
RPE	random priming and extension
T cell	T Lymphocyte
TCR	T cell receptor
TCR Cβ KO	Chicken line lacking alpha beta T cells
TCR Cγ KO	Chicken line lacking gamma delta T cells
scRNA-seq	Single cell RNA-sequencing
UMAP	Uniform Manifold Approximation and Projection
WT	Wild type
WTA	Whole transcriptomic analysis
°C	Degrees Celsius

7   Ethics approval and consent to participate

Animal experiments were approved by the government of Upper Bavaria, Germany (ROB-55.2-
2532.Vet_02-17-101 & 55.2-1-54-2532-104-2015). Experiments were performed according to the
German Welfare Act and European Union Normative for Care and Use of Experimental Animals.

8   Consent for publication

Not applicable

9   Data Availability Statement

Raw data for this study can be found on SRA under the accession number PRJNA1226579.

10 Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

11 Funding

This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) in the framework of the Research Unit ImmunoChick (FOR5130) project SCHU 2446/6-1 (project number: 434524619; awarded to BS) and an Emmy-Noether research fellowship (DFG Schu2446/3-1 awarded to BS).

12 Author Contributions

TvH performed and analyzed the experiments and wrote the paper. BS supervised the work and wrote the paper. MB and SPF helped with the analysis of the data. AS and CW performed experiments. All authors approved the submitted version.

13 Acknowledgments:

ChatGPT (https://chat.openai.com/) was used to improve conciseness of language and coding, with no contribution to the scientific content.

14 References

Abdelwhab, E. M. and T. C. Mettenleiter (2023). "Zoonotic Animal Influenza Virus and Potential Mixing Vessel Hosts." <u>Viruses</u> **15**(4).

Arstila, T. P., O. Vainio and O. Lassila (1994). "Central role of CD4+ T cells in avian immune response." <u>Poult Sci</u> **73**(7): 1019-1026.

Bailes, S. M., J. J. Devers, J. D. Kirby and D. D. Rhoads (2007). "An inexpensive, simple protocol for DNA isolation from blood for high-throughput genotyping by polymerase chain reaction or restriction endonuclease digestion." <u>Poult Sci</u> **86**(1): 102-106.

Berndt, A. and U. Methner (2001). "Gamma/delta T cell response of chickens after oral administration of attenuated and non-attenuated Salmonella typhimurium strains." <u>Vet Immunol Immunopathol</u> **78**(2): 143-161.

Berndt, A., J. Pieper and U. Methner (2006). "Circulating gamma delta T cells in response to Salmonella enterica serovar enteritidis exposure in chickens." <u>Infect Immun</u> **74**(7): 3967-3978.

Bodewes, R., T. M. Bestebroer, E. van der Vries, J. H. Verhagen, S. Herfst, M. P. Koopmans, R. A. Fouchier, V. M. Pfankuche, P. Wohlsein, U. Siebert, W. Baumgartner and A. D. Osterhaus (2015). "Avian Influenza A(H10N7) virus-associated mass deaths among harbor seals." <u>Emerg Infect Dis</u> **21**(4): 720-722.

Burkhardt, N. B., D. Elleder, B. Schusser, V. Krchlikova, T. W. Gobel, S. Hartle and B. Kaspers (2022). "The Discovery of Chicken Foxp3 Demands Redefinition of Avian Regulatory T Cells." <u>J Immunol</u> **208**(5): 1128-1138.

Chen, C. H., T. W. Gobel, T. Kubota and M. D. Cooper (1994). "T cell development in the chicken." <u>Poult Sci</u> **73**(7): 1012-1018.

Chen, C. H., J. T. Sowder, J. M. Lahti, J. Cihak, U. Losch and M. D. Cooper (1989). "TCR3: a third T-cell receptor in the chicken." <u>Proc Natl Acad Sci U S A</u> **86**(7): 2351-2355.

Coltey, M., R. P. Bucy, C. H. Chen, J. Cihak, U. Losch, D. Char, N. M. Le Douarin and M. D.
Cooper (1989). "Analysis of the first two waves of thymus homing stem cells and their T cell
progeny in chick-quail chimeras." J Exp Med **170**(2): 543-557.
Cooper, M. D. and M. N. Alder (2006). "The evolution of adaptive immune systems." Cell **124**(4):
815-822.
Eschke, M., P. F. Moore, H. Chang, G. Alber and S. M. Keller (2023). "Canine peripheral blood
TCRalphabeta T cell atlas: Identification of diverse subsets including CD8A(+) MAIT-like cells by
combined single-cell transcriptome and V(D)J repertoire analysis." Front Immunol **14**: 1123366.
Farrell, D. (2011). "The role of poultry in human nutrition." FAO Poultry Production.
Fenzl, L., T. W. Gobel and M. L. Neulen (2017). "gammadelta T cells represent a major
spontaneously cytotoxic cell population in the chicken." Dev Comp Immunol **73**: 175-183.
Gaugitsch, H. W., E. Hofer, N. E. Huber, E. Schnabl and T. Baumruker (1991). "A new superfamily
of lymphoid and melanoma cell proteins with extensive homology to Schistosoma mansoni antigen
Sm23." Eur J Immunol **21**(2): 377-383.
Hao, Y., S. Hao, E. Andersen-Nissen, W. M. Mauck, 3rd, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk,
C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T.
Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P. Smibert
and R. Satija (2021). "Integrated analysis of multimodal single-cell data." Cell **184**(13): 3573-3587
e3529.
Hayday, A. C., S. Roberts and E. Ramsburg (2000). "gammadelta cells and the regulation of mucosal
immune responses." Am J Respir Crit Care Med **162**(4 Pt 2): S161-163.
Holderness, J., J. F. Hedges, A. Ramstead and M. A. Jutila (2013). "Comparative biology of
gammadelta T cell function in humans, mice, and domestic animals." Annu Rev Anim Biosci **1**: 99-
124.
Lamichhane, S., J. S. Mo, G. Sharma, T. Y. Choi and S. C. Chae (2021). "MicroRNA 452 regulates
IL20RA-mediated JAK1/STAT3 pathway in inflammatory colitis and colorectal cancer." Inflamm
Res **70**(8): 903-914.
Luhtala, M. (1998). "Chicken CD4, CD8alphabeta, and CD8alphaalpha T cell co-receptor
molecules." Poult Sci **77**(12): 1858-1873.
Maxwell, M., R. Soderlund, S. Hartle and E. Wattrang (2024). "Single-cell RNA-seq mapping of
chicken peripheral blood leukocytes." BMC Genomics **25**(1): 124.
Mountford, J., A. Gheyas, L. Vervelde and J. Smith (2022). "Genetic variation in chicken interferon
signalling pathway genes in research lines showing differential viral resistance." Anim Genet **53**(5):
640-656.
Mousset, C. M., W. Hobo, R. Woestenenk, F. Preijers, H. Dolstra and A. B. van der Waart (2019).
"Comprehensive Phenotyping of T Cells Using Flow Cytometry." Cytometry A **95**(6): 647-654.
Rahman, M. T., M. A. Sobur, M. S. Islam, S. Ievy, M. J. Hossain, M. E. El Zowalaty, A. T. Rahman
and H. M. Ashour (2020). "Zoonotic Diseases: Etiology, Impact, and Control." Microorganisms **8**(9).
Raphael, I., S. Nalawade, T. N. Eagar and T. G. Forsthuber (2015). "T cell subsets and their signature
cytokines in autoimmune and inflammatory diseases." Cytokine **74**(1): 5-17.
Sabsabi, M. A., A. Kheimar, Y. You, D. von La Roche, S. Hartle, T. W. Gobel, T. von Heyl, B.
Schusser and B. B. Kaufer (2024). "Unraveling the role of gammadelta T cells in the pathogenesis of
an oncogenic avian herpesvirus." mBio: e0031524.
Salter, A. M. (2018). "The effects of meat consumption on global helth." Revue Scientifique et
Technique **37**: 47-55.
Sowder, J. T., C. L. Chen, L. L. Ager, M. M. Chan and M. D. Cooper (1988). "A large subpopulation
of avian T cells express a homologue of the mammalian T gamma/delta receptor." J Exp Med **167**(2):
315-322.

Tang, F., C. Barbacioru, Y. Wang, E. Nordman, C. Lee, N. Xu, X. Wang, J. Bodeau, B. B. Tuch, A.
Siddiqui, K. Lao and M. A. Surani (2009). "mRNA-Seq whole-transcriptome analysis of a single
cell." Nat Methods **6**(5): 377-382.
von Heyl, T., R. Klinger, D. Aumann, C. Zenner, M. Alhussien, A. Schlickenrieder, K. Lengyel, H.
K. Vikkula, T. Mittermair, H. Sid and B. Schusser (2023). "Loss of alphabeta but not gammadelta T
cells in chickens causes a severe phenotype." Eur J Immunol **53**(12): e2350503.

15 Tables

Table 1. Quality control metrics of BD Rhapsody[TM] Scanner

Timepoint	Number of wells with viable cells at cell load	Cell multiplet rate	Bead load efficiency
ED18	39080	7,2%	94,6%
d3	45884	9,8%	97,9%
d14	80075	15,8%	98,2%

496 **Table 2. Sequencing quality of cells and sample multiplexing provided by Novogene:**

	Cells				Sample Multiplexing	
Day	Putative Cells	Reads Passing Quality Filter	Reads from putative cells	Mean Reads per cell	Valid Reads aligned to Sample Tags	Reads from putative cells
ED18	41,998	97,33%	93,87%	13,258.31	5,611,637	65,20%
d3	37,975	96,98%	90,83%	14,650.78	7,944,922	68,93%
d14	54,683	96,83%	92,96%	12,617.67	6,633,932	71,24%

498 **Table 3. Cellcount of integrated clusters**

clusters	Cellcount	Percentage (%)
0	13848	16,96
1	13830	16,94
2	10453	12,80
3	7330	8,98
4	5363	6,57
5	5350	6,55

6	3539	4,34
7	3477	4,26
8	3328	4,08
9	3139	3,85
10	2883	3,53
11	2318	2,84
12	2149	2,63
13	1570	1,92
14	1401	1,72
15	1229	1,51
16	290	0,36
17	138	0,17
Total	81636	100

500 **Table 4. Lineage defining genes for cluster annotation**

Cluster	Biological Process	Genes
0	Effector T cells	CD28, ICOS, CD8A
1	Naïve αβ T cells	FOXO1, KLF2, IL7R, P2RY8, CD38, RORA
2	Differentiating αβ T cells	RAG1, RAG 2, CD4, CD8A
3	mature γδ T cells	PDE1A, IL20RA, CD82, TARP
4	Differentiating γδ T cells	CTLA4, SOX13, GATA3, IL7R, TARP
5	Naive γδ T cells	FOXO1, KLF2, IL7R, P2RY8, CD38, RORA
6	Proliferating γδ T cells	TARP, CDCA7, MCM2,3,4,5,6, TOP2A, CDK1, PPIA
7	Cytolytic T cells	GPR18, GRNLY, GRMM, PTPN22
8	Activated γδ T cells	CHAC1, ASNS, MTHFD2, SLC1A4, PSAT1, CD28, CD44, TARP
9	Division active precursor T cells	CDK1, TOP2A, CCNBI, BUB1B, CCNA2, CDC20, PLK1, CENPF, ASPM, KIF11, KIFF20A, TACC3
10	Interferon activated T cells	CD8A, CD28, STAT1, BACH2, IFNGR1, IL7R
11	Proliferating αβ T cells	CDCA7, MCM2,4,5,6, TOP2A, CDK1
12	Division active T cells	FOXM1, MELK, MCM10, TOP2A, NCAPG2, DTL
13	Thrombocytes	GP1BB, GP9, F13A1
14	Proliferating αβ T cells	CDCA7, MCM2,4,5,6, TOP2A, CDK1
15	Activated γδ T cells	IL2RA (=CD25), IL2RB,

		IL1R1, IL18R1
16	Erythrocytes	HBE1, SLC4A1, EPB42, SPTB, DMTN
17	Myeloid cells	ANXA1+, CD74, PDCD1LG2, BLB1, BLB2

16   Figure Legends

Figure 1. Experimental setup for the single cell analysis of chicken CD3+ T cells. (A) PBMC and thymocytes of three chickens from each genotype WT, TCR Cβ KO and TCR Cγ KO were used. CD3+T cells were isolated by fluorescence-activated cell sorting (FACS) and applied to 5' single-cell RNA sequencing (5' scRNA-seq) for whole transcriptome analysis (WTA) (created with BioRender). (B) Flow cytometric analysis of CD3+ T cells from Thymus and PBMCs of WT, TCR Cβ KO and TCR Cγ KO chickens n=3 at ED18, d3, d14 (Mean +/- SD are shown). (C) The gating strategy for isolation of CD3+ T cells by FACS. One representative dotplot of each timepoint (ED1, d3, d14) and each genotype (WT, TCR Cβ KO and TCR Cγ KO) is shown.

Figure 2. (A) Single-cell RNA sequencing of CD3□ T cells isolated from the thymus and PBMCs of WT, TCR Cβ KO and TCR Cγ KO chickens at ED18, d3, and d14 post hatch revealed distinct gene clusters in the integrated UMAP plots. UMAPs showing CD3+ T cell clusters annotated according to specific gene markers in Table 4. (B) Percentages (%) of cells in each Cluster (0-17) of integrated data from all three timepoints.

Figure 3. Identification of T cell subsets among chicken CD3+ clusters on (A) ED18, (B) d3 and (C) d14 post hatch from WT, TCR Cβ KO and TCR Cγ KO chickens from thymus and PBMCs. UMAPs showing CD3+ T cell clusters annotated according to specific gene markers in Table 4.

Figure 4. Identification of markers for γδ T cell subsets among chicken CD3+ cells. (A) Dot plot representing the expression of a selection of γδ T cell associated genes in subgroups grouped by timepoint, organ and genotype. The radius of the dot corresponds to the percentage of cells in each cluster expressing the gene, and color intensity corresponds to scaled expression values (average_log2 fold change). (B) Feature plot of potential γδ T cell markers in integrated UMAP cluster. The color intensity corresponds to scaled expression values.

Figure 5. Identification of markers for αβ T cell subsets among chicken CD3+ cells from PBMCs. (A) Dot plot representing the expression of a selection of αβ T cell associated genes in in subgroups grouped by timepoint, organ and genotype. The radius of the dot corresponds to the percentage of cells in each cluster expressing the gene, and color intensity corresponds to scaled expression values (average_log2 fold change). (B) Feature plot of potential αβ T cell markers in integrated UMAP cluster. The color represents the expression level (average_log2 fold change).

Figure 6. Interferon-activated γδ T cell clusters. A) Volcano plot of differentially expressed genes (DEGs) between TCR Cβ KO and WT chickens in PBMCs at d14 post hatch. The log2(FC) indicates the mean fold-change (FC) in expression levels for each gene, and each dot represents one gene. DEGs identified in pairwise comparisons between TCR Cβ KO and WT chickens are defined as p < 0.05, FC > |2|. DEGs with higher and lower expression are represented by red dots, while genes with no substantial change in expression are indicated as grey dots. B) String Network of DEGs from TCR Cβ KO compared to WT chickens involved in interferon pathway. C) Heatmap of clusters from

543   thymus and PBMCs of WT, TCR Cβ KO and TCR Cγ KO chickens at d14 showing genes involved
544   in the interferon pathway. The color represents the gene expression level (average_log2 fold change).
545
546   **Figure 7.** Cytolytic γδ T cell clusters. A) Heatmap of Clusters from thymus and PBMCs of WT, TCR
547   Cβ KO and TCR Cγ KO chickens 14 days after hatch showing genes involved in cytolytic activity.
548   The color represents the gene expression level (average_log2 fold change). B) String Network of
549   DEGs involved in cytolytic activity.

A

B

A

B

C

A

B

A

B