# Critical Period Plasticity is Associated with Resilience to Short Unpredictable Stress

**Robert Williams, Jr[1†], Charlie Van Den Oord[2†], Erica N. Lee[1†], Samuel C. Fedde[1], Gia L.**
**Oscherwitz[3], Adema Ribic[1,*]**

[1]Department of Psychology, College of Arts and Sciences, University of Virginia, Charlottesville,
VA

[2]Department of Biology, College of Arts and Sciences, University of Virginia, Charlottesville, VA

[3]School of Engineering, University of Virginia, Charlottesville, VA

[†]Equal contribution

**\* Correspondence:**
Corresponding Author
adema.ribic@virginia.edu

**Keywords: plasticity, SynCAM 1, stress, resilience, hyperthermia**

**Abstract**

Low resilience to stressful events can increase the risk of anxiety and depression. Resilience
decreases with age, parallel to drastic changes in the quality of brain plasticity from juvenile to old
age, suggesting that the type of plasticity found in the maturing brain promotes resilience. To
indirectly test this, we administered short unpredictable stress to adult male and female wild type
(WT) C57BL/6 mice, as well as to two groups of mice characterized by heightened cortical plasticity:
adolescent C57BL/6 WT mice and adult mice that lack SynCAM 1 (Synaptic Cell Adhesion
Molecule 1), a critical plasticity brake in the mature brain. We found that short unpredictable stress
robustly increased core body temperature in all groups of mice, indicative of stress-induced
hyperthermia (SIH) and confirming the efficacy of the stress paradigm. However, depressive-like
behavior as measured though tail suspension test was increased in adult WT mice only, supporting
that the type of plasticity found in the immature brains of adolescent WT and adult SynCAM 1
knockout (KO) mice promotes resilience to stress. All three groups of mice showed a mild increase
in locomotor activity after stress, suggesting that the quality of plasticity does not correlate with
resilience to anxiety-like phenotypes. Our study hence provides indirect evidence for the protective
role of developmental plasticity during stress and points to new mechanisms that promote resilience
to stress-induced depression.

**Introduction**

Stressful life events are strongly linked to the onset of depression (McEwen, 2012; McEwen et
al., 2015; Czéh et al., 2016), which is one of the leading causes of disability in the USA (WHO). Yet,
stress does not increase the risk of depression in individuals that are resilient: capable of successfully

adapting to stress exposure (Faye et al., 2018; McEwen and Morrison, 2013; Russo et al., 2012). Stress resilience is thought to decline with age, parallel to changes in the quality of brain plasticity, which transitions from robust critical period plasticity during in the developing brain to a restricted, top-down modulated process in the adult brain (Ribic, 2020). While the link between plasticity and resilience is still unclear, a previous study demonstrated that chronic restraint stress results in a reversible reduction in the length of dendritic branches in the prefrontal cortex (PFC) of young, but not aged rats, in whom the reduction persists (Bloss et al., 2010). Heightened brain plasticity during postnatal development, as well as interventions that change the quality of plasticity in adulthood, can hence promote resilience to stress (McEwen, 2016, 2012). In support of this notion, antidepressants markedly increase plasticity in the adult brain and change its quality to a more juvenile/adolescent-like (Maya Vetencourt et al., 2008; Phoumthipphavong et al., 2016). Environmental enrichment, a classical behavioral paradigm that also "rejuvenates" the adult cortex (Greifzu et al., 2016, 2014), mitigates the effects of stress across the lifespan (Macartney et al., 2022; Lehmann and Herkenham, 2011; Hegde et al., 2020). However, both antidepressants and environmental enrichment have systemic effects on the brain, and it is still unclear if they promote resilience and mitigate the effects of stress through altered quality of brain plasticity.

In mice, a species of choice for genetic manipulations of plasticity, stress during late postnatal development has adverse effects on adult behavior and physiology (Peleg☐Raibstein and Feldon, 2011; Brydges et al., 2014). However, stress administered during the juvenile period does not negatively impact behavior during adolescence, a period in which plasticity is still heightened, supporting the protective role of developmental plasticity during periods of stress (Peleg☐Raibstein and Feldon, 2011). Interestingly, genetic mouse models in which developmental windows of plasticity do not close show altered behavioral responses to adverse experience (Ribic, 2020), from impairments in fear learning (Park et al., 2016; Banerjee et al., 2017) and juvenile-like responses during fear conditioning paradigms (Bhagat et al., 2016; Gogolla et al., 2009), to facilitated fear erasure (Yang et al., 2016) and fear-conditioned response (Miwa et al., 2006). While these studies indicate a complex association between plasticity and adverse experiences, they also point to a protective role of critical period plasticity after fear-inducing events and suggest that reintroducing juvenile/adolescent-like plasticity in adult brain could promote stress resilience.

To begin to address this, we tested the depressive-like and anxiety-like behaviors after a week-long bout of unpredictable stress in mice during early adolescence (postnatal days 28-35), an age at which cortical areas still display critical period plasticity (Ribic, 2020). We further tested if adult mice with extended critical period plasticity through deletion of SynCAM 1 (Synaptic Cell Adhesion Molecule 1)(Ribic et al., 2019) respond to stress in a manner similar to adolescent mice. SynCAM 1 is an immunoglobulin domain-containing Type I transmembrane protein that organizes excitatory synapses (Perez de Arce et al. 2015). SynCAM 1 protein is expressed throughout the brain, with expression gradually increasing during postnatal development and plateauing in adulthood (Ribic et al. 2019; Robbins et al. 2010; Thomas et al. 2008; Fogel et al. 2007). Constitutive loss of SynCAM 1 enhances hippocampal long-term depression *ex vivo* (Robbins et al. 2010) and, much like antidepressants and environmental enrichment, shifts the quality of *in vivo* plasticity in the adult visual cortex to a juvenile/adolescent-like state (Ribic et al. 2019). Further, adult SynCAM 1 knockout (KO) are resilient to fear conditioning (Park et al. 2016), suggesting that loss of SynCAM 1 plays a protective role during traumatic experiences.

Using a battery of physiological and behavioral assays, in this study we describe the effects of short-term unpredictable stress on the physiology, as well as depressive-like and anxiety-like behaviors of young and adult wild type (WT) male and female C57BL/6 mice and adult male and

female SynCAM 1 KO mice, to test the association between the quality of plasticity (adult vs
juvenile/adolescent like) and behavioral resilience to stress.

**Methods**

*Mice.* All mice were maintained on C57BL/6 background (The Jackson Laboratory, Bar Harbor, ME)
on a reverse 12:12 light:dark cycle, with food and water *ad libitum.* Young animals from both sexes
were used during the 4[th] and 5[th] week after birth. Adult animals from both sexes were used from 2-4
months of age. SynCAM 1 KO mice (Fujita et al., 2006) and their WT littermates were maintained
on a C57BL/6 background and used at 2-4 months of age. Animals were randomly assigned to
experimental groups and littermates were group housed for the duration of experiments, except
during social isolation. Animals from multiple litters were included in the study to avoid any litter
effects. Animals were treated in accordance with the University of Virginia Institutional Animal Care
and Use Committee guidelines.

*Unpredictable stress.* Mice underwent unpredictable stress over a course of one week, which
consisted of the following stressors: physical restraint in a 50 ml falcon tube for adult mice or 20 ml
plastic scintillation vial for young mice (3 times a week for 1 hr), placement in a cage with no
bedding (2 times a week for 2 hrs) or wet bedding (250 ml of distilled water added to a standard cage
with bedding 2 times a week for 2 hrs), tilting the cage at a 20º angle (2 times a week for 2 hrs),
shaking the cage lightly on an orbital shaker (150 rpm, 2 times a week for 2 hrs) and exposure to 37
centigrade temperature (40% humidity, 1 time for 10 mins). For males, social defeat (2 times a week
for the time needed to defeat) was used as a social stressor, and social isolation (2 times a week for
12 hrs) was used for females. Social defeat was terminated as soon as intruder (test) mice display
signs of social avoidance of the resident (dominant) mouse to avoid any physical wounding. Stressors
were pseudorandomized (random order that was kept constant between all cohorts of mice). Before
the stress week, all mice underwent a control week during which body weight and core body
temperatures were measured at the same times as during stress week.

*Physiological measurements.* Animals were weighed daily at the beginning of the dark cycle. Core
body temperature was measured using a rectal probe (Kent Scientific, Torrington, CT) inserted 2 cm
deep and read as soon as the temperature stabilized. Temperature measurements were taken before
the onset of stress and 15-20 minutes after the onset of stress. For control animals, temperature
measurements were taken 2 successive times, 15-20 minutes apart, at the same times of day as during
stress. Mice were returned to their home cage in between measurements. All temperature differences
were averaged per animal before the statistical comparison.

*Adrenal gland isolation.* For adrenal gland isolation, mice were anesthetized with a mixture of
ketamine and xylazine and transcardially perfused with warm 0.1 M phosphate buffer, followed by
warm 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA). Kidneys and adrenals
were isolated and postfixed overnight in 4% paraformaldehyde and washed overnight in 0.1 M
phosphate buffer. Glands were isolated using a stereomicroscope (Olympus, Tokyo, Japan) and
weighed using a precision scale (Mettler Toledo, Columbus, OH). Weight from both glands was
averaged and divided by the animal's weight before the statistical comparison.

*Tail suspension.* Mice were suspended by their tail on a bar 55 cm above a platform for 6 minutes
and recorded using a high frame rate camera (GoPro, San Mateo, CA), as previously described (Can
et al., 2012). Videos were manually scored by experimenters blind to the experimental group being
analyzed. Mice were considered immobile if no movement was detected for 3 seconds.

*Open field.* Mice were habituated to the room 30 minutes before the behavioral monitoring started.
Individual mice were released in a 50x50x50 (all in cm) plexiglass arena with an overhead high
frame rate camera (GoPro), and their activity was recorded for 10 minutes. Data was analyzed using
MATLAB (Natick, MA)(Zhang et al., 2020).

*Experimental design and data analysis.* The experimenters were blind to the genotype or group of
animals used during data analysis. Between-subject design was used for physiological measures to
allow for the collection of adrenal glands, and within subject design was used for behavioral
measures as previous studies suggest no effect of repeated behavioral assays (such as tail suspension)
on mouse behavior (Cnops et al. 2022). All data was first compared using 3-way ANOVA (factors:
stress, plasticity level and genotype) using DATAtab (Graz, Austria), and followed up using 2-way
ANOVA in JASP (JASP Team, 2024) if no 3-way interaction was detected. For 2-way ANOVAs, all
data was first tested for interactions between stress and genotype as factors, as well as for the effect
of genotype alone if 2-way interaction was not detected. After that, the data was tested for
interactions between stress and age, followed by tests for interactions between stress and plasticity,
with adolescent WT mice and SynCAM 1 KO mice grouped together into a "high plasticity" group.
All data are reported as mean ± SEM or mean ± SD, as indicated, where N represents number of
animals used. Target power for all sample sizes was 0.8. In all cases, alpha was set to 0.05.

**Results**

**Short bout of unpredictable stress induces robust hyperthermia**

As age correlates with neuronal resilience to stress-induced loss of dendritic complexity
(Bloss et al., 2010), we hypothesized that the type of plasticity present in the young brain promotes
resilience to stress. To test this, we administered a short bout of unpredictable stress to young mice
during the 5[th] postnatal week, when the levels of cortical developmental plasticity are high (Gordon
and Stryker, 1996). We did the same to adult (> 2 months old) mice that do not express SynCAM 1
(Synaptic Cell Adhesion Molecule 1), as they display critical period-like cortical plasticity (Ribic et
al., 2019). Our stress paradigm was based on the well-established chronic unpredictable mild stress
paradigm (Willner et al., 1987), where male and female adult and adolescent wild type (WT), as well
as adult SynCAM 1 knock out (KO) mice underwent a pseudo-randomized array of sex-specific and
sex-nonspecific stressors over 1 week (Figure 1A). The sex-specific stressors included social
isolation for female mice, i.e. group-housed female mice were separated to individual cages for five
hours at a time, and social defeat for male mice, i.e. male mice are placed in the resident cage of an
aggressor male mouse until they displayed signs of defeat (Golden et al., 2011). The sex nonspecific
stressors included restraint, wet bedding, no bedding, cage tilt, cage shaking (at 150 rpm) and heat
exposure (37ºC). The length of stress exposure was kept at 1 week to limit it to the peak of critical
period plasticity in the cortex (typically the 5[th] postnatal week)(Gordon and Stryker, 1996; Ribic,
2020)

To test if our stress paradigm indeed induced a physiological stress response, we compared 3
different physiological measures of stress response using a between-subject design (adrenal gland
weight, body weight and core body temperature) across different groups of mice: control and stressed
WT adults, young WT mice and SynCAM 1 KO mice (Figure 1). For adrenal glands, we normalized
their weight to the total body weight of each mouse to account for size differences between young
and adult mice, as well as between males and females. A 2-way ANOVA was performed to evaluate
the effects of stress and quality of plasticity on adrenal gland weight, where adolescent WT and adult
SynCAM 1 KO mice were assigned to a group with "high plasticity" (Tables 1 and 2). We found no
significant interaction between stress and genotype ($F_{(1,80)}$=2.095, p=0.152) and no significant effect

of genotype alone (p=0.072) on adrenal gland weight. When we tested the interaction between stress
and plasticity, we found it to be significant (Table 2). However, further analysis of data revealed that
age significantly interacts with stress (p=0.002), and that stress significantly increased adrenal gland
weight in adolescent mice of both sexes (Figure 1C; adult mice control vs stress p=0.967, young mice
control vs stress p=0.003; Supplementary table 1), but not in adult mice, in agreement with a
previous study that demonstrated no changes in adrenal weights after 1 week of unpredictable stress
(Luo et al., 2025). As previously reported (Bielohuby et al., 2007), adrenal glands were larger in
females than in males across all groups (Supplementary Table 1).

It is well established that different types of stress paradigms in rodents result in reduced body
weight (Patterson and Abizaid, 2013). As young mice at baseline are significantly smaller than adult
mice of both genotypes (young males=15.95±0.6 g, N=10; young females=13.46±0.23 g, N=13; adult
males=22.25±0.42 g, N=11; adult females=17.10±0.44 g, N=12; SynCAM 1 KO males=21.34±0.71
g, N=7; SynCAM1 KO females=17.5±0.85 g, N=6), we compared the difference between body
weights at the beginning and at the end of the week in control and stressed mice instead of comparing
the body weights themselves (Figure 1D). As with adrenal weights, we found no interaction between
stress and genotype ($F_{(1,116)}$, p=0.65) and no effect of genotype alone (p=0.951) on changes in body
weight during one week. While the interaction between stress and plasticity was not significant either
(Table 2), the interaction between stress and age was (p=0.029), with young stressed mice showing a
significant reduction in body weight gain compared to control mice regardless of sex (Figure 1D;
adult mice control vs stress p=0.495, young mice control vs stress p=0.004; Supplementary table 1)

Stress induces a robust elevation in core body temperature within 15 minutes, a phenomenon
known as stress-induced hyperthermia (SIH)(Vinkers et al., 2008). As our stress paradigm resulted in
age-specific adrenal gland weight (Figure 1B) and body weight (Figure 1C) changes, we tested if SIH
is present 15-20 minutes after the introduction of each stressor in our paradigm in all groups of mice
to ensure that adult mice indeed had a physiological response to our stress paradigm. Rectal readings
of core body temperatures can induce stress as well (Zethof et al., 1994), so we compared the
differences in temperature (ΔT) before and after the onset of stress to ΔT between two rectal
temperature measurements 15-20 minutes apart taken from control mice not exposed to short
unpredictable stress (controls; Figure 1D). As expected, we found a robust elevation of core body
temperature in stressed mice of both ages and genotypes that was significantly higher than the ΔT of
control mice (Tables 1 and 2). While the interaction between stress and genotype was also significant
($F_{(1,102)}$=9.264, p=0.003), post-hoc tests (Tukey) revealed no significant differences between WT and
KO control mice, or between WT and KO stressed mice, and no effect of genotype alone (p=0.903).
Interactions between stress and age, as well as stress and sex were not significant (p=0.338 and
p=0.68, respectively). Altogether, our results demonstrate that a short bout of unpredictable stress
induces a robust elevation of core body temperature in all mice, as well as an increase in adrenal
gland weight and a reduction in body weight gain in young mice, demonstrating an efficacy of our
stress paradigm and indicating age-specific effects of short unpredictable stress on bodily response.

**Stress does not result in depressive-like behaviors in mice with open cortical critical periods**

Once we established that our stress paradigm induces a robust physiological response (Figure
1), we tested how mice in whom developmental critical periods of plasticity are still open, namely
young wild type (WT) and adult SynCAM 1 KO mice, behaviorally respond to stress compared to
adult WT mice. We used two well-established behavioral tests to measure depressive-like and
anxiety-like behaviors: tail suspension test (TST) and open field test (OFT; respectively)(Figure 2
and Supplementary Figure 1)(Cryan et al., 2005; Can et al., 2012; Seibenhener and Wooten, 2015).

TST is a measure of learned helplessness-like behavior in rodents that quantifies the fraction
of time spent immobilized after exposure to a stressful stimulus, as well as the latency to
immobilization. The 2-way interaction between stress and plasticity was significant for the fraction of
time spent immobilized (p=0.000005; Table 2) and borderline significant for the latency to first
immobilization (p=0.051; Table 2), with post-hoc (Tukey) tests revealing a significant, almost
uniform increase in the fraction of time spent immobilized and a significant reduction in time lapsed
to first immobilization only in adult WT mice (Figures 2A and 2B; Tables 1 and 2). Interactions
between stress and age, as well as stress and sex were not significant (% immobilization: p=0.098
and p=0.55, respectively; latency to first immobilization: p=0.156 and p=0.56, respectively).
Genotype had an effect % immobilization (p=0.017) and post-hoc tests revealed that the difference in
immobility between stressed SynCAM 1 KO mice compared to WT mice drives this effect. Genotype
had no effect on latency to first immobilization (p=0.18).

To test if our stress paradigm increases anxiety-like behaviors, we compared the preference of
mice to stay close to the walls of the open field arena (thigmotaxis, Tables 1 and 2; Supplementary
Figure 1A) as well as their locomotor activity (Tables 1 and 2; Supplementary Figure 1B).
Interestingly, we found that short unpredictable stress had no impact on thigmotaxis, while distance
travelled in the open field arena was mildly increased in stressed mice from all subgroups, in
agreement with previous findings (Sequeira☐Cordero et al., 2019). While the interactions between
stress and genotype, as well as stress and age were not significant (p=0.55 and p=0.967,
respectively), level of plasticity and genotype alone had significant effects on distance travelled in the
open field arena, with SynCAM 1 KO mice traversing the longest distance (Tukey post-hoc; stressed
KO vs WT mice, p=0.019), in agreement with increased locomotor activity in these mice (Giza et al.,
2013). Locomotor activity in all mouse groups did not correlate with the fraction of time spent
immobilized in TST, suggesting that increased locomotor activity of SynCAM 1 KO mice is not the
cause of their low immobility in TST assay (Pearson r adult=-0.07, p=0.76; young=0.04, p=0.89;
SynCAM 1 KO=-0.02, p=0.91). Altogether, our results demonstrated that short unpredictable stress
robustly increases depressive-like behaviors in adult mice, but not in young mice or adult mice with
critical period-like cortical plasticity.

## Discussion

Since Hubel and Wiesel reported a temporally restricted window of plasticity in the kitten
cortex during which neurons robustly respond to changes in sensory input (Hubel and Wiesel, 1970),
many studies have attempted to restore that type of plasticity to the adult brain aiming to mitigate the
damage caused by a myriad of stressors (Ribic, 2020; Hensch and Quinlan, 2018; Levelt and
Hübener, 2012). In support of this longstanding notion, antidepressants can restore juvenile-like
plasticity to the adult cortex (Maya Vetencourt et al., 2008), but it is still unclear if they promote
recovery from stress through changing the quality of plasticity in the adult brain. Our study attempted
to shed more light on this issue by testing if short unpredictable stress results in depressive-like
behavior in adolescent mice in whom the cortical window of plasticity is still open, as well as in adult
mice in whom the plasticity windows are extended into adulthood through deletion of SynCAM 1
(Ribic et al., 2019). We found that this is indeed the case, supporting that interventions that change
the quality of plasticity in the adult brain can facilitate stress resilience (Nestler and Russo, 2024;
McEwen, 2016; Fuchs et al., 2004).

The duration and type of stressors, along with age and sex of animals, can have a significant
impact on physiological measures of stress, as evident in our results as well (Luo et al., 2025;
Monteiro et al., 2015). Our stress paradigm abrogated weight gain in young mice, but not in adults,

indicating the sensitivity of bodily homeostasis to stress in maturing mice. Similarly, adrenal gland
weight was increased in young mice only, suggesting that the hypothalamic-pituitary-adrenal (HPA)
axis is uniquely sensitive to stress in young mice and may mediate the abrogated weight gain during
stress in these mice. While our stress paradigm did not impact weight gain and adrenal weights of
adult mice, all 3 groups of mice displayed robust stress-induced hyperthermia, indicating that our
protocol was stressful to them. In agreement, adult mice displayed an increase in depressive-like
behaviors as measured with tail suspension test (TST). Interestingly, adolescent mice and adult
SynCAM 1 KO mice did not show any changes in immobilization after stress, in support of the
notion that juvenile/adolescent-like plasticity promotes resilience to stress. The increase in
immobilization in WT mice was almost uniform, while young WT and adult SynCAM 1 KO mice
displayed significant variability that was not sex-dependent, suggesting a gradation of resilience in
these groups of mice. Future studies need to address if the variability in depressive-like measures
after stress in young WT and adult SynCAM 1 KO mice correlates with the variability in the level of
cortical plasticity.  While it is possible that increased locomotor activity of SynCAM 1 KO mice
results in reduced immobilization during TST, we found no correlation between immobility and
locomotor activity in the open field arena, indicating that young mice and adult SynCAM 1 mice
indeed are resilient to short unpredictable stress. Based on our and other studies (Hu et al., 2025),
Future studies can now address the impact of longer stress duration or more severe stressors on the
behavior of adolescent and adult SynCAM 1 KO mice, and whether juvenile/adult-like plasticity
provides a lasting protection from stress.

Unlike immobilization, locomotor activity was mildly impacted by stress, indicated by
increased distance travelled in the open field arena in all three groups of mice. While post-hoc
comparisons of controlled and stressed mice within different subgroups were not statistically
significant, our results are in line with previous research that demonstrated stress-associated
hyperlocomotion (Sequeira☐Cordero et al., 2019). Distance travelled positively correlates with
corticosterone levels in mice head-fixed above a treadmill, which itself is stressful (Juczewski et al.,
2020), suggesting that increased locomotion is a stress response. Locomotor activity is traditionally
classified as anxiety-like behavior, indicating that juvenile/adolescent-like plasticity may not promote
resilience to anxiety after stress. However, future studies need to address this issue using paradigms
that can assess anxiety-like phenotype independent of locomotion, like novelty suppressed feeding
(NSF)(Koskinen and Hovatta, 2023).

While the type of plasticity in our study correlates with resilience to short unpredictable
stress, it is possible that the resilience of SynCAM 1 KO mice is due to another mechanism.
SynCAM 1 gene expression is not restricted to the brain (Fujita et al., 2006) and, while SynCAM 1
protein is localized to the membrane, it can trigger multiple signaling cascades that promote neuronal
resilience to stress (Cheadle and Biederer, 2012; Murakami et al., 2014). However, we find this
unlikely, as SynCAM 1 KO mice display overall accelerated learning in spatial tasks and juvenile-
like responses to noxious stimuli (Park et al., 2016; Robbins et al., 2010). SynCAM 1 is a potent
plasticity brake in the adult brain and even a transient knock-down of its expression can restore
juvenile/adolescent like plasticity in the adult primary visual cortex (Ribic et al., 2019). Our study is
hence in line with the notion that critical period-like plasticity is protective of stress and provides a
rationale for further inquiries into the link between resilience and different types of plasticity.

Our study is limited in its scope, but our findings unambiguously demonstrate the efficacy of
short unpredictable stress in inducing depressive-like behaviors in adult mice. The short duration and
ease of administering our protocol will undoubtedly facilitate further studies of short unpredictable
stress and its long-term effects on brain and behavior. Importantly, our study provides indirect

evidence that heightened brain plasticity promotes resilience to stress, and provides a rationale for
future, more in-depth mechanistic inquiries into the relationship between plasticity and stress.

**Conflict of Interest**

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

**Author Contributions**

AR and RW conceived and designed the study. RW, CVDO, ENL, SCF, GLO and AR collected and
analyzed the data. AR wrote the manuscript with input from all authors.

**Funding**

This study was funded through the Department of Psychology at the University of Virginia.

**Acknowledgments**

The authors would like to thank the Ribic Lab for critical reading of the manuscript.

**Data Availability Statement**

The datasets generated and analyzed in this study will be provided upon reasonable request.

**References**

Banerjee, S. B., Gutzeit, V. A., Baman, J., Aoued, H. S., Doshi, N. K., Liu, R. C., and Ressler, K. J.
(2017). Perineuronal nets in the adult sensory cortex are necessary for fear learning. *Neuron* 95,
169-179.e3. doi:10.1016/j.neuron.2017.06.007.

Bhagat, S. M., Butler, S. S., Taylor, J. R., McEwen, B. S., and Strittmatter, S. M. (2016). Erasure of
fear memories is prevented by Nogo Receptor 1 in adulthood. *Mol. Psychiatry* 21, 1281–1289.
doi:10.1038/mp.2015.179.

Bielohuby, M., Herbach, N., Wanke, R., Maser-Gluth, C., Beuschlein, F., Wolf, E., and Hoeflich, A.
(2007). Growth analysis of the mouse adrenal gland from weaning to adulthood: time- and gender-
dependent alterations of cell size and number in the cortical compartment. *Am. J. Physiol.*
*Endocrinol. Metab.* 293, E139-46. doi:10.1152/ajpendo.00705.2006.

Bloss, E. B., Janssen, W. G., McEwen, B. S., and Morrison, J. H. (2010). Interactive effects of stress
and aging on structural plasticity in the prefrontal cortex. *J. Neurosci.* 30, 6726–6731.
doi:10.1523/JNEUROSCI.0759-10.2010.

Brydges, N. M., Jin, R., Seckl, J., Holmes, M. C., Drake, A. J., and Hall, J. (2014). Juvenile stress
enhances anxiety and alters corticosteroid receptor expression in adulthood. *Brain Behav.* 4, 4–13.
doi:10.1002/brb3.182.

Can, A., Dao, D. T., Terrillion, C. E., Piantadosi, S. C., Bhat, S., and Gould, T. D. (2012). The tail
suspension test. *J. Vis. Exp.*, e3769. doi:10.3791/3769.

Cheadle, L., and Biederer, T. (2012). The novel synaptogenic protein Farp1 links postsynaptic
cytoskeletal dynamics and transsynaptic organization. *J. Cell Biol.* 199, 985–1001.
doi:10.1083/jcb.201205041.

Cryan, J. F., Mombereau, C., and Vassout, A. (2005). The tail suspension test as a model for
assessing antidepressant activity: review of pharmacological and genetic studies in mice. *Neurosci.*
*Biobehav. Rev.* 29, 571–625. doi:10.1016/j.neubiorev.2005.03.009.

Czéh, B., Fuchs, E., Wiborg, O., and Simon, M. (2016). Animal models of major depression and their
clinical implications. *Prog Neuropsychopharmacol Biol Psychiatry* 64, 293–310.
doi:10.1016/j.pnpbp.2015.04.004.

Faye, C., Mcgowan, J. C., Denny, C. A., and David, D. J. (2018). Neurobiological mechanisms of
stress resilience and implications for the aged population. *Curr. Neuropharmacol.* 16, 234–270.
doi:10.2174/1570159X15666170818095105.

Fuchs, E., Czéh, B., Kole, M. H. P., Michaelis, T., and Lucassen, P. J. (2004). Alterations of
neuroplasticity in depression: the hippocampus and beyond. *Eur. Neuropsychopharmacol.* 14
Suppl 5, S481-90. doi:10.1016/j.euroneuro.2004.09.002.

Fujita, E., Kouroku, Y., Ozeki, S., Tanabe, Y., Toyama, Y., Maekawa, M., Kojima, N., Senoo, H.,
Toshimori, K., and Momoi, T. (2006). Oligo-astheno-teratozoospermia in mice lacking
RA175/TSLC1/SynCAM/IGSF4A, a cell adhesion molecule in the immunoglobulin superfamily.
*Mol. Cell. Biol.* 26, 718–726. doi:10.1128/MCB.26.2.718-726.2006.

Giza, J. I., Jung, Y., Jeffrey, R. A., Neugebauer, N. M., Picciotto, M. R., and Biederer, T. (2013). The
synaptic adhesion molecule SynCAM 1 contributes to cocaine effects on synapse structure and
psychostimulant behavior. *Neuropsychopharmacology* 38, 628–638. doi:10.1038/npp.2012.226.

Gogolla, N., Caroni, P., Lüthi, A., and Herry, C. (2009). Perineuronal nets protect fear memories
from erasure. *Science* 325, 1258–1261. doi:10.1126/science.1174146.

Golden, S. A., Covington, H. E., Berton, O., and Russo, S. J. (2011). A standardized protocol for
repeated social defeat stress in mice. *Nat. Protoc.* 6, 1183–1191. doi:10.1038/nprot.2011.361.

Gordon, J. A., and Stryker, M. P. (1996). Experience-dependent plasticity of binocular responses in
the primary visual cortex of the mouse. *J. Neurosci.* 16, 3274–3286. doi:10.1523/JNEUROSCI.16-
10-03274.1996.

Greifzu, F., Kalogeraki, E., and Löwel, S. (2016). Environmental enrichment preserved lifelong
ocular dominance plasticity, but did not improve visual abilities. *Neurobiol. Aging* 41, 130–137.
doi:10.1016/j.neurobiolaging.2016.02.014.

Greifzu, F., Pielecka-Fortuna, J., Kalogeraki, E., Krempler, K., Favaro, P. D., Schlüter, O. M., and
Löwel, S. (2014). Environmental enrichment extends ocular dominance plasticity into adulthood
and protects from stroke-induced impairments of plasticity. *Proc Natl Acad Sci USA* 111, 1150–
1155. doi:10.1073/pnas.1313385111.

Hegde, A., Suresh, S., and Mitra, R. (2020). Early-life short-term environmental enrichment
counteracts the effects of stress on anxiety-like behavior, brain-derived neurotrophic factor and
nuclear translocation of glucocorticoid receptors in the basolateral amygdala. *Sci. Rep.* 10, 14053.
doi:10.1038/s41598-020-70875-5.

Hensch, T. K., and Quinlan, E. M. (2018). Critical periods in amblyopia. *Vis. Neurosci.* 35, E014.
doi:10.1017/S0952523817000219.

Hubel, D. H., and Wiesel, T. N. (1970). The period of susceptibility to the physiological effects of
unilateral eye closure in kittens. *J Physiol (Lond)* 206, 419–436.
doi:10.1113/jphysiol.1970.sp009022.

Hu, W., Jiang, L., Wang, Q., Hu, Q., Zhong, T., Wu, J., Chen, X., and Liu, T. (2025). Chronic
unpredictable stress during adolescence exerts sex-specific effects on depressive-like behavior and
neural activation triggered by tail suspension test. *Behav. Brain Res.* 477, 115314.
doi:10.1016/j.bbr.2024.115314.

Juczewski, K., Koussa, J. A., Kesner, A. J., Lee, J. O., and Lovinger, D. M. (2020). Stress and
behavioral correlates in the head-fixed method: stress measurements, habituation dynamics,
locomotion, and motor-skill learning in mice. *Sci. Rep.* 10, 12245. doi:10.1038/s41598-020-69132-
6.

Koskinen, M.-K., and Hovatta, I. (2023). Genetic insights into the neurobiology of anxiety. *Trends
Neurosci.* 46, 318–331. doi:10.1016/j.tins.2023.01.007.

Lehmann, M. L., and Herkenham, M. (2011). Environmental enrichment confers stress resiliency to
social defeat through an infralimbic cortex-dependent neuroanatomical pathway. *J. Neurosci.* 31,
6159–6173. doi:10.1523/JNEUROSCI.0577-11.2011.

Levelt, C. N., and Hübener, M. (2012). Critical-period plasticity in the visual cortex. *Annu. Rev.
Neurosci.* 35, 309–330. doi:10.1146/annurev-neuro-061010-113813.

Luo, Y., Zhang, Y., Zhao, Y., Chen, F., Li, X., Liu, X., Shakir, M. Z., Jiang, N., and Shan, C. (2025).
Chronic Unpredictable Mild Stress-Induced Anxiety is Linked to Inflammatory Responses and
Disruptions in Tryptophan Metabolism in Male C57BL/6 N Mice. *Behav. Brain Res.*, 115506.
doi:10.1016/j.bbr.2025.115506.

Macartney, E. L., Lagisz, M., and Nakagawa, S. (2022). The relative benefits of environmental
enrichment on learning and memory are greater when stressed: A meta-analysis of interactions in
rodents. *Neurosci. Biobehav. Rev.* 135, 104554. doi:10.1016/j.neubiorev.2022.104554.

Maya Vetencourt, J. F., Sale, A., Viegi, A., Baroncelli, L., De Pasquale, R., O'Leary, O. F., Castrén,
E., and Maffei, L. (2008). The antidepressant fluoxetine restores plasticity in the adult visual
cortex. *Science* 320, 385–388. doi:10.1126/science.1150516.

McEwen, B. S., Bowles, N. P., Gray, J. D., Hill, M. N., Hunter, R. G., Karatsoreos, I. N., and Nasca,
C. (2015). Mechanisms of stress in the brain. *Nat. Neurosci.* 18, 1353–1363. doi:10.1038/nn.4086.

McEwen, B. S., and Morrison, J. H. (2013). The brain on stress: vulnerability and plasticity of the
prefrontal cortex over the life course. *Neuron* 79, 16–29. doi:10.1016/j.neuron.2013.06.028.

McEwen, B. S. (2012). Brain on stress: how the social environment gets under the skin. *Proc Natl*
*Acad Sci USA* 109 Suppl 2, 17180–17185. doi:10.1073/pnas.1121254109.

McEwen, B. S. (2016). In pursuit of resilience: stress, epigenetics, and brain plasticity. *Ann. N. Y.*
*Acad. Sci.* 1373, 56–64. doi:10.1111/nyas.13020.

Miwa, J. M., Stevens, T. R., King, S. L., Caldarone, B. J., Ibanez-Tallon, I., Xiao, C., Fitzsimonds, R.
M., Pavlides, C., Lester, H. A., Picciotto, M. R., et al. (2006). The prototoxin lynx1 acts on
nicotinic acetylcholine receptors to balance neuronal activity and survival in vivo. *Neuron* 51,
587–600. doi:10.1016/j.neuron.2006.07.025.

Monteiro, S., Roque, S., de Sá-Calçada, D., Sousa, N., Correia-Neves, M., and Cerqueira, J. J.
(2015). An efficient chronic unpredictable stress protocol to induce stress-related responses in
C57BL/6 mice. *Front. Psychiatry* 6, 6. doi:10.3389/fpsyt.2015.00006.

Murakami, S., Sakurai-Yageta, M., Maruyama, T., and Murakami, Y. (2014). Trans-homophilic
interaction of CADM1 activates PI3K by forming a complex with MAGuK-family proteins MPP3
and Dlg. *PLoS ONE* 9, e110062. doi:10.1371/journal.pone.0110062.

Nestler, E. J., and Russo, S. J. (2024). Neurobiological basis of stress resilience. *Neuron* 112, 1911–
1929. doi:10.1016/j.neuron.2024.05.001.

Park, K. A., Ribic, A., Laage Gaupp, F. M., Coman, D., Huang, Y., Dulla, C. G., Hyder, F., and
Biederer, T. (2016). Excitatory synaptic drive and feedforward inhibition in the hippocampal CA3
circuit are regulated by syncam 1. *J. Neurosci.* 36, 7464–7475. doi:10.1523/JNEUROSCI.0189-
16.2016.

Patterson, Z. R., and Abizaid, A. (2013). Stress induced obesity: lessons from rodent models of
stress. *Front. Neurosci.* 7, 130. doi:10.3389/fnins.2013.00130.

Peleg-Raibstein, D., and Feldon, J. (2011). Differential effects of post-weaning juvenile stress on the
behaviour of C57BL/6 mice in adolescence and adulthood. *Psychopharmacology (Berl)* 214, 339–
351. doi:10.1007/s00213-010-1991-8.

Phoumthipphavong, V., Barthas, F., Hassett, S., and Kwan, A. C. (2016). Longitudinal effects of
ketamine on dendritic architecture in vivo in the mouse medial frontal cortex. *eNeuro* 3.
doi:10.1523/ENEURO.0133-15.2016.

Ribic, A., Crair, M. C., and Biederer, T. (2019). Synapse-Selective Control of Cortical Maturation
and Plasticity by Parvalbumin-Autonomous Action of SynCAM 1. *Cell Rep.* 26, 381-393.e6.
doi:10.1016/j.celrep.2018.12.069.

Ribic, A. (2020). Stability in the Face of Change: Lifelong Experience-Dependent Plasticity in the
Sensory Cortex. *Front. Cell. Neurosci.* 14, 76. doi:10.3389/fncel.2020.00076.

Robbins, E. M., Krupp, A. J., Perez de Arce, K., Ghosh, A. K., Fogel, A. I., Boucard, A., Südhof, T.
C., Stein, V., and Biederer, T. (2010). SynCAM 1 adhesion dynamically regulates synapse number
and impacts plasticity and learning. *Neuron* 68, 894–906. doi:10.1016/j.neuron.2010.11.003.

Russo, S. J., Murrough, J. W., Han, M.-H., Charney, D. S., and Nestler, E. J. (2012). Neurobiology of
resilience. *Nat. Neurosci.* 15, 1475–1484. doi:10.1038/nn.3234.

Seibenhener, M. L., and Wooten, M. C. (2015). Use of the Open Field Maze to measure locomotor
and anxiety-like behavior in mice. *J. Vis. Exp.*, e52434. doi:10.3791/52434.

Sequeira-Cordero, A., Salas-Bastos, A., Fornaguera, J., and Brenes, J. C. (2019). Behavioural
characterisation of chronic unpredictable stress based on ethologically relevant paradigms in rats.
*Sci. Rep.* 9, 17403. doi:10.1038/s41598-019-53624-1.

Vinkers, C. H., van Bogaert, M. J. V., Klanker, M., Korte, S. M., Oosting, R., Hanania, T., Hopkins,
S. C., Olivier, B., and Groenink, L. (2008). Translational aspects of pharmacological research into
anxiety disorders: the stress-induced hyperthermia (SIH) paradigm. *Eur. J. Pharmacol.* 585, 407–
425. doi:10.1016/j.ejphar.2008.02.097.

Willner, P., Towell, A., Sampson, D., Sophokleous, S., and Muscat, R. (1987). Reduction of sucrose
preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant.
*Psychopharmacology (Berl)* 93, 358–364. doi:10.1007/BF00187257.

Yang, W.-Z., Liu, T.-T., Cao, J.-W., Chen, X.-F., Liu, X., Wang, M., Su, X., Zhang, S.-Q., Qiu, B.-
L., Hu, W.-X., et al. (2016). Fear erasure facilitated by immature inhibitory neuron transplantation.
*Neuron* 92, 1352–1367. doi:10.1016/j.neuron.2016.11.018.

Zethof, T. J., Van der Heyden, J. A., Tolboom, J. T., and Olivier, B. (1994). Stress-induced
hyperthermia in mice: a methodological study. *Physiol. Behav.* 55, 109–115. doi:10.1016/0031-
9384(94)90017-5.

Zhang, C., Li, H., and Han, R. (2020). An open-source video tracking system for mouse locomotor
activity analysis. *BMC Res. Notes* 13, 48. doi:10.1186/s13104-020-4916-6.

**Figure Legends**

**Figure 1. Short unpredictable stress induces robust hyperthermia. A)** Experimental
schematics. Mice of both sexes (adult, postnatal day/P 60 and older; young, P28; adult SynCAM 1
KO) underwent one week of short unpredictable stress with indicated stressors. Control mice of
matching ages were subjected to the same measurements but not exposed to the stressors. Body
weights and core temperatures were measured daily, and adrenal glands were isolated from both
groups of mice after 1 week. **B)** For behavioral assays, mice underwent behavioral assays
immediately prior to the first stressor exposure and after the exposure to the last stressor. **C)** Adrenal
gland weight was significantly increased in young mice subjected to stress, but not in adult mice. **D)**
Short unpredictable abrogated weight gain in young, but not in adult mice. **E)** Short unpredictable
stress resulted in robust elevation of core body temperature in all groups of mice. Medians and
quartiles of the data are indicated by lines and individual points represent mice.

**Figure 2. Young mice and adult SynCAM 1 KO mice are resilient to stress-induced**
**increase learned helplessness-like behavior. A)** Only stressed adult WT mice show increased
immobility after stress in tail suspension test and a significant reduction in time lapsed to first
immobilization **(B).** Medians and quartiles of the data are indicated by lines and individual points and
lines represent mice.

**Supplementary Figure 1. Short unpredictable stress has no impact on anxiety-like**
**behaviors. A)** Short unpredictable stress did not impact thigmotaxis in open field test. **B)** While
stressed mice from all groups travelled more in the open field arena, within-group post-hoc
comparisons were not statistically significant. Medians and quartiles of the data are indicated by lines
and individual points and lines represent mice.

**A** Physiological assays

Adult: >P60
Young: P28
SynCAM 1 KO: >P60

Day 1 — Control — Day 7
Short unpredictable stress

Body weight
Core body temperature
Body weight
Adrenal gland isolation

Restraint
Cage tilt
Shaking
Wet/no bedding
Heat
Social isolation (♀)
Social defeat (♂)

**B** Behavioral assays

Adult: >P60
Young: P28
SynCAM 1 KO: >P60

Day 1 — Short unpredictable stress — Day 7

Open field
Tail suspension
Open field
Tail suspension

**C** Adrenal gland weight (% of body weight)

ns — Adult
** — Young
ns — SynCAM 1 KO

Control
Stress

**D** Δ body weight (g)

Control
Stress

ns — Adult
** — Young
ns — SynCAM 1 KO

**E** Δ core body temperature (°C)

*** — Adult
** — Young
** — SynCAM 1 KO

Control
Stress