Critical Period Plasticity is Associated with Resilience to Short Unpredictable Stress

Robert Williams, Jr[1], Charlie Van Den Oord[2†], Erica N. Lee[1†], Samuel C. Fedde[1], Gia L.
Oscherwitz[3], Adema Ribic[1,*]

[1]Department of Psychology, College of Arts and Sciences, University of Virginia, Charlottesville,
VA

[2]Department of Biology, College of Arts and Sciences, University of Virginia, Charlottesville, VA

[3]School of Engineering, University of Virginia, Charlottesville, VA

[†]Equal contribution

* Correspondence:
Corresponding Author
adema.ribic@virginia.edu

Keywords: plasticity, SynCAM 1, stress, resilience, hyperthermia

Abstract

Exposure to stress can increase the risk of depression in susceptible individuals, but not in resilient
ones. Resilience to stress decreases with age, parallel to drastic changes in the quality of brain
plasticity from juvenile to old age, suggesting that the type of plasticity found in the maturing brain
promotes resilience. To indirectly test this, we administered short unpredictable stress to adult male
and female mice, as well as to adolescent mice and mice that lack SynCAM 1 and display extended
adolescent-like, critical period plasticity in the brain. We found that short unpredictable stress
robustly increased core body temperature in all groups of mice, indicative of stress-induced
hyperthermia (SIH) and confirming the efficacy of the stress paradigm. However, depressive-like
behavior as measured though tail suspension test was increased in adult mice only, supporting that
the type of plasticity found in the maturing brain promotes resilience to stress. All three groups of
mice showed a mild increase in locomotor activity after stress, suggesting that the quality of
plasticity does not correlate with resilience to anxiety-like phenotypes. Our study hence provides
further evidence for the protective role of developmental plasticity during stress and points to new
mechanisms that promote resilience to stress-induced depression.

Introduction

Stressful life events are strongly linked to the onset of depression (McEwen, 2012; McEwen et
al., 2015; Czéh et al., 2016), which is one of the leading causes of disability in the USA (WHO). Yet,
stress does not increase the risk of depression in individuals that are resilient: capable of successfully
adapting to stress exposure (Faye et al., 2018; McEwen and Morrison, 2013; Russo et al., 2012).
Stress resilience is thought to decline with age, parallel to changes in the quality of brain plasticity,

which transitions from robust critical period plasticity during in the developing brain to a restricted,
top-down modulated process in the adult brain (Ribic, 2020). While the link between plasticity and
resilience is still unclear, a previous study demonstrated that chronic restraint stress results in a
reversible reduction in the length of dendritic branches in the prefrontal cortex (PFC) of young, but
not aged rats, in whom the reduction persists (Bloss et al., 2010). Heightened brain plasticity during
postnatal development, as well as interventions that change the quality of plasticity in adulthood, can
hence promote resilience to stress (McEwen, 2016, 2012).

In support of this notion, antidepressants markedly increase plasticity in the adult brain and
change its quality to a more juvenile/adolescent-like (Maya Vetencourt et al., 2008;
Phoumthipphavong et al., 2016). Environmental enrichment, a classical behavioral paradigm that
also "rejuvenates" the adult cortex (Greifzu et al., 2016, 2014), mitigates the effects of stress across
the lifespan (Macartney et al., 2022; Lehmann and Herkenham, 2011; Hegde et al., 2020). However,
both antidepressants and environmental enrichment have systemic effects on the brain, and it is still
unclear if they promote resilience and mitigate the effects of stress through altered quality of brain
plasticity.

In mice, a species of choice for genetic manipulations of plasticity, stress during late postnatal
development has adverse effects on adult behavior and physiology (Peleg☐Raibstein and Feldon,
2011; Brydges et al., 2014). However, stress administered during the juvenile period does not
negatively impact behavior during adolescence, a period in which plasticity is still heightened,
supporting the protective role of developmental plasticity during periods of stress (Peleg☐Raibstein
and Feldon, 2011). Interestingly, genetic mouse models in which developmental windows of
plasticity do not close show altered behavioral responses to adverse experience (Ribic, 2020), from
impairments in fear learning (Park et al., 2016; Banerjee et al., 2017) and juvenile-like responses
during fear conditioning paradigms (Bhagat et al., 2016; Gogolla et al., 2009), to facilitated fear
erasure (Yang et al., 2016) and fear-conditioned response (Miwa et al., 2006). While these studies
indicate a complex association between plasticity and adverse experiences, they also point to a
protective role of critical period plasticity after fear-inducing events and suggest that reintroducing
juvenile/adolescent-like plasticity in adult brain could promote stress resilience.

To begin to address this, we tested the depressive-like and anxiety-like behaviors after a week-
long bout of unpredictable stress in mice during early adolescence (postnatal days 28-35), an age at
which cortical areas still display critical period plasticity (Ribic, 2020). We further tested if adult
mice with extended critical period plasticity through deletion of SynCAM 1 (Synaptic Cell Adhesion
Molecule 1)(Ribic et al., 2019) respond to stress in a manner similar to adolescent mice. Our results
demonstrate that 1 week of unpredictable stress induces a robust physiological response regardless of
age or the level of plasticity in the brain. However, adolescent wild type (WT) and adult SynCAM 1
knock-out (KO) mice are behaviorally resilient to unpredictable stress, unlike adult WT mice that
show increased learned helplessness-like behavior after stress. Our study hence provides new
evidence in support of the protective role of juvenile/adolescent-like plasticity in the aging brain and
suggests that interventions that reintroduce such plasticity in aged brain can promote stress resilience
(McEwen, 2016).

Methods

Mice and data analysis. All mice were maintained on C57BL/6 background (The Jackson
Laboratory, Bar Harbor, ME) on a reverse 12:12 light:dark cycle, with food and water *ad libitum*.
Young animals from both sexes were used during the 4th and 5th week after birth. Adult animals from

both sexes were used from 2-4 months of age. SynCAM 1 KO mice (Fujita et al., 2006) and their WT
littermates were maintained on a C57BL/6 background and used at 2-4 months of age. Animals were
randomly assigned to experimental groups and littermates were group housed for the duration of
experiments, except during social isolation. Animals were treated in accordance with the University
of Virginia Institutional Animal Care and Use Committee guidelines. The experimenters were blind
to the genotype or group of animals used during data analysis. Both within-subject and between-
subject designs were used to allow for the collection of adrenal glands, with physiological and
behavioral measures compared using 2-way ANOVA (GraphPad, San Diego, CA). All data are
reported as mean ± SEM, where N represents number of animals and infants used, unless indicated
otherwise. Target power for all sample sizes was 0.8. In all cases, alpha was set to 0.05.

Unpredictable stress. Mice underwent unpredictable stress over a course of one week, which
consisted of the following stressors: physical restraint in a 50 ml falcon tube for adult mice or 20 ml
plastic scintillation vial for young mice (3 times a week for 1 hr), placement in a cage with no
bedding (2 times a week for 2 hrs) or wet bedding (250 ml of distilled water added to a standard cage
with bedding 2 times a week for 2 hrs), tilting the cage at a 20º angle (2 times a week for 2 hrs),
shaking the cage lightly on an orbital shaker (150 rpm, 2 times a week for 2 hrs) and exposure to 37
centigrade temperature (40% humidity, 1 time for 10 mins). For males, social defeat (2 times a week
for the time needed to defeat) was used as a social stressor, and social isolation (2 times a week for
12 hrs) was used for females. Social defeat was terminated as soon as intruder (test) mice display
signs of social avoidance of the resident (dominant) mouse to avoid any physical wounding. Stressors
were pseudorandomized (random order that was kept constant between all cohorts of mice). Before
the stress week, all mice underwent a control week during which body weight and core body
temperatures were measured at the same times as during stress week.

Physiological measurements. Animals were weighed daily at the beginning of the dark cycle. Core
body temperature was measured using a rectal probe (Kent Scientific, Torrington, CT) inserted 2 cm
deep and read as soon as the temperature stabilized. Temperature measurements were taken before
the onset of stress and 15-20 minutes after the onset of stress. For control animals, temperature
measurements were taken 2 successive times, 15-20 minutes apart, at the same times of day as during
stress. Mice were returned to their home cage in between measurements. All temperature differences
were averaged per animal before the statistical comparison.

Adrenal gland isolation. For adrenal gland isolation, mice were anesthetized with a mixture of
ketamine and xylazine and transcardially perfused with warm 0.1 M phosphate buffer, followed by
warm 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA). Kidneys and adrenals
were isolated and postfixed overnight in 4% paraformaldehyde and washed overnight in 0.1 M
phosphate buffer. Glands were isolated using a stereomicroscope (Olympus, Tokyo, Japan) and
weighed using a precision scale (Mettler Toledo, Columbus, OH). Weight from both glands was
averaged and divided by the animal's weight before the statistical comparison.

Tail suspension. Mice were suspended by their tail on a bar 55 cm above a platform for 6 minutes
and recorded using a high frame rate camera (GoPro, San Mateo, CA), as previously described (Can
et al., 2012). Videos were manually scored by experimenters blind to the experimental group being
analyzed. Mice were considered immobile if no movement was detected for 3 seconds.

Open field. Mice were habituated to the room 30 minutes before the behavioral monitoring started.
Individual mice were released in a 50x50x50 (all in cm) plexiglass arena with an overhead high
frame rate camera (GoPro), and their activity was recorded for 10 minutes. Data was analyzed using
MATLAB (Natick, MA)(Zhang et al., 2020).

Results

Short bout of unpredictable stress induces robust hyperthermia

As age correlates with neuronal resilience to stress-induced loss of dendritic complexity (Bloss et al., 2010), we hypothesized that the type of plasticity present in the young brain promotes resilience to stress. To test this, we administered a short bout of unpredictable stress to young mice during the 5^{th} postnatal week, when the levels of cortical developmental plasticity are high (Gordon and Stryker, 1996). We did the same to adult (> 2 months old) mice that do not express SynCAM 1 (Synaptic Cell Adhesion Molecule 1), as they display critical period-like cortical plasticity (Ribic et al., 2019). Our stress paradigm was based on the well-established chronic unpredictable mild stress paradigm (Willner et al., 1987), where male and female adult and adolescent wild type, as well as adult SynCAM 1 knock out (KO) mice underwent a pseudo-randomized array of sex-specific and sex-nonspecific stressors over 1 week (Figure 1A). The sex-specific stressors included social isolation for female mice, i.e. group-housed female mice were separated to individual cages for five hours at a time, and social defeat for male mice, i.e. male mice are placed in the resident cage of an aggressor male mouse until they displayed signs of defeat (Golden et al., 2011). The sex nonspecific stressors included restraint, wet bedding, no bedding, cage tilt, cage shaking (at 150 rpm) and heat exposure (37ºC). The length of stress exposure was kept at 1 week to limit it to the peak of critical period plasticity in the cortex (typically the 5^{th} postnatal week)(Gordon and Stryker, 1996; Ribic, 2020)

To test if our stress paradigm indeed induced a physiological stress response, we compared 3 different physiological measures of stress response using a between-subject design (adrenal gland weight, body weight and core body temperature) across different groups of mice: control and stressed adults, young mice and SynCAM 1 KO mice (Figure 1A-C). For adrenal glands, we normalized their weight to the total body weight of each mouse to account for size differences between young and adult mice, as well as between males and females. We found that the interaction between stress and age/plasticity level was not significant, with neither of the factors having an impact on adrenal gland weight (Figure 1B and Table 1; 2-way ANOVA, p=0.098, $F_{(2, 61)}$=2.42; p_{stress}=0.67, $F_{(1, 61)}$=0.2; $p_{age/plasticity}$=0.078, $F_{(2, 61)}$=2.66), in agreement with a previous study that demonstrated no changes in adrenal weights after 1 week of unpredictable stress (Luo et al., 2025). However, the young mice displayed a trend towards higher values in the stressed group, which became significant after separating the animals by sex (Figure 1C and Table 1; 2-way ANOVA, stress x sex p=0.51, $F_{(1,15)}$=0.47; p_{stress}=0.003, $F_{(1,15)}$=12.6; p_{sex}<0.0001, $F_{(1,15)}$=58.75; Sidak multiple comparison test males p=0.19, females p=0.004). As previously reported (Bielohuby et al., 2007), adrenal glands were larger in females than in males across all groups (Table 1).

It is well established that different types of stress paradigms in rodents result in reduced body weight (Patterson and Abizaid, 2013). As young mice at baseline are significantly smaller than adult mice of both genotypes (young males=18.15±0.8 g, N=3; young females=13.65±0.37 g, N=5; adult males=22.25±0.42 g, N=11; adult females=17.10±0.44 g, N=12; SynCAM 1 KO males=21.34±0.71 g, N=7; SynCAM1 KO females=17.5±0.85 g, N=6), we compared the difference between body weights at the beginning and at the end of the week in control and stressed mice instead of comparing the body weights themselves (Figure 1D). We found that one week of stress did not result in reduced weight gain in any of the groups (Figure 1D; ΔBW adult control=0.33±0.13 g, N=33; adult stress=0.21±0.097 g, N=25; young control=1.5±0.25 g, N=8; young stress=1.33±0.26 g, N=9; SynCAM 1 KO control=0.91±0.13 g, N=16; SynCAM 1 KO stress 0.43±0.33 g, N=14; 2-way ANOVA stress x age/plasticity p=0.56, $F_{(2,99)}$=0.58; p_{stress}=0.12, $F_{(1,99)}$=2.45; $p_{age/plasticity}$<0.0001,

$F_{(2,99)=}15.58$). The effect of age/plasticity was also significant, post-hoc tests (Holm Sidak) revealed
that it was mostly due to weight changes in young WT mice, as expected (control adult vs young,
p=0.0004; adult vs SynCAM 1 KO, p=0.037; young vs SynCAM 1 KO, p=0.2; stress adult vs young,
p=0.0007; adult vs SynCAM 1 KO, p=0.76; young vs SynCAM 1 KO, p=0.02).

Stress induces a robust elevation in core body temperature within 15 minutes, a phenomenon
known as stress-induced hyperthermia (SIH)(Vinkers et al., 2008). As our stress paradigm resulted in
very small changes in standard measures of bodily stress response, namely adrenal gland weight
(Figure 1B) and body weight (Figure 1C), we tested if SIH is present 15-20 minutes after the
introduction of each stressor in our paradigm (Figure 1D). Rectal readings of core body temperatures
can induce stress as well (Zethof et al., 1994), so we compared the differences in temperature (ΔT)
before and after the onset of stress to ΔT between two rectal temperature measurements 15-20
minutes apart taken from control mice not exposed to short unpredictable stress (controls; Figure
1D). As expected, we found a robust elevation of core body temperature in stressed mice of both ages
and genotypes that was significantly higher than the ΔT of control mice (ΔT adult control=0.04±0.08
ºC, N=24; adult stress=0.54±0.06 ºC, N=19; young control=0.22±0.13 ºC, N=13; young
stress=0.66±0.06 ºC, N=12; SynCAM 1 KO control=-0.07±0.09 ºC, N=6; SynCAM 1 KO
stress=0.41±0.04 ºC, N=32; 2-way ANOVA stress x age/plasticity p=0.95, $F_{(2,100)=}0.05$;
$p_{stress}<0.0001$, $F_{(2,100)=}42.75$; $p_{age/plasticity}=0.02$, $F_{(2,100)=}4.05$). While the effect of age/plasticity was also
significant, post-hoc tests (Holm Sidak) revealed no significant differences between adult, young and
SynCAM 1 KO mice within control or stress groups. Altogether, our results demonstrate that a short
bout of unpredictable stress induces a robust elevation of core body temperature and an increase in
adrenal gland weight in young female mice, indicating the validity of our paradigm.

Stress does not result in depressive-like behaviors in mice with open cortical critical periods

Once we established that our stress paradigm induces a robust physiological response (Figure
1), we tested how mice in whom developmental critical periods of plasticity are still open, namely
young wild type (WT) and adult SynCAM 1 KO mice, behaviorally respond to stress compared to
adult WT mice. We used two well-established behavioral tests to measure depressive-like and
anxiety-like behaviors: tail suspension test (TST) and open field test (OFT; respectively)(Figure
2)(Cryan et al., 2005; Can et al., 2012; Seibenhener and Wooten, 2015).

TST is a measure of learned helplessness-like behavior in rodents that quantifies the fraction
of time spent immobilized after exposure to a stressful stimulus, as well as the latency to
immobilization. As expected, stressed adult WT mice show a significant increase in the fraction of
time spent immobilized (Figure 2A; 2-way repeated measures (RM) ANOVA, stress x age/plasticity
p<0.0001, $F_{(2,45)}=25.24$; $p_{stress}=0.04$, $F_{(1,45)}=4.11$; $p_{age/plasticity}<0.0001$, $F_{(2,45)}=18.3$; adult WT
control=31.79±2.79, stress=53.73±3.5, N=17, p<0.0001; young control=21.74±3.37,
stress=16.75±3.22, N=13, p=0.42; SynCAM 1 KO control=26.66±3.29 s, stress=21±3.44, N=18,
p=0.18; all values in %, Sidak's post-hoc) and a significant reduction in time lapsed to first
immobilization (Figure 2B; 2-way repeated measures (RM) ANOVA, stress x age/plasticity p=0.054,
$F_{(2,45)}=3.12$; $p_{stress}=0.002$, $F_{(1,45)}=10.41$; $p_{age/plasticity}=0.0003$, $F_{(2,45)}=9.75$; adult WT
control=82.71±10.44 s, stress=37.18±3.63 s, N=17, p=0.0007; young control=108.54±12.04 s,
stress=103.46±14.42 s, N=13, p=0.98; SynCAM 1 KO control=75.72±5.55 s, stress=60.06±9.36 s,
N=18, p=0.42; Sidak's post-hoc). The effect of age/plasticity level was present in both measures,
mostly reflecting the differences in adult WT mice compared to other groups (fraction of time
immobilized adult vs young and adult vs SynCAM 1 KO, p<0.0001; young vs SynCAM 1 KO,

p=0.65; latency to first immobilization adult vs young, P<0.0001; adult vs SynCAM 1 KO, p=0.21;
young vs SynCAM 1 KO, p=0.007).

To test if our stress paradigm increases anxiety-like behaviors, we compared the preference of
mice to stay close to the walls of the open field arena (thigmotaxis, Figure 2C) as well as their
locomotor activity (Figure 2D). Interestingly, we found that short unpredictable stress did not impact
thigmotaxis (Figure 2C; 2-way repeated measures (RM) ANOVA, stress x age/plasticity p=0.65,
$F_{(2,62)}$=0.44; p_{stress}=0.82, $F_{(1,62)}$=0.05; $p_{age/plasticity}$=0.07, $F_{(2,62)}$=2.79; adult WT control=0.83±0.01,
stress=0.82±0.01, N=29; young control=0.76±0.01, stress=0.78±0.02, N=9; SynCAM 1 KO
control=0.81±0.02, stress=0.81±0.02, N=21). However, as previously reported (Sequeira☐Cordero et
al., 2019), stressed mice from all groups travelled more in the open field arena, with SynCAM 1 KO
mice traversing the longest distance (Figure 2D; 2-way repeated measures (RM) ANOVA, stress x
age/plasticity p=0.72, $F_{(2,62)}$=0.32; p_{stress}=0.02, $F_{(1,62)}$=5.42; $p_{age/plasticity}$=0.0014, $F_{(2,62)}$=7.35; adult WT
control=24.26±0.98 m, stress=26.39±1.53 m, N=29, p=0.51; young control=29.78±1.95 m,
stress=32.66±1.88 m, N=9, p=0.72; SynCAM 1 KO control=30.05±2.18 m, stress=34.14±2.21 m,
N=21, p=0.07). SynCAM 1 KO mice crossed more distance when compared to adult mice in both
control and stress groups (adult vs SynCAM 1 KO control p=0.03, stress=0.002), but not compared
to young mice (young vs SynCAM 1 KO control p=0.99, stress p=0.96), confirming previous reports
of increased locomotor activity in these mice (Giza et al., 2013). Locomotor activity in all mouse
groups did not correlate with the fraction of time spent immobilized in TST, suggesting that
increased overall locomotor activity in young and SynCAM 1 KO mice is not the cause of their
resilience to stress in TST assay (Pearson r adult=-0.07, p=0.76; young=0.04, p=0.89; SynCAM 1
KO=-0.02, p=0.91). Altogether, our results demonstrated that short unpredictable stress robustly
increases depressive-like behaviors in adult mice, but not in young mice or adult mice with critical
period-like cortical plasticity.

Discussion

Since Hubel and Wiesel reported a temporally restricted window of plasticity in the kitten
cortex during which neurons robustly respond to changes in sensory input (Hubel and Wiesel, 1970),
many studies have attempted to restore that type of plasticity to the adult brain aiming to mitigate the
damage caused by a myriad of stressors (Ribic, 2020; Hensch and Quinlan, 2018; Levelt and
Hübener, 2012). In support of this longstanding notion, antidepressants can restore juvenile-like
plasticity to the adult cortex (Maya Vetencourt et al., 2008), but it is still unclear if they promote
recovery from stress through changing the quality of plasticity in the adult brain. Our study attempted
to shed more light on this issue by testing if short unpredictable stress results in depressive-like
behavior in adolescent mice in whom the cortical window of plasticity is still open, as well as in adult
mice in whom the plasticity windows are extended into adulthood through deletion of SynCAM 1
(Ribic et al., 2019). We found that this is indeed the case, supporting that interventions that change
the quality of plasticity in the adult brain can facilitate stress resilience (Nestler and Russo, 2024;
McEwen, 2016; Fuchs et al., 2004).

The duration and type of stressors, along with age and sex of animals, can have a significant
impact on physiological measures of stress, as evident in our results as well (Luo et al., 2025;
Monteiro et al., 2015). Our stress paradigm did not result in abrogated weight gain in any of the
mouse groups, likely because of the short duration of our protocol, as well as the stressors
themselves. Adrenal gland weight was, however, increased in young female mice and all 3 groups of
mice displayed robust stress-induced hyperthermia, indicating that our protocol was stressful to them.
In agreement, adult mice displayed an increase in depressive-like behaviors as measured with tail

suspension test (TST). Interestingly, adolescent mice and adult SynCAM 1 KO mice did not show
any changes in immobilization after stress, in support of the notion that juvenile/adolescent-like
plasticity promotes resilience to stress. While it is possible that increased locomotor activity of young
and adult SynCAM 1 KO mice results in reduced immobilization during TST, we find it unlikely as
longer stress duration in adolescent mice results in increased immobilization in female mice (Hu et
al., 2025). Further, we found no correlation between immobility and locomotor activity in the open
field arena, indicating that young mice and adult SynCAM 1 mice indeed are resilient to short
unpredictable stress. Based on our and other studies (Hu et al., 2025), Future studies can now address
the impact of longer stress duration or more severe stressors on the behavior of adolescent and adult
SynCAM 1 KO mice, and whether juvenile/adult-like plasticity provides a lasting protection from
stress.

Unlike immobilization, locomotor activity was mildly impacted by stress, indicated by
increased distance travelled in the open field arena in all three groups of mice. While within-group
post-hoc comparisons were not statistically significant, our results are in line with previous research
that demonstrated stress-associated hyperlocomotion (Sequeira□Cordero et al., 2019). Distance
travelled positively correlates with corticosterone levels in mice head-fixed above a treadmill, which
itself is stressful (Juczewski et al., 2020), suggesting that increased locomotion is a stress response.
Locomotor activity is traditionally classified as anxiety-like behavior, indicating that
juvenile/adolescent-like plasticity may not promote resilience to anxiety after stress. However, future
studies need to address this issue using paradigms that can assess anxiety-like phenotype independent
of locomotion, like novelty suppressed feeding (NSF) (Koskinen and Hovatta, 2023).

While the type of plasticity in our study correlates with resilience to short unpredictable
stress, it is possible that the resilience of SynCAM 1 KO mice is due to another mechanism.
SynCAM 1 gene expression is not restricted to the brain (Fujita et al., 2006) and, while SynCAM 1
protein is localized to the membrane, it can trigger multiple signaling cascades that promote neuronal
resilience to stress (Cheadle and Biederer, 2012; Murakami et al., 2014). However, we find this
unlikely, as SynCAM 1 KO mice display overall accelerated learning in spatial tasks and juvenile-
like responses to noxious stimuli (Park et al., 2016; Robbins et al., 2010). SynCAM 1 is a potent
plasticity brake in the adult brain and even a transient knock-down of its expression can restore
juvenile/adolescent like plasticity in the adult primary visual cortex (Ribic et al., 2019). Our study is
hence in line with the notion that critical period-like plasticity is protective of stress and provides a
rationale for further inquiries into the link between resilience and different types of plasticity.

Our study is limited in its scope, but our findings unambiguously demonstrate the efficacy of
short unpredictable stress in inducing depressive-like behaviors in adult mice. The short duration and
ease of administering our protocol will undoubtedly facilitate further studies of short unpredictable
stress and its long-term effects on brain and behavior. Importantly, our study provides novel evidence
that supports the notion that heightened brain plasticity promotes resilience to stress, and provides a
rationale for future, more in-depth mechanistic inquiries into the relationship between plasticity and
stress.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Author Contributions

AR and RW conceived and designed the study. RW, CVDO, ENL, SCF, GLO and AR collected and
analyzed the data. AR wrote the manuscript with input from all authors.

Funding

This study was funded through the Department of Psychology at the University of Virginia.

Acknowledgments

The authors would like to thank the Ribic Lab for critical reading of the manuscript.

Data Availability Statement

The datasets generated and analyzed in this study will be provided upon reasonable request.

References

Banerjee, S. B., Gutzeit, V. A., Baman, J., Aoued, H. S., Doshi, N. K., Liu, R. C., and Ressler, K. J.
(2017). Perineuronal nets in the adult sensory cortex are necessary for fear learning. *Neuron* 95,
169-179.e3. doi:10.1016/j.neuron.2017.06.007.

Bhagat, S. M., Butler, S. S., Taylor, J. R., McEwen, B. S., and Strittmatter, S. M. (2016). Erasure of
fear memories is prevented by Nogo Receptor 1 in adulthood. *Mol. Psychiatry* 21, 1281–1289.
doi:10.1038/mp.2015.179.

Bielohuby, M., Herbach, N., Wanke, R., Maser-Gluth, C., Beuschlein, F., Wolf, E., and Hoeflich, A.
(2007). Growth analysis of the mouse adrenal gland from weaning to adulthood: time- and gender-
dependent alterations of cell size and number in the cortical compartment. *Am. J. Physiol.*
Endocrinol. Metab. 293, E139-46. doi:10.1152/ajpendo.00705.2006.

Bloss, E. B., Janssen, W. G., McEwen, B. S., and Morrison, J. H. (2010). Interactive effects of stress
and aging on structural plasticity in the prefrontal cortex. *J. Neurosci.* 30, 6726–6731.
doi:10.1523/JNEUROSCI.0759-10.2010.

Brydges, N. M., Jin, R., Seckl, J., Holmes, M. C., Drake, A. J., and Hall, J. (2014). Juvenile stress
enhances anxiety and alters corticosteroid receptor expression in adulthood. *Brain Behav.* 4, 4–13.
doi:10.1002/brb3.182.

Can, A., Dao, D. T., Terrillion, C. E., Piantadosi, S. C., Bhat, S., and Gould, T. D. (2012). The tail
suspension test. *J. Vis. Exp.*, e3769. doi:10.3791/3769.

Cheadle, L., and Biederer, T. (2012). The novel synaptogenic protein Farp1 links postsynaptic
cytoskeletal dynamics and transsynaptic organization. *J. Cell Biol.* 199, 985–1001.
doi:10.1083/jcb.201205041.

Cryan, J. F., Mombereau, C., and Vassout, A. (2005). The tail suspension test as a model for
assessing antidepressant activity: review of pharmacological and genetic studies in mice. *Neurosci.*
Biobehav. Rev. 29, 571–625. doi:10.1016/j.neubiorev.2005.03.009.

Czéh, B., Fuchs, E., Wiborg, O., and Simon, M. (2016). Animal models of major depression and their
clinical implications. *Prog Neuropsychopharmacol Biol Psychiatry* 64, 293–310.
doi:10.1016/j.pnpbp.2015.04.004.

Faye, C., Mcgowan, J. C., Denny, C. A., and David, D. J. (2018). Neurobiological mechanisms of
stress resilience and implications for the aged population. *Curr. Neuropharmacol.* 16, 234–270.
doi:10.2174/1570159X15666170818095105.

Fuchs, E., Czéh, B., Kole, M. H. P., Michaelis, T., and Lucassen, P. J. (2004). Alterations of
neuroplasticity in depression: the hippocampus and beyond. *Eur. Neuropsychopharmacol.* 14
Suppl 5, S481-90. doi:10.1016/j.euroneuro.2004.09.002.

Fujita, E., Kouroku, Y., Ozeki, S., Tanabe, Y., Toyama, Y., Maekawa, M., Kojima, N., Senoo, H.,
Toshimori, K., and Momoi, T. (2006). Oligo-astheno-teratozoospermia in mice lacking
RA175/TSLC1/SynCAM/IGSF4A, a cell adhesion molecule in the immunoglobulin superfamily.
Mol. Cell. Biol. 26, 718–726. doi:10.1128/MCB.26.2.718-726.2006.

Giza, J. I., Jung, Y., Jeffrey, R. A., Neugebauer, N. M., Picciotto, M. R., and Biederer, T. (2013). The
synaptic adhesion molecule SynCAM 1 contributes to cocaine effects on synapse structure and
psychostimulant behavior. *Neuropsychopharmacology* 38, 628–638. doi:10.1038/npp.2012.226.

Gogolla, N., Caroni, P., Lüthi, A., and Herry, C. (2009). Perineuronal nets protect fear memories
from erasure. *Science* 325, 1258–1261. doi:10.1126/science.1174146.

Golden, S. A., Covington, H. E., Berton, O., and Russo, S. J. (2011). A standardized protocol for
repeated social defeat stress in mice. *Nat. Protoc.* 6, 1183–1191. doi:10.1038/nprot.2011.361.

Gordon, J. A., and Stryker, M. P. (1996). Experience-dependent plasticity of binocular responses in
the primary visual cortex of the mouse. *J. Neurosci.* 16, 3274–3286. doi:10.1523/JNEUROSCI.16-
10-03274.1996.

Greifzu, F., Kalogeraki, E., and Löwel, S. (2016). Environmental enrichment preserved lifelong
ocular dominance plasticity, but did not improve visual abilities. *Neurobiol. Aging* 41, 130–137.
doi:10.1016/j.neurobiolaging.2016.02.014.

Greifzu, F., Pielecka-Fortuna, J., Kalogeraki, E., Krempler, K., Favaro, P. D., Schlüter, O. M., and
Löwel, S. (2014). Environmental enrichment extends ocular dominance plasticity into adulthood
and protects from stroke-induced impairments of plasticity. *Proc Natl Acad Sci USA* 111, 1150–
1155. doi:10.1073/pnas.1313385111.

Hegde, A., Suresh, S., and Mitra, R. (2020). Early-life short-term environmental enrichment
counteracts the effects of stress on anxiety-like behavior, brain-derived neurotrophic factor and
nuclear translocation of glucocorticoid receptors in the basolateral amygdala. *Sci. Rep.* 10, 14053.
doi:10.1038/s41598-020-70875-5.

Hensch, T. K., and Quinlan, E. M. (2018). Critical periods in amblyopia. *Vis. Neurosci.* 35, E014.
doi:10.1017/S0952523817000219.

Hubel, D. H., and Wiesel, T. N. (1970). The period of susceptibility to the physiological effects of
unilateral eye closure in kittens. *J Physiol (Lond)* 206, 419–436.
doi:10.1113/jphysiol.1970.sp009022.

Hu, W., Jiang, L., Wang, Q., Hu, Q., Zhong, T., Wu, J., Chen, X., and Liu, T. (2025). Chronic
unpredictable stress during adolescence exerts sex-specific effects on depressive-like behavior and
neural activation triggered by tail suspension test. *Behav. Brain Res.* 477, 115314.
doi:10.1016/j.bbr.2024.115314.

Juczewski, K., Koussa, J. A., Kesner, A. J., Lee, J. O., and Lovinger, D. M. (2020). Stress and
behavioral correlates in the head-fixed method: stress measurements, habituation dynamics,
locomotion, and motor-skill learning in mice. *Sci. Rep.* 10, 12245. doi:10.1038/s41598-020-69132-
6.

Koskinen, M.-K., and Hovatta, I. (2023). Genetic insights into the neurobiology of anxiety. *Trends*
Neurosci. 46, 318–331. doi:10.1016/j.tins.2023.01.007.

Lehmann, M. L., and Herkenham, M. (2011). Environmental enrichment confers stress resiliency to
social defeat through an infralimbic cortex-dependent neuroanatomical pathway. *J. Neurosci.* 31,
6159–6173. doi:10.1523/JNEUROSCI.0577-11.2011.

Levelt, C. N., and Hübener, M. (2012). Critical-period plasticity in the visual cortex. *Annu. Rev.*
Neurosci. 35, 309–330. doi:10.1146/annurev-neuro-061010-113813.

Luo, Y., Zhang, Y., Zhao, Y., Chen, F., Li, X., Liu, X., Shakir, M. Z., Jiang, N., and Shan, C. (2025).
Chronic Unpredictable Mild Stress-Induced Anxiety is Linked to Inflammatory Responses and
Disruptions in Tryptophan Metabolism in Male C57BL/6 N Mice. *Behav. Brain Res.*, 115506.
doi:10.1016/j.bbr.2025.115506.

Macartney, E. L., Lagisz, M., and Nakagawa, S. (2022). The relative benefits of environmental
enrichment on learning and memory are greater when stressed: A meta-analysis of interactions in
rodents. *Neurosci. Biobehav. Rev.* 135, 104554. doi:10.1016/j.neubiorev.2022.104554.

Maya Vetencourt, J. F., Sale, A., Viegi, A., Baroncelli, L., De Pasquale, R., O'Leary, O. F., Castrén,
E., and Maffei, L. (2008). The antidepressant fluoxetine restores plasticity in the adult visual
cortex. *Science* 320, 385–388. doi:10.1126/science.1150516.

McEwen, B. S., Bowles, N. P., Gray, J. D., Hill, M. N., Hunter, R. G., Karatsoreos, I. N., and Nasca,
C. (2015). Mechanisms of stress in the brain. *Nat. Neurosci.* 18, 1353–1363. doi:10.1038/nn.4086.

McEwen, B. S., and Morrison, J. H. (2013). The brain on stress: vulnerability and plasticity of the
prefrontal cortex over the life course. *Neuron* 79, 16–29. doi:10.1016/j.neuron.2013.06.028.

McEwen, B. S. (2012). Brain on stress: how the social environment gets under the skin. *Proc Natl*
Acad Sci USA 109 Suppl 2, 17180–17185. doi:10.1073/pnas.1121254109.

McEwen, B. S. (2016). In pursuit of resilience: stress, epigenetics, and brain plasticity. *Ann. N. Y.*
Acad. Sci. 1373, 56–64. doi:10.1111/nyas.13020.

Miwa, J. M., Stevens, T. R., King, S. L., Caldarone, B. J., Ibanez-Tallon, I., Xiao, C., Fitzsimonds, R.
M., Pavlides, C., Lester, H. A., Picciotto, M. R., et al. (2006). The prototoxin lynx1 acts on
nicotinic acetylcholine receptors to balance neuronal activity and survival in vivo. *Neuron* 51,
587–600. doi:10.1016/j.neuron.2006.07.025.

Monteiro, S., Roque, S., de Sá-Calçada, D., Sousa, N., Correia-Neves, M., and Cerqueira, J. J.
(2015). An efficient chronic unpredictable stress protocol to induce stress-related responses in
C57BL/6 mice. *Front. Psychiatry* 6, 6. doi:10.3389/fpsyt.2015.00006.

Murakami, S., Sakurai-Yageta, M., Maruyama, T., and Murakami, Y. (2014). Trans-homophilic
interaction of CADM1 activates PI3K by forming a complex with MAGuK-family proteins MPP3
and Dlg. *PLoS ONE* 9, e110062. doi:10.1371/journal.pone.0110062.

Nestler, E. J., and Russo, S. J. (2024). Neurobiological basis of stress resilience. *Neuron* 112, 1911–
1929. doi:10.1016/j.neuron.2024.05.001.

Park, K. A., Ribic, A., Laage Gaupp, F. M., Coman, D., Huang, Y., Dulla, C. G., Hyder, F., and
Biederer, T. (2016). Excitatory synaptic drive and feedforward inhibition in the hippocampal CA3
circuit are regulated by syncam 1. *J. Neurosci.* 36, 7464–7475. doi:10.1523/JNEUROSCI.0189-
16.2016.

Patterson, Z. R., and Abizaid, A. (2013). Stress induced obesity: lessons from rodent models of
stress. *Front. Neurosci.* 7, 130. doi:10.3389/fnins.2013.00130.

Peleg-Raibstein, D., and Feldon, J. (2011). Differential effects of post-weaning juvenile stress on the
behaviour of C57BL/6 mice in adolescence and adulthood. *Psychopharmacology (Berl)* 214, 339–
351. doi:10.1007/s00213-010-1991-8.

Phoumthipphavong, V., Barthas, F., Hassett, S., and Kwan, A. C. (2016). Longitudinal effects of
ketamine on dendritic architecture in vivo in the mouse medial frontal cortex. *eNeuro* 3.
doi:10.1523/ENEURO.0133-15.2016.

Ribic, A., Crair, M. C., and Biederer, T. (2019). Synapse-Selective Control of Cortical Maturation
and Plasticity by Parvalbumin-Autonomous Action of SynCAM 1. *Cell Rep.* 26, 381-393.e6.
doi:10.1016/j.celrep.2018.12.069.

Ribic, A. (2020). Stability in the Face of Change: Lifelong Experience-Dependent Plasticity in the
Sensory Cortex. *Front. Cell. Neurosci.* 14, 76. doi:10.3389/fncel.2020.00076.

Robbins, E. M., Krupp, A. J., Perez de Arce, K., Ghosh, A. K., Fogel, A. I., Boucard, A., Südhof, T.
C., Stein, V., and Biederer, T. (2010). SynCAM 1 adhesion dynamically regulates synapse number
and impacts plasticity and learning. *Neuron* 68, 894–906. doi:10.1016/j.neuron.2010.11.003.

Russo, S. J., Murrough, J. W., Han, M.-H., Charney, D. S., and Nestler, E. J. (2012). Neurobiology of
resilience. *Nat. Neurosci.* 15, 1475–1484. doi:10.1038/nn.3234.

Seibenhener, M. L., and Wooten, M. C. (2015). Use of the Open Field Maze to measure locomotor
and anxiety-like behavior in mice. *J. Vis. Exp.*, e52434. doi:10.3791/52434.

Sequeira-Cordero, A., Salas-Bastos, A., Fornaguera, J., and Brenes, J. C. (2019). Behavioural
characterisation of chronic unpredictable stress based on ethologically relevant paradigms in rats.
Sci. Rep. 9, 17403. doi:10.1038/s41598-019-53624-1.

Vinkers, C. H., van Bogaert, M. J. V., Klanker, M., Korte, S. M., Oosting, R., Hanania, T., Hopkins,
S. C., Olivier, B., and Groenink, L. (2008). Translational aspects of pharmacological research into
anxiety disorders: the stress-induced hyperthermia (SIH) paradigm. *Eur. J. Pharmacol.* 585, 407–
425. doi:10.1016/j.ejphar.2008.02.097.

Willner, P., Towell, A., Sampson, D., Sophokleous, S., and Muscat, R. (1987). Reduction of sucrose
preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant.
Psychopharmacology (Berl) 93, 358–364. doi:10.1007/BF00187257.

Yang, W.-Z., Liu, T.-T., Cao, J.-W., Chen, X.-F., Liu, X., Wang, M., Su, X., Zhang, S.-Q., Qiu, B.-
L., Hu, W.-X., et al. (2016). Fear erasure facilitated by immature inhibitory neuron transplantation.
Neuron 92, 1352–1367. doi:10.1016/j.neuron.2016.11.018.

Zethof, T. J., Van der Heyden, J. A., Tolboom, J. T., and Olivier, B. (1994). Stress-induced
hyperthermia in mice: a methodological study. *Physiol. Behav.* 55, 109–115. doi:10.1016/0031-
9384(94)90017-5.

Zhang, C., Li, H., and Han, R. (2020). An open-source video tracking system for mouse locomotor
activity analysis. *BMC Res. Notes* 13, 48. doi:10.1186/s13104-020-4916-6.

   Figures and Figure Legends

Figure 1. Short unpredictable stress induces robust hyperthermia. A) Experimental
schematics. Mice of both sexes (adult, postnatal day/P 60 and older; young, P21-P35; adult SynCAM

1 KO) underwent one week of short unpredictable stress with indicated stressors. Both within-subject
(top) and between-subject design (bottom) was used to accommodate for isolation of adrenal glands.
B) Adrenal gland weight was not significantly increased after stress in any of the groups (2-way
ANOVA, p=0.098, $F_{(2, 61)}$=2.42; p_{stress}=0.67, $F_{(1, 61)}$=0.2; $p_{age/plasticity}$=0.078, $F_{(2, 61)}$=2.66). **C)** After
separating the mice by sex, young female mice showed an increase in adrenal gland weight after
stress (2-way ANOVA, stress x sex p=0.51, $F_{(1,15)}$=0.47; p_{stress}=0.003, $F_{(1,15)}$=12.6; p_{sex}<0.0001,
$F_{(1,15)}$=58.75; Sidak multiple comparison test males p=0.19, females p=0.004). **D)** Short
unpredictable stress did not abrogate weight gain in any of the groups (2-way ANOVA stress x
age/plasticity p=0.56, $F_{(2,99)}$=0.58; p_{stress}=0.12, $F_{(1,99)}$=2.45; $p_{age/plasticity}$<0.0001, $F_{(2,99)}$=15.58). **E)** Short
unpredictable stress resulted in robust elevation of core body temperature in all groups of mice (2-
way ANOVA stress x age/plasticity p=0.95, $F_{(2,100)}$=0.05; p_{stress}<0.0001, $F_{(2,100)}$=42.75;
$p_{age/plasticity}$=0.02, $F_{(2,100)}$=4.05). Medians and quartiles of the data are indicated by lines and individual
points represent mice.

Figure 2. Young mice and adult SynCAM 1 KO mice are resilient to stress-induced
increase learned helplessness-like behavior. A) Only stressed adult WT mice show increased
immobility after stress in tail suspension test (2-way repeated measures (RM) ANOVA, stress x
age/plasticity p<0.0001, $F_{(2,45)}$=25.24; p_{stress}=0.04, $F_{(1,45)}$=4.11; $p_{age/plasticity}$<0.0001, $F_{(2,45)}$=18.3; adult

WT, p<0.0001; young WT, p=0.42; SynCAM 1 KO, p=0.18; Sidak's post-hoc) and a significant reduction in time lapsed to first immobilization **(B)**(2-way repeated measures (RM) ANOVA, stress x age/plasticity p=0.054, $F_{(2,45)}$=3.12; p_{stress}=0.002, $F_{(1,45)}$=10.41; $p_{age/plasticity}$=0.0003, $F_{(2,45)}$=9.75; adult WT, p=0.0007; young WT, p=0.98; SynCAM 1 KO, p=0.42; Sidak's post-hoc). **C)** Short unpredictable stress did not impact thigmotaxis in open field test (2-way repeated measures (RM) ANOVA, stress x age/plasticity p=0.65, $F_{(2,62)}$=0.44; p_{stress}=0.82, $F_{(1,62)}$=0.05; $p_{age/plasticity}$=0.07, $F_{(2,62)}$=2.79). **D)** While stressed mice from all groups travelled more in the open field arena, within-group post-hoc comparisons were not statistically significant (2-way repeated measures (RM) ANOVA, stress x age/plasticity p=0.72, $F_{(2,62)}$=0.32; p_{stress}=0.02, $F_{(1,62)}$=5.42; $p_{age/plasticity}$=0.0014, $F_{(2,62)}$=7.35; adult WT, p=0.51; young WT, p=0.72; SynCAM 1 KO, p=0.07).