An organotrophic *Sideroxydans* reveals potential iron oxidation marker genes

Rene L. Hoover[1,2], Kirsten Küsel[3,4], Clara S. Chan[1,2,5]

1 Microbiology Graduate Program, University of Delaware, Newark, Delaware, USA
2 Department of Earth Sciences, University of Delaware, Newark, Delaware, USA
3 Aquatic Geomicrobiology, Institute of Biodiversity, Friedrich Schiller University, Jena, Germany
4 Cluster of Excellence Balance of the Microverse, Friedrich Schiller University, Jena, Germany
5 Delaware Biotechnology Institute, Newark, Delaware, USA

Abstract

To understand the ecophysiology and the role of iron-oxidizing bacteria (FeOB) in various ecosystems, we need to identify marker genes of the iron oxidation pathway to track activity *in situ*. The Gallionellaceae *Sideroxydans* sp. CL21, an autotrophic iron-oxidizing bacteria isolated from a peatland, is unusual amongst FeOB isolates in its genomic potential to utilize organic compounds as energy sources. Therefore, it offers the unique opportunity to determine genes expressed under litho- versus organotrophic conditions. We demonstrated the growth of *Sideroxydans* sp. CL21 on organic substrates (lactate and pyruvate) and inorganic substrates (Fe(II), magnetite, thiosulfate, and S(0)). Thus, cells were capable of lithoautotrophic, organotrophic, and potentially organoheterotrophic growth. Surprisingly, when lactate-grown cells were given Fe(II), mid-log phase cells were unable to oxidize iron, while late-log phase cells oxidized iron. To elucidate iron oxidation pathways, we compared gene expression between mid-log (non-iron-oxidizing) and late-log (iron-oxidizing) lactate-grown cells. Genes for iron oxidases (*cyc2, mtoA*) were highly expressed at both time points, so did not correspond to iron oxidation capability, making them unsuitable marker genes of iron oxidation activity by themselves. However, genes encoding periplasmic and inner membrane cytochromes were significantly upregulated in cells capable of iron oxidation. These genes include *mtoD*, *cymA/imoA*, and a cluster of Fe(II)-responsive genes (*ircABCD*). These findings suggest Gallionellaceae regulate their iron oxidation pathways in multiple stages, with iron oxidase-encoding genes proactively expressed. Other genes encoding electron carriers are upregulated only when iron oxidation is needed, which makes these genes (i.e. *ircABCD*) good prospective indicators of iron oxidation ability.

Importance

FeOB are widespread in the environment and we suspect that they play key roles in nutrient and other elemental cycles. However, with no isotopic marker, we lack the ability to

monitor FeOB activity, prompting us to search for genetic markers. Previous work suggests that
expression of iron oxidase genes does not directly correspond to iron oxidation activity in
Gallionellaceae and little was known about the other genes in the pathway. Here we study a
unique FeOB isolate that possesses organotrophic capabilities and demonstrate its potential for
mixotrophic growth on lactate and Fe(II). Its ability to oxidize iron is regulated, allowing us to
discover potential iron oxidation pathway genes with expression that corresponds to iron
oxidation activity. If these genes can be further validated as iron oxidation marker genes, they
will enable us to delineate autotrophic and organoheterotrophic FeOB impacts on carbon cycling
in wetlands and other natural and engineered environments.

Introduction

Iron oxidation is a widespread process in the environment, playing key roles in water and
soil health as Fe(III) oxyhydroxides can adsorb various nutrients and metals (1). Despite the
pervasiveness and importance of iron oxidation, there remains much uncertainty as to how much
of it is abiotic versus biotic, i.e. catalyzed by iron-oxidizing bacteria (FeOB). This distinction
(biotic vs. abiotic) is key to understanding how iron oxidation affect other nutrient cycles, which
requires 1) a deeper understanding of C metabolisms of FeOB and 2) a method to link microbial
iron oxidation activity in the environment to C and other metabolic transformations. Although
FeOB are widely thought to be primarily autotrophic, accumulating evidence suggests a
considerable number of FeOB may be mixotrophic (2–6) or organo(hetero)trophic (7, 8),
expanding their potential effects on C cycling. Because these FeOB can use alternate energy
sources (9–12), they are not always oxidizing iron, so we require a method of detecting active
iron oxidation. To link microbial iron oxidation to biogeochemical effects, we need gene markers
of iron oxidation activity.
Studying facultative FeOB allows us to discern genes specific to iron oxidation. An
isolate of the Gallionellaceae, *Sideroxydans* sp. CL21, presents a new opportunity to learn about
FeOB metabolism, iron oxidation mechanisms, and potential marker genes. *Sideroxydans* sp.
CL21 is rare amongst FeOB isolates in possessing genes for organoheterotrophy, notably lactate
permease and dehydrogenase. It was isolated from the Schlöppnerbrunnen fen (13), a semi-acidic
(pH 4.5 to 5.5), iron-rich peatland in Northern Germany (14–16). *Sideroxydans* sp. CL21 is
closely related to the well-studied *Sideroxydans lithotrophicus* ES-1 (9), but differs from ES-1 in
that its genome is ~20% larger and encodes genes for organoheterotrophy (17, 18). *Sideroxydans*
sp. CL21 has been shown to grow on the non-iron substrates H_2 and thiosulfate (18). It has also
been grown on the reduced iron substrates FeS and Fe(0) (13, 18), but neither of these are pure
Fe(II) substrates as Fe(0) evolves H_2, so growth on Fe(II) alone has not been demonstrated.
Sideroxydans sp. CL21 has also been grown in the presence of mixed substrates, including
lactate/FeS and lactate/Fe(0) (13, 18). However, previous cultures did not grow on lactate alone
(13, 18). Thus, while the growth in the presence of combined organic and inorganic electron

donors suggests a potential for mixotrophy, further work is needed to clarify exactly which

substrates promote *Sideroxydans* sp. CL21 growth and support organoheterotrophy.

Markers of iron oxidation activity could be the genes within iron oxidation pathways, but

our knowledge of these pathways is still in development. The *Sideroxydans* sp. CL21 genome

encodes several genes related to iron oxidation and associated pathways. Its genome encodes two

known iron oxidases, 1) the monoheme cytochrome-porin Cyc2 and 2) the decaheme

cytochrome MtoA and associated porin MtoB (17–19). In fact, the genome has three copies of

cyc2 and an *mtoAB,* plus an additional copy of *mtoA* next to an unnamed porin. While the

presence of *cyc2* or *mtoA* indicate iron oxidation capability, it is unclear if they are suitable

genetic markers of iron oxidation *activity*. Ideally, a gene marker would exhibit little to no

expression when there is no iron oxidation activity and be upregulated during iron oxidation,

with expression level scaling with activity. Yet, in the closely related *Sideroxydans*

lithotrophicus ES-1, one copy of *cyc2* is highly expressed in both iron-oxidizing and thiosulfate-

oxidizing cells, while the other two copies have low expression during thiosulfate oxidation, but

are upregulated when oxidizing iron (10). In marine hydrothermal environments, the *cyc2* gene is

highly expressed by various iron-oxidizing Zetaproteobacteria, and expression increases in

microcosms amended with Fe(II) (20). In the facultative FeOB Zetaproteobacteria *Ghiorsea*

bivora TAG-1, $cyc2_{TAG\text{-}1\_1}$ and $Cyc2_{TAG\text{-}1\_1}$ are expressed highly under both H_2- and iron-

oxidizing conditions (21). Together, this suggests *cyc2*/Cyc2 expression may not necessarily

correspond to iron oxidation activity. There is less information on *mtoAB/*MtoAB, though these

iron oxidase genes/proteins are upregulated in *S. lithotrophicus* ES-1 grown on Fe(II) minerals

(22, 23). *Sideroxydans* sp. CL21 also encodes two gene clusters, predicted to encode porin-

cytochrome complexes (PCC), each with two multiheme cytochromes ($PCC3_{copy1}$, cytochromes

have 24 and 14 hemes; $PCC3_{copy2}$, cytochromes have 24 and 12 hemes). PCC3 is hypothesized to

also play a role in extracellular electron transport and possibly iron oxidation (24). Additional

putative components of the iron oxidation pathway (*mtoD*, *cymA/imoA*, Slit $1321\text{-}1324_{ES\text{-}1}$) are

less well studied; these include genes for inner membrane and periplasmic electron carriers. In

all, *Sideroxydans* sp. CL21 has a wide variety of genes that are known or predicted to participate

in iron oxidation, giving multiple possibilities for marker genes.

Here, we constrain potential iron oxidation marker genes by studying differential gene

expression in *Sideroxydans* sp. CL21. We first optimized the cultivation conditions by testing

Sideroxydans sp. CL21's growth on organic acids alone, plus other Fe(II) and non-iron

substrates. Cells grew quickly and to the highest cell density on organic acids. We then tested the

ability of these organic-grown cells to oxidize iron and found that lactate-grown cells exhibited a

difference in iron-oxidizing phenotype between mid-log (no iron oxidation activity) and late-log

(iron oxidation activity). This enabled us to compare gene expression between iron-oxidizing and

non-iron-oxidizing conditions to detect genes whose expression corresponds to iron oxidation

potential. The genes identified help expand our knowledge of the iron oxidation pathway and

provide a list of candidate marker genes for iron oxidation.

Results

Physiology and growth on Fe(II), S, and organic substrates

Sideroxydans sp. CL21 grew on a range of individual organic and inorganic electron donors (Fig. 1). *Sideroxydans* sp. CL21 grew well on dissolved, ferrous iron (FeCl$_2$), reaching a cell density of 1.25 x 10^7 cells/mL in 15 days (Fig. 1A). This growth on FeCl$_2$ was improved when media included 2.5 mM sodium citrate to complex iron, reaching a density of 2.76 x 10^7 cells/mL (Fig. 1A). Attempts to culture *Sideroxydans* sp. CL21 on 5 mM sodium citrate alone showed no growth beyond that of a no-substrate control (Fig. 1A), demonstrating *Sideroxydans* sp. CL21 cannot conserve energy from citrate. As Fe(II) was the only available electron donor, these results confirm that *Sideroxydans* sp. CL21 can grow as a lithoautotrophic iron oxidizer.

Sideroxydans sp. CL21 was cultured on commercial and biogenic magnetite to assess its growth on solid iron minerals. Cells grew better on the biogenic versus commercial magnetite (Fig. 1B). This may be due to the lower crystallinity and higher solubility of biogenic magnetite, which sheds more Fe(II) into solution than commercial magnetite (25). The biogenic magnetite, which was produced from ferrihydrite by the iron-reducer *Shewanella*, may also contain extracellular metabolites that could enhance iron oxidation activity and growth in *Sideroxydans* sp. CL21 (26).

Sideroxydans sp. CL21 was also tested for growth on sulfur substrates. The genome includes *dsrAB* genes that encode a reverse dissimilatory sulfite reductase (rDSR) (19). It has previously been cultured on FeS and in agarose-stabilized media in gradient tubes with either thiosulfate or thiosulfate plus H$_2$ or lactate (18). Since agarose may provide a source of C (27, 28), we used liquid media without an organic carbon source to confirm whether cells could grow using individual sulfur substates as their sole electron donor. In these experiments *Sideroxydans* sp. CL21 grew on thiosulfate and elemental sulfur (Fig. 1B), reaching densities near 10^7 cells/mL. However, growth was slower (18-19 days) and cell density was lower than that of Fe(II)-grown cells (Fig. 1A, 1B). Nonetheless, this shows *Sideroxydans* sp. CL21 is capable of two additional lithoautotrophic metabolisms using thiosulfate and S(0) as electron donors.

Previous studies showed growth of *Sideroxydans* sp. CL21 in multi-substrate experiments where lactate was combined with inorganic substrates (FeS, Fe(0), H$_2$, NaS$_2$O$_3$) in agarose gradient tubes (18), yet cells were never successfully grown on lactate alone using these methods. Our data show that *Sideroxydans* sp. CL21 grew in liquid culture with either 5 mM lactate (Fig. 1C, 2A) or 5 mM pyruvate (Fig. 1D, 2B) as the sole electron donor. It is unclear why growth on lactate was positive in our study but previously negative, but it could be due to the more controlled oxygen concentrations in the headspace of these serum bottle liquid cultures versus a gradient tube. Cells cultured on lactate grew to an order of magnitude higher density than those cultured on Fe(II), reaching 4.8 x 10^8 cells/mL in 14 days (Fig. 1C). Together, these results show *Sideroxydans* sp. CL21 can grow organotrophically, and likely organoheterotrophically, on lactate or pyruvate.

The potential for mixotrophic energy metabolism

Sideroxydans sp. CL21 cells have the genomic potential to carry out multiple energy metabolisms simultaneously (mixotrophy), but this requires further experimental proof. Our approach was to determine whether *Sideroxydans* sp. CL21 cells grown on short chain organic acids could also oxidize ferrous iron in a short-term assay. We first harvested lactate-grown cells on days 6, 9, and 14 then added a 400 μM "spike" of $FeCl_2$, and measured Fe^{2+} concentration over time. On Day 6, live cells (7.1×10^6 cells/mL) did not oxidize iron within the 60-minute experiment. On Day 9 (3.3×10^7 cells/mL), the modest amount of iron oxidation was mostly driven by a single bottle, 6A, which had a higher cell density than the other replicates, making it closer to late-log phase (Fig. 3A, Fig. S1). In contrast, with live cells (4.8×10^8 cells/mL) harvested on Day 14, iron was completely oxidized within 15 minutes (Fig. 3A) despite the presence of >1 mM of lactate remaining in the cultures (Fig. 2A). This difference in response between mid- and late-log cultures indicates iron oxidation in *Sideroxydans* sp. CL21 is regulated. We also tested pyruvate-grown cells (Fig. 1D, 2B) and found *Sideroxydans* sp. CL21 was primed to oxidize iron on both Day 6 (4.4×10^7 cells/mL) and Day 14 (1.1×10^8 cells/mL) (Fig. 3B). Together, these results demonstrate that *Sideroxydans* sp. CL21 has the ability to oxidize iron while growing on organic acids, indicating that cells are capable of mixotrophic energy metabolism. Furthermore, because iron oxidation is regulated, we are able to use these samples to investigate gene expression differences between iron-oxidizing and non-iron-oxidizing conditions.

Transcriptomics

Lactate-grown cultures were harvested for transcriptomics after aliquots were taken for the Fe(II)-spike experiments on Days 6, 9, and 14 (Fig. 1C, 2A). We present the transcriptome data as Z-scores, calculated from the normalized, regularized log (rlog) transformed read data. This gives a measure of a gene's expression on a log scale relative to the mean expression (Z-score = 0) of all genes at a given time point.

Expression of lactate and carbon fixation genes

The *Sideroxydans* sp. CL21 genome contains six genes for lactate oxidation. These genes are co-located in the genome and encode a lactate response regulator (LldR), L-lactate dehydrogenase (LldE), L-lactate dehydrogenase (LldG), L-lactate dehydrogenase (LldF), D-lactate dehydrogenase (Dld), and lactate permease (LldP) (Fig. 4A). The genes *lldR*, *dld*, and *lldP* set *Sideroxydans* sp. CL21 apart from other closely related Gallionellaceae isolates *S. lithotrophicus ES-1* and *S. emersonii* MIZ01 that are unable to oxidize lactate (Fig. 4A). As expected, all six of the *Sideroxydans* sp. CL21 lactate oxidation genes were consistently expressed throughout the experiment (Fig. 4B, Table S1). None of the genes were significantly

up- or down-regulated between the timepoints (Fig. 4B, Table S1), which fits with the consistent
availability of lactate in the media (Fig. 2A).
When cultured with solely inorganic substrates, *Sideroxydans* sp. CL21 grows
lithoautotrophically fixing CO_2 via ribulose bisphosphate carboxylase (RuBisCo) and the
Calvin–Benson–Bassham (CBB) cycle (26). High concentrations of organic carbon might be
expected to promote heterotrophy and suppress autotrophy, yet lactate-grown *Sideroxydans* sp.
CL21 still expressed genes for both Form I and Form II RuBisCo at all timepoints (Fig. S1B).
Expression of Form I RuBisCo was relatively low while expression of Form II RuBisCo was
high and greater than mean total expression at all three timepoints (Fig. S1B, Table S1). Form II
RuBisCo has a higher affinity for oxygen (29), so its higher expression relative to Form I would
be consistent with the micro-oxic culturing conditions used in our experiments. While further
metabolic evidence is needed, high expression of the gene encoding Form II RuBisCo suggests
Sideroxydans sp. CL21 could continue to fix CO_2 while oxidizing lactate.

Expression of iron oxidation genes

The different iron oxidation activities of lactate cultures prompted us to investigate
whether this ability corresponds to expression of iron oxidation genes. *Sideroxydans* sp. CL21
has multiple iron oxidase genes, three copies of *cyc2* and two copies of *mtoA*. Though there was
variation in expression between the copies, overall both *cyc2* and *mtoA* were highly expressed on
all three days (Fig. 5A, 5B). Compared to Day 6 (no iron oxidation), expression of *cyc2* or *mtoA*
genes was not statistically significantly up- or down-regulated on Day 14, when iron oxidation
was rapid (Fig. 5A, 5B, Table S1). Thus, expression of known iron oxidase genes did not
correspond to cells' ability to oxidize iron.
Both known iron oxidases, Cyc2 and MtoAB, are predicted to be outer membrane, porin-
cytochromes, allowing them to oxidize Fe(II) outside the cell. Though functionally unverified,
porin-cytochrome c complex 3 (PCC3) is predicted to be an outer membrane porin-cytochrome
complex similar to, but larger than, the MtoAB iron oxidase complexes (24). PCC3 is widely
distributed among various FeOB genomes (19, 24), making PCC3 another potential iron
oxidation mechanism. *Sideroxydans* sp. CL21 has two copies of the PCC3 gene cluster. Each
consists of four genes predicted to encode an extracellular multiheme cytochrome c (MHC), a
porin, a periplasmic MHC, and an inner membrane protein. Analysis of the Z-scores, where
mean expression is 0, showed both copies of PCC3 were expressed higher than the mean (Z-
score ~1; Fig. S1), yet not as highly as *cyc2* or *mtoA* (Z-score ~3; Fig. 5A, 5B, Fig. S1) across all
three time points. From Day 6 to Day 14, PCC3 copy 1 was downregulated while PCC3 copy 2
was upregulated, though the only significant (p-value <0.01) change was to the inner membrane
protein of PCC3 copy 2 (Fig. 6, Fig. S1, Table S1).

Iron oxidases are just one part of FeOB extracellular electron transport pathways, which
require additional cytochromes to convey electrons to the terminal oxidase or quinone pool.
Sideroxydans sp. CL21 has genes encoding two *c*-type cytochromes, the monoheme MtoD and

tetraheme CymA/ImoA, co-located in the genome next to *mtoA* copy 1 (Fig. 5D). Based on
genomic context and previous genetic/biochemical characterization, these proteins may form part
of the Mto iron oxidation pathway, moving electrons from the MtoA iron oxidase to the electron
acceptor and/or quinone pool (30–33). Here, *mtoD* and *cymA/imoA* were upregulated at late log
(Day 14) relative to Day 6, with *mtoD* being significantly upregulated (p-value <0.001) (Fig. 5,
Fig. 6). In this case, the iron oxidation capability corresponds best with expression of the
periplasmic electron carrier gene *mtoD*.
Sideroxydans sp. CL21 also has homologs to the four "Fe(II)-responsive" genes
identified in *S. lithotrophicus* ES-1 (Slit_1321-Slit_1324 (Zhou et al. 2021)). We refer to these
genes as "*ircABCD"* for "**i**ron **r**esponsive **c**luster." The *irc* genes encode two *c*-type
cytochromes, diheme (IrcA) and a monoheme (IrcB), both predicted to be periplasmic. They also
encode a hypothetical protein (IrcC) and a *b*-type cytochrome (IrcD), with predicted localization
in the inner membrane (Fig. 6). These genes were upregulated when *S. lithotrophicus* ES-1 was
grown on Fe(II) versus thiosulfate (10) and also highly expressed in *Sideroxydans* sp. CL21
when grown with Fe(0) (26). In our lactate experiment, expression of these genes increased as
cells approached late-log phase (Day 14) and all were significantly upregulated on Day 14
relative to Day 6 (Fig. 5C, Fig. 6). This significant upregulation corresponds to cells' ability to
oxidize iron on Day 14.
Overall, *Sideroxydans* sp. CL21 seems to be concurrently expressing genes for its known
and putative iron oxidases (*cyc2*, *mtoA*, and PCC3) regardless of whether cells are capable of
iron oxidation. However, expression of genes for periplasmic and inner membrane proteins
(*mtoD*, *ircABCD*, inner membrane component of PCC3 copy 2) varies. These genes were
significantly upregulated in cells that showed iron oxidation activity, making them potential
markers of iron oxidation activity.

Discussion

In this study, we demonstrated that *Gallionellaceae Sideroxydans* sp. CL21 exhibits
remarkable metabolic versatility. *Sideroxydans* sp. CL21 can utilize a wide range of substrates,
including short-chain organic acids such as lactate and pyruvate, as well as oxidize various
soluble inorganic substrates, such as Fe(II) and thiosulfate. Additionally, it can metabolize
mineral substrates, including magnetite and elemental sulfur (S(0)). Overall, our findings show
that *Sideroxydans* sp. CL21 is capable of organotrophy and lithoautotrophy, with the potential
for mixotrophy and organoheterotrophy. This metabolic flexibility likely enhances its
competitiveness and enables it to contribute to a wide range of biogeochemical transformations.
Understanding the ecophysiology and biogeochemical roles of facultative iron oxidizers like
Sideroxydans sp. CL21 requires more than just detecting their presence in environmental
samples. We need a means to decipher whether these FeOB are actively oxidizing iron.
Developing methods to directly track microbial iron oxidation activity requires (1) the

identification of genes encoding the iron oxidation pathway and (2) determining which of these
genes can serve as reliable markers of microbial iron-oxidizing activity.

To be a marker of activity, a gene needs to increase in expression with iron oxidation and
have minimal to no expression when cells are unable to oxidize iron. Validating marker genes
requires study of facultative FeOB, of which there are few in culture, and there ideally needs to
be a condition under which cells *cannot* oxidize iron. Previous studies on *S. lithotrophicus* ES-1
showed differential gene expression of cells grown on Fe(II)-citrate and thiosulfate, but
thiosulfate-grown cells were able to oxidize iron quickly with no lag after the substrate was
switched to Fe(II). In contrast, *Sideroxydans* sp. CL21 grown on lactate was surprisingly unable
to oxidize iron until late-log phase (Day 14), giving an opportunity to compare gene expression
of iron-oxidizing and non-iron-oxidizing conditions.

271 Relationship between iron oxidation gene expression and activity

Although significant progress has been made in assigning function to the key iron
oxidases such as Cyc2 and MtoA (10, 20, 33–35), the pathway is likely more complex, and
additional iron oxidases may yet be discovered. Genes encoding the larger multiheme
cytochrome complexes PCC3_1 and PCC3_2 are expressed at higher than mean levels. PCC3s
are predicted to be large outer membrane cytochrome (OMC)-porin complexes (Fig. 6) (24), and
while their function has not been tested, as OMCs, they could in principle oxidize iron. Thus,
based on the *Sideroxydans* sp. CL21 transcriptome, there are at least four possible iron oxidases
(Cyc2, MtoA, and the PCC3_1 and PCC3_2 cytochromes) with various proteins that could direct
electrons to terminal oxidases or reverse electron transport pathways.

One might expect the expression of iron oxidase genes to correlate with iron oxidation,
but our results show that the well-characterized iron oxidase genes *cyc2* and *mtoA* are always
highly expressed in our experiments, and thus do not correspond to iron oxidation ability. This
differs from *S. lithotrophicus* ES-1, which has very low expression of *mtoA* on dissolved Fe^{2+} but
upregulates *mtoA* when grown on solid Fe(II)-containing minerals smectite and magnetite (10,
22, 23) . We showed that *Sideroxydans* sp. CL21 grows on magnetite, and though magnetite was
not present in lactate-grown cultures, expression of *mtoA* may still indicate a readiness to take up
electrons from solid minerals. *Sideroxydans* sp. CL21 is more like *S. lithotrophicus* ES-1 in its
high expression of *cyc2*. In *S. lithotrophicus* ES-1, expression of *cyc2* (one of three copies) is
high even when cells are cultured on thiosulfate (10). Similarly, *Sideroxydans* sp. CL21 highly
expresses two of its three *cyc2* copies when grown on lactate. Expression of *cyc2* is also
generally high in marine iron-oxidizing Zetaproteobacteria (20). However, in *S. lithotrophicus*
ES-1 additional copies of *cyc2* are iron-responsive (10). In theory, Cyc2 production could be
post-transcriptionally regulated. However, this is not the case in *S. lithotrophicus* ES-1, since
proteome data shows Cyc2 is highly expressed on Fe(II) (dissolved Fe^{2+} and magnetite (23)) and
thiosulfate (25). One explanation for the overall high expression is that these FeOB maintain
readiness or, in the case of *Sideroxydans* sp. CL21, partial readiness to oxidize iron. In any case,

because *cyc2* expression does not correspond to iron oxidation activity, *cyc2* (alone) is not an
accurate indicator of biotic iron oxidation.
Beyond iron oxidases, iron oxidation activity also requires electron carrier proteins in the
periplasm and inner membrane. Currently, there is limited evidence supporting other essential
components of the pathway, particularly periplasmic electron carriers and inner membrane
proteins (Table S3). Our findings contribute to filling this gap by providing evidence for the
involvement of genes encoding predicted periplasmic and inner membrane proteins. In fact, the
genes that were most upregulated in cells that showed iron oxidation (Day 14) were periplasmic
c-type cytochrome genes *ircA*, *ircB*, and *mtoD* (Fig. 6), followed by the inner membrane protein
genes *ircC*, *ircD*, and the PCC3_2 inner membrane component. There is accumulating evidence
for the involvement of *irc* genes in iron oxidation. Homologs are widespread among
Gallionellaceae and found in several other FeOB including *Leptothrix cholodnii* SP-6,
Rhodoferax spp., and several Zetaproteobacteria (Table S2; (19, 21)). In *S. lithotrophicus* ES-1,
irc genes are upregulated on iron-oxidizing cultures versus thiosulfate-oxidizing ones (10). The
periplasmic di- and mono-heme *c*-type cytochrome genes (*ircA* and *ircB*) are homologs of *grcJ*
and *grcI* in *Ghiorsea bivoria* TAG-1. In *G. bivoria* TAG-1, GrcJ is significantly upregulated
under iron-oxidizing vs. H_2-oxidizing conditions and expression of *grcJ* increases over time
when cells were grown with Fe(II) (21). In lactate-grown *Sideroxydans* sp. CL21, upregulated
expression of *ircABCD* corresponded to the cells' potential to oxidize iron in late-log cultures
(Day 14). Thus, it appears that cells increase expression of electron carrier genes like *irc*, *grc*,
and *mtoD* when cells are ready to oxidize iron, making them the best candidates thus far for
markers of iron oxidation potential.
The results to date suggest that the iron oxidation pathway is not regulated as a complete
set in the neutrophilic FeOB Gallionellaceae. Instead, at least one OMC (one Cyc2) is
constitutively expressed, which could serve multiple purposes. Cyc2 may operate as an Fe(II)
sensor to tell the cell when it is time to increase expression of the iron oxidation pathway. An
issue with this pathway is that OMCs are complicated to produce; they must be secreted to the
periplasm, where the heme cofactor is attached and then inserted into the outer membrane with
proper folding. Each of these three steps involves separate machinery, so this requires substantial
space in the inner membrane and periplasm, which may interfere with the electron transfer
pathway. Thus, expressing Cyc2 and other OMCs ahead of iron oxidation may be a strategy for a
quicker iron oxidation response. In contrast to the OMC, many of the periplasmic and inner
membrane components are smaller and would be easier to produce when they are required. This
strategy also allows the organism to express different periplasmic and inner membrane proteins
to match different iron substrates (Fe(II), magnetite, chelated Fe(II)), which have varied redox
potentials (36, 37), as is the case in *Geobacter* (38).
Thus, we conclude that the Gallionellaceae may implement a multistage production of the
iron oxidation pathway, with a subset of the OMCs expressed proactively, followed by additional
OMCs and the rest of the pathway once iron oxidation is required. This may be further tested as
additional facultative FeOB are isolated. In the meantime, we will need to carefully parse the

expression profiles to interpret FeOB activity in the environment, though the *irc/grc* genes are
promising markers.

C utilization among FeOB

Sideroxydans sp. CL21 is distinct among the current Gallionellaceae isolates in its ability
to grow on organic acids. However, other uncultured mixotrophic or organo(hetero)trophic
FeOB are likely present in C-rich environments. A pangenomic analysis of *Gallionellaceae*
isolates and environmental MAGs reveals that the potential for lactate utilization is rare but not
unique, as it has been observed in some genomes from the Crystal Bog humic lake (Wisconsin)
and freshwater lakes near Kuujjuarapik-Whapmagoostui (Canada) (19, 39–41). Active organic
substrate utilization by FeOB fundamentally alters their role in the carbon cycle. While FeOB are
traditionally considered autotrophs that fix CO_2, the ability to grow organotrophically suggests
they may also contribute to the degradation of organic carbon in situ. The ability to grow on
organics may provide a significant ecological advantage in specific habitats, especially wetlands
and peatlands. In peatlands, mixotrophic FeOB face competition with both aerobic and anaerobic
microorganisms for organic intermediates such as lactate and pyruvate. However, FeOB exhibit a
unique advantage by coupling the consumption of these substrates to aerobic respiration under
microaerophilic conditions. This metabolic flexibility of FeOB enables them to thrive in
environments like peatlands, where fluctuating water tables create dynamic oxic-anoxic
gradients, offering transient microaerophilic niches. Thus, such versatility may be key to their
survival and dominance in specific ecosystems, and play integral roles in biogeochemical iron
and C cycling in dynamic environments.

Implications of FeOB organotrophy for wetland C cycling

By demonstrating the versatility of FeOB metabolism and its genetic basis, our findings
provide a foundation for understanding the broader biogeochemical roles of FeOB in wetland
ecosystems and how they connect iron and C cycles. It is already recognized that organotrophic
iron-reducing bacteria can compete for organic carbon, thus shunting carbon away from
methanotrophs, which could reduce wetland methane emissions (42–46). FeOB help promote
this process by replenishing the supply of Fe(III). If the FeOB are autotrophic, then iron
oxidation drives the regeneration of organic C, partly reversing the effect of organotrophic iron
reduction and muting the effects on methanogenesis. But if the FeOB are organotrophic, they
would join the competition for organics and, if sufficiently abundant and active, contribute to the
suppression of methanogenesis. These are theoretical effects; evaluating the actual effects of
FeOB organotrophy requires specifically monitoring FeOB C metabolism in the wetland
environment and soil incubations. The potential iron oxidation gene markers would give us a
way to link iron oxidation activity to organotrophic metabolisms via genome-resolved
metatranscriptomics, paving the way toward deeper insights into FeOB influences on wetland
biogeochemistry and interactions with wetland communities. This knowledge is essential for

predicting how iron and C cycling—and associated greenhouse gas fluxes—will respond to environmental changes.

Methods

Growth experiments

For all growth experiments, *Sideroxydans* sp. CL21 was cultured in triplicate, in 150 mL serum bottles with 80 mL of modified Wolfe's minimal medium (MWMM; containing 1.0 g NH_4Cl, 0.5 g $MgSO_4 \cdot 7H2O$, 0.2 g $CaCl_2$, and 0.05 g K_2HPO_4 per liter). All cultures were amended with a 0.1% vol/vol addition each of ATCC trace vitamins and trace minerals. To keep micromolar concentrations of oxygen, headspaces were flushed daily with low-oxygen gas mix (2% O_2, 20% CO_2, 76% N_2). A Firesting oxygen spot sensor (Pyro Science) was used to measure and monitor oxygen concentrations. With the exception of lactate and pyruvate cultures, all bottles were incubated on their sides in a dark culturing cabinet at room temperature. Lactate and pyruvate bottles were incubated upright in a New Brunswick Innova 42 shaking incubator at 20°C, 75 RPM. Growth for all cultures was measured by direct cell counting whereby cells were stained with Syto-13 then counted manually in a Petroff-Hauser counting chamber under fluorescence microscopy.

Lactate cultures included a one-time addition of sodium DL-lactate (CAS 72-17-3) for a concentration of 5 mM and were buffered with 10mM of 2-(N-morpholino)ethanesulfonic acid (MES) at pH 6.0. Likewise, citrate (CAS 6132-04-3), pyruvate (CAS 113-24-6), thiosulfate (CAS 10102-17-7), and S(0) cultures were buffered with MES at pH 6.0 and given a one-time dose of sodium citrate dihydrate, sodium pyruvate, sodium thiosulfate pentahydrate, or elemental sulfur powder for a starting concentration of 5 mM. Biogenic and commercial magnetite cultures were each given a single addition of magnetite (1 g/L). Biogenic magnetite was synthesized by incubating *Shewanella oneidensis* MR-1 in Luria-Bertani (LB) media with two-line ferrihydrite for 10-12 days in an anerobic chamber. After incubation a magnet was used to hold the magnetite in place while the supernatant was decanted. Minerals were washed 3x with deoxygenated water, then dried in an anaerobic desiccator.

$FeCl_2$ cultures were buffered with 50 mM of MES at pH 6.0. These cultures were given a daily addition of $FeCl_2$ ranging from 250 μM on Day 0 to 500 μM on Day 14. The $FeCl_2$ + citrate experiment was conducted under these same conditions plus a one-time addition of 2.5 mM sodium citrate.

Organic acids measurements

Samples from lactate and pyruvate cultures were collected, filtered with a 0.22 μm syringe filter, and stored at 4°C. Concentration of organic acids (lactate, pyruvate) were

measured by high-performance liquid chromatography (HPLC). Standards for formate and
acetate were also included to check for partial oxidation products, but neither were detected.
Measurements were made on a Shimadzu HPLC with a Photodiode Array Detector (model
MPD-40) with a Alltech Prevail Organic Acid 5 µm 150 x 4.6 mM column and guard column.
Organic acids were eluted at room temperature with 25 mM KH_2PO_4 at pH 2.5, adjusted with
phosphoric acid, at 1 mL/minute. Absorbance was read at 210 nm to quantify compounds based
on peak area. Instrument control and data collection were performed using LabSolutions
software (v. 5.106 SP1). Peak areas were exported to Microsoft Excel where calibration curves
were generated by linear regression of authentic standard peak areas and used to calculate
concentrations.

Transcriptome experiment

Sideroxydans sp. CL21 was cultured in 5 mM lactate as described above. Bottles were
inoculated with stationary phase, lactate-adapted cells. Cell counts and lactate measurements
were taken on days 0, 3, 6, 9, 14. On days 6, 9, and 14, aliquots of cultures were used in Fe(II)
spike assays (described below) and bottles were sacrificed in biological sextuplets for
transcriptome sequencing. For all sacrifices, a 1:10 vol/vol of stop solution (buffer-saturated
phenol:absolute ethanol at a 1:9 vol/vol) was injected into the bottles to stop transcription. Cells
were then harvested by filtering onto a 0.22 µm polyethersulfone (PES) filter and stored at -80°C
until RNA extraction.

Fe(II) spike assay

Fe(II) spike assays were performed at different timepoints to test *Sideroxydans* sp.
CL21's response to Fe(II) when grown on lactate and pyruvate. For these assays, 10 mL of
culture were extracted from each of the six inoculated serum bottles. Of this, 5 mL was incubated
in 5 mM sodium azide for 20 minutes to create the azide-killed control. The other 5 mL was
untreated and served as the "live cell" samples. A 5 mL aliquot was taken from six uninoculated
bottles for the uninoculated control. All samples were bubbled with a 2% O_2, 20% CO_2, 78% N_2
gas mix throughout the experiment to keep oxygen levels consistent and low. Each sample, live
and control, was spiked with 20 µL of 100 mM $FeCl_2$ for a 400 µM starting concentration.
Samples were collected at 0, 5, 15, 30, and 60 minutes and preserved in a 1:1 mix with 40 mM
sulfamic acid. After all samples were collected, a spectrophotometric ferrozine assay was used to
measure ferrous iron concentrations (47, 48).

RNA extraction and sequencing

RNA was extracted from cells collected on PES filters using a RNeasy PowerSoil RNA
kit (Qiagen). Extracted RNA was treated with three RNase inhibitors (1 µL Invitrogen
SUPERase, 3 µL RNaseOUT, 1.25 µL Ambion RNase Inhibitor) to prevent degradation. Total

RNA was quantified using a Qubit RNA HS assay kit from Invitrogen and quality was assessed
with an Agilent fragment analyzer. A Zymo RNA Clean & Concentrator™-5 kit was used to
remove excess salts and DNA from all samples. After clean-up, three samples with the best yield
and quality from each timepoint (Day 6 - 12.6, 13.2, 29.6 ng/µL; Day 9 – 34.6, 43.0, 25.6 ng/µL;
Day 14 – 82.0, 20.0, 92.0 ng/µL), along with sample 6A from Day 9 (33.4 ng/µL) were selected
for sequencing. Ribosomal RNA depletion and library preparation were done using a Zymo-Seq
RiboFree® Total RNA Library Kit. Sixty-four ng of total RNA from each samples was
sequenced on an Illumina NextSeq 2000 using a P2 100-cycle protocol yielding up to 40 GB of
data (up to 400 million single-end reads). All clean-up, library prep and sequencing was
performed by the University of Delaware DNA Sequencing & Genotyping Center.

Transcriptome QC and analysis

Raw transcriptome reads were quality checked using FastQC v0.11.9 (49). Reads were
trimmed and adaptor sequences were removed using TrimGalore! v.0.6.6 (50) and Cutadapt v4.4
(51). A minimum quality score of 28 and minimum length of 75 bp were applied to filter out
poor reads. Post-trim, the read quality was reassessed with FastQC v0.11.9 (49) and SortMeRNA
v4.3.6 (52) was run on all samples to remove any rRNA sequences before mapping.
Reads were mapped to the closed, RefSeq annotated genome of *Sideroxydans* sp. CL21
(GCF_902459525.1) using Bowtie 2 v2.5.1 (53) and Samtools v1.10 (54). The average
alignment rate per sample was 97%, with the exception of Day 6, Sample 3B which was 73%.
Read counts were reported using Htseq v2.0.2 (55). After comparing the total number of reads
mapped for each sample and the overall alignment rates, we determined that Day 6, Sample 3B
was an outlier with potential contamination and too few quality reads to be useful. To be
thorough, we ran and compared the DGE analysis (described below) with and without Day 6
Sample 3B. Exclusion of Day 6 Sample 3B did not affect the overall findings or change whether
the differential expression of our genes of interest was significant. Instead, removing the outlier
decreased noise and made the dataset more precise.
Differential gene expression was analyzed in R v4.4.0 (56) with RStudio v2023.06.0
Build 421 (57) using the DESeq2 v1.26.0 (58) package. Data was normalized using DESeqs
default algorithm which uses an estimation of size factors to control for differences in
sequencing depth. P-values were determined with the Wald Test and adjusted for false discovery
rate (FDR) using the Benjamini-Hochberg (BH) correction. An alpha of 0.01 was set as the cut-
off for significance. Reports of normalized count data for each sample were generated using the
regularized logarithm (rlog) transformation (58). This rlog counts data was used to calculate Z-
score to analyze the variance of individual genes across samples. To test validity of these
transformations, we checked the expression of housekeeping genes for DNA gyrase and RNA
polymerase (*gyrA, gyrB*, and *rpoA*). Their even expression over all timepoints indicates the rlog
and Z-score transformations were successful (Fig. S1A, Table S1).

Identification and analysis of the Fe(II)-responsive (*irc*) gene cluster

Homologs of the Fe(II)-responsive (*irc*) gene cluster were identified by using cblaster (59) to search the NCBI database for BLAST hits (e-value cut-off < 1E-10) to the *irc* genes of *Sideroxydans* sp. CL21 that were located within 20,000 bp of each other in their respective genomes. Positive hits needed to contain at least three of the genes, excluding the Hsp33 chaperone. Results were visualized using clinker (60). PSORTb 3.0 (61), Gneg-mPLoc (62), PredictProtein (63), and InterProScan (64, 65) were used to predict subcellular localization and domains of the *irc* genes.

Data availability

Transcriptomic data were uploaded to NCBI BioProject PRJNA1216980.

Acknowledgements

This work was funded by the National Science Foundation (EAR-1833525 to C.S.C., MCB-1817651 to C.S.C.) and the Office of Naval Research (N00014-17-1-2640 C.S.C.). R.L.H. was also supported by fellowships from University of Delaware Graduate College and the Microbiology Program/Unidel Foundation. Support from the University of Delaware Center for Bioinformatics and Computational Biology Core Facility (RRID:SCR_017696), including the use of the BIOMIX computer cluster, was made possible through funding from Delaware INBRE (NIH P20GM103446), the State of Delaware, and the Delaware Biotechnology Institute. KK was supported by the Collaborative Research Centre Chemical Mediators in Complex Biosystems (SFB 1127 ChemBioSys, Project Number 239748522), which is funded by the Deutsche Forschungsgemeinschaft.

We thank Thomas Hanson for his help with the HPLC analyses, Christian Sinjari for his assistance culturing *Sideroxydans* sp. CL21 on magnetite, Jessica Keffer for feedback and editing, and Mark Shaw for sample clean-up and sequencing.

We declare no conflict of interest.

Figure Captions

FIG 1. Growth of *Sideroxydans* sp. CL21 on: (A) FeCl$_2$, FeCl$_2$ + citrate, and citrate compared to a no substrate control; (B) thiosulfate, S(0), biogenic and commercial magnetite; (C) lactate, and

514   (D) pyruvate. Red arrows in panels C indicate days cells were sampled for the Fe(II) spike
515   experiments and transcriptome sequencing. Red arrows in panel D indicate days cells were
516   sampled for Fe(II) spike experiments. Error bars represent one standard deviation between three
517   replicates.
518
519   **FIG 2.** Consumption of (A) lactate and (B) pyruvate over time by live *Sideroxydans* sp. CL21 in
520   growth experiments (Fig. 1C, 1D) versus uninoculated controls. Arrows indicate sampling points
521   for short-term iron oxidation assays (Fig 3). Error bars represent one standard deviation between
522   three replicates.
523
524   **FIG 3.** Results of Fe(II) spike assays for lactate- and pyruvate-grown cells. Aliquots of live,
525   lactate-grown *Sideroxydans* sp. CL21 from the transcriptome experiment (Fig. 1C, 2A) rapidly
526   oxidized iron on Day 14 compared to Day 6 or Day 9 (A). Cells from pyruvate growth
527   experiments (Fig. 1D, 2B) rapidly oxidized iron on both Day 6 and Day 14 (B). Error bars
528   represent one standard deviation between three replicates.
529
530   **FIG 4.** A) The six lactate oxidation genes encoded by *Sideroxydans* sp. CL21 and B) their
531   patterns of expression on days 6, 9, and 14. Expression is represented by Z-score, which shows
532   the expression of each gene relative to the mean gene expression (0) at the time point. Error bars
533   represent the spread of the interquartile range.
534
FIG 5. Boxplots of Z-score normalized expression for each time point with heatmaps showing
the Log2 Fold Change in expression between Day 9 and Day 14 versus Day 6. Expression in the
boxplots is represented by Z-score, which shows the expression of each gene relative to the mean
gene expression (0) at the time point. Z-score was calculated from rlog normalized count data
from DESeqs2. Heatmaps above the boxplots represent the Log2 Fold Change in average gene
expression between Day 9 and Day 14 versus Day 6, with statistically significant p-values noted
with asterisks (*). Data is shown for: (A) *cyc2* iron oxidase genes; (B) *mtoA* iron oxidase, *mtoB*
porin, and *mtoD*, *imoA/cymA* periplasmic *c*-type cytochrome genes; and (C) *ircABCD* Fe(II)-
responsive genes. Error bars represent the spread of the interquartile range. A gene map (D)
shows the co-location of genes in the CL21 genome. Slashes breaking the line indicate genes are
distant and "hyp." denotes a gene encoding a hypothetical protein.
546
547   **FIG 6.** Cartoon schematic of iron oxidases, outer membrane cytochrome, porin, periplasmic, and
548   inner membrane proteins colored by log2 fold change (L2FC) in normalized, rlog transformed
549   gene expression between Day 14 and Day 6. Positive numbers (purple) were upregulated on Day
550   14 relative to Day 6. Asterisks indicate the gene encoding a particular protein was significantly
551   upregulated.
552

Supplementals

Figures

- **Figure S1** – Line graphs of iron oxidation response of individual bottles on A) Day 6, B) Day 9, and C) Day 14.
- **Figure S2** – Boxplot/heatmap of A) housekeeping genes, B) RuBisCo, C) PCC3 copy 1, D) PCC3 copy 2

Tables

- **Table S1** – Differential expression, normalized expression, and Z-score data for key genes
- **Table S2** – *ircABCD* homologs in FeOB and FeRB
- **Table S3** – Table of existing evidence for the involvement of various genes and proteins in iron oxidation and/or extracellular electron transport.

References

1.   Borch T, Kretzschmar R, Kappler A, Cappellen PV, Ginder-Vogel M, Voegelin A,
Campbell K. 2010. Biogeochemical redox processes and their impact on contaminant
dynamics. Environ Sci Technol 44:15–23.

2.   Barros MEC, Rawlings DE, Woods DR. 1984. Mixotrophic growth of a *Thiobacillus*
ferrooxidans strain. Appl Environ Microbiol 47:593–595.

3.   Akob DM, Hallenbeck M, Beulig F, Fabisch M, Küsel K, Keffer JL, Woyke T, Shapiro N,
Lapidus A, Klenk H-P, Chan CS. 2020. Mixotrophic iron-oxidizing *Thiomonas* isolates
from an acid mine drainage-affected creek. Appl Environ Microbiol 86:e01424-20.

4.   Fleming EJ, Woyke T, Donatello RA, Kuypers MMM, Sczyrba A, Littmann S, Emerson D.
2018. Insights into the fundamental physiology of the uncultured Fe-oxidizing bacterium
Leptothrix ochracea. Appl Environ Microbiol 84:e02239-17.

5.   Hallbeck L, Pedersen K. 1991. Autotrophic and mixotrophic growth of *Gallionella*
ferruginea. Microbiology, 137:2657–2661.

6.   Tothero GK, Hoover RL, Farag IF, Kaplan DI, Weisenhorn P, Emerson D, Chan CS. 2024.
Leptothrix ochracea genomes reveal potential for mixotrophic growth on Fe(II) and organic
carbon. Appl Environ Microbiol 0:e00599-24.

7.   Sobolev D, Roden EE. 2004. Characterization of a neutrophilic, chemolithoautotrophic
Fe(II)-oxidizing β -Proteobacterium from freshwater wetland sediments. Geomicrobiol J
21:1–10.

8.   Eggerichs T, Wiegand M, Neumann K, Opel O, Thronicker O, Szewzyk U. 2020. Growth
of iron-oxidizing bacteria *Gallionella ferruginea* and *Leptothrix cholodnii* in oligotrophic
environments: Ca, Mg, and C as limiting factors and *G. ferruginea* necromass as C-source.
Geomicrobiol J 37:190–199.

9.   Emerson D, Moyer C. 1997. Isolation and Characterization of novel iron-oxidizing bacteria
that grow at circumneutral pH. Appl Environ Microbiol 63:9.

10.   Zhou N, Keffer JL, Polson SW, Chan CS. 2021. Unraveling Fe(II)-oxidizing mechanisms
in a facultative Fe(II) oxidizer, *Sideroxydans lithotrophicus* strain ES-1, via culturing,
transcriptomics, and reverse transcription-quantitative PCR. Appl Environ Microbiol
88:e01595-21.

11.   Kato S, Itoh T, Iino T, Ohkuma M. 2022. *Sideroxyarcus emersonii* gen. nov. sp. nov., a
neutrophilic, microaerobic iron- and thiosulfate-oxidizing bacterium isolated from iron-rich
wetland sediment. Int J Syst Evol Microbiol 72.

12.   Garry M, Farasin J, Drevillon L, Quaiser A, Bouchez C, Le Borgne T, Coffinet S, Dufresne
A. 2024. *Ferriphaselus amnicola* strain GF-20, a new iron: and thiosulfate-oxidizing
bacterium isolated from a hard rock aquifer. FEMS Microbiol Ecol 100:fiae047.

13.   Lüdecke C, Reiche M, Eusterhues K, Nietzsche S, Küsel K. 2010. Acid-tolerant
microaerophilic Fe(II)-oxidizing bacteria promote Fe(III)-accumulation in a fen. Environ
Microbiol 12:2814–2825.

14.   Küsel K, Blöthe M, Schulz D, Reiche M, Drake HL. 2008. Microbial reduction of iron and
porewater biogeochemistry in acidic peatlands. Biogeosciences 5:1537-1549.

15.   Hädrich A, Taillefert M, Akob DM, Cooper RE, Litzba U, Wagner FE, Nietzsche S,
Ciobota V, Rösch P, Popp J, Küsel K. 2019. Microbial Fe(II) oxidation by *Sideroxydans*
lithotrophicus ES-1 in the presence of Schlöppnerbrunnen fen-derived humic acids. FEMS
Microbiol Ecol 95:fiz034.

16.   Kügler S, Cooper RE, Wegner C-E, Mohr JF, Wichard T, Küsel K. 2019. Iron-organic
matter complexes accelerate microbial iron cycling in an iron-rich fen. Sci Total Environ
646:972–988.

17.   Cooper RE, Wegner C-E, McAllister SM, Shevchenko O, Chan CS, Küsel K. 2020. Draft
genome sequence of *Sideroxydans* sp. strain CL21, an Fe(II)-oxidizing bacterium.
Microbiol Resour Announc 9:e01444-19.

18.   Cooper RE, Finck J, Chan C, Küsel K. 2023. Mixotrophy broadens the ecological niche
range of the iron oxidizer *Sideroxydans* sp. CL21 isolated from an iron-rich peatland.
FEMS Microbiol Ecol 99:fiac156.

19.   Hoover RL, Keffer JL, Polson SW, Chan CS. 2023. Gallionellaceae pangenomic analysis
reveals insight into phylogeny, metabolic flexibility, and iron oxidation mechanisms.
mSystems 8:e00038-23.

20.   McAllister SM, Polson SW, Butterfield DA, Glazer BT, Sylvan JB, Chan CS. 2020.
Validating the Cyc2 Neutrophilic Iron oxidation pathway using meta-omics of
Zetaproteobacteria iron mats at marine hydrothermal vents. mSystems 5:e00553-19

21.   Barco RA, Merino N, Lam B, Budnik B, Kaplan M, Wu F, Amend JP, Nealson KH,
Emerson D. 2024. Comparative proteomics of a versatile, marine, iron-oxidizing
chemolithoautotroph. Environ Microbiol 26:e16632.

22.   Zhou N, Kupper RJ, Catalano JG, Thompson A, Chan CS. 2022. Biological oxidation of
Fe(II)-bearing smectite by microaerophilic iron oxidizer *Sideroxydans lithotrophicus* using
dual Mto and Cyc2 iron oxidation pathways. Environ Sci Technol 56:17443-17453.

23. Keffer JL, Zhou N, Rushworth DD, Yu Y, Chan CS. 2024. Microbial magnetite oxidation
via MtoAB porin-multiheme cytochrome complex in *Sideroxydans lithotrophicus* ES-1.
bioRxiv https://doi.org/10.1101/2024.09.20.614158. In press at AEM.

24. He S, Barco RA, Emerson D, Roden EE. 2017. Comparative genomic analysis of
neutrophilic iron(II) oxidizer genomes for candidate genes in extracellular electron transfer.
Front Microbiol 8:1584.

25. Zhou N. 2022. The molecular mechanisms and organic effects on microaerophilic Fe(II)
oxidation by *Sideroxydans Lithotrophicus* ES-1. Doctoral dissertation, University of
Delaware.

26. Cooper RE, Wegner C-E, Kügler S, Poulin RX, Ueberschaar N, Wurlitzer JD, Stettin D,
Wichard T, Pohnert G, Küsel K. 2020. Iron is not everything: unexpected complex
metabolic responses between iron-cycling microorganisms. The ISME Journal 14:2675–
2690.

27. Ohta Y, Nogi Y, Miyazaki M, Li Z, Hatada Y, Ito S, Horikoshi K. 2004. Enzymatic
properties and nucleotide and amino acid sequences of a thermostable β-agarase from the
novel marine isolate, JAMB-A94. Biosci Biotechnol Biochem 68:1073–1081.

28. Bannikova GE, Lopatin SA, Varlamov VP, Kuznetsov BB, Kozina IV, Miroshnichenko
ML, Chernykh NA, Turova TP, Bonch-Osmolovskaya EA. 2008. The thermophilic bacteria
hydrolyzing agar: Characterization of thermostable agarase. Appl Biochem Microbiol
44:366–371.

29. Badger MR, Bek EJ. 2008. Multiple Rubisco forms in proteobacteria: their functional
significance in relation to CO_2 acquisition by the CBB cycle. J Exp Bot 59:1525–1541.

30. Beckwith CR, Edwards MJ, Lawes M, Shi L, Butt JN, Richardson DJ, Clarke TA. 2015.
Characterization of MtoD from *Sideroxydans lithotrophicus*: a cytochrome c electron
shuttle used in lithoautotrophic growth. Front Microbiol 6:332.

31. Jain A, Kalb MJ, Gralnick JA. 2022. Reconstructing electron transfer components from an
Fe(II) oxidizing bacterium. Microbiology 168:001240.

32. Jain A, Coelho A, Madjarov J, Paquete CM, Gralnick JA. 2022. Evidence for quinol
oxidation activity of ImoA, a novel NapC/NirT family protein from the neutrophilic Fe(II)-
oxidizing bacterium *Sideroxydans lithotrophicus* ES-1. mBio 13:e02150-22.

33. Liu J, Wang Z, Belchik SM, Edwards MJ, Liu C, Kennedy DW, Merkley ED, Lipton MS,
Butt JN, Richardson DJ, Zachara JM, Fredrickson JK, Rosso KM, Shi L. 2012.
Identification and Characterization of MtoA: a decaheme c-type cytochrome of the
neutrophilic Fe(II)-oxidizing bacterium *Sideroxydans lithotrophicus* ES-1. Front Microbio
3:37.

34. Keffer JL, McAllister SM, Garber AI, Hallahan BJ, Sutherland MC, Rozovsky S, Chan CS.
2021. Iron oxidation by a fused cytochrome-porin common to diverse iron-oxidizing
bacteria. mBio 12:e01074-21.

35. Kappler A, Bryce C, Mansor M, Lueder U, Byrne JM, Swanner ED. 2021. An evolving
view on biogeochemical cycling of iron. Nat Rev Microbiol 19:360–374.

36. Huang J, Jones A, Waite TD, Chen Y, Huang X, Rosso KM, Kappler A, Mansor M,
Tratnyek PG, Zhang H. 2021. Fe(II) Redox chemistry in the environment. Chem Rev 121:
8161-8233.

37. Levar CE, Rollefson DR, Bond. 2012. Microbial Metal Respiration: From Geochemistry to
Potential Applications. Springer Berlin Heidelberg p29-48.

38. Joshi K, Chan CH, Bond DR. 2021. Geobacter sulfurreducens inner membrane cytochrome
CbcBA controls electron transfer and growth yield near the energetic limit of respiration.
Mol Microbiol 116:1124–1139.

39. Rohwer RR, Kirkpatrick M, Garcia SL, Kellom M, McMahon KD, Baker BJ. 2025. Two
decades of bacterial ecology and evolution in a freshwater lake. Nat Microbiol 10:246–257.

40. Oliver T, Varghese N, Roux S, Schulz F, Huntemann M, Clum A, Foster B, Foster B, Riley
R, LaButti K, Egan R, Hajek P, Mukherjee S, Ovchinnikova G, Reddy TBK, Calhoun S,
Hayes RD, Rohwer RR, Zhou Z, Daum C, Copeland A, Chen I-MA, Ivanova NN, Kyrpides
NC, Mouncey NJ, del Rio TG, Grigoriev IV, Hofmeyr S, Oliker L, Yelick K,
Anantharaman K, McMahon KD, Woyke T, Eloe-Fadrosh EA. 2024. Coassembly and
binning of a twenty-year metagenomic time-series from Lake Mendota. Sci Data 11:966.

41. Buck M, Garcia SL, Fernandez L, Martin G, Martinez-Rodriguez GA, Saarenheimo J,
Zopfi J, Bertilsson S, Peura S. 2021. Comprehensive dataset of shotgun metagenomes from
oxygen stratified freshwater lakes and ponds. Sci Data 8:131.

42. Hädrich A, Heuer VB, Herrmann M, Hinrichs K-U, Küsel K. 2012. Origin and fate of
acetate in an acidic fen. FEMS Microbiol Ecol 81:339–354.

43. Roden EE, Wetzel RG. 1996. Organic carbon oxidation and suppression of methane
production by microbial Fe(III) oxide reduction in vegetated and unvegetated freshwater
wetland sediments. Limnol Oceanogr 41:1733–1748.

44. Roden EE, Wetzel RG. 2003. Competition between Fe(III)-reducing and methanogenic
bacteria for acetate in iron-rich freshwater sediments. Microb Ecol 45:252–258.

45. Reiche M, Torburg G, Küsel K. 2008. Competition of Fe(III) reduction and methanogenesis
in an acidic fen. FEMS Microbiol Ecol 65:88–101.

46.   Teh YA, Dubinsky EA, Silver WL, Carlson CM. 2008. Suppression of methanogenesis by
698   dissimilatory Fe(III)-reducing bacteria in tropical rain forest soils: implications for
699   ecosystem methane flux. Glob Change Biol 14:413–422.

700   47.   Stookey LL. 1970. Ferrozine—a new spectrophotometric reagent for iron. Anal Chem
701   42:779–781.

702   48.   Riemer J, Hoepken HH, Czerwinska H, Robinson SR, Dringen R. 2004. Colorimetric
703   ferrozine-based assay for the quantitation of iron in cultured cells. Anal Biochem 331:370–
704   375.

705   49.   Babraham Bioinformatics. FastQC a quality control tool for high throughput sequence data.
706   http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

707   50.   Krueger F. Trim Galore. The Babraham Institute.
708   https://github.com/FelixKrueger/TrimGalore

709   51.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
710   reads. 1. EMBnet.journal 17:10–12.

711   52.   Kopylova E, Noé L, Touzet H. 2012. SortMeRNA: fast and accurate filtering of ribosomal
712   RNAs in metatranscriptomic data. Bioinformatics 28:3211–3217.

713   53.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods
714   9:357–359.

715   54.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
716   R, 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map
717   format and SAMtools. Bioinforma Oxf Engl 25:2078–2079.

718   55.   Anders S, Pyl PT, Huber W. 2015. HTSeq—a Python framework to work with high-
719   throughput sequencing data. Bioinformatics 31:166–169.

720   56.   R Core Team. 2024. R: A Language and Environment for Statistical Computing. R
721   Foundation for Statistical Computing, Vienna, Austria.

722   57.   Posit team. 2023. RStudio: Integrated Development Environment for R (1.4.1717). Posit
723   Software, PBC, Boston, MA.

724   58.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
725   for RNA-seq data with DESeq2. Genome Biol 15:550.

726   59.   Gilchrist CLM, Booth TJ, van Wersch B, van Grieken L, Medema MH, Chooi Y-H. 2021.
727   cblaster: a remote search tool for rapid identification and visualization of homologous gene
728   clusters. Bioinforma Adv 1:vbab016.

60.   Gilchrist CLM, Chooi Y-H. 2021. clinker & clustermap.js: automatic generation of gene
cluster comparison figures. Bioinformatics 37:2473–2475.

61.   Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, Dao P, Sahinalp SC, Ester M, Foster
LJ, Brinkman FSL. 2010. PSORTb 3.0: improved protein subcellular localization prediction
with refined localization subcategories and predictive capabilities for all prokaryotes.
Bioinformatics 26:1608–1615.

62.   Shen H-B, Chou K-C. 2010. Gneg-mPLoc: A top-down strategy to enhance the quality of
predicting subcellular localization of Gram-negative bacterial proteins. J Theor Biol
264:326–333.

63.   Bernhofer M, Dallago C, Karl T, Satagopam V, Heinzinger M, Littmann M, Olenyi T, Qiu
J, Schütze K, Yachdav G, Ashkenazy H, Ben-Tal N, Bromberg Y, Goldberg T, Kajan L,
O'Donoghue S, Sander C, Schafferhans A, Schlessinger A, Vriend G, Mirdita M, Gawron
P, Gu W, Jarosz Y, Trefois C, Steinegger M, Schneider R, Rost B. 2021. PredictProtein -
predicting protein structure and function for 29 years. Nucleic Acids Res 49:W535–W540.

64.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,
Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-
Y, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics 30:1236–1240.

65.   Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, Nuka G,
Paysan-Lafosse T, Qureshi M, Raj S, Richardson L, Salazar GA, Williams L, Bork P,
Bridge A, Gough J, Haft DH, Letunic I, Marchler-Bauer A, Mi H, Natale DA, Necci M,
Orengo CA, Pandurangan AP, Rivoire C, Sigrist CJA, Sillitoe I, Thanki N, Thomas PD,
Tosatto SCE, Wu CH, Bateman A, Finn RD. 2021. The InterPro protein families and
domains database: 20 years on. Nucleic Acids Res 49:D344–D354.

A) Lactate-grown cells

B) Pyruvate-grown cells

Legend: Lactate (live culture)   Lactate (uninoc. control)   Pyruvate (live culture)   Pyruvate (uninoc. control)

A) Lactate-grown cells

Day 6 — Day 9 — Day 14

Fe(II) remaining (µM) vs Time (minutes)

B) Pyruvate-grown cells

Day 6 — Day 14

Fe(II) remaining (µM) vs Time (minutes)

Legend:
- Live cell cultures
- Azide killed controls
- Uninoculated controls

A

lldR, Lactate-responsive reg.
lldE, L-lactate dehyd.
lldG, L-lactate dehyd.
lldF, L-lactate dehyd.
dld, D-lactate dehydrogenase
lldP, Lactate permease

Sideroxydans sp. CL21

S. lithotrophicus ES-1

S. emersonii MIZ01

B

lldR, Lactate response regulator

lldE, L-lactate dehydrogenase

lldG, L-lactate dehydrogenase

lldF, L-lactate dehydrogenase

dld, D-lactate dehydrogenase

lldP, Lactate permease

z−score

Day 6   Day 9   Day 14