Targeted and random mutagenesis of cassava brown streak disease susceptibility

factors reveal molecular determinants of disease severity

ZJ Daniel Lin[1,3], Myia K. Stanton[1], Gabriela L. Hernandez[1], Elizabeth J. De Meyer[1], Zachary

von Behren[1], Katherine Benza[1,2], Helene Tiley[1], Emerald Hood[1], Greg Jensen[1], Kerrigan B.

Gilbert[1], James C. Carrington[1], Rebecca S. Bart[1]

1. Donald Danforth Plant Science Center, Saint Louis, MO 63132, USA

2. Washington University, Saint Louis, MO 63130, USA

3. corresponding author: ZDL: dlin@danforthcenter.org

Abstract

Cassava brown streak disease (CBSD) is caused by cassava brown streak viruses (CBSVs)

from the family *Potyviridae*. Potyvirid viral genome-linked protein (VPg) recruitment of host

eukaryotic translation initiation factor 4E (eIF4E) proteins is a critical step in the viral life cycle.

CBSV VPg interacts with all five cassava eIF4E-family members. Simultaneously knocking out

eIF4E-family genes *nCBP-1* and *nCBP-2*, in cultivar 60444, strongly reduces CBSD root

symptoms and viral titer but does not result in complete resistance, likely due to gene family

redundancy. To test for redundancy, we generated single and double mutants for each clade

of the *eIF4E* gene family in farmer preferred cultivar TME419. Double mutants for the

eIF(iso)4E and *nCBP* clades both exhibited reduced symptom severity, with *ncbp-1 ncbp-2*

having the strongest phenotype. A yeast two-hybrid screen for nCBP-2 mutants that lose VPg

affinity identified fifty-one mutants, including an L51F mutant. This finding is consistent with

one of the recovered cassava mutants that had a 6 amino acid deletion, including L51, in

nCBP-2 and showed a reduction in symptoms relative to wild type. The data presented here

suggest that generating mutations corresponding to L51F of nCBP-2 in multiple or all five

cassava eIF4E proteins may lead to stronger resistance to CBSD while avoiding pleiotropic

effects.

Keywords
Cassava brown streak disease, *Potyviridae*, eIF4E, viral genome-linked protein (VPg), gene
editing, yeast two-hybrid

Introduction: 717
Results: 1,845
Discussion: 1,519
Materials & Methods: 1,263

Introduction

Cassava is one of the most important starch crops grown in the world (*Agricultural production statistics 2000–2022*, 2023). In sub-Saharan Africa, cassava is particularly important for food security, but its production is threatened by viral diseases (Rey & Vanderschuren, 2017). In particular, cassava brown streak virus (CBSV) and Ugandan cassava brown streak virus (UCBSV), closely related ipomoviruses belonging to the family *Potyviridae*, are of major concern (Tomlinson *et al.*, 2018; Robson *et al.*, 2023). Independent or simultaneous infection by these viruses cause cassava brown streak disease (CBSD). Symptoms of CBSD include feathery chlorosis of leaves, brown streaking of stems, and necrosis of storage roots. Cassava varieties with varying levels of resistance have been identified and are being deployed, but none of the varieties currently available to farmers are fully immune (Sheat & Winter, 2023; Robson *et al.*, 2023). A highly effective RNAi-based transgenic approach has been developed but faces regulatory hurdles in many target countries (Odipio *et al.*, 2014; Beyene *et al.*, 2016; Wagaba *et al.*, 2016; Tomlinson *et al.*, 2018).

Viruses of *Potyviridae* are positive sense, single stranded RNA viruses (Valli *et al.*, 2015). Many of these viruses have been studied extensively as a large number are agronomically important pests (Revers & García, 2015; Yang *et al.*, 2021). In many potyvirus-plant pathosystems, members of the host *eIF4E* gene family function as susceptibility factors and are necessary for completion of the viral life cycle (Robaglia & Caranta, 2006; Naderpour *et al.*, 2010; Hart & Griffiths, 2013; Bastet *et al.*, 2017; Gomez *et al.*, 2019; Gao *et al.*, 2020). This gene family encodes mRNA cap-binding proteins that facilitate recruitment of translation machinery to transcripts (Browning & Bailey-Serres, 2015). The plant *eIF4E*-family can be broken into three clades: *eIF4E*, *eIF(iso)4E*, and *nCBP* (Gomez *et al.*, 2019)*.* Previous work in *Arabidopsis* demonstrated that turnip mosaic virus (TuMV) requires a functional copy of *eIF(iso)4E* for disease (Duprat *et al.*, 2002). Since then, potyvirus resistance loci in many crops have been mapped to *eIF4E*-family genes and alleles conferring potyvirus resistance often harbor missense mutations (Bastet *et al.*, 2017). Additional studies have demonstrated that potyvirid viral protein genome-linked (VPg) interacts with eIF4E-family proteins (Wittmann *et al.*, 1997; Schaad *et al.*, 2000) confirming that the interacting *eIF4E*-family member functions as a susceptibility factor (Yeam *et al.*, 2007; Charron *et al.*, 2008). Conversely, viral isolates that break *eIF4E*-family based resistance have been found to have accumulated compensating mutations in their VPg (Charron *et al.*, 2008; Gallois *et al.*, 2010; Li *et al.*, 2016; Lebaron *et al.*, 2016; Takakura *et al.*, 2018). VPg is covalently linked to the 5' end of the viral genome. It has been hypothesized that VPg interacts with eIF4E-family proteins to recruit translational machinery to synthesize the viral polypeptide (Sanfaçon, 2015) and complete the viral life cycle. Solving the structure of VPg-susceptibility factor protein complex could facilitate

engineering of susceptibility factor alleles that cannot be co-opted by the pathogen. However,

co-crystal structures have yet to be produced and the VPg protein complex with eIF4E-family

S factors falls below the current minimum size limit required for cryogenic electron microscopy.

To date, engineering synthetic *eIF4E* resistance alleles has involved transferring resistance-

conferring mutations from one pathosystem to another with some degree of success (Kim *et*

al., 2014; Bastet *et al.*, 2018, 2019; Zafirov *et al.*, 2021).

We previously found that simultaneous mutations in two cassava eIF4E-family

members, *nCBP-1* and *nCBP-2,* strongly attenuated CBSD storage root necrosis and

accumulation of CBSV in storage roots of cultivar 60444 (Gomez *et al.*, 2019). Knocking out

nCBP-2 alone resulted in an intermediate phenotype, while the *ncbp-1* single mutant exhibited

no differences from wild type. Aerial symptoms persisted in the *ncbp-1 ncbp-2* double mutant,

although with less severe stem brown streaking. This incomplete resistance suggests that VPg

may interact broadly with the remaining cassava eIF4E-family proteins. Indeed, CBSV VPg

could associate with all five cassava eIF4E-family proteins by coimmunoprecipitation (Gomez

et al., 2019). As full CBSD resistance may require engineering all cassava *eIF4E*-family

alleles to evade CBSV VPg, we employed targeted and random mutagenesis of nCBP-2 to

identify structural determinants of VPg interaction. Furthermore, to better understand the

degree of functional redundancy exhibited by cassava eIF4E-family proteins in CBSD, we here

expand our CRISPR/Cas enabled genetic approach to characterize the roles of cassava eIF4E

and eIF(iso)4E clades in CBSD.

Results

Generation of cassava *eIF4E*-family mutants

We previously reported that a cassava *ncbp-1 ncbp-2* double mutant in the 60444

background displayed increased, but incomplete, resistance to CBSV challenge (Taylor *et al.*,

2012; Gomez *et al.*, 2019). While VPg interacted most strongly with nCBP-1 and nCBP-2 in

yeast two-hybrid assays, all five eIF4E-family proteins interacted with VPg in co-

immunoprecipitation assays. These results suggest that there is likely functional redundancy

among this family of proteins. To test this idea, we used CRISPR/Cas9-mediated genome

editing to generate new cassava *eIF4E*-family mutants in TME419, a farmer preferred cassava

cultivar. gRNAs were designed to target the first exon of each target gene and cassava

transformations were performed as previously described (Fig. S2) (Taylor *et al.*, 2012; Gomez

et al., 2019). A total of 156 recovered lines from 21 transformations were characterized for

possible mutations at the target site(s) using amplicon sequencing. These efforts yielded

independent single mutant lines for all members of the cassava *eIF4E* gene family, along with

eif(iso)4e-1 eif(iso)4e-2 and *ncbp-1 ncbp-2* double mutants (Table 1, S3). All but one mutant

line used in this study are predicted to have frameshifting mutations that result in loss of protein

function. The lone exception is one of the *ncbp-1 ncbp-2* double mutants where both *nCBP-1*
alleles and one allele of *nCBP-2* have frameshifting mutations and the remaining *nCBP-2*
allele has a deletion that results in loss of amino acids 45 through 51. This allele is designated
nCBP-2$^{K45\_L51del}$ and the mutant line is designated *ncbp-1 nCBP-2$^{K45\_L51del}$* (Table 1, S3). No
differences in growth and development were observed between any of the mutant lines and
wild type (Fig. S3-4).
119

CBSD phenotypes of TME419 *eIF4E*-family mutants

To determine the contribution of cassava *eIF4E* family genes to development of CBSD,
mutant and wild type TME419 were challenged with CBSV isolate Naliendele in greenhouse
trials where stem brown streaking and leaf feathery chlorosis were tracked for approximately
three months. The specific CBSV isolate was chosen as it produces strong aerial symptoms
quickly after inoculation and reliably results in strong storage root necrosis (Wagaba *et al.*,
2013). At the end of the viral challenges, storage roots were excavated from soil, sectioned,
and scored for severity of necrosis as a measure of spread along storage root length and
diameter (Gomez *et al.*, 2019). These same root samples were used to measure viral titer by
quantitative real-time PCR (qRT-PCR). For aboveground symptoms, *eif4e*, *eif(iso)4e-1,*
eif(iso)4e-2, and *ncbp-1* single mutants had no reduction in disease severity relative to wild
type, while *ncbp-2* and both *eif(iso)4e* and *ncbp* double mutants exhibited small but significant
reductions (Figure 1a). For storage root necrosis symptoms in the single mutants, the *ncbp-1*
and *ncbp-2* mutants displayed a decrease in symptoms compared to wild type with the latter
showing the strongest phenotype (Figure 1b). Each of the double mutants showed a decrease
in symptom with the *ncbp-1 ncbp-2* mutant, where all *nCBP* alleles have frameshifting
mutations, having the greatest reduction in necrosis; virtually no necrosis was observed
(Figure 1b, d). We further measured viral accumulation in storage roots of wild type, *ncbp-1*,
ncbp-2, *ncbp-1 ncbp-2*, and *ncbp-1 nCBP-2$^{K45\_L51del}$* lines using qPCR analysis and observed
that lines with frameshifting mutations in *ncbp-2* had significantly lower virus titer (Figure 1c).
Taken together, these data suggest that while nCBP-2 is the primary contributor to root
disease severity, other eIF4E-family members may also contribute throughout the plant.
142

Purified CBSV VPg and cassava eIF4E-family proteins interact *in vitro*

To further explore functional redundancy within the eIF4E-family with respect to CBSD
and, specifically, interaction with VPg, we used a biochemical approach. Using co-
immunoprecipitation, we previously demonstrated that CBSV VPg and cassava eIF4E-family
proteins form complexes in whole cell lysates using co-immunoprecipitation (Gomez *et al.*,
2019). However, these experiments did not address whether the proteins are directly
interacting or if another complex member is present and facilitating the protein-protein

association. Thus, we tested if VPg directly associates with eIF4E-family proteins in the

absence of additional factors using a GST pull-down experiment. GST-tagged cassava eIF4E-

family proteins and VPg, tagged with both 6xHis and 3xFLAG, were expressed in and purified

from *E. coli* (Figure S5). We found that CBSV VPg was pulled down by GST-tagged eIF4E,

nCBP-1, and nCBP-2 in comparison to GST alone (Figure 2a, b). Pulldown of VPg by GST-

tagged eIF(iso)4E-1 and eIF(iso)4E-2 was also tested but we observed inconsistent

enrichment of VPg by GST-eIF(iso)4E-1 & -2, relative to GST alone, across experimental

replicates.

Targeted nCBP-2 mutations fail to disrupt interaction with VPg

If CBSV VPg can coopt all cassava eIF4E-family proteins for completing the viral

lifecycle, engineering cassava to be resistant to CBSV may require identifying alleles of *eIF4E-*

family genes encoding mutations that disrupt VPg binding but still allow for translation-initiation

activity. In other plant-potyvirid pathosystems, specific mutations within eIF4E-family members

have proved sufficient to block interaction with VPg while maintaining normal protein function

(Figure 2c) (Ashby *et al.*, 2011; Kim *et al.*, 2014). One well studied example is TuMV VPg and

Arabidopsis eIF(iso)4E. In this system, mutation of a highly conserved tryptophan (W92L) and

semi-conserved lysine (K149E) in *Arabidopsis* eIF(iso)4E disrupts binding of TuMV VPg in

pull-down and co-immunoprecipitation experiments. Overexpression of this mutant *eIF(iso)4E*

allele in Chinese cabbage confers resistance to TuMV (Miyoshi *et al.*, 2006; Kim *et al.*, 2014).

To test the hypothesis that specific *eIF4E* gene family alleles might avoid VPg binding, we

took a yeast two-hybrid approach using different alleles of *nCBP-2*. We recapitulated the

W92L/K149E mutations from Arabidopsis eIF(iso)4E in cassava nCBP-2, resulting in nCBP-

$2^{W118L/N170E}$ (Figure 2d). Whereas the eIF(iso)$4E^{W92L/K149E}$ no longer associated with TuMV VPg

in a co-immunoprecipitation experiment (Figure 2e), we observed that CBSV VPg was

similarly present in both wild type nCBP-2 and nCBP-$2^{W118L/N170E}$ pull downs (Figure 2f). VPg

- nCBP-$2^{W118L/N170E}$ interaction was also observed through yeast two-hybrid assay (Figure 2g,

h).

TuMV and CBSV belong to two different genera of *Potyviridae*, *Potyvirus* and

Ipomovirus. It is possible that viruses from these distinct genera have evolved mechanistic

differences in how their VPg proteins interact with host eIF4E-family *S* factors. To determine

VPg protein conservation between potyviruses and ipomoviruses, VPg sequences from five

major potyvirus clades were aligned with VPg sequences from seven of the eight known

ipomovirus species (TomMMoV, SPMMV, SqVYV, CocMoV, CVYV, UCBSV, CBSV) (Figure

S6). Across all VPg sequences analyzed, several highly conserved residues are observed, as

are striking genus-specific differences; four stretches of amino acids are present only in one

genus or the other (pink boxes). One of these stretches coincides with an α-α loop in PVY VPg

thought to directly interact with the ventral side of eIF4E (Coutinho de Oliveira *et al.*, 2019); in
ipomoviruses this region is eight amino acids shorter. Additionally, a highly conserved
aspartate residue (D77) necessary for TuMV VPg interaction with *Arabidopsis* eIF(iso)4E
(Léonard *et al.*, 2000), is not present in ipomoviruses, indicating this residue may have a less
conserved role in interactions between ipomovirus VPgs and eIF4E-family proteins.
192

193 A yeast two-hybrid screen identifies nCBP-2 mutants that lose VPg affinity

Since mutations from the known Arabidopsis resistance allele did not block nCBP-2 –
VPg interaction, we performed a yeast two-hybrid screen for nCBP-2 mutants that lose affinity
for VPg (Figure 3a). Mutant *nCBP-2* amplicons were generated by error-prone PCR and
cloned into the bait vector by *in vivo* recombination in EGY48 yeast already containing both
lacZ reporter and VPg prey vectors (Figure 3a). From roughly 1000 yeast transformants, 51
nCBP-2 mutants were isolated that lost interaction with VPg (Table 2, Table S4). Forty-eight
unique amino acid substitutions were observed across forty-six unique nCBP-2 mutants. Of
the unique mutants, 27 had mutations affecting a single residue or the introduction of a
frameshifting deletion while 19 contained two or three mutated residues. Nine residues were
isolated more than once, with up to three unique amino acid substitutions identified. Mutants
were then filtered to identify residues with varying degrees of conservation (Figure S7) across
cassava eIF4E-family proteins resulting in 15 being re-cloned and validated for loss of VPg
interaction in yeast using a quantitative β-galactosidase activity assay. All but two mutants,
nCBP-2^{L147H} and nCBP-2^{R196S}, were confirmed to confer total loss of β-galactosidase activity
in yeast and suggest commensurate loss of VPg affinity (Figure 3b, 5b, S8). The R196S
mutant yeast did not have any reduction of β-galactosidase activity relative to yeast expressing
wild-type nCBP-2, while L147H mutant yeast exhibited an intermediate reduction in β-
galactosidase activity. We further validated by western blot analysis that the yeast used for
two-hybrid experiments expressed nCBP-2 variants and CBSV VPg (Figure S8).
213

214 Translation initiation activity of isolated nCBP-2 mutants

Previous work with *Arabidopsis eif4e* or *eif(iso)4e* mutants suggested that both genes
were functionally redundant for translation initiation. However, *eif4e* mutants exhibit subtle
reductions in fertility and biomass in addition to delayed flowering time (Bastet *et al.*, 2017,
2018). Similarly, simultaneous knockout of tomato *eIF4E1* and *eIF4E2* incurs a growth penalty
(Gauffier *et al.*, 2016). T93C is a yeast *eif4e* mutant with chromosomal *eIF4E* replaced with
LEU2 (Altmann *et al.*, 1989). This yeast strain can survive on galactose media due to a
plasmid-based *eIF4E* gene that is galactose inducible but is repressed in the presence of
glucose (Figure 4a). Coding sequences of nCBP-2 mutants confirmed to lose affinity for CBSV
VPg by quantitative yeast two-hybrid were cloned into pG1.1 under control of the constitutive

yeast GPD promoter and transformed into T93C. Transfer of these T93C transformants from

galactose- to glucose-containing media revealed that only three nCBP-2 mutants rescue loss

of *eif4e*: nCBP-2$^{A20T/H94Q/L200P}$, nCBP-2^{L51F}, and nCBP-2^{S138P} (Figure 4bc, 5cd).

Mutating the conserved HPL motif of nCBP-2 can affect VPg binding and disease

Among the three nCBP-2 mutants that retained translation initiation activity, nCBP-

2^{L51F} stood out as having a mutation in the highly conserved HPL motif. In yeast, this motif

participates in docking with eIF4G (Grüner *et al.*, 2016, 2018). In addition, this amino acid is

deleted in the nCBP-2$^{K45\_L51del}$ cassava mutant line that displayed increased resistance to

CBSD (Figure 5a). The HPL motif is found in many plant eIF4E-family proteins and as such,

a mutation of the leucine may be suitable for engineering resistance to other potyvirids. We

tested the nCBP-2$^{K45\_L51del}$ mutant for interaction with VPg by quantitative yeast two-hybrid.

Yeast expressing this mutant exhibited no β-galactosidase activity, like yeast expressing

nCBP-2^{L51F} (Figure 5b, S9a). We confirmed this implied loss of VPg interaction by observing

strongly reduced VPg binding in GST pulldowns with both mutant proteins (Figure 6).

However, comparing wild-type nCBP-2, nCBP^{L51F} and nCBP-2$^{K45\_L51del}$ for translation initiation

activity in T93C yeast (Δ*eif4e*) showed that while nCBP-2^{L51F} was able to rescue loss of *eIF4E*

in T93C it was to a lesser degree than wild type nCBP-2, and nCBP-2$^{K45\_L51del}$ was not able to

complement T93C (Figure 4d, e). Leucine 51 of nCBP-2 corresponds to leucine 28 within

Arabidopsis eIF(iso)4E. We hypothesized that recapitulating the L51F substitution in

Arabidopsis eIF(iso)4E would have the similar effects on TuMV VPg binding. Using the same

quantitative yeast two-hybrid approach as above we observed that the L28F mutation strongly

disrupted eIF(iso)4E interaction with TuMV VPg (Figure 7a, S9b). However, it is not able to

rescue T93C *eif4e* mutant yeast (Figure 7b, c).

Discussion

Potyviridae is the largest plant-infecting family of RNA viruses comprised of 249

viruses across 12 genera (Walker *et al.*, 2022). The genus *Potyvirus* is perhaps the best

studied as it contains 206 of the 249 known viruses belonging to the family *Potyviridae*. In

comparison, only eight species of viruses belonging to the genus *Ipomovirus* have been

identified and much less is known about molecular ipomovirus-host interactions (Dombrovsky

et al., 2014). While the genome structure and genes encoded by potyviruses and ipomoviruses

are similar, and thus it is reasonable to expect a degree of conservation in virulence strategies,

it is also likely that differences have arisen given the evolutionary divergence of the two

genera. For CBSV in particular, we note that while eIF4E-family proteins are clear

susceptibility factors, the breadth of family members that contribute to disease appears

broader than typically found for potyviruses (Bastet *et al.*, 2017). Furthermore, while

resistance-conferring amino acid substitutions in eIF4E-family proteins have been effectively transferred between potyvirus-pathosystems, we find in this study that *Arabidopsis* eIF(iso)4E mutations that break TuMV VPg interaction fail to do so for the cassava nCBP-2 interaction with CBSV VPg. These observations highlight the need for further characterization of ipomovirus-host molecular interactions.

Structure-function analysis of VPg interaction with eIF4E-family susceptibility factors would strongly benefit from crystal or cryo-EM structures of the complex; these have yet to be successfully generated (Coutinho de Oliveira *et al.*, 2019). Most recently, an NMR approach solved the structure of PVY VPg (Coutinho de Oliveira *et al.*, 2019). The PVY VPg and human eIF4E interaction interface was also mapped through monitoring chemical shift perturbations of either protein during NMR spectroscopy. It is unclear how well these findings will translate to diverse potyvirid VPg interactions with cognate plant eIF4E-family proteins, especially for the plant specific eIF(iso)4E clade and considering significant divergence of ipomovirus VPg sequences from those of potyviruses. Thus far, attempts to design *eIF4E*-family resistance alleles have relied on inference-based approaches (German-Retana *et al.*, 2008; Ashby *et al.*, 2011; Kim *et al.*, 2014; Bastet *et al.*, 2018, 2019; Zafirov *et al.*, 2021). In contrast, here we utilize a loss-of-affinity screen of randomly mutagenized nCBP-2 variants to identify key residues necessary for interaction with CBSV VPg. Only three nCBP-2 amino acid residues that we found to be necessary for interaction with VPg had previously been described in potyvirus-plant pathosystems. These amino acid residues, D106, V164, and R168 in nCBP-2, are highly conserved across eIF4E-family proteins. In pepper, the *pvr1*[2] allele of *eIF4E* is responsible for resistance to isolates of PVY and TEV, and the amino acid corresponding to nCBP-2 D106 is one of three causal amino acid substitutions (Kang *et al.*, 2005). That aspartate in pvr1[2] is substituted with asparagine (D to N). For cassava nCBP-2, two separate D106 substitutions were observed; D106V and D106H, the latter of which co-occurred with an A41T substitution. Only the nCBP-2[D106V] mutant was tested for translation initiation activity and was found deficient in rescuing T93C *eif4e* yeast. It is unclear if a D106N mutation in nCBP-2, as found in pvr1[2], would similarly break VPg interaction while preserving its translation initiation activity. nCBP-2 V164, where mutation to leucine disrupts CBSV VPg interaction, corresponds to V167 in eIF4E of pea. The pea eIF4E[V167A] variant confers resistance to PSbMV and can rescue *eif4e* yeast. In contrast, cassava nCBP-2[V164L] does not rescue T93C *eif4e* yeast. Whether the V164A mutation in nCBP-2 would have the same effects as in pea eIF4E is an open question. Lastly, for nCBP-2 R168, the previously described NMR structural study demonstrated that corresponding residues in human eIF4E and wheat eIF(iso)4E are involved in the interaction with *Potato virus Y* VPg (Coutinho de Oliveira *et al.*, 2019). Like nCBP-2[D016V] and nCBP-2[V164L], nCBP-2[R168C] also failed to rescue T93C.

Of the fourteen nCBP-2 loss-of-VPg-interaction mutants that we tested for translation
initiation activity, eleven were nonfunctional. The three nCBP-2 variants that retained function
were nCBP-2$^{A20T/H94Q/A148V}$, nCBP-2^{L51F}, and nCBP-2^{S138P}. For nCBP-2 $^{A20T/H94Q/A148V}$, A20
resides outside of the conserved core of eIF4E family proteins while H94 and A148 are present
in alpha helices found on the dorsal side of nCBP-2; further work is required to identify which
individual or combination of these amino acids are required for CBSV VPg interaction. S138
is similarly located in a dorsal alpha helix while L51 is closer to the dorsal side and located N-
terminal of a beta strand. Evidence from a study on PVY VPg interaction with human eIF4E
indicates that VPg interfaces with the ventral side of eIF4E (Coutinho de Oliveira *et al.*, 2019).
If this also applies to CBSV VPg interaction with nCBP-2, the four aforementioned amino acid
residues likely affect VPg binding in an indirect manner. Of particular importance for this study,
the L51F substitution impacts the HPL motif, which was also affected in our cassava *ncbp-1*
nCBP-2$^{K45\_L51del}$ line. This enabled assessment of the importance of that motif for CBSD
symptom severity. Unlike nCBP-2^{L51F}, nCBP-2$^{K45\_L51del}$ did not rescue T93C *eif4e* deficient
yeast. Surprisingly, despite inability to strongly interact with VPg and complement *eif4e*
activity, the *ncbp-1 nCBP-2$^{K45\_L51del}$* root disease phenotype was intermediate to wild type and
ncbp-1 ncbp-2, indicating that this mutant version of *nCBP-2* is sufficient for the development
of some disease symptoms. A similar result has also been observed in pea *eIF4E* mutants
where of eight mutants unable to rescue *eif4e* mutant yeast, two were still able to confer
susceptibility to PsbMV (Ashby *et al.*, 2011). In such mutants, it may be that low levels of
residual translation initiation activity are sufficient for completion of the viral life cycle.

In cassava the eIF4E-family has five members and we previously demonstrate that all
five proteins interact with CBSV VPg by co-immunoprecipitation (Gomez *et al.*, 2019). We now
find, through pulldown experiments with purified proteins, that nCBP-1, nCBP-2, and eIF4E
can directly interact with CBSV VPg. This validates our approach for identifying nCBP-2
residues crucial for VPg interaction. In contrast, our pulldown experiments with VPg and
eIF(iso)4E-1 and -2 were uninformative. It may be that the cassava eIF(iso)4E proteins have
lower affinity for CBSV VPg than the nCBPs and eIF4E, precluding detection of interaction via
GST pulldown. This is supported by our earlier work using yeast two-hybrid to dissect CBSV
VPg and cassava eIF4E-family interactions; there, the combination of CBSV VPg with
eIF(iso)4Es produced weaker reporter activity than when VPg was combined with nCBPs
(Gomez *et al.*, 2019). A more granular understanding of differences in cassava eIF4E-family
protein affinity for CBSV VPg may be achieved using techniques like surface plasmon
resonance or isothermal titration calorimetry (Upadhyay et al., 2024).

To determine the contribution of each *eIF4E*-family gene to CBSV symptoms, and
therefore the biological relevance of interaction with VPg, we have used CRISPR/Cas9 to
generate single mutants of all five genes, as well as some double mutants, in cassava variety

TME419. Notably, the *ncbp* double mutant has near total reduction of storage root necrosis,
commensurate with strong reductions in viral titer. Of the two *nCBP* genes, *nCBP-2* appears
to be the main driver for the improved disease response as evidenced by the results of the
single *nCBP* mutants, and that *nCBP-2* is more highly expressed than *nCBP-1* (Gomez *et al.*,
2019). As noted in our previous work, the expression difference between *nCBP-2* and *nCBP-1*
1 is even more pronounced in storage roots and may be the reason for differential disease
response in roots and aerial tissues of *ncbp-2* and *ncbp-1 ncbp-2* mutants (Gomez *et al.*,
2019). The disease data are consistent between mutants in both 60444 (Gomez *et al.*, 2019)
and TME419. Beyond the *nCBP* clade, the *eIF(iso)4E* clade appears to have a small
contribution to disease as aerial symptom severity and storage root necrosis is slightly reduced
when both *eIF(iso)4E-1* and *eIF(iso)4E-2* are simultaneously knocked out. This finding also
argues against the *ncbp-1 ncbp-2* root disease phenotype being simply due to additively
knocking out family members. It remains to be seen if higher order mutant combinations would
have an even greater effect on reducing CBSD disease severity. It will also be important to
test the utility of *ncbp-1 ncbp-2* mediated resistance against the natural diversity of cassava
brown streak viruses in field trials.
A key consideration going forward is that any benefits conferred by mutations in eIF4E-
family proteins with respect to viral resistance may also be accompanied by growth and
developmental penalties (Gauffier *et al.*, 2016, Bastet *et al.*, 2017). Of the four residues
identified in *nCBP-2* mutants that appear to avoid this penalty, three (L51, H94, and A148) are
conserved within all cassava eIF4E-family proteins and the fourth (S138) is conserved within
three members. Targeted base editing of some or all of these residues across the entire eIF4E-
family may allow for improved CBSV response while maintaining essential protein translation
functions. Indeed, we demonstrate here that transferring the L51F mutation from *nCBP-2* into
Arabidopsis eIF(iso)4E, as an L28F mutation, results in loss of affinity for TuMV VPg. As base
editing approaches mature alongside improvements with gene targeting, combining these
approaches with loss-of-VPg interaction screens may facilitate isolation and implementation
of novel *eIF4E*-family resistance alleles. This may allow for countering the edge viruses hold
in the host-pathogen evolutionary arms race.
363
Materials and Methods
Plasmid vector construction
gRNAs for CRISPR/Cas9-based mutagenesis of cassava *eIF4E*-family genes were
selected using the CRISPOR webtool (Concordet & Haeussler, 2018). Cas-OFFinder was
then used to screen gRNAs for off-target activity against a TME419 reference based genome
assembly produced using data (SRX1393295) from Bredeson et al. and the AM560-2 v6.0
reference genome (Table S1) (Bae *et al.*, 2014; Bredeson *et al.*, 2016). gRNAs with off-targets

containing less than 2-mismatches were excluded from consideration. Finally, gRNAs were
cloned into the pDIRECT 21C CRISPR binary vector as previously described by (Čermák *et*
al., 2017).

Plant transformation, genotyping, and growth parameters

Cassava cultivar TME419 was transformed as previously described (Veley *et al.*,
2023). DNA was extracted from recovered transformants using the CTAB procedure. PCR
was then used to amplify 240 to 428 bp regions encompassing the gRNA binding sites. Purified
amplicons were sent to GENEWIZ for amplicon sequencing. Amplicon sequencing data was
analyzed using the CRIS.py python program (Connelly & Pruett-Miller, 2019). Plantlets of
transgenic lines were then maintained in tissue culture (Taylor *et al.*, 2012).

Viral challenge and disease phenotyping

Cassava disease challenges with CBSV isolate Naliendele, including inoculation,
aerial and below ground disease severity assessments, and qPCR-based quantification of
storage root viral titer, were carried out as described previously (Gomez *et al.*, 2019). For data
analysis, wild type disease severity was compared across all challenges and no significant
differences were found (Figure S1). Data for all challenges were then aggregated to increase
sample sizes for statistical analysis.

Co-immunoprecipitation and GST-pulldown experiments

Co-immunoprecipitation was performed as described previously (Gomez *et al.*, 2019).
For GST-pulldown experiments, cassava *eIF4E*-family genes in pENTR/D-TOPO, from
Gomez et al., were cloned into pDEST15 (Invitrogen) by Gateway recombination (Invitrogen).
pDEST15 encoding cassava *eIF4E*-family genes and pGEX-4T-1 (encoding unfused GST,
Cytiva) were transformed into BL21-AI E. coli (Invitrogen) and transformants were used to
start overnight seed cultures. Seed cultures were then used to inoculate 250 mL of LB broth.
Once inoculated cultures were grown to an OD600 of 0.4, IPTG (pGEX-4T-1) or arabinose
(pDEST15) were added to final concentrations of 0.5 mM or 0.2%, respectively, to induce
recombinant protein production. Induced cells were incubated for four hours at 28°C. Cells
were then pelleted and resuspended in 4 mL of lysis buffer (125 mM Tris, 150 mM NaCl, 1
mM DTT, 5 mM $MgCl_2$, 10 mM $CaCl_2$, pH 7.4) supplemented with 1X cOmplete protease
inhibitor (EDTA-free, Roche). Lysozyme (GoldBio) was added to a concentration of 1 mg/mL
and the mixture was allowed to incubate at room temperature for 15 minutes. 50 uL DNAse I
(Thermo) was then added to the mixture, followed by sonication on ice to ensure complete
lysis and elimination of DNA induced viscosity. The resulting cell lysate was centrifuged at
4°C, maximum speed, for 20 minutes. 3.5 mL of clarified lysate was transferred to a new tube

and EDTA was added to a final concentration of 1 mM. GST and GST-eIF4E proteins were
then purified from clarified lysate using glutathione magnetic agarose beads (Pierce)
according to manufacturer's protocol. Glutathione was removed from GST and GST-fusion
protein eluates by buffer exchange with Amicon Ultra centrifugal filters (Millipore). CBSV
6xHis-VPg-3xFLAG-6xHIS was produced and purified for pulldown experiments as described
in (Gomez *et al.*, 2019). GST pulldown was performed by mixing 3 ug of GST or GST-eIF4E
proteins with 3 ug of VPg and 10 uL of resuspended glutathione magnetic agarose beads
(Pierce) in buffer A (125 mM Tris, 150 mM NaCl, 10 mM DTT, 1 mM EDTA, pH 7.4)
supplemented with 1X cOmplete protease inhibitor (EDTA-free, Roche). Samples were
incubated at 4°C for one hour. Beads were then washed four times with buffer A and proteins
were eluted with Laemmli sample buffer for analysis by western blotting. Antibodies used in
this study are described in Table S2.

For comparison of VPg pulldown by nCBP-2 variants across experiments, nCBP-2 and
VPg eluate band intensities were first measured using Image Lab (Bio-Rad). Within an
experiment, eluate VPg band intensities were first adjusted to correct for differing pulldown
efficiencies of each nCBP-2 variant using the following equation where VPg_X is VPg band
intensity for sample X and GST_X is full length GST fusion protein band intensity for sample X:

$$VPg_{X,\ adjusted} = VPg_X\ adjusted\ for\ GST\ fusion\ protein\ pulldown\ efficiency = \frac{VPg_X}{\left(\frac{GST_X}{GST_{nCBP-2}}\right)}$$

Adjusted VPg intensities were then expressed relative to the VPg intensity of the wild-type
nCBP-2 sample for that experiment:

$$Normalized\ VPg_X = \frac{VPg_{X,\ adjusted}}{VPg_{nCBP-2,\ adjusted}} \times 100$$

Yeast two-hybrid assays

All yeast two-hybrid experiments were performed using the EGY48 strain harboring
the pSH18-34 *lacZ* reporter plasmid along with pJG4-5 and pEG202 bait and prey plasmids
(Golemis & Khazak, 1997). pJG4-5 *CBSV VPg* bait plasmid and pEG202 *nCBP-2* prey
plasmids were created by using In-Fusion cloning (Takara) to insert CBSV VPg or nCBP-2
coding sequences into pJG4-5 or pEG202 linearized with EcoRI and XhoI. For qualitative
yeast two-hybrid experiments, EGY48 pSH18-34 yeast transformed with bait and prey
plasmids were first selected for on SD –ura –trp –his media supplemented with 2% glucose
(Gietz & Schiestl, 2007). Transformants were then plated onto SD –ura –trp –his media
supplemented with 2% galactose, 1% raffinose, 1X BU salts, and X-gal. Blue/white screening
was performed three to four days later. Quantitative yeast two-hybrid was performed using a
spectrophotometric β-galactosidase activity assay (Amberg *et al.*, 2006).

To screen for nCBP-2 mutants that lose affinity for CBSV VPg, pEG202 *nCBP-2* was

used as template for mutagenic PCR (Genemorph II, Agilent). Primers immediately flanking

the second and penultimate codon of *nCBP-2*, preserving the start and stop codons during

mutagenic PCR, were used with the low mutation rate protocol from the Genmorph II kit

(Agilent). pEG202 empty vector was then digested with EcoRI and SacI and the fragment

containing the yeast *HIS* marker was gel purified (Qiagen). A third pEG202 fragment

containing the yeast 2μ origin of replication was produced by PCR with ends that overlap the

nCBP-2 mutant amplicons and pEG202 *HIS* marker fragment by 30 base pairs; co-

transformation of all three fragments into yeast result in assembly of an intact pEG202 *nCBP-*

2 plasmid through yeast recombinational cloning. Mutant *nCBP-2* amplicons, pEG202 *HIS*

fragment, and pEG202 2μ fragment were mixed in a 4:1:1 molar ratio and co-transformed into

EGY48 yeast harboring the pSH18-34 and pJG4-5 *CBSV VPg* plasmids. Transformed yeast

were directly plated onto SD –ura –trp –his media supplemented with 2% galactose, 1%

raffinose, 1X BU salts, and X-gal. Four days post transformation, white or light blue colonies

were restreaked onto a new plate for further visual assessment of reduced β-galactosidase

activity. The yeast were then checked by PCR for the presence of full length *nCBP-2* insert in

pEG202 using vector backbone specific primers. *nCBP-2* mutant alleles were then amplified

from insert positive yeast using Q5 Hot Start polymerase (NEB) and analyzed by Sanger

sequencing.

Yeast *eif4e* complementation assay

3xFLAG and nCBP-2 or mutant nCBP-2 coding sequences were cloned into pG1.1

linearized with NcoI and BamHI using In-Fusion (Takara) to produce pG1.1 *3xFLAG-nCBP-2*.

3xFLAG was amplified from pB7-HFC (Huang *et al.*, 2016) while *nCBP-2* variants were

amplified from pEG202 yeast two-hybrid plasmids. pG1.1 plasmids were then transformed into

the conditional yeast *eif4e* mutant strain T93C (*eif4e*::LEU2 *ura3 trp1 leu2* [pGAL1-eIF4E

URA3]) (Altmann *et al.*, 1989). Transformants were plated on SD –ura –trp dropout media

containing 2% galactose and 1% raffinose and allowed to grow for five to seven days. Colonies

of transformed yeast were then resuspended in sterile water and a ten-fold dilution series

ranging from OD600 = 10^{-1} through 10^{-4} was plated on SD –ura –trp dropout media containing

2% glucose to examine for rescue by pG1.1 encoded *eIF4E*-family gene variants. Plates were

imaged after 5-7 days.

Acknowledgements

We would like to thank the DDPSC PGF team for their hard work in caring for our

plants. We thank Claire Albin for her assistance with CBSV challenges and Nigel Taylor for

allowing us to use Taylor Lab greenhouse space. Yeast strain T93C, pG1.1 At_eIF4E1, and

pG1.1 empty vector were generously provided by Dr. Karen S. Browning. Z.D.L. would like to
thank Susan Lin, Wei-Yang Lin, Chun-Hwa Lin, and Jocelyn Lin for their love and support.
Z.D.L. was funded by USDA NIFA award 2019-67012-29626 and the Bill & Melinda Gates
Foudnation (OPP1125410, R.B.S., S.E.J., J.C.C.). E.J.D., Z.V.B., H.T., and E.H. were funded
by the DDPSC NSF-REU program, DBI-1659812, -2050394, -1659812, and -2244275,
respectively.

Competing Interests
None to declare.

Author Contributions
Conceptualization: Z.D.L.; Methodology: Z.D.L., J.C.C., R.S.B.; Investigation: Z.D.L.,
M.K.S., G.L.H., E.D.M., Z.V.B., K.B., H.T., E.H., G.J., K.B.G.; Visualization: Z.D.L.; Writing –
original draft: Z.D.L.; Writing – reviewing & editing: Z.D.L., K.B.G., J.C.C., R.S.B.

Data Availability
The source data for all results presented in this study are provided in accompanying
supplementary files.

References

Agricultural production statistics 2000–2022. **2023**. Rome, Italy: Food & Agriculture
Organization of the United Nations (FAO).

Altmann M, **Sonenberg N**, **Trachsel H**. **1989**. Translation in Saccharomyces cerevisiae:
initiation factor 4E-dependent cell-free system. *Molecular and cellular biology* **9**: 4467–4472.

Amberg DC, **Burke DJ**, **Strathern JN**. **2006**. Assay of β-galactosidase in yeast:
Permeabilized cell assay. *CSH protocols* **2006**: db.prot4158.

Ashby JA, **Stevenson CEM**, **Jarvis GE**, **Lawson DM**, **Maule AJ**. **2011**. Structure-based
mutational analysis of eIF4E in relation to sbm1 resistance to pea seed-borne mosaic virus
in pea. *PloS one* **6**: e15873.

Bae S, **Park J**, **Kim J-S**. **2014**. Cas-OFFinder: a fast and versatile algorithm that searches
for potential off-target sites of Cas9 RNA-guided endonucleases. *Bioinformatics (Oxford,*
England) **30**: 1473–1475.

Bastet A, **Lederer B**, **Giovinazzo N**, **Arnoux X**, **German-Retana S**, **Reinbold C**, **Brault V**,
Garcia D, **Djennane S**, **Gersch S**, *et al.* **2018**. Trans-species synthetic gene design allows

resistance pyramiding and broad-spectrum engineering of virus resistance in plants. *Plant biotechnology journal* **16**: 1569–1581.

Bastet A, **Robaglia C**, **Gallois J-L**. **2017**. EIF4E resistance: Natural variation should guide gene editing. *Trends in plant science* **22**: 411–419.

Bastet A, **Zafirov D**, **Giovinazzo N**, **Guyon-Debast A**, **Nogué F**, **Robaglia C**, **Gallois J-L**. **2019**. Mimicking natural polymorphism in eIF4E by CRISPR-Cas9 base editing is associated with resistance to potyviruses. *Plant biotechnology journal* **17**: 1736–1750.

Beyene G, **Chauhan RD**, **Ilyas M**, **Wagaba H**, **Fauquet CM**, **Miano D**, **Alicai T**, **Taylor NJ**. **2016**. A virus-derived stacked RNAi construct confers robust resistance to cassava brown streak disease. *Frontiers in plant science* **7**: 2052.

Bredeson JV, **Lyons JB**, **Prochnik SE**, **Wu GA**, **Ha CM**, **Edsinger-Gonzales E**, **Grimwood J**, **Schmutz J**, **Rabbi IY**, **Egesi C**, *et al.* **2016**. Sequencing wild and cultivated cassava and related species reveals extensive interspecific hybridization and genetic diversity. *Nature biotechnology* **34**: 562–570.

Browning KS, **Bailey-Serres J**. **2015**. Mechanism of cytoplasmic mRNA translation. *The Arabidopsis book* **13**: e0176.

Čermák T, **Curtin SJ**, **Gil-Humanes J**, **Čegan R**, **Kono TJY**, **Konečná E**, **Belanto JJ**, **Starker CG**, **Mathre JW**, **Greenstein RL**, *et al.* **2017**. A multipurpose toolkit to enable advanced genome engineering in plants. *The plant cell* **29**: 1196–1217.

Charron C, **Nicolaï M**, **Gallois J-L**, **Robaglia C**, **Moury B**, **Palloix A**, **Caranta C**. **2008**. Natural variation and functional analyses provide evidence for co-evolution between plant eIF4E and potyviral VPg. *The Plant journal: for cell and molecular biology* **54**: 56–68.

Concordet J-P, **Haeussler M**. **2018**. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic acids research* **46**: W242–W245.

Connelly JP, **Pruett-Miller SM**. **2019**. CRIS.Py: A versatile and high-throughput analysis program for CRISPR-based genome editing. *Scientific reports* **9**: 4194.

Coutinho de Oliveira L, **Volpon L**, **Rahardjo AK**, **Osborne MJ**, **Culjkovic-Kraljacic B**, **Trahan C**, **Oeffinger M**, **Kwok BH**, **Borden KLB**. **2019**. Structural studies of the eIF4E-VPg complex reveal a direct competition for capped RNA: Implications for translation. *Proceedings of the National Academy of Sciences of the United States of America* **116**: 24056–24065.

Dombrovsky A, **Reingold V**, **Antignus Y**. **2014**. Ipomovirus--an atypical genus in the family

Potyviridae transmitted by whiteflies. *Pest management science* **70**: 1553–1567.

Duprat A, **Caranta C**, **Revers F**, **Menand B**, **Browning KS**, **Robaglia C**. **2002**. The

Arabidopsis eukaryotic initiation factor (iso)4E is dispensable for plant growth but required

for susceptibility to potyviruses. *The Plant journal: for cell and molecular biology* **32**: 927–

934.

Gallois J-L, **Charron C**, **Sánchez F**, **Pagny G**, **Houvenaghel M-C**, **Moretti A**, **Ponz F**,

Revers F, **Caranta C**, **German-Retana S**. **2010**. Single amino acid changes in the turnip

mosaic virus viral genome-linked protein (VPg) confer virulence towards Arabidopsis

thaliana mutants knocked out for eukaryotic initiation factors eIF(iso)4E and eIF(iso)4G. *The*

Journal of general virology **91**: 288–293.

Gao L, **Luo J**, **Ding X**, **Wang T**, **Hu T**, **Song P**, **Zhai R**, **Zhang H**, **Zhang K**, **Li K**, *et al.*

2020. Soybean RNA interference lines silenced for eIF4E show broad potyvirus resistance.

Molecular plant pathology **21**: 303–317.

Gauffier C, **Lebaron C**, **Moretti A**, **Constant C**, **Moquet F**, **Bonnet G**, **Caranta C**, **Gallois**

J-L. **2016**. A TILLING approach to generate broad-spectrum resistance to potyviruses in

tomato is hampered by eIF4E gene redundancy. *The Plant journal: for cell and molecular*

biology **85**: 717–729.

German-Retana S, **Walter J**, **Doublet B**, **Roudet-Tavert G**, **Nicaise V**, **Lecampion C**,

Houvenaghel M-C, **Robaglia C**, **Michon T**, **Le Gall O**. **2008**. Mutational analysis of plant

cap-binding protein eIF4E reveals key amino acids involved in biochemical functions and

potyvirus infection. *Journal of virology* **82**: 7601–7612.

Gietz RD, **Schiestl RH**. **2007**. Frozen competent yeast cells that can be transformed with

high efficiency using the LiAc/SS carrier DNA/PEG method. *Nature protocols* **2**: 1–4.

Golemis EA, **Khazak V**. **1997**. Alternative yeast two-hybrid systems. The interaction trap

and interaction mating. *Methods in molecular biology (Clifton, N.J.)* **63**: 197–218.

Gomez MA, **Lin ZD**, **Moll T**, **Chauhan RD**, **Hayden L**, **Renninger K**, **Beyene G**, **Taylor NJ**,

Carrington JC, **Staskawicz BJ**, *et al.* **2019**. Simultaneous CRISPR/Cas9-mediated editing

of cassava eIF4E isoforms nCBP-1 and nCBP-2 reduces cassava brown streak disease

symptom severity and incidence. *Plant biotechnology journal* **17**: 421–434.

Grüner S, **Peter D**, **Weber R**, **Wohlbold L**, **Chung M-Y**, **Weichenrieder O**, **Valkov E**,

Igreja C, **Izaurralde E**. **2016**. The structures of eIF4E-eIF4G complexes reveal an extended
interface to regulate translation initiation. *Molecular cell* **64**: 467–479.

Grüner S, **Weber R**, **Peter D**, **Chung M-Y**, **Igreja C**, **Valkov E**, **Izaurralde E**. **2018**.
Structural motifs in eIF4G and 4E-BPs modulate their binding to eIF4E to regulate translation
initiation in yeast. *Nucleic acids research* **46**: 6893–6908.

Hart JP, **Griffiths PD**. **2013**. A series of eIF4E alleles at the Bc-3 locus are associated with
recessive resistance to Clover yellow vein virus in common bean. *Theoretical and applied*
genetics **126**: 2849–2863.

Huang H, **Alvarez S**, **Bindbeutel R**, **Shen Z**, **Naldrett MJ**, **Evans BS**, **Briggs SP**, **Hicks**
LM, **Kay SA**, **Nusinow DA**. **2016**. Identification of evening complex associated proteins in
Arabidopsis by affinity purification and mass spectrometry. *Molecular & cellular proteomics:*
MCP **15**: 201–217.

Kang B-C, **Yeam I**, **Frantz JD**, **Murphy JF**, **Jahn MM**. **2005**. The pvr1 locus in Capsicum
encodes a translation initiation factor eIF4E that interacts with Tobacco etch virus VPg. *The*
Plant journal: for cell and molecular biology **42**: 392–405.

Kelley LA, **Mezulis S**, **Yates CM**, **Wass MN**, **Sternberg MJE**. **2015**. The Phyre2 web portal
for protein modeling, prediction and analysis. *Nature protocols* **10**: 845–858.

Kim J, **Kang W-H**, **Hwang J**, **Yang H-B**, **Dosun K**, **Oh C-S**, **Kang B-C**. **2014**. Transgenic
Brassica rapa plants over-expressing eIF(iso)4E variants show broad-spectrum Turnip
mosaic virus (TuMV) resistance. *Molecular plant pathology* **15**: 615–626.

Lebaron C, **Rosado A**, **Sauvage C**, **Gauffier C**, **German-Retana S**, **Moury B**, **Gallois J-L**.
2016. A new eIF4E1 allele characterized by RNAseq data mining is associated with
resistance to potato virus Y in tomato albeit with a low durability. *The Journal of general*
virology **97**: 3063–3072.

Léonard S, **Plante D**, **Wittmann S**, **Daigneault N**, **Fortin MG**, **Laliberté JF**. **2000**. Complex
formation between potyvirus VPg and translation eukaryotic initiation factor 4E correlates
with virus infectivity. *Journal of virology* **74**: 7730–7737.

Li H, **Kondo H**, **Kühne T**, **Shirako Y**. **2016**. Barley yellow mosaic virus VPg is the
determinant protein for breaking eIF4E-mediated recessive resistance in barley plants.
Frontiers in plant science **7**: 1449.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R,
Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in
2024. Nucleic acids research **52**: W521–W525.

Miyoshi H, Suehiro N, Tomoo K, Muto S, Takahashi T, Tsukamoto T, Ohmori T,
Natsuaki T. 2006. Binding analyses for the interaction between plant virus genome-linked
protein (VPg) and plant translational initiation factors. *Biochimie* **88**: 329–340.

Naderpour M, Lund OS, Larsen R, Johansen E. 2010. Potyviral resistance derived from
cultivars of Phaseolus vulgaris carrying bc-3 is associated with the homozygotic presence of
a mutated eIF4E allele. *Molecular plant pathology* **11**: 255–263.

Odipio J, Ogwok E, Taylor NJ, Halsey M, Bua A, Fauquet CM, Alicai T. 2014. RNAi-
derived field resistance to Cassava brown streak disease persists across the vegetative
cropping cycle. *GM crops & food* **5**: 16–19.

Revers F, García JA. 2015. Molecular biology of potyviruses. *Advances in virus research*
92: 101–199.

Rey C, Vanderschuren H. 2017. Cassava mosaic and brown streak diseases: Current
perspectives and beyond. *Annual review of virology* **4**: 429–452.

Robaglia C, Caranta C. 2006. Translation initiation factors: a weak link in plant RNA virus
infection. *Trends in plant science* **11**: 40–45.

Robson F, Hird DL, Boa E. 2023. Cassava brown streak: A deadly virus on the move. *Plant*
pathology.

Sanfaçon H. 2015. Plant translation factors and virus resistance. *Viruses* **7**: 3392–3419.

Schaad MC, Anderberg RJ, Carrington JC. 2000. Strain-specific interaction of the tobacco
etch virus NIa protein with the translation initiation factor eIF4E in the yeast two-hybrid
system. *Virology* **273**: 300–306.

Sheat S, Winter S. 2023. Developing broad-spectrum resistance in cassava against viruses
causing the cassava mosaic and the cassava brown streak diseases. *Frontiers in plant*
science **14**: 1042701.

Takakura Y, Udagawa H, Shinjo A, Koga K. 2018. Mutation of a Nicotiana tabacum L.
eukaryotic translation-initiation factor gene reduces susceptibility to a resistance-breaking
strain of Potato virus Y. *Molecular plant pathology* **19**: 2124–2133.

Taylor N, **Gaitán-Solís E**, **Moll T**, **Trauterman B**, **Jones T**, **Pranjal A**, **Trembley C**,
Abernathy V, **Corbin D**, **Fauquet CM**. **2012**. A high-throughput platform for the production
and analysis of transgenic cassava (Manihot esculenta) plants. *Tropical plant biology* **5**:
127–139.

Tomlinson KR, **Bailey AM**, **Alicai T**, **Seal S**, **Foster GD**. **2018**. Cassava brown streak
disease: historical timeline, current knowledge and future prospects. *Molecular plant*
pathology **19**: 1282–1294.

Upadhyay V, Lucas A, Patrick C, Mallela KMG. 2024. Isothermal titration calorimetry and
surface plasmon resonance methods to probe protein-protein interactions. Methods (San
Diego, Calif.) **225**: 52–61.

Valli A, **García JA**, **López-Moya JJ**. **2015**. Potyviridae. *Encyclopedia of Life Sciences*: 1–
10.

Veley KM, **Elliott K**, **Jensen G**, **Zhong Z**, **Feng S**, **Yoder M**, **Gilbert KB**, **Berry JC**, **Lin Z-**
JD, **Ghoshal B**, *et al.* **2023**. Improving cassava bacterial blight resistance by editing the
epigenome. *Nature communications* **14**: 85.

Wagaba H, Beyene G, Trembley C, Alicai T, Fauquet CM, Taylor NJ. 2013. Efficient
transmission of cassava brown streak disease viral pathogens by chip bud grafting. BMC
research notes **6**: 516.

Wagaba H, **Beyene G**, **Aleu J**, **Odipio J**, **Okao-Okuja G**, **Chauhan RD**, **Munga T**, **Obiero**
H, **Halsey ME**, **Ilyas M**, *et al.* **2016**. Field level RNAi-mediated resistance to cassava brown
streak disease across multiple cropping cycles and diverse East African Agro-ecological
locations. *Frontiers in plant science* **7**: 2060.

Walker PJ, **Siddell SG**, **Lefkowitz EJ**, **Mushegian AR**, **Adriaenssens EM**, **Alfenas-**
Zerbini P, **Dempsey DM**, **Dutilh BE**, **García ML**, **Curtis Hendrickson R**, *et al.* **2022**.
Recent changes to virus taxonomy ratified by the International Committee on Taxonomy of
Viruses (2022). *Archives of virology* **167**: 2429–2440.

Wittmann S, **Chatel H**, **Fortin MG**, **Laliberté JF**. **1997**. Interaction of the viral protein
genome linked of turnip mosaic potyvirus with the translational eukaryotic initiation factor
(iso) 4E of Arabidopsis thaliana using the yeast two-hybrid system. *Virology* **234**: 84–92.

Yang X, **Li Y**, **Wang A**. **2021**. Research advances in potyviruses: From the laboratory bench
to the field. *Annual review of phytopathology* **59**: 1–29.

Yeam I, **Cavatorta JR**, **Ripoll DR**, **Kang B-C**, **Jahn MM**. **2007**. Functional dissection of
naturally occurring amino acid substitutions in eIF4E that confers recessive potyvirus
resistance in plants. *The plant cell* **19**: 2913–2928.

Zafirov D, **Giovinazzo N**, **Bastet A**, **Gallois J-L**. **2021**. When a knockout is an Achilles'
heel: Resistance to one potyvirus species triggers hypersusceptibility to another one in
Arabidopsis thaliana. *Molecular plant pathology* **22**: 334–347.

Supporting information

Fig. S1 Wild type aerial and storage root disease scores across all challenges
Fig. S2 gRNAs used in this study
Fig. S3 *eIF4E*-family mutant cassava plants
Fig. S4 Representative *eif(iso)4e-1 eif(iso)4e-2* double mutant and wild type TME419 six
weeks post transfer to soil
Fig. S5 Purified proteins for GST pulldown assays
Fig. S6 Alignment of ipomovirus and potyvirus VPg's show significant divergence in AA
sequences
Fig. S7 Alignment of cassava eIF4E-family proteins
Fig. S8 Quantitative yeast two-hybrid assay confirms loss-of-affinity screen results
Fig. S9 Western blot analysis of yeast strains used for yeast two-hybrid experiments
Table S1 TME419 gRNA off-target sites
Table S2 Antibodies used in this study
Table S3 Gene editing outcomes for TME419 eIF4E-family mutants
Table S4 All nCBP-2 loss-of-VPg-affinity mutants identified by yeast two-hybrid
689

Figure legends

Fig. 1 Endpoint CBSD storage root disease severity in cassava eIF4E-family mutants
a, b) Endpoint aerial disease (a) and storage root necrosis (b) observed in cassava eIF4E-
family single and double mutants after challenge with CBSV isolate Naliendele. Data
presented are an aggregate from all experimental replicates. Leaf and stem disease
symptoms are evaluated separately on a 0-5 scale and then added together for a total aerial
symptom score. Storage root necrosis severity was scored on a 1 to 5 scale, 1 being
asymptomatic and 5 corresponding to severe necrosis throughout all storage roots of a
plant. Mann-Whitney U test was used to detect statistical differences with an alpha = 0.05. *

≤ 0.05, ** ≤ 0.01, *** ≤ 0.001. Asterisks above boxplots denote differences from wild type,

while asterisks above brackets indicate differences between indicated genotypes.

c) qRT-PCR quantification of storage root CBSV viral levels from challenged mutants. CBSV

HAM1 was quantified relative to cassava PP2A to obtain HAM1 normalized relative

quantities (NRQs). HAM1 NRQs for each genotype are presented as relative to the mean of

wild type. Statistical differences were detected with the Mann-Whitney U test as described in

(a, b).

d) Representative storage root sections from wild type and ncbp-1 ncbp-2 #213 plants

challenged with CBSV. Scale bar denotes 1 cm.

Figure 2. Interaction of GST-cassava eIF4E family proteins and CBSV VPg *in vitro*

a, b) GST-pulldown assays were performed with CBSV VPg-6xHis-3xFLAG and GST or

GST-tagged cassava eIF4E-family proteins. Proteins were expressed in and purified from *E.*

coli. Human (Hs) eIF4E was also tested in (a).

c) Model for eIF4E-mediated resistance against potyvirids. Cartoon depicts potyvirid

genome-linked VPg protein bound to eIF4E-family protein in order to cause disease, while

mutations that disrupt this interaction results in resistance.

d) Coimmunoprecipitation of YFP or YFP- eIF4E-family proteins with TuMV 6xHis-VPg-

6xHIS-3xFLAG. VPg was expressed in and purified from *E. coli* and then mixed with extract

from *N. benthamiana* leaves expressing YFP or YFP-fusion proteins.

e) *Arabidopsis* eIF(iso)4E residues 92 through 150 aligned with homologous regions of

cassava eIF4E-family proteins. W92 and K150, marked red, are necessary for interaction

with TuMV VPg. * indicates conservation across all aligned proteins, : indicates conservation

of amino acids with strongly similar properties, and . indicates conservation of amino acids

with weakly similar properties.

f) GST-pulldown assay as in a) and b) with CBSV VPg incubated with either GST, GST-

nCBP-2, and GST-nCBP-2$^{W118L/N170E}$.

g) Comparison of nCBP-2 and nCBP-2$^{W118L/N170E}$ interaction with CBSV VPg by quantitative

yeast two-hybrid assay.

h) Western blot analysis of nCBP-2 and VPg expression in yeast from g).

Fig. 3 A yeast two-hybrid loss-of-affinity screen identifies nCBP-2 residues necessary

for VPg interaction

a) Workflow for using a yeast two-hybrid approach to identify nCBP-2 mutants that lose

affinity for CBSV Naliendele VPg. The resultant pEG202 prey vector encodes nCBP-2 fused

to the LexA DNA binding domain. pJG4-5 encodes CBSV VPg fused to the B42

transcriptional activation domain. The EGY48 yeast strain used harbors the pSH18-34

reporter plasmid containing *lacZ* driven by the LexA operator.

b) Ventral and dorsal views of cassava nCBP-2 structure produced using the Phyre2 server

(Kelley *et al.*, 2015). nCBP-2 structure of amino acids 49 through 226 was predicted using

wheat eIF4E (PDB ID: 2IDR) as a template. Mutated amino acid residues in nCBP-2 mutants

that lose VPg affinity and either rescue or fail to rescue *eif4e* yeast are highlighted in green

and pink, respectively. W118 and N170, homologous with TuMV VPg interacting residues of

Arabidopsis eIF(iso)4E, are highlighted in blue.

Fig. 4 T93C *eif4e* yeast complementation by *nCBP-2* mutants

a) Schematic for testing translation initiation activity of *nCBP-2* variants expressed from the

pG1.1 plasmid in T93C *eif4e* conditional mutant yeast.

b) Left panels: T93C yeast transformed with pG1.1 empty vector or pG1.1 encoding various

eIF4E-proteins grown on SD dropout media supplemented with galactose allowing for

conditional expression of yeast eIF4E from the pGAL1-eIF4E plasmid. Right panels: same

yeast strains from left panel grown on dropout media supplemented with glucose, preventing

complementation by the pGAL1-eIF4E plasmid.

c) Western blot analysis of 3xFLAG tagged nCBP-2 variant expression in yeast strains used

in a). Coomassie stained membrane is presented for assessing loading control.

Fig. 5 Effects of HPL motif mutation on nCBP-2 affinity for CBSV VPg and translation

initiation

a) Alignment of human eIF4E protein sequence with variants of cassava nCBP-2. The

conserved HPL motif is boxed in red and the L51F mutation in nCBP-2 is highlighted in cyan.

Asterisks denote conserved amino acid residues.

b) Quantitative yeast two-hybrid analysis of CBSV VPg interaction with nCBP-2 variants.

c) T93C *eif4e* yeast transformed with pG1.1 encoding *Arabidopsis eIF4E1* or cassava *nCBP-*

2 variants under the control of a constitutive promoter. nCBP-2 variants are N-terminally

tagged with 3xFLAG. The T93C yeast strain harbors a plasmid encoding galactose-inducible

and glucose-repressible yeast *eIF4E*.

d) Western blot analysis of nCBP-2 expression in yeast used in c). A segment of coomassie

stained membrane is presented for loading control.

Fig. 6 nCBP-2$^{K45\_L51del}$ and nCBP-2^{L51F} exhibit reduced affinity for CBSV VPg, *in vitro*

a) GST pulldown experiment assessing the ability of GST-nCBP-2 variants to bind CBSV

VPg. Full-length protein eluate band intensities are presented underneath the eluate western

blots and are expressed relative to the sample with the strongest band intensity.

b) Analysis of eluate VPg band intensity across three GST-pulldown experiments. Band

intensities are normalized to account for differing amounts of GST-fusion protein pulled down

and expressed relative to VPg intensity in the GST-nCBP-2 sample.

Fig. 7 Effects of mutating the HPL motif leucine in *Arabidopsis* eIF(iso)4E

a) Quantitative yeast two-hybrid analysis of TuMV VPg interaction with wild type *Arabidopsis*

eIF(iso)4E alongside W92L/K149E and L28F mutants.

b) T93C *eif4e* yeast complementation assay with wild-type and mutant *eIF4E*-family genes

from *Arabidopsis* and cassava.

c) Western blot analysis of eIF(iso)4E and nCBP-2 variants from cells used in (b).

Table 1. TME419 CRISPR mutant genotypes

Line #	CRISPR target	Gene/allele	INDEL	Protein nomenclature
6-95	*eIF4E*	*A*	-2	S21*fsX1
		B	–	"
172	*iso4E1+2*	*iso4E1 A*	-1	A11PfsX17
		iso4E1 B	-2	A11HfsX9
		iso4E2 A	-2	I7RfsX29
		iso4E2 B	1	E8RfsX36
92	*iso4E1+2*	*iso4E1 A*	-2	A11HfsX9
		iso4E1 B	-1	A11PfsX17
		iso4E2 A	-2	I7RfsX29
		iso4E2 B	1	E8RfsX36
10-9	*iso4E2*	*A*	1	E8RfsX36
		B	–	"
3-29	*iso4E1*	*A*	-1	A11PfsX17
		B	–	"
93	*nCBP1*	*A*	1	S23YfsX4
		B	-2	S23TfsX3
4-53	*nCBP2*	*A*	1	T23NfsX8
		B	–	"
213	*nCBP1+2*	*nCBP1 A*	1	S23YfsX4
		nCBP1 B	–	"
		nCBP2 A	-4	T22HfsX10
		nCBP2 B	-2	T23IfsX7
40	*nCBP1+2*	*nCBP1 A*	1	S23YfsX4
		nCBP1 B	1	P24TfsX3
		nCBP2 A	-21	K45_L51del
		nCBP2 B	-1	A46LfsX31

Table 2. Summary of nCBP-2 mutants found to lose affinity for CBSV VPg

No. loss-of-affinity mutants	Nucleotide substitutions per mutant (mean)	Nucleotide deletions per mutant (mean)	Unique alleles	No. unique AA substitutions	Frameshift mutants (unique/total)	Stop-gain mutants	Redundant AA substitution occurrences	AA residues with multiple occurrences of substitutions
51	1.37	0.15	46	48 [a]	7/8	7	7	8

[a] non-synonymous mutations downstream of frameshifting INDELs or stop-gain mutations are not counted

Fig. 1 Endpoint CBSD storage root disease severity in cassava *eIF4E*-family mutants

a, b) Endpoint aerial disease (a) and storage root necrosis (b) observed in cassava *eIF4E*-family single and double mutants after challenge with CBSV isolate Naliendele. Data presented are an aggregate from all experimental replicates. Leaf and stem disease symptoms are evaluated separately on a 0-5 scale and then added together for a total aerial symptom score. Storage root necrosis severity was scored on a 1 to 5 scale, 1 being asymptomatic and 5 corresponding to severe necrosis throughout all storage roots of a plant. Mann-Whitney U test was used to detect statistical differences with an alpha = 0.05. * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001. Asterisks above boxplots denote differences from wild type, while asterisks above brackets indicate differences between indicated genotypes.
c) qRT-PCR quantification of storage root CBSV viral levels from challenged mutants. CBSV *HAM1* was quantified relative to cassava *PP2A* to obtain *HAM1* normalized relative quantities (NRQs). *HAM1* NRQs for each genotype are presented as relative to the mean of wild type. Statistical differences were detected with the Mann-Whitney U test as described in (a, b).
d) Representative storage root sections from wild type and *ncbp-1 ncbp-2* #213 plants challenged with CBSV. Scale bar denotes 1 cm.

Figure 2. Interaction of GST-cassava eIF4E family proteins and CBSV VPg *in vitro*

a, b) GST-pulldown assays were performed with CBSV VPg-6xHis-3xFLAG and GST or GST-tagged cassava eIF4E-family proteins. Proteins were expressed in and purified from *E. coli*. Human (Hs) eIF4E was also tested in (a).

c) Model for eIF4E-mediated resistance against potyvirids. Cartoon depicts potyvirid genome-linked VPg protein bound to eIF4E-family protein in order to cause disease, while mutations that disrupt this interaction results in resistance.

d) Coimmunoprecipitation of YFP or YFP- eIF4E-family proteins with TuMV 6xHis-VPg-6xHIS-3xFLAG. VPg was expressed in and purified from *E. coli* and then mixed with extract from *N. benthamiana* leaves expressing YFP or YFP-fusion proteins.

e) *Arabidopsis* eIF(iso)4E residues 92 through 150 aligned with homologous regions of cassava eIF4E-family proteins. W92 and K150, marked red, are necessary for interaction with TuMV VPg. * indicates conservation across all aligned proteins, : indicates conservation of amino acids with strongly similar properties, and . indicates conservation of amino acids with weakly similar properties.

f) GST-pulldown assay as in a) and b) with CBSV VPg incubated with either GST, GST-nCBP-2, and GST-nCBP-2$^{W118L/N170E}$.

g) Comparison of nCBP-2 and nCBP-2$^{W118L/N170E}$ interaction with CBSV VPg by quantitative yeast two-hybrid assay.

h) Western blot analysis of nCBP-2 and VPg expression in yeast from g).

Fig. 3 A yeast two-hybrid loss-of-affinity screen identifies nCBP-2 residues necessary for VPg interaction

a) Workflow for using a yeast two-hybrid approach to identify nCBP-2 mutants that lose affinity for CBSV Naliendele VPg. The resultant pEG202 prey vector encodes nCBP-2 fused to the LexA DNA binding domain. pJG4-5 encodes CBSV VPg fused to the B42 transcriptional activation domain. The EGY48 yeast strain used harbors the pSH18-34 reporter plasmid containing *lacZ* driven by the LexA operator. b) Ventral and dorsal views of cassava nCBP-2 structure produced using the Phyre2 server (Kelley LA et al., 2015). nCBP-2 structure of amino acids 49 through 226 was predicted using wheat eIF4E (PDB ID: 2IDR) as a template. Mutated amino acid residues in nCBP-2 mutants that lose VPg affinity and either rescue or fail to rescue *eif4e* yeast are highlighted in green and pink, respectively. W118 and N170, homologous with TuMV VPg interacting residues of *Arabidopsis* eIF(iso)4E, are highlighted in blue.

Fig. 4 T93C *eif4e* yeast complementation by *nCBP-2* mutants

a) Schematic for testing translation initiation activity of *nCBP-2* variants expressed from the pG1.1 plasmid in T93C *eif4e* conditional mutant yeast.
b) Left panels: T93C yeast transformed with pG1.1 empty vector or pG1.1 encoding various eIF4E-proteins grown on SD dropout media supplemented with galactose allowing for conditional expression of yeast eIF4E from the pGAL1-eIF4E plasmid. Right panels: same yeast strains from left panel grown on dropout media supplemented with glucose, preventing complementation by the pGAL1-eIF4E plasmid.
c) Western blot analysis of 3xFLAG tagged nCBP-2 variant expression in yeast strains used in a). Coomassie stained membrane is presented for assessing loading control.

Fig. 5 Effects of HPL motif mutation on nCBP-2 affinity for CBSV VPg and translation initiation

a) Alignment of human eIF4E protein sequence with variants of cassava nCBP-2. The conserved HPL motif is boxed in red and the L51F mutation in nCBP-2 is highlighted in cyan. Asterisks denote conserved amino acid residues.

b) Quantitative yeast two-hybrid analysis of CBSV VPg interaction with nCBP-2 variants.

c) T93C *eif4e* yeast transformed with pG1.1 encoding *Arabidopsis eIF4E1* or cassava *nCBP-2* variants under the control of a constitutive promoter. nCBP-2 variants are N-terminally tagged with 3xFLAG. The T93C yeast strain harbors a plasmid encoding galactose-inducible and glucose-repressible yeast *eIF4E*.

d) Western blot analysis of nCBP-2 expression in yeast used in c). A segment of coomassie stained membrane is presented for loading control.

Fig. 6 nCBP-2$^{K45\_L51del}$ and nCBP-2^{L51F} exhibit reduced affinity for CBSV VPg, *in vitro*

a) GST pulldown experiment assessing the ability of GST-nCBP-2 variants to bind CBSV VPg. Full-length protein eluate band intensities are presented underneath the eluate western blots and are expressed relative to the sample with the strongest band intensity.
b) Analysis of eluate VPg band intensity across three GST-pulldown experiments. Band intensities are normalized to account for differing amounts of GST-fusion protein pulled down and expressed relative to VPg intensity in the GST-nCBP-2 sample.

Fig. 7 Effects of mutating the HPL motif leucine in *Arabidopsis* eIF(iso)4E

a) Quantitative yeast two-hybrid analysis of TuMV VPg interaction with wild type *Arabidopsis* eIF(iso)4E alongside W92L/K149E and L28F mutants.
b) T93C *eif4e* yeast complementation assay with wild-type and mutant *eIF4E*-family genes from *Arabidopsis* and cassava.
c) Western blot analysis of eIF(iso)4E and nCBP-2 variants from cells used in (b).