Physical Network Constraints Define the
Lognormal Architecture of the Brain's Connectome

Ben Piazza[1,*], Dániel L. Barabási[2,*], André Ferreira Castro[3],
Giulia Menichetti[1,4,5,], Albert-László Barabási[1,6,✉]

[1] Network Science Institute and Department of Physics, Northeastern University, Boston, MA, USA
[2] Eric and Wendy Schmidt Center, Broad Institute of MIT and Harvard, Cambridge, USA, 02142
[3] Technical University Munich, School of Life Sciences, Munich.
[4] Channing Division of Network Medicine and Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA
[5] Harvard Data Science Initiative, Harvard University, 114 Western Avenue, 02134, MA, USA
[6] Department of Network and Data Science, Central European University, Budapest, Hungary
* These authors contributed equally
✉ Correspondence should be addressed to: barabasi@gmail.com

Abstract

The brain has long been conceptualized as a network of neurons connected by synapses. However, attempts to describe the connectome using established network science models have yielded conflicting outcomes, leaving the architecture of neural networks unresolved. Here, by performing a comparative analysis of eight experimentally mapped connectomes, we find that their degree distributions cannot be captured by the well-established random or scale-free models. Instead, the node degrees and strengths are well approximated by lognormal distributions, although these lack a mechanistic explanation in the context of the brain. By acknowledging the physical network nature of the brain, we show that neuron size is governed by a multiplicative process, which allows us to analytically derive the lognormal nature of the neuron length distribution. Our framework not only predicts the degree and strength distributions across each of the eight connectomes, but also yields a series of novel and empirically falsifiable relationships between different neuron characteristics. The resulting multiplicative network represents a novel architecture for network science, whose distinctive quantitative features bridge critical gaps between neural structure and function, with implications for brain dynamics, robustness, and synchronization.

Since Ramón y Cajal revealed the first images of neurons in 1888, the brain has been conceptualized as a weighted network, whose nodes represent individual neurons and weighted links encode the number of synapses connecting them (Fig. 1a). However, it took more than a century to map this network, starting with the map of the *C. elegans* nematode four decades ago[1], and culminating with the recent reconstruction of the full brain of the fruit fly[2-4]. While an ultimate understanding of brain function also requires data on neuronal activity, a quantitative understanding of the connectome's wiring principles is crucial for detecting alterations underlying brain disorders, from autism to schizophrenia. Indeed, advances in network science have demonstrated how the structural features define multiple dynamical characteristics of a network, from its robustness to node and link failures[5,6], to its resilience[7], information spread[8], controllability[9,10], and synchronization[11]. These advances have inspired multiple attempts to elucidate the structural characteristics of neural networks[12], with conflicting outcomes: while the wiring of the *C. elegans* connectome was reported to be consistent with a randomly growing network[13], co-activity networks extracted from human fMRI data have repeatedly demonstrated a scale-free organization[14-17], two irreconcilable architectures. The recent proliferation of connectome data has also yielded puzzling outcomes: although the degree distributions do not exhibit a clear power-law scaling[18-20], the observed prevalence of highly connected neurons[19-22] remains a hallmark typically associated with scale-free networks[23].

To resolve these conflicting outcomes, here we compare data from eight connectomes spanning five organisms, arriving at the conclusion that the traditional classes of models previously explored by network science are unable to account for the observed structural features of the connectome. Specifically, we show that the degree and the strength

distributions of the connectome are inconsistent with both random[24] and scale-free[23] models, but both quantities are well approximated by a lognormal distribution. However, lognormals are the result of multiplicative processes whose origin remains unknown in the development of the brain. We then show that the lognormal architecture is rooted in the physical nature of the neurons: in contrast with the assumption of a separation between nodes and links (Fig. 1a), valid for the much studied social and technological networks, the brain consists of spatially extended neurites that simultaneously serve as both nodes and links (Fig. 1b). Consequently, the large-scale architecture of the connectome is driven primarily by the length of its individual neurons, which determine both a neuron's synapse number (node strength) and its degree. We then use a Galton-Watson branching process to analytically derive the neuronal length distribution, not only predicting the degree and the strength distributions, but obtaining falsifiable predictions that can account for the empirical data across all eight connectomes, as well as the unique features of individual neural classes. The observed multiplicative network with a lognormal degree distribution defines a novel architectural class in network science, integrating features of both random and scale-free networks and redefining canonical properties such as robustness and synchronization.

Datasets

We integrated data from eight connectome mapping efforts that offer synaptic connectivity at the resolution of single neurons for five organisms (SI 1): (1) the complete *C. elegans* connectome[1,25] (*C. elegans*), (2) a neural-level map of the entire brain of the *Drosophila melanogaster* larva's first instar phase[21] (Fly Larva), and two efforts to map the adult *Drosophila*'s brain, (3) one reconstructing a single hemisphere (Hemi-Brain)[26] and (4) another capturing neurons from both hemispheres[2–4] (FlyWire), (5) a map of the *Drosophila* adult's ventral nerve cord[27] (MANC), (6) the brainstem of the larval zebrafish[28] (Zebrafish), (7) a millimeter-scale volume of mouse primary visual cortex spanning from layer 1 to white matter[29,30] (Mouse), and (8) a human surgical sample from the temporal lobe of the cerebral cortex[31] (Human). While the size of the profiled brains spans eight orders of magnitude, from 302 neurons in the roundworm to an estimated 10^{10} human neurons[32], most of the available connectomes represent incomplete samples: although they capture all 302 neurons in *C. elegans*[1,25] and all 129,278 neurons in the adult Fly brain[2–4], they map only 2,587 neurons of an estimated total of 100,000 in the larval zebrafish[28], 52,059 out of 70 million neurons in the mouse brain[30], and 46,378 out of 86 billion neurons in the human brain[31] (Table S1).

We harmonize all eight connectome maps by correcting formatting differences and rescaling spatial measurements based on imaging resolution and sample thickness (SI 1.3), enabling direct comparisons of neuron size and connectivity across datasets. For incomplete connectomes we distinguish completely mapped neurons from those that are only partially mapped (SI 1.3-1.5). We limit our study to complete neurons, while utilizing all reported synapses that complete neurons make (SI 1).

The Degree Distribution of the Connectome is Neither Random nor Scale-Free

The degree distribution, *P(k)*, representing the probability that a neuron synapses with *k* other neurons, is a fundamental network property[33], as we must know (and control for) *P(k)* to interpret any higher-order features, from network robustness[5], to community structure[34,35] or motifs[36]. While the functional form of *P(k)* is determined by multiple diverse local processes[37,38], real networks tend to belong to one of two broad classes (SI 2.10): (1) Networks with a bounded *P(k)*, such as the Poisson degree distribution observed for random networks[24] or a simple exponential emerging in randomly growing networks (Fig. 1c). The exponentially decaying tail[37] of such bounded networks limits the variance in node sizes, excluding hubs; (2) Networks observed in many real systems, from protein interactions to the WWW, are scale-free, with a *P(k)* whose tail is often approximated by a power law[23] (Fig. 1e), allowing for high degree nodes, including hubs.

To elucidate the connectome's architecture, we began by measuring the degree distribution for each dataset. We find that each connectome is characterized by a unique *P(k)* distribution with rather different average (peak position) and different width (Fig. 2a). Yet, all reconstructions share a unimodal shape on a log-linear plot (*P(k)* vs *log k*), leading to the hypothesis that apart from dataset-dependent logarithmic average μ_k = *<log k>* and logarithmic width σ_k (standard deviation of *P(log k)*) (Fig. 2a), the degrees may be drawn from the same distribution. To test this hypothesis, we employed a standard procedure of statistical physics, plotting $P(\bar{k})$, where $\bar{k} = exp(\frac{log\,k - \mu_k}{\sigma_k})$, a linear transformation in the log space that corrects for differences in μ_k and $\sigma_{k'}$ but leaves the functional form of *P(k)* unchanged. We find that the

rescaled degrees from each organism collapse onto a single $P(\bar{k})$ distribution (Fig. 2b), supporting the hypothesis that, irrespective of the organism, the degrees are extracted from a single universal distribution.

The connectome is best described as a weighted network, as a link between neurons i and j is formally an integer s_{ij}, representing the number of distinct synapses connecting them.

In weighted networks, each node i is characterized by the node strength, $S_i = \sum_j s_{ij}$, representing the total number of synapses neuron i has. We find $P(S)$ to be again characterized by a symmetric and unimodal peak in a log-linear scale with different average and width (Fig. 2b). Plotting $P(\bar{S})$, where $\bar{S} = exp(\frac{log\, S - \mu_S}{\sigma_S})$ (Fig. 2d), we again find that the rescaled distributions collapse onto a common $P(\bar{S})$ (Fig. 2d).

Taken together, the existence of a common $P(\bar{k})$ and $P(\bar{S})$ is unexpected, considering that the eight datasets encompass five different species with varying levels of mapping completeness, diverse cell types, distinct synaptic connections, and a range of neurotransmitters. On one hand, our findings suggest the existence of an organism-independent developmental mechanism that shapes the connectome's structural features. Equally importantly, the observed $P(k)$ and $P(S)$ are incompatible with the canonical degree distributions in network science (Fig. 1f,h). Indeed, the empirically observed $P(k)$ cannot be fit by a Poisson distribution (random static network), a simple exponential $e^{-\alpha k}$ (randomly growing networks), or by a power law $P(k) \sim k^{-\gamma}$ (scale-free network) (Fig. 1g,h,i). Kolmogorov-Smirnoff tests (Fig. 2d,h) indicate that all these canonical distributions have large

KS statistics (see SI 2.5-2.8 for additional statistical measures). Instead, the clear unimodal and symmetric nature of the data in a log-linear plot (Fig. 2b,e) suggests that both *P(k)* and *P(S)* follow a lognormal distribution. Consistent with this hypothesis, we find that a fit to a lognormal has systematically low KS statistics across most datasets (Fig. 2b,f, see also SI 2.5-2.8 for additional statistical evidence).

While lognormal distributions provide reasonable fits to correlations in neural and brain activity[39], as well as synapse sizes[40–42], their significance to network structure remains less understood. The puzzle arises from how networks grow: the incremental addition of nodes yields a slow (polynomial or logarithmic) growth in degrees that usually gives rise to either a power law or an exponential *P(k)*. While certain models of socio-economic systems allow for a lognormal *P(k)*, they require rather specific preferential attachment functions[43] or additional parameter families[44] (SI 2.10). By contrast, lognormal distributions naturally arise from multiplicative processes[45], that yield exponential dynamics — fundamentally different from the incremental, node-by-node growth observed in real networks, including the brain. Consequently, the origin of lognormal distributions in the connectome remains unknown (SI 2.10). At the same time, while the empirical evidence presented in Figs. 1-2 is evocative, it is insufficient to conclusively establish the lognormal nature of the connectome[46]. A definitive assessment requires us to unveil the nature of the multiplicative process that can mechanistically explain the observed lognormal architecture.

The Connectome is a Physical Network

The brain is often conceptualized as a hub-and-spoke network, where nodes represent the somas of each neuron, connected by directed and weighted links that represent the number of synaptic connections between dendrite and axon pairs (Fig. 1a). However, in the connectome we lack a clear separation between the nodes and the links, as links (synapses) require spatial proximity between the axons and the dendrites (Fig. 1b). In other words, the brain is a physical network[47], hence its description demands a modeling framework that acknowledges the spatially extended physical nature of each neuron[41,48,49]. In physical networks, the longer a neuron is, the more chances it has to reach other neurons (proximity)[2,50], and to synapse with them. Formally, this suggests that

$$S_i \sim \rho_i L_i , \qquad\qquad (1)$$

where L_i is the neuron's total length and ρ_i is the local synapse density[51] of neuron *i*. To test the validity of hypothesis (1), we plotted S_i vs L_i for each neuron (Fig. 3a), finding that while for small L_i we observe deviations from (1), in each of the eight datasets the scaling holds for large L_i neurons, an asymptotic scaling sufficient for the subsequent theory.

Given (1), a lognormal form for *P(S)* is only possible if the neuronal lengths also follow a lognormal distribution. This prompted us to measure the total length, L_i, of each fully reconstructed neuron's skeleton (Fig. 1b, SI 3.1), finding a unimodal and symmetric *P(L)*

distribution on a log-linear plot (Fig. 2g). Plotting P$(\bar{L})$ vs $\bar{L} = exp(\frac{log\ L - \mu_L}{\sigma_L})$, we find that each

organism's length distribution collapses onto the same functional form (Fig. 2h).

Equation (1) also requires us to account for variations in synaptic densities, ρ_i. We

therefore determined the *local synapse density,* ρ_i of each fully reconstructed neuron, making

sure that we limit the measurement only to the regions of the neuron that contain synapses (SI

3.2). We find that *P(ρ)* for the different connectomes again unimodal (Fig. 2j) and under

$\bar{\rho} = exp(\frac{log\ \rho - \mu_\rho}{\sigma_\rho})$ rescaling collapses onto a single P$(\bar{\rho})$ curve (Fig. 2k). Finally, we fitted

multiple candidate distributions to *P(L)* and *P(ρ)*, finding that the lognormal form offers the

lowest KS parameter across all datasets (Fig. 2i,l and SI 2.7 for additional statistical tests).

Falsifying the Lognormal Connectome

Equation (1) offers falsifiable predictions that go beyond the scaling law of Fig. 3a.

Indeed, let us assume that consistent with empirical data, *S* and *L* follow the lognormal

distribution

$$P(x) = LN(\mu, \sigma) = \frac{1}{x\sigma\sqrt{2\pi}}e^{-\frac{1}{2}(\frac{log\ x - \mu}{\sigma})^2}, \tag{2}$$

where $\mu = <log(x)>$ is the scale parameter, and $\sigma > 0$ is the shape parameter[52],

corresponding to the standard deviation of $P(log(x))$. However, if L_i and ρ_i both follow

lognormal distributions with parameters $P(L) = LN(\mu_L, \sigma_L)$ and $P(\rho) = LN(\mu_\rho, \sigma_\rho)$, then one

can prove that their product (1) must also follow a lognormal distribution $P(S) = LN(\mu_S, \sigma_S)$,

with modified parameters (SI 2.1),

$$\mu_S = \mu_L + \mu_\rho, \tag{3}$$

$$\sigma_S = (\sigma_L^2 + \sigma_\rho^2 + 2r_{L,\rho}\sigma_L\sigma_\rho)^{1/2}. \tag{4}$$

Equations (3) and (4) offer empirically falsifiable predictions, indicating that $P(S)$ is uniquely determined by $P(L)$ and $P(\rho)$. To test their validity, we measured the (μ, σ) parameters for each connectome (see SI Table 2-3), finding a remarkable agreement between the empirically observed (μ_S, σ_S), and the values predicted by (3) (Fig. 3b) and (4) (Fig. 3c).

The final test of Eqs. (1)-(3) is offered by Fig. 3d-f, where we plot for three connectomes the empirically observed $P(S)$ distribution (symbols), along with a lognormal function whose parameters (μ_S, σ_S) are predicted by Eqs. (3) and (4) (Fig. 3d-f, continuous line), finding that the predicted $P(S)$ offers a good approximation for the empirically observed $P(S)$ (Fig. 3d-f, symbols).

Degree Distribution

As the number of synapses between neurons i and j is driven by the number of total synapses each neuron has (S_i and S_j), it is tempting to write $k_i \sim S_i \sim \rho_i L_i$. However, as we have multiple synapses between two neurons ($s_{ij} \geq 1$), we need to formalize the connectome as a weighted network, where k_i depends on S_i. We find that the degrees are well approximated by

$$k_i = \alpha S_i^\beta, \tag{5}$$

where $\beta < 1$ (Fig. 3g), a sublinear dependence frequently observed in weighted networks[53,54].

Yet, if S follows a lognormal distribution with parameters $P(S) = LN(\mu_S, \sigma_S)$, one can prove that $P(k)$ must also follow a lognormal distribution with the modified parameters

$$\mu_k = \ln a + \beta\mu_S, \qquad (6)$$

$$\sigma_k^2 = \beta^2\sigma_S^2. \qquad (7)$$

To falsify the predictions (6) and (7), we plot in Fig. 3h,i the empirically observed (μ_k, σ_k) against the values predicted by (6) and (7), finding an agreement for most datasets. As a final test, we plot the empirically observed $P(k)$ (symbols), together with the lognormal function whose parameters (μ_k, σ_k) are predicted by (7) and (8), confirming that the analytically predicted $P(k)$ offers an excellent approximation to the empirically observed $P(k)$ (Fig. 3j-l).

In summary, Figs. 2 and 3 indicate that $P(S)$, $P(k)$, $P(\rho)$, and $P(L)$ are stereotyped measures, displaying a remarkable consistency across the different connectomes. Further, Eqs. (3), (4), (6), and (7), predict tight and empirically validated relationships between the independently measured neural characteristics of S, k, ρ and L, offering direct and falsifiable evidence for the validity of lognormals. We lack, however, a theoretical argument for the mechanistic origin of the lognormal distribution. To identify the origin of the observed distributions, we next focus on the neuronal length distribution $P(L)$, given that according to Eqs. (1) and (5), $P(L)$ determines $P(S)$ and $P(k)$.

The Mechanistic Origin of the Lognormal Connectome

Neuronal growth is driven by complex genetically encoded developmental mechanisms[55], and is guided by multiple families of signaling and adhesion molecules[56–60]. Yet, the stereotypical form of *P(L)* suggests that the mechanisms governing neuronal length are robust to genetic variability, hinting at the existence of a common mechanistic origin. Morphologically, every neuron is an incomplete tree (Fig. 1b and 4c, See also SI 4.1), and under rather general conditions the growth of such trees is formally described by a Galton-Watson branching process, introduced in 1874 to describe the extinction of social groups[61] (SI 4.2) The process generates a random binary tree with 2^n possible branches after n generations (Fig. 4a), with a total length of $L_n=2^n<\ell>$, where $<\ell>$ is the average length of the individual branches. Although the growth if each branch is governed by chemical signals, cell-type-specific processes, and partner feedback[62], a mathematically robust description of neuronal branching can encapsulate these details into a single survival probability p_s that represents the likelihood of a branches' continued growth (Fig. 4a). In this case, the total length of a neuron after n generations follows (SI 4.4)

$$L_n = 2^n<\ell> \prod_{j=1}^{n} p_s^j \qquad (8)$$

where we acknowledge that the survival probability can vary as the neuron grows, hence we assign a different family of distributions $\{p_s^j\}$, with parameter families $\{\mu_p^j, \sigma_p^j\}$ to each layer j (Fig. 4a). As (8) describes a multiplicative process (Fig. 4a,b), the multiplicative central limit theorem allows us to analytically derive the tree size distribution $P(L_n/\ell)$, finding that each tree will be sampled from a lognormal length distribution (2) with parameters (SI 4.4),

$$\mu_L = n\left(\mu_p + \log 2\right) \tag{9}$$

$$\sigma_L^2 = n\sigma_p^2. \tag{10}$$

where n represents the layer number at which the tree terminates, still to be determined. In others words, the size distribution of the neural trees is determined by the survival probability distribution defined by $\{\mu_p^j, \sigma_p^j\}$, which govern the growth of the individual branches.

Predictions (9) and (10) are directly falsifiable, as for each neuron we can measure the layer-dependent survival probability $p_s^j = \frac{1}{2}\frac{X_j}{X_{j-1}}$, where X_j is the observed number of branches in layer j (Fig. 4c,d), and use maximum-likelihood estimation to extract $\{p_s^j\}$ for each neuron i from the geometry of its skeleton. We performed this procedure for all FlyWire neurons, confirming that p_s^j indeed depends on the layer number, being high near the soma (small j), but rapidly converging to $p_s^j = 1/2$ for large j (Fig. 4e). Ultimately Eqs. (8), (9) and (10) close the theoretical framework necessary to understand the neural-level architecture of the connectome, predicting that the multiplicative process governing the growth of the individual neuronal trees is responsible for the emergence of the lognormal $P(L)$, which not only drives $P(S)$ and $P(k)$, but allows us to explicitly derive its value for the functional form of the branch survival probabilities p_s^j, as we show next.

Predicting the characteristic of individual neural classes

The more than 8,000 subtypes of neurons identified in the FlyWire dataset[2] illustrate the exceptional diversity of individual neurons (SI 1.3), prompting us to ask if the theoretical framework developed above is capable of capturing these morphological differences. Differences between classes are well described by each class's unique *P(S)* and *P(L)* distributions, characterizing each of the 29 distinct neural classes identified for the adult fly (Fig. 5a,e, see also Fig. S4 for *P(k)* and *P(ρ)*). First we verified that upon rescaling, the *P(S)* and *P(L)* distributions characterizing the different neural classes once again collapse into a single stereotyped distribution (Fig. 5b,f), best approximated by a lognormal (Fig. 5c,g). At the same time, we find that each neuronal class is characterized by a different survival probability p_s^j function, underpinning the different developmental processes governing them (Fig. 5d). To test the applicability of our theory to specific classes, we next used the empirically measured class-dependent survival probability functions p_s^j (Fig. 5d) as input to Eqs. (9) and (10), allowing us to predict the μ_L and σ_L parameters that govern the tree size distribution $P(L/\ell)$ for each neuron class. In Fig. 5d, we plot μ_L predicted by Eq. (9) against the empirically observed neuronal length $\mu_L = <log\ (L/\ell)>$ for each of the 29 classes. The slope of the observed linear dependence allows us to empirically determine the characteristic layer number *n* in Eq. (8) at which the neurons cease their growth, finding to be $n \simeq 20$ for all classes (Fig. 5h). Most importantly, the agreement between the predicted and measured $\mu_L = <log\ (L/\ell)>$ not only confirms the predictive power of Eqs. (9) and (10), but also indicates that our theoretical

framework, developed for the whole connectome, captures the characteristics of individual neuronal classes as well.

Multiplicative Networks and Network Science

The observed lognormal degree distribution points to a network architecture in which degrees emerge through multiplicative rather than additive processes. Such *multiplicative networks* are unexplored in network science, exhibiting properties distinct from both random and scale-free networks. Consider, for example, the expected maximum degree, k_{max}, which in a random network grows as *log N*, where *N* is the network size. Hence, for both *C. elegans* ($N \simeq 300$) and the human brain ($N \simeq 10^{11}$), such a log-based growth predicts an implausibly modest and comparable maximum degree (Fig. 1i). By contrast, in a hypothetical scale-free human brain the largest hub scales more steeply (e.g. $k_{max} \sim N^{\frac{1}{\gamma-1}}$), which can exceed $k_{max} \sim 10^7$ for humans, an unrealistically large number of distinct connections per neuron. In a *multiplicative* network, however, the maximum degree follows (Fig 1i)

$$k_{max} \sim exp(\mu_k + \sigma_k \sqrt{2 log N}), \tag{11}$$

predicting a reasonable $k_{max} \simeq 10^3$ for $N \simeq 10^{11}$, and indicating that a multiplicative architecture can accommodate the existence of hub neurons while keeping their degree within biologically realistic limits.

The novel features of multiplicative networks are reflected by the relative degree heterogeneity κ = *<k²>/<k>*, which governs many network-based processes, from robustness[63] to resilience[7] and synchronization[11,64]. In a random network κ = *<k> + 1,* which is independent of

the network size. By contrast, for a scale-free network κ grows polynomially with N, following κ ~ $N^{(3-\gamma)/(\gamma-1)}$. In a lognormal network, however, $\kappa = <k>e^{3\sigma_k^2/2}$. Notably, κ does not depend on N, but its exponential dependence on σ_k can make it arbitrarily large. In other words, while a scale-free network's robustness and synchronizability vary significantly with size, the corresponding properties of a multiplicative network depend only on changes in σ_k— and hence on the *developmental processes* shaping the variability of degrees.

Finally, numerous empirical studies report a "rich-club" organization in brain networks—both in structural connectomes[65] and fMRI-based functional maps[66,67]. As we show in SI 4.7, such rich club behavior emerges naturally from the sublinear scaling (6), which predicts $S_i \sim k_i^{1/\beta}$, implying that high-degree nodes carry disproportionately large weights.

Discussion and Conclusions

The connectome has long posed a quantitative challenge for network science, often yielding conflicting conclusions about its underlying architecture. Here, we reconcile these discrepancies by showing that the physical nature of neurons imposes features absent in previously studied non-physical social and technological networks. Accounting for this physicality enables us to analytically derive the empirically observed *P(k), P(S)*, and *P(L)* distributions, each following a lognormal distribution. Most importantly, we provide a directly testable mechanism for the emergence of these distributions, along with empirically falsifiable equations (4), (5), (7), (8), (10), and (11), that reveal previously unknown non-intuitive relationships among measurable neuronal characteristics, that can only be valid if the underlying distributions follow a

lognormal. Finally, we show that the lognormal connectome leads to a *multiplicative* network architecture, whose robustness and synchronizability are shaped not by the network's size but by the intrinsic heterogeneity of neurons rooted in developmental processes.

We expect the theoretical framework introduced here to be enriched by the inclusion of additional developmental processes that govern neuron and synapse formation, thereby deepening our understanding of multiplicative networks and leading to accurate generative models. As an initial step in this direction, in SI 2.10 we discuss the Connectome Configuration Model, which tests our central hypothesis that the brain's architecture is driven by the physical properties of individual neurons. While the model intentionally simplifies many aspects of neuronal structure and connectivity, it successfully reproduces several key empirical observations, helping identify features that require more precise treatment of physicality. Indeed, although two neurons must be proximal to form a synapse, they also need to excrete the correct combination of surface membrane proteins to initiate synapse formation[68]. Consequently, we need concurrent advances in understanding the packing of physical networks[19,69], and the combinatorial expression of transcription factors[59,60], to construct a biologically accurate generative model.

Our study has several limitations that can only be resolved by more complete datasets. First, for mammalian brains, we currently have access to only a small fraction of the connectome—0.07% of estimated neurons for mice and 0.00005% for humans. While we mitigated this incompleteness by focusing on fully reconstructed neurons, the available maps systematically omit neurons whose size exceeds the mapped region. Accordingly, we examined

the length distributions obtained from injection-based techniques (SI 1.5) and found that they too fit a lognormal pattern, albeit with higher μ_L values than those in Fig. 2. In other words, the current data gaps underestimate the (μ_k, σ_k) parameters, along with the heterogeneity parameter κ, and data incompleteness is also responsible for the observed log-skewness in *P(L)*, *P(S)* and *P(k)* (SI 3.5).

Finally, understanding the connectome's statistical properties is fundamental not only for elucidating its structural organization, but also for advancing models of brain dynamics. Insights into these architectural principles are crucial for linking genetic variations and developmental constraints to alterations in neural wiring that underlie specific brain disorders, like Fetal Alcohol Syndrome. This knowledge will foster a deeper integration of structural and functional neuroscience, with broad implications for both basic research and clinical practice.

Figure 1. The Connectome and Network Science.

(a) The traditional network representation of the connectome views the neurons as separate nodes (circles), connected by links whose weights correspond to the number of synapses

between them. **(b)** The connectome is a physical network, as each neuron has a three dimensional spatial structure, as illustrated here for three neurons from the fruit fly (Flyware dataset). Each neuron is characterized by the length of its skeleton (L_i), degree (k_i), total number of synapses (S_i), and synapse density (ρ_i). **(c)** On log-linear scale a static random network has a narrow Poisson degree distribution, whose $<k>$, $<k^2>$, σ_k, expected maximum degree k_{max} and degree heterogeneity κ are analytically known, and listed below. The plot also shows $P(k)$ for a growing network with random attachment (Model A), that has an exponential $P(k)$[37]. **(d)** A lognormal distribution shown on a log-linear axis, along with its analytically calculated parameters. **(e)** The degree distribution of a scale-free network shown on a log-log plot, along with its analytically calculated parameters. **(f-h)** Compared to the exponential, Poisson and scale-free distributions, the lognormal form offers the best approximation for the degree distribution of the zebrafish **(f)**, the adult fly **(g)** and mouse **(h)**. **(i)** The scaling of the maximum degree, k_{max} with N (brain size) for the three basic network models, a random network (green), a scale-free network with degree exponent $\gamma = 2.5$, and a lognormal network. For comparison, we show as vertical bars the expected number of neurons N in the five brains explored here. **(j)** The scaling of the relative heterogeneity parameter κ with neuron number for the three basic network models.

Figure 2. Neuron Degree and Strengths Follow Stereotypic Forms
(a) The degree distribution for the eight datasets shown on a log-linear plot. The data points are log-binned to increase the statistical significance of each data point (SI 2.9). **(b)** The eight degree distributions in (a) plotted on the same figure after normalizing with their respective average (μ_k) and variance (σ_k). The fact that the distributions collapse into a single curve suggests that the values are sampled from the same distribution. As a guide to the eye, we also show as a continuous line a lognormal distribution with μ=1 and σ=1. **(c)** The KS statistics for fitting the individual datasets shown in (a) to a Poisson, exponential, power law and lognormal distributions. We find that for each dataset, the lognormal has the lowest KS statistics, indicating that it offers the best fit to the data. **(d-f)** The strength distribution for the neurons, representing the total number of synapses each neuron has, for the eight datasets shown on a log-linear plot (d), along with their rescaled values (e), and KS parameters (f). **(g-i)** The log-binned neuron skeleton length *L* distribution for the eight datasets shown on a log-linear plot, along with their rescaled values (h) and KS parameters (i). **(j-l)** The log-binned local synapse density distribution for neurons (see SI 3.2 for definition) for the eight datasets shown on a log-linear plot, along with their rescaled values (k), and KS statistics (l). See SI 3.2 for the lacking information on *C. elegans* in panel j.

Figure 3. Predicting Dependencies between Neuron Lengths, Synapse Densities, and Degrees.
(a) Scatter plot showing the strength S_i for each neuron i in function of its length L_i, on a log-log plot. We show data for seven of the eight datasets in different colors, offering evidence for the scaling of Eq. (1). We excluded the human dataset as the low cell count prevented a sufficient

diversity for fitting S vs k. **(b)** A test of Eq. (3), that connects the empirically observed scale parameter μ_S for the neuron strength (horizontal axis) with the scale parameter predicted by Eq. 3 (vertical axis). **(c)** A direct test of Eq. (4), connecting the empirical observed shape parameter σ_S for the strength (horizontal axis) with the shape parameter predicted by Eq. 4 (vertical axis). **(d,e,f)** The observed strength distributions P(S) plotted along with a lognormal distribution with the analytically predicted (μ_S, σ_S) parameters. **(g)** Scatter plot showing the degree k_i for each neuron i in function of its length S_i, on a log-log plot. We show data for each of the eight datasets in different colors, offering evidence for the sublinear scaling of Eq. (5). **(h)** A direct test of Eq. (6), that connects the empirically observed scale parameter μ_k for the neuron degrees (horizontal axis) with the scale parameter predicted by Eq. 6 (vertical axis). **(i)** A direct test of Eq. (7), connecting the empirical observed shape parameter σ_k for the neuronal (horizontal axis) with the shape parameter predicted by Eq. 7 (vertical axis). **(j,k,l)** The empirically observed degree distributions $P(k)$ (symbols) plotted along with a lognormal distribution with the analytically predicted (μ_k, σ_k) parameters.

a

$j{=}1$ p_s^1

$j{=}2$ p_s^2

$j{=}3$ p_s^3

$j{=}4$ p_s^4

b

$$L_n = 2^n \langle l \rangle \prod_{j=1}^n p_s^j$$

$$P(L_n) = \frac{1}{L_n \sigma_n \sqrt{2\pi}} e^{-\frac{(\log(L_n) - \mu_n)^2}{2\sigma_n^2}}$$

$$\mu_L = \sum_{j=1}^n \mu_p^j + n \log(2)$$

$$\sigma_L^2 = \sum_{j=1}^n (\sigma_p^j)^2$$

c

d

Layer j

FlyWire Neuron 720575940638731427

e

$p_s^j = \frac{1}{2}\frac{X_j}{X_{j-1}}$

σ_s^j

p_s^c

Layer j

Figure 4. Multiplicative Process Governs Neuron Lengths and Predicts the Lognormal Architecture.

(a) Each neuron is an incomplete tree whole growth we model via a Galton-Watson branching process. At each branching point at layer j the branch can either create two new nodes with survival probability p_s^j, or cease permanently its growth, with probability $1{-}p_s^j$. The figure illustrates this process for a small tree that survives for $n = j_{max} = 4$ layers, creating a tree of length L_4. **(b)** The model predicts that a neuron's total length L_n at any larger n follows a multiplicative process that, thanks to the central limit theorem, converges to a lognormal distribution whose parameters are governed by the layer dependent survival probability $\{p_s^j\}$, specifically the parameters $\{\mu_p^j, \sigma_p^j\}$. **(c)** The skeleton of FlyWire ALLN neuron 720575940638731427, each segment being colored according to its layer number j in the tree representation shown in panel d. **(d)** The incomplete tree behind the neuron shown in (c),

extracted from its skeleton. **(e)** The empirically measured layer dependent average branching probability p_s^j over all FlyWire neurons (black line) along with its variability across neurons (gray), shown in function of the layer number *j*. We find the average branching probability to be high near the soma (small *j*), but it rapidly decreases and stabilizes at a probability just above 0.5, i.e. always staying in the $p_s > 0.5$ regime where the analytically derived lognormal distribution, shown in (b), is valid.

Figure 5. FlyWire cell classes follow lognormal distributions. To test whether the developed analytical framework applies to individual neuron classes, we focused on the 29 neuronal classes defined by the FlyWire project[3]. **(a)** Strength distribution for neurons from five FlyWire cell classes. The difference between these classes is captured by the different average synapse numbers, spanning from median $S = e^\mu$ = 24.8 synapses for visual neurons to median $S = e^\mu$ = 177.7 synapses for gustatory neurons. **(b)** The five distributions in (a) collapsed after normalizing with their respective μ_S and σ_S value, indicating that despite the significant difference in $\mu_S = <log\ S>$ and σ_S, the values are stereotypical, being sampled from a single distribution. As a guide to the eye, we also show as a continuous line a lognormal distribution with μ=1 and σ=1. **(c)** KS statistics obtained by fitting the individual datasets shown in (a) to a Poisson, exponential, power law and log-normal distributions, indicating that for each neuronal class, the lognormal has the lowest KS statistics. **(d)** The empirically measured branching probability p_s^j for five FlyWire cell classes versus the layer number j. Each cell type has its unique

and different p_s^j function, some approaching 0.5 faster, hence expecting a shorter layer number, and smaller neuronal length L. **(e)** The neuron skeleton length distribution for the five representative FlyWire cell types shown in (a). The differences between these classes is illustrated by the significant differences in the average neuron lengths $L \sim e^\mu$, spanning from median $e^\mu = 144.5$ microns for visual neurons to median $e^\mu = 972.6$ microns for central neurons. **(f)** The five distributions in (e) after normalizing with their respective μ and σ value, along with a lognormal distribution with $\mu=1$ and $\sigma=1$. **(g)** The KS statistics for fitting the individual datasets shown in (e) to a Poisson, exponential, power-law and log-normal distributions. For each dataset the lognormal has the lowest KS statistics. **(h)** Mean measured tree size $log\,(L/\ell)$ (vertical axis) vs the tree size predicted by Eq. (9) (horizontal axis), where we used as input to Eq. (9) the empirically measured survival function p_s^j shown in (d). The symbols cover all 29 classes, highlighting the 5 classes shown in (a) and (e). Points are weighted linearly according to the relative frequency of each neuron class. The linear dependence between the predicted (horizontal axis) and the measured (vertical axis) tree size confirms that the neuron length L is primarily driven by survival branching probability p_s^j, as predicted by Eqs. (8) and (9). The slope of the fit (n, indicated in red) reveals the characteristic layer number $n \simeq 20$ in Eq. (8) at which the neurons cease their growth.

References

1. White, J. G., Southgate, E., Thomson, J. N. & Brenner, S. The structure of the nervous system of the nematode Caenorhabditis elegans. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **314,** 1–340 (1986).

2. Dorkenwald, S., Matsliah, A., Sterling, A. R., Schlegel, P., Yu, S.-C., McKellar, C. E., Lin, A., Costa, M., Eichler, K., Yin, Y., Silversmith, W., Schneider-Mizell, C., Jordan, C. S., Brittain, D., Halageri, A., Kuehner, K., Ogedengbe, O., Morey, R., Gager, J., Kruk, K., Perlman, E., Yang, R., Deutsch, D., Bland, D., Sorek, M., Lu, R., Macrina, T., Lee, K., Alexander Bae, J., Mu, S., Nehoran, B., Mitchell, E., Popovych, S., Wu, J., Jia, Z., Castro, M., Kemnitz, N., Ih, D., Bates, A. S., Eckstein, N., Funke, J., Collman, F., Bock, D. D., Jefferis, G. S. X., Sebastian Seung, H., Murthy, M. & the FlyWire Consortium. Neuronal wiring diagram of an adult brain. *Nature* **634,** 124–138 (2024).

3. Schlegel, P., Yin, Y., Bates, A. S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D. S., Gkantia, M., dos Santos, M., Munnelly, E. J., Badalamente, G., Capdevila, L. S., Sane, V. A., Pleijzier, M. W., Tamimi, I. F. M., Dunne, C. R., Salgarella, I., Javier, A., Fang, S., Perlman, E., Kazimiers, T., Jagannathan, S. R., Matsliah, A., Sterling, A. R., Yu, S.-C., McKellar, C. E., FlyWire Consortium, Costa, M., Sebastian Seung, H., Murthy, M., Hartenstein, V., Bock, D. D. & Jefferis, G. S. X. Whole-brain annotation and multi-connectome cell typing quantifies circuit stereotypy in Drosophila. *Nature* **634,** 139–152 (2024).

4. Zheng, Z., et. al. A Complete Electron Microscopy Volume of the Brain of Adult Drosophila melanogaster. *Cell* **174,** 730–743.e22 (2018).

5. Albert, R., Jeong, H. & Barabási, A. L. Error and attack tolerance of complex networks.

Nature **406,** 378–382 (2000).

6. Cohen, R. & Havlin, S. Scale-free networks are ultrasmall. *Phys. Rev. Lett.* **90,** 058701 (2003).

7. Gao, J., Barzel, B. & Barabási, A.-L. Universal resilience patterns in complex networks. *Nature* **530,** 307–312 (2016).

8. Pastor-Satorras, R. & Vespignani, A. Epidemic spreading in scale-free networks. *Phys Rev Lett* **86,** 3200–3203 (2001).

9. Liu, Y.-Y. & Barabási, A.-L. Control principles of complex systems. *Rev. Mod. Phys.* **88,** (2016).

10. Gu, S., Pasqualetti, F., Cieslak, M., Telesford, Q. K., Yu, A. B., Kahn, A. E., Medaglia, J. D., Vettel, J. M., Miller, M. B., Grafton, S. T. & Bassett, D. S. Controllability of structural brain networks. *Nat Commun* **6,** 8414 (2015).

11. Nishikawa, T., Motter, A. E., Lai, Y.-C. & Hoppensteadt, F. C. Heterogeneity in oscillator networks: are smaller worlds easier to synchronize? *Phys. Rev. Lett.* **91,** 014101 (2003).

12. Barabási, D. L., et. al. Neuroscience Needs Network Science. *J Neurosci* **43,** 5989–5995 (2023).

13. Amaral, L. A., Scala, A., Barthelemy, M. & Stanley, H. E. Classes of small-world networks. *Proc. Natl. Acad. Sci. U. S. A.* **97,** 11149–11152 (2000).

14. Sporns, O. & Zwi, J. D. The small world of the cerebral cortex. *Neuroinformatics* **2,** 145–162 (2004).

15. Ed Bullmore, Anna Barnes, Danielle S Bassett, Alex Fornito, Manfred Kitzbichler, David Meunier, John Suckling. Generic aspects of complexity in brain imaging data and other biological systems. *Neuroimage* **47,** 1125–1134 (2009).

16. van den Heuvel, M. P., Stam, C. J., Boersma, M. & Hulshoff Pol, H. E. Small-world and scale-free organization of voxel-based resting-state functional connectivity in the human brain. *Neuroimage* **43,** 528–539 (2008).

17. Eguíluz, V. M., Chialvo, D. R., Cecchi, G. A., Baliki, M. & Apkarian, A. V. Scale-free brain functional networks. *Phys Rev Lett* **94,** 018102 (2005).

18. Yang, R., Vishwanathan, A., Wu, J., Kemnitz, N., Ih, D., Turner, N., Lee, K., Tartavull, I., Silversmith, W. M., Jordan, C. S., David, C., Bland, D., Sterling, A., Goldman, M. S., Aksay, E. R. F., Seung, H. S. & Eyewirers. Cyclic structure with cellular precision in a vertebrate sensorimotor neural circuit. *Curr Biol* **33,** 2340–2349.e3 (2023).

19. Salova, A. & Kovács, I. A. Combined topological and spatial constraints are required to capture the structure of neural connectomes. *Network Neuroscience* (2025). at <http://arxiv.org/abs/2405.06110>

20. Lin, A., Yang, R., Dorkenwald, S., Matsliah, A., Sterling, A. R., Schlegel, P., Yu, S.-C., McKellar, C. E., Costa, M., Eichler, K., Bates, A. S., Eckstein, N., Funke, J., Jefferis, G. S. X. & Murthy, M. Network statistics of the whole-brain connectome of Drosophila. *Nature* **634,** 153–165 (2024).

21. Winding, M., Pedigo, B. D., Barnes, C. L., Patsolic, H. G., Park, Y., Kazimiers, T., Fushiki, A., Andrade, I. V., Khandelwal, A., Valdes-Aleman, J., Li, F., Randel, N., Barsotti, E., Correia, A.,

Fetter, R. D., Hartenstein, V., Priebe, C. E., Vogelstein, J. T., Cardona, A. & Zlatic, M. The connectome of an insect brain. *Science* **379,** eadd9330 (2023).

22. Betzel, R., Puxeddu, M. G. & Seguin, C. Hierarchical communities in the larval *Drosophila* connectome: Links to cellular annotations and network topology. *bioRxiv* (2023). doi:10.1101/2023.10.25.562730

23. Barabási, A. L. & Albert, R. Emergence of scaling in random networks. *Science* **286,** 509–512 (1999).

24. Erdős, P. & Rényi, A. On the evolution of random graphs. *Publications of the Mathematical Institute of the Hungarian Academy of Sciences* **5,** 17–61 (1960).

25. Cook, S. J., Jarrell, T. A., Brittin, C. A., Wang, Y., Bloniarz, A. E., Yakovlev, M. A., Nguyen, K. C. Q., Tang, L. T.-H., Bayer, E. A., Duerr, J. S., Bülow, H. E., Hobert, O., Hall, D. H. & Emmons, S. W. Whole-animal connectomes of both Caenorhabditis elegans sexes. *Nature* **571,** 63–71 (2019).

26. Scheffer, L. K., et. al. A connectome and analysis of the adult central brain. *Elife* **9,** (2020).

27. Azevedo, A., Lesser, E., Phelps, J. S., Mark, B., Elabbady, L., Kuroda, S., Sustar, A., Moussa, A., Khandelwal, A., Dallmann, C. J., Agrawal, S., Lee, S.-Y. J., Pratt, B., Cook, A., Skutt-Kakaria, K., Gerhard, S., Lu, R., Kemnitz, N., Lee, K., Halageri, A., Castro, M., Ih, D., Gager, J., Tammam, M., Dorkenwald, S., Collman, F., Schneider-Mizell, C., Brittain, D., Jordan, C. S., Dickinson, M., Pacureanu, A., Seung, H. S., Macrina, T., Lee, W.-C. A. & Tuthill, J. C. Connectomic reconstruction of a female Drosophila ventral nerve cord. *Nature* **631,** 360–368 (2024).

28. Vishwanathan, A., Ramirez, A. D., Wu, J., Sood, A., Yang, R., Kemnitz, N., Ih, D., Turner, N., Lee, K., Tartavull, I., Silversmith, W. M., Jordan, C. S., David, C., Bland, D., Goldman, M. S., Aksay, E. R. F., Seung, H. S. & the Eyewirers. Predicting modular functions and neural coding of behavior from a synaptic wiring diagram. *bioRxiv* (2020). doi:10.1101/2020.10.28.359620

29. MICrONS Consortium, et. al. Functional connectomics spanning multiple areas of mouse visual cortex. *bioRxiv* (2021). doi:10.1101/2021.07.28.454025

30. Schneider-Mizell, C. M., et. al. Cell-type-specific inhibitory circuitry from a connectomic census of mouse visual cortex. *bioRxiv* (2024). doi:10.1101/2023.01.23.525290

31. Shapson-Coe, A., et. al. A petavoxel fragment of human cerebral cortex reconstructed at nanoscale resolution. *Science* **384,** eadk4858 (2024).

32. Herculano-Houzel, S. & Lent, R. Isotropic fractionator: a simple, rapid method for the quantification of total cell and neuron numbers in the brain. *J. Neurosci.* **25,** 2518–2521 (2005).

33. Caldarelli, G. *Scale-Free Networks: Complex Webs in Nature and Technology*. (OUP Oxford, 2007).

34. Newman, M. E. J. & Girvan, M. Finding and evaluating community structure in networks. *Phys Rev E* **69,** 026113 (2004).

35. Fortunato, S. Community detection in graphs. *Phys. Rep.* **486,** 75–174 (2010).

36. Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D. & Alon, U. Network motifs:

simple building blocks of complex networks. *Science* **298,** 824–827 (2002).

37. Barabási, A.-L. *Network Science*. (Cambridge University Press, 2016).

38. Dorogovtsev, S. N. & Mendes, J. F. F. *The Nature of Complex Networks*. (2025).

39. Buzsáki, G. & Mizuseki, K. The log-dynamic brain: how skewed distributions affect network operations. *Nat. Rev. Neurosci.* **15,** 264–278 (2014).

40. Loewenstein, Y., Kuras, A. & Rumpel, S. Multiplicative dynamics underlie the emergence of the log-normal distribution of spine sizes in the neocortex in vivo. *J Neurosci* **31,** 9481–9488 (2011).

41. Horvát, S., Gămănuț, R., Ercsey-Ravasz, M., Magrou, L., Gămănuț, B., Van Essen, D. C., Burkhalter, A., Knoblauch, K., Toroczkai, Z. & Kennedy, H. Spatial Embedding and Wiring Cost Constrain the Functional Layout of the Cortical Network of Rodents and Primates. *PLOS Biology* **14,** e1002512 (2016).

42. Song, S., Sjöström, P. J., Reigl, M., Nelson, S. & Chklovskii, D. B. Highly Nonrandom Features of Synaptic Connectivity in Local Cortical Circuits. *PLOS Biology* **3,** e68 (2005).

43. Sheridan, P. & Onodera, T. A Preferential Attachment Paradox: How Preferential Attachment Combines with Growth to Produce Networks with Log-normal In-degree Distributions. *Sci Rep* **8,** 2811 (2018).

44. Redner, S. Citation statistics from 110 years of *physical review*. *Phys. Today* **58,** 49–54 (2005).

45. Mitzenmacher, M. A brief history of generative models for power law and lognormal distributions. *Internet Math.* **1,** 226–251 (2004).

46. Serafino, M., Cimini, G., Maritan, A., Rinaldo, A., Suweis, S., Banavar, J. R. & Caldarelli, G. True scale-free networks hidden by finite size effects. *Proc Natl Acad Sci U S A* **118,** (2021).

47. Dehmamy, N., Milanlouei, S. & Barabási, A.-L. A structural transition in physical networks. *Nature* **563,** 676–680 (2018).

48. Ercsey-Ravasz, M., Markov, N. T., Lamy, C., Van Essen, D. C., Knoblauch, K., Toroczkai, Z. & Kennedy, H. A predictive network model of cerebral cortical connectivity based on a distance rule. *Neuron* **80,** 184–197 (2013).

49. Pósfai, M., Szegedy, B., Bačić, I., Blagojević, L., Abért, M., Kertész, J., Lovász, L. & Barabási, A.-L. Impact of physicality on network structure. *Nature Physics* (2023). doi:10.1038/s41567-023-02267-1

50. Pete, G., Timár, Á., Stefánsson, S. Ö., Bonamassa, I. & Pósfai, M. Physical networks as network-of-networks. *Nat Commun* **15,** 4882 (2024).

51. Wittner, L., Henze, D. A., Záborszky, L. & Buzsáki, G. Three-dimensional reconstruction of the axon arbor of a CA3 pyramidal cell recorded and filled in vivo. *Brain Struct Funct* **212,** 75–83 (2007).

52. Menichetti, G. & Barabási, A.-L. Nutrient concentrations in food display universal behaviour. *Nature Food* **3,** 375–382 (2022).

53. Goh, K. I., Kahng, B. & Kim, D. Universal behavior of load distribution in scale-free

networks. *Phys Rev Lett* **87,** 278701 (2001).

54. Barrat, A., Barthélemy, M., Pastor-Satorras, R. & Vespignani, A. The architecture of complex weighted networks. *Proc Natl Acad Sci U S A* **101,** 3747–3752 (2004).

55. Fishell, G. & Kepecs, A. Interneuron Types as Attractors and Controllers. *Annu. Rev. Neurosci.* **43,** 1–30 (2020).

56. Sanes, J. R. & Zipursky, S. L. Synaptic Specificity, Recognition Molecules, and Assembly of Neural Circuits. *Cell* **181,** 536–556 (2020).

57. Kerstjens, S., Michel, G. & Douglas, R. J. Constructive connectomics: How neuronal axons get from here to there using gene-expression maps derived from their family trees. *PLoS Comput. Biol.* **18,** e1010382 (2022).

58. Flaherty, E. & Maniatis, T. The role of clustered protocadherins in neurodevelopment and neuropsychiatric diseases. *Curr. Opin. Genet. Dev.* **65,** 144–150 (2020).

59. Barabási, D. L. & Barabási, A.-L. A Genetic Model of the Connectome. *Neuron* **105,** 435–445.e5 (2020).

60. Kovács, I. A., Barabási, D. L. & Barabási, A.-L. Uncovering the genetic blueprint of the nervous system. *Proc. Natl. Acad. Sci. U. S. A.* **117,** 33570–33577 (2020).

61. Harris, T. E. *The Theory of Branching Processes*. (Springer, 1963).

62. Matsumoto, N., Barson, D., Liang, L. & Crair, M. C. Hebbian instruction of axonal connectivity by endogenous correlated spontaneous activity. *Science* **385,** eadh7814

(2024).

63. Cohen, R., Erez, K., ben-Avraham, D. & Havlin, S. Resilience of the internet to random breakdowns. *Phys Rev Lett* **85,** 4626–4628 (2000).

64. Restrepo, J. G., Ott, E. & Hunt, B. R. Onset of synchronization in large networks of coupled oscillators. *Phys Rev E* **71,** 036151 (2005).

65. Towlson, E. K., Vértes, P. E., Ahnert, S. E., Schafer, W. R. & Bullmore, E. T. The rich club of the C. elegans neuronal connectome. *J Neurosci* **33,** 6380–6387 (2013).

66. Bassett, D. S. & Bullmore, E. Small-world brain networks. *Neuroscientist* **12,** 512–523 (2006).

67. Li, Q., Del Ferraro, G., Pasquini, L., Peck, K. K., Makse, H. A. & Holodny, A. I. Core language brain network for fMRI language task used in clinical applications. *Netw Neurosci* **4,** 134–154 (2020).

68. Südhof, T. C. Towards an Understanding of Synapse Formation. *Neuron* **100,** 276–293 (2018).

69. Zhang, X.-Y., Moore, J. M., Ru, X. & Yan, G. Geometric Scaling Law in Real Neuronal Networks. *Phys Rev Lett* **133,** 138401 (2024).

Acknowledgements

D.L.B. is supported by funding from the Eric and Wendy Schmidt Center at the Broad Institute of MIT and Harvard. A.-L.B. is supported by the NSF award No 2243104 – COMPASS and by the European Union's Horizon 2020 research and innovation program No 810115 – DYNASNET. We thank Daria Koshkina for designing figures.

Author Contributions

B.P., D.L.B. and A.F.C. collected and cleaned the data. B.P. and D.L.B. analyzed the data. B.P. and G.M. conducted theoretical analyses. B.P., D.L.B., G.M. and A.-L.B. wrote, reviewed and edited the manuscript.

Competing Interests

A.-L.B. is the scientific founder of Scipher Medicine, Inc., which applies network medicine to biomarker development.

Supplementary Information

Supplementary Information is available for this paper.

Correspondence

Correspondence and requests for materials should be addressed to A.-L.B.