The phagosome-mediated anti-bacterial immunity is governed by the proton-activated chloride channel in peritoneal macrophages

Henry Yi Cheng[1], Jiachen Chu[1], Nathachit Limjunyawong[2,3], Jianan Chen[1], Yingzhi Ye[1], Kevin Hong Chen[1], Nicholas Koylass[1], Shuying Sun[1,2], Xinzhong Dong[2,4], Zhaozhu Qiu[1,2,5*]

Affiliations

[1]Department of Physiology, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[2]Solomon H. Snyder Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[3]Center of Research Excellence in Allergy and Immunology, Research Department, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

[4]Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[5]Department of Neurosurgery, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

*Corresponding author: zhaozhu@jhmi.edu

Summary

PAC channel mediates phagosome maturation during bacterial infection in macrophages. The deletion of PAC promotes phagosome-mediated interferon signaling and inflammasome-mediated gasdermin D secretion during bacterial infection in peritoneal macrophages.

Abstract

Phagosome degradation is an evolutionarily conserved and highly effective innate immune response against pathogen infections. The success of this process relies on the ability of phagocytes to regulate the maturation of phagosomes. However, the underlying molecular mechanisms and its roles in shaping downstream immune activation remain poorly understood. Here, we identify the proton-activated chloride (PAC) channel as a key negative regulator of phagosome maturation. PAC deletion enhanced phagosomal acidification and protease activities, leading to augmented bacterial killing in large peritoneal macrophages (LPMs) upon peritoneal *Escherichia coli* infection in mice. Surprisingly, phagosome bacterial degradation also stimulated STING-IRF3-interferon responses and inflammasome activation in LPMs, both of which are enhanced upon PAC deletion. The increased inflammasome activation and pyroptosis induced an unexpected release of cleaved gasdermin D, which localized to the surface of bacteria in the peritoneum and further contributed to their killing. Finally, enhanced bacterial clearance by PAC-deficient LPMs reduced proinflammatory immune cell infiltration and overall peritoneal inflammation, resulting in improved survival in mice. Our study thus provides new insights into the molecular mechanism of phagosome maturation and the dynamics of host defense response following phagosome-mediated bacterial degradation in peritoneal macrophages. It also highlights the potential of targeting the PAC channel as a therapeutic strategy for treating bacterial infections.

Introduction

Phagocytosis represents a fundamental cellular process through which eukaryotic cells internalize large (>0.5 µm) particles, such as microbes, cell corpses and debris, for eventual degradation (Kinchen and Ravichandran, 2008). This process is the first-line host defense in phagocytes, playing a pivotal role in combating various pathogenic infections (Ince et al., 1987). The effectiveness of this defense mechanism hinges on the ability of phagocytes to exert control over the phagosome maturation process, most importantly, through regulating acidity (Beertsen et al., 2008). Upon budding from the plasma membrane, nascent phagosomes undergo a series of maturation steps, acquiring essential components for their acidification. The vesicular proton pump V-ATPase utilizes the energy derived from ATP hydrolysis to transport protons (H^+) into the phagosomal lumen (Lukacs et al., 1990). Such vectorial transport of H^+ generates a transmembrane electrical potential difference that would restrain the activity of the pump if left uncompensated. Chloride (Cl^-) is the major counterion responsible for shunting this luminal positive membrane potential, thus facilitating vesicular acidification (Di et al., 2006). The CLC transporters mediate the accumulation of Cl^- within endo-lysosomes, neutralizing the positive charge gradient through their $2Cl^-$ to $1H^+$ exchange activity (Graves et al., 2008; Li et al., 2002). However, their precise contributions to phagosomal acidification and maturation have remained a subject of debate. For example, CLC-3, an early endosomal Cl^- exchanger, was linked to phagosomal Cl^- accumulation and acidification, but this role was not reported in other studies (Li et al., 2002; Moreland et al., 2006; Wu et al., 2023). Emerging evidence suggests that the phagosomal Cl^- concentration itself may also regulate various protease activities. For instance, CLC-7, a lysosomal Cl^- exchanger, has been shown to enhance protease activities in phagosomes and promote their fusion with lysosomes, without affecting steady-state phagosomal pH (Wong et al., 2017; Wu et al., 2023). These studies highlight the multifaceted roles of Cl^- beyond electrical balancing.

In addition to mediating the killing of microorganisms, phagosomes are also a signaling hub important for innate and adaptive immune responses (Goodridge et al., 2011; Underhill and Goodridge, 2012). For example, innate immune receptors and associated signaling proteins are localized to maturing phagosomes, exhibiting various microorganism-sensing features (Husebye et al., 2010). The antigens of digested pathogens are also processed and interact with the major histocompatibility complex (MHC-I) within the phagolysosome compartments (Nair-Gupta et al., 2014). While most studies focus on phagosomal membrane-bound receptors, emerging evidence suggests that the phagosomal contents, such as peptidoglycan from the gram-positive bacterial cell wall, may also be released after digestion and activate the cytoplasmic pattern recognition receptors (PRRs) (Wolf et al., 2016). However, how the phagosome-mediated degradation and release of bacterial components shape innate immune responses *in vivo* remain underexplored.

Through unbiased RNA interference screens, a novel acid- or proton-activated Cl^- (PAC) channel encoded by *PACC1*, also known as *TMEM206*, has been recently identified as an endosomal Cl^- channel (Osei-Owusu et al., 2021; Ullrich et al., 2019; Yang et al., 2019). By releasing Cl^- from endosomal lumen under increasing acidic environment, the PAC channel prevents hyperacidification, thus playing an opposite role to the endo-lysosomal CLC exchangers. Notably, PAC is also localized to macropinosomes, facilitating their shrinkage by mediating Cl^- efflux (Zeziulia et al., 2022). Despite the importance of Cl^-, it remains unknown if the PAC channel plays a role in phagosome maturation and how it may contribute to pathogen degradation and subsequently shape the host defense response against infections.

In this study, we identify the PAC channel as an important negative regulator for phagosome maturation in macrophages. Using a peritoneal *Escherichia coli* infection model, we revealed a critical role for PAC in bacterial degradation in large peritoneal macrophages (LPMs). The

enhanced bacterial killing in PAC-deficient LPMs activated STING-IRF3-IFNβ axis and inflammasomes through the increased release of bacterial ligands from phagosomes, which further drive the macrophage bactericidal activity. With a novel depletion-adoptive transfer model, we demonstrated crucial roles of PAC and inflammasome-activated gasdermin D in LPM-mediated bacterial clearance. Moreover, we determined how the resolution of peritoneal infection was facilitated by the augmented phagosome bacterial degradation in PAC-deficient LPMs through comprehensive and unbiased analyses, including secretome, RNA-seq, and high-dimensional mass cytometry. Our findings therefore provide new insights into the intricate mechanisms underlying phagosome maturation upon bacterial infection and the dynamics of the innate immune responses triggered by pathogen degradation.

Results

The PAC channel localizes to phagosomes and regulates bacterial clearance in macrophages

To identify ion channels and transporters that regulate phagosome maturation, we analyzed a published proteome dataset on phagosomes isolated from mouse macrophages (Guo et al., 2015b) and generated a list of 48 potential candidates (Fig. 1 A and Fig. S1 A). We further cross-referenced with the GeneCards database and narrowed down to 22 proteins that also localize to endo-lysosome compartments thus potentially involved in phagosome maturation. We then mapped the mRNA expression levels of these genes across immune cells with the ImmGen database (Heng et al., 2008). Interestingly, *Pacc1*, encoding the newly identified proton-activated Cl⁻ channel (Ullrich et al., 2019; Yang et al., 2019), was among the highest and most specifically expressed genes across the four types of macrophages (Fig. 1 B and C), including splenic macrophages which are specialized phagocytes relying on phagosomes to degrade their cargo (Flannagan et al., 2009). The proteome and expression analyses suggest a role for the PAC channel in phagosome maturation and macrophage functions. To validate PAC's phagosomal localization, we performed immunofluorescence analysis on human THP1-differentiated macrophages with a specific monoclonal antibody for human PAC protein. In the resting state, PAC was predominantly localized to early and recycling endosomes, consistent with previous findings in other cell types (Osei-Owusu et al., 2021; Zeziulia et al., 2022) (Fig. S1 B and C). To examine various phagocytosis processes, we applied different substrates, including inorganic silica beads, zymosan particles, and live GFP-expressing *E. coli* (*E. coli*-GFP), and observed the localization of PAC to phagosomes with all three types of cargo (Fig. 1 D and E and Fig. S1 D). Live cell imaging of Raw264.7 macrophages expressing PAC-mCherry also revealed the association of PAC with the dynamic internalization of *E. coli-GFP* (Fig. 1 F and Fig. S1 E). These data are consistent with the

phagosome proteome dataset and a previous report on mouse bone marrow-derived macrophages (BMDMs) treated with dead *E. coli* particles (Zeziulia et al., 2022).

To further examine the function of PAC in phagosome maturation and macrophage functions, we cultured BMDMs from PAC whole-body knockout (*Pacc1*$^{-/-}$; KO) mice and validated the loss of PAC-specific acid-induced Cl$^-$ currents ($I_{Cl, H}$) by patch clamp recordings (Fig. S1 F). PAC deletion did not alter BMDM differentiation from bone-marrow hematopoietic stem cells treated with either L929 cell conditioned media or M-CSF supplement (Fig. S1 G). Given the critical role of macrophages in clearing bacterial infection, we next evaluate the bactericidal activity of *Pacc1*$^{-/-}$ BMDMs by infecting them with *E. coli*. Interestingly, while *Pacc1*$^{-/-}$ BMDMs internalized a similar number of *E. coli* as control cells, the surviving bacteria within the KO cells were less than WT cells after 1-2 hours of digestion, indicating an enhanced bactericidal activity in macrophages without PAC channel (Fig. 1 G).

To further examine the role of PAC in bacterial clearance *in vivo*, we adopted a peritoneal *E. coli* infection model (Fig. 1 H). In this model, immune cells can be obtained from peritoneal lavage immediately after infection and subjected to fluorescence-activated cell sorting (FACS) analyses and functional studies. Large peritoneal macrophages (LPMs) and small peritoneal macrophages (SPMs) are two resident macrophages in the peritoneum, characterized as the CD11b$^+$, F4/80^{high}, and I-A/I-E^{low} population, and CD11b$^+$, F4/80^{low}, and I-A/I-E^{high} population, respectively (Fig. 1 I). LPMs in particular are the major phagocytes of the first-line defense against peritoneal infection (Vega-Perez et al., 2021). To examine the role of PAC in bacterial killing, we crossed *Pacc1*$^{F/F}$ mice with *Cx3cr1* promoter-driven Cre mice to selectively delete the gene in all resident macrophages and monocytes (*Cx3cr1*-PAC$^{-/-}$). The loss of PAC channel activity in LPMs isolated from *Cx3cr1*-PAC$^{-/-}$ mice was validated by patch clamp recordings (Fig. 1 J). The mutant mice

had similar numbers of LPMs, SPMs, and monocytes in the peritoneum compared to their $Pacc1^{F/F}$ control littermates, suggesting that the PAC channel is not required for their general development (Fig. S2 A and B). As expected, neutrophil and various lymphocyte (CD4 T cells, CD8 T cells, B cells, and NK cells) populations in the peritoneum were also normal since Cre expression is mainly limited to macrophages (Fig. S2 B-E). For real-time *in vivo* tracking of bacterial clearance, we peritoneally infected mice with *E. coli*-GFP and analyzed GFP intensity in cells isolated from the peritoneal lavage. Given that PAC deletion does not affect initial phagocytosis and bacterial uptake (Fig. 1 K), GFP fluorescence within various immune cells can serve as an indicator of their bactericidal activity. Consistent with the enhanced bacterial clearance *in vitro*, we observed a more rapid decrease in GFP intensity in LPMs from $Cx3cr1$-PAC$^{-/-}$ mice compared to controls (Fig. 1 K). This effect was specific for LPMs as the bacterial clearance in neutrophils was comparable between $Cx3cr1$-PAC$^{-/-}$ and control mice. No *E. coli*-GFP signals were observed above the background for SPMs, consistent with their non-phagocyte status (Fig. S2 G).

$Cx3cr1$-Cre line was generated through a knock-in strategy, leading to the deletion of one allele of endogenous $Cx3cr1$, a chemokine receptor (Yona et al., 2013). Although the immune system development was grossly normal in $Cx3cr1$-PAC$^{-/-}$ mice, we further validated our findings by crossing $Pacc1^{F/F}$ mice with *LysM*-Cre (Clausen et al., 1999), another popular Cre line specific for myeloid cell lineages to generate *LysM*-PAC$^{-/-}$ mice. Consistently, *LysM*-PAC$^{-/-}$ mice also had normal numbers of peritoneal myeloid populations (Fig. S2 A and C) and lymphoid populations (Fig. S2 D and F), confirming that PAC is not essential for their development. We next performed peritoneal infection of *E. coli*-GFP and observed comparable decreases in GFP intensity in LPMs from *LysM*-PAC$^{-/-}$ mice, similar to those from $Cx3cr1$-PAC$^{-/-}$ mice (Fig. 1 K and Fig. S2 G). Taken together, these results from both *in vitro* and *in vivo* experiments with two independent Cre

lines demonstrate that PAC-deficient macrophages exhibit an enhanced bacterial clearance capability.

PAC downregulation is required for LPS-induced phagosome maturation

To further explore the role of PAC in bacterial infections, we analyzed *PACC1* expression in published transcriptome databases of different infectious diseases (Bertrams et al., 2020; Scicluna et al., 2020). Interestingly, we found that *PACC1* was significantly downregulated in the peripheral blood mononuclear cells (PBMCs) of septic and pneumonia patients (Fig. 2 A). To test if this happens in macrophages, we treated BMDMs with lipopolysaccharides (LPS), the main surface membrane components of infectious bacteria that trigger immune cell activation (Poltorak et al., 1998). Strikingly, LPS treatment markedly reduced *Pacc1* mRNA levels as well as the PAC channel activity (Fig. 2 B and C). This LPS-mediated PAC downregulation was also observed in human THP1 cells at both mRNA and protein levels (Fig. S3 A and B).

Given its role in vesicular acidification (Osei-Owusu et al., 2021; Zeziulia et al., 2022), we tested whether the PAC channel regulates phagosomal acidification by generating PAC KO THP1 cells using CRISPR-Cas9 approach (Fig. S3 C). We next performed ratiometric measurement of phagosomal pH using zymosan particles conjugated with pH-sensitive PhrodoGreen and pH-insensitive Alexa Flour-633 (AF633) fluorescence dyes. Interestingly, loss of PAC resulted in ~0.5 unit lower pH in phagosomes at steady-state compared to WT THP1 cells (Fig. 2 D). We further validated this finding in BMDMs and LPMs. Consistent with *E. coli* uptake results (Fig. 1 G), *Pacc1*$^{-/-}$ BMDMs also had a normal uptake of zymosan particles (Fig. 2 E). However, phagosomes were more acidic during the maturation process in the KO BMDMs compared to WT control cells (Fig. 2 F). Additionally, we observed a similar decrease in phagosomal pH in LPMs isolated from *Cx3cr1*-PAC$^{-/-}$ mice after peritoneally injecting pH-sensing zymosan particles (Fig. 2 G, left).

Given that the activities of phagosomal proteases are dependent on acidic pH (Kinchen and Ravichandran, 2008), we measured them by a similar ratiometric method using zymosan beads double-conjugated with bovine serum albumin (BSA) self-quenched DQ green (BSA-DQ) fluorescence dye and AF633. The protease activities were indicated by the ratio of the green fluorescence detected after BSA is degraded by phagosomal proteases to internal control AF633. Consistent with enhanced phagosomal acidification, we also observed a modest but significant increase in protease activities in LPMs isolated from $Cx3cr1$-PAC$^{-/-}$ mice after peritoneally injecting BSA-DQ particles (Fig. 2 G, right). Together, these results demonstrate that the PAC channel plays a negative role in phagosome acidification and maturation.

LPS-primed macrophages exhibit enhanced phagosome degradation, suggesting that LPS promotes phagosome acidification and maturation (Wu et al., 2023). Indeed, BMDMs primed with LPS had more acidic phagosomes with increased overall protease activities (Fig. 2 H and I, black). Without LPS treatment, $Pacc1^{-/-}$ BMDMs already exhibited a similarly more acidic phagosomal pH and higher protease activities at the baseline (Fig. 2 H and I, red), indicating that loss of PAC mimics the effect of LPS on phagosome maturation. Importantly, treating $Pacc1^{-/-}$ BMDMs with LPS did not generate further responses, suggesting that the PAC channel may mediate the effect of LPS priming on phagosome maturation. To test this hypothesis, we overexpressed the PAC protein in Raw264.7 macrophages (Fig. 2 J). Consistent with a negative role in phagosome maturation, PAC overexpression alkalized phagosomal pH and inhibited overall protease activities (Fig. 2 K and L). Similar to BMDMs, LPS priming promoted phagosome activation in vector-transfected Raw264.7 macrophages (Fig. 2 K and L, black). However, these effects were abolished in PAC-overexpressing cells (Fig. 2 K and L, blue), further suggesting that PAC downregulation is required for LPS-mediated phagosome activation. In addition to phagosome maturation, LPS

also activates macrophage toll-like receptor 4 (TLR4) and its downstream IRF3 and NFκB signaling pathways (Poltorak et al., 1998). Interestingly, PAC deletion did not alter LPS-induced expression of IRF3-regulated *IFNβ* and NFκB-mediated transcripts (*IL-6*, *TNFα*, and *IL-1β*) in THP-1 and BMDMs (Fig. S3 D). These results indicate that PAC does not regulate LPS-induced TLR4 activation.

Enhanced phagosome bacterial degradation promotes interferon responses in PAC-deficient LPMs

To further investigate potential downstream responses following enhanced bacterial degradation, we performed RNA-sequencing (RNA-seq) analysis on LPMs isolated from *Cx3cr1*-PAC$^{-/-}$ and control mice after peritoneal *E. coli* infection. Interestingly, the gene enrichment analysis (GSEA) showed that the signatures of "killing of cells of another organism" and "MHC-I protein binding" were upregulated in PAC-deficient LPMs compared to control cells (Fig. 3 A). Given the antigen processing of MHC-I complexes occurs in phagosomes, these results are consistent with enhanced phagosome function. The gene ontology (GO) analysis further revealed upregulation of interferon responses in PAC-deficient LPMs (Fig. 3 B and Fig. S4 A and B), with increased expression of many IFN-responsive genes, including several *Ifi*, *Gbp*, and *Oas* family genes (Fig. 3 C). To validate the enhanced IFN responses, we infected LPMs with *E. coli in vitro* and performed a comprehensive secretome analysis (Fig. S4 C and D). In accordance with the RNA-seq data, *Pacc1*$^{-/-}$ LPMs showed increased release of IFNβ and many other cytokines compared to control LPMs (Fig. 3 D and Fig. S4 E). Specifically, the KO cells released more chemokines (CXCL1, CCL3, and CCL4) and alarmins (IL-1α, IL-23, and IL-33) (Fig. S4 F and G), which may contribute to LPM activation and host defense against bacterial infection (Chan et al., 2012; Tilstam et al.,

2021). Interestingly, we also observed increased releases of IL-1β and IL-18 in *Pacc1*$^{-/-}$ LPMs, indicating enhanced inflammasome activation (Fig. S4 H).

With increased IFNβ production in LPMs from *Cx3cr1*-PAC$^{-/-}$ mice after peritoneal *E. coli* infection, we further examined its downstream signaling pathways. The IFNβ-induced phosphorylation of STAT1 and downstream IRF1 expression drive the transcription of IFN responsive genes during bacterial infections (Man et al., 2016). Consistent with RNA-seq and secretome results, immunoblotting revealed increases in STAT1 phosphorylation and IRF1 expression in PAC-deficient LPMs compared to control cells (Fig. 3 E). Immunostaining further showed elevations of their translocation into nucleus (Fig. 3 F). The increased expression of downstream GBP and OAS family proteins in *Cx3cr1*-PAC$^{-/-}$ LPMs was also validated by immunoblotting and FACS analyses (Fig. 3 G and Fig. S4 I). These data suggest that the increased activation of IFNβ-STAT1-IRF1 axis in PAC-deficient LPMs drives the expression of crucial macrophage anti-microbial genes, such as *Gbp* and *Oas*, that participate in pathogen sensing, microbial killing and inflammasome activation (Fisch et al., 2023; Hornung et al., 2014; Leisching et al., 2019; Man et al., 2016).

Enhanced phagosome bacterial degradation induces IFNβ expression through STING-IRF3 activation in PAC-deficient LPMs

We next investigated the potential mechanism underlying the enhanced IFNβ production in PAC-deficient LPMs upon peritoneal *E. coli* infection. During bacterial infection, the production of IFNβ is regulated by cell surface TLR4 and cytoplasmic pattern recognition receptors (PRRs) (Honda et al., 2006; Manzanillo et al., 2012). Given that no significant change in TLR4-induced genes was observed in *Pacc1*$^{-/-}$ BMDMs upon LPS treatment (Fig. S3 D), we reasoned that the

increased IFNβ production in PAC-deficient LPMs may be driven by augmented activation of cytoplasmic PRRs. Interestingly, the STING (stimulator of interferon genes) receptor has been reported to drive macrophage IFN production in various bacterial infections (Andrade et al., 2016; Manzanillo et al., 2012; Moretti et al., 2017; Ruangkiattikul et al., 2017), presumably by sensing bacterial cyclic dinucleotides (CDNs) in the cytoplasm. Therefore, we hypothesized that the enhanced phagosome bacterial degradation in PAC-deficient LPMs drives STING hyperactivation, thereby promoting downstream IFNβ production. To test this hypothesis, we performed immunoblotting and evaluated the phosphorylation of STING and the downstream transcription factor IRF3. Indeed, we observed the activation of STING-IRF3 signaling in control LPMs upon peritoneal *E. coli* infection. More importantly, STING-IRF3 activation was further enhanced in *Cx3cr1*-PAC$^{-/-}$ LPMs (Fig. 3 H). We next assayed NFκB p65 activation which is also downstream of STING as well as other cytoplasmic PRRs (Abdullah and Knolle, 2014). Consistently, the phosphorylation and translocation of p65 were also augmented in *Cx3cr1*-PAC$^{-/-}$ LPMs (Fig. S4 J and K). To further examine the role of phagosome degradation in interferon responses, we treated mice with E64d, a phagosome pan-cathepsin inhibitor, which potently blocked the bactericidal activity of LPMs in the peritoneal *E. coli* infection model (Fig. 3 I). E64d treatment also abolished the activation of STING-IRF3 pathway in both control and *Cx3cr1*-PAC$^{-/-}$ LPMs (Fig. 3 J), thereby eliminating IFNβ production (Fig. 3 K). Notably, E64d treatment increased total STING protein expression (Fig. 3 J), likely due to its effect on STING lysosomal turnover properties (Kuchitsu et al., 2023). Even though STING was stabilized upon E64d treatment, no phosphorylated STING was detected, further demonstrating the dependency of STING activation on phagosome bacterial degradation. Together, our findings provide an underappreciated mechanism downstream of phagosome degradation, in which STING-IRF3 axis is activated by

phagosome bacterial degradation and triggers the IFNβ-STAT1-IRF1 activation to induce host defense genes expression in LPMs. The enhanced phagosome degradation in PAC-deficient macrophages further promotes the activation loop and creates a hyperactivated status in response to *E. coli* infection.

Augmented phagosome bacterial degradation induces inflammasome activation and pyroptosis in PAC-deficient LPMs

LPMs are known to undergo pyroptosis, but not apoptosis or necroptosis during peritoneal *E. coli* infection (Vega-Perez et al., 2021). Consistently, we observed the releases of IL-1β, IL-18, and lactate dehydrogenase (LDH) from control LPMs after *E. coli* infection *in vitro* (Fig. 4 A and B and Fig. S4 H). The treatment of necrosulfonamide (NSA), a gasdermin D inhibitor, abolished IL-1β release further confirming inflammasome activation (Fig. S5 A). The releases of IL-1β, IL-18, and LDH were all increased in *Pacc1*$^{-/-}$ LPMs compared to control cells, indicating that loss of PAC leads to enhanced inflammasome activation and pyroptosis (Fig. 4 A and B and Fig. S4 H). We next examined the LPM inflammasome status *in vivo* by isolating and characterizing them after peritoneal *E. coli* infection. Consistently, ~40% LPMs isolated from control mice were Annexin V and 7-AAD double positive, indicating ongoing cell death. More LPMs underwent cell death in *Cx3cr1*-PAC$^{-/-}$ mice (Fig. 4 C). Accordingly, more gasdermin D pores were observed on their cell surface by non-permeabilized immunostaining (Fig. 4 D) and FACS analysis (Fig. S5 B), indicating increased pyroptotic cell death. Inflammasome activation entails caspase-1 (CASP1) cleavage into active CASP1 p20, further leading to the cleavage of gasdermin D into pore-forming N-terminus gasdermin D (gasdermin D-NT) (Barnett et al., 2023). Consistent with increased gasdermin D pore formation, immunoblotting analysis revealed that the cleavage of CASP1 and its downstream target gasdermin D were elevated in LPMs from *Cx3cr1*-PAC$^{-/-}$ mice after

peritoneal *E. coli* infection (Fig. 4 E). The increased CASP1 activation was also validated by FACS analysis using fluorescein-conjugated CASP1 inhibitor FLICA (FLICA-FAM), which specifically binds to the active CASP1 cleavage site (Fig. S5 C).

To investigate the mechanism underlying the increased inflammasome activation in PAC-deficient LPMs, we first evaluated the IL-1β secretion with two canonical inflammasome activators, ATP and nigericin, after LPS priming. Surprisingly, no difference between control and *Pacc1*$^{-/-}$ LPMs was observed, suggesting that PAC regulates inflammasome activation specifically in the context of bacterial infection, likely involving phagosome degradation (Fig. S5 D). To test this hypothesis, we treated LPMs with E64d, which, as expected, had no effect on the canonical ATP-induced IL-1β secretion (Fig. S5 E). However, E64d treatment reduced *E. coli*-induced IL-1β release in control LPMs. More importantly, E64d entirely abolished the potentiation of IL-1β release observed in *Pacc1*$^{-/-}$ LPMs (Fig. 4 F). These data suggest that the augmented phagosome bacterial degradation in PAC-deficient LPMs leads to the enhanced inflammasome activation. Inflammasomes are formed by different sensors that are activated by various cytoplasmic bacterial ligands, such as LPS, flagellin, and bacterial DNA (Barnett et al., 2023). Given that we did not observe *E. coli* escaping from phagosomes, the phagosome degradation-mediated inflammasome activation indicates a potential release of these bacterial ligands from phagosomes to the cytoplasm. Among various ligands, flagellin can be readily detected by antibody-based assays. Therefore, we first performed immunofluorescence analysis on flagellin in LPMs isolated after peritoneal *E. coli* infection. As expected, large flagellin signals were co-localized with LAMP1, indicating bacteria within phagolysosomes. Interestingly, we also observed many small flagellin-positive puncta in the cytoplasm of both control and *Cx3cr1*-PAC$^{-/-}$ LPMs, which did not colocalize with LAMP1$^+$ phagolysosomes (Fig. S5 F). Consistently, flagellin proteins were also detected by immunoblotting

in the cytosolic fraction of LPM lysates (Fig. 4 G). In accordance with enhanced phagosome degradation, the level of cytosolic flagellin was higher in *Cx3cr1*-PAC$^{-/-}$ LPMs compared to control cells (Fig. 4 G). More importantly, peritoneal injection of E64d before *E. coli* infection blocked flagellin release in the cytoplasm and eliminated the elevation of cell death in *Cx3cr1*-PAC$^{-/-}$ LPMs (Fig. 4 H and I). These results suggest that the phagosome bacterial degradation promotes inflammasome activation and pyroptosis through the release of bacterial ligands from phagosomes in LPMs.

Previous studies have shown that reconstituted gasdermin D-NT proteins forms large pores on liposomes containing cardiolipin, a phospholipid found in bacterial membranes (Ding et al., 2016; Liu et al., 2016). Moreover, co-culturing *E. coli* with purified gasdermin D-NT or supernatants from inflammasome-activated cells kills bacteria, suggesting a direct bactericidal activity (Liu et al., 2016). To test if the augmented pyroptosis in PAC-deficient LPMs directly facilitate peritoneal bacterial killing, we first examined the release of gasdermin D by LPMs upon *E. coli* infection. Consistent with increased inflammasome activation, *Pacc1*$^{-/-}$ LPMs secreted more gasdermin D to the supernatants than control cells (Fig. S5 G). Importantly, treating fresh *E. coli* culture with antibiotic-free supernatants from infected *Pacc1*$^{-/-}$ LPMs more strongly inhibited bacterial growth than antibiotic-free supernatants from infected control LPMs. To further validate the bactericidal role of the released gasdermin D, we also treated *E. coli* with antibiotic-free supernatants from infected *Gsdmd*$^{-/-}$ LPMs, which showed no inhibition in bacterial growth (Fig. 4 J and Fig. S5 H). These data suggest that extracellular gasdermin D exhibits direct bactericidal activity *in vitro*. To test this *in vivo*, we collected peritoneal fluids from mice after peritoneal *E. coli* infection and observed a higher gasdermin D concentration in *Cx3cr1*-PAC$^{-/-}$ mice than controls (Fig. 4 K). Interestingly, we also observed more gasdermin D puncta on the surface of bacteria isolated from

the peritoneum of the *Cx3cr1*-PAC$^{-/-}$ mice (Fig. 4 L), further supporting gasdermin D pore-forming as a potential bactericidal mechanism.

LPM depletion-adoptive transfer model reveals a role for gasdermin D in bacterial clearance

We next sought to further examine the role of LPM inflammasome activation and pyroptosis in bacterial clearance by focusing on their main effector gasdermin D. However, the absence of conditional KO mouse models for gasdermin D prevents the evaluation of its specific role in LPMs during peritoneal bacterial infection. To circumvent this limitation, we developed a novel depletion-adoptive transfer model for LPMs (Fig. 4 M and Fig. S5 J). Clodronate-containing liposomes can effectively deplete cavity resident macrophages, such as peritoneal and pleural macrophages (van Rooijen and Sanders, 1997). Indeed, peritoneal injection of clodronate liposomes specifically depleted LPMs without affecting SPMs (Fig. S5 I). To further minimize the depletion-induced inflammation, we waited 12 days before the adoptive transfer. We examined the peritoneal cell composition after the depletion, which showed none to minimal neutrophil recruitment, indicating little inflammation, and modest monocyte recruitment, likely to replenish the depleted LPMs (Fig. S5 K). We labeled MACS-isolated LPMs from donor mice with CellTrace dyes to track their viability after the adoptive transfer. FACS analysis demonstrated a clear gating for the transferred cell with good viability (Fig. S5 L). Strikingly, the adoptive transfer of WT control LPMs significantly protected mice from lethality caused by peritoneal *E. coli* infection (Fig. S5 M), thereby validating this new model. With this new tool, we first tested if PAC truly plays a LPM cell-autonomous role in bacterial clearance. We adoptively transferred *Pacc1*$^{-/-}$ LPMs to LPM-depleted mice and observed an augmented anti-bacterial activity of the KO LPMs compared to control LPMs after peritoneal *E. coli* infection. This included a decrease in intracellular *E. coli*-GFP signals in LPMs (Fig. 4 N), indicating an increased bactericidal activity,

and a reduction in peritoneal *E. coli* number (Fig. 4 O), suggesting a better infection resolution. These data corroborate the findings based on *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$ mice, supporting a role in bacterial clearance for the PAC channel specifically in LPMs. They further validate the depletion-adoptive transfer model as a reliable method to investigate the function of LPMs.

To examine the role of LPM pyroptosis in bacterial clearance, we next adoptively transferred LPMs isolated from *Gsdmd*$^{-/-}$ mice to LPM-depleted mice and then performed peritoneal infection with *E. coli*-GFP (Fig. 4 M). As expected, *Gsdmd* deletion protected the transferred LPMs from pyroptotic cell death during infection (Fig. 4 P). Intracellular GFP signals were similar between transferred control and *Gsdmd*$^{-/-}$ LPMs, suggesting that gasdermin D in LPMs does not affect *E. coli* uptake or killing (Fig. 4 Q). However, interestingly, we observed an increase in *E. coli* number within peritoneal fluids in mice transferred with *Gsdmd*$^{-/-}$ LPMs (Fig. 4 R). This data indicates that gasdermin D in LPMs is involved in promoting extracellular bacterial clearance, consistent with a direct bacterial killing activity previously proposed (Ding et al., 2016; Liu et al., 2016) and further supported by our findings (Fig. 4 J-L and Fig. S5 G and H). Together, these results suggest that enhanced phagosome bacterial degradation in PAC-deficient LPMs promotes inflammasome activation and gasdermin D release, which further facilitates bacterial removal and disease resolution of *E. coli* infection.

PAC deletion in macrophages mitigates peritoneal inflammation and promotes resolution of peritoneal *E. coli* infection

The balance between cellular activation for anti-microbial activities and subsequent inflammation are crucial determinants for host survival during infections (Guo et al., 2015a). The hyperactivation of IFN and inflammasomes in PAC-deficient LPMs prompted us to examine the overall peritoneal inflammation status in the KO mice. To achieve that, we first performed a comprehensive

secretome analysis with peritoneal fluids after peritoneal *E. coli* infection (Fig. S4 D). Surprisingly, we observed a consistent reduction in several inflammatory cytokines and chemokines in both *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$ mice compared to controls, including IL-6, macrophage migration inhibitory factor (MIF), and CCL8 (Fig. 5 A and B and Fig. S6 A and B). In addition, CXCL2 and IL-1β levels were also decreased in *LysM*-PAC$^{-/-}$ mice. MIF is a major driver for peritoneal inflammation in a peritonitis-induced sepsis model (Tilstam et al., 2021). IL-6 and IL-1β are key proinflammatory cytokines, while CCL8 and CXCL2 as major chemoattractants for neutrophils and monocytes. Only one chemokine, CXCL9, was increased in *Cx3cr1*-PAC$^{-/-}$ mice. CXCL9 has been shown to protect mice against colonic bacterial infection although the underlying mechanism is unclear (Reid-Yu et al., 2015). We also observed a very modest increase of IL-33 in *LysM*-PAC$^{-/-}$ mice. Given its very low concentration, the significance of such an increase remains unclear. Overall, the secretome analysis revealed a less inflammatory environment in the peritoneum of macrophage-specific PAC KO mice after *E. coli* infection. These results suggest that although PAC-deficient LPMs were hyperactivated upon bacterial infection, enhanced bacterial clearance may dampen overall peritoneal inflammation due to reduced activation of other peritoneal cells, such as SPMs as well as recruited proinflammatory neutrophils and monocytes.

To evaluate the contribution of infiltrated immune cells to peritoneal inflammation, we employed an unbiased high-dimensional mass cytometry (termed cytometry by Time-Of-Flight, cyTOF) analysis to examine changes in immune cell populations upon peritoneal infection. Given the chemokine receptor nature of CX3CR1, we focused our analysis on *LysM*-PAC$^{-/-}$ instead of *Cx3cr1*-PAC$^{-/-}$ mice. We isolated cells from the peritoneal lavage after *E. coli* infection and subjected them to cyTOF analysis (Fig. S7 A). We first unmasked major peritoneal cell populations using algorithmic clustering, which discerned nine major immune populations, including LPMs,

SPMs, neutrophils, dendritic cells, monocytes, mast cells, NK cells, T cells (CD4, CD8, and γδT cells), and B cells (Fig. 5 C and D and Fig. S7 B and C). As expected, *E. coli* infection in control mice resulted in dramatic increases in neutrophils and monocytes, the two major infiltrated immune cells during peritoneal bacterial infection. Consistent with a less inflammatory environment, the recruitment of neutrophils as well as highly proliferative Ly6C^{high} monocytes was markedly reduced in *LysM*-PAC$^{-/-}$ mice compared to controls (Fig. 5 E-G and Fig. S7 D and E). The decreased recruitment of these infiltrating cells was also validated with FACS analysis (Fig. S7 F). Together with the secretome analysis, these results suggest that enhanced bacterial clearance by PAC-deficient LPMs reduces the subsequent recruitment of proinflammatory cells, thereby further mitigating peritoneal inflammation.

To further evaluate the role of PAC in infection resolution *in vivo*, we conducted a time-course analysis of bacterial load in peritoneal fluids after peritoneal *E. coli* infection. Consistent with the enhanced bactericidal activity in LPMs, both *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$ mice exhibited a reduced peritoneal bacterial load compared to controls (Fig. 6 A). Uncontrolled peritoneal bacterial infection can lead to systematic sepsis and even death (Brun-Buisson et al., 1995). In line with the decreased bacterial load and dampened peritoneal inflammation, both macrophage-specific PAC KO mice showed alleviated bacteremia in the tail blood and were largely protected from sepsis-induced mortality (Fig. 6 B and C). In addition to WT *E. coli* strain, we also observed enhanced bacterial clearance and improved survival for peritoneal infection of a clinically relevant carbapenem-resistant strain of *E. coli* (cpm-resistant *E. coli*) (Fig. 6 D and E). These data demonstrate that enhanced phagosome bacterial degradation together with downstream IFN and inflammasome activation in PAC-deficient macrophages promotes disease resolution and reduces mortality against peritoneal *E. coli* infections *in vivo*.

Discussion

Our study identifies that PAC, a newly identified chloride channel, as a critical negative regulator for phagosome maturation and anti-bacterial immunity. PAC-deficient macrophages exhibited increased phagosomal acidification and protease activation, leading to enhanced bactericidal activity and improved mouse survival upon peritoneal *E. coli* infection (Fig. 7). These findings not only highlight the importance of the PAC channel in phagosome maturation and bacterial clearance, but also provide an ideal model to examine the role of phagosome-mediated degradation in shaping host defense responses against bacterial infection.

We find that enhanced phagosome bacterial degradation in PAC-deficient LPMs boosted inflammasome activation. This is likely due to the increased release of bacterial components, such as flagellin, to the cytoplasm, which can activate cytoplasmic NOD-like receptors NAIP5-NLRC4 (Paidimuddala et al., 2023). Augmented phagosome bacterial degradation in PAC-deficient LPMs also stimulated STING-IRF3, another key innate immune pathway, leading to IFNβ production. Interestingly, a recent preprint reported that evolutionarily diverse bacteria, including *E. coli*, stimulate IFN responses in macrophages after phagosome-mediated bacteriolysis in a membrane bound-TLR-independent, but STING-dependent manner. The authors further provided evidence that two CDN transporters LRRC8A (as known as SWELL1) and SLC46A2 are responsible for the phagosome-mediated cytoplasmic activation of STING (Stephanie A. Ragland, 2022). STING and various other cytoplasmic PRRs are traditionally thought to be only activated by intracellular pathogens, such as viruses and bacteria escaping phagosomes (Abdullah and Knolle, 2014; Man et al., 2016). However, our findings and the new study highlight the importance of phagosome-mediated bacteriolysis in innate immune activation by engaging cytoplasmic PRRs with the released bacterial ligands (CDNs, flagellin, and nucleic acids) from phagosomes. Future research

will determine if and how various bacterial components are transported from phagosomes to the cytoplasm and whether they activate their respective receptors and signaling pathways. Beyond *E. coli*, phagosome-medicated degradation of other types of pathogens or cargo may also use a similar mechanism to engage with cytoplasmic PRRs to elicit various immune responses.

Besides the conventional phagocytic bacterial killing, our comprehensive analysis on PAC-deficient LPMs in the peritoneal *E. coli* infection model reveals two new anti-bacterial mechanisms that are downstream of elevated phagosome degradation. First, the activation of STING-IRF3 signaling pathway drives the production of IFNβ, which upregulates the expression of key anti-microbial factors, including GBP and OAS family proteins (Fisch et al., 2023; Hornung et al., 2014; Leisching et al., 2019; Man et al., 2016). Second, enhanced inflammasome activation and pyroptosis in LPMs further facilitate bacterial clearance by releasing gasdermin D to the peritoneum, which forms pores on the bacterial surface and kill them (Ding et al., 2016; Liu et al., 2016). Additional mechanisms of bacterial removal may also exist. For example, the pyroptotic LPMs may form bacterial traps from cell corpses, which eventually promote efferocytosis by other phagocytes (Jorgensen et al., 2016).

The popular *Cx3cr1*-Cre and *LysM*-Cre mice we used here mediate gene deletion in all macrophages, including SPMs, monocytes, and potentially other cell types (Clausen et al., 1999; Yona et al., 2013). To circumvent this limitation, we successfully developed a new depletion-adoptive transfer model for LPMs. This novel approach complemented our genetic approach using conditional KO mice, validating a role for PAC specifically in LPMs in facilitating bacterial clearance. It also overcame the challenge laid by the lack of gasdermin D conditional KO mice and revealed a direct involvement of LPM inflammasome activation in peritoneal bacterial killing.

Thus, this new depletion-adoptive transfer model will be useful for the field to investigate the biology of LPMs.

Our RNA-seq analysis of *E. coli*-infected LPMs revealed that the upregulated genes were mainly associated with signatures of bacterial sensing and killing instead of conventional inflammation pathways. This observation aligns with the notion that LPMs serve as the dedicated phagocytes, while SPMs and the recruited neutrophils and monocytes are major drivers for peritoneal inflammation during bacterial infections (Louwe et al., 2021; Teh et al., 2022; Tilstam et al., 2021; Vega-Perez et al., 2021). The expression of these anti-bacterial molecular signatures, but not proinflammatory genes (i.e. *Nos2*, *Tnf* or *Il6*) were further increased in PAC-deficient LPMs when phagosome-mediated bacterial degradation was enhanced. This is consistent with the finding that *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$ mice not only exhibited a better LPM-mediated bacterial clearance during disease resolution, but also showed mitigated peritoneal inflammation. These results further suggest that the peritoneal bacterial load is the major determinant for peritoneal inflammation, likely by regulating the activation of SPMs and other infiltrated immune cells.

The negative role of PAC in phagosome acidification in macrophages is consistent with a model previously proposed for endosomes that PAC is activated by acidic pH and releases the counterion Cl$^-$ from organelle lumen, thereby preventing further acidification (Osei-Owusu et al., 2021). The increase in phagosomal protease activities in PAC-deficient macrophages is likely due to more efficient phagosomal acidification. Loss of the PAC channel may also lead to a higher Cl$^-$ concentration, which could elevate protease activities as well (Wu et al., 2023). In addition to proteases, the activity of other hydrolytic enzymes in phagosomes, such as glycosidases, lipases, and nucleases, is likely also enhanced in PAC-deficient macrophages. LPS treatment typically

upregulates gene expression. However, PAC expression was downregulated by LPS, which in turn is required for LPS-induced phagosome activation. This provides an elegant mechanism for macrophages to facilitate phagosome maturation in response to bacterial infections, while preserving the housekeeping function of PAC in maintaining endosomal pH homeostasis under normal physiological conditions (Osei-Owusu et al., 2021; Zeziulia et al., 2022). Although macrophages are efficient in killing most bacteria, some pathogens, such as *Mycobacterium* and *Legionella pneumophilia*, can survive and adapt after phagocytosis (Horwitz and Maxfield, 1984; Queval et al., 2017). They achieve this mainly by hijacking the host acidification machinery and preventing phagosomal acidification. In the future, it will be interesting to test if the effector proteins from these bacteria can directly target and, perhaps, activate the PAC channel to interfere with acidification.

PAC deletion presents beneficial effects in enhancing macrophage bactericidal activities against both WT and cpm-resistant *E. coli* infections, leading to reduced lethality in mice. By promoting phagosome-mediated bacterial degradation, the effect of PAC deletion likely extends beyond *E. coli*, to other clinically relevant bacteria, especially those sensitive to phagosome degradation, such as *Streptococcus pneumoniae* (Ip et al., 2010). Our study thus suggests the PAC channel as a potential therapeutical target for the treatment of bacterial infectious diseases. Due to the lack of molecular identity until recently, there is no potent and selective PAC inhibitor available. We previously developed a high-throughput cell-based fluorescence quenching assay for the PAC channel and applied it to successfully identify its underlying gene. This assay will offer a promising avenue for screening the PAC inhibitors in the future, which can be tested for their potential in combating bacterial infections.

Acknowledgments

We thank Dr. Andrea Cox for providing human THP1 cells, Dr. Won Jin Ho and Johns Hopkins Sidney Kimmel Comprehensive Cancer Center (SKCCC) Flow/Mass Cytometry Core for the assistance on cyTOF experiments, as well as Dr. Nathan Archer and members of the Qiu lab for valuable discussion.

Funding

National Institutes of Health grant R35GM124824 (ZQ)

National Institutes of Health grant R01NS118014 (ZQ)

National Institutes of Health grant RF1NS134549 (ZQ)

McKnight Scholar award (ZQ)

Klingenstein-Simon Scholar award (ZQ)

Sloan Research Fellowship in Neuroscience (ZQ)

Karen Toffler Scholar Program (YY)

Author contributions

Conceptualization: HYC, JChu, ZQ

Methodology: HYC, JChu, JChen, YY, SY, NL, KHC, XD

Investigation: HYC., JChu, JChen

Visualization: HYC

Funding acquisition: ZQ

Project acquisition: HYC, JChu, ZQ

Supervision: ZQ

Writing – original draft: HYC, ZQ

Writing – review & editing: HYC, ZQ

Competing interests: The authors declare that they have no competing interests.

Materials and Methods

Contact for reagent and resource sharing

Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Zhaozhu Qiu (zhaozhu@jhmi.edu).

Material availability

PAC KO THP-1 (gRNA1 and gRNA2), PAC overexpressing Raw264.7, and PAC-mCherry expressing Raw264.7 cells were generated in this study. *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$ mice were generated in this study. All will be made available from the lead contact upon request.

Experimental Model and Subject Details

Mice

The use of all mice in the study was approved by the Johns Hopkins University Animal Care and Use Committee, under approved protocol MO22M17. All mice were housed in a specific pathogen free (SPF) environment, under standard 12-hour light/12-hour dark cycle with *ad libitum* access to food and water. PAC$^{F/F}$ mice were generated in the laboratory as previously described(Yang et al., 2019), while *Gsdmd*$^{-/-}$ (C57BL/6N-*Gsdmd*^{em4Fcw}/J), *Cx3cr1*-Cre [B6J.B6N(Cg)-*Cx3cr1*$^{tm1.1(cre)Jung}$/J] and *LysM*-Cre [B6.129P2-*Lyz2*$^{tm1(cre)Ifo}$/J] were purchased from The Jackson Laboratory (strain #032410, #025524 and #004781, respectively). *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$ mice were generated by crossing *Cx3cr1*-Cre or *LysM*-Cre mice with PAC$^{F/F}$ mice, with *Cx3cr1*$^{Cre/WT}$;PAC$^{F/F}$ and *LysM*$^{Cre/WT}$;PAC$^{F/F}$ considered as *Cx3cr1*-PAC$^{-/-}$ and *LysM*-PAC$^{-/-}$, respectively. Adult male or female mice between 8-12 weeks of age were used for experiments as specified with age- and sex-matched littermates as controls.

Bacterial Culture

Wildtype (WT) *Escherichia coli* (*E. coli*), GFP-expressing *E. coli*, and cpm-resistant *E. coli* strains were purchased from American Type Culture Collection (ATCC, #25922, #25922GFP, and #BAA-2469 respectively). All handling of the infectious bacterial strains strictly adhered to instructions provided by the Johns Hopkins Biosafety Office. The use of bacterial strains in the study was approved by Johns Hopkins Institutional Biosafety Committee (IBC).

Bacterial cultures were prepared following the instructions provided by ATCC. Briefly, wildtype *E. coli* was grown in LB broth. *E. coli*-GFP was grown in LB broth supplemented with carbamacilin (100 μg/mL). Cpm-resistant *E. coli* was grown in Trypticase Soy Broth (TSB) supplemented with Imipenem (25 μg/mL). All cultures were incubated aerobically at 37°C overnight. Subsequently, bacteria were subcultured (1:50) in respective fresh media for 4 hours for the determination of colony formation units (CFU) that were used for the subsequent infection experiments.

Cell Culture

Details of BMDM culture have been previously reported (Zeziulia et al., 2022). Briefly, BMDMs were freshly isolated and cultured from adult C57BL6/J mice (8-12 weeks old). Bone marrow was flushed from both femurs and tibiae with a 26-gauge needle syringe into a 50 mL conical tube. Hematopoietic stem cells were collected through 300xg 10 min of centrifugation at 4ºC. BMDMs were differentiated by culturing stem cells in 10 cm petri dishes in L929-differentiation medium or full RPMI 1640 medium (10% HI-FBS, 2mM L-glutamine, 10mM MEM non-essential amino acids, 100 units/mL penicillin, and 100 mg/mL streptomycin) supplemented with mouse recombinant M-CSF (40 ng/mL) for 6 days. L929-differentiation medium was made from 70%

RPMI 1640 medium (20% heat-inactivated fetal bovine serum (HI-FBS), 100 units/mL penicillin, and 100 mg/mL streptomycin) supplemented with 30% L929 mouse fibroblast-conditioned medium. During the differentiation, 2-3 mL of freshly made respective medium was supplemented every 2 days. On day 6, matured BMDMs were then subcultured using macrophage detachment buffer (10mM EDTA in PBS) for following experiments.

Raw264.7 mouse macrophage cell line was purchased through ATCC (TIB-71) and cultured following manufacturer's instructions. Briefly, cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% HI-FBS, 100 units/mL penicillin, and 100 mg/mL streptomycin at 37°C in humidified 95% CO_2 incubator. THP-1 human monocytes, generously provided by Dr. Andrea Cox (Johns Hopkins University), were cultured in full RPMI 1640 medium at 37°C in a humidified 95% CO_2 incubator. THP-1 macrophages were differentiated by treating the cells with 5 ng/mL of PMA for 24 hours, followed by an additional 24-hour resting in RPMI 1640 medium before subsequent experiments.

Method Details

Genotyping and Characterization of Mice

The extraction of genomic DNA from tail samples was performed with Extracta DNA Prep kit (Quanta Bio). Samples were digested with 20 μL of digestion buffer for 30 min at 95ºC, and stopped with 20 μL of stopping buffer. 1 μL of genomic DNA sample was further used as template for PCR using AccuStart II GelTrack PCR SuperMix (Quanta Bio). PCR products were electrophoretically separated with a 2-3% agarose gel and visualized with SYBR safe DNA gel stain (Thermo Fisher Scientific). Genotype was determined using the following primers:

Cx3cr1-Cre:

Common primer: 5'-GCA GGG AAA TCT GAT GCA AG-3'

Mutant primer: 5'-GAC ATT TGC CTT GCT GGA C-3'

Wildtype primer: 5'-CCT CAG TGT GAC GGA GAC AG-3'

LysM-Cre:

Common primer: 5'-AAG GAG GGA CTT GGA GGA TG-3'

Mutant primer: 5'-ACC GGT AAT GCA GGC AAA T-3'

Wildtype primer: 5'-GTC ACT CAC TGC TCC CCT GT-3'

Gsdmd$^{-/-}$:

Forward primer: 5'-ACA CCC CTG AAC CCA TAT CC-3'

Reverse primer: 5'-GCA CTC ACC CCA TTT AGA GC-3'

Casp1$^{-/-}$:

Common primer: 5'-ATG GCA CAC CAC AGA TAT CGG-3'

Mutant primer: 5'-GAG ACA TAT AAG GGA GAA GGG-3'

Wildtype primer: 5'-TGC TAA AGC GCA TGC TCC AGA CTG-3'

LoxP sites at *Pacc1* exon 2:

Forward primer: 5'-GAA GCC AGG CCA TTC TTT TT-3'

Reverse primer: 5'-GCT CAA GGA AAC CAC TGA GG-3'

Peritoneal Bacterial Infection

The CFU of each bacterium freshly grown after 4 hours was determined by LB plate culture. For the peritoneal infection, bacterium was resuspended in sterile PBS. Mice were peritoneally

infected by the injection of 200 µL of bacterial solution with indicated CFU and monitored daily for health and survival following institutional guidance.

Peritoneal Lavage

Mice were euthanized before the procedure. Peritoneal lavage was carried out by injecting 7 mL of sterile PBS into the peritoneal cavity, followed by gentle abdominal massage for homogenization. Afterward, a skin incision was made and a 26-guage syringe was used to collect peritoneal fluid. 5 mL of the injected PBS from each mouse was collected and transferred into a 15 mL conical tube. Centrifugation at 300xg for 5-10 minutes at 4°C was performed to isolate peritoneal cells or obtain peritoneal fluid for subsequent analyses. ACK lysis buffer (Quality Biological) was used for the removal of red blood cells. For peritoneal fluid secretome analyses, 1 mL of PBS was injected, and 0.6-7 mL of the injected fluid was subsequently harvested.

Flow Cytometry (FACS) Analysis

Single cell suspension was prepared from peritoneal lavage or cell culture and resuspend in FACS buffer (2% FBS, 2 mM EDTA in sterile PBS). Live versus Dead cells were stained in PBS using Live/Dead Fixable Aqua Dead Cell Stain (Thermo Fisher Scientific). Fc receptors were then blocked by incubation with TrueStain FcX anti-CD16/32 antibody (BioLegend) at 4°C for 20 minutes, and the cells were subsequently stained with fluorophore-conjugated antibodies against the mouse antigens listed in the Supplemental Material Table, for 20 minutes at 4°C. For primary antibodies without fluorophore, subsequent fluorophore-conjugated secondary antibodies were used with the same staining approach. For the time-traced bacterial clearance analysis, the mean fluorescence intensity (MFI) of GFP within LPM or neutrophils was acquired following specified infection time. FACS analysis of intracellular GBP5 was carried out using Cyto-Fast Fix/Perm

Buffer set (BioLegend), following manufacturer's instructions. Detection of the cell death was performed, after cell surface labeling, using Annexin V and 7-AAD staining (BioLegend), following manufacturer's instructions. Data was acquired on a CytoFLEX LX (Beckman Coulter) and analyzed using FlowJo (BD).

Mouse LPM depletion-adoptive transfer model

Male WT mice of 8-12 weeks old were subjected to LPM depletion with peritoneal injection of 100 uL PBS-loaded or clodronate-loaded liposomes (Liposoma). 12 days post depletion, LPMs isolated from WT, $Pacc1^{-/-}$, or $Gsdmd^{-/-}$ mice were counted, labeled with CellTrace CSFE or CTV dyes (Invitrogen), and adoptively transferred to the peritoneum of LPM-depleted mice. After 24 hours of transfer, mice were peritoneally infected with *E.coli*-GFP to evaluate LPM bactericidal activities and disease resolution. Otherwise, 12 days post depletion or 24 hours after LPM transfer, mouse peritoneal cells were acquired by peritoneal lavage for the FACS analysis determination of cell composition.

Secretome ELISA Analysis

Peritoneal fluid prepared from peritoneal lavage, or LPM culture medium post infection was subjected to LegendPlex multi-analyte ELISA assay (BioLegend) or individual ELISAs following manufacturer's instruction. For LPM supernatants, freshly isolated LPMs were plated at a density of 5×10^5 cells/well in 12-well plates. For *E. coli* infection, bacterium at MOI 20 was simultaneously plated in 12-well plates. Both peritoneal fluids and cell supernatants were collected, centrifugated at 12000xg to remove dead cells and debris before subjected to ELISA assays. Specifically, pre-defined LegendPlex panels of Mouse Cytokine Release Syndrome Panel, Mouse Cytokine Panel 2, and Mouse Proinflammatory Chemokine Panel 2 were selected for analyses.

Additionally, IL-18 duoset ELISA (R&D Systems), IFNβ duoset ELISA (R&D Systems), MIF Legend MAX ELISA (BioLegend), and CXCL1 duoset ELISA (R&D Systems) were utilized for the purpose of comprehensiveness. For the gasdermin D release quantification, Mouse gasdermin D ELISA Kit (Abcam) was used to determine gasdermin D release in LPM supernatant following manufacturer's instruction.

LPM Purification and Isolation

LPMs were isolated from peritoneal fluids by negative selection after immunomagnetic depletion of B cells, SPMs, T cells, and neutrophils with peritoneum mouse macrophage isolation kit (Miltenyi Biotec). After immunomagnetic depletion, LPM preparations had a purity > 90%. For immunoblotting analysis, isolated LPM was immediately lysed in cell lysis buffer. For immunofluorescence analysis, isolated LPM was cultured on coverslips in full RPMI 1640 medium at 37°C for an additional 20 min for adhesion before fixation. For secretome analysis, isolated LPM was immediately infected with *E. coli* in full RPMI 1640 medium at 37°C for 24 hours before supernatant collection.

Gene Transduction and CRISPR Deletion

PAC KO THP-1 cells were generated by CRISPR-Cas9 technology. Guide RNA1 (GGACCGAGAAGACGTTCTTC) and guide RNA 2 (GTGGATCGCTATGATGCCCC) targeting exon 3 of human *PACC1*/*TMEM206* (PAC) was cloned into LentiCRISPR-v2-PuroR plasmid, which was a gift from Dr. F. Zhang (Addgene plasmid #98290). Lentiviral particles containing PAC sgRNA1 or sgRNA2 were packaged using the third-generation lentiviral system and used to infect THP-1 cells. 24 hours after infection, the medium was changed to fresh RPMI culture medium. Cells were then treated with puromycin (1 μg/ml) for 3-5 days to select for

successfully transduced cells. PAC protein expression was validated in puromycin-resistant cells using protein immunoblotting or whole-cell recording for PAC-specific currents.

PAC overexpressing and mCherry-tagged PAC overexpressing Raw264.7 cells were generated using plasmids as previously described (Osei-Owusu et al., 2021) and packaged into lentiviral particles using the third-generation lentiviral system. 24 hours after infection, the medium was changed to fresh DMEM culture medium. Cells were then FACS sorted for mCherry-positive cells. PAC protein expression was validated in sorted cells using whole-cell recording for PAC-specific currents.

In vitro Bacterial Clearance Assay

BMDMs were counted and cultured in 6-well plates at a density of 1.2×10^6 cells per well. Infection of wildtype *E. coli* was conducted at an MOI of 20 for 1 hour at 37°C. Then the medium was changed to DMEM containing 100 μg/mL gentamycin to remove extracellular bacteria. Following 1 or 2 hours of digestion, cells were washed with ice-cold PBS and lysed with 0.3% (vol/vol) Triton X-100 for 5 minutes. Cell lysates containing undigested bacteria were serially diluted with PBS and inoculated on LB plates followed by CFU determination after overnight incubation at 37°C. The uptake of the bacteria was determined by culturing bacteria from cell lysates immediately after 1 hour of infection.

Measurement of Phagosome pH and Protease Activity

Analysis of phagosome pH was measured by a ratiometric method described previously (Osei-Owusu et al., 2021). Dual-labeled zymosan particles were employed for these measurements. PhrodoGreen and AF633-labeled zymosan particles were used to assess pH, while DQGreen BSA and AF633-labeled zymosan particles were employed to evaluate protease activity. The

conjugation of dual-labeled zymosan particles follows the Amine-Reactive Probe Labeling Protocol (Thermo Fisher Scientific). Subsequently, BMDM or Raw264.7 cells were treated with dual-labeled zymosan particles as indicated. Single cell suspensions were obtained from detached cells using macrophage detachment buffer and subjected to FACS analysis. For phagosome pH measurement, singlet cells positive with both PhrodoGreen and AF633 were gated, and the fluorescence intensity was quantified. A standard curve was generated using the Intracellular pH Calibration Buffer kit (Thermo Fisher Scientific) with different pH values (5.5, 6.5, and 7.5) in the presence of 10 mM nigericin and valinomycin to equilibrium the pH with the calibration buffers. The phagocytic index was calculated by subtracting AF633 MFI from baseline autofluorescence value. The PhrodoGreen/AF633 ratio was used to normalize phagocytic uptake and used for eventual ratiometric calculation. For protease activity, the DQGreen BSA/AF633 ratio was used to normalize phagocytic uptake, and the value was used to compare the protease activity between experimental groups.

For pH and protease activity measurements in peritoneal cells, mice were peritoneally injected with either pH or protease dual-labeled zymosans at an MOI of 20 for 30 minutes. Peritoneal lavage was then conducted to acquire peritoneal cells, which were subjected to surface staining. FACS analysis was used to quantify MFI of PhrodoGreen/AF633 or DQGreen BSA/AF633 within gated LPM, and these values were used for further calculations.

Immunostaining and Confocal Microscopy

Immunostaining was performed as previously described (Osei-Owusu et al., 2021). THP-1 macrophages were differentiated on coverslips 48 hours before fixation. Isolated LPM macrophages were incubated on coverslips for 20 minutes for adherence. Peritoneal *E. coli* was collected by centrifugating peritoneal fluid 3000xg 5 minutes onto coverslips. For phagosomal

PAC, Rab5, Rab7, and Lamp1 imaging, cells were fixed with methanol for 10 minutes at -20°C followed by acetone permeabilization 1 minute at -20°C. For EEA1, Rab11, flagellin, IRF1, pSTAT1, and pp65 imaging, cells were fixed in 4% PFA for 20 minutes at room temperature (RT) followed by 0.3% Triton X-100 permeabilization 30 minutes at RT. Cells further underwent blocking by 5% bovine serum albumin (BSA) in PBS for 60 minutes at RT and incubated with indicated antibodies in 5% BSA buffer overnight at 4°C. For surface F4/80 and gasdermin D staining, cells were 4% PFA fixed without permeabilization and blocked with 5% BSA for 60 minutes before primary antibody incubation. After washing with PBS three times, samples were incubated with Alexa Fluor–conjugated secondary antibodies (1:500; Invitrogen) for 1 hour and DAPI (1 μg/mL) 10 minutes at RT. Coverslips were washed with PBS prior to mounting with Aqua-poly Mount (Polysciences, Inc). Mounted samples were cured overnight at RT and imaged. Single plane or Z-stack images were taken under Zeiss LSM900 confocal microscopes with 60x objective lens. Images were analyzed with ImageJ or ZEN software.

Patch Clamp Electrophysiology

Whole-cell patch clamp recordings were performed as previously described (Osei-Owusu et al., 2021). Cells were recorded in an extracellular solution (ECS) containing: 145mM NaCl, 1.5mM $CaCl_2$, 2mM $MgCl_2$, 2mM KCl, 10mM HEPES, and 10mM glucose (pH adjusted to pH 7.3 with NaOH and osmolality was 300–310 mOsm/kg). Acidic ECS with pH 4.6 was made with the same ionic composition without HEPES but with 5mM Na_3-citrate as buffer and the pH was adjusted using citric acid. The acidic solution was perfused using a gravity perfusion system. Recording pipettes (2–4MU) were filled with internal solution containing: 135mM CsCl, 2mM $CaCl_2$, 1mM $MgCl_2$, 4mM MgATP, 0.5mM $Na3$ -GTP, and 5mM EGTA (pH adjusted to 7.2 with CsOH and osmolality was 280–290 mOsm/kg).

All recordings were done at room temperature with MultiClamp 700B amplifier and 1550B digitizer (Molecular Devices). Data acquisition was performed with pClamp 10.7 software (Molecular Device), filtered at 2 kHz and digitized at 10 kHz. Voltage ramp pulses were applied every 5 s from -100 to +100 mV at a holding potential of 0 mV. Voltage step pulses were applied every 3 s for a duration of 1 s, from -100 to +100 mV at a holding potential of 0 mV in a 20-mV increment. All data were analyzed using Clampfit 10.7 and GraphPad Prism software was used for all statistical analyses.

LPM RNA sequencing

Mice were subjected to peritoneal infection for 30 minutes and underwent peritoneal lavage to acquire peritoneal cells. Cells were washed with ice-cold PBS and collected in FACS buffer. Cells further underwent surface staining for LPM gating. Subsequently, Sony MA900 Multi-Application Cell Sorter was used for isolating LPM cells. 3 samples from PBS- and infection-treated mice from control or $Cx3cr1$-PAC$^{-/-}$ mice were collected. Total RNA was extracted using the RNeasy Micro kit (Qiagen). Total RNA quality was determined by the A_{260}/A_{280} and A_{260}/A_{230} as well as Agilent 2100 Bioanalyzer.

RNA sequencing service was carried out by Novogene Company. In brief, libraries were constructed using the NEBNext Ultra II RNA library Prep Kit for Illumina (New England Biolabs) and sequenced on an illumina NovaSeq 6000 System. Reads were mapped to GRCm38/mm10 reference mouse genome with Hisat2 v2.0.5. FeatureCounts v1.5.0-p3 was used to count the reads numbers mapped to each gene. Fragments per kilobase per million mapped reads (FPKM) of each gene was calculated based on the length of the gene and reads count mapped to this gene. Further differential expression analysis, Gene Set Enrichment analysis, Gene Ontology analysis, and

KEGG pathway analysis were carried out by clusterProfiler R package. Data were visualized using GraphPad Prism and NovoMagic (Novogene).

High-Dimensional Mass Cytometry (CyTOF)

Mice were subjected to peritoneal infection for 4 hours and underwent peritoneal lavage to acquire peritoneal cells. Cells were first washed with ice-cold MAXPAR PBS (Fluidigm) then stained with 1 mL of 1 μM Cell-ID cisplatin-^{194}Pt (Fluidigm) for 3 minutes at RT to exclude dead cells. Next, cells were prepared and stained following Maxpar Cell Surface Staining with Fresh Fix protocol (Fluidigm). In brief, cells were Fc blocked and stained with antibody panel as provided in the Supplemental Material Table for 30 minutes at RT. A fresh fix was then conducted with incubating cells with freshly made 1.6% formaldehyde solution (Thermo Fisher Scientific) for 10 minutes at RT. Lastly, cells were transferred to intercalation solution (125 nM of Cell-ID intercalator-Ir in Maxpar Fix and Perm Buffer) before submitted to Johns Hopkins SKCCC Flow/Mass Cytometry Core for data acquisition. Of note, 168Er conjugated Tim-4 and 156Gd conjugated SIRPα antibodies were made from conjugating isotopes to Tim-4 and SIRPα antibodies (BioLegend) using Maxpar X8 antibody labeling kits (Fluidigm).

OMIQ analysis and workflow

cyTOF FCS files acquired and exported from the Johns Hopkins SKCCC Flow/Mass Cytometry Core were uploaded to OMIQ. Live, singlet and CD45-positive cells were gated before subsampling into a maximum equal distribution across groups. After subsampling, dimensional reduction methods (tSNE, opt-SNE, and UMAP) were performed using default parameters, followed by PhenoGraph, SPADE, and FlowSOM clustering. Data was further analyzed with EdgeR to determine significantly different clusters, or by generating heatmaps with Euclidean

clustering. OMIQ was also used for performing manual gating strategies for identified populations and the quantification of population sizes. Data were visualized using OMIQ or GraphPad Prism.

Phagosome proteome and database gene expression analysis

The phagosome proteome dataset was downloaded from ProteomeXchange (PXD001293). The list of proteins was curated by selecting transmembrane proteins from the reported proteome before filtering out the unknown proteins or proteins that have been reported with substrates other than ions. Additionally, proteins that were only detected in either BMDM or Raw264.7 cells or with zero signal intensity in any samples were also excluded. For the GeneCards database analysis, each of proteins within the phagosome proteome list was surveyed on the GeneCards database with corresponding gene for the subcellular localization confidence. Genes with confidence above 3 on endosomes or lysosomes were used for the following analysis. Gene expression RNA-Seq values for the Immunological Genome Project (Heng et al., 2008) (ImmGen) was downloaded from the online portal. The expression values were used for the subsequent z-score calculation across various cell types or the direct comparison between genes. Dataset for GSE134364 or GSE94916 was downloaded from the NCBI Gene Expression Omnibus (GEO). The expression values were acquired from GEO2R portal and were used for the statistical analysis.

Inflammasome Activation

Freshly isolated LPMs were plated at a density of 5 x 10^5 cells/well in 12-well plates. For *E. coli* infection, bacterium at MOI 20 was simultaneously plated in 12-well plates. For ATP and nigericin activation, cells were activated by LPS (1 µg/mL) for 4 hours upon plating before ATP (4 mM) or nigericin (5 µM) treatment for 30 minutes. Subsequently, cell supernatants were collected,

centrifugated at 12000xg to remove dead cells and debris, and analyzed for IL-1β, IL-18, or LDH release as indicated.

E64d phagolysome cathepsin inhibition

E64d cathepsin inhibitor was acquired from Selleck Chemicals. For *in vivo* E64d inhibition in LPM cells, E64d (0.15 mg/kg) in 100 μL PBS was peritoneally injected 30 minutes before *E. coli* injection. For *in vitro* E64d inhibition, isolated LPM was pretreated with 5μM of E64d for 30 minutes before subsequent *E. coli* infection.

Quantitative Real-Time PCR

Total RNA was isolated using TRIzol reagent (Life Technologies). 1-2 μg of total RNA was used to generate the cDNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). The reaction was run in QuantStudio 6 Real-Time PCR Systems using 0.2 μl of cDNA in a 15 μl reaction. For the amplification of PAC mRNA, IDT PrimeTime Predesigned qPCR probes targeting mouse *Tmem206* exon 1-2 were used in combination with PerfeCTa qPCR SuperMix (Quanta Bio) according to manufacturer's instruction. Primers targeting other genes were provided in the Supplemental Material Table and used in combination with SYBR green master mix (Applied Biosystems) according to manufacturer's instruction. The quantification was collected as CT value. Calibrations and normalizations were performed using the $2^{-\Delta\Delta CT}$ method. *Gapdh* or *Actb* was used as the reference gene.

Protein Immunoblotting

Cytosolic cell extracts were prepared using hypotonic buffer to exclude organelles and nucleus (250 mM sucrose, 20 mM HEPES, 10 mM KCl, 1.5 mM $MgCl_2$, 1 mM EGTA, 1 mM EDTA, 1 mM

Pefabloc). Cytosolic extracts were then precipitated by adding trichloroacetic acid (final concentration 5%). After 10 min on ice, the proteins were pelleted by centrifugation at $20,800 \times g$ for 15 min. The pellets were then resuspended in RIPA buffer (Cell Signaling Technology). Whole cell lysates of THP-1 cell or isolated LPMs were prepared in RIPA buffer with protease inhibitor cocktails (Roche). Lysates in RIPA buffer were cleared of debris by centrifugation at 12000xg for 10 minutes at 4°C and the concentration of protein lysates was determined by BCA assay kit (Thermo Fisher Scientific). Lysates were then collected and made into same concentrations with water and Blot 4x LDS sample buffer (Invitrogen) before boiling at 75°C for 10 minutes prior to SDS-PAGE electrophoresis. Novex 4-20% gradient gel (Invitrogen) was used for electrophoretic separation according to manufacturer's instruction. For subsequent antibody detection, nitrocellulose membranes were used for protein transfer. Primary antibodies were made as indicated by manufacturers and incubated with nitrocellulose membranes overnight at 4°C. Following primary antibody incubation, samples were incubated with HRP-conjugated anti-mouse, anti-rabbit, or anti-rat secondary antibody (Proteintech) for 1 hour at RT for the visualization.

Statistics

All data were analyzed using GraphPad Prism software. Images were analyzed with ImageJ (U.S. National Institutes of Health) and ZEN software (ZEISS). Data were presented as mean with standard error of the mean (SEM) as indicated in figure legends. Survival curve data was presented as a Kaplan-Maier plot with a log-rank (Mantel-Cox) test used to compare susceptibility between different groups. Bar plots, dot plots, donut graphs, survival plots, volcano plots, and heatmaps were plotted using GraphPad Prism. FACS dot plots, histograms, FlowSOM trees, SPADE trees, and UMAPs were plotted using FlowJo (BD) or OMIQ. Graphical abstract and illustrations were made using Biorender.com. The sample sizes were indicated in figures and legends. Statistical

significance was determined using t-tests (two-tailed), one-way ANOVA, or two-way ANOVA followed by multiple comparison tests. A *p* value less than 0.05 was considered statistically significant.

Reference

Abdullah, Z., and P.A. Knolle. 2014. Scaling of immune responses against intracellular bacterial infection. *EMBO J* 33:2283-2294.

Andrade, W.A., A. Firon, T. Schmidt, V. Hornung, K.A. Fitzgerald, E.A. Kurt-Jones, P. Trieu-Cuot, D.T. Golenbock, and P.A. Kaminski. 2016. Group B Streptococcus Degrades Cyclic-di-AMP to Modulate STING-Dependent Type I Interferon Production. *Cell Host Microbe* 20:49-59.

Barnett, K.C., S. Li, K. Liang, and J.P. Ting. 2023. A 360 degrees view of the inflammasome: Mechanisms of activation, cell death, and diseases. *Cell* 186:2288-2312.

Beertsen, W., M. Willenborg, V. Everts, A. Zirogianni, R. Podschun, B. Schroder, E.L. Eskelinen, and P. Saftig. 2008. Impaired phagosomal maturation in neutrophils leads to periodontitis in lysosomal-associated membrane protein-2 knockout mice. *J Immunol* 180:475-482.

Bertrams, W., K. Griss, M. Han, K. Seidel, A. Klemmer, A. Sittka-Stark, S. Hippenstiel, N. Suttorp, F. Finkernagel, J. Wilhelm, T. Greulich, C.F. Vogelmeier, J. Vera, and B. Schmeck. 2020. Transcriptional analysis identifies potential biomarkers and molecular regulators in pneumonia and COPD exacerbation. *Sci Rep* 10:241.

Brun-Buisson, C., F. Doyon, J. Carlet, P. Dellamonica, F. Gouin, A. Lepoutre, J.C. Mercier, G. Offenstadt, and B. Regnier. 1995. Incidence, risk factors, and outcome of severe sepsis and septic shock in adults. A multicenter prospective study in intensive care units. French ICU Group for Severe Sepsis. *JAMA* 274:968-974.

Chan, J.K., J. Roth, J.J. Oppenheim, K.J. Tracey, T. Vogl, M. Feldmann, N. Horwood, and J. Nanchahal. 2012. Alarmins: awaiting a clinical response. *J Clin Invest* 122:2711-2719.

Clausen, B.E., C. Burkhardt, W. Reith, R. Renkawitz, and I. Forster. 1999. Conditional gene targeting in macrophages and granulocytes using LysMcre mice. *Transgenic Res* 8:265-277.

Di, A., M.E. Brown, L.V. Deriy, C. Li, F.L. Szeto, Y. Chen, P. Huang, J. Tong, A.P. Naren, V. Bindokas, H.C. Palfrey, and D.J. Nelson. 2006. CFTR regulates phagosome acidification in macrophages and alters bactericidal activity. *Nat Cell Biol* 8:933-944.

Ding, J., K. Wang, W. Liu, Y. She, Q. Sun, J. Shi, H. Sun, D.C. Wang, and F. Shao. 2016. Pore-forming activity and structural autoinhibition of the gasdermin family. *Nature* 535:111-116.

Fisch, D., M.M. Pfleiderer, E. Anastasakou, G.M. Mackie, F. Wendt, X. Liu, B. Clough, S. Lara-Reyna, V. Encheva, A.P. Snijders, H. Bando, M. Yamamoto, A.D. Beggs, J. Mercer, A.R. Shenoy, B. Wollscheid, K.M. Maslowski, W.P. Galej, and E.M. Frickel. 2023. PIM1 controls GBP1 activity to limit self-damage and to guard against pathogen infection. *Science* 382:eadg2253.

Flannagan, R.S., G. Cosio, and S. Grinstein. 2009. Antimicrobial mechanisms of phagocytes and bacterial evasion strategies. *Nat Rev Microbiol* 7:355-366.

Goodridge, H.S., C.N. Reyes, C.A. Becker, T.R. Katsumoto, J. Ma, A.J. Wolf, N. Bose, A.S. Chan, A.S. Magee, M.E. Danielson, A. Weiss, J.P. Vasilakos, and D.M. Underhill. 2011. Activation of the innate immune receptor Dectin-1 upon formation of a 'phagocytic synapse'. *Nature* 472:471-475.

Graves, A.R., P.K. Curran, C.L. Smith, and J.A. Mindell. 2008. The Cl-/H+ antiporter ClC-7 is the primary chloride permeation pathway in lysosomes. *Nature* 453:788-792.

Guo, H., J.B. Callaway, and J.P. Ting. 2015a. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med* 21:677-687.

Guo, M., A. Hartlova, B.D. Dill, A.R. Prescott, M. Gierlinski, and M. Trost. 2015b. High-resolution quantitative proteome analysis reveals substantial differences between phagosomes of RAW 264.7 and bone marrow derived macrophages. *Proteomics* 15:3169-3174.

Heng, T.S., M.W. Painter, and C. Immunological Genome Project. 2008. The Immunological Genome Project: networks of gene expression in immune cells. *Nat Immunol* 9:1091-1094.

Honda, K., A. Takaoka, and T. Taniguchi. 2006. Type I interferon [corrected] gene induction by the interferon regulatory factor family of transcription factors. *Immunity* 25:349-360.

Hornung, V., R. Hartmann, A. Ablasser, and K.P. Hopfner. 2014. OAS proteins and cGAS: unifying concepts in sensing and responding to cytosolic nucleic acids. *Nat Rev Immunol* 14:521-528.

Horwitz, M.A., and F.R. Maxfield. 1984. Legionella pneumophila inhibits acidification of its phagosome in human monocytes. *J Cell Biol* 99:1936-1943.

Husebye, H., M.H. Aune, J. Stenvik, E. Samstad, F. Skjeldal, O. Halaas, N.J. Nilsen, H. Stenmark, E. Latz, E. Lien, T.E. Mollnes, O. Bakke, and T. Espevik. 2010. The Rab11a GTPase controls Toll-like receptor 4-induced activation of interferon regulatory factor-3 on phagosomes. *Immunity* 33:583-596.

Ince, C., B. Thio, B. van Duijn, J.T. van Dissel, D.L. Ypey, and P.C. Leijh. 1987. Intracellular K+, Na+ and Cl- concentrations and membrane potential in human monocytes. *Biochim Biophys Acta* 905:195-204.

Ip, W.K., A. Sokolovska, G.M. Charriere, L. Boyer, S. Dejardin, M.P. Cappillino, L.M. Yantosca, K. Takahashi, K.J. Moore, A. Lacy-Hulbert, and L.M. Stuart. 2010. Phagocytosis and phagosome acidification are required for pathogen processing and MyD88-dependent responses to Staphylococcus aureus. *J Immunol* 184:7071-7081.

Jorgensen, I., Y. Zhang, B.A. Krantz, and E.A. Miao. 2016. Pyroptosis triggers pore-induced intracellular traps (PITs) that capture bacteria and lead to their clearance by efferocytosis. *J Exp Med* 213:2113-2128.

Kinchen, J.M., and K.S. Ravichandran. 2008. Phagosome maturation: going through the acid test. *Nat Rev Mol Cell Biol* 9:781-795.

Kuchitsu, Y., K. Mukai, R. Uematsu, Y. Takaada, A. Shinojima, R. Shindo, T. Shoji, S. Hamano, E. Ogawa, R. Sato, K. Miyake, A. Kato, Y. Kawaguchi, M. Nishitani-Isa, K. Izawa, R. Nishikomori, T. Yasumi, T. Suzuki, N. Dohmae, T. Uemura, G.N. Barber, H. Arai, S. Waguri, and T. Taguchi. 2023. STING signalling is terminated through ESCRT-dependent microautophagy of vesicles originating from recycling endosomes. *Nat Cell Biol* 25:453-466.

Leisching, G., V. Cole, A.T. Ali, and B. Baker. 2019. OAS1, OAS2 and OAS3 restrict intracellular M. tb replication and enhance cytokine secretion. *Int J Infect Dis* 80S:S77-S84.

Li, X., T. Wang, Z. Zhao, and S.A. Weinman. 2002. The ClC-3 chloride channel promotes acidification of lysosomes in CHO-K1 and Huh-7 cells. *Am J Physiol Cell Physiol* 282:C1483-1491.

Liu, X., Z. Zhang, J. Ruan, Y. Pan, V.G. Magupalli, H. Wu, and J. Lieberman. 2016. Inflammasome-activated gasdermin D causes pyroptosis by forming membrane pores. *Nature* 535:153-158.

Louwe, P.A., L. Badiola Gomez, H. Webster, G. Perona-Wright, C.C. Bain, S.J. Forbes, and S.J. Jenkins. 2021. Recruited macrophages that colonize the post-inflammatory peritoneal niche convert into functionally divergent resident cells. *Nat Commun* 12:1770.

Lukacs, G.L., O.D. Rotstein, and S. Grinstein. 1990. Phagosomal acidification is mediated by a vacuolar-type H(+)-ATPase in murine macrophages. *J Biol Chem* 265:21099-21107.

Man, S.M., R. Karki, M. Sasai, D.E. Place, S. Kesavardhana, J. Temirov, S. Frase, Q. Zhu, R.K.S. Malireddi, T. Kuriakose, J.L. Peters, G. Neale, S.A. Brown, M. Yamamoto, and T.D. Kanneganti. 2016. IRGB10 Liberates Bacterial Ligands for Sensing by the AIM2 and Caspase-11-NLRP3 Inflammasomes. *Cell* 167:382-396 e317.

Manzanillo, P.S., M.U. Shiloh, D.A. Portnoy, and J.S. Cox. 2012. Mycobacterium tuberculosis activates the DNA-dependent cytosolic surveillance pathway within macrophages. *Cell Host Microbe* 11:469-480.

Moreland, J.G., A.P. Davis, G. Bailey, W.M. Nauseef, and F.S. Lamb. 2006. Anion channels, including ClC-3, are required for normal neutrophil oxidative function, phagocytosis, and transendothelial migration. *J Biol Chem* 281:12277-12288.

Moretti, J., S. Roy, D. Bozec, J. Martinez, J.R. Chapman, B. Ueberheide, D.W. Lamming, Z.J. Chen, T. Horng, G. Yeretssian, D.R. Green, and J.M. Blander. 2017. STING Senses Microbial Viability to Orchestrate Stress-Mediated Autophagy of the Endoplasmic Reticulum. *Cell* 171:809-823 e813.

Nair-Gupta, P., A. Baccarini, N. Tung, F. Seyffer, O. Florey, Y. Huang, M. Banerjee, M. Overholtzer, P.A. Roche, R. Tampe, B.D. Brown, D. Amsen, S.W. Whiteheart, and J.M. Blander. 2014. TLR signals induce phagosomal MHC-I delivery from the endosomal recycling compartment to allow cross-presentation. *Cell* 158:506-521.

Osei-Owusu, J., J. Yang, K.H. Leung, Z. Ruan, W. Lu, Y. Krishnan, and Z. Qiu. 2021. Proton-activated chloride channel PAC regulates endosomal acidification and transferrin receptor-mediated endocytosis. *Cell Rep* 34:108683.

Paidimuddala, B., J. Cao, G. Nash, Q. Xie, H. Wu, and L. Zhang. 2023. Mechanism of NAIP-NLRC4 inflammasome activation revealed by cryo-EM structure of unliganded NAIP5. *Nat Struct Mol Biol* 30:159-166.

Poltorak, A., X. He, I. Smirnova, M.Y. Liu, C. Van Huffel, X. Du, D. Birdwell, E. Alejos, M. Silva, C. Galanos, M. Freudenberg, P. Ricciardi-Castagnoli, B. Layton, and B. Beutler. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science* 282:2085-2088.

Queval, C.J., O.R. Song, J.P. Carralot, J.M. Saliou, A. Bongiovanni, G. Deloison, N. Deboosere, S. Jouny, R. Iantomasi, V. Delorme, A.S. Debrie, S.J. Park, J.C. Gouveia, S. Tomavo, R. Brosch, A. Yoshimura, E. Yeramian, and P. Brodin. 2017. Mycobacterium tuberculosis Controls Phagosomal Acidification by Targeting CISH-Mediated Signaling. *Cell Rep* 20:3188-3198.

Reid-Yu, S.A., B.R. Tuinema, C.N. Small, L. Xing, and B.K. Coombes. 2015. CXCL9 contributes to antimicrobial protection of the gut during citrobacter rodentium infection independent of chemokine-receptor signaling. *PLoS Pathog* 11:e1004648.

Ruangkiattikul, N., A. Nerlich, K. Abdissa, S. Lienenklaus, A. Suwandi, N. Janze, K. Laarmann, J. Spanier, U. Kalinke, S. Weiss, and R. Goethe. 2017. cGAS-STING-TBK1-IRF3/7 induced interferon-beta contributes to the clearing of non tuberculous mycobacterial infection in mice. *Virulence* 8:1303-1315.

Scicluna, B.P., F. Uhel, L.A. van Vught, M.A. Wiewel, A.J. Hoogendijk, I. Baessman, M. Franitza, P. Nurnberg, J. Horn, O.L. Cremer, M.J. Bonten, M.J. Schultz, T. van der Poll, D. Molecular, and c. Risk Stratification in Sepsis. 2020. The leukocyte non-coding RNA landscape in critically ill patients with sepsis. *Elife* 9:

Stephanie A. Ragland, J.C.K. 2022. Phagosome-mediated activation of STING during diverse bacterial infections. *BioRxiv* 2022.10.28.514268.

Teh, Y.C., M.Y. Chooi, D. Liu, I. Kwok, G.C. Lai, L. Ayub Ow Yong, M. Ng, J.L.Y. Li, Y. Tan, M. Evrard, L. Tan, K.H. Liong, K. Leong, C.C. Goh, A.Y.J. Chan, N.B. Shadan, C.K. Mantri, Y.Y. Hwang, H. Cheng, T. Cheng, W. Yu, H.L. Tey, A. Larbi, A. St John, V. Angeli, C. Ruedl, B. Lee, F. Ginhoux, S.L. Chen, L.G. Ng, J.L. Ding, and S.Z. Chong. 2022. Transitional premonocytes emerge in the periphery for host defense against bacterial infections. *Sci Adv* 8:eabj4641.

Tilstam, P.V., W. Schulte, T. Holowka, B.S. Kim, J. Nouws, M. Sauler, M. Piecychna, G. Pantouris, E. Lolis, L. Leng, J. Bernhagen, G. Fingerle-Rowson, and R. Bucala. 2021. MIF but not MIF-2 recruits inflammatory macrophages in an experimental polymicrobial sepsis model. *J Clin Invest* 131:

Ullrich, F., S. Blin, K. Lazarow, T. Daubitz, J.P. von Kries, and T.J. Jentsch. 2019. Identification of TMEM206 proteins as pore of PAORAC/ASOR acid-sensitive chloride channels. *Elife* 8:

Underhill, D.M., and H.S. Goodridge. 2012. Information processing during phagocytosis. *Nat Rev Immunol* 12:492-502.

van Rooijen, N., and A. Sanders. 1997. Elimination, blocking, and activation of macrophages: three of a kind? *J Leukoc Biol* 62:702-709.

Vega-Perez, A., L.H. Villarrubia, C. Godio, A. Gutierrez-Gonzalez, L. Feo-Lucas, M. Ferriz, N. Martinez-Puente, J. Alcain, A. Mora, G. Sabio, M. Lopez-Bravo, and C. Ardavin. 2021. Resident macrophage-dependent immune cell scaffolds drive anti-bacterial defense in the peritoneal cavity. *Immunity* 54:2578-2594 e2575.

Wolf, A.J., C.N. Reyes, W. Liang, C. Becker, K. Shimada, M.L. Wheeler, H.C. Cho, N.I. Popescu, K.M. Coggeshall, M. Arditi, and D.M. Underhill. 2016. Hexokinase Is an Innate Immune Receptor for the Detection of Bacterial Peptidoglycan. *Cell* 166:624-636.

Wong, C.O., S. Gregory, H. Hu, Y. Chao, V.E. Sepulveda, Y. He, D. Li-Kroeger, W.E. Goldman, H.J. Bellen, and K. Venkatachalam. 2017. Lysosomal Degradation Is Required for Sustained Phagocytosis of Bacteria by Macrophages. *Cell Host Microbe* 21:719-730 e716.

Wu, J.Z., M. Zeziulia, W. Kwon, T.J. Jentsch, S. Grinstein, and S.A. Freeman. 2023. ClC-7 drives intraphagosomal chloride accumulation to support hydrolase activity and phagosome resolution. *J Cell Biol* 222:

Yang, J., J. Chen, M. Del Carmen Vitery, J. Osei-Owusu, J. Chu, H. Yu, S. Sun, and Z. Qiu. 2019. PAC, an evolutionarily conserved membrane protein, is a proton-activated chloride channel. *Science* 364:395-399.

Yona, S., K.W. Kim, Y. Wolf, A. Mildner, D. Varol, M. Breker, D. Strauss-Ayali, S. Viukov, M. Guilliams, A. Misharin, D.A. Hume, H. Perlman, B. Malissen, E. Zelzer, and S. Jung. 2013. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity* 38:79-91.

Zeziulia, M., S. Blin, F.W. Schmitt, M. Lehmann, and T.J. Jentsch. 2022. Proton-gated anion transport governs macropinosome shrinkage. *Nat Cell Biol* 24:885-895.

Figure 1. PAC is a phagosome chloride channel mediating macrophage bactericidal activity.

A, Donut plots depicting ion channels and transporters presented within phagosome proteome (ProteomeXchange: PXD001293).

B, Heatmap depicting relative expression of genes filtered by endo-lysosome localization at GeneCards database among different immune cells analyzed in ImmGen.

C, Quantification of gene expression level of genes from (**B**) within splenic macrophages analyzed in ImmGen.

D-E, Representative images of PMA-treated THP1 cell treated with PhrodoGreen-labeled zymosan bioparticles (**D**) or GFP-expressing *E. coli* (**E**) (green) and stained with PAC (red). Data shown as representative of at least 20 cells. Scale bar as indicated.

F, Quantification of fluorescence intensities from live cell imaging of PAC-mCherry at 9 min in Extended Data Fig. 1e. Data shown as representative of at least 20 cells.

G, Quantification of bacterial load of BMDM bacterial clearance assay. All conditions and genotypes have n = 8 independent experiments.

H, Schematic illustration of peritoneal infection model.

I, Gating strategy of neutrophils, LPM, and SPM in the peritoneal fluid.

J, Representative I-V relationship (top) and current density (bottom) of whole-cell PAC current induced by pH 4.6 in FACS sorted control or *Cx3cr1*-PAC$^{-/-}$ LPMs. control, n = 10 cells; *Cx3cr1*-PAC$^{-/-}$, n = 9 cells.

K, Histogram (left) and quantification (right) of *in vivo* bacterial clearance assay from control, *Cx3cr1*-PAC$^{-/-}$ (top), and *LysM*-PAC$^{-/-}$ (bottom) mice. For histograms, *E. coli*-GFP fluorescence in LPM was plotted at indicated time post infection. At least 500,000 cells were collected per condition. For quantification, mean fluorescence intensity (MFI) in LPM was quantified at indicated time post infection. Each timepoint or genotype has at least n > 4 mice from independent experiments.

All control groups represent *Pacc1$^{F/F}$* genotype. Data are reported as mean ± SEM. Unpaired t-test for **G**. Two-way ANOVA with Sidak's test for **K**. **p < 0.01, ****p < 0.0001.

Figure 2. PAC downregulation is required for LPS-induced phagosome maturation.

A, Quantification of *PACC1* expression in healthy and septic PBMCs (left) or healthy and pneumonia PBMCs (right) analyzed in GSE134364 (left) or GSE94916 (right).

B, *Pacc1* mRNA expression in BMDMs post 0-6 hours of LPS (100 ng/mL) treatment. Data was collected from n = 3 independent experiments.

C, Representative I-V relationship (left) and current density (right) of whole-cell PAC current induced by pH 4.6 in BMDMs with PBS or 4-hour LPS (100 ng/mL) treatment. PBS: n = 14 cells from 3 mice. LPS: n = 9 cells from 3 mice.

D, Quantification of phagosome pH in control or *PACC1*$^{-/-}$ THP1 cells (KO 1 and KO 2, with CRISPR gRNA1 or gRNA2 transduced, respectively). Cells were measured 30 min post internalization. Each genotype has n = 6 independent experiments.

E, Quantification of phagocytosis activity in control or *Pacc1*$^{-/-}$ BMDMs. Cells were challenged with zymosan for 0-120 min. Each timepoint or genotype has n > 4 independent experiments.

F, Quantification of phagosome pH in control or *Pacc1*$^{-/-}$ BMDMs. Cells were challenged with zymosan for 0-120 min. For treatments longer than 30 min, zymosan was washed out to prevent continuous phagocytosis. Each timepoint or genotype has n > 4 independent experiments.

G, Quantification of steady-state phagosome pH (left) or protease activity (right) from LPMs within peritoneal fluids after 30 min of injection. Each condition has at least 500,000 cells from n = 3 independent experiments.

H-I, Quantification of steady-state and LPS (20 ng/mL) primed phagosome pH (**H**) or protease activity (**I**). Each condition has at least 20,000 cells from n > 3 independent experiments.

J, Representative I-V relationship (left) and current density (right) of whole-cell PAC current induced by pH 4.6 in vector or PAC overexpressing (PAC) Raw264.7 cells. Vector, n = 5 cells; PAC, n = 4 cells.

K-L, Quantification of steady-state and LPS (20 ng/mL) primed phagosome pH (**K**) or protease activity (**L**). Each condition has at least 20,000 cells from n > 3 independent experiments.

All control groups represent *Pacc1*$^{F/F}$ genotype. Data are reported as mean ± SEM. Unpaired t-test for **A, C, G**. One-way ANOVA with Fisher's LSD test for **B, D, F**. Two-way ANOVA with Sidak's test for **H, I, K, L**. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 3. Enhanced phagosome bacterial degradation promotes interferon responses in PAC-deficient LPMs.

A, GSEA signatures identified in RNA-seq analysis. NES, normalized enrichment score.

B, List of highlighted GO terms and KEGG pathways identified in RNA-seq analysis.

C, Heatmap depicting highlighted genes within interferon responses within GO terms in (**B**).

D, Quantification of secreted IFNα and IFNβ from isolated control or *Pacc1*$^{-/-}$ LPMs after *E. coli* infection. Each genotype has cells from n = 4 mice in independent experiments.

E, Immunoblotting of LPM expression of pSTAT1 (Y701), STAT1, IRF1, and GAPDH. LPMs from 2 mice after 60 min of peritoneal infection were pooled and used as 1 sample. Each condition and genotype have n = 3 independent experiments.

F, Representative images (left) and quantification (right) of LPMs after 60 min of peritoneal infection staining IRF1 or pSTAT1 (Y701) with DAPI. Quantification represents the nucleus to cytosol ratio. Control LPM for IRF1, n = 45 cells; *Cx3cr1*-PAC$^{-/-}$ LPM for IRF1, n = 53 cells; control LPM for pSTAT1, n = 90 cells; *Cx3cr1*-PAC$^{-/-}$ LPM for pSTAT1, n = 74 cells.

G, Immunoblotting of LPM expression of GBP2, GBP5, OAS1, OAS2, OAS3, and GAPDH. Isolated LPMs from 2 mice after 60 min of peritoneal infection were pooled and used as 1 sample. Each condition and genotype have n = 3 independent experiments.

H, Immunoblotting of LPM expression of pSTING (S365), STING, pIRF3 (S396), IRF3 and GAPDH. LPMs from 2 mice after 15 min of peritoneal infection were pooled and used as 1 sample. Each condition and genotype have n = 3 independent experiments.

I, Quantification of *in vivo* bacterial clearance assay from control and *Cx3cr1*-PAC$^{-/-}$ mice pretreated with E64d (0.15 mg/kg) before peritoneal infection. Mean fluorescence intensity (MFI) of *E. coli*-GFP in LPM was quantified from mice after 60 min of infection. Each treatment or genotype has at least n > 5 mice from independent experiments.

J, Immunoblotting of LPM expression of pSTING (S365), STING, pIRF3 (S396), IRF3 and GAPDH. LPMs from 2 mice pretreated with E64d (0.15 mg/kg) following 15 min of peritoneal infection were pooled and used as 1 sample. Each condition and genotype have n = 3 independent experiments.

K, Quantification of secreted IFNβ from isolated control or *Pacc1*$^{-/-}$ LPMs pretreated with E64d (5 μM) following *E. coli* infection. Each genotype has cells from n = 4 mice in independent experiments. N.D, not detected.

All control groups represent *Pacc1*$^{F/F}$ genotype. Data are reported as mean ± SEM. Unpaired t-test for **D**, **F**. Two-way ANOVA with Sidak's test for **I**, **K**. *p < 0.05, **p < 0.01, ****p < 0.0001.

Figure 4. Augmented phagosome bacterial degradation facilitates inflammasome activation and pyroptosis in PAC-deficient LPMs.

A-B, Quantification of secreted IL-1β and IL-18 (**A**) or LDH release (**B**) from isolated control or *Pacc1*$^{-/-}$ LPMs after *E. coli* infection. Each genotype has n = 5 mice in independent experiments.

C, FACS analysis (left) and quantification (right) of annexin-V$^+$ and 7-AAD$^+$ LPMs within peritoneal fluids after 30 min of infection. Each genotype has at least n > 8 mice in independent experiments.

D, Representative images (left) and quantification (right) of LPMs after peritoneal infection staining gasdermin D (white) and F4/80 (red) with DAPI. 15 min control, n = 98 cells; 15 min *Cx3cr1*-PAC$^{-/-}$, n = 109; 60 min control, n = 99 cells, 60 min *Cx3cr1*-PAC$^{-/-}$, n = 108.

E, Immunoblotting of LPM expression of gasdermin D, Caspase 1, and GAPDH. LPMs from 4 mice after 30 min of peritoneal infection were pooled and used as 1 sample. Each condition and genotype have n = 3 independent experiments.

F, Quantification of secreted IL-1β from isolated control or *Pacc1*$^{-/-}$ LPMs pretreated with E64d (5 µM) following *E. coli* infection. Each genotype has n = 4 mice in independent experiments.

G-H, Immunoblotting of LPM cytosolic expression of flagellin and GAPDH. LPMs from 2 mice (**G**) or mice pretreated with E64d (0.15 mg/kg) (**H**) before 15 min of peritoneal infection were pooled and used as 1 sample. Each condition and genotype have n = 3 independent experiments.

I, Quantification of annexin-V$^+$ and 7-AAD$^+$ LPMs pretreated with E64d (0.15 mg/kg) within peritoneal fluids after 60 min of *E. coli* infection. Each genotype has at least n > 6 mice in independent experiments.

J, Quantification of bacterial load cultured with antibiotic-free supernatants from *E. coli*-infected control, *Pacc1*$^{-/-}$, or *Gsdmd*$^{-/-}$ LPM for 6 h at 37C. Blank, antibiotic-free RPMI medium.

K, Quantification of secreted gasdermin D from control or *Cx3cr1*-PAC$^{-/-}$ mice after 60 min of peritoneal infection. Each genotype has n = 5 mice in independent experiments.

L, Representative images (left) and quantification (right) of isolated *E. coli* after 60 min of peritoneal infection staining gasdermin D (green) and DAPI. Quantification represents the percentage of gasdermin D punta-positive bacteria from each mouse. Each genotype has at least n > 4 mice in independent experiments.

M, Schematic illustration of depletion-adoptive transfer model.

N-O, Quantification of intra-LPM GFP intensity (**N**) and peritoneal bacterial load (**O**) within recipient mice transferred with control or *Pacc1*$^{-/-}$LPM. Each genotype has n > 6 mice in independent experiments.

P-R, Quantification of LPM cellularity (**P**), intra-LPM GFP intensity (**Q**) and peritoneal bacterial load (**R**) within recipient mice transferred with control or *Gsdmd*$^{-/-}$LPM. Each genotype has n = 6 mice in independent experiments.

Control represents *Pacc1*$^{F/F}$ genotype in **A-O** or C57BL/6 wildtype in **P-R**. Data are reported as mean ± SEM. Unpaired t-test for **A, B, C, K, L, N, O, P, Q, R**. Two-way ANOVA with Sidak's test for **D**, **F, I**. One-way ANOVA with Fisher's LSD test for **J**. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 5. PAC deletion in macrophages mitigates peritoneal inflammation after peritoneal _E. coli_ infection.

A, Volcano plot depicting differentially secreted proteins in _Cx3cr1_-PAC$^{-/-}$ (left) or _LysM_-PAC$^{-/-}$ (right) peritoneal fluids after 4 hours of peritoneal infection. Highlighted proteins represent top differentially secreted proteins with p < 0.05 and average log-transformed fold-change > 0.25.

B, Quantification of highlighted proteins in (**A**), excluding proteins with absolute concentration below 100 pg/mL. Each genotype has n = 6 mice in independent experiments.

C, UMAP color-coded for immune cell types based on the expression of cell-type enriched surface markers before and after infection. NK, natural killer cells. DCs, dendritic cells.

D, FlowSOM analyzed colored tree for immune cell types based on the expression of cell-type enriched surface markers post-infection in control (left) or _LysM_-PAC$^{-/-}$ (right) peritoneal fluids.

E, Donut plots depicting composition of peritoneal immune cell types with respect to genotypes and treatments quantified with manual gating.

F, Manual gating strategy of indicated immune cell types in cyTOF dataset. DN, double negative.

G, Quantification of neutrophil and Ly6C^{high} monocyte populations through manual gating strategy. Each condition and genotype have at least n > 3 mice in independent experiments.

All control groups represent *Pacc1$^{F/F}$* genotype. Data are reported as mean ± SEM. Unpaired t-test for **B**. Two-way ANOVA with Sidak's test for **G**. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Figure 6. PAC deletion in macrophages promotes resolution of peritoneal *E. coli* infection.

A, Quantification of bacterial load in peritoneal fluids after wildtype *E. coli* peritoneal infection in control, *Cx3cr1*-PAC$^{-/-}$ (top), and *LysM*-PAC$^{-/-}$ (bottom) mice. Each timepoint or genotype has at least n > 4 mice in independent experiments.

B, Quantification of bacterial load in tail blood after wildtype *E. coli* peritoneal infection in control, *Cx3cr1*-PAC$^{-/-}$ (top), and *LysM*-PAC$^{-/-}$ (bottom) mice. Each timepoint or genotype has at least n > 5 mice in independent experiments.

C, Survival after wildtype *E. coli* peritoneal infection at 2 x 10^7 CFU/mL; WT n = 19. *Cx3cr1*-PAC$^{-/-}$ n = 13. *LysM*-PAC$^{-/-}$ n = 12.

D, Quantification of bacterial load in peritoneal fluids after cpm-resistant *E. coli* peritoneal infection in control, *Cx3cr1*-PAC$^{-/-}$ (top), and *LysM*-PAC$^{-/-}$ (bottom) mice. Each timepoint or genotype has at least n > 4 mice in independent experiments.

E, Survival after cpm-resistant *E. coli* peritoneal infection at 10^8 CFU/mL; WT n = 8. *Cx3cr1*-PAC$^{-/-}$ n = 9. *LysM*-PAC$^{-/-}$ n = 10.

All control groups represent *Pacc1*$^{F/F}$ genotype. Data are reported as mean ± SEM. Unpaired t-test for **A, B, D**. Mantel-Cox test and Gehan-Breslow-Wilcoxon test for **C, E**. *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 7. Summary diagram. PAC regulates phagosome maturation and host defense against bacteria in peritoneal macrophages.