Predicting the functional impact of single nucleotide variants in *Drosophila melanogaster* with FlyCADD

Julia Beets[1*], Julia Höglund[2,3], Bernard Y. Kim[4], Jacintha Ellers[1], Katja M. Hoedjes[1*], Mirte Bosse[1,3*]

[1] Amsterdam Institute for Life and Environment, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
[2] Centre for Palaeogenetics, Dept of Zoology, Stockholm University, Stockholm, Sweden
[3] Animal Breeding & Genomics, Wageningen University and Research, Wageningen, The Netherlands
[4] Department of Ecology and Evolutionary Biology, Princeton University, United States

Corresponding authors: j.beets@vu.nl, k.m.hoedjes@vu.nl, m.bosse@vu.nl

Running head: SNP impact prediction in the fruit fly

Keywords: single nucleotide polymorphism, *Drosophila melanogaster*, impact prediction, annotations, genotype-to-phenotype, GWAS, genome editing, evolutionary constraint, purifying selection

Abstract

Understanding how genetic variants drive phenotypic differences is a major challenge in molecular biology. Single nucleotide polymorphisms form the vast majority of genetic variation and play critical roles in complex, polygenic phenotypes, yet their functional impact is poorly understood from traditional gene-level analyses. In-depth knowledge on the impact of single nucleotide polymorphisms has broad applications in health and disease, population genomic or evolution studies. The wealth of genomic data and available functional genetic tools make *Drosophila melanogaster* an ideal model species for studies at single nucleotide resolution. However, to leverage these resources and potentially combine it with the power of functional genetics to enhance genotype-phenotype research, it is essential to develop techniques to predict functional impact and causality of single nucleotide variants.

Here, we present FlyCADD, a functional impact prediction tool for single nucleotide variants in *D. melanogaster*. FlyCADD, based on the Combined Annotation-Dependent Depletion (CADD) framework, integrates over 650 genomic features - including conservation scores, GC content, and DNA secondary structure - into a single metric reflecting a variant's predicted impact on evolutionary fitness. FlyCADD provides impact prediction scores for any single nucleotide variant on the *D. melanogaster* genome. We demonstrate FlyCADD's utility with some examples of application, including the ranking of phenotype-associated variants to prioritize variants for follow-up studies, evaluation of naturally occurring polymorphisms, and refining of CRISPR-Cas9 experimental design. FlyCADD provides a powerful framework for interpreting impact of any single nucleotide variant in *D. melanogaster*, thereby improving our understanding of genotype-phenotype connections.

Article summary

Single nucleotide polymorphisms (SNPs), the most common form of genomic variation, play key roles in micro-evolution and adaptation. In *Drosophila melanogaster*, many SNPs have been associated to phenotypes through association studies, yet functional validation remains challenging and experimental evidence for functional impact is rare. Here, we present FlyCADD, an impact prediction tool that integrates high-quality *D. melanogaster* genome annotations into a single score reflecting predicted impact of a SNP. FlyCADD can be applied to distinguish causal from neutral variants, for variant prioritization prior to functional studies, and to enhance interpretation of natural variation, thereby improving our understanding of genotype-phenotype relationships.

Introduction

Unravelling the causative relationships between phenotypes, genes and their variants represents a significant challenge in molecular biology, especially at the resolution of single nucleotides (Cano-Gamez and Trynka 2020; Gallagher and Chen-Plotkin 2018). Investigating functional effects of genetic variation advances our understanding of evolutionary adaptation, population dynamics, disease susceptibility, and other fundamental biological processes (Ungerer et al. 2008). The wealth of genotyping data has enabled researchers to explore population genomics, evolutionary adaptation, and the genetic basis of health and disease. For example, the identification of loci associated with Type 2 diabetes, as well as loci linked to fitness strategies such as insect pupation (Visscher et al. 2017; Xue et al. 2018; Zhang et al. 2020). Despite the unprecedented resolution and availability of genotype data, however, uncovering causality and interpreting genetic variation remains challenging.

Species typically harbor millions of single nucleotide polymorphisms (SNPs) in their genomes, for example there are over five million SNPs identified in *Drosophila melanogaster* (Auton et al. 2015; Nunez et al. 2024; Wang et al. 2015). Genomic studies, such as genome-wide association studies (GWAS) and Evolve and Resequence (E&R) studies, have identified SNPs associated with traits of interest (Uffelmann et al. 2021; Visscher et al. 2017). This led to the discovery of many candidate loci associated to a broad range of phenotypes (Wangler et al. 2017). Elucidating the functional effects of individual single nucleotide variants is critical for understanding how genetic variation shapes phenotypes (de Visser and Krug 2014). Yet, the functional impact of individual SNPs is only rarely elucidated after identification of phenotype-associated variants (Hoedjes et al. 2022; Katzenberger et al. 2015; Perlmutter et al. 2024; Schaid et al. 2018; Visscher et al. 2017).

There are several technical and biological challenges that complicate interpretation of genetic variation at single nucleotide level. Many of the SNPs identified in association studies are genetically linked variants that are not under selection themselves but merely show allele frequency (AF) changes by hitchhiking with loci that are under selection (Cano-Gamez and Trynka 2020; Gallagher and Chen-Plotkin 2018; Smith and Haigh 1974). Other factors that can complicate the interpretation of the effect(s) of SNPs are if several SNPs interplay through epistasis, if SNPs have additive effects, or if the impact of a SNP variant is affected by factors such as environment, population structure, diet or sex (Gallagher and Chen-Plotkin 2018; Huang et al. 2012; Yashin et al. 2012). Because of these factors, statistical discovery of SNPs does not always reflect the environmental and genomic context in which variants occur, nor the causality between variants and phenotypes. For understanding any genotype-phenotype link it is imperative to distinguish causal SNPs from linked neutral loci, yet distinguishing functional alleles from linked neutral variants among phenotype-associated SNPs remains a significant challenge (Franssen et al. 2015; Smith and Haigh 1974; Wang et al. 2022). Developing techniques and tools to assess the functional impact of single nucleotide polymorphisms (SNPs) is therefore essential for advancing genotype-phenotype research.

Functional genetics such as gene knockdown or knockout, through RNAi or CRISPR-Cas9 for example, and precise genome editing can give insight in the functional impact of SNPs. Various

experimental approaches have been applied to study functional impact of SNPs (Akhund-Zade et al. 2017; Hoedjes et al. 2023; Hoedjes et al. 2022; Mokashi et al. 2021; Parker et al. 2020; Wolf et al. 2023). These techniques are often applied at the gene-level, meaning that they focus on (nearby) genes associated with the SNP of interest, not taking into account the fact that a single gene may harbour multiple single nucleotide variants that can have different phenotypic effects (Hoedjes et al. 2023; Hoedjes et al. 2022; Mokashi et al. 2021; Perlmutter et al. 2024). Hence, this gene-level approach might not identify effects of the targeted SNPs but rather all phenotypic effects that a gene might have (Hoedjes et al. 2023; Hoedjes et al. 2022). Different SNPs within the same gene could potentially have different functional impact or pleiotropic effects, so it is important to understand the functional impact of SNPs instead of genes to understand phenotypic variation within species (Zhang and Yang 2015)(Hoedjes et al. 2023; Mokashi et al. 2021). Precise genome editing, for example using CRISPR-Cas9, enables targeted investigations at single nucleotide resolution, but these methods are costly and time-consuming so only few SNPs can be tested, and experimental evidence remains scarce (de Visser and Krug 2014; Hoedjes et al. 2023; Mokashi et al. 2021; Perlmutter et al. 2024). Therefore, it is critical to be able to prioritize the most promising candidate SNPs prior to functional studies.

Computational approaches can be applied to narrow down a set of candidate SNPs, allowing focused functional studies without spending costly time and resources on SNPs without functional impact themselves. Existing computational approaches for predicting the effects of SNPs include SIFT, PolyPhen, VEP, SnpEff, ANNOVAR, AlphaMissense, ProteoCast and GPN-MSA (Abakarova et al. 2025; Adzhubei et al. 2010; Benegas et al. 2023; Cheng et al. 2023; Cingolani et al. 2012; McLaren et al. 2016; Ng and Henikoff 2001; Wang et al. 2022). These tools, whether focusing on genome-wide patterns, (nearby) genes, or coding sequence variants, are based on a narrow set of genomic annotations—primarily conservation scores, genomic structure, and locus proximity (Wang et al. 2022). Predictions of SNP impact typically integrate insights from multiple tools to take advantage of various annotations (Gibert et al. 2017; Hall et al. 2019). Yet, many additional annotations that could provide critical insights into SNP functionality remain underutilized. These annotations provide essential insights for pinpointing the functional significance of genetic variants at single-nucleotide resolution (de Visser and Krug 2014; Öztürk-Çolak et al. 2024).

A promising computational approach that combines a much more extensive set of annotations to predict functional impact of genetic variants is the Combined Annotation-Dependent Depletion (CADD) framework (Kircher et al. 2014). The CADD framework is a first of its kind tool integrating many annotations into a single metric reflecting functional impact of any SNP. One of its unique features is that it is not restricted to SNPs in coding regions, details on related genes or specific groups of SNPs. CADD scores reflect the predicted impact of a variant, correlating with its biological consequences and selective constraints. CADD was first applied to the human genome and subsequently CADD models were specifically designed for chicken, mouse and pig genomes (Groß et al. 2020a; Groß et al. 2018; Groß et al. 2020b; Kircher et al. 2014; Schubach et al. 2024). These CADD models made it possible to score functional impact of SNPs for detection of pathogenic variants, ranking SNPs based on predicted impact, prioritizing them for functional testing, disease studies or enhancing animal breeding

(Boshove et al. 2023; Derks et al. 2021; Schubach et al. 2024; Speak et al. 2024; Wang et al. 2022).

D. melanogaster is a powerful model organism to study the genotype-phenotype link due to its rich repertoire of gene editing approaches, the wealth of genotyping data, high-quality genomes and single-nucleotide resolution annotations, enabled by large-scale sampling efforts and technological advancements (Kim et al. 2024; Nunez et al. 2024; Öztürk-Çolak et al. 2024; Wangler et al. 2017). Functional genomic studies are still rare, but the few done so far show unique and strong phenotypic effects of individual SNPs (Hoedjes et al. 2023; Mokashi et al. 2021; Perlmutter et al. 2024). To direct effort and resources to the most promising SNPs, high quality, whole genome annotations at single-nucleotide resolution are valuable tools to make substantiated hypothesis prior to functional studies about functional effects of SNPs as this location-based information provides crucial insights into their function(s). Integrating diverse annotations enables a more comprehensive perspective on how genetic variation translates into phenotypic diversity in *D. melanogaster*.

Here, we introduce FlyCADD, the first adaptation of the CADD framework to an insect genome, applied to *D. melanogaster*. FlyCADD combines the available high-quality annotations to provide a robust and versatile tool for studying the functional impact of SNPs by providing impact prediction scores for all possible genomic variants in the genome of *D. melanogaster*. Combined annotations include conservation scores, gene structures, regulatory elements, GC content, and DNA secondary structure, among others. Combining in total 691 features into a single impact prediction metric, FlyCADD enables researchers to prioritize SNPs based on their predicted functional impact across different contexts and in any genomic region of the *D. melanogaster* genome. SNPs identified in the genome of *D. melanogaster* natural populations, experimental strains or disease models can now be computationally studied based on predicted functional impact, with the potential to complement experimental studies or prioritize SNPs for further (functional) studies. This model represents a significant step forward in integrating computational predictions into the study of genotype-phenotype link in *D. melanogaster*.

We describe the FlyCADD impact prediction model and demonstrate that impact predictions for SNPs can improve interpretation of candidate SNPs and experimental design. To illustrate the usability of FlyCADD, we applied the FlyCADD impact prediction scores 1) to study naturally segregating SNPs, 2) to enhance interpretation of GWAS-identified SNP, 3) to rank SNPs prior to functional studies, and 4) to improve experimental design for genome editing approaches. To facilitate variant impact analysis, we make FlyCADD impact prediction scores available for all possible single nucleotide variants in the *D. melanogaster* Release 6 reference genome. Additionally, we provide scripts and pre-processed annotations for annotating novel variants. Both the precomputed scores and the locally executable pipeline can be accessed on Zenodo (10.5281/zenodo.14887338).

Methods

FlyCADD is a logistic regressor trained on 691 genomic features to assign an impact prediction score to any single nucleotide variant in the *D. melanogaster* genome. Impact refers to a mutation that on an evolutionary time scale is likely not tolerated at that position due to an associated fitness cost. We build upon the existing CADD models for human, pig and mouse by adapting the pipeline to fit *D. melanogaster* through incorporation of *D. melanogaster*-specific annotations, sequences, alignments and the reference genome (Groß et al. 2018; Groß et al. 2020b; Kircher et al. 2014). The annotations include conservation scores, marked regulatory elements and DNA secondary structure among others (see File S2). The FlyCADD model was created across chromosomes 2L, 2R, 3L, 3R, 4, and X based on the Release 6 reference genome (GCA_000001215.4). FlyCADD is aimed at scoring the impact of single nucleotide variants, no other types of variants were included in training, testing or application of the model. An outline of FlyCADD development and application is illustrated in Figure 1.

Figure 1: Overview of the FlyCADD pipeline indicating resources (blue) and information extracted (pink) to obtain the training datasets (teal) for development (left) of the FlyCADD model and application (right) of the model to any single nucleotide variant in the *D. melanogaster* genome.

Overview of FlyCADD model training

Briefly, the model consists of a logistic regressor trained to distinguish between two sets of training data: nearly-fixed derived variants - assumed to be depleted of impactful variants - and simulated variants – containing the full spectrum of impactful variants due to the absence of purifying selection. Ancestral sequence reconstruction was obtained from a Cactus whole-genome alignment of 166 *Drosophila* species in multiple alignment format (MAF)(Armstrong et al. 2020). This alignment was made available for FlyCADD development and represents an earlier version of the 298-way alignment described in (Kim et al. 2024). The 166-way alignment specific to this study is available on Zenodo (10.5281/zenodo.14887338). While both alignments were generated using the same methodology, the more recent version includes additional species and should be used for future analyses.

The set of derived variants was constructed by comparing the last common ancestral sequence of *D. melanogaster* (melanogaster subgroup) and *D. tani* (montium subgroup) to the Release 6 reference genome (GCA_000001215.4). Details of ancestral sequence reconstruction are described below (Methods "Obtaining training datasets"). Derived alleles

that are fixed or nearly-fixed (AF ≥ 0.9) in *D. melanogaster* populations across Europe and
America with a different allele relative to the reconstructed ancestral genome are assumed to
be depleted of deleterious variants due to purifying selection and belong to the derived
variant set (proxy-benign variants). The second set of single nucleotide variants, containing
simulated variants, was obtained by simulating *de novo* variants using mutation rates based
on comparison between the ancestral and reference sequence. This set of simulated variants
is thought to contain the full spectrum of deleteriousness, because natural selection has not
acted upon these variants. Therefore, compared to the derived set of variants, the simulated
set is enriched for deleterious alleles. In the following section ("Obtaining training datasets"),
construction of these training datasets and variant simulation are described in more detail.

The second step in the CADD pipeline was to annotate the derived and simulated variants with
a range of *Drosophila*-specific annotations. These annotations included a total of 38 individual
annotations such as sequence information, conservation scores, chromatin states, and
combinations thereof, resulting in a set of 691 features (see File S2). The FlyCADD model was
trained as a logistic regressor to classify new, annotated variants as belonging to either the
derived or simulated class. For each single nucleotide variant on the reference genome, an
impact prediction score representing the probability of the variant belonging to the simulated
variants class is readily available. The impact score is a measure for predicted functional
impact to fitness between 0 (low functional impact) and 1 (high functional impact). All
annotations and impact prediction scores are 1-based. The pipeline for FlyCADD development
can be found on GitHub (https://github.com/JuliaBeets/FlyCADD).

Obtaining training datasets

A Cactus 166-way alignment of *Drosophila* species was provided for ancestral sequence
reconstruction, prior to publication of the 298-way alignment (Kim et al. 2024). The Cactus
alignment software reconstructs ancestral genomes for each node, calling individual bases as
the maximum likelihood estimate from a Jukes-Cantor substitution model (Armstrong et al.
2020). The 166-way alignment including ancestral sequence reconstructions used in this work
is available on Zenodo (10.5281/zenodo.14887338). The ancestral sequence was obtained by
extracting the reconstructed sequence relative to coordinates in Release 6 of the reference
genome for the last common ancestor between *D. melanogaster* and *D. tani* within the 166-
way alignment (see File S1). This node was chosen as the reconstructed ancestral sequence
was largest in proportion to the reference genome (~ 16 %) and represents no excessive bias
towards specific chromosomes or coding sequences (see File S1). More recent ancestral
reconstruction yielded too few different alleles compared to the reference genome, whilst
more distant ancestral reconstruction yielded more bias towards coding regions and certain
chromosomes, specifically chromosome 3 (see File S1). Considering these aspects of bias is
important to ensure derived and simulated variants to be evenly distributed across the
genome.

Derived variants were identified as (nearly) fixed variants in the reference sequence with a
different allele in the reconstructed ancestral genome. To construct this set of variants, alleles
in the reference genome differing from that of the ancestral genome were identified.
Subsequently, to ensure that the set of derived variants includes (nearly) fixed variants,

variants with an allele frequency below 0.9 were removed. The *D. melanogaster* populations
used to define allele frequencies included 46 genomes from 12 *D. melanogaster* populations
in Europe and America (Rech et al. 2022) and > 200 Drosophila Genetic Reference Panel 2
(DGRP) lines (Mackay et al. 2012b). This resulted in a set of 2.733.695 derived variants, which
is assumed to contain benign or neutral variants.

An equal number of variants was obtained through simulation using the variant simulator of
the original CADD model to create a set of simulated variants that is enriched for impactful
variants compared to the set of derived variants (Kircher et al. 2014). Mutation rates and CpG
sites per chromosome were calculated based on comparison between the reconstructed
ancestral sequence and the reference genome and applied to simulate variants that are
absent in the reference genome but could have naturally occurred based on mutation rates in
the absence of purifying selection and free from the influence of biological selection. The final
mutation rates applied in simulation can be found on GitHub
(https://github.com/JuliaBeets/FlyCADD). The set of simulated *de novo* mutations was then
randomly trimmed to match the number of derived variants.

The final numbers of derived and simulated variants per chromosome are summarized in Table
1. The Y chromosome was excluded from model training due to the lack of sufficiently
reconstructed ancestral sequence needed to construct the datasets of derived and simulated
variants. In total, the training dataset contained 5.467.390 single nucleotide variants, equally
divided over the set of derived variants and the set of simulated variants.

Table 1: Final numbers of derived and simulated SNPs used in training and testing the logistic regression model.

	Derived	Simulated
Chromosome 2L	594.922	662.282
Chromosome 2R	471.645	435.022
Chromosome 3L	563.599	577.720
Chromosome 3R	658.177	659.379
Chromosome 4	25.216	15.959
Chromosome X	420.148	383.345
Total	2.733.695	2.733.695

Annotations

The training dataset was annotated with a total of 691 features (38 distinct annotations and
relevant combinations thereof, see File S2). The Ensembl Variant Effect Predictor (VEP, v105.0)
provided 13 sequence-based annotations in offline mode (McLaren et al. 2016). These were
supplemented with Grantham-scores, secondary structure predictions, repeat annotations,
regulatory elements, PhyloP, PhastCons, GERP, chromatin state, miRNA, coding regions,
transcription factor binding sites, and cis-regulatory motifs. Details for all features can be
found in File S2, and sources of the annotations are described below.

Grantham scores were incorporated based on the Grantham-matrix that annotates the
chemical difference induced by the amino acid change upon variation (Grantham 1974).
Secondary structure predictions for the reference genome were obtained using the *getshape*()

function from DNAshapeR (Chiu et al. 2016). The included features were minor groove width
(MGW), roll, helix twist (HelT), propeller twist (ProT), and electrostatic potential (EP).

Coordinates for eight repeat types (DNA repeat elements, low complexity repeats (LCR), long
interspersed nuclear elements (LINE), long terminal repeat elements (LTR), rolling circle (RC),
satellite repeats, simple repeats, and RNA repeats) were retrieved from the RepeatMasker
Track of the UCSC Genome Browser (v460, last updated 2014-08-28)(Perez et al. 2025).

ReMap records of regulatory elements, PhyloP base-wise and PhastCons element
conservation scores, were obtained from the UCSC Genome Browser (v460, last updated
2014-08-28)(Perez et al. 2025). ReMap records were incorporated in FlyCADD as the density
of records per position. PhyloP and PhastCons scores were based on different multi-species
alignment sizes: Multiz alignments of 27 insects (PhyloP/PhastCons) and 124 insects
(PhyloP124/PhastCons124). Additionally, PhyloP scores (PhyloP_all) and GERP-like scores
(GerpRS_all and GERPN_all) were computed from the 166-way Cactus alignment with PHAST
(Hubisz et al. 2010), using the implementation of the tools in the halPhyloPMP.py script
provided as part of the HAL toolkit (Hickey et al. 2013), excluding the *D. melanogaster*
sequence to avoid bias.

Chromatin states for BG3 and S2 cells, miRNA encoding sequence coordinates, and
coordinates for coding sequences were retrieved from FlyBase (Release FB2023_03)(Öztürk-
Çolak et al. 2024). Coordinates for cis-regulatory motifs, predicted cis-regulatory motifs, and
transcription factor binding sites were retrieved from REDfly (v9.6.2)(Keränen et al. 2022).

Several annotations were combined with VEP annotations to create the final set of 691
features (see File S2). These include all combinations of amino acid changes, variant
consequences, conservation scores, ReMap scores, and distances to coding sequences. All
features were scaled by their standard deviation based on variation within the training
dataset. In the final annotated dataset, derived variants were annotated with "1" and
simulated variants with "0".

Logistic regression model

A Turi Create Logistic Classifier (v6.4.1) was trained on 90% of the annotated dataset,
containing variants in two categories—"derived" and "simulated"-, reserving the remaining
10% for model testing. After cross-validation, the classifier was configured with L2
penalization set to 1.0, using the Newton solver, and a maximum of 100 iterations with
tc.logistic_classifier.create(). A trained model was obtained for the full genome, excluding the
Y chromosome due to its absence in the training dataset. The model assigns a weight to each
feature after training, representing the direction in which the feature contributes to the
FlyCADD score. A lower, negative weight represents contribution to the probability of variants
being impactful, resulting in higher FlyCADD-scores, whilst a more positive weight implies
more contribution of the feature towards the probability of variants being neutral / benign,
lowering the FlyCADD-score. There is no validation dataset available with known deleterious
or impactful SNPs in *D. melanogaster*. Therefore, the model was evaluated using a built-in
testing functionality of the logistic regression function applied to the held-out 10 % of the
training dataset, ensuring that none of these SNPs were used for training.

Computing FlyCADD scores

The FlyCADD model can be applied to any given variant on the *D. melanogaster* genome that is annotated with the same set of features and scaled by the standard deviation per feature within the training data. With *loaded_model.predict*(), FlyCADD generates a raw probability score for each variant, indicating its likelihood of belonging to the simulated class. FlyCADD scores range from 0, indicating no predicted functional impact, to 1, indicating high predicted functional impact. The FlyCADD model was applied to all possible single nucleotide variants on the *D. melanogaster* reference genome autosomes and X chromosome, resulting in posterior probabilities for each variant belonging to the simulated variants set and thus being impactful. Since the precomputed impact prediction scores for all possible genomic variants including those not observed in natural populations, predictions are unbiased by allele frequencies, function or location of known variants and solely based on variant annotations and model training. The precomputed impact prediction scores, trained model and scripts to score novel variants are readily available on Zenodo (10.5281/zenodo.14887338).

Variant impact at codon positions

Positions of nucleotides within the codon reflect differing biological importance for these nucleotides and therefore provide an opportunity to evaluate the impact prediction scores (Crick 1966). Genome-wide codon positions were retrieved from Ensembl (v111) and filtered for unique transcripts on the sense strand, resulting in transcripts of 6.894 genes. Genes with multiple transcripts were excluded to avoid interaction effects. The distributions of minimum FlyCADD scores within a codon were compared per position amongst all transcripts and within each gene using the Mann-Whitney U-test. Statistical analyses were performed using scipy.stats (v1.12.0) in Python3, with Bonferroni corrected p-values (Virtanen et al. 2020).

Allele frequency of natural variants

To investigate the distribution of FlyCADD scores for naturally occurring SNPs with differing allele frequency, positions and allele frequencies for over 4.8 million SNPs were obtained from the VCF file (version dest.all.PoolSNP.001.50.24Aug2024) of the *Drosophila* Evolution over Space and Time (DEST) 2.0 resource, the most comprehensive genomic resource for *D. melanogaster* genomes worldwide (Nunez et al. 2024). Only samples with label "PASS" were included in this analysis to avoid inclusion of low-quality variants.

Results

FlyCADD, a logistic regression model trained on over five million annotated variants, predicts the functional impact of SNPs in *D. melanogaster*. Impact scores were precomputed for all possible single nucleotide variants on the Release 6 reference genome (excluding the Y chromosome), with higher scores reflecting greater predicted functional impact of the SNP. FlyCADD scores range from 0 to 1, where lower scores (e.g., < 0.50) suggest a benign or neutral effect, while higher scores (e.g., > 0.50) indicate potentially impactful functional impact of the SNP. These scores allow researchers to rank SNPs based on their potential impact and assess the functional relevance of candidate SNPs. Here, we evaluate these scores based on the accuracy metrics of the model, contributions of different annotations, biologically relevant impact predictions and we show the tool's applicability across diverse use cases.

High accuracy achieved by training the logistic regression model

The performance of the FlyCADD model was evaluated using a held-out 10 % of the training dataset (see Methods). The overall performance for FlyCADD is ~ 0.83 (area under the ROC curve) with an accuracy of ~ 0.76, reflecting the proportion of correct predictions. It is important to note that FlyCADD's accuracy evaluation relies on a binary classification approach, where variants in the testing dataset were labeled strictly as 0 (neutral) or 1 (impactful), similar to the variants in the training dataset. However, functional impact is a continuous spectrum, and some variants labeled as 0 may still have (minor) effects. A prediction could be interpreted as misclassification during testing despite reflecting a biologically meaningful ranking, therefore, the accuracy of the model is likely to be higher when considering an impact gradient or using validation datasets for model evaluation. Validation datasets with SNPs of known pathogenicity are not available for *D. melanogaster*.

Key predictors are combined features

The strength of the CADD framework lies in its ability to integrate both individual and combined annotations, allowing for more precise distinction between variants. FlyCADD incorporated 38 individual annotations (of which 20 categorical) and combinations thereof (see File S2). Out of the total 691 features used for annotation by FlyCADD, 581 were combinations of individual annotations. The model assigns weights to these features, reflecting their contribution to the impact prediction. Across all features, 354 features were contributing to classification of variants as neutral or benign (with a positive weight), and 337 features were more contributing towards the probability of being impactful (with a negative weight). The impact prediction score is a combined score, taking into account all 691 features of a SNP. The weights for all 691 features can be found in File S3.

Figure 2 shows the top ten features that most strongly influence the model's ability to predict whether a variant is neutral or benign (Figure 2a) or impactful (Figure 2b). These features enhance predictive power but do not independently determine the predicted functional effect of a SNP. The strongest predictors of impact (Figure 2b) were the features indicating stop codon gain ("Consequence_SG"; weight = - 6.69) and stop codon loss dependent on position within the coding sequence ("SL_relCDSpos"; weight = - 2.66). The strongest predictors of

neutrality were the features indicating stop codon loss ("Consequence_SL"; weight = 4.23) and stop codon gain dependent on position within the coding sequence ("SG_relCDSpos"; weight = 3.76, Figure 2a).

When considering the twenty most influential features in the model (Figure 2), it is visible from the composed feature names that most key features are combinations of annotations. These combinations involve the annotations on conservation scores ("_PhyloP"), proximity to coding sequence ("_relCDSpos") and protein-level consequences (synonymous and non-synonymous, "SN_"/"NS_"). The individual annotations amongst the most influential features of FlyCADD are position within the protein ("protPos"), protein-level consequences stop loss, non-synonymous, synonymous and stop gain ("Consequence_"), stop codon as alternative allele (nAA_*) and conservation score (PhyloP).

Moreover, the influence of combined features in the model can be opposite to the effects of the individual annotations. For example, stop codon loss ("Consequence_SL") as a single annotation has a positive weight, whereas stop codon loss in combination with position in the coding sequence ("SL_relCDSpos") has a negative weight which indicates contribution to impact. A stop codon loss can therefore not only increase the probability of being neutral, but depending on the position in the coding sequence can also contribute to impact. A similar dependency was seen among stop codon gain ("Consequence_SG") and its position within the coding sequence ("SG_relCDSpos")(Figure 2). This pattern is consistent with biological expectations as stop codon polymorphisms in *D. melanogaster* are highly impactful due to changes to protein length, and especially deleterious in certain regions of protein closer to the end of the full length protein (Lee and Reinhardt 2012). These findings highlight the value of combining features into a single metric reflecting functional impact to differentiate between otherwise similar variants such as stop codon polymorphisms.

Figure 2: Feature weights for the ten highest (a) and ten lowest (b) weighted features. Positive feature weights indicate stronger contribution to the ability of the model to classifying a variant as neutral or benign, while negative feature weights indicate contribution to the probability of a variant being impactful. Features names "Consequence_" reflect the predicted protein-level consequence derived from the Ensembl Variant Effect Predictor. Abbreviations: (rel)CDSpos: (relative) position within the coding sequence; (rel)Protpos: (relative) position within the protein; E: glutamine; L: leucine; nAA: alternative amino acid; NS: non-synonymous substitution; oAA: original amino acid; SG: stop-gain; SL: stop-loss; SN: synonymous; *: stop codon. Detailed explanations of all features can be found in File S2.

Genome-wide FlyCADD scores reflect biological relevance of polymorphisms

Next, we analyzed the precomputed FlyCADD scores for all possible single nucleotide variants on the *D. melanogaster* reference genome. We first categorized the variants by chromosomal region (Figure 3). The genome-wide distribution of FlyCADD scores shows that predicted high-impact variants (> 0.95) are the most prevalent, while predicted low-impact variants (0.00 – 0.1) are the least frequent – a pattern that remains consistent across chromosomes (Figure 3). The limited number of low-impact scores (< 0.1) likely reflects that many of the theoretically possible SNPs in this dataset do not actually segregate in natural populations. Variants with such low predicted impact reflect a pattern of annotations by FlyCADD similar to the nearly-fixed, derived variants, as these are the proxy-benign variants in training. The theoretically possible SNPs in the set of all possible SNPs on the reference genome do not reflect such patterns as they are likely not variants that will become nearly-fixed or occur at all, and are therefore they are predicted to be more impactful.

Figure 3: Distribution of FlyCADD scores in the set of precomputed scores containing all possible single nucleotide variants on chromosomes 2L, 2R, 3L, 3R, 4 and X of the *D. melanogaster* reference genome.

Subsequently, we assessed whether FlyCADD scores vary among variants when categorized by predicted consequence to the protein (based on VEP), such as gain of a stop codon or synonymous variants, to provide insights into FlyCADD's ability to predict functional impact across classes of variants (Figure 4). For example, it is known that non-synonymous mutations and stop codon polymorphisms are impactful as they alter the protein. In line with our hypothesis, synonymous (SN) variants have lower FlyCADD scores, whereas non-synonymous (NS) variants have higher scores, potentially due to their amino acid changing properties (Figure 4). However, there are SN variants present in the dataset with higher predicted impact and, alternatively, NS variants with lower predicted impact are present (Figure 4). The range of predicted impact for both these types of variants is large and overlapping. Applying FlyCADD to NS or SN point mutations can help refine interpretation of the functional impact of these variants instead of solely relying on a single protein-level consequence prediction. Variants with other consequences showed a broader FlyCADD score distribution compared to SN or NS variants, emphasizing the additional information provided through integration of >

650 features. Intergenic (IG) and non-coding (NC) variants, for example, exhibit widely spread
FlyCADD scores, suggesting that insights into their functional impact benefit from combining
annotations into a single FlyCADD impact prediction score instead of relying on single features.
FlyCADD enables ranking of any single nucleotide variant, including those in less characterized
regions like intergenic regions, where functional insights are often limited.

Figure 4: Distribution of FlyCADD scores for all possible single nucleotide variants on the *D. melanogaster*
reference genome categorized by predicted protein-level consequences from VEP. UP: upstream; DN:
downstream; U5: 5 prime UTR; U3: 3 prime UTR; S: splice site; SN: synonymous; NS: non-synonymous; NC: non-
coding exon change; IG: intergenic; I: intronic; CS: canonical splice; SL: stop loss; SG: stop gain; O: unknown.

FlyCADD scores refine impact prediction within gene structure

Genes are composed of distinct functional regions, such as introns, exons and transcription
factor binding sites. We found that FlyCADD scores reflect the underlying genomic
arrangement by recognizing, amongst others, exonic variants with generally higher FlyCADD
scores compared to intronic variants when considering all possible single nucleotide variants
within genes. This is illustrated in Figure 5 for the gene *white* where, as expected, variants in
exons generally receive higher FlyCADD scores compared to intronic variants. However, both
regions contain SNPs that cover the full range of FlyCADD scores. By interpreting SNP impact
based on combined annotations rather than solely on gene structure, FlyCADD enables
functional impact prediction at single-nucleotide resolution rather than interpreting impact
of all variants within a functional domain as equivalent. Traditional approaches often infer SNP
impact based on proximity to genes or annotated domains, overlooking functional differences
between variants within the same region. FlyCADD allows researchers to distinguish between
neutral and functionally relevant variants within these functional regions.

Figure 5: Mean FlyCADD scores (purple) per position for all possible single nucleotide variants in the gene *white*, including 500 bp upstream and downstream of the gene. Exons of the gene are marked dark grey, introns marked in light grey.

Nucleotides at each codon position show differing predicted impact

To investigate the performance of FlyCADD impact predictions at single nucleotide resolution, we examined variation in FlyCADD scores across codon positions, which differ in functional importance due to their unique roles in DNA secondary structure, transcription, translation and mutation accumulation (Crick 1966). The third codon position is generally least impactful, as variation at this site rarely results in amino acid changes (Crick 1966).

We compared the distribution of minimum FlyCADD scores of all possible variants on the three codon positions across 6.894 genes (Figure 6a), using the lowest-scoring variant at each position to provide a conservative estimate of functional impact and avoid bias from outliers. Variants at the first and second codon position both have significantly higher FlyCADD scores compared to the third nucleotide, which is in line with expectations. The third nucleotide has least functional impact, followed by the first nucleotide, whereas the second nucleotide has most functional impact. Additionally, we compared the distribution of FlyCADD scores of the three codon positions per gene (Figure 6b). Analysis shows that the first nucleotide has significantly higher FlyCADD scores compared to the third nucleotide in 50.4 % of the genes and the second nucleotide in 62.2 % of the genes (Figure 6b). The first codon position was scored significantly higher by FlyCADD compared to the second nucleotide in 12 % of the genes. This is consistent with observations across all genes and indicate that FlyCADD scores reflect biologically expected patterns of functional impact at the single-nucleotide level.

Figure 6: Distribution of minimum FlyCADD scores of variants at each of the positions within the codon (a) across 6.894 transcripts and (b) compared within each gene. The proportion of genes where the distribution of FlyCADD scores significantly differed between two nucleotide positions after Bonferroni corrected Mann-Whitney U tests is shown in panel b.

Use cases for FlyCADD scores

Approaches such as GWAS and E&R often yield large numbers of candidate SNPs associated with a phenotype of interest, but these often include false positive and false negative hits due to linkage or technical limitations, such as statistical power (Wangler et al. 2017). In addition, experimental validation of candidate SNPs is costly, time-consuming and not yet feasible for large panels of SNPs. The FlyCADD model generates impact prediction scores reflecting fitness-reducing effect of SNPs, making it a promising tool for genetic research by facilitating the prioritization and interpretation of identified SNPs across a range of studies and topics. Here, we outline four key applications of FlyCADD to study *D. melanogaster* genomic variation at single nucleotide resolution, each illustrated with a specific example demonstrating the utility of impact prediction scores in various research contexts.

Impact of SNPs in natural populations of D. melanogaster

Natural populations of *D. melanogaster* exhibit extensive genetic diversity, with large numbers of segregating alleles continuously being identified (Nunez et al. 2024). While this natural variation is a driver of adaptation, genetic variation without functionally relevant impact is likely to circulate in populations as well. Methods for identification of functional loci from population genomic resources are often not focussed on functional impact of individual variants, but rather on sets of SNPs or nearby genomic regions (Harr et al. 2002; Jha et al. 2015). The FlyCADD model can filter out hitchhiking loci or variants with negligible functional effects from population genomics studies, and identify naturally occurring, high-impact SNPs. Furthermore, FlyCADD scores offer a basis for identifying ranges to classify variants by their potential impact, providing insights into the tolerability of variation within natural populations.

The genomic resource "*Drosophila* Evolution over Space and Time" (DEST) 2.0 contains high-quality whole-genome sequence information of 530 *D. melanogaster* populations worldwide (Nunez et al. 2024). When applying FlyCADD to > 4.5 million naturally segregating SNPs from

this dataset, we observed a mean FlyCADD score of 0.45, with an increased mean impact prediction score (0.50) observed for rare alleles (AF < 0.05)(Figure 7). When excluding rare alleles, the mean FlyCADD score is 0.36. This data is indicative of a higher predicted functional impact for rare alleles. Naturally occurring variants are showing a distribution of FlyCADD scores across the full range (0 – 1), irrespective of AF. Outliers with FlyCADD scores exceeding 0.80 (high impact) are more commonly observed among rare alleles, while becoming less frequent as AF increases. The presence of both low (< 0.50) and high (> 0.50) impact variants highlights the potential value of FlyCADD in differentiating functionally relevant variants from neutral loci within population genomic resources like DEST.

Based on the distribution of FlyCADD scores of naturally occurring SNPs (Figure 7), we propose a range for interpreting functional impact of naturally occurring SNPs. Variants with scores between 0 and approximately 0.40 appear to be generally tolerated in natural populations as they are frequently present in natural populations, whereas those with scores above 0.50 emerge as candidates for further study due to their high potential for functional impact. However, the degree to which functional impact influences populations may be shaped by ecological and selective pressures.

Figure 7: FlyCADD scores of naturally occurring SNPs, identified in the DEST2.0 genomic resource, across different allele frequency bins (bin size 0.025). Allele frequency represents the frequency of the non-reference allele (as mapped to the reference genome). The boxplots display the median, interquartile range and spread of FlyCADD scores. The mean FlyCADD score for variants per allele frequency bin is marked purple.

FlyCADD-based ranking compared to p-value ranking of SNPs from GWAS

Currently, p-values are the most common metric used for prioritization of phenotype-associated candidate SNPs resulting from GWAS. This approach gives a first insight into candidate loci, but identification of causal SNPs from these results is troublesome and subject to varying cut-off values (Cano-Gamez and Trynka 2020; Korte and Farlow 2013). Solely relying on p-values for ranking GWAS SNPs is prone to confounding factors such as linked variants, statistical limitations and environmental factors, which are well-known to influence these p-values (Chen et al. 2021; Fadista et al. 2016; Korte and Farlow 2013). FlyCADD scores can complement the p-value ranking derived from association studies by providing an impact prediction score per SNP and potentially offer additional insight into evolutionary fitness consequences of candidate SNPs.

A GWAS study by Katzenberger et al. provides an opportunity to apply FlyCADD scores in addition to p-values for assessment of functional impact of SNPs (Katzenberger et al. 2015). The study investigated the genetic basis of death after induced traumatic brain injury to individuals of DGRP lines. GWAS resulted in 216 candidate SNPs significantly associated to death induced by traumatic brain injury. Although the use of CRISPR technology was suggested to understand how each of these SNPs would affect the phenotype, no functional validation was performed in this study (Katzenberger et al. 2015).

We applied FlyCADD to complement p-values in ranking the candidate SNPs based on predicted functional impact. Comparison of the GWAS p-values to the FlyCADD-scores of significantly associated SNPs did not reveal a significant correlation between the two metrics (Figure 8a). Applying FlyCADD scores showed that variants with significant p-values can even have very low (< 0.2) predicted impact, and alternatively, variants with lower yet significant p-values can have high predicted impact (Figure 8a). For example, in the fifteen most significant candidate SNPs, ten variants have a predicted high (> 0.5) impact, and five variants have a predicted low (< 0.3) impact (Figure 8a). Nine of these SNPs are in a 2000 bp intergenic region on chromosome 2L with two of these SNPs having low predicted impact and seven having high predicted impact (Figure 8b). This suggests the presence of both causal (potentially those with high FlyCADD scores) and hitchhiking (potentially those with low FlyCADD scores) SNPs among this panel of candidate SNPs. In addition, one GWAS-identified SNP on chromosome 2L was not in the top 15 most significantly associated SNPs, yet it shows a moderate predicted impact and should not be overlooked by solely relying on p-value ranking in functional analysis of candidate loci (Figure 8b). Therefore, FlyCADD impact prediction scores apply an additional layer of information with regards to fitness-associated traits applicable to SNPs detected through GWAS.

Figure 8: FlyCADD score and p-value compared for (a) 216 SNPs associated to death after brain trauma in GWAS by Katzenberger et al. (Katzenberger et al. 2015). The fifteen SNPs with the lowest p-values are highlighted in purple. (b) Nine of these SNPs with the lowest p-values are in a 2000 bp intergenic region of chromosome 2L. One SNP ranked 36[th] lowest p-value is in the same region with moderate predicted impact (0.38, blue).

Ranking to identify causal SNP(s) or eliminate hitchhiking SNPs before experimental studies

Prioritization of SNPs using impact prediction scores helps eliminate hitchhiking loci and focus
follow-up studies on functionally relevant variants, which will ultimately clarify the genetic
architecture of complex traits. FlyCADD scores can be applied to either rank SNPs prior to
functional studies or to confirm or interpret results of functional studies. To show how to
prioritization using FlyCADD scores can be applied to GWAS-identified SNPs and how this
compares to experimental studies, we apply FlyCADD to the results of a GWAS on a fitness-
affecting phenotype.

Natural variation in pigmentation has been observed among *D. melanogaster* populations,
which can affect fitness by influencing temperature tolerance, mate choice, energy allocation
and visibility among others (Dembeck et al. 2015; Freoa et al. 2023). Moreover, experimental
evolution demonstrated that darker *D. melanogaster* populations showed a higher fecundity
and significantly reduced lifespan compared to the less pigmented populations (Rajpurohit et
al. 2016). Bastide et al. performed a GWAS identifying naturally occurring SNPs associated to
female abdominal pigmentation, which resulted in seventeen significant candidate SNPs and
hundreds of non-significant SNPs associated to a darker phenotype (Bastide et al. 2013). The
three most significantly associated SNPs were located in the cis-regulatory region of *tan*, a
pleiotropic gene involved in pigmentation, evolution of this trait, and other roles such as vision
or mating behaviour, suggesting that these SNPs affect pigmentation through regulation of *tan*
(Gibert et al. 2017; Massey et al. 2019). Transgenic lines with each one out of eight
combinations of these three SNPs were screened to determine SNP functional impact *in vivo*
(Gibert et al. 2017). This demonstrated that all three SNPs (X:9227096, X:9227061, X:9226889)
showed an additive, complex epistatic effect on pigmentation with most of the variance in
pigmentation explained by X:9227096. In addition, the allele at X:9227061, associated to the
dark phenotype in GWAS, resulted in less abdominal pigmentation in transgenic lines. This
effect is in the opposite direction of what the GWAS initially suggested, which Gibert et al.
attributed to linkage of the SNP with the other candidate SNPs as epistasis showed a less
strong effect (Gibert et al. 2017).

We applied FlyCADD to complement the GWAS results and *in vivo* study by ranking these SNPs
based on predicted functional impact. The FlyCADD impact prediction scores for the SNPs are
as follows: 0.30 for X:9227096, 0.04 for X:9227061, 0.41 for X:9226889 (Figure 9). With
FlyCADD scores between 0.30 and 0.50, two of these SNPs are predicted to have a moderate
phenotypic impact. The low FlyCADD score of X:9227061 (0.04) suggests that its association
to pigmentation in the GWAS was likely due to linkage with functionally relevant SNPs rather
than a direct phenotypic effect. This aligns with the findings of Gibert et al. where the SNP
showed opposite phenotypic effects compared to the GWAS prediction (Gibert et al. 2017).
The additional GWAS-identified SNPs (triangles in Figure 9) can now be ranked to focus
functional studies on those with highest FlyCADD scores. The prioritization of candidate SNPs
using FlyCADD scores facilitates targeted functional assays, focusing on top-ranked SNPs by
filtering out (possibly linked) SNPs with lower functional impact scores such as X:9227061.

Figure 9: Mean FlyCADD scores per position for all possible single nucleotide variants in the regulatory domain associated to *tan*, with the FlyCADD score of the GWAS identified SNPs in purple (circles functionally validated).

Evaluation of off-target effects in functional studies

Genome editing using CRISPR-Cas9 or other gene-editing technologies is widely used for functional genomics *in vivo* or using cell lines (Adli 2018). However, these techniques can introduce off-target effects, including unwanted point mutations, that may confound experimental results (Zhang et al. 2015). In some cases, intentional modifications in the target regions are required to increase efficiency of genome editing (Perlmutter et al. 2024; Zhang et al. 2015). FlyCADD can be applied in genome editing experiments to evaluate the potential functional impact of intentional off-target modifications during experimental design, as well as the potential functional impact of unintended off-target effects introduced during genome editing. This can increase the confidence that observed phenotypic effects are due to the on-target modification and cannot be attributed to (intentional or unintended) off-target effects.

In their study, Perlmutter et al. applied CRISPR-Cas9 to assess functional differences between two alleles of a non-synonymous polymorphism in the gene *Metchnikowin* (*Mtk*) in *D. melanogaster* (Perlmutter et al. 2024). *Mtk* is involved in microbe defense, response to wounding, neuropathology, and fitness among others (Levashina et al. 1998; Perlmutter et al. 2024; Swanson et al. 2020). Perlmutter et al. focussed on infection and life-history assessments of CRISPR-Cas9 edited and control lines (Perlmutter et al. 2024). In addition to the desired single nucleotide modification, multiple silent mutations were introduced in the target region to disrupt the protospacer adjacent motif (PAM) sites essential to Cas9 to increase the genome editing efficiency. The authors deemed the effect of these intended point mutations neglectable as these were synonymous mutations and limited expression differences were observed (Perlmutter et al. 2024).

We applied FlyCADD to determine whether the introduced silent mutations could have altered the results of the phenotypic screening. The SNP of interest (2R:15409024) has a predicted moderate functional impact with a FlyCADD score of 0.47, and the introduced silent mutations have FlyCADD scores of 0.52 (2R:15409016), 0.34 (2R:15409019) and 0.10 (2R:15409033). These impact prediction scores reveal that even though the intentional point mutations are silent in transcription, they are predicted to have impact on fitness, with one of these silent

mutations (2R:15409016) having a predicted impact higher than the predicted impact of the
SNP interest. These intended "silent" point mutations could potentially have impact on the
measured fitness-related phenotypes and therefore alter the phenotypic outcomes of the
study. FlyCADD scores can now be applied during experimental design to more precisely place
intended mutations and thereby avoid phenotypic effects caused by predetermined
mutations.

Additionally, it is possible to apply FlyCADD after genome editing by applying the scores to
sequencing data of the obtained lines to estimate whether observed phenotypic differences
are influenced by unintended off-target modifications. Perlmutter et al. identified 69 SNPs
differing between the edited and unedited lines, most of which were deemed cases of
mapping uncertainty (Perlmutter et al. 2024). Of these, however, 20 SNPs have a FlyCADD
scores indicating a probability of being impactful (> 0.50), therefore it is valuable to further
assess mapping quality for these SNPs to make sure these SNPs do not confound the
phenotypic results observed. The authors identified one missense SNP (2R:16854417) as a
confirmed off-target mutation segregating between the edited and unedited lines (Perlmutter
et al. 2024). This SNP has a very high FlyCADD score (0.99), indicating that the SNP is very likely
to impact fitness and potentially influenced the experimental results. This information on
predicted variant impact allows for a more cautious interpretation of phenotypic
measurements.

Discussion

FlyCADD is a versatile new tool for genetics research in *D. melanogaster*, offering valuable insights into the functional impact of SNPs across the entire genome. By integrating over 650 genomic features into a single impact prediction score, it enables researchers to prioritize SNPs, assess natural variation, evaluate off-target effects of genome editing, and clarify patterns of causal and neutral variants based on predicted functional impact. The CADD framework is based on the assumption that (nearly) fixed derived variants are depleted of impactful mutations, whilst purifying selection has not affected simulated variants, and therefore, the set of simulated variants contains variants on the full spectrum of impact (Kircher et al. 2014). It is important to note that while high impact scores indicate functional impact, they do not necessarily imply a harmful or deleterious fitness effect. FlyCADD scores reflect the probability of a variant belonging to the set of simulated variants, serving as a proxy for functional impact. Variants with a high predicted impact score have a higher probability of having an impact on fitness, either harmful or beneficial. FlyCADD can advance our understanding of the functional importance of SNPs and genotype-phenotype connections in *D. melanogaster* by providing impact scores for every SNP throughout the genome.

Ensuring model reliability through training and validation

The CADD framework works best when rich genomic information is available, such as in model species. The available sequencing datasets, high-quality reference genome, species alignments and nucleotide-based annotations for *D. melanogaster* made it feasible to develop FlyCADD tailored to *D. melanogaster* based on the existing CADD framework (Groß et al. 2020b; Hoskins et al. 2015; Kim et al. 2024; Kircher et al. 2014; Mackay et al. 2012a; Nunez et al. 2024; Rech et al. 2022). Ancestral sequence reconstruction was performed on a node that resulted in an ancestral sequence with minimized bias towards specific chromosomes or coding sequence, though challenges remained in reconstruction of heterochromatic regions such as the Y chromosome, repeat regions and centromeric regions as they are often lacking in genome assemblies (Chang and Larracuente 2018). Therefore, FlyCADD is trained on the autosomes and X chromosome and no scores are present for variants on the Y chromosome. The reconstructed ancestral genome was sufficiently large to develop a robust predictive model trained on more than five million annotated single nucleotide variants. Our data shows that the CADD approach can be effectively extended to insect genomes, especially when high-quality genomic resources are present.

Our analyses indicate that FlyCADD has a high accuracy, which can likely be attributed to the high-quality reference genome, multi-species alignment and annotations incorporated into FlyCADD. The ROC-AUC of FlyCADD is 0.83 with a prediction accuracy of 0.76 on the held-out test dataset. These numbers indicate that FlyCADD performance is higher compared to the existing mouse (ROC-AUC of 0.67) and pig (ROC-AUC of 0.68) CADD models (Groß et al. 2018; Groß et al. 2020b). mCADD (mouse) and humanCADD (hCADD, (Rentzsch et al. 2018; Schubach et al. 2024)), tested on available validation datasets containing variants of known pathogenicity rather than held-out training data, reported AUC scores of > 0.95. In contrast, there is no validation dataset available for functional impact of SNPs in *D. melanogaster*.

Therefore, direct comparison of accuracy is difficult as the CADD models are species-specific.
Species-specific models are important as, for example, mammal and insect genomes differ
widely, and therefore, predictability of functional impact of SNPs and importance of different
genomic features differ between species (Huber et al. 2017). FlyCADD as species-specific
application of the CADD framework tailored to *D. melanogaster* is beneficial for high-quality
genotype-phenotype research.

Species-specific functional predictions

Previous CADD models demonstrated that annotations derived from the sequence contribute
most to the predictions and that species-specific annotations are enhancing performance
(Groß et al. 2018). Out of a total of 691 features in FlyCADD, the most important features were
derived from the sequence and were combinations of positional, functional or conservation
scores, emphasizing the importance of composite features over single annotations for *D.*
melanogaster in contrast to mouse (mCADD) and pig (pCADD) models, where conservation
was the primary predictive factor (Groß et al. 2018; Groß et al. 2020b). Species-specific
application of the CADD framework is important to preserve these differences between
species and accurately predict functional impact.

It is currently unknown whether species-specific CADD models can reliably be applied to
(closely) related species, for example applying FlyCADD to *D. suzukii*, *D. simulans* or *D. yakuba*.
ChickenCADD (chCADD, (Groß et al. 2020a)) has been applied to the pink pigeon genome, but
no validation was performed to assess the accuracy of chCADD on this pink pigeon genome
(Speak et al. 2024). The availability of genome editing techniques and high-quality genomic
resources in *D. melanogaster* and related *Drosophila* species provides opportunities to
functionally validate impact prediction scores from FlyCADD within and among related
species. Future research could evaluate FlyCADD's transferability to other *Drosophila* species
through score lift over or by directly scoring novel SNPs, as the scripts and impact prediction
scores of FlyCADD are publicly available.

Genome-wide functional insights from FlyCADD

CADD applied to the human genome is widely used for analysis of disease-related
polymorphisms, while pCADD and chCADD have been applied to agriculture and conservation
genomics (Derks et al. 2021; Groß et al. 2020b; Speak et al. 2024). By applying FlyCADD to
different use cases, we demonstrated several examples of its value in prioritizing SNPs from
GWAS prior to follow-up studies, designing and evaluating of genome editing experiments and
assessment of natural variation in the fruit fly. FlyCADD enables researchers to examine
functional impact of candidate loci associated to complex traits at the SNP level across the
entire *D. melanogaster* genome and in different research contexts, offering insights into the
genetic basis of phenotypic variation. Given that linkage between causal SNPs and hitchhiking
loci is common, both in natural and experimental populations, FlyCADD can help distinguish
functionally impactful SNPs from neutral loci based on combined annotations, especially in
non-coding regions where traditional methods of assessing functional impact of SNPs fall
short, such as intergenic regions (Smit-McBride et al. 1988; Smith and Haigh 1974). We have
shown that FlyCADD scores can easily be applied as an additional layer of information on

candidate SNPs and as scores are available across the entire genome, no variant or region is overlooked. Currently, FlyCADD can only be used to score functional impact of point mutations and not structural variants or other types of genomic variants. Future expansion of FlyCADD towards predicting functional impact of other types of genomic variants is possible, however. For example, the CADD framework has been applied to score functional impact of structural variants in the human genome (Kleinert and Kircher 2022).

In conclusion, we present FlyCADD, an impact prediction tool for SNPs in the genome of *D. melanogaster.* We have demonstrated its use in assessing functionally relevant SNPs in natural populations, ranking variants prior to functional studies, enhancing interpretation of GWAS results and refining genome editing study design and evaluation. In addition to the selected use cases, FlyCADD impact prediction scores can be used on a genome-wide scale in many more research contexts to study the genetic basis of phenotypes in *D. melanogaster*. Prioritization of promising SNPs or elimination of predicted low-impact SNPs based on FlyCADD impact prediction scores will help researchers reduce time, cost, and effort of follow-up studies by preventing unnecessary validation of hitchhiking loci that are not likely to impact the phenotype of interest. This targeted approach directs resources toward functionally relevant SNPs, making (experimental or computational) validation of impact more feasible.

We have made FlyCADD impact prediction scores readily available on Zenodo, providing both precomputed scores for all possible single nucleotide variants on the *D. melanogaster* reference genome and a locally executable pipeline for scoring novel variants of interest (10.5281/zenodo.14887338). By extending functional SNP prediction beyond coding regions and known functional domains, FlyCADD presents a valuable addition to the existing genetic toolbox of *Drosophila*. The combination of FlyCADD with the powerful functional genetic tools for *D. melanogaster* provides unique opportunities to create a better understanding of the genotype-phenotype connections underlying phenotypic variation.

Data availability

File S1 describes the Cactus 166-way multi-species alignment that was used for ancestral sequence reconstruction and details on the obtained ancestral sequence, including two alternative ancestral sequences that showed excessive biases. File S2 gives an overview of the annotations and combinations thereof used by FlyCADD. File S3 shows the weight of all features incorporated in the trained logistic regression model, describing the contribution of each feature to the impact prediction.

All supporting data and resources related to FlyCADD are publicly available. The pipeline of FlyCADD development is available at GitHub (https://github.com/JuliaBeets/FlyCADD). Precomputed scores, a locally executable FlyCADD pipeline (including the trained logistic regression model files, annotation files and scripts) for scoring novel variants, the multi-species alignment file, the reconstructed ancestral sequence, generated derived and simulated variants, and FlyCADD scores at all codon positions of unique transcripts in the *D. melanogaster* genome can be found on Zenodo (10.5281/zenodo.14887338).

Acknowledgements

The authors acknowledge BAZIS HPC cluster computing facilities at the Vrije Universiteit Amsterdam. The authors thank the *D. melanogaster* research community for their contributions in generating essential resources, annotations, and sequencing datasets that formed the basis for developing FlyCADD. We acknowledge the *Drosophila* Evolutionary Population Genomics Consortium (DrosEU) for generating and maintaining essential sequencing datasets and genomic resources, including the DEST(2.0) resource.

The authors have declared no competing interest.

Funding

This project was supported by the 2022 Early Career Support Grant of the Amsterdam Institute of Life and Environment (A-LIFE) at Vrije Universiteit Amsterdam awarded to MB and KMH. JH was supported by Swedish Research Council (2022-00209_VR).

References

Abakarova M, Freiberger MI, Lierhmann A, Rera M, Laine E. 2025. Proteome-wide prediction of the functional impact of missense variants with proteocast. bioRxiv.2025.2002.2009.637326. doi:10.1101/2025.02.09.637326.

Adli M. 2018. The crispr tool kit for genome editing and beyond. Nature Communications. 9(1):1911. doi:10.1038/s41467-018-04252-2.

Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. 2010. A method and server for predicting damaging missense mutations. Nat Methods. 7(4):248-249. doi:10.1038/nmeth0410-248.

Akhund-Zade J, Bergland AO, Crowe SO, Unckless RL. 2017. The genetic basis of natural variation in drosophila (diptera: Drosophilidae) virgin egg retention. J Insect Sci. 17(1). doi:10.1093/jisesa/iew094.

Armstrong J, Hickey G, Diekhans M, Fiddes IT, Novak AM, Deran A, Fang Q, Xie D, Feng S, Stiller J et al. 2020. Progressive cactus is a multiple-genome aligner for the thousand-genome era. Nature. 587(7833):246-251. doi:10.1038/s41586-020-2871-y.

Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, Chakravarti A, Clark AG, Donnelly P, Eichler EE et al. 2015. A global reference for human genetic variation. Nature. 526(7571):68-74. doi:10.1038/nature15393.

Bastide H, Betancourt A, Nolte V, Tobler R, Stöbe P, Futschik A, Schlötterer C. 2013. A genome-wide, fine-scale map of natural pigmentation variation in drosophila melanogaster. PLoS Genet. 9(6):e1003534. doi:10.1371/journal.pgen.1003534.

Benegas G, Albors C, Aw AJ, Ye C, Song YS. 2023. Gpn-msa: An alignment-based DNA language model for genome-wide variant effect prediction. bioRxiv.2023.2010.2010.561776. doi:10.1101/2023.10.10.561776.

Boshove A, Derks MFL, Sevillano CA, Lopes MS, Son Mv, Knol EF, Dibbits BW, Harlizius B. 2023. Large scale sequence-based screen for recessive variants allows for identification and monitoring of rare deleterious variants in pigs. PLOS Genetics. 20. doi:10.1371/journal.pgen.1011034.

Cano-Gamez E, Trynka G. 2020. From gwas to function: Using functional genomics to identify the mechanisms underlying complex diseases. Frontiers in Genetics. 11. doi:10.3389/fgene.2020.00424.

Chang C-H, Larracuente AM. 2018. Heterochromatin-enriched assemblies reveal the sequence and organization of the drosophila melanogaster y chromosome. Genetics. 211(1):333-348. doi:10.1534/genetics.118.301765.

Chen Z, Boehnke M, Wen X, Mukherjee B. 2021. Revisiting the genome-wide significance threshold for common variant gwas. G3 Genes|Genomes|Genetics. 11(2):jkaa056. doi:10.1093/g3journal/jkaa056.

Cheng J, Novati G, Pan J, Bycroft C, Žemgulytė A, Applebaum T, Pritzel A, Wong LH, Zielinski M, Sargeant T et al. 2023. Accurate proteome-wide missense variant effect prediction with alphamissense. Science. 381(6664):eadg7492. doi:10.1126/science.adg7492.

Chiu TP, Comoglio F, Zhou T, Yang L, Paro R, Rohs R. 2016. Dnashaper: An r/bioconductor package for DNA shape prediction and feature encoding. Bioinformatics. 32(8):1211-1213. doi:10.1093/bioinformatics/btv735.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, snpeff. Fly. 6(2):80-92. doi:10.4161/fly.19695.

Crick FH. 1966. Codon--anticodon pairing: The wobble hypothesis. J Mol Biol. 19(2):548-555. doi:10.1016/s0022-2836(66)80022-0.

de Visser JAGM, Krug J. 2014. Empirical fitness landscapes and the predictability of evolution. Nature Reviews Genetics. 15(7):480-490. doi:10.1038/nrg3744.

Dembeck LM, Huang W, Magwire MM, Lawrence F, Lyman RF, Mackay TFC. 2015. Genetic architecture of abdominal pigmentation in drosophila melanogaster. PLOS Genetics. 11(5):e1005163. doi:10.1371/journal.pgen.1005163.

Derks MFL, Gross C, Lopes MS, Reinders MJT, Bosse M, Gjuvsland AB, de Ridder D, Megens H-J, Groenen MAM. 2021. Accelerated discovery of functional genomic variation in pigs. Genomics. 113(4):2229-2239. doi:10.1016/j.ygeno.2021.05.017.

Fadista J, Manning AK, Florez JC, Groop L. 2016. The (in)famous gwas p-value threshold revisited and updated for low-frequency variants. Eur J Hum Genet. 24(8):1202-1205. doi:10.1038/ejhg.2015.269.

Franssen SU, Nolte V, Tobler R, Schlötterer C. 2015. Patterns of linkage disequilibrium and long range hitchhiking in evolving experimental drosophila melanogaster populations. Molecular Biology and Evolution. 32(2):495-509. doi:10.1093/molbev/msu320.

Freoa L, Chevin L-M, Christol P, Méléard S, Rera M, Véber A, Gibert J-M. 2023. Drosophilids with darker cuticle have higher body temperature under light. Scientific Reports. 13(1):3513. doi:10.1038/s41598-023-30652-6.

Gallagher MD, Chen-Plotkin AS. 2018. The post-gwas era: From association to function. Am J Hum Genet. 102(5):717-730. doi:10.1016/j.ajhg.2018.04.002.

Gibert J-M, Blanco J, Dolezal M, Nolte V, Peronnet F, Schlötterer C. 2017. Strong epistatic and additive effects of linked candidate snps for drosophila pigmentation have implications for analysis of genome-wide association studies results. Genome Biology. 18(1):126. doi:10.1186/s13059-017-1262-7.

Grantham R. 1974. Amino acid difference formula to help explain protein evolution. Science. 185(4154):862-864. doi:10.1126/science.185.4154.862.

Groß C, Bortoluzzi C, de Ridder D, Megens HJ, Groenen MAM, Reinders M, Bosse M. 2020a. Prioritizing sequence variants in conserved non-coding elements in the chicken genome using chcadd. PLoS Genet. 16(9):e1009027. doi:10.1371/journal.pgen.1009027.

Groß C, de Ridder D, Reinders M. 2018. Predicting variant deleteriousness in non-human species: Applying the cadd approach in mouse. BMC Bioinformatics. 19(1):373. doi:10.1186/s12859-018-2337-5.

Groß C, Derks M, Megens H-J, Bosse M, Groenen MAM, Reinders M, de Ridder D. 2020b. Pcadd: Snv prioritisation in sus scrofa. Genetics Selection Evolution. 52(1):4. doi:10.1186/s12711-020-0528-9.

Hall BS, Barnett YA, Crofts JJ, Chuzhanova N. 2019. Identification of novel genes associated with longevity in <i>drosophila melanogaster</i> - a computational approach. Aging. 11(23):11244-11267. doi:10.18632/aging.102527.

Harr B, Kauer M, Schlötterer C. 2002. Hitchhiking mapping: A population-based fine-mapping strategy for adaptive mutations in <i>drosophila</i> <i>melanogaster</i>. Proceedings of the National Academy of Sciences. 99(20):12949-12954. doi:10.1073/pnas.202336899.

Hickey G, Paten B, Earl D, Zerbino D, Haussler D. 2013. Hal: A hierarchical format for storing and analyzing multiple genome alignments. Bioinformatics. 29(10):1341-1342. doi:10.1093/bioinformatics/btt128.

Hoedjes KM, Kostic H, Flatt T, Keller L. 2023. A single nucleotide variant in the pparγ-homolog eip75b affects fecundity in drosophila. Molecular Biology and Evolution. 40(2):msad018. doi:10.1093/molbev/msad018.

Hoedjes KM, Kostic H, Keller L, Flatt T. 2022. Natural alleles at the doa locus underpin evolutionary changes in drosophila lifespan and fecundity. Proc Biol Sci. 289(1986):20221989. doi:10.1098/rspb.2022.1989.

Hoskins RA, Carlson JW, Wan KH, Park S, Mendez I, Galle SE, Booth BW, Pfeiffer BD, George RA, Svirskas R et al. 2015. The release 6 reference sequence of the drosophila melanogaster genome. Genome Res. 25(3):445-458. doi:10.1101/gr.185579.114.

Huang W, Richards S, Carbone MA, Zhu D, Anholt RR, Ayroles JF, Duncan L, Jordan KW, Lawrence F, Magwire MM et al. 2012. Epistasis dominates the genetic architecture of drosophila quantitative traits. Proc Natl Acad Sci U S A. 109(39):15553-15559. doi:10.1073/pnas.1213423109.

Huber CD, Kim BY, Marsden CD, Lohmueller KE. 2017. Determining the factors driving selective effects of new nonsynonymous mutations. Proceedings of the National Academy of Sciences. 114(17):4465-4470. doi:10.1073/pnas.1619508114.

Hubisz MJ, Pollard KS, Siepel A. 2010. Phast and rphast: Phylogenetic analysis with space/time models. Briefings in Bioinformatics. 12(1):41-51. doi:10.1093/bib/bbq072.

Jha AR, Miles CM, Lippert NR, Brown CD, White KP, Kreitman M. 2015. Whole-genome resequencing of experimental populations reveals polygenic basis of egg-size variation in drosophila melanogaster. Molecular Biology and Evolution. 32(10):2616-2632. doi:10.1093/molbev/msv136.

Katzenberger RJ, Chtarbanova S, Rimkus SA, Fischer JA, Kaur G, Seppala JM, Swanson LC, Zajac JE, Ganetzky B, Wassarman DA. 2015. Death following traumatic brain injury in drosophila is associated with intestinal barrier dysfunction. eLife. 4:e04790. doi:10.7554/eLife.04790.

Keränen SVE, Villahoz-Baleta A, Bruno AE, Halfon MS. 2022. Redfly: An integrated knowledgebase for insect regulatory genomics. Insects. 13(7):618. doi:10.3390/insects13070618.

Kim BY, Gellert HR, Church SH, Suvorov A, Anderson SS, Barmina O, Beskid SG, Comeault AA, Crown KN, Diamond SE et al. 2024. Single-fly genome assemblies fill major phylogenomic gaps across the drosophilidae tree of life. PLOS Biology. 22(7):e3002697. doi:10.1371/journal.pbio.3002697.

Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. 2014. A general framework for estimating the relative pathogenicity of human genetic variants. Nature Genetics. 46(3):310-315. doi:10.1038/ng.2892.

Kleinert P, Kircher M. 2022. A framework to score the effects of structural variants in health and disease. Genome Res. 32(4):766-777. doi:10.1101/gr.275995.121.

Korte A, Farlow A. 2013. The advantages and limitations of trait analysis with gwas: A review. Plant Methods. 9(1):29. doi:10.1186/1746-4811-9-29.

Lee YC, Reinhardt JA. 2012. Widespread polymorphism in the positions of stop codons in drosophila melanogaster. Genome Biol Evol. 4(4):533-549. doi:10.1093/gbe/evr113.

Levashina EA, Ohresser S, Lemaitre B, Imler J-L. 1998. Two distinct pathways can control expression of the gene encoding the drosophila antimicrobial peptide metchnikowin11edited by m. Yaniv. Journal of Molecular Biology. 278(3):515-527. doi:10.1006/jmbi.1998.1705.

Mackay TF, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire MM, Cridland JM et al. 2012a. The drosophila melanogaster genetic reference panel. Nature. 482(7384):173-178. doi:10.1038/nature10811.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire MM, Cridland JM et al. 2012b. The drosophila melanogaster genetic reference panel. Nature. 482(7384):173-178. doi:10.1038/nature10811.

Massey JH, Akiyama N, Bien T, Dreisewerd K, Wittkopp PJ, Yew JY, Takahashi A. 2019. Pleiotropic effects of ebony and tan on pigmentation and cuticular hydrocarbon composition in drosophila melanogaster. Front Physiol. 10:518. doi:10.3389/fphys.2019.00518.

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F. 2016. The ensembl variant effect predictor. Genome Biology. 17(1):122. doi:10.1186/s13059-016-0974-4.

Mokashi SS, Shankar V, Johnstun JA, Huang W, Mackay TFC, Anholt RRH. 2021. Systems genetics of single nucleotide polymorphisms at the drosophila obp56 locus. bioRxiv.2021.2006.2028.450219. doi:10.1101/2021.06.28.450219.

Ng PC, Henikoff S. 2001. Predicting deleterious amino acid substitutions. Genome Res. 11(5):863-874. doi:10.1101/gr.176601.

Nunez JCB, Coronado-Zamora M, Gautier M, Kapun M, Steindl S, Ometto L, Hoedjes KM, Beets J, Wiberg RAW, Mazzeo GR et al. 2024. Footprints of worldwide adaptation in structured populations of d. Melanogaster through the expanded dest 2.0 genomic resource. bioRxiv.2024.2011.2010.622744. doi:10.1101/2024.11.10.622744.

Öztürk-Çolak A, Marygold SJ, Antonazzo G, Attrill H, Goutte-Gattat D, Jenkins VK, Matthews BB, Millburn G, dos Santos G, Tabone CJ, FlyBase C. 2024. Flybase: Updates to the drosophila genes and genomes database. Genetics. 227(1):iyad211. doi:10.1093/genetics/iyad211.

Parker GA, Kohn N, Spirina A, McMillen A, Huang W, Mackay TFC. 2020. Genetic basis of increased lifespan and postponed senescence in drosophila melanogaster. G3 Genes|Genomes|Genetics. 10(3):1087-1098. doi:10.1534/g3.120.401041.

Perez G, Barber GP, Benet-Pages A, Casper J, Clawson H, Diekhans M, Fischer C, Gonzalez JN, Hinrichs AS, Lee CM et al. 2025. The ucsc genome browser database: 2025 update. Nucleic Acids Res. 53(D1):D1243-d1249. doi:10.1093/nar/gkae974.

Perlmutter JI, Chapman JR, Wilkinson MC, Nevarez-Saenz I, Unckless RL. 2024. A single amino acid polymorphism in natural metchnikowin alleles of drosophila results in systemic immunity and life history tradeoffs. PLOS Genetics. 20(3):e1011155. doi:10.1371/journal.pgen.1011155.

Rajpurohit S, Richardson R, Dean J, Vazquez R, Wong G, Schmidt PS. 2016. Pigmentation and fitness trade-offs through the lens of artificial selection. Biol Lett. 12(10). doi:10.1098/rsbl.2016.0625.

Rech GE, Radío S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H, Jamilloux V, Quesneville H, González J. 2022. Population-scale long-read sequencing uncovers transposable elements associated with gene expression variation and adaptive signatures in drosophila. Nat Commun. 13(1):1948. doi:10.1038/s41467-022-29518-8.

Rentzsch P, Witten D, Cooper GM, Shendure J, Kircher M. 2018. Cadd: Predicting the deleteriousness of variants throughout the human genome. Nucleic Acids Research. 47(D1):D886-D894. doi:10.1093/nar/gky1016.

Schaid DJ, Chen W, Larson NB. 2018. From genome-wide associations to candidate causal variants by statistical fine-mapping. Nat Rev Genet. 19(8):491-504. doi:10.1038/s41576-018-0016-z.

Schubach M, Maass T, Nazaretyan L, Röner S, Kircher M. 2024. Cadd v1.7: Using protein language models, regulatory cnns and other nucleotide-level scores to improve genome-wide variant predictions. Nucleic Acids Res. 52(D1):D1143-d1154. doi:10.1093/nar/gkad989.

Smit-McBride Z, Moya A, Ayala FJ. 1988. Linkage disequilibrium in natural and experimental populations of drosophila melanogaster. Genetics. 120(4):1043-1051. doi:10.1093/genetics/120.4.1043.

Smith JM, Haigh J. 1974. The hitch-hiking effect of a favourable gene. Genetical Research. 23(1):23-35. doi:10.1017/S0016672300014634.

Speak SA, Birley T, Bortoluzzi C, Clark MD, Percival-Alwyn L, Morales HE, van Oosterhout C. 2024. Genomics-informed captive breeding can reduce inbreeding depression and the genetic load in zoo populations. Molecular Ecology Resources. 24(7):e13967. doi:10.1111/1755-0998.13967.

Swanson LC, Rimkus SA, Ganetzky B, Wassarman DA. 2020. Loss of the antimicrobial peptide metchnikowin protects against traumatic brain injury outcomes in drosophila melanogaster. G3 (Bethesda). 10(9):3109-3119. doi:10.1534/g3.120.401377.

Uffelmann E, Huang QQ, Munung NS, de Vries J, Okada Y, Martin AR, Martin HC, Lappalainen T, Posthuma D. 2021. Genome-wide association studies. Nature Reviews Methods Primers. 1(1):59. doi:10.1038/s43586-021-00056-9.

Ungerer MC, Johnson LC, Herman MA. 2008. Ecological genomics: Understanding gene and genome function in the natural environment. Heredity. 100(2):178-183. doi:10.1038/sj.hdy.6800992.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J et al. 2020. Scipy 1.0: Fundamental algorithms for scientific computing in python. Nature Methods. 17(3):261-272. doi:10.1038/s41592-019-0686-2.

Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, Yang J. 2017. 10 years of gwas discovery: Biology, function, and translation. The American Journal of Human Genetics. 101(1):5-22. doi:10.1016/j.ajhg.2017.06.005.

Wang D, Li J, Wang Y, Wang E. 2022. A comparison on predicting functional impact of genomic variants. NAR Genom Bioinform. 4(1):lqab122. doi:10.1093/nargab/lqab122.

Wang F, Jiang L, Chen Y, Haelterman NA, Bellen HJ, Chen R. 2015. Flyvar: A database for genetic variation in drosophila melanogaster. Database. 2015:bav079. doi:10.1093/database/bav079.

Wangler MF, Hu Y, Shulman JM. 2017. Drosophila and genome-wide association studies: A review and resource for the functional dissection of human complex traits. Dis Model Mech. 10(2):77-88. doi:10.1242/dmm.027680.

Wolf S, Abhyankar V, Melo D, Ayroles JF, Pallares LF. 2023. From gwas to signal validation: An approach for estimating genetic effects while preserving genomic context. bioRxiv. doi:10.1101/2023.03.09.531909.

Xue A, Wu Y, Zhu Z, Zhang F, Kemper KE, Zheng Z, Yengo L, Lloyd-Jones LR, Sidorenko J, Wu Y et al. 2018. Genome-wide association analyses identify 143 risk variants and putative regulatory mechanisms for type 2 diabetes. Nature Communications. 9(1):2941. doi:10.1038/s41467-018-04951-w.

Yashin AI, Wu D, Arbeev KG, Ukraintseva SV. 2012. Polygenic effects of common single-nucleotide polymorphisms on life span: When association meets causality. Rejuvenation Res. 15(4):381-394. doi:10.1089/rej.2011.1257.

Zhang J, Yang J-R. 2015. Determinants of the rate of protein sequence evolution. Nature Reviews Genetics. 16(7):409-420. doi:10.1038/nrg3950.

Zhang W, Reeves GR, Tautz D. 2020. Identification of a genetic network for an ecologically relevant behavioural phenotype in drosophila melanogaster. Mol Ecol. 29(3):502-518. doi:10.1111/mec.15341.

Zhang XH, Tee LY, Wang XG, Huang QS, Yang SH. 2015. Off-target effects in crispr/cas9-mediated genome engineering. Mol Ther Nucleic Acids. 4(11):e264. doi:10.1038/mtna.2015.37.