Cross-family Signaling: PDGF Mediates VEGFR Activation and Endothelial Function

Xinming Liu[1], Yingye Fang[1], P.I. Imoukhuede[1]

[1]University of Washington, Department of Bioengineering, Seattle, WA.

Abstract

A recently discovered but unexplored mechanism of angiogenesis regulation is cross-family
binding between platelet-derived growth factors (PDGFs) and vascular endothelial growth factor
receptors (VEGFRs), which suggests a novel therapy for addressing vascular dysregulation. This
study elucidates the role of PDGFs in endothelial cell (EC) signaling and functions, focusing on
VEGFR activation. Using human dermal microvascular ECs (HDMECs) with double knockout of
PDGFRα/β and human brain microvascular ECs (HBMECs), we show three key findings:
(1) PDGF-AA and -BB induced VEGFR1 phosphorylation, peaking at 2-fold increases at low
concentrations (0.5 ng/mL), while PDGF-AB stimulated a 2-fold rise in VEGFR2 phosphorylation.
(2) Downstream effectors PLCγ1, Akt, and FAK were activated by all three PDGFs at levels
comparable to VEGF-A, achieving approximately 70% of VEGF-A's effects. (3) PDGF-BB
significantly enhanced EC proliferation (up to 240%) and migration (up to 170%), with lower
PDGF concentrations (0.5–5 ng/mL) eliciting stronger effects than higher concentrations (50–100
ng/mL). Overall, PDGF subtypes differentially induce VEGFR phosphorylation, downstream
effector activation, and angiogenic hallmarks such as proliferation and migration, revealing novel
mechanisms for regulating endothelial function.

Introduction

Angiogenesis, the formation of new blood vessels from existing vascular systems, is central to
several physiological conditions and supports tissue growth in development, wound healing, or
cancer [1]. The process is primarily regulated by vascular endothelial growth factors (VEGFs) and
platelet-derived growth factors (PDGFs), which bind to their respective receptors, VEGFRs and
PDGFRs, which we describe as a uni-family model, initiating downstream angiogenic signaling.
Dysregulated angiogenesis can lead to over 70 vascular diseases and cancers [2, 3]. Current
therapies for excessive angiogenesis targeting VEGFs have demonstrated limited efficacy [4]. The
anti-VEGF-A antibody resulted in transitory improvements, such as tumor stasis and shrinkage.
However, after a period of weeks to months, the tumors typically resume their path of growth, and
the disease progresses [5]. Surprisingly, tumors resistant to anti-VEGF treatment showed PDGF
overexpression [4–6]. This finding suggests the involvement of other molecules in the pathological
process beyond the commonly recognized uni-family VEGF-VEGFR signaling axis.

A novel paradigm of cross-family signaling has been proposed and identified between VEGF and
PDGF families [7]. The two families share similarities structurally and functionally for both
ligands (VEGFs and PDGFs) and receptors (VEGFRs and PDGFRs) [8–11]. The ligands bind to
and activate their respective receptors in a similar way, followed by dimerization and tyrosine
phosphorylation [12–14]. Studies in literature have established evidence of cross-family signaling.
The bindings of VEGF-to-PDGFR and PDGF-to-VEGFR were both discovered with high
affinities [7], indicating a complex interplay between these growth factor families. VEGF-A has

shown the ability to activate PDGFRs in a dose-dependent manner, independently of VEGFRs
[15]. Investigations in human dermal fibroblasts lacking VEGFRs have revealed that VEGF-A can
modulate surface PDGFR levels and inhibit PDGF ligand interactions [15, 16]. Furthermore,
PDGF-CC, a homodimer structurally similar to VEGF-A [17], has displayed the capacity to
stimulate EC migration at a level comparable to VEGF-A [18]. Treatment of ECs with PDGFs has
been associated with reduced VEGFR2 expression [19]. Computational models have predicted that
PDGFs contribute significantly to VEGFR2 ligation in both healthy conditions and breast cancer
[7]. Taken together, these data underscore the potential impact of VEGF-PDGF cross-family
signaling in angiogenesis regulation. However, the resulting functional effects of PDGF-to-
VEGFR binding have not been fully elucidated.

In this study, we aimed to reveal the role of PDGFs in regulating endothelial signaling and function.
To achieve this, we used ECs that do not express PDGFRs on the plasma membrane to exclude
the effects of PDGF signaling through PDGFRs. As comparative control, we also used HDFs
lacking VEGFRs but expressing PDGFRs to elucidate the functional effects of PDGF signaling
through VEGFRs and PDGFRs. To exclude the possible VEGF upregulation in the treatment
medium, we validated the VEGF-A protein concentrations in the cell culture supernatant followed
by PDGF stimulations compared with untreated conditions. Our comprehensive investigation
assessed the activation of VEGFRs and their downstream effectors, i.e., PLCγ1, Akt, and FAK,
and examined the resulting functional roles in angiogenesis, such as cell proliferation and
migration. Through our research, we aim to shed light on alternative mechanisms that could
potentially address drug resistance issues observed in anti-VEGF therapies, offering new insights
into overcoming vascular dysregulation.

Material and Methods

Cell culture

Primary human brain microvascular endothelial cells (HBMECs) (Cell Systems, ACBRI 376) were cultured
in Complete Classic Medium (4Z0-500) supplemented with CultureBoost and Bac-Off Antibiotic (Cell
Systems, 4Z0-644). Primary human dermal microvascular endothelial cells (HDMECs) (ATCC, CRL-4060)
PDGFRA and *PDGFRB* double knock out (DKO) (to eliminate PDGFRα and β expression, respectively)
via CRISPR/Cas9 at the Genome Engineering & Stem Cell Center at Washington University in St. Louis,
MO. The efficiency of DKO was validated by next-generation sequencing (NGS) analysis (cite my qFlow
paper). HDMECs (*PDGFRA$^{-/-}$* and *PDGFRB$^{-/-}$*) were cultured in Vascular Cell Basal Medium (ATCC,
PCS100030) supplemented with microvascular endothelial cell growth kit -VEGF (i.e., with VEGF; ATCC,
PCS110041) and penicillin streptomycin (10,000 U/mL) (Thermo Fisher Scientific, 15-140-122). Human
dermal fibroblasts (HDFs) were purchased from ATCC (PCS-201-010) and cultured in DMEM (Thermo
Fisher Scientific, 11995065) with 5% FBS (Sigma-Aldrich, 12306C) and Penicillin Streptomycin (10,000
U/mL) (Thermo Fisher Scientific, 15-140-122). Cells in passages 3–6 were used and maintained in a
humidified incubator at 37 °C and with 5% CO_2. Low-serum medium used in this study for cell starvation
was made from EBM-2 (Lonza, CC-3156) supplemented with 1% FBS, Bac-Off Antibiotic (Cell Systems,
4Z0-644), and necessary components minus growth factors (hydrocortisone, ascorbic acid, GA-1000,
heparin).

Figure 1. ECs expressed few to no PDGFRα or PDGFRβ on either plasma membrane (A) and low levels in the whole cell (B). The numbers of PDGFRs resulted from nonspecific binding between PDGFR antibodies and PDGFRs. Error bars represent the standard error of mean (SEM). (*p<0.05, **p<0.01, ***p<0.001 as determined by one-way ANOVA and multi-comparison post-hoc Tukey's test.) The same method was used to calculate all p-values in this study.

Proliferation assay

Cells were seeded in a 96-well plate and grown to approximately 50% confluence and serum-starved
overnight using low-serum medium. Cells were stimulated with specific ligands, PDGF-AA, -AB, -BB, or
VEGF-A (R&D Systems, 221-AA-050, 222-AB-050, 220-BB-050, 293-VE-050/CF; same materials were
used throughout the study) at specific concentrations for 24 hours. XTT (2,3-Bis-(2-Methoxy-4-Nitro-5-
Sulfophenyl)-2H-Tetrazolium-5-Carboxanilide) (Invitrogen, X6493) was dissolved in EBM-2 (Lonza, CC-

3156) at 1 mg/mL and activated by PMS (N-methyl dibenzopyrazine methyl sulfate) (0.0075 mg/mL). The
reagent was added to each well and mixed with pipetting. The plates were incubated at 37 °C for 3 hours in
the dark. The absorbance was read at 450 and 650 nm. The final OD value is $[Abs_{450nm}(Test) -$
$Abs_{450nm}(Blank)] - Abs_{660nm}(Test)$ and normalized to the untreated control.

Figure 2. VEGFR1 was significantly phosphorylated by low concentrations of PDGF-AA and –BB in HBMECs, whereas no PDGFs induced VEGFR2 phosphorylation (Ty1175) in HBMECs. HBMECs were starved with low-serum medium for 6 hours and stimulated with growth factors at the indicated concentrations for 30 min (VEGFR1) or 10 min (VEGFR2). The receptor activation was picked up either by ELISA detecting the pan-tyrosine phosphorylation of VEGFR1 **(A–C)** or VEGFR2 **(D)**, or by immunoblotting and probed with primary antibody targeting VEGFR2 phosphorylation at tyrosine site 1175 **(E)**.

91 *Migration assay*

92 Cells were cultured in T-75 flasks until they reached approximately 90% confluence. Subsequently, the
93 cells were serum-starved overnight using low-serum medium. Transwell inserts (Falcon, 353097) were
94 coated with 2.45 μg/mL Collagen type I (Corning, 354236) in PBS and incubated overnight at 4°C. After
95 starvation, the cells were detached and seeded in the top chamber of the transwell inserts at a density of 1

Figure 3. High concentrations of PDGF-BB induced VEGFR1 phosphorylation in HDMECs (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$), whereas low concentrations of PDGF-AB activated VEGFR2. (A–G) HDMEC (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) were starved with low-serum medium for 6 hours and stimulated with growth factors at the indicated concentrations for 30 min (VEGFR1) or 10 min (VEGFR2). **(A–C)** The fold changes of VEGFR1 phosphorylation induced were assessed via ELISA. **(D–G)** The fold changes of VEGFR1 phosphorylation induced were assessed via immunoblotting. **(H)** The fold changes of PDGFRβ phosphorylation in HDFs induced by 30-min VEGF-A or PDGF-BB treatment were measured via ELISA.

$\times 10^5$ cells/well. The bottom chamber contained growth factors at the specified concentrations in low-serum
medium. The cultures were then incubated for 24 hours at 37 °C with 5% CO_2. Following incubation, the
unmigrated cells on the upper surface of the chamber were gently removed using a cotton swab. The inserts
were subsequently washed with PBS. The cells that had migrated through the membrane to the lower side
of the insert were fixed with 4% paraformaldehyde, permeabilized with 0.1% Tween-20, and stained with
a 0.01% Hoechst 3342 solution (Invitrogen, H3570). After a final wash with PBS, the inserts were imaged

under the DAPI channel (EVOS M5000). Cell migration was quantified using ImageJ (1.52a) as the number
of migrated cells to the downside of the inserts.

Enzyme-Linked Immunosorbent Assay (ELISA) / Immunology for phosphorylation detection

Cells were cultured till approximately 90% confluence and processed serum starvation for 6 hours using
low-serum medium. The indicated growth factors were added into the medium and incubated for 30 minutes
for detecting phosphor-VEGFR1 and phosphor-PDGFRβ or 10 minutes for detecting phosphor-VEGFR2.
Ttreatment medium was aspirated to terminate the stimulation, and the plates were washed with ice-cold
phosphate-buffered saline (PBS). For ELISA, cell lysates were collected with lysis buffer (R&D Systems,
DYC002) with protease and phosphatase inhibitor cocktail (Sigma-Aldrich, P8340). After being briefly
vortexed and centrifuged at 12,000$\times g$ for 10 min at 4 °C, the cell lysates went through the protocol of
Human Phospho-VEGFR1/Flt-1 DuoSet IC ELISA kit (R&D Systems, DYC4170), Human Phospho-
VEGFR2/KDR DuoSet IC ELISA kit (R&D Systems, DYC1766), or Human Phospho-PDGFRβ DuoSet
IC ELISA kit (R&D Systems, DYC1767) to assess the phosphorylation levels of VEGFR1, VEGFR2, or
PDGFRβ, respectively. The protein mass was quantified through bicinchoninic acid assay (BCA) (Pierce
Biotech, 23227). The results were normalized to untreated control samples and statistically analyzed.

For immunoblotting, the cells were scraped in ice-cold 1$\times$ RIPA lysis buffer (EMD Millipore, 20-188)
supplemented with protease and phosphatase inhibitor cocktails (Sigma-Aldrich, 11836170001 and
4906837001). The cell lysates were cleared by centrifuge at 12,000$\times g$ for 10 min at 4 °C, mixed with 6 $\times$
SDS buffer (Boston BioProducts, BP-111R), denatured at 95 °C for 10 min, resolved on 4–15% SDS-PAGE
gels (Bio-Rad, 4561084), and transferred to nitrocellulose membranes (Bio-Rad). The membranes were
blocked with 5% non-fat dry milk dissolved in 1$\times$ Tris Buffered Saline (TBS) for 1 hour at room
temperature before adding primary antibodies (Supplementary Table 1) overnight at 4 °C. Secondary
antibodies were added and incubated for 20 min at room temperature. Protein signals were detected using
the LI-COR Odyssey CLx Imager (LI-COR Biosciences) and quantified using ImageJ Software
(2.14.0/1.54f) or Image Studio 6.0 (LI-COR Biosciences). The protein mass was quantified through
bicinchoninic acid assay (BCA) (Pierce Biotech, 23227).

Enzyme-Linked Immunosorbent Assay (ELISA) for VEGF-A secretion detection

Cells were cultured till approximately 90% confluence and processed serum starvation for 6 hours using
low-serum medium. The indicated growth factors were added into the medium and incubated for 10 min,
30 min, or 24 hours. The cell culture supernatant was collected and centrifuged at 1000$\times g$ for 5 min. The
concentration of VEGF-A in the cell culture supernatant was detected with Human VEGF ELISA Kit -
Quantikine (R&D Systems, SVE00) following the instructions. The results were normalized to untreated
control samples and statistically analyzed.

Quantitative flow cytometry

ECs were seeded in T-75 flasks and grown to approximately 80% confluence. The cells were washed with
Ca^{2+}/Mg^{2+}-free PBS, harvested in Corning CellStripper solution, a nonenzymatic cell-dissociation solution,
and incubated for 15 min at 37 °C and 5% CO_2. The disassociation was stopped by adding Ca^{2+}/Mg^{2+}-free
PBS, and the cell suspension was centrifuged twice at 400$\times g$ for 5 min at 4 °C. The cells were resuspended
at 2–4 $\times10^6$ cells/mL in ice-cold stain buffer (PBS, bovine serum albumin, sodium azide). We added 25-μL
aliquots of a single-cell suspension for plasma membrane staining and 50 μL for whole-cell staining to 96-
well deep-well plates. The appropriate amount of human VEGFR1 phycoerythrin (PE)-conjugated antibody
(R&D Systems, FAB321P), VEGFR2 PE-conjugated antibody (BioLegend, 359904), VEGFR3 PE-
conjugated antibody (R&D Systems, FAB3492P), PDGFRα PE-conjugated antibody (R&D Systems,

FAB1264P), or PDGFRβ PE-conjugated antibody (R&D Systems, FAB1263P) was added to each sample
well except the fluorescence-minus-one (FMO) control. The saturating concentrations of PE-conjugated
antibodies to detect receptors on the plasma membrane were previously determined at 14 μg/mL for
VEGFR1, 2, and 3; and 9.4 μg/mL for PDGFRα and PDGFRβ [16, 20]. For whole-cell receptor detection,
the saturating concentration of VEGFR1 and VEGFR2 antibodies was determined at the beginning of the
study to be 20 ug/mL for both. Samples were incubated in dark for 40 min at 4 °C and washed twice with
stain buffer by centrifugation at 400×g for 5 min at 4 °C. Samples for plasma membrane receptor
measurement were washed and resuspended in 100 µL stain buffer before analysis.

Samples for whole-cell receptor measurement were fixed using 2% paraformaldehyde for 20 min in the
dark, washed, and permeabilized (0.5% Tween-20) for 20 min in the dark. Samples were washed twice with
stain buffer containing 0.1% Tween-20. PE-conjugated antibodies were added to the samples for staining
for 40 min in the dark, followed by three washes with stain buffer containing 0.1% Tween-20. Washed
samples for whole-cell receptor measurement were resuspended in 50 µL stain buffer. The readouts were
acquired by a Cytoflex S flow cytometer (Beckman Coulter, IN) and CytExpert software (Beckman Coulter,
IN). Kaluza software (Beckman Coulter, IN) was used for data analysis. Quantibrite PE beads (BD
Biosciences, 340495) were collected and analyzed under the same compensation and voltage settings as
cell fluorescence data. Quantibrite PE beads are a mixture of beads with four different concentrations of
conjugated PE molecules: low (474 PE molecules/bead), medium-low (5359 PE molecules/bead), medium-
high (23,843 PE molecules/bead), and high (62,336 PE molecules/bead). A calibration curve that translated
PE geometric mean to the number of bound molecules was determined using linear regression, $y = mx +$
c, where y is $\log_{10}$ fluorescence and x is $\log_{10}$ PE molecules per bead. Receptor levels were calculated as
described previously [20–23].

Figure 4. PDGFs activated adaptor proteins at levels comparable to those activated by VEGF-A. HDMECs (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) were treated with PDGF-AA, -AB, or -BB at the concentrations of 5 and 50 ng/mL for 15 min. The effects of PDGF treatment on phosphorylating PLCγ1 at tyrosine site 783 **(A-D)**, Akt at serine site 473 **(E-H)**, and FAK at tyrosine site 397 **(I-L)** were assessed via immunoblotting.

 Statistics

All data were acquired in at least three independent experiments. Ensemble averaged data are represented
by mean ± standard error of the mean (SEM). One-way analysis of variance (ANOVA), multiple
comparison post-hoc Tukey's test, and power analysis of pair-comparison were performed.

Results

Selection of VEGFR-expressing cell lines for investigating PDGF-induced signaling and functions

We selected human brain microvascular endothelial cells (HBMECs) to investigate the PDGF-induced
VEGFR signaling and its effects on endothelial functions. This selection was based on our quantification
of VEGFR expression across five human endothelial cell lines, which showed that HBMECs have the
highest surface expression of VEGFR1 while exhibiting similar levels of VEGFR2 (Figure 1C). Further, to
eliminate potential PDGF signaling through PDGFRs, we engineered a *PDGFRA* and *PDGFRB* double
knockout (PDGFR-DKO) human dermal microvascular endothelial cell (HDMEC) line using
CRISPR/Cas9. HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$) expressed < 100 PDGFRα/cell and < 100
PDGFRβ/cell on the plasma membrane (Figure 1A), which likely resulted from nonspecific binding as
described previously [22, 23]. Similarly, HBMECs exhibited negligible amounts of plasma membrane
PDGFRs (< 300 PDGFRs/cell) when compared to HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$) (p < 0.05).
The whole-cell PDGFR expressions were about 10 times higher than the expression levels on the plasma
membrane, likely because of increased nonspecific binding during whole-cell staining (Figure 1B). As a
control, we performed the same PDGF treatments on human dermal fibroblast (HDF) cells, which naturally
express PDGFRs but lack VEGFRs on their plasma membranes [24], to confirm the PDGF protein activity.

PDGF-AA and -BB induced VEGFR1 phosphorylation, whereas PDGF-AB phosphorylated
VEGFR2

We initially assessed angiogenic receptor activation by quantifying VEGFR1 and VEGFR2
phosphorylation through ELISA and immunoblotting under growth factor stimulation. We treated
HBMECs and HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$) with PDGF-AA, -BB, and -AB at increasing
concentrations from 0 to 100 ng/mL. HBMECs showed an overall descending trend as PDGF-AA and -BB
treatment concentrations increased (Figure 2A, 2C). We observed that 0.5 ng/mL PDGF-AA and PDGF-
BB treatments significantly increased VEGFR1 phosphorylation and presented a peak by a 1.8 and 2.2-fold
increase, respectively, compared with untreated HBMECs (p<0.001), which was similar to the changes of
VEGF-A treatment (Figure 2A, 2C). PDGF-AB did not have an effect on VEGFR1 phosphorylation (Figure
2B). We did not observe any PDGF-induced VEGFR2 phosphorylation in HBMECs, regardless of low or
high concentrations of PDGF (Figure 2D, 2E).

In HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$), VEGFR1 phosphorylation was induced to the greatest level
to 180% by 25 ng/mL PDGF-BB treatment (p < 0.001) (Figure 3C). PDGF-AB demonstrated a 1.8-fold
increase at 10 ng/mL (p < 0.05) (Figure 3B), whereas PDGF-AA did not show any significant effects
(Figure 3A). Phosphorylation of VEGFR2 at tyrosine site 1175 was highly induced by 5 ng/mL PDGF-
AB to 200% elevation after 15-min stimulation (Figure 3F), whereas the other two PDGFs showed no
significant impacts (Figure 3E, 3G). VEGF-A induced the activation to a greater extent to 360%–400%
increase at a relatively high concentration of 50 ng/mL, aligning with the observations from ELISA
conducted in HBMECs (Figure 3D, 2D).

In HDFs, PDGFRβ phosphorylation was assessed to technically validate PDGF treatment. PDGF-BB
induced robust PDGFRβ phosphorylation (Figure 3H). The significant effects were detectable at 5 ng/mL
and peaked at 100 ng/mL, where the phosphorylation level was 8-fold higher compared with untreated cells.
In contrast, treatment with 100 ng/mL VEGF or 0.5 ng/mL PDGF-BB resulted in low PDGFRβ
phosphorylation, approximately 1.5-fold higher than control levels (Figure 3H).

Figure 5. Low concentrations of PDGF significantly induced HBMEC proliferation up to 240%.
HBMECs were treated with PDGF-AA, -AB, -BB, or VEGF-A with indicated concentration for 24
hours and subjected to proliferation assay. XTT (2,3-Bis-(2-Methoxy-4-Nitro-5-Sulfophenyl)-2H-
Tetrazolium-5-Carboxanilide) was used to quantify cell numbers by detecting cell viability. The
absorbance was read at 450 and 650 nm. OD value for each condition was normalized by untreated

PDGFs activated adaptor proteins in HDMECs to a level comparable to that of VEGF-A

The effects of PDGF-AA, -AB, or -BB (5 and 50 ng/mL) on downstream protein activation (i.e., PLC γ1,
Akt, and FAK) were assessed in HDMECs (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) via immunoblotting. Both
PDGF-AA and -AB concentrations significantly phosphorylated PLCγ1 at tyrosine site 783 ($p < 0.001$) by
similar fold changes to 200%–225% within 30-min treatment, which is comparable to the effects by 50
ng/mL VEGF-A (Figure 4A–4C). In contrast, 5 ng/mL of VEGF-A induced PLCγ1 phosphorylation to over
300% increase ($p < 0.001$) (Figure 4A). PDGF-BB, however, showed a subtle phosphorylation elevation
to1.2-fold at 50 ng/mL after 60-min incubation ($p < 0.05$) (Figure 4D). The phosphorylation of Akt at serine
site 473 showed the highest increase of all investigated growth factors to a 6–15-fold elevation ($p < 0.001$)
(Figure 4E–4H). Notably, PDGF-AA and -AB at 5 ng/mL showed Akt activation at a similar level to VEGF-
A at 50 ng/mL, whereas PDGF-BB showed the highest level to 15-fold elevation at 5 ng/mL after 15-min
treatment (Figure 4E–4H). Growth factors showed the lowest impacts on the activation of FAK at tyrosine
site 397 with a 120–200% increase (Figure 4I–4L). A high concentration of 50 ng/mL PDGF-AB displayed

a 2-fold increase in FAK phosphorylation (p < 0.001), comparable to that of 50 ng/mL VEGF-A after 30-
min treatment (p < 0.05) (Figure 4I, 4K). PDGF-BB showed a minor increase at 1.2 times (p < 0.05),
whereas PDGF-AA did not significantly phosphorylate FAK at the site (p = 0.86) (Figure 4J, 4L).
229
PDGFs showed significant capacities on stimulating EC proliferation comparable to VEGF-A

The activation of PLCγ1, Akt, and FAK has been found to modulate VEGF-induced EC proliferation,
migration, and survival [25–28], which served as the foundation for exploring PDGF-induced EC
proliferation. Serum-starved cells were treated with 0.5 ng/mL to 200 ng/mL VEGF-A or PDGF. Notably,
all studied PDGFs (PDGF-AA, -AB, and -BB) significantly stimulated EC proliferation. HBMECs treated
with PDGF-AA or -AB exhibited up to a 150% increase in proliferation when the concentrations were
between 0.5 ng/mL and 50 ng/mL (p < 0.01) (Figure 5A, 5B), while PDGF-BB exhibited an inverse dose-
response relationship (Figure 5C): the low PDGF-BB concentration, 0.5 ng/mL, induced the highest
HBMEC proliferation, resulting in a remarkable 240% increase compared with untreated cells (p < 0.001)
(Figure 5C). As expected, VEGF-A treatment served as a positive control, demonstrating a substantial
increase in cell growth, reaching up to 200% at concentrations ranging from 0.5 ng/mL to 100 ng/mL
(Figure 5D). HDMEC (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) showed significant proliferation increases relative to

Figure 6. HDMEC (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) showed a comparable increase in cell proliferation to around 150% induced by PDGF or VEGF-A across all investigated concentrations.

Figure 7. HDF exhibited a more remarkable proliferation increase to 280% by PDGF-BB or 250% by VEGF-A at higher concentrations.

untreated conditions across all investigated concentrations of PDGF or VEGF-A ($p < 0.001$). Notably this
increased proliferation was similar across all concentrations (Figure 6). While PDGFs can induce EC
proliferation, the PDGF-BB-mediated proliferation response in HDFs was more pronounced ($p < 0.001$)
with up to a 280% increase in proliferation relative to the untreated control across the 5 ng/mL to 100 ng/mL
concentration range (Figure 7A). Notably, VEGF-A mediated a significant proliferation response in HDFs
($p < 0.001$), relative to the untreated control, at concentrations between 10 ng/mL to 200 ng/mL, promoting
proliferation by 250% (Figure 7B).

Figure 8. PDGFs induced significant and comparable enhancement in HBMEC migration across a wide range of concentrations. HBMECs were seeded in the top chamber of transwell inserts and stimulated with PDGF-AA, -AB, -BB, or VEGF-A at the indicated concentrations in low-serum medium in the lower chamber for 24 hours. The cells that migrated through the membrane to the downside of the insert were fixed, permeabilized, stained, imaged, and counted on the following day.

PDGF induced EC migration to 70% of the level stimulated by VEGF-A

To investigate the impact of PDGF on cell migration, we quantified the number of migrated cells through
the transwell inserts stimulated by growth factors across the 0.5 ng/mL to 100 ng/mL concentration range
for 24 hours. All investigated PDGF significantly triggered EC migration. In HBMECs, we observed that
25 ng/mL PDGF-AA or 10 ng/mL PDGF-AB induced the highest level of migration to 170% and 160%
compared with untreated cells, respectively ($p < 0.001$) (Figure 8A, 8B). PDGF increased HBMEC

migration to 130% at 0.5 ng/mL (p < 0.001) (Figure 8C). In HDMEC ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$), low
concentrations of PDGF-AA and -AB resulted in the most remarkable migration with a 130% enhancement
relative to the untreated control at 5 ng/mL and 0.5 ng/mL, respectively (p < 0.01 and p < 0.001, respectively)
(Figure 9A, 9B). PDGF-BB exhibited less migration with increasing concentrations in both EC types, with
the lowest concentration, 0.5 ng/mL PDGF-BB resulting in a significant, 1.3-fold increase in migration
relative to the untreated control in HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$) (p < 0.001) (Figure 9C).
Conversely, increasing PDGF-BB concentrations correlated with increasing migration of HDFs, starting
from 5 ng/mL and peaking with a 2.8-fold increase in migration at 100 ng/mL (p < 0.001) (Figure 9D).
VEGF-A served as a positive control with a substantial increase of up to 230% in both ECs (p < 0.001) but
failed to show significant changes in HDFs (p = 0.09).

PDGF-induced EC responses are independent of VEGF-A upregulation

To ascertain the potential for VEGF-A protein to have induced the PDGF-changes observed via PDGF-
mediated upregulation of VEGF-A [4,43], we conducted an ELISA to quantify VEGF-A concentrations in
the cell culture supernatant following varying durations of PDGF-BB treatment. Our analysis revealed no
statistically significant changes in VEGF-A levels between the untreated control and PDGF-BB-treated
experimental cohorts (Figure 10B). Interestingly, within the VEGF-A positive subgroup, a progressive
decline in VEGF-A concentration was noted over the course of incubation (Figure 10A). The trajectory of
VEGF-A concentration reduction was accurately delineated by a regression curve (Figure 10C), with a
calculated half-life of 32.33 minutes, suggesting the temporal stability of VEGF-A in the culture media
after VEGF-A treatment.

Discussion

Our study elucidates the effects of cross-family PDGF-to-VEGFR binding in regulating
endothelial signaling and functions. We found that (1) PDGFs played a critical role in augmenting EC
angiogenic hallmarks, i.e., receptor and downstream protein phosphorylation, cell proliferation, and
migration, through non-PDGF receptors, to a similar level compared to VEGF-A. (2) Low concentrations
of PDGF-BB induced the angiogenic capacity of ECs to the greatest extent, whereas high concentrations
showed a more potent effect in HDF. Other PDGFs did not exhibit such trends. (3) The angiogenic effects
of PDGF-BB in HBMECs were more prominent than in HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$). Other
PDGFs showed similar impacts between EC types. (4) VEGF-A protein upregulation was not observed
upon PDGF-BB stimulation, further validating the effects of PDGF-to-VEGFR signaling in ECs.

PDGF-induced endothelial signaling and function

Cross-family interaction has been identified between PDGF and VEGFR as a novel angiogenic mechanism
[7], the physiological functions are, however, yet unexplored. Here, we examined the angiogenic hallmarks
of cross-family PDGF-to-VEGFR binding in regulating endothelial signaling and functions, i.e., receptor
and effector activation, cell proliferation, and migration. We employed ECs that lack PDGF receptors
(PDGFRs) on their plasma membranes to isolate the effects of possible PDGF-BB signaling through
VEGFR2 from its signaling through PDGFR and also employed human fibroblasts that lack VEGFRs but
have PDGFRs on their plasma membranes as a comparative control. To exclude the possible effects of
upregulated VEGF-A transcription induced by PDGF treatment [29, 30], we observed no significant
changes in VEGF-A protein concentrations in the cell culture supernatant after varying durations of PDGF-

BB treatment, further confirming that the observed effects resulted from PDGF signaling through non-PDGFRs.

Figure 9. Migrative effects of PDGFs in HDMECs (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) and HDF. (A-C) Low concentrations of PDGF significantly induced HDMEC (*PDGFRA*$^{-/-}$ and *PDGFRB*$^{-/-}$) migration by approximately 1.3-fold, comparable to HBMEC. **(D)** Higher concentrations (> 5 ng/mL) of PDGF-BB promoted a significant increase in HDF migration by 2.8-fold.

Low concentrations of PDGF showed more pronounced effects

Our detailed analyses revealed that low concentrations of PDGFs induced EC functional changes at a higher level. Specifically, treatment of 0.5 ng/mL PDGF-AA or -BB induced the highest VEGFR1 phosphorylation concentration by approximately 2-fold in HBMECs, whereas high concentrations at 50–100 ng/mL yielded the lowest receptor activation. Around 5 ng/mL of PDGF-AA and -AB activated downstream proteins, PLCγ1, Akt, and VEGFR2 at the investigated sites at a higher level than the concentration of 50 ng/mL. A low concentration of PDGF-BB at 0.5 ng/mL exhibited stronger activation of Akt and FAK compared to a high concentration at 50 ng/mL. Similar results were observed in the examinations of angiogenic hallmarks, where the highest impacts of PDGF treatment appeared at relatively low concentrations. Notably, PDGF in human blood serum is measured to be 17.5 ng/mL [31],

approximately 35-fold higher than 0.5 ng/mL, the concentration producing the most substantial angiogenic effects, which may explain why the functions of PDGF were long overlooked.

Figure 10. The concentration of VEGF-A exhibited no significant alteration upon PDGF-BB stimulation. (A and B) Subsequent to a 6-hour period of serum deprivation (1% FBS), HBMECs underwent treatment with growth factors as detailed. Post-incubation, the cell culture supernatant underwent centrifugation and analysis as per the protocols outlined in the Human VEGF ELISA Kit – Quantikine (R&D Systems, SVE00). **(C)** A fitting curve was generated to depict the correlation between VEGF-A concentration and in-vitro incubation duration. The determination of VEGF-A half-life time was computed utilizing the formula $t = -t1 \cdot \ln 2$, yielding a value of 26.76 minutes.

PDGF-to-VEGFR1 signaling

Our observations suggest a compelling link between PDGF-induced signaling through VEGFRs, downstream proteins, and EC angiogenic hallmarks. PLCγ1, Akt, and FAK are downstream proteins after VEGFR2 activation induced by VEGF [25]. They are found to involve the signaling that regulates EC

proliferation, survival, and migration [25–28]. In this study, we found that PDGFs could significantly
activate VEGFRs and these proteins and induced EC angiogenic hallmarks. It is suggested that PDGF may
signal through VEGFR1 and downstream proteins to regulate EC angiogenesis.

Our observations suggest that ECs may engage in novel signaling pathways between PDGF and non-PDGF
receptors on the plasma membrane. PDGF-BB was suggested to induce VEGFR1 and VEGFR2
phosphorylation through VEGFR2 binding, since PDGF-BB-to-VEGFR1 binding was not previously
observed, but PDGF-BB-to-VEGFR2 binding was seen [7]. In addition, VEGFR1 primarily
heterodimerizes with VEGFR2 to regulate VEGF activity [13, 32]. Most VEGFR1 are in heterodimer forms
with VEGFR2 on the plasma membrane of ECs [33]. These findings suggest the need to explore VEGFR1-
VEGFR2 heterodimerization in PDGF-mediated cross-family interactions.

VEGFR1 has been typically considered a "decoy" receptor in angiogenesis due to its high binding affinity
but limited kinase activity [34, 35]. However, an association between increased VEGFR1 levels and
angiogenesis hallmarks was recently found [21, 24, 36–38], challenging the decoy paradigm. VEGFR1 is
upregulated in tumors and ischemia, but VEGFR2 remains stable [21, 38]. Here, the VEGFR activity on
ECs induced by PDGFs is the first time shown. PDGFs could significantly activate and phosphorylate
VEGFR1 and VEGFR2 in ECs.

PDGFs function variously in different cells

PDGF-BB showed more pronounced angiogenic effects in HBMECs compared with HDMECs
($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$). Treatment with PDGF-BB led to a 220% increase in VEGFR1
phosphorylation in HBMECs, contrasting with a 180% increase in HDMECs ($PDGFRA^{-/-}$ and
$PDGFRB^{-/-}$). Moreover, cell proliferation in HBMECs surged by 240% compared to 150% in HDMECs
following PDGF-BB treatment, potentially influenced by the twofold higher concentration of VEGFR1 on
the plasma membrane of HBMECs (700 ± 70 VEGFR1/cell) compared to HDMECs (300 ± 40
VEGFR1/cell) [cite]. Moreover, heparin can inhibit growth factor effects on cell proliferation [39]. The
medium culturing HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$) contain a significant amount of heparin
(0.0015 mg/mL).

The more remarkable effects of PDGF at higher concentrations in HDFs were observed compared with the
results in ECs with PDGF-BB at lower concentrations. PDGF-BB increased VEGFR1 phosphorylation by
around 2-fold at 0.5 ng/mL and 8-fold at 100 ng/mL in ECs and HDFs, respectively. Cell proliferation rate
was elevated by 240% in HBMECs at 0.5 ng/mL compared with 280% in HDFs by 100 ng/mL of PDGF-
BB treatment. Cell migration was promoted to 130% enhancement by 0.5 ng/mL PDGF-BB in HBMECs
and HDMECs ($PDGFRA^{-/-}$ and $PDGFRB^{-/-}$), whereas HDFs reached the peak migration capacity by 100
ng/mL PDGF-BB of 2.8-fold increase. The data agreed with previous publications that strong cell
proliferation induction only occurs with PDGF concentrations > 5 ng/mL and arises as the concentration
increases [40]. PDGF-BB at 25 ng/mL was found to increase fibroblast proliferation by approximately 2-
fold [41], comparable to our findings. The cell migration response is negligible at low concentrations and
arises starting from 1 ng/mL [40]. PDGF-BB at 10 ng/mL increases fibroblast migration to around 200%
[42], similar to our result. The less remarkable PDGF-induced impacts in ECs aligned with the differential
receptor expression on these cells, as HDFs express PDGFRs on the plasma membrane, but ECs do not. In
such case, PDGF displayed less activity on ECs than HDFs due to the lack of plasma-membrane PDGFRs.

Our study highlighted significant VEGF-A-induced effects in HDFs lacking VEGFRs. VEGF-A
significantly phosphorylated VEGFR1 and enhanced cell proliferation of HDFs, but not on cell migration.
The validated impacts of VEGF-A in HDFs that lack VEGFRs but express PDGFRs on the plasma

membrane, agreed with the published studies that VEGF-A could signal through PDGFRs [15] and
demonstrated the effects of cross-family signaling between VEGF-A to PDGFRβ.

Conclusion

By uncovering the angiogenic functions of PDGF-VEGFR interactions, we can better understand the cross-
family angiogenic pathway, which may reveal alternate mechanisms for addressing vascular dysregulation.
The pathways that downregulate VEGFR1 may be responsible for limiting excessive angiogenesis (cite).
Proteomics is a reasonable attempt to systematically explore the underlying mechanisms. Our findings also
provide essential data for computational modeling and systems biology to refine the signaling pathways.

References

365   1.   Abhinand CS, Raju R, Soumya SJ, Arya PS, Sudhakaran PR (2016) VEGF-A/VEGFR2 signaling
366   network in endothelial cells relevant to angiogenesis. J Cell Commun Signal 10:347–354

367   2.   Weddell JC, Imoukhuede PI (2018) Computational systems biology for the VEGF family in
368   angiogenesis. In: Encyclopedia of Cardiovascular Research and Medicine. Elsevier, pp 659–676

369   3.   Egginton S (2010) Angiogenesis - may the force be with you! J Physiol 588:4615–4616

370   4.   Francia G, Emmenegger U, Kerbel RS (2009) Tumor-associated fibroblasts as ``Trojan horse''
371   mediators of resistance to anti-VEGF therapy. Cancer Cell 15:3–5

372   5.   Bergers G, Hanahan D (2008) Modes of resistance to anti-angiogenic therapy. Nat Rev Cancer
373   8:592–603

374   6.   Crawford Y, Kasman I, Yu L, Zhong C, Wu X, Modrusan Z, Kaminker J, Ferrara N (2009) PDGF-C
375   mediates the angiogenic and tumorigenic properties of fibroblasts associated with tumors
376   refractory to anti-VEGF treatment. Cancer Cell 15:21–34

377   7.   Mamer SB, Chen S, Weddell JC, Palasz A, Wittenkeller A, Kumar M, Imoukhuede PI (2017)
378   Discovery of High-Affinity PDGF-VEGFR Interactions: Redefining RTK Dynamics. Sci Rep 7:.
379   https://doi.org/10.1038/s41598-017-16610-z

380   8.   Li X, Eriksson U (2003) Novel PDGF family members: PDGF-C and PDGF-D. Cytokine Growth Factor
381   Rev 14:. https://doi.org/10.1016/S1359-6101(02)00090-4

382   9.   Abu-Duhier FM, Goodeve AC, Care RS, Gari M, Wilson GA, Peake IR, Reilly JT (2003) Mutational
383   analysis of class III receptor tyrosine kinases (C-KIT, C-FMS, FLT3) in idiopathic myelofibrosis. Br J
384   Haematol 120:464–470

385   10.   Smith GA, Fearnley GW, Tomlinson DC, Harrison MA, Ponnambalam S (2015) The cellular
386   response to vascular endothelial growth factors requires co-ordinated signal transduction,
387   trafficking and proteolysis. Biosci Rep 35:. https://doi.org/10.1042/BSR20150171

388   11.   Heldin C-H, Lennartsson J (2013) Structural and Functional Properties of Platelet-Derived Growth
389   Factor and Stem Cell Factor Receptors. Cold Spring Harb Perspect Biol 5:.
390   https://doi.org/10.1101/cshperspect.a009100

391   12.   Muller YA, Li B, Christinger HW, Wells JA, Cunningham BC, de Vos AM (1997) Vascular endothelial
392   growth factor: Crystal structure and functional mapping of the kinase domain receptor binding
393   site. Proceedings of the National Academy of Sciences 94:.
394   https://doi.org/10.1073/pnas.94.14.7192

395   13.   Koch S, Claesson-Welsh L (2012) Signal transduction by vascular endothelial growth factor
396   receptors. Cold Spring Harb Perspect Med 2:a006502–a006502

397   14.   Heldin C-H (2013) Targeting the PDGF signaling pathway in tumor treatment. Cell Communication
398   and Signaling 11:. https://doi.org/10.1186/1478-811X-11-97

399   15.   Ball SG, Shuttleworth CA, Kielty CM (2007) Vascular endothelial growth factor can signal through
400   platelet-derived growth factor receptors. J Cell Biol 177:489–500

401   16.   Chen S, Guo X, Imarenezor O, Imoukhuede PI (2015) Quantification of VEGFRs, NRP1, and
402   PDGFRs on Endothelial Cells and Fibroblasts Reveals Serum, Intra-Family Ligand, and Cross-Family
403   Ligand Regulation. Cell Mol Bioeng 8:. https://doi.org/10.1007/s12195-015-0411-x

404   17.   Reigstad LJ, Sande HM, Fluge Ø, Bruland O, Muga A, Varhaug JE, Martnez A, Lillehaug JR (2003)
405   Platelet-derived growth factor (PDGF)-C, a PDGF family member with a vascular endothelial
406   growth factor-like structure. J Biol Chem 278:17114–17120

407   18.   Li X, Tjwa M, Moons L, Fons P, Noel A, Ny A, Zhou JM, Lennartsson J, Li H, Luttun A, Pontén A,
408   Devy L, Bouché A, Oh H, Manderveld A, Blacher S, Communi D, Savi P, Bono F, Dewerchin M,
409   Foidart J-M, Autiero M, Herbert J-M, Collen D, Heldin C-H, Eriksson Ulf and Carmeliet P (2005)
410   Revascularization of ischemic tissues by PDGF-CC via effects on endothelial cells and their
411   progenitors. J Clin Invest 115:118–127

412   19.   Liu T, Ma W, Xu H, Huang M, Zhang D, He Z, Zhang L, Brem S, O'Rourke DM, Gong Y, Mou Y,
413   Zhang Z, Fan Y (2018) PDGF-mediated mesenchymal transformation renders endothelial
414   resistance to anti-VEGF treatment in glioblastoma. Nat Commun 9:.
415   https://doi.org/10.1038/s41467-018-05982-z

416   20.   Imoukhuede PI, Popel AS (2011) Quantification and cell-to-cell variation of vascular endothelial
417   growth factor receptors. Exp Cell Res 317:955–965

418   21.   Imoukhuede PI, Popel AS (2014) Quantitative fluorescent profiling ofVEGFRs reveals tumor cell
419   and endothelial cell heterogeneity in breast cancer xenografts. Cancer Med 3:225–244

420   22.   Chen S, Weddell J, Gupta P, Conard Grace and Parkin J, Imoukhuede PI (2017) QFlow cytometry-
421   based receptoromic screening: A high-throughput quantification approach informing biomarker
422   selection and nanosensor development. In: Methods in Molecular Biology. Springer New York,
423   New York, NY, pp 117–138

424   23.   Fang Y, Malik M, England SK, Imoukhuede PI (2022) Absolute quantification of plasma membrane
425   receptors via quantitative flow cytometry. In: Methods in Molecular Biology. Springer US, New
426   York, NY, pp 61–77

427   24.   Chen S, Imoukhuede PI (2019) Single-cell receptor quantification of an in vitro coculture
428   angiogenesis model reveals VEGFR, NRP1, Tie2, and PDGFR regulation and endothelial
429   heterogeneity. Processes (Basel) 7:356

430   25.   Olsson A-K, Dimberg A, Kreuger J, Claesson-Welsh L (2006) VEGF receptor signalling ? in control
431   of vascular function. Nat Rev Mol Cell Biol 7:359–371

432   26.   Abedi H, Zachary I (1997) Vascular endothelial growth factor stimulates tyrosine phosphorylation
433   and recruitment to new focal adhesions of focal adhesion kinase and paxillin in endothelial cells. J
434   Biol Chem 272:15442–15451

27. Le Boeuf F, Houle F, Huot J (2004) Regulation of vascular endothelial growth factor receptor 2-
mediated phosphorylation of focal adhesion kinase by heat shock protein 90 and src kinase
activities. J Biol Chem 279:39175–39185

28. Fujio Y, Walsh K (1999) Akt mediates cytoprotection of endothelial cells by vascular endothelial
growth factor in an anchorage-dependent manner. J Biol Chem 274:16349–16354

29. Finkenzeller G, Marmé D, Weich HA, Hug H (1992) Platelet-derived growth factor-induced
transcription of the vascular endothelial growth factor gene is mediated by protein kinase C.
Cancer Res 52:4821–4823

30. Uemura A, Fruttiger M, D'Amore Patricia A and De Falco S, Joussen AM, Sennlaub Florian and
Brunck LR, Johnson KT, Lambrou George N and Rittenhouse KD, Langmann T (2021) VEGFR1
signaling in retinal angiogenesis and microinflammation. Prog Retin Eye Res 84:100954

31. Bowen-Pope DF, Malpass TW, Foster DM, Ross R (1984) Platelet-derived growth factor in vivo:
levels, activity, and rate of clearance. Blood 64:458–469

32. Cudmore MJ, Hewett PW, Ahmad S, Wang K-Q, Cai M, Al-Ani B, Fujisawa T, Ma B, Sissaoui S,
Ramma W, Miller MR, Newby DE, Gu Y, Barleon B, Weich H, Ahmed A (2012) The role of
heterodimerization between VEGFR-1 and VEGFR-2 in the regulation of endothelial cell
homeostasis. Nat Commun 3:

33. Cai M, Wang K, Murdoch CE, Gu Yuchun and Ahmed A (2017) Heterodimerisation between
VEGFR-1 and VEGFR-2 and not the homodimers of VEGFR-1 inhibit VEGFR-2 activity. Vascul
Pharmacol 88:11–20

34. Carmeliet P, Moons L, Luttun A, Vincenti V, Compernolle V, De Mol M, Wu Y, Bono F, Devy L, Beck
H, Scholz D, Acker T, DiPalma T, Dewerchin M, Noel A, Stalmans I, Barra A, Blacher S,
Vandendriessche Thierry and Ponten A, Eriksson U, Plate KH, Foidart J-M, Schaper W, Charnock-
Jones DS, Hicklin DJ, Herbert J-M, Collen D, Persico MG (2001) Synergism between vascular
endothelial growth factor and placental growth factor contributes to angiogenesis and plasma
extravasation in pathological conditions. Nat Med 7:575–583

35. Sawano A, Iwai S, Sakurai Y, Ito M, Shitara K, Nakahata T, Shibuya M (2001) Flt-1, vascular
endothelial growth factor receptor 1, is a novel cell surface marker for the lineage of monocyte-
macrophages in humans. Blood 97:785–791

36. Fang Y, Chen L, Imoukhuede PI (2023) Toward blood-based precision medicine: Identifying age-
sex-specific vascular biomarker quantities on circulating vascular cells. Cell Mol Bioeng 16:189–
204

37. Weddell JC, Imoukhuede PI (2014) Quantitative characterization of cellular membrane-receptor
heterogeneity through statistical and computational modeling. PLoS One 9:e97271

38. Imoukhuede PI, Dokun AO, Annex BH, Popel AS (2013) Endothelial cell-by-cell profiling reveals
the temporal dynamics of VEGFR1 and VEGFR2 membrane localization after murine hindlimb
ischemia. Am J Physiol Heart Circ Physiol 304:H1085–H1093

39. Herrick WG, Rattan S, Nguyen T V, Grunwald MS, Barney CW, Crosby AJ, Peyton SR (2015)
Smooth muscle stiffness sensitivity is driven by soluble and insoluble ECM chemistry. Cell Mol
Bioeng 8:333–348

40. De Donatis A, Comito G, Buricchi F, Vinci MC, Parenti A, Caselli A, Camici G, Manao G, Ramponi G,
Cirri P (2008) Proliferation versus migration in platelet-derived growth factor signaling. J Biol
Chem 283:19948–19956

41. Takamura N, Renaud L, da Silveira WA, Feghali-Bostwick C (2021) PDGF promotes dermal
fibroblast activation via a novel mechanism mediated by signaling through MCHR1. Front
Immunol 12:

42. Li L, Heldin C-H, Heldin P (2006) Inhibition of platelet-derived growth factor-BB-induced receptor
activation and fibroblast migration by hyaluronan activation of CD44. J Biol Chem 281:26512–
26519

484