TITLE

Rescuing bacterial genome replication: essential functions to repair a double-strand break

and restart DNA synthesis

AUTHORS

Charles Winterhalter[*], Stepan Fenyk, Heath Murray[**]

AFFILIATION

Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle

Upon Tyne, NE2 4AX, UK

AUTHOR LIST FOOTNOTES

*Correspondence: Charles.winterhalter@newcastle.ac.uk

**Correspondence: Heath.murray@newcastle.ac.uk

HIGHLIGHTS

- Single-strand discontinuities inactivate the bacterial replisome
- Nicks on the leading or lagging strand template differentially affect fate of the helicase
- Double-strand breaks are repaired via homologous recombination and elicit the replication restart pathway
- Resection often generates a DNA strand with a 5′-end that must be processed to ensure productive helicase reloading

SUMMARY

Many antibiotics indirectly generate reactive oxygen species (ROS) that can damage bacterial genomes. Oxidised nucleobases become genotoxic when they are targeted for repair through excision, generating a single-strand discontinuity that can be converted to a double-strand break (DSB) by an oncoming replication fork. Because the genomic location of nucleobase oxidation is stochastic, investigating the fate of DNA replication machinery (replisome) at single-strand discontinuities has been limited. Here we have addressed this issue by expressing Cas9 nickases in *Bacillus subtilis* to create site specific single-strand discontinuities in a bacterial chromosome. We find that nicks in either leading or lagging strand arrest bacterial replication fork progression and generate a DSB that requires repair using homologous recombination to allow replication restart. These discoveries provoke reassessment of the fundamental mechanism of bacterial homologous recombination and provide insights to the development of alternative antimicrobials by identifying a specific pathway that can potentiate ROS-dependent bacterial killing.

KEYWORDS

DNA, replication, nick, helicase, fork, homologous repair, recombination, PriA, restart.

INTRODUCTION

Emergence of antibiotic resistant bacterial pathogens represents an urgent threat to human health and food production[1]. Understanding how bacteria respond to antibiotics is necessary to generate effective treatments. While antibiotics generally target specific essential bacterial systems (e.g. cell wall synthesis, translation, transcription) to inhibit bacterial proliferation, there is growing evidence that diverse antibiotics share the potential to disrupt bacterial physiology, thereby generating a second challenge to pathogens[2]. Work in several bacterial species indicates that a range of antibiotics ultimately generate reactive oxygen species (ROS) which can damage biomolecules including DNA[3,4]. Importantly, antibiotics that generate ROS elicit the bacterial DNA damage response, indicating the presence of double-strand breaks (DSB) under these conditions[4-7].

The potency of ROS-mediated DNA damage has been suggested to arise from conflicts between genome repair and replication systems[4-6,8]. Oxidised nucleobases located within the genome must be excised as part of the requisite repair pathway, leading to transient single-strand discontinuities in the DNA template[9]. If a bacterial replication fork arrives at a single-strand discontinuity, the encounter generates a DSB[10]. Elegant experiments using bacteriophage λ demonstrated that an engineered nick in either strand of the viral genome generated replication-dependent DSBs (Fig. 1A)[11]. While active DNA replication is necessary to generate DSBs at a nick, the fate of the bacterial DNA replication machinery (replisome) at single-strand discontinuities has not been determined.

The replisome is a multicomponent complex of proteins and enzymes that unwinds the chromosome and coordinates synthesis of new DNA polymers on the leading and lagging strands[12,13]. The most stable component of the replisome and most pivotal for its processivity is helicase[14,15]. Cellular replicative helicases are toroid enzymes that unwind DNA by encircling and translocating along a single client nucleic acid strand, acting as a wedge to disrupt base pairing[16]. Studies of bacterial replisomes *in vitro* and *in vivo* have revealed that the machinery is intrinsically dynamic, with most factors observed to bind and unbind interaction partners multiple times per replication cycle[17,18]. This behaviour enables

the replisome to overcome a range of potential roadblocks and remain associated with the

template, competent to continue DNA synthesis[19]. However, recent studies in eukaryotic

systems have indicated that single-strand discontinuities generate DSBs and arrest DNA

replication by impairing helicase progression[20,21].

Determining how bacteria react and respond when a replication fork encounters a

single-strand discontinuity is crucial to understanding how ROS inhibits bacterial

proliferation. Here we have investigated this question by expressing Cas9 nickases (nCas9)

in the model bacterium *Bacillus subtilis*. The results show that a nick located in either

template strand blocks downstream DNA synthesis, and that resumption of DNA replication

is dependent upon a core set of homologous recombination proteins and the PriA-dependent

replication restart system necessary to reload the replicative helicase. Unexpectedly,

interrogation of the DNA repair pathway suggests that processing of the DSB often leaves

the 5′-terminated DNA strand intact, which can hybridise to the homologous chromosome

during recombination and perturb replication restart. We ascribe the enigmatic RecG

helicase as the factor that remodels these recombination products containing a 5′-terminated

DNA strand, thereby generating the replication fork structure recognised by PriA for

productive helicase loading. These findings have critical implications for the fundamental

mechanism of homologous recombination in bacteria and the aim of potentiating ROS-

dependent antibiotic toxicity.

RESULTS

Single-strand discontinuities in a bacterial chromosome arrest DNA synthesis

Oxidised nucleobases generated by ROS can be located randomly throughout a genome, making it a challenge to directly analyse conflicts with DNA replication during base excision repair. To mimic the activity of oxidised base repair, an experimental system capable of generating a site-specific single-strand discontinuity was developed in *B. subtilis* using Cas9 nuclease variants and single-guide RNAs (sgRNA)[20-23]. Cas9 carrying single amino acid substitutions (Cas9^{D10A} or Cas9^{H840A}) created nickases (nCas9), while the double mutant (Cas9$^{D10A/H840A}$) created a catalytically inactive enzyme (dCas9)[24]. Each *cas9* allele was integrated into the *B. subtilis* chromosome under the control of a xylose-inducible promoter (Fig. 1B)[25]. The location of sgRNA hybridisation is indicated by coordinates between the origin (0°) and terminus (~180°) on either the right (+) or left (-) replicore of the chromosome. To avoid complications arising from potential activation of prophage following DNA damage in the laboratory strain of *B. subtilis*, a derivative lacking these genetic elements (Δ6) was used throughout this study[26].

Cell viability was quantified using spot-titre assays, where strains are grown to saturation in liquid medium without xylose, before samples were serially diluted and plated onto solid medium with or without xylose to enumerate the number of colony forming units (CFUs). It was observed that the Cas9^{D10A} nickase elicited a stronger phenotype than Cas9^{H840A} upon induction with xylose (Fig. S1A-B)[24]. To achieve comparable nicking efficiency at different sites in the chromosome, the *cas9^{D10A}* allele was used throughout the remainder of this study, in conjunction with distinct sgRNAs targeted to the template of either the leading or the lagging strand (in non-essential regions of the genome). Using sgRNAs to target leading and lagging strands at loci on both left and right chromosome arms, it was observed that expression of nCas9 resulted in severe growth inhibition (Figs. 1C, S1C). In contrast, expression of dCas9 targeted to the same locations did not affect the number of CFUs (Figs. 1C, S1D), indicating that DNA nicking was necessary for the observed growth defect.

Next chromosome replication was investigated using whole genome marker

frequency analysis (MFA). Expression of Cas9 variants was induced for one hour in

exponentially growing cultures. Genomic DNA (gDNA) was harvested for next-generation

sequencing, reads were mapped onto the reference genome sequence, and data is

displayed with the chromosome origin (*oriC*) at the center of the graphs (Fig. S2). It was

observed that nicks in either the leading or lagging strand resulted in a drop in sequencing

coverage from the nick site to the terminus region, whereas expression of dCas9 with the

same sgRNAs did not appear to inhibit DNA replication (Fig. 1D). Because cells expressing

nCas9 are unable to synthetise DNA downstream of the nick, this likely explains the strong

growth inhibition observed in spot-titre assays (Fig. 1C).

Previous results indicated that a replication fork encountering a single-strand

discontinuity generates a DSB[11]. To determine whether DSBs were being formed in *B.*

subtilis following nCas9 induction, localisation of the reporter mNeonGreen-RecA (mNG-

RecA) was observed in live cells using fluorescence microscopy. It has been shown that

following artificial endonuclease cleavage to create a DSB, fluorescently labelled RecA

recombinase assembles into bundles[27-29]. Prior to induction of nCas9, mNG-RecA is

observed evenly distributed over the nucleoid (Fig. S3A). Following expression of nCas9 for

one hour during exponential growth, mNG-RecA bundles were detected in 58.5% of cells

when the leading strand template was nicked and 48% of cells when the lagging strand

template was nicked (Figs. 1E-F, S3B). In contrast, expression of dCas9 did not significantly

promote mNG-RecA bundling (Figs. 1E-F, S3B). These results are consistent with the model

that collision of a replication fork with a nick in either strand of the parental DNA duplex

generates a DSB.

Single-strand discontinuities in a bacterial chromosome inactivate the replisome

MFA indicated that under conditions expressing nCas9, DNA synthesis does not proceed

beyond the nick. We wondered whether the replisome becomes arrested at a single-strand

discontinuity, or rather whether the replication fork collapses. If the latter event occurs, then

the bacterial replication restart system would be required to reload helicase and promote

replisome assembly[30,31]. In *B. subtilis* there is only one known restart system composed of

the essential proteins PriA and the helicase loaders DnaD, DnaB, and DnaI (note that in *B.*

subtilis the replicative helicase is DnaC)[32-36].

Chromatin immunoprecipitation (ChIP) followed by quantitative PCR (qPCR) was

used to determine the enrichment of helicase and replication restart proteins surrounding a

nick in either the leading or lagging strand template. Following induction of nCas9 for one

hour in exponentially growing cells, samples were harvested for ChIP. Along with sites

flanking the nick region, protein enrichment was also probed at the origin *oriC* as positive

control and normalised to a site located on the opposite chromosome arm equidistant from

oriC (+90, Fig. 2A). ChIP showed that helicase is specifically enriched upstream of nicks

located in either template strand (Figs. 2B, S4). This upstream helicase enrichment was not

detected when dCas9 was expressed, indicating that the Cas9-sgRNA nucleoprotein

complex alone does not stall replication fork progression (Figs. 2B, S4).

Interestingly, significant enrichment of helicase was observed downstream of the nick

site only when cleavage occurred in the template used for leading strand synthesis (Figs.

2B, S4). Because the bacterial replicative helicase translocates along a single strand in the

5′→3′ direction, ChIP results suggest that when the template for the lagging strand is nicked

helicase runs-off the DNA substrate, whereas when the template for the leading strand is

nicked helicase progresses downstream of the single-strand discontinuity (Fig. 2G).

Consistent with the ChIP enrichment observed *in vivo* (Fig. 2A-B) and replisome inactivation

(Fig. 1D), reconstituted unwinding assays *in vitro* using bacterial helicase on model

replication fork substrates have shown that if a flap is absent from the substrate (i.e.

mimicking a nick), helicase transitions from encircling a single DNA strand onto double-

stranded DNA (dsDNA) and slides along the duplex, no longer able to unwind[37].

The enrichment of helicase upstream of nick sites is consistent with either helicase

stalling or helicase reloading via the replication restart pathway. To distinguish between

these models, ChIP was used to probe for the presence of replication restart proteins

surrounding the nicks. Following nCas9 induction, enrichment of PriA, DnaD, DnaB and

DnaI was detected upstream (but not downstream) of the sgRNA target sequences, while

these proteins were absent from this region when dCas9 was expressed (Fig. 2C-F). These

results suggest that nicks in either the leading or lagging strand template elicit the replication

restart pathway for helicase reloading, whether helicase falls off the template (lagging strand

template nick) or translocates downstream (leading strand template nick) in an inactive state

(Fig. 2G).

A weakened nCas9 system permits DNA replication restart

While replication restart proteins were enriched upstream of nick sites (Fig. 2B-F), viability

assays and MFA both indicated that downstream DNA synthesis was inhibited (Fig. 1C-D).

We speculated that persistent nCas9 cleavage activity competes with productive DNA

replication restart. Therefore, to allow interrogation of processes occurring downstream of

DNA cleavage and replisome collapse, a weaker nCas9 system was developed. An *ssrA*

degradation tag[38] was fused to *cas9^{D10A}* to promote nCas9 proteolysis, and expression of

both *nCas9-ssrA* and *sgRNAs* were placed under the control of xylose-inducible promoters

(Fig. S5A). In this background, induction of nCas9 did not significantly inhibit cell viability or

chromosome content (Fig. S5B-C) and MFA showed that DNA synthesis continued after a

one-hour xylose induction, consistent with low tolerable levels of nickase activity (Fig. 2H).

This weakened nicking system was employed for all experiments described below.

PriA is required to restart DNA replication at single-strand DNA discontinuities

The master replication restart factor *priA* is essential for growth of *B. subtilis.* To ascertain

whether PriA-dependent replication restart is required following chromosome nicking, a

titratable PriA complementation system was developed (Fig. S6A). Here conditions were

established to express PriA-ssrA at a low level, sufficient to maintain *B. subtilis* viability while

limiting the amount of PriA activity within a cell (Fig. S6B-C). When nCas9-ssrA and sgRNA

were induced with limiting PriA-ssrA, spot-titre analyses showed that cell growth was

inhibited in the absence of the endogenous *priA* copy (Fig. 2I-J). Yet in this background,

higher expression of PriA-ssrA rescued viability (Fig. S6D). Following induction of nCas9

and sgRNA for one hour during exponential growth, MFA showed that limited levels of PriA-

ssrA could not sustain chromosome synthesis downstream of the nick site (Fig. 2K).

Therefore, these data indicate that PriA is essential to restart DNA replication following

replisome inactivation at single-strand DNA discontinuities.

The DSB resulting from a single-strand discontinuity is repaired using a core set of

homologous recombination proteins

Cytological analysis indicated that a nick in the template of either the leading or lagging

strand generated a DSB (Fig. 1E). Strong evidence indicates that in bacteria such DSBs are

repaired by homologous recombination, thereby recreating a replication fork to allow

resumption of DNA synthesis[39]. To determine the factors required to repair DSBs following

nCas9 nicking, a targeted reverse genetic analysis was employed, focused on genes

previously implicated in homologous recombination and other DNA repair processes (Table

S1). Using this approach, spot-titre analyses revealed that eight non-essential genes were

required for normal growth upon nick induction: DNA end-processing genes *addA*/*addB* and

homologous recombination genes *recF*/*recO*/*recR*/*recA*/*recG*/*recU* (Fig. 3A and S7A).

Functional complementation of each synthetic lethal deletion confirmed that these gene

products are necessary to support growth in cells expressing nCas9 (Fig. S7B). Surprisingly,

the screen revealed that several factors implicated in DNA repair are not essential under

these conditions, such as those classed for end-processing (RecQ, RecS, RecJ)[7,40],

maintaining chromosome structure ("SMC-like" RecN, SbcCD, SbcEF)[41,42], recombination

mediation (RecD2, RecX)[43-45] and Holliday junction migration (RuvA, RuvB)[40] (Fig. S8).

Focusing on the eight mutants that produced a clear growth defect, MFA was used to

analyse DNA replication following induction of nCas9 and sgRNA in exponentially growing

cells. The results showed that DNA end-processing (*addA*/*addB*) and recombination mutants

(*recF*/*recO*/*recR*/*recA*/*recG*) are defective in chromosome replication downstream of the nick

site, with the only exception being *recU* (Fig. 3B; note subtle differences in replication

profiles of parental strains containing deletions of homologous recombination genes,

consistent with previous findings[46,47]). These observations support the hypothesis that

homologous recombination is necessary to repair the DSB and restart DNA replication.

Based on the proposed activities of these proteins, the following model was

developed (Fig. 3C). AddAB helicase/nuclease processes the DSB to generate a single-

stranded substrate with a 3′-end required for recombination[48,49]; RecFOR displaces the

single-stranded binding protein (SSB) from single-stranded DNA substrates produced by

AddAB, thereby allowing RecA to bind[50,51]; RecA is the recombinase that binds to single-

stranded DNA (ssDNA) with a free 3′-end, identifies homologous DNA on the non-broken

chromosome, and performs strand invasion and exchange[52,53]; RecU resolves the Holliday

junction generated by RecA[54]. The role of RecG *in vivo* is enigmatic[55,56] and discussed

further below.

The SSB C-terminal tail is required for DNA repair following replisome inactivation

The requirement of the RecFOR system for repair of a seDSB strongly suggested that the

essential single-strand binding protein (SSB) was also involved. To interrogate this

hypothesis, recruitment of SSB was probed using ChIP following induction of nCas9 and

sgRNA in exponentially growing cells. The results showed that SSB is enriched upstream of

either leading or lagging strand nicks (Fig. 4A). SSB enrichment ~3 kb upstream from the

nick site was dependent upon AddAB but independent of the remaining factors required for

homologous recombination (Fig. 4B), indicating that AddAB is necessary to generate ssDNA

for SSB.

SSB forms a homotetramer that binds ssDNA non-specifically via its N-terminal

domain. Each SSB monomer harbours a long disordered C-terminal domain that contains a

protein interaction sequence at the terminus (the C-terminal tail or CTT) (Fig. S9A-B)[57].

While the *B. subtilis* SSB-CTT is dispensable for cell viability, it is required for efficient repair

of DNA lesions created by either UV irradiation or Mitomycin C treatment [42,58].

To determine whether the SSB-CTT is required for repair of a DSB following

replisome inactivation at a nick, an IPTG-inducible SSB complementation system was

generated in a Δ*ssbA* strain, allowing expression of either the wild-type protein or a variant

lacking the last six amino acids (SSB$^{\Delta CTT}$, Fig. S9C). Spot-titre analyses confirmed that

SSB$^{\Delta CTT}$ can complement a Δ*ssbA* mutant similar to wild-type SSB and that viability was

dependent upon IPTG (Fig. S9D), and immunoblots showed that the inducible SSB proteins

were being expressed at levels comparable to the endogenous protein (Fig. S9E).

The SSB complementation system was then moved into strains expressing nCas9

and sgRNA (Fig. S9F). Spot titre analyses showed that wild-type SSB was able to support

viability following induction of nCas9 and sgRNA, whereas the SSB$^{\Delta CTT}$ variant could not

(Fig. 4C). MFA confirmed that chromosome replication from the nick site to the terminus is

arrested when SSB lacks its C-terminal tail (Fig. 4D). Therefore, the results indicate that the

SSB-CTT is necessary for DNA repair following replisome inactivation at a nick.

The SSB-CTT is required to recruit RecO to the site of DNA repair during homologous

recombination

The SSB-CTT acts as a protein interaction hub[42] that directly interacts with several factors

found to be necessary for repair of a DSB, including RecO, RecG, and PriA (Figs. S9G, 2I-K,

3A-B). To investigate the essential role of the SSB-CTT, ChIP was employed to probe the

recruitment of SSB variants, His-RecO, and His-RecA upstream of nick sites. ChIP showed

that the SSB$^{\Delta CTT}$ variant remained enriched upstream of nicks at similar levels to the wild-

type protein (Fig. 4E). However, enrichment in His-RecO and His-RecA was lost in cells

harbouring SSB$^{\Delta CTT}$ (Fig. 4E), indicating that the SSB:RecO interaction depends on the SSB-

CTT (Fig. S9H). In cells encoding wild-type SSB, enrichment of His-RecA was dependent

upon both AddAB and RecFOR (Fig. 4F), consistent with the model that RecFOR is

necessary for RecA loading onto ssDNA substrates bound by SSB. Taken together, these

results indicate that an essential role of the SSB-CTT is to recruit RecO to repair sites

following replisome inactivation at single-strand DNA discontinuities.

What is the essential role of RecG for restart of DNA replication?

While the role of most factors required for resuming DNA replication following replisome

inactivation at a single-strand discontinuity could be rationalised, the role of RecG was

unclear. RecG has been implicated in a myriad of DNA transactions including reversal of

replication forks, migration of Holliday junctions, regulation of RecA strand exchange,

processing of flaps generated when DNA replication forks converge, and stabilisation of D-

loops[56,59-62].

Examination of the MFA in a Δ*recG* mutant showed that there was an increase in

sequence read depth upstream of a nick targeted to either the leading or lagging strand

template (Fig. 5A), suggesting that DNA synthesis increases in this region in the absence of

RecG. This observation was reminiscent of results reported in *Escherichia coli* during DSB

repair in a Δ*recG* mutant, where it was proposed that RecG remodels intermediate DNA

structures formed during homologous recombination to ensure PriA is correctly orientated to

load helicase (Fig. 5B, in the absence of RecG helicase reloading occurs on the opposite

strand normally used for DNA replication, thus directing inappropriate DNA synthesis from

the terminus towards *oriC*)[47]. This hypothesis diverges from the generally accepted model

for homologous recombination, in which resection of a DSB involves degradation of the

strand with a 5′-end, thereby generating a single-stranded 3′-end substrate for recombinases

(e.g. RecA, Fig. S10). The revised model instead suggests that following DNA end-

processing, the strand containing a 5′-end remains intact and can hybridise the displaced

strand (D-loop) formed as a consequence of RecA strand invasion[47]. It was suggested that

this intermediate DNA structure was the substrate for RecG, such that the enzyme would

remodel the strands by unwinding the annealed 5′-end until the annealed 3′-end abutted the

fork junction, thus generating the correct substrate for PriA to reload helicase (Fig. S11)[63].

Biochemical data using purified proteins has shown that RecG and PriA can perform these

activities on model replication fork substrates[63,64]. Therefore, we sought to test this

hypothesis *in vivo* using the nCas9 system.

313 RecG is required to remodel replication forks formed when a DSB contains a 5′-end

The genetic analysis above indicated that the AddAB helicase/nuclease was responsible of

end processing of the DSB (Fig. 3A). While both AddA and AddB subunits harbour nuclease

activity, the two enzymes cleave distinct strands of the DNA duplex following unwinding by

AddA helicase activity (AddA cleaves the strand terminating with a 3′-end, AddB cleaves the

strand terminating with a 5′-end)[49]. Therefore, it was hypothesised that inactivating either

AddA or AddB nuclease activity would bias DSB processing to favour the generation of

either a 3′-tail (AddA nuclease mutant) or a 5′-tail (AddB nuclease mutant).

To begin, ChIP was used to confirm that AddAB was acting directly on the DSB

generated following replisome inactivation at a nick. AddA and AddB were independently

fused to the fluorescent protein *mNeonGreen* and enrichment of each subunit was

determined. The results showed that mNG-AddA and mNG-AddB are both enriched

upstream of nicks, consistent with their direct role in end-processing under these conditions

(Fig. 5C).

Next, the enzymatic activities of AddA and AddB were reduced through substitution

of catalytic residues[49]. Spot titre analyses showed that the helicase defective AddA^{K36A}

(AddA$^{\Delta HEL}$) was unable to sustain bacterial growth upon induction of nicks, whereas

nuclease defective variants of either AddA^{D1172A} (AddA$^{\Delta NUC}$) or AddB^{D961A} (AddB$^{\Delta NUC}$) grew

similar to a strain with wild-type AddAB (Figs. 5D, S12A). Notably, the double nuclease

variant AddA$^{\Delta NUC}$AddB$^{\Delta NUC}$ also supported normal growth (Fig. 5D), suggesting that under

these conditions in live cells the DSB may not require resection. MFA confirmed that DNA

synthesis was blocked downstream of a nick in the strain expressing AddA$^{\Delta HEL}$, while the

strains expressing AddA$^{\Delta NUC}$, AddB$^{\Delta NUC}$ or AddA$^{\Delta NUC}$AddB$^{\Delta NUC}$ were comparable to wild-type

(Figs. 5E, S12B-C). Moreover, ChIP demonstrated that SSB is not enriched upstream of

nicks in a strain expressing AddA$^{\Delta HEL}$, whereas SSB was enriched in AddA$^{\Delta NUC}$ and

AddB$^{\Delta NUC}$ backgrounds (Fig. 5F). Together these results indicate that AddAB helicase

activity is directly required to process a DSB generated after replisome inactivation at a nick,

and that nuclease activity can be abolished without compromising DNA replication restart.

Finally, the Δ*recG* mutant was moved into strains expressing either AddA$^{\Delta NUC}$ or

AddB$^{\Delta NUC}$ and nicking was induced (Fig. 6A-B). Spot-titre assays showed that RecG

remained essential in the strains containing AddB$^{\Delta NUC}$, where degradation of the strand

terminating with a 5′-end is inhibited (Fig. 6C). Strikingly, in the AddA$^{\Delta NUC}$ strain where

degradation of the strand terminating with a 3′-end is inhibited, the absence of RecG was

well tolerated (Fig. 6C-D). These results are consistent with the model that RecG is required

to remodel recombination intermediates that contain a single-strand with a 5′-end annealed

to a D-loop. Importantly, the critical role of RecG in strains with single-strand discontinuities

(Fig. 3A) implies that wild-type processing of a DSB often generates a single-strand with a

5′-end, which is inconsistent with most current models of bacterial homologous

recombination.

DISCUSSION

In this study we have characterised the fate of the bacterial replisome *in vivo* after it encounters a single-strand discontinuity in the DNA template (Fig. 1A). The results indicate that a nick in the template for either the leading or lagging strand arrests DNA synthesis, generates a DSB, and elicits replication restart following repair of the DNA break through homologous recombination (Figs. 1-3). These findings are consistent with the observation that a nick in a replicating bacteriophage chromosome generates DSBs[11] and with recent reports of the impact of nicks in eukaryotic replication systems[20,21,23]. Importantly, while replisomes are thought to be generally intolerant to many types of DNA damage[65], it appears that the bacterial replication machinery cannot readily bypass single-strand interruptions as these lesions affect helicase progression (Figs. 2, 7(I)).

Repair of the DSB to facilitate replication restart requires homologous recombination (Fig. 3). Genetic analysis combined with ChIP indicates that a single pathway is utilised under these experimental conditions (Fig. 7): AddAB unwinds the DNA; SSB binds the ssDNA; RecFOR counteracts SSB and allows RecA to bind ssDNA; RecA promotes strand invasion of the homologous chromosome and generates a D-loop, which often anneals to a single-strand with a 5′-end; RecG regresses the fork and simultaneously separates the annealed strand with a 5′-end; PriA binds to the remodelled fork where it interacts with the strand harbouring a 3′-end and reloads helicase to promote replisome assembly and DNA replication restart. The observed lack of redundancy between proteins involved in recombinational DNA repair suggests that the pathway identified here is the major route for fixing this type of collapsed replication fork in *B. subtilis*[41,66].

Building upon these results, we investigated specific protein activities to help elucidate their roles during recombinational repair of a DSB. Below we discuss these findings in what we propose is chronological order.

The replicative helicase is inactivated after it encounters a nick

In bacteria the replicative helicase encircles and tracks along the template for lagging strand

synthesis. ChIP data suggests that there are distinct fates of the helicase depending on the

location of a nick (Figs. 2A-B, S4). As expected for a nick in the lagging strand template,

helicase enrichment was only observed at the cleavage site and upstream, consistent with

the model that helicase runs-off the template when it encounters the single-strand

discontinuity (Figs. 2G, 7(I)). In contrast, when the nick was in the leading strand template,

helicase was also enriched downstream of the cleavage site, suggesting that the enzyme

continued translocation. We propose that upon encountering a nick in the leading strand

template, helicase slides onto the dsDNA template in an inactive conformation that cannot

perform further DNA unwinding (Figs. 2G, 7(I)). This model is consistent with *in vitro*

experiments using *Thermus aquaticus* helicase, where after unwinding up to a nick on the

leading strand template, the enzyme can continue to translocate with the two strands of DNA

passing through the central channel with no resultant unwinding[37]. Whether the bacterial

helicase is actively removed from dsDNA under these conditions, akin to what has been

observed for the eukayotic CMG helicase in *Xenopus* extracts[20], requires further

investigation.

AddAB helicase activity is necessary and sufficient for recombinational repair

AddAB and related enzymes have been shown to bind blunt end dsDNA or dsDNA with

short overhangs (4-20 nucleotides with either a 5′- or a 3′-end)[49,67]. When the template for

leading strand synthesis is nicked, the DNA polymerase likely runs-off the end of the

substrate to produce a blunt or nearly blunt end, suitable for recognition by AddAB.

However, when the template for lagging strand synthesis is nicked, a substrate with a 3′-tail

would likely be produced. Because the reverse genetic screen suggested that no additional

nucleases are required for this recombinational repair pathway (Figs. 3, S8), we speculate

that the high local concentration of replication factors would facilitate priming near the end of

the 3′-tail, thereby filling the ssDNA with an Okazaki fragment to create a suitable substrate

for AddAB.

The master recombination factor RecA requires ssDNA to promote homologous

pairing. A long-standing question is how this ssDNA is generated *in vivo*. Most current

models suggest that enzymes resect a DSB to generate a single-strand terminating with a

3′-end[7,31]. Here we tested this model by mutating the nuclease activities of AddA and AddB.

The results showed that neither nuclease activity is required for enzyme function under the

experimental conditions tested (Figs. 5D-E, S12B-C). In contrast, AddAB helicase activity

was required for recombinational repair, consistent with a previous study of the ancestral

AdnAB resection machinery in *Mycobacteria tuberculosis*[68]. These results suggest that the

key role of AddAB is not to digest the dsDNA ends, but rather to separate the strands and

create the ssDNA substrate for SSB (Figs. 5F, 7(II-III)).

The local concentration of SSB is high at the site of DNA replication[69,70], potentially

facilitating its rapid association with the ssDNA emerging from AddAB. In fact, *in vitro* the

AddA^{NUC}AddB^{NUC} nuclease-dead enzyme does not unwind dsDNA in the absence of SSB[71],

suggesting that AddAB unwinding and SSB binding ssDNA may be coupled. We propose

that AddAB nuclease activity represents a secondary method of generating a ssDNA

substrate for SSB, following initial AddAB helicase activity required to separate the DNA

duplex.

Interaction of the SSB C-terminal tail with RecO is necessary for replication repair

The SSB-CTT is an established protein interaction hub that has been shown to directly bind

several of the essential replication and repair factors required at nicks, including RecO[72-75],

RecG[76,77], and PriA[58,78]. However, the critical activities performed by the SSB-CTT *in vivo*

have not yet been established[42]. Based on the results above, we conclude that the SSB-CTT

interaction with RecO is necessary for replication repair. RecO is part of the RecFOR system

required for RecA loading onto ssDNA *in vitro*[79] and RecA-GFP foci formation *in vivo*[80].

Because RecO activity occurs before the activities of RecG and PriA (i.e. RecA binding

ssDNA is blocked, Fig. 4F), this precludes further conclusions regarding the importance of

these protein:protein interactions at this time.

In vitro the RecO:RecR complex facilitates loading of RecA onto SSB-coated

ssDNA[81,82]. Because RecO, RecR, and RecF were all required for replication repair, we

propose that the essential role of RecF here is to recruit RecR and generate the RecO:RecR

complex. This model is consistent with a hand-off mechanism for RecR observed *in vitro*[81].

RecF is a dimeric ATPase that adopts a clamp conformation around dsDNA[51]. Recent

studies in *E. coli* have shown that recruitment of RecF to sites of repair is mediated by the

DNA polymerase sliding clamp processivity factor DnaN[82]. As was the case for SSB[70], the

local concentration of DnaN is high near the sites of DNA replication collapse[69], thereby

potentially facilitating RecF recruitment. Importantly for this model, since both SSB and

DnaN are enriched at a replication fork, we predict that there is a mechanism to regulate

RecO:RecR complex formation until it is required, thus safeguarding the SSB-coated strand

from promiscuous RecA loading (Figs. 4B, 4F, 7(III-V)).

RecG remodels a repaired replication fork by unwinding dsDNA with an exposed 5′-

end

The biological role of RecG has been a long-standing question[55,56]. *In vitro* RecG is a DNA

translocase that unwinds a myriad of DNA substrates, including Holliday junctions, D-loops,

and model replication forks[47,60-62,64]. Here MFA indicates that a *ΔrecG* mutant cannot

efficiently restart DNA replication at a nick directed to either the leading or the lagging strand

template (Figs. 3B, 5A-B), consistent with a role in fork remodelling[46]. Based on genetic

experiments and supported by biochemical assays *in vitro*, a model has been presented for

RecG binding to a replication fork and unwinding dsDNA on the lagging strand template to

allow correct orientation of PriA for helicase loading[47,63]. Critically, this model requires that

the single strand with a 5′-end from the DSB can hybridise with ssDNA of the D-loop on the

homologous chromosome.

How this predicted intermediate structure is constructed remains to be determined,

but the implication is that the separated strands of the DSB are either the same length or the

strand with a 5′-end is longer (Figs. 6A, 6D, 7(VI-VII)). This model contrasts with most

current views of bacterial DSB end-processing where a single strand with a 3′-end is

produced for RecA (Fig. S10)[7,31,83]. Moreover, because RecG is essential in strains

expressing Cas9 nickases, it suggests that during wild-type DSB end-processing the

separated strands often contain a 5′-end that requires remodelling by RecG (Fig. 7(VII)).

These unanticipated results, together with those of the accompanying manuscript analysing

resolution of collapsed eukaryotic replication forks[84], necessitate further evaluation of DSB

management following replisome collapse across all domains of life.

Holliday junction migration and resolution are not necessary for replication restart

Branch migration of a Holliday junction is generally considered a hallmark of homologous

recombination, therefore it was notable that the canonical Holliday junction translocase

RuvAB was not required for replication repair at nicks (Fig. S8D). In fact, the *ΔrecU ΔruvAB*

knockout attenuated the penetrance of the *ΔrecU* mutant upon introduction of single-strand

discontinuities, suggesting that either another enzyme can functionally complement RuvAB

activities or that branch migration is not required for this repair pathway.

While genetic analysis shows that RecU is required for wild-type growth following

expression of Cas9 nickases, MFA indicates that RecU is not necessary for replication

restart *per se* (Fig. 3). These results are consistent with the model that RecU resolves the

Holliday junction formed during recombinational repair[85]. We hypothesise that the growth

defect observed in the *ΔrecU* mutant may be caused by aberrant chromosome

segregation[85], as the recombined DNA molecules need to be separated prior to completing

cell division. It is known that multiple systems can facilitate chromosome separation

following replication (e.g. Topoisomerase IV, XerD)[86,87], potentially explaining why the

growth defect of the *ΔrecU* mutant under these experimental conditions was less severe

compared to the other recombination mutants identified.

Potentiating ROS-mediated antibiotic toxicity by perturbing DNA repair

In this work, the nicks generated by Cas9 enzymes were used as proxies for single-strand

discontinuities formed during removal of damaged DNA from a bacterial genome. We find

that when a bacterial replication fork encounters a single strand discontinuity it becomes

inactivated, likely contributing to the toxic effect of ROS-dependent damage induced by

antimicrobials. By identifying the essential factors and activities required to repair the DSB

formed after a replisome encounters a nick, we have provided a set of molecular targets

whose disruption could potentiate the activity of antibiotics that produce ROS-dependent

DNA damage.

RESOURCE AVAILABILITY

Lead contact

For additional information and requests for resources and reagents, please contact the lead contact: Charles Winterhalter (charles.winterhalter@newcastle.ac.uk).

Materials availability

Genetic materials generated in this study can be obtained directly from the lead contact.

Data and code availability

R code (DOI: 10.5281/zenodo.14793122) and protein structures for SSB:ssDNA and SSB:RecO complexes (DOI: 10.5281/zenodo.14793919) are available from Zenodo.

ACKNOWLEDGEMENTS

This work was supported by a Wellcome Trust Early-Career Award [226338/Z/22/Z] to CW and Wellcome Trust Discovery Award [225811/Z/22/Z] to HM. We would also like to express our gratitude to Frances Davison for technical assistance, James Grimshaw for microscope maintenance and Frederic Schramm for insightful discussions prior to submission.

AUTHOR CONTRIBUTIONS

CW and HM contributed to the conception/design of the work. CW and SF generated results presented in the manuscript. CW created figures. CW and HM wrote/edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES

During the preparation of this work, the authors used AlphaFold 3 to generate SSB:ssDNA and SSB:RecO protein models. After using this tool, the authors reviewed the models and edited their content as needed and take full responsibility for the content of the publication.

530 **STAR Methods**

531 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Cas9 monoclonal	Merck	SAB4200701
Anti-FtsZ polyclonal	Eurogentec	Lucet *et al.* (2000)[88]
Anti-PriA polyclonal	Eurogentec	This study
Anti-SSB polyclonal	Eurogentec	This study
Anti-mNeonGreen monoclonal	ProteinTech	32F6
HRP-conjugated Goat Anti-Rabbit IgG (H+L)	ProteinTech	SA00001-2
HRP-conjugated Donkey Anti-Sheep IgG (whole molecule)	Merck	A3415
HRP-conjugated Goat Anti-Mouse IgG, Fcγ Fragment Specific	ProteinTech	SA00001-1-A
Anti-DnaD polyclonal	Eurogentec	Winterhalter *et al.* (2022)[33]
Anti-DnaB polyclonal	Eurogentec	This study
Anti-DnaI polyclonal	Eurogentec	This study
Anti-DnaC polyclonal	Eurogentec	This study
Anti-His monoclonal	Qiagen	34660
Bacterial strains		
See Table S2		
Chemicals		
Oxoid™ Nutrient agar	Thermo Scientific	CM0003B
Oxoid™ Tryptone	Thermo Scientific	LP0042B
Sodium chloride (NaCl)	VWR	27810.364P
Oxoid™ Yeast Extract Powder	Thermo Scientific	LP0021B
Ampicillin sodium salt	Merck	A9518
Erythromycin	Merck	E5389
Lincomycin hydrochloride	Merck	L6004
Chloramphenicol	Merck	C0378
Kanamycin disulfate salt	Merck	K1876
Spectinomycin dihydrochloride pentahydrate	Melford	S23000
Zeocin™ Selection Reagent	Invitrogen	R25001
D-(+)-Xylose	Merck	W360600
Isopropyl-β-D-thio-galactopyranoside (IPTG)	Melford	I56000
Ribonuclease A	Merck	R6513

Lysozyme	Merck	L6876
Ammonium sulphate	VWR	21333.296
di-Potassium hydrogen phosphate	VWR	26930.293
Potassium dihydrogen phosphate	VWR	26936.293P
tri-Sodium citrate dihydrate	VWR	27833.294P
Magnesium sulphate heptahydrate ($MgSO_4$)	VWR	25165.260
Ammonium iron(III) citrate	Merck	F5879
D-(+)-Glucose	VWR	101176K
L-Tryptophan	Formedium	DOC0189
Casein Peptone	Merck	22090
Ethanol	Fisher Scientific	12478740
Oxoid™ Phosphate Buffered Saline Tablets	Thermo Scientific	BR0014G
cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail	Roche	4693159001
DL-Dithiothreitol (DTT)	Melford	D11000
Methanol	Fisher Scientific	M/4000/17
TRIS base	Melford	T60040
Glycine	Merck	G7126
Tricine	Thermo Scientific	J64437.36
Sodium dodecyl sulfate (SDS)	Merck	75746
Ethylenediaminetetraacetic acid (EDTA)	Merck	03609
TWEEN® 20	Merck	P7949
Dried skimmed milk	Tesco	267377676
Calcium chloride dihydrate ($CaCl_2$)	Merck	C3306
Manganese(II) Sulfate Tetrahydrate ($MnSO_4$)	Fisher Scientific	M/2300/50
L-Glutamic acid potassium salt monohydrate	Merck	G1501
Sodium azide	Merck	S8032
Water, DEPC-Treated, DNASE/RNASE free, Fisher BioReagents™	Fisher Scientific	BP561-1
Formaldehyde solution	Merck	252549
Sucrose	Merck	S0389
Triton™ X-100	Merck	T8787
Protease from *Streptomyces griseus* (pronase)	Merck	P5147
Lithium chloride (LiCl)	Merck	L9650
IGEPAL® CA-630	Merck	I8896
Sodium deoxycholate monohydrate	Merck	D5670
Recombinant Proteinase K Solution	Invitrogen	AM2548
Commercial kits and equipment		

Q5® High-Fidelity DNA Polymerase	NEB	M0491L
KOD Xtreme™ Hot Start DNA Polymerase	Merck	71975
QIAprep Spin Miniprep Kit	Qiagen	27106
NEBuilder® HiFi DNA Assembly Cloning Kit	NEB	E5520S
QIAquick PCR Purification Kit	Qiagen	28106
DNeasy Blood and Tissue Kit	Qiagen	69504
Ultrasonic processors, Vibra-Cell™	Jencons-PLS	VC130
NuPAGE™ Bis-Tris Midi Protein Gels, 4 to 12%, 1.0 mm	Invitrogen	WG1402BX10
NuPAGE™ MES SDS Running Buffer (20X)	Invitrogen	NP0002
NuPAGE™ LDS Sample Buffer (4X)	Invitrogen	NP0007
Amersham™ Hybond® P membrane, PVDF	Merck	GE10600023
Wypall X60 cloths	Camlab	1206641
Trans-Blot Turbo Transfer System	Bio-rad	1704150
Clarity Western ECL Substrate	Bio-rad	1705061
ImageQuant™ LAS 4000 mini	GE Healthcare	28-9607-61 AB
DNA Prep IPB and buffers	Illumina	20049005
DNA Prep PCR and buffers	Illumina	20015826
DNA Prep CD indexes	Illumina	20015882
Qubit™ 1X dsDNA High Sensitivity (HS) kit	Invitrogen	Q33231
Qubit™ 4 Fluorometer	Invitrogen	Q33238
iSeq 100 i1 Reagent v2 sequencing chips	Illumina	20009555
iSeq 100 i1 Reagent v2 cartridges	Illumina	20009584
iSeq 100 Sequencing System	Illumina	20021532
GFP filter (EX470/40 DM495lpxr, EM525/50)	Chroma	49002
DAPI filter (EX350/50, DM400lp, EM460/50)	Chroma	49000
mCherry filter (EX560/40, DM585lprx, EM630/75)	Chroma	49008
Dynabeads™ Protein G for Immunoprecipitation	Invitrogen	10009D
Luna® Universal qPCR Master Mix	NEB	M3003E
QIAgility robotic workstation	Qiagen	9001908
Rotor-Disc 100	Qiagen	981311
Rotor-Disc Heat Sealer	Qiagen	9019725
Rotor-Disc Heat Sealing Film	Qiagen	981601
Rotor-Gene Q	Qiagen	9001560
Deposited data		
R script for WGS data processing	This study	10.5281/zenodo.14793122
AlphaFold 3 protein models	This study	10.5281/zenodo.14793919

Oligonucleotides		
See Table S4		
Recombinant DNA		
See Table S3		
Software and algorithms		
CLC Genomics Workbench	Qiagen	
CorelDRAW® Graphics Suite	Alludo	
Fiji	ImageJ	Schindelin *et al.* (2012)[89]
Microsoft Office Suite	Microsoft	
NIS-Elements	Nikon	
Primer3Plus	Online tool	Untergasset *et al.* (2007)[90]
Pymol Molecular Graphics	Schrödinger, Inc.	Delano (2002)[91]
Rotor-Gene Q software v2.0.2	Qiagen	
Rsamtools	Bioconductor	
RStudio Desktop	Posit	

Experimental model and details

The following bacterial organisms were used: *E. coli* and *B. subtilis.* Unless otherwise stated

in method details, strains were grown at 37°C using nutrient agar (NA) or lysogeny broth

(LB: 10 g/L tryptone, 10 g/L NaCl, 5 g/L yeast extract) for routine selection and maintenance

of bacterial strains. Supplements were added as required: ampicillin (100 μg/ml),

erythromycin (1 μg/ml) in conjunction with lincomycin (25 μg/ml), chloramphenicol (5 μg/ml),

kanamycin (5 μg/ml), spectinomycin (50 μg/ml), zeocin (10 μg/ml), xylose (1% w/v) and

IPTG (0.05 mM, 0.1 mM or 1 mM).

Method details

Bacterial strains

Bacterial strains used in this study are listed in Table S2.

E. coli plasmid construction

E. coli transformations for constructs harbouring *priA* were performed in CW198 via heat-
shock following the Hanahan method[92] and propagated in LB with appropriate antibiotics at
37°C. DH5α [*F- Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17(rk-, mk+) phoA*
supE44 thi-1 gyrA96 relA1 λ-] was used for other plasmids construction. Plasmids were
purified using QIAprep spin miniprep kits and recombinant DNA confirmed by sequencing
(Table S3). Descriptions, where necessary, are provided below.

pCW367, pCW368, pCW478, pCW479, pCW510, pCW717, pCW734, pCW736 and
pCW850 were generated by Quickchange mutagenesis using mutagenic primer pairs
containing an overlap of 10-15 bp. PCR products were treated with DpnI for at least 3 hours
to digest template DNA and 10 μl were used for *E. coli* transformation.

pCW300, pCW301, pCW327, pCW460, pCW471, pCW472, pCW575, pCW734,
pCW736 and pCW766 were generated by Gibson assembly using the NEB Hi-Fi cloning kit
according to manufacturer instructions following purification of individual PCR fragments
using QIAquick PCR purification kits.

Linear recombinant DNA assembly

For recombinant DNA assembly without *E. coli* propagation, pCW573, pCW574, pCW577,
pCW578, pCW605, pCW640, pCW672, pCW702, pCW720, pCW732, pCW758, pCW759,
pCW760, pCW777, pCW779, pCW781, pCW783, pCW785, pCW802, pCW803, pCW804,
pCW809, pCW834, pCW848, pCW854, pCW855, pCW857 and pCW859 were assembled
linearly using a multistep *in vitro* assembly process followed by nested PCR amplification
prior to *B. subtilis* transformation. Briefly, individual PCR fragments corresponding to
homology regions and insert DNA were amplified, purified and DNA ends ligated using the
NEB Hi-Fi cloning kit. Raw Gibson assembly products were then amplified using nested
primer pairs annealing towards the end of homology regions and PCR products were used
for *B. subtilis* transformation. Following isolation of *B. subtilis* colonies resistant to the
appropriate antibiotic markers, genomic DNA was extracted using the DNeasy blood and
tissue kit, regions corresponding to recombinant DNA were amplified via PCR using

oligonucleotides annealing outside homology fragments to ensure double crossover had

occurred and results were confirmed via PCR product sequencing. Note that CW2045 and

derivatives harbour the additional *addA*[D682E] mutation necessary to propagate *addB*[D961A]-

addA[D1172A] in live cells, which is surface exposed and structurally unlikely to impact AddAB

end-processing activities (Fig. S12E).

B. subtilis strain construction

Transformation of competent *B. subtilis* cells was performed using an optimized two-step

starvation procedure as previously described[93]. Briefly, recipient strains were grown

overnight at 37°C in transformation medium (Spizizen salts: 0.2% w/v ammonium sulphate,

1.4% w/v dipotassium phosphate, 0.6% w/v potassium phosphate, 0.1% w/v tri-sodium

citrate, 0.02% w/v magnesium sulphate supplemented with 1 µg/ml ammonium iron(III)

citrate, 6 mM $MgSO_4$, 0.5% w/v glucose, 0.02 mg/ml tryptophan and 0.02% w/v casein

peptone) supplemented with IPTG where required. Overnight cultures were diluted 1:20 into

fresh transformation medium supplemented with IPTG where required and grown at 37°C for

2.5-3 hours with continual shaking. An equal volume of prewarmed starvation medium

(Spizizen salts supplemented with 6 mM $MgSO_4$ and 0.5% w/v glucose) was added and the

culture was incubated at 37°C for two hours with continual shaking. DNA was added to 350

µl cells and the mixture was incubated at 37°C for one hour with continual shaking. 20-200

µl of each transformation was plated onto selective media supplemented with IPTG where

required and incubated at 37°C for 24-48 hours. Results were validated by propagating cells

on relevant antibiotic markers and sequencing.

Spot titre assays

Cells were grown in LB overnight in the presence of 0.5% w/v glucose. Overnight cultures

were diluted in fresh LB medium using 10-fold serial dilutions and 5 µL aliquots were spotted

onto NA plates with or without xylose. NA plates were supplemented with IPTG when

required as indicated on individual figure panels. All plates were incubated for 24 hours or up

to 48 hours when indicated. Experiments were performed independently at least three times

and representative data are shown.

Immunoblot analyses

Cultures were grown overnight in LB and diluted 1:100 in fresh medium the next morning.

Diluted cells were grown to an absorbance at 600 nm (A_{600}) of 0.2-0.3, induced with 1% w/v

xylose or 1 mM IPTG where required and further incubated for one hour. Cells were

harvested using centrifugation, resuspended in 1x PBS supplemented with one tenth of a

peptidase inhibitor tablet and sonicated (40 amp) twice for 12 seconds with 3 second pulses

at 4°C. Samples were adjusted to 1X LDS buffer and 125 mM DTT, fixed for 5 minutes at

95°C and centrifuged for 2 minutes prior to loading the supernatant on polyacrylamide gels.

Proteins were separated by electrophoresis using NuPAGE 4-12% Bis-Tris gradient gels run

in MES SDS buffer and transferred to methanol-activated Hybond-P PVDF membranes

using Wypall X60 cloths for blotting in a semi-dry apparatus (Bio-rad Trans-Blot Turbo,

transfer buffer: 300 mM Tris-HCl, 300 mM glycine, 140 mM tricine, 0.05% w/v SDS, 2.5 mM

EDTA). Membranes were washed twice using PBST (1X PBS supplemented with 0.1% v/v

Tween-20) prior to blocking for 90 minutes at room temperature (blocking buffer: PBST

supplemented with 7.5% w/v milk). Individual polypeptides were probed using the following

primary antibodies diluted in blocking buffer following overnight incubation at 4°C: anti-Cas9

(1:1000), anti-FtsZ (1:5000), anti-PriA (1:1000), anti-SSB (1:1000) and anti-mNeonGreen

(1:1000). Detection was performed with anti-rabbit (PriA, SSB), anti-sheep (FtsZ) or anti-

mouse (Cas9, mNeonGreen) horseradish peroxidase-linked secondary antibodies in

combination with the Clarity ECL substrate using an ImageQuant LAS 4000 mini digital

imaging system. Detection of Cas9, FtsZ, PriA, SSB and mNeonGreen was within a linear

range. Experiments were independently performed at least twice and representative data

are shown.

Whole genome sequencing

Cells were grown in rich chemically defined medium (RCDM: Spizizen salts supplemented

with 1 μg/ml ammonium iron(III) citrate, 0.1 mM $CaCl_2$, 0.13 mM $MnSO_4$, 6 mM $MgSO_4$,

0.5% w/v glucose, 0.1% w/v glutamate, 0.02 mg/ml tryptophan and 0.02% w/v casein

peptone) overnight and diluted 1:100 in fresh RCDM the next morning. Diluted cells were

grown to an absorbance at 600 nm (A_{600}) of 0.2-0.3, induced with 1% w/v xylose and further

incubated for one hour. Growth was arrested by adding 0.05% w/v sodium azide to cultures

followed by vigorous mixing, cells were collected by centrifugation, the supernatant

discarded, and pellets were flash frozen in liquid nitrogen before genomic DNA extraction

using the DNeasy blood and tissue kit and a final elution volume of 50 μl in ultrapure water.

Individual gDNA libraries were prepared using the DNA Prep tagmentation kit and pre-paired

i5/i7 indices prior to pooling and DNA quantification with the 1X dsDNA high sensitivity kit

using a Qubit instrument. Pooled libraries were diluted to 100 pM in resuspension buffer and

20 μl were loaded onto iSeq 100 i1 Reagent v2 sequencing chips for whole-genome

sequencing using an iSeq100 sequencer.

Following whole-genome sequencing and demultiplexing embedded within the

iSeq100 indexed sequencing workflow, .FASTQ files were extracted and loaded onto the

CLC Genomics Workbench for initial data processing. Paired .FASTQ files were mapped

onto the *B. subtilis* Δ6 reference genome using the 'Map Reads to Reference' function and a

.BAM file was created from mapped sequencing reads for each sequenced genome. An in-

house script written in R using the Rsamtools library was employed to deconvolute .BAM

files to validate even read distribution for sense/antisense strands prior to calculating density

coverage and extracting cumulated densities per chromosome position.

Fluorescence microscopy

Strains were grown overnight at 37°C in RCDM and the following day, cultures were diluted

1:100 into fresh imaging medium (RCDM where glucose is replaced by 0.5% v/v glycerol) in

the presence of 0.1 mM IPTG for induction of mNeonGreen-RecA expression where

required. Cells were allowed to grow to an A_{600} of 0.3, induced with 1% w/v xylose and

further incubated for 90 minutes. DAPI (5 µg/ml) and Nile red (1 µg/ml) stains were used to

visualise the nucleoid and cell membrane following a 10 minute incubation on a benchtop

incubator prior to imaging.

Cells were mounted on ~1.25% agar pads (in sterile ultrapure water) and a 0.13- to

0.17-mm glass coverslip was placed on top. Microscopy was performed on an inverted

epifluorescence microscope (Nikon Ti) fitted with a Plan Apochromat Objective 100x/1.40

NA Oil Ph3. Light was transmitted from a CoolLED pE-300 white light source through a

Sutter Instruments liquid light guide and images were collected using a Photometrics Prime

camera. Chroma fluorescence filter sets were used with 100 ms GFP, 250 ms DAPI and 50

ms mCherry exposure times at 100% LED power. Digital images were acquired using NIS

Elements and analysed via the Fiji software[89]. Quantification of mNeonGreen-RecA features

was manually curated from the count of 100 individual cells. All experiments were

independently performed at least twice and representative data are shown.

Chromatin immunoprecipitation

Chromatin immunoprecipitation (ChIP) was performed as previously described with minor

modifications[32]. Strains were grown overnight at 37°C in RCDM and the following day,

cultures were diluted 1:100 into fresh medium and allowed to grow to an A_{600} of 0.2-0.3.

Cells were induced with 1% w/v xylose and further incubated for one hour prior to ChIP.

Samples were adjusted to 1X PBS and crosslinked with 1% v/v formaldehyde for 10 minutes

on a roller at room temperature, then quenched with 0.1 M glycine. Cells were pelleted at

4°C, washed three times with ice-cold 1X PBS (pH 7.3), frozen in liquid nitrogen and stored

at -80°C. Frozen cell pellets were resuspended in 500 µl of lysis buffer (50 mM NaCl, 10 mM

Tris-HCl pH 8.0, 20% w/v sucrose, 10 mM EDTA, 100 µg/ml RNase A, one quarter of a

peptidase inhibitor tablet, 4 mg/ml lysozyme) and incubated at 37°C for 30 min to degrade

the cell wall. Protoplasts were supplemented with 500 µl of immunoprecipitation buffer (300

mM NaCl, 100 mM Tris-HCl pH 7.0, 2% v/v Triton X-100, one quarter of a peptidase inhibitor

tablet) to lyse cells and the mixture was incubated at 37°C for a further 10 minutes before

cooling on ice for 5 minutes. Lysis and immunoprecipitation buffer volumes were multiplied

by the number of antibodies to probe per sample. DNA samples were sonicated (40 amp)

three times for 2 minutes at 4°C to obtain an average fragment size of 500 to 1000 base

pairs. Cell debris were removed by centrifugation at 4°C and the supernatant transferred to a

fresh Eppendorf tube. To determine the relative amount of DNA immunoprecipitated

compared to the total amount of DNA, 100 μl of supernatant was removed, treated with 0.5

mg/ml pronase at 37°C for four hours then stored on ice. To immunoprecipate protein-DNA

complexes, 800 μl of the remaining supernatant was incubated with individual antibodies (2

μl anti-SSB, 2 μl anti-PriA, 2 μl anti-DnaD, 2 μl anti-DnaB, 2 μl anti-DnaI, 2 μl anti-DnaC, 2 μl

anti-mNeonGreen or 4 μl anti-His) for 90 minutes at room temperature. 750 μg of protein G

Dynabeads were equilibrated by washing with bead buffer (1X PBS, 0.01% v/v Tween 20),

resuspended in 25 μl of bead buffer and incubated with the sample supernatant for 1 hour at

room temperature. Immunoprecipitated complexes were collected by applying the mixture to

a magnet and washed with the following buffers in the respective order: 0.5X

immunoprecipitation buffer for 15 min, 0.5X immunoprecipitation buffer supplemented with

500 mM NaCl for 15 min, stringent wash buffer (250 mM LiCl, 10 mM Tris-HCl pH 8.0, 0.5%

v/v Igepal, 0.5% w/v sodium deoxycholate, 10 mM EDTA) for 20 min. Finally, protein-DNA

complexes were washed a further three times with TET buffer (10 mM Tris-HCl pH 8.0, 1

mM EDTA, 0.01% v/v Tween 20) and resuspended in 100 μl of TEN buffer (10 mM Tris-HCl

pH 8.0, 1 mM EDTA, 300 mM NaCl). Formaldehyde crosslinks of both the

immunoprecipitate and total DNA were reversed by incubation at 65°C for 18 hours in the

presence of 1 mg/ml proteinase K. DNA was then removed from the magnetic beads,

cleaned using QIAquick PCR purification columns and used for quantitative PCR analyses.

Quantitative PCR

Quantitative PCR (qPCR) was performed using the Luna qPCR master mix to measure

relative amounts of DNA bound by SSB, PriA, DnaD, DnaB, DnaI, DnaC, mNeonGreen-

AddA, mNeonGreen-AddB, His-RecO and His-RecA at specific genomic locations (e.g. near

the nick site located at -90° on the chromosome or the origin *oriC* compared to a control site

located at +90° on the chromosome). All PCR reactions were assembled using the QIAgility

robotic workstation in 20 µl reaction volumes in a Rotor-Disc 100 and qPCRs were run on a

Rotor-Gene Q instruments. Standard curves were obtained using the Rotor-Gene Q

software to calculate the efficiency of each primer pair, which varied ~5% between sets.

Oligonucleotides were designed to amplify specific genomic regions using the Primer3Plus

tool[90], were typically 20-25 bases in length (Table S4) and amplified a ~100 bp PCR product.

Individual fold enrichment ratios were obtained as follows: first, every Ct value was

converted to $1/2^{Ct}$ and technical triplicates were averaged to generate a single enrichment

value; second, genomic location specific enrichment was normalised by corresponding

values obtained at +90° on the chromosome. Error bars indicate the standard error of the

mean for three biological replicates.

Protein structure prediction and representation

Protein models for SSB:ssDNA and SSB:RecO were generated using AlphaFold 3 and

polypeptides corresponding to individual protein entities as input[94]. Main and alternative

models were manually examined and Model 0 was chosen to highlight key features using

the Pymol Molecular Graphics 2.1 software[91]. Structural highlights of the AddAB complex

were derived from the crystal structure PDB: 4CEH[95].

 FIGURE TITLES AND LEGENDS

Figure 1. Single-strand discontinuities arrest DNA synthesis and generate DSBs.

(A) Diagram illustrating the impact of leading and lagging strand discontinuities on a

replication fork, with the bacterial replicative helicase translocating along the template for the

lagging strand.

(B) Genetic system employed to create a site-specific single-strand discontinuity in the *B.*

subtilis chromosome. Black boxes correspond to antibiotic resistance markers, pink to Cas9

variants and green to sgRNA. Chromosome annotations refer to the replication origin (0),

terminus (180) and sgRNA target sequence (yellow box).

(C) Spot titre analysis of strains engineered to express (+xylose) either nCas9 or dCas9.

Individual sgRNAs were designed to target both strands at loci located at either -90° (top

panel) or +90° (bottom panel).

(D) MFA using whole genome sequencing of strains engineered to express either nCas9 or

dCas9. Individual sgRNAs were designed to target both strands at loci located at either -90°

(left panel) or +90° (right panel). The frequency of sequencing reads was plotted against

genome position.

(E) Representative fluorescence microscopy images of the DSB reporter mNG-RecA in

strains engineered to express (+xylose) either nCas9 or dCas9 (sgRNAs target -90°).

Fluorescent RecA organises into bundles in the presence of DSBs. Green signal

corresponds to mNG-RecA fluorescence, grey scale images show corresponding phase

contrast images.

(F) Quantification of fluorescence microscopy images of the DSB reporter mNG-RecA in

strains engineered to express (+xylose) either nCas9 or dCas9 (sgRNA target -90°).

Detection of spots and bundles corresponds to distinct foci or filament-like structures

annotated in panel (E). Error bars indicate the standard error of the mean.

Figure 2. Replication restart is necessary to resume DNA synthesis following

replisome inactivation at a single-strand discontinuity.

(A) Diagram illustrating genomic locations probed via ChIP-qPCR at loci located at either -

90° or +90°. The sgRNA targeted Cas9 protein variants to the locus at -90°. The enrichment

signal at the +90° locus was used for fold change normalisation. Lagging strand and leading

strand templates are represented in black and grey lines, respectively (numbers refer to

polarity). Blue arrows indicate leading strand synthesis and red arrows indicate lagging

strand synthesis.

(B-F) ChIP-qPCR analyses of helicase (DnaC) and replication restart proteins (PriA, DnaD,

DnaB, DnaI) in strains engineered to express either nCas9 or dCas9. The sgRNAs targeted

Cas9 proteins to a locus located at -90° and fold change was normalised to enrichment

values obtained at the +90° locus. Error bars indicate the standard error of the mean.

(G) Diagram illustrating the fate of helicase (red hexagon) upon encountering a single-strand

discontinuity in either the leading or lagging strand template.

(H) MFA using whole genome sequencing in strains engineered to express lower levels of

sgRNAs and Cas9 proteins (weak nicking system). The frequency of sequencing reads was

plotted against genome position.

(I) Genetic system employed to express low levels of PriA in a background encoding the

weak nicking system.

(J) Spot titre analysis of strains with a low level of PriA engineered to express nCas9

(+xylose). sgRNAs were targeted to a locus at -90°.

(K) MFA using whole genome sequencing in strains with a low level of PriA engineered to

express (+xylose) nCas9. The frequency of sequencing reads was plotted against genome

position.

Figure 3. The repair of single-strand discontinuities requires homologous

recombination.

(A) Spot titre analysis of homologous recombination mutants in strains engineered to

express (+xylose) nCas9. sgRNAs were targeted to a locus at -90° and control strains

correspond to parental strains.

(B) MFA using whole genome sequencing of homologous recombination mutants in strains

engineered to express nCas9. The frequency of sequencing reads was plotted against

genome position.

(C) Diagram illustrating proposed molecular events required for homologous recombination

to repair the DSB created after a replication fork encounters a single-strand discontinuity in

the leading strand template.

Figure 4. The SSB-CTT is essential to recruit RecO following replisome inactivation at

a single-strand discontinuity.

(A) ChIP-qPCR analyses of SSB in strains engineered to express nCas9. The sgRNAs

targeted nCas9 to a locus located at -90° and fold change was normalised to enrichment

values obtained at the +90° locus. Error bars indicate the standard error of the mean.

(B) ChIP-qPCR analyses of SSB in strains with homologous recombination mutants

engineered to express nCas9. qPCR primers amplified either a site located at -3 kb from the

sgRNA target (-90°) or a control locus (+90°) used for normalisation. Error bars indicate the

standard error of the mean.

(C) Spot titre analysis of SSB variants in strains engineered to express (+xylose) nCas9. Top

panel shows the genetic system employed in this experiment. sgRNAs were targeted to a

locus at -90°.

(D) MFA using whole genome sequencing in strains with SSB variants engineered to

express nCas9. The frequency of sequencing reads was plotted against genome position.

(E) ChIP-qPCR analyses of SSB, His-RecO and His-RecA proteins in strains engineered to

express nCas9. qPCR primers amplified either a site located at -3 kb from the sgRNA target

(-90°) or a control locus (+90°) used for normalisation. Error bars indicate the standard error

of the mean.

(F) ChIP-qPCR analyses of His-RecA in strains with homologous recombination mutants

engineered to express nCas9. Top panel shows the genetic system employed in this

experiment. qPCR primers amplified either a site located at -3 kb from the sgRNA target (-

90°) or a control locus (+90°) used for normalisation. Error bars indicate the standard error of

the mean.

Figure 5. DNA end resection via AddAB is not required to recruit SSB following

replisome inactivation at a single-strand discontinuity.

(A) MFA using whole genome sequencing in a Δ*recG* mutant strain engineered to express

nCas9. The frequency of sequencing reads was plotted against genome position (same data

as in Fig. 3B Δ*recG* panel). A region flanking the sgRNA binding site is enlarged.

(B) Diagram illustrating incorrect helicase reloading leading to reverse restart on a repaired

replication fork in a Δ*recG* mutant. Helicase is shown as a red hexagon and RecG is shown

in green; protein movement is indicated by colour-corresponding arrows. Lagging strand and

leading strand templates are represented in black and grey lines, respectively. Blue arrows

indicate leading strand synthesis.

(C) ChIP-qPCR analyses of mNG-AddA and mNG-AddB in strains engineered to express

nCas9. Left panel shows the genetic system employed in this experiment. The sgRNAs

targeted nCas9 to a locus located at -90° and fold change was normalised to enrichment

values obtained at the +90° locus. Error bars indicate the standard error of the mean.

(D) Spot titre analysis in strains with either AddA$^{\Delta HEL}$, AddA$^{\Delta NUC}$ or AddB$^{\Delta NUC}$ engineered to

express (+xylose) nCas9. sgRNAs were targeted to a locus at -90° and control strains

correspond to parental strains.

(E) MFA using whole genome sequencing of strains with either AddA$^{\Delta HEL}$ or

AddA$^{\Delta NUC}$/AddB$^{\Delta NUC}$ engineered to express nCas9. The frequency of sequencing reads was

plotted against genome position. Top panel shows the genetic system employed in this

experiment.

(F) ChIP-qPCR analyses of SSB in strains with AddA and AddB variants engineered to

express nCas9. Alleles refer to wild-type (WT), knockouts (Δ), and point mutations

inactivating nuclease (ΔNUC) or helicase (ΔHEL) activities. qPCR primers amplified either a

site located at -3 kb from the sgRNA target (-90°) or a control locus (+90°) used for

normalisation. Error bars indicate the standard error of the mean.

Figure 6. RecG is dispensable when DSB processing degrades the strand with a 5′-

end.

(A) Diagram illustrating differential resection of a DSB in the presence of either the AddA or

AddB nuclease activity. Leading and lagging strand templates are indicated by grey and blue

lines, respectively (numbers indicate polarity). Dotted lines indicate nuclease cleavage.

(B) Genetic system employed to investigate the role of RecG in DSB repair following

replisome inactivation at a single-strand discontinuity.

(C) Spot titre analysis in a Δ*recG* mutant strain with either AddA$^{\Delta NUC}$ or AddB$^{\Delta NUC}$ engineered

to express (+xylose) nCas9. sgRNAs were targeted to a locus at -90° and control strains

correspond to parental strains encoding the weak nicking system.

(D) Diagram illustrating differential helicase reloading events on a repaired replication fork

when AddA or AddB nuclease activity is inactivated in the absence of RecG. Helicase is

shown as a red hexagon and PriA in yellow; protein movement is indicated by colour-

corresponding arrows. Lagging strand and leading strand templates are represented in black

and grey lines, respectively. Blue arrows indicate leading strand synthesis.

Figure 7. Model for DSB repair and helicase reloading following replisome inactivation

at a single-strand discontinuity.

Revised model illustrating molecular events that are required or dispensable for homologous

repair of a DSB created after a replication fork encounters a single-strand discontinuity

located on either the leading or lagging strand template.

SUPPLEMENTARY FIGURE TITLES AND LEGENDS

Supplementary Figure 1. Impact of Cas9 variants on *B. subtilis* cell growth.

(A) Spot titre analysis of strains expressing (+ xylose) either Cas9^{D10A} nickase, Cas9^{H840A}

nickase, or Cas9$^{D10A-H840A}$ catalytically inactive nuclease in the presence of the same sgRNA.

Left panel shows the genetic system employed in this experiment. sgRNAs were targeted to

a locus at -90°.

(B) Immunoblot of Cas9 variants in the presence of a sgRNA targeted to a locus at -90°..

Cas9 proteins were induced (+xylose) during the exponential growth phase for 1 hour before

cells were harvested for protein extraction. The tubulin homolog FtsZ was used as a loading

control.

(C) Spot titre analysis of a strain expressing (+ xylose) the Cas9^{D10A} nickase with different

sgRNAs targeting different locations along the left chromosome arm.

(D) Spot titre analysis of strains expressing (+ xylose) the Cas9$^{D10A-H840A}$ catalytically inactive

nuclease. Left panel shows the genetic system employed in this experiment for sgRNAs

targeted to a locus at either -90° or +90°.

Supplementary Figure 2. MFA using whole genome sequencing of the parental strain

***B. subtilis* Δ6.**

The frequency of sequencing reads was plotted against genome position. Note that

chromosome physical boundaries indicated on x-axis may not represent termination events

occurring in the cell (*terminus* region highlighted in yellow).

Supplementary Figure 3. Live cell fluorescence microscopy using the mNG-RecA

reporter of DNA damage.

(A) Phase contrast image and fluorescence signal from the mNG-RecA reporter in wild-type

B. subtilis under standard laboratory growth conditions.

(B) Representative fluorescence microscopy images of mNG-RecA in strains engineered to

express either nCas9 or dCas9 (sgRNA target -90°). Fluorescent RecA organises into

bundles in the presence of DSBs. Green signal corresponds to mNG-RecA fluorescence,

grey scale images show corresponding phase contrast images. Cas9 proteins were induced

(+xylose) during the exponential growth phase for 1 hour before cells were collected for

imaging.

Supplementary Figure 4. The replicative helicase DnaC is recruited near a single-

strand discontinuity.

ChIP-qPCR analyses of helicase (DnaC) in strains engineered to express either nCas9 or

dCas9. The sgRNAs targeted Cas9 proteins to a locus located at -90°. The replication origin

oriC was used as positive control and fold change was normalised to enrichment values

obtained at the +90° locus. Error bars indicate the standard error of the mean.

Supplementary Figure 5. The weak nicking system does not arrest cell growth.

(A) Schematic showing differences between the strong and weak nicking systems to

introduce a single-strand discontinuity.

(B) Spot titre analysis of strains expressing (+ xylose) the strong of weak nicking system in

the presence of the same sgRNA. Control indicates a strain encoding the Cas9[D10A-H840A]

catalytically inactive nuclease. sgRNAs targeted to a locus located at -90°.

(C) Live cell fluorescence microscopy using nucleoid and membrane stains in strains

expressing either the strong or the weak nicking system. Grey scale images correspond to

phase contrast, blue signal corresponds to DAPI staining (nucleoid), red signal corresponds

to Nile red staining (membrane) and merge corresponds to DAPI/Nile red composite images.

Cas9 proteins were induced (+xylose) during the exponential growth phase for 90 minutes

before cells were collected for imaging.

Supplementary Figure 6. Complementation system employed to investigate PriA

essentiality following replisome inactivation at single-strand discontinuities.

(A) Genetic system employed for *priA* complementation via the IPTG inducible *priA-ssrA*

cassette placed at an ectopic locus in wild-type *B. subtilis*.

(B) Spot titre analysis of strains expressing (+ IPTG) the *priA-ssrA* complementation system.

(C) Immunoblot of PriA variants expressed at variable IPTG concentrations. The tubulin

homolog FtsZ was used as a loading control.

(D) Spot titre analysis of strains with a very low or high level of PriA engineered to express

nCas9 (+xylose). sgRNAs were targeted to a locus at -90°.

Supplementary Figure 7. DSB repair following replisome inactivation at a single-

strand discontinuity is achieved via homologous recombination.

(A) Spot titre analysis of homologous recombination mutants in strains engineered to

express (+xylose) nCas9. sgRNAs were targeted to a locus at -90° and control strains

correspond to parental strains. Plates were the same as shown in Fig. 3A when incubated

up to 48 hours.

(B) Spot titre analysis of homologous recombination mutants complemented with epitope-

tagged ectopic gene copies in strains engineered to express (+xylose) nCas9. sgRNAs were

targeted to a locus at -90° and control strains correspond to parental strains.

(C) Diagram illustrating proposed molecular events required for homologous recombination

to repair the DSB created after a replication fork encounters a single-strand discontinuity in

the lagging strand template.

Supplementary Figure 8. DSB repair following replisome inactivation at a single-

strand discontinuity relies on a minimal set of non-essential genes.

(A-D) Spot titre analysis of DNA repair mutants grouped by known/proposed functions in

strains engineered to express (+xylose) nCas9. sgRNAs were targeted to a locus at -90°

and control strains correspond to parental strains.

Supplementary Figure 9. Complementation system employed to investigate the

function of the SSB-CTT following replisome inactivation at single-strand

discontinuities.

(A) Schematic of SSB primary structure with corresponding functions. Numbers indicate

amino acid residue boundaries between the N-terminal domain (NTD), C-terminal domain

(CTD) and C-terminal tail (CTT).

(B) Model *of B. subtilis* SSB homotetramer in the presence of single-strand DNA (AlphaFold

3). The black backbone corresponds to a homopolymer of 60 thymidine bases and SSB-CTT

residues are highlighted in red.

(C) Genetic system employed for *ssbA* complementation via an IPTG inducible cassette

placed at an ectopic locus in wild-type *B. subtilis.*

(D) Spot titre analysis of strains expressing (+ IPTG) *ssbA* variants using the

complementation system.

(E) Immunoblot of SSB variants expressed using the *ssbA* complementation system. The

tubulin homolog FtsZ was used as a loading control.

(F) Genetic system employed to express SSB variants in a background encoding the weak

nicking system.

(G) Diagram illustrating potential essential interactions of the SSB-CTT.

(H) Model of a complex between an SSB homotetramer and RecO from *B. subtilis*

(AlphaFold 3). Cyan corresponds to SSB with CTT residues coloured in red and RecO is

displayed in beige.

Supplementary Figure 10. Traditional homologous repair models require end

resection to generate 3′-tailed DNA.

Diagram illustrating textbook molecular events required for homologous recombination to

repair the DSB created after a replication fork encounters a single-strand discontinuity in

either the leading or the lagging strand template.

Supplementary Figure 11. PriA recognises D-loops where the 3′-OH group of the

nascent leading strand abuts the fork junction.

Diagram illustrating the substrate recognised by PriA to orchestrate correct helicase

reloading.

Supplementary Figure 12. End-processing activities required following replisome

inactivation at a single-strand discontinuity.

(A) Immunoblot of AddA variants expressed from *mNG* epitope-tagged fusions. The tubulin

homolog FtsZ was used as a loading control.

(B-C) MFA using whole genome sequencing of strains with either AddA$^{\Delta HEL}$, AddA$^{\Delta NUC}$ or

AddB$^{\Delta NUC}$ engineered to express nCas9. The frequency of sequencing reads was plotted

against genome position. AddA$^{\Delta HEL}$ samples were the same as shown in Fig. 5E.

(D) Crystal structure of the *B. subtilis* AddAB complex with locations of catalytic sites and the

AddA^{D682} residue indicated (PDB template: 4CEH). The black backbone corresponds to

DNA,

AddA is displayed in orange and AddB in beige.

SUPPLEMENTAL INFORMATION

Documents S1-S12. Figures S1-S12, related to main Figures.

Table S1. Excel file too large to fit in a PDF containing the list of repair gene knockouts

tested in the genetic screen, related to Figures 3 and S8.

Table S2. Excel file too large to fit in a PDF containing genotypes from strains generated in

this study, related to the Key resources table.

Table S3. Excel file too large to fit in a PDF containing genotypes and sequences of

recombinant DNA used to construct strains in this study, related to the Key resources table

and Table S2.

Table S4. Excel file too large to fit in a PDF containing oligonucleotide sequences used for

qPCR in this study, related to the Key resources table.

REFERENCES

1. Murray, C.J.L., Ikuta, K.S., Sharara, F., Swetschinski, L., Aguilar, G.R., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E., et al. (2022). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. Lancet *399*, 629-655. 10.1016/S0140-6736(21)02724-0.

2. Butler, M.S., Vollmer, W., Goodall, E.C.A., Capon, R.J., Henderson, I.R., and Blaskovich, M.A.T. (2024). A Review of Antibacterial Candidates with New Modes of Action. Acs Infect Dis *10*, 3440-3474. 10.1021/acsinfecdis.4c00218.

3. Dwyer, D.J., Kohanski, M.A., and Collins, J.J. (2009). Role of reactive oxygen species in antibiotic action and resistance. Curr Opin Microbiol *12*, 482-489. 10.1016/j.mib.2009.06.018.

4. Charbon, G., Bjorn, L., Mendoza-Chamizo, B., Frimodt-Moller, J., and Lobner-Olesen, A. (2014). Oxidative DNA damage is instrumental in hyperreplication stress-induced inviability of. Nucleic Acids Research *42*, 13228-13241. 10.1093/nar/gku1149.

5. Kohanski, M.A., Dwyer, D.J., Hayete, B., Lawrence, C.A., and Collins, J.J. (2007). A common mechanism of cellular death induced by bactericidal antibiotics. Cell *130*, 797-810. 10.1016/j.cell.2007.06.049.

6. Fan, X.Y., Tang, B.K., Xu, Y.Y., Han, A.X., Shi, K.X., Wu, Y.K., Ye, Y., Wei, M.L., Niu, C., Wong, K.W., et al. (2018). Oxidation of dCTP contributes to antibiotic lethality in stationary-phase mycobacteria. Proc Natl Acad Sci U S A *115*, 2210-2215. 10.1073/pnas.1719627115.

7. Lenhart, J.S., Schroeder, J.W., Walsh, B.W., and Simmons, L.A. (2012). DNA repair and genome maintenance in *Bacillus subtilis*. Microbiol Mol Biol Rev *76*, 530-564. 10.1128/MMBR.05020-11.

8. Gaimster, H., Stevens, D., Hubbard, J., Mickiewicz, K., Murray, H., and Winterhalter, C. (2024). Conflicts between the DNA replication and repair machineries promote cell death in Gram-positive bacteria. bioRxiv, 2024.2010.2024.619994. 10.1101/2024.10.24.619994.

9. Lindahl, T., and Wood, R.D. (1999). Quality control by DNA repair. Science *286*, 1897-1905. DOI 10.1126/science.286.5446.1897.

10. Michel, B., Sinha, A.K., and Leach, D.R.F. (2018). Replication Fork Breakage and Restart in Escherichia coli. Microbiol Mol Biol Rev *82*. 10.1128/MMBR.00013-18.

11. Kuzminov, A. (2001). Single-strand interruptions in replicating chromosomes cause double-strand breaks. Proc Natl Acad Sci U S A *98*, 8241-8246. 10.1073/pnas.131009198.

12. Kornberg, A., and Baker, T.A. (1992). DNA replication, 2 Edition (W.H. Freeman and Co).

13. O'Donnell, M. (2006). Replisome architecture and dynamics in Escherichia coli. J Biol Chem *281*, 10653-10656. 10.1074/jbc.R500028200.

14. Kim, D.E., Narayan, M., and Patel, S.S. (2002). T7 DNA helicase: a molecular motor that processively and unidirectionally translocates along single-stranded DNA. J Mol Biol *321*, 807-819. 10.1016/s0022-2836(02)00733-7.

15. Xu, Z.Q., and Dixon, N.E. (2018). Bacterial replisomes. Curr Opin Struct Biol *53*, 159-168. 10.1016/j.sbi.2018.09.006.

16. Fernandez, A.J., and Berger, J.M. (2021). Mechanisms of hexameric helicases. Crit Rev Biochem Mol Biol *56*, 621-639. 10.1080/10409238.2021.1954597.

17. Beattie, T.R., Kapadia, N., Nicolas, E., Uphoff, S., Wollman, A.J., Leake, M.C., and Reyes-Lamothe, R. (2017). Frequent exchange of the DNA polymerase during bacterial chromosome replication. Elife *6*. 10.7554/eLife.21763.

18. Graham, J.E., Marians, K.J., and Kowalczykowski, S.C. (2017). Independent and Stochastic Action of DNA Polymerases in the Replisome. Cell *169*, 1201-1213 e1217. 10.1016/j.cell.2017.05.041.

19.   Yeeles, J.T., and Marians, K.J. (2011). The Escherichia coli replisome is inherently
DNA damage tolerant. Science *334*, 235-238. 10.1126/science.1209111.
20.   Vrtis, K.B., Dewar, J.M., Chistol, G., Wu, R.A., Graham, T.G.W., and Walter, J.C.
(2021). Single-strand DNA breaks cause replisome disassembly. Mol Cell *81*, 1309-
1318 e1306. 10.1016/j.molcel.2020.12.039.
21.   Pavani, R., Tripathi, V., Vrtis, K.B., Zong, D., Chari, R., Callen, E., Pankajam, A.V.,
Zhen, G., Matos-Rodrigues, G., Yang, J., et al. (2024). Structure and repair of
replication-coupled DNA breaks. Science *385*, eado3867. 10.1126/science.ado3867.
22.   Jinek, M., East, A., Cheng, A., Lin, S., Ma, E.B., and Doudna, J. (2013). RNA-
programmed genome editing in human cells. Elife *2*. ARTN e00471
10.7554/eLife.00471.
23.   Xu, Y., Morrow, C.A., Laksir, Y., Holt, O.M., Taylor, K., Tsiappourdhi, C., Collins, P.,
Jia, S., Andreadis, C., and Whitby, M.C. (2025). DNA nicks in both leading and
lagging strand templates can trigger break-induced replication. Mol Cell *85*, 91-106
e105. 10.1016/j.molcel.2024.10.026.
24.   Lee, J., Lim, K., Kim, A., Mok, Y.G., Chung, E., Cho, S.I., Lee, J.M., and Kim, J.S.
(2023). Prime editing with genuine Cas9 nickases minimizes unwanted indels.
Nature Communications *14*. 10.1038/s41467-023-37507-8.
25.   Peters, J.M., Colavin, A., Shi, H.D., Czarny, T.L., Larson, M.H., Wong, S., Hawkins,
J.S., Lu, C.H.S., Koo, B.M., Marta, E., et al. (2016). A Comprehensive, CRISPR-
based Functional Analysis of Essential Genes in Bacteria. Cell *165*, 1493-1506.
10.1016/j.cell.2016.05.003.
26.   Westers, H., Dorenbos, R., van Dijl, J.M., Kabel, J., Flanagan, T., Devine, K.M.,
Jude, F., Seror, S.J., Beekman, A.C., Darmon, E., et al. (2003). Genome engineering
reveals large dispensable regions in Bacillus subtilis. Mol Biol Evol *20*, 2076-2090.
10.1093/molbev/msg219.
27.   Simmons, L.A., Grossman, A.D., and Walker, G.C. (2007). Replication is required for
the RecA localization response to DNA damage in Bacillus subtilis. Proc Natl Acad
Sci U S A *104*, 1360-1365. 10.1073/pnas.0607123104.
28.   Lesterlin, C., Ball, G., Schermelleh, L., and Sherratt, D.J. (2014). RecA bundles
mediate homology pairing between distant sisters during DNA break repair. Nature
506, 249-253. 10.1038/nature12868.
29.   Kidane, D., and Graumann, P.L. (2005). Dynamic formation of RecA filaments at
DNA double strand break repair centers in live cells. J Cell Biol *170*, 357-366.
10.1083/jcb.200412090.
30.   Smits, W.K., Goranov, A.I., and Grossman, A.D. (2010). Ordered association of
helicase loader proteins with the Bacillus subtilis origin of replication in vivo. Mol
Microbiol *75*, 452-461. 10.1111/j.1365-2958.2009.06999.x.
31.   Michel, B., and Sandler, S.J. (2017). Replication Restart in Bacteria. J Bacteriol *199*.
10.1128/JB.00102-17.
32.   Winterhalter, C., Pelliciari, S., Stevens, D., Fenyk, S., Marchand, E., Cronin, N.B.,
Soultanas, P., Costa, T.R.D., Ilangovan, A., and Murray, H. (2023). The DNA
replication initiation protein DnaD recognises a specific strand of the Bacillus subtilis
chromosome origin. Nucleic Acids Res *51*, 4322-4340. 10.1093/nar/gkad277.
33.   Winterhalter, C., Stevens, D., Fenyk, S., Pelliciari, S., Marchand, E., Soultanas, P.,
Ilangovan, A., and Murray, H. (2022). SirA inhibits the essential DnaA:DnaD
interaction to block helicase recruitment during Bacillus subtilis sporulation. Nucleic
Acids Res. 10.1093/nar/gkac1060.
34.   Matthews, L.A., and Simmons, L.A. (2020). Regulation of DNA Binding and High-
Order Oligomerization of the DnaB Helicase Loader. J Bacteriol *202*.
10.1128/JB.00286-20.
35.   Bruand, C., Farache, M., McGovern, S., Ehrlich, S.D., and Polard, P. (2001). DnaB,
DnaD and DnaI proteins are components of the Bacillus subtilis replication restart
primosome. Molecular Microbiology *42*, 245-255. DOI 10.1046/j.1365-
2958.2001.02631.x.

36.   Kaguni, J.M. (2011). Replication initiation at the Escherichia coli chromosomal origin.
1118   Curr Opin Chem Biol *15*, 606-613. 10.1016/j.cbpa.2011.07.016.
1119   37.   Kaplan, D.L. (2000). The 3'-tail of a forked-duplex sterically determines whether one
1120   or two DNA strands pass through the central channel of a replication-fork helicase. J
1121   Mol Biol *301*, 285-299. 10.1006/jmbi.2000.3965.
1122   38.   Griffith, K.L., and Grossman, A.D. (2008). Inducible protein degradation in Bacillus
1123   subtilis using heterologous peptide tags and adaptor proteins to target substrates to
1124   the   protease   ClpXP.   Mol   Microbiol   *70*,   1012-1025.   10.1111/j.1365-
1125   2958.2008.06467.x.
1126   39.   Cox, M.M., Goodman, M.F., Kreuzer, K.N., Sherratt, D.J., Sandler, S.J., and Marians,
1127   K.J. (2000). The importance of repairing stalled replication forks. Nature *404*, 37-41.
1128   40.   Yasmin, T., Azeroglu, B., Cockram, C.A., and Leach, D.R.F. (2021). Distribution of
1129   Holliday junctions and repair forks during Escherichia coli DNA double-strand break
1130   repair. PLoS Genet *17*, e1009717. 10.1371/journal.pgen.1009717.
1131   41.   Sanchez, H., Kidane, D., Castillo Cozar, M., Graumann, P.L., and Alonso, J.C.
1132   (2006). Recruitment of Bacillus subtilis RecN to DNA double-strand breaks in the
1133   absence of DNA end processing. J Bacteriol *188*, 353-360. 10.1128/JB.188.2.353-
1134   360.2006.
1135   42.   Costes, A., Lecointe, F., McGovern, S., Quevillon-Cheruel, S., and Polard, P. (2010).
1136   The C-terminal domain of the bacterial SSB protein acts as a DNA maintenance hub
1137   at   active   chromosome   replication   forks.   PLoS   Genet   *6*,   e1001238.
1138   10.1371/journal.pgen.1001238.
1139   43.   Ramos, C., Hernandez-Tamayo, R., Lopez-Sanz, M., Carrasco, B., Serrano, E.,
1140   Alonso, J.C., Graumann, P.L., and Ayora, S. (2022). The RecD2 helicase balances
1141   RecA activities. Nucleic Acids Research *50*, 3432-3444. 10.1093/nar/gkac131.
1142   44.   Cárdenas, P.P., Carrasco, B., Soufo, C.D., César, C.E., Herr, K., Kaufenstein, M.,
1143   Graumann,   P.L.,   and   Alonso,   J.C.   (2012).   RecX   Facilitates   Homologous
1144   Recombination by Modulating RecA Activities. Plos Genetics *8*. ARTN e1003126
1145   10.1371/journal.pgen.1003126.
1146   45.   Mortier-Barrière, I., Velten, M., Dupaigne, P., Mirouze, N., Piétrement, O., McGovern,
1147   S., Fichant, G., Martin, B., Noirot, P., Le Cam, E., et al. (2007). A key presynaptic
1148   role in transformation for a widespread bacterial protein:: DprA conveys incoming
1149   ssDNA to RecA. Cell *130*, 824-836. 10.1016/j.cell.2007.07.038.
1150   46.   Azeroglu, B., and Leach, D.R.F. (2017). RecG controls DNA amplification at double-
1151   strand   breaks   and   arrested   replication   forks.   FEBS   Lett   *591*,   1101-1113.
1152   10.1002/1873-3468.12583.
1153   47.   Azeroglu, B., Mawer, J.S., Cockram, C.A., White, M.A., Hasan, A.M., Filatenkova,
1154   M., and Leach, D.R. (2016). RecG Directs DNA Synthesis during Double-Strand
1155   Break Repair. PLoS Genet *12*, e1005799. 10.1371/journal.pgen.1005799.
1156   48.   Symington, L.S. (2014). End resection at double-strand breaks: mechanism and
1157   regulation. Cold Spring Harb Perspect Biol *6*. 10.1101/cshperspect.a016436.
1158   49.   Yeeles, J.T., and Dillingham, M.S. (2007). A dual-nuclease mechanism for DNA
1159   break processing by AddAB-type helicase-nucleases. J Mol Biol *371*, 66-78.
1160   10.1016/j.jmb.2007.05.053.
1161   50.   Cox, M.M., Goodman, M.F., Keck, J.L., van Oijen, A., Lovett, S.T., and Robinson, A.
1162   (2023). Generation and Repair of Postreplication Gaps in Escherichia coli. Microbiol
1163   Mol Biol Rev *87*, e0007822. 10.1128/mmbr.00078-22.
1164   51.   Nirwal, S., Czarnocki-Cieciura, M., Chaudhary, A., Zajko, W., Skowronek, K.,
1165   Chamera, S., Figiel, M., and Nowotny, M. (2023). Mechanism of RecF-RecO-RecR
1166   cooperation in bacterial homologous recombination. Nat Struct Mol Biol *30*, 650-660.
1167   10.1038/s41594-023-00967-z.
1168   52.   Bell, J.C., and Kowalczykowski, S.C. (2016). RecA: Regulation and Mechanism of a
1169   Molecular Search Engine: (Trends in Biochemical Sciences, June 2016, Vol. 41, No.
1170   6, 491--507). Trends Biochem Sci *41*, 646. 10.1016/j.tibs.2016.05.006.

53. Lin, Z.G., Kong, H.Z., Nei, M., and Ma, H. (2006). Origins and evolution of the gene
family:: Evidence for ancient gene duplication and endosymbiotic gene transfer. P
Natl Acad Sci USA *103*, 10328-10333. 10.1073/pnas.0604232103.
54. West, S.C. (1997). Processing of recombination intermediates by the RuvABC
proteins. Annu Rev Genet *31*, 213-244. 10.1146/annurev.genet.31.1.213.
55. Bianco, P.R. (2015). I came to a fork in the DNA and there was RecG. Prog Biophys
Mol Biol *117*, 166-173. 10.1016/j.pbiomolbio.2015.01.001.
56. Lloyd, R.G., and Rudolph, C.J. (2016). 25 years on and no end in sight: a
perspective on the role of RecG protein. Curr Genet *62*, 827-840. 10.1007/s00294-
016-0589-z.
57. Shereda, R.D., Kozlov, A.G., Lohman, T.M., Cox, M.M., and Keck, J.L. (2008). SSB
as an organizer/mobilizer of genome maintenance complexes. Crit Rev Biochem Mol
Biol *43*, 289-318. 10.1080/10409230802341296.
58. Lecointe, F., Serena, C., Velten, M., Costes, A., McGovern, S., Meile, J.C., Errington,
J., Ehrlich, S.D., Noirot, P., and Polard, P. (2007). Anticipating chromosomal
replication fork arrest: SSB targets repair DNA helicases to active forks. EMBO J *26*,
4239-4251. 10.1038/sj.emboj.7601848.
59. Carrasco, B., Torres, R., Moreno-Del Alamo, M., Ramos, C., Ayora, S., and Alonso,
J.C. (2024). Processing of stalled replication forks in Bacillus subtilis. FEMS
Microbiol Rev *48*. 10.1093/femsre/fuad065.
60. Sanchez, H., Carrasco, B., Cozar, M.C., and Alonso, J.C. (2007). Bacillus subtilis
RecG branch migration translocase is required for DNA repair and chromosomal
segregation. Mol Microbiol *65*, 920-935. 10.1111/j.1365-2958.2007.05835.x.
61. Torres, R., Gandara, C., Carrasco, B., Baquedano, I., Ayora, S., and Alonso, J.C.
(2021). DisA Limits RecG Activities at Stalled or Reversed Replication Forks. Cells
10. 10.3390/cells10061357.
62. Torres, R., Romero, H., Rodriguez-Cerrato, V., and Alonso, J.C. (2017). Interplay
between Bacillus subtilis RecD2 and the RecG or RuvAB helicase in recombinational
repair. DNA Repair (Amst) *55*, 40-46. 10.1016/j.dnarep.2017.05.004.
63. Tanaka, T., and Masai, H. (2006). Stabilization of a stalled replication fork by
concerted actions of two helicases. J Biol Chem *281*, 3484-3493.
10.1074/jbc.M510979200.
64. McGlynn, P., Al-Deib, A.A., Liu, J., Marians, K.J., and Lloyd, R.G. (1997). The DNA
replication protein PriA and the recombination protein RecG bind D-loops. J Mol Biol
270, 212-221. 10.1006/jmbi.1997.1120.
65. Sun, B., Pandey, M., Inman, J.T., Yang, Y., Kashlev, M., Patel, S.S., and Wang, M.D.
(2015). T7 replisome directly overcomes DNA damage. Nat Commun *6*, 10260.
10.1038/ncomms10260.
66. Fernandez, S., Ayora, S., and Alonso, J.C. (2000). Bacillus subtilis homologous
recombination: genes and products. Res Microbiol *151*, 481-486. 10.1016/s0923-
2508(00)00165-0.
67. Unciuleac, M.C., and Shuman, S. (2010). Double strand break unwinding and
resection by the mycobacterial helicase-nuclease AdnAB in the presence of single
strand DNA-binding protein (SSB). J Biol Chem *285*, 34319-34329.
10.1074/jbc.M110.162925.
68. Gupta, R., Unciuleac, M.C., Shuman, S., and Glickman, M.S. (2017). Homologous
recombination mediated by the mycobacterial AdnAB helicase without end resection
by the AdnAB nucleases. Nucleic Acids Res *45*, 762-774. 10.1093/nar/gkw1130.
69. Su'etsugu, M., and Errington, J. (2011). The replicase sliding clamp dynamically
accumulates behind progressing replication forks in *Bacillus subtilis* cells. Mol Cell
41, 720-732. 10.1016/j.molcel.2011.02.024.
70. Reyes-Lamothe, R., Sherratt, D.J., and Leake, M.C. (2010). Stoichiometry and
architecture of active DNA replication machinery in Escherichia coli. Science *328*,
498-501. 10.1126/science.1185757.

71. Yeeles, J.T., van Aelst, K., Dillingham, M.S., and Moreno-Herrero, F. (2011).
Recombination hotspots and single-stranded DNA binding proteins couple DNA
translocation to DNA unwinding by the AddAB helicase-nuclease. Mol Cell *42*, 806-
816. 10.1016/j.molcel.2011.04.012.
72. Umezu, K., and Kolodner, R.D. (1994). Protein interactions in genetic recombination
in Escherichia coli. Interactions involving RecO and RecR overcome the inhibition of
RecA by single-stranded DNA-binding protein. J Biol Chem *269*, 30005-30013.
73. Hobbs, M.D., Sakai, A., and Cox, M.M. (2007). SSB protein limits RecOR binding
onto single-stranded DNA. J Biol Chem *282*, 11058-11067.
10.1074/jbc.M611007200.
74. Inoue, J., Honda, M., Ikawa, S., Shibata, T., and Mikawa, T. (2008). The process of
displacing the single-stranded DNA-binding protein from single-stranded DNA by
RecO and RecR proteins. Nucleic Acids Res *36*, 94-109. 10.1093/nar/gkm1004.
75. Manfredi, C., Carrasco, B., Ayora, S., and Alonso, J.C. (2008). Bacillus subtilis RecO
nucleates RecA onto SsbA-coated single-stranded DNA. J Biol Chem *283*, 24837-
24847. 10.1074/jbc.M802002200.
76. Bonde, N.J., Henry, C., Wood, E.A., Cox, M.M., and Keck, J.L. (2023). Interaction
with the carboxy-terminal tip of SSB is critical for RecG function in E. coli. Nucleic
Acids Res *51*, 3735-3753. 10.1093/nar/gkad162.
77. Buss, J.A., Kimura, Y., and Bianco, P.R. (2008). RecG interacts directly with SSB:
implications for stalled replication fork regression. Nucleic Acids Research *36*, 7029-
7042. 10.1093/nar/gkn795.
78. Kozlov, A.G., Jezewska, M.J., Bujalowski, W., and Lohman, T.M. (2010). Binding
Specificity of Single-Stranded DNA Binding Protein for the χ Subunit of DNA pol III
Holoenzyme and PriA Helicase. Biochemistry-Us *49*, 3555-3566. 10.1021/bi100069s.
79. Morimatsu, K., and Kowalczykowski, S.C. (2003). RecFOR proteins load RecA
protein onto gapped DNA to accelerate DNA strand exchange: a universal step of
recombinational repair. Mol Cell *11*, 1337-1347. 10.1016/s1097-2765(03)00188-6.
80. Lenhart, J.S., Brandes, E.R., Schroeder, J.W., Sorenson, R.J., Showalter, H.D., and
Simmons, L.A. (2014). RecO and RecR are necessary for RecA loading in response
to DNA damage and replication fork stress. J Bacteriol *196*, 2851-2860.
10.1128/JB.01494-14.
81. Henry, C., Mbele, N., and Cox, M.M. (2023). RecF protein targeting to postreplication
(daughter strand) gaps I: DNA binding by RecF and RecFR. Nucleic Acids Res *51*,
5699-5713. 10.1093/nar/gkad311.
82. Henry, C., Kaur, G., Cherry, M.E., Henrikus, S.S., Bonde, N.J., Sharma, N., Beyer,
H.A., Wood, E.A., Chitteni-Pattu, S., van Oijen, A.M., et al. (2023). RecF protein
targeting to post-replication (daughter strand) gaps II: RecF interaction with
replisomes. Nucleic Acids Res *51*, 5714-5742. 10.1093/nar/gkad310.
83. Smith, G.R. (2012). How RecBCD enzyme and Chi promote DNA break repair and
recombination: a molecular biologist's view. Microbiol Mol Biol Rev *76*, 217-228.
10.1128/MMBR.05026-11.
84. Conwell, S., Dahmen, S., Cranford, M., Long, D., Cortez, D., and Dewar, J. (2025).
Resolution of collapsed forks is separate from completion of DNA synthesis.
Submitted.
85. Ayora, S., Carrasco, B., Doncel-Perez, E., Lurz, R., and Alonso, J.C. (2004). Bacillus
subtilis RecU protein cleaves Holliday junctions and anneals single-stranded DNA.
Proc Natl Acad Sci U S A *101*, 452-457. 10.1073/pnas.2533829100.
86. Aussel, L., Barre, F.X., Aroyo, M., Stasiak, A., Stasiak, A.Z., and Sherratt, D. (2002).
FtsK Is a DNA motor protein that activates chromosome dimer resolution by
switching the catalytic state of the XerC and XerD recombinases. Cell *108*, 195-205.
87. Kato, J.-I., Nishimura, Y., Imamura, R., Niki, H., Hiraga, S., and Suzuki, H. (1990).
New topoisomerase essential for chromosome segregation in E. coli. Cell *63*, 393-
404.

88.   Lucet, I., Feucht, A., Yudkin, M.D., and Errington, J. (2000). Direct interaction
1280   between the cell division protein FtsZ and the cell differentiation protein SpoIIE.
1281   EMBO J. *19*, 1467-1475.
1282   89.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
1283   Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
1284   platform   for   biological-image   analysis.   Nat.   Methods   *9*,   676-682.
1285   10.1038/nmeth.2019.
1286   90.   Untergasser, A., Nijveen, H., Rao, X., Bisseling, T., Geurts, R., and Leunissen, J.A.
1287   (2007). Primer3Plus, an enhanced web interface to Primer3. Nucleic Acids Res *35*,
1288   W71-74. 10.1093/nar/gkm306.
1289   91.   Delano, W.L. (2002). The PyMOL Molecular Graphics System.
1290   92.   Hanahan, D., Jessee, J., and Bloom, F.R. (1991). Plasmid transformation of
1291   Escherichia coli and other bacteria. Methods Enzymol *204*, 63-113. 10.1016/0076-
1292   6879(91)04006-a.
1293   93.   Anagnostopoulos, C., and Spizizen, J. (1961). Requirements for transformation in
1294   *Bacillus subtilis*. J. Bacteriol. *81*, 741-746.
1295   94.   Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger,
1296   O., Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure
1297   prediction of biomolecular interactions with AlphaFold 3. Nature *630*, 493-500.
1298   10.1038/s41586-024-07487-w.
1299   95.   Krajewski, W.W., Fu, X., Wilkinson, M., Cronin, N.B., Dillingham, M.S., and Wigley,
1300   D.B. (2014). Structural basis for translocation by AddAB helicase-nuclease and its
1301   arrest at chi sites. Nature *508*, 416-419. 10.1038/nature13037.
1302

GRAPHICAL ABSTRACT

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

A

3′-terminal DNA degradation: AddB nuclease-deficient

5′-terminal DNA degradation: AddA nuclease-deficient

B

+ xylose

Nicking system // *ΔrecG* // *addB^{ΔNUC}* *addA^{ΔNUC/ΔHEL}* *kan*

C

	24 hours	48 hours
	- xylose / + xylose	- xylose / + xylose

Leading strand: Control, *ΔrecG*, *addA^{ΔNUC} ΔrecG*, *addB^{ΔNUC} ΔrecG*

Lagging strand: Control, *ΔrecG*, *addA^{ΔNUC} ΔrecG*, *addB^{ΔNUC} ΔrecG*

10^{-1} 10^{-2} 10^{-3} 10^{-4} 10^{-5} 10^{-6}

D

3′-terminal DNA degradation (AddA nuclease only)

5′-terminal DNA degradation (AddB nuclease only)

PriA

Helicase reloading on 5′-terminal DNA

Correct helicase reloading at D-loop

No leading strand 3′-OH: reverse restart

D-loop abuts 3'-OH: productive replication restart (role of RecG)

FIGURE 6

(I) Fork collapse:
Generation of a DSB and detection via AddAB

(II) End processing:
No strict requirement for DNA end resection

(III) SSB loading:
AddAB helicase activity required for SSB recruitment

(IV) Pre-synapsis:
Dependent on AddAB and RecO recruitment via SSB-CTT

(V) RecA recruitment:
Dependent on AddAB, RecFOR and SSB-CTT:RecO interaction

(VI) Strand exchange:
D-loop formation and presence of a 5′-tail

(VII) Fork regression:
RecG remodels the lagging strand, holliday junction cleavage dispensable

(VIII) PriA loading:
Recruitment of restart proteins for helicase reloading

(IX) Replication restart:
Helicase loaded on lagging strand and DNA synthesis resumes

FIGURE 7

SUPPLEMENTARY FIGURE 1

SUPPLEMENTARY FIGURE 2

SUPPLEMENTARY FIGURE 3

SUPPLEMENTARY FIGURE 4

A

Bacillus subtilis chromosome

| zeo | *nCas9** | // | cat | sgRNA |

| Nicking system | Strong | P_{XYL}-nCas9 | P_{VEG}-sgRNA |
| | Weak | P_{XYL}-nCas9-ssrA | P_{XYL}-sgRNA |

B

- xylose   + xylose

10^{-1} 10^{-2} 10^{-3} 10^{-4} 10^{-5} 10^{-6} 10^{-1} 10^{-2} 10^{-3} 10^{-4} 10^{-5} 10^{-6}

Leading strand
- Control
- Strong
- Weak

Lagging strand
- Control
- Strong
- Weak

C

| Phase contrast | Nucleoid | Membrane | Merge |

Wild-type

Leading strand
- Strong
- Weak

Lagging strand
- Strong
- Weak

SUPPLEMENTARY FIGURE 5

A

PriA complementation

B

	No IPTG	+ 0.05 mM IPTG	+ 1 mM IPTG

Wild-type -

Wild-type ssrA

Δ ssrA

C

Endogenous *priA* allele	WT	WT	Δ
Ectopic *priA* allele	-	ssrA	ssrA
IPTG (mM)	0 0.05 1	0 0.05 1	0 0.05 1

anti-PriA   ← PriA/PriA-ssrA
            ← Non specific

anti-FtsZ

D

Nicking system

PriA-ssrA induction (IPTG)

priA allele   - xylose   + xylose

Leading strand Wild-type / Δ   No IPTG

Lagging strand Wild-type / Δ

Leading strand Wild-type / Δ   + 1 mM

Lagging strand Wild-type / Δ

SUPPLEMENTARY FIGURE 6

A

DNA repair mutants	Leading strand		Lagging strand		
	- xylose (48 h)	+ xylose (48 h)	- xylose (48 h)	+ xylose (48 h)	
Control					
ΔaddA					End processing
ΔaddB					
ΔrecF					Early HR
ΔrecO					
ΔrecR					
ΔrecA					Synapsis
ΔrecG					Late HR
ΔrecU					

Homologous recombination

B

DNA repair mutants	Leading strand		Lagging strand	
	- xylose	+ xylose	- xylose	+ xylose
Control				
ΔaddA mNG-addA				
ΔaddB mNG-addB				
ΔrecF his-recF				
ΔrecO his-recO				
ΔrecR his-recR				
ΔrecA his-recA				
ΔrecG his-recG				

C

Lagging strand discontinuity homologous repair

B A AddA-AddB end-processing machinery **SSB tetramer** **OR F** RecOR:RecF complex **RecA filament** **RecU resolvase**

Collapsed fork
Leading strand
Lagging strand

End processing
B
A

SSB loading
End resection
A
B

Pre-synapsis
OR F

RecA recruitment
OR F

Strand exchange

Post-synapsis
RecU

SUPPLEMENTARY FIGURE 7

SUPPLEMENTARY FIGURE 8

SUPPLEMENTARY FIGURE 9

SUPPLEMENTARY FIGURE 10

Repaired replication fork following introduction of a leading strand discontinuity

Nascent leading strand

Repaired replication fork following introduction of a lagging strand discontinuity

Nascent leading strand

Correct substrate for PriA loading

SUPPLEMENTARY FIGURE 11

SUPPLEMENTARY FIGURE 12