TITLE

Rescuing the bacterial replisome at a nick requires recombinational repair and helicase

reloading

AUTHOR LIST

Charles Winterhalter[1*], Kathryn J Stratton[1], Stepan Fenyk[1], Heath Murray[1**]

*Correspondence: Charles.winterhalter@newcastle.ac.uk

**Correspondence: Heath.murray@newcastle.ac.uk

AFFILIATION

[1]Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle

Upon Tyne, NE2 4AX, UK

ABSTRACT

DNA damage occurs in all cells and must be repaired to maintain genome integrity. Many

DNA lesions are targeted for removal by repair systems that excise the damage, thereby

generating a temporary single-strand discontinuity in the chromosome. If DNA repair has not

been completed prior to a round of genome duplication, the single-strand discontinuity (nick

or gap) can be converted to a double-strand break (DSB) by an oncoming replication fork.

Because the genomic location of nucleobase damage is stochastic, investigating the fate of

replication machinery (replisome) at DNA repair sites with single-strand discontinuities has

been limited. Here we have addressed this issue by expressing Cas9 nickases in *Bacillus*

subtilis to create site specific single-strand discontinuities in a bacterial chromosome. We

find that a nick in either leading or lagging strand arrests DNA replication, while the fate of

the replicative helicase is distinct and depends upon the strand nicked. Genetic,

biochemical, and single cell analyses indicate that replisome/nick encounters generate a

single-end DSB which requires recombinational repair to enable PriA-dependent replication

restart. Together this work defines the physiologically relevant pathway used by *B. subtilis* to

reinitiate DNA synthesis following replication fork inactivation at a single-strand discontinuity.

Highlights

• Single-strand discontinuities inactivate the bacterial replisome

• Nicks on the leading or lagging strand template differentially affect fate of the

helicase

• Replication forks are repaired via recombinational repair

• The single strand binding protein (SSB) acidic tail is essential for replication restart

• DNA replication restart is PriA-dependent

Keywords

DNA, replication, repair, nick, helicase, fork, homologous recombination, PriA, restart.

INTRODUCTION

Intracellular DNA can become damaged and/or distorted through a multitude of endogenous and exogenous sources (e.g. post-replication mismatches, base oxidation, cyclobutane pyrimidine dimers)[1, 2]. Damaged and distorted DNA can be mutagenic, resulting in permanent changes to genetic information of progeny. To avoid these negative consequences on genome stability, specialised DNA repair systems recognise and remove the damaged (base/nucleotide excision repair) or distorted (mismatch repair) DNA.

As part of their repair mechanism, these systems excise the damaged or mismatched base, causing the formation of a nick or single-strand gap in the affected DNA strand[3]. Critically, if a replication fork progresses towards such a single-strand discontinuity before repair is complete, the encounter can generate a single-end double-strand break (seDSB)[4]. Elegant experiments using bacteriophage λ demonstrated that an engineered nick in either strand of the viral genome generated replication-dependent DSBs (Fig. 1A)[5]. While active DNA replication was necessary to generate DSBs at a nick[5], the fate of the bacterial DNA replication machinery (replisome) at single-strand discontinuities has not been determined.

The replisome is a multicomponent complex of proteins and enzymes that unwinds the chromosome and coordinates synthesis of new DNA polymers on the leading and lagging strands[6, 7]. The most stable component of the replisome, and most pivotal for its processivity, is helicase[8, 9]. Cellular replicative helicases are toroid enzymes that unwind DNA by encircling and translocating along a single client nucleic acid strand, acting as a wedge to disrupt base pairing[10]. Studies of bacterial replisomes *in vitro* and *in vivo* have revealed that the machinery is intrinsically dynamic, with most factors observed to bind and unbind interaction partners multiple times per replication cycle[11, 12]. This behaviour enables the replisome to overcome a range of potential roadblocks and remain associated with the template, competent to continue DNA synthesis[13]. However, recent studies in eukaryotic systems have indicated that single-strand discontinuities generate DSBs and arrest DNA replication by impairing helicase progression[14, 15].

Most bacterial chromosomes contain a single replication origin (*oriC*) that promotes

assembly of two independent replication forks for bidirectional DNA synthesis[16, 17].

Consequently, replisomes initiated from *oriC* must continue until they complete replication of

their respective chromosome arm[18, 19, 20, 21]. In contrast, eukaryotic chromosomes contain

multiple replication origins, therefore no single replication fork is essential to complete

genome duplication[15, 16, 22]. If a bacterial replisome encounters a single-strand discontinuity

and generates a DSB, it has been proposed that rescue of the broken chromosome occurs

through homologous recombination and that reassembly of the replisome follows helicase

loading at the repaired replication fork[23].

Determining how bacteria react and respond when a replication fork encounters a

single-strand discontinuity is crucial to understanding how these organisms ensure genome

stability. Here we have investigated this question by expressing Cas9 nickases (nCas9) in

the model bacterium *Bacillus subtilis*. The results show that a nick located in either template

strand blocks downstream DNA synthesis. Resumption of DNA replication is dependent

upon a core set of recombinational repair proteins and the PriA-dependent replication restart

system to reload the replicative helicase.

RESULTS

Single-strand discontinuities in a bacterial chromosome arrest DNA synthesis

Damaged and mismatched nucleobases can be located randomly throughout a genome,

making it a challenge to directly analyse conflicts between DNA replication and repair. To

mimic the formation of a single-strand discontinuity produced during DNA repair, an

experimental system capable of generating a nick was developed in *B. subtilis* using Cas9

nuclease variants and single-guide RNAs (sgRNA)[14, 15, 22, 24]. Cas9 carrying single amino

acid substitutions (Cas9^{D10A} or Cas9^{H840A}) creates nickases that can cleave the

phosphodiester bond of one DNA strand (nCas9), while the double mutant (Cas9$^{D10A/H840A}$)

creates a catalytically inactive enzyme (dCas9)[25]. Each *cas9* allele was integrated into the *B.*

subtilis chromosome under the control of a xylose-inducible promoter (Fig. 1B)[26]. The

location of sgRNA hybridisation is indicated by coordinates between the origin (0°) and

terminus (~180°) on either the right (+) or left (-) arm of the chromosome. To avoid

complications arising from potential activation of prophage following DNA damage in the

laboratory strain of *B. subtilis*, a derivative lacking these genetic elements (Δ6) was used

throughout this study[27].

Cell viability following nicking was quantified using spot-titre assays, where strains

are grown to saturation in liquid medium without xylose, before samples were serially diluted

and plated onto solid medium with or without xylose to enumerate the number of colony

forming units (CFUs). It was observed that the Cas9^{D10A} nickase elicits a stronger phenotype

than Cas9^{H840A} upon induction with xylose (Supplementary Figure 1A-B)[25]. To achieve

comparable nicking efficiency at different sites in the chromosome, the *cas9^{D10A}* allele was

used throughout the remainder of this study, in conjunction with distinct sgRNAs targeted to

non-essential regions of the genome. Using sgRNAs to target either leading or lagging

strands at loci on the left or right chromosome arms, it was observed that expression of

nCas9 resulted in severe growth inhibition (Fig. 1C, Supplementary Figure 1C). In contrast,

expression of dCas9 targeted to the same locations did not affect the number of CFUs (Fig.

1C, Supplementary Figure 1D), indicating that DNA nicking is necessary to cause growth

inhibition.

Next chromosome replication was investigated using whole genome marker

frequency analysis (MFA). Expression of nCas9 was induced for one hour in exponentially

growing cultures. Genomic DNA (gDNA) was harvested for next-generation sequencing,

reads were mapped onto the reference genome sequence, and data is displayed with the

chromosome origin (*oriC*) at the centre of the graphs (Supplementary Figure 2). Introduction

of nicks in either the leading or the lagging strand resulted in a drop in sequencing coverage

from the nick site to the terminus region, whereas expression of dCas9 with the same

sgRNAs did not appear to inhibit DNA replication (Fig. 1D). Copy number at the nick position

was low in the presence of nCas9, potentially reflecting DNA degradation[28]. Consistent with

a DNA replication-dependent phenotype, spot-titre analyses showed that nick induction in

late-exponential phase does not result in loss of viability, whereas induction in early-

exponential phase results in severe growth defects (Supplementary Figure 1E). Therefore,

because cells expressing nCas9 are unable to synthetise DNA downstream of the nick,

incomplete chromosome synthesis can explain the strong growth inhibition observed in spot-

titre assays (Fig. 1C).

Previous results indicated that a bacterial replication fork encountering a single-

strand discontinuity generates a DSB[5]. To determine whether DNA damage was generated

following nCas9 induction, localisation of an ectopic mNeonGreen-RecA reporter (mNG-

RecA) was observed in live cells using fluorescence microscopy (endogenous *recA* is

present in these merodiploid strains). It has been shown in *B. subtilis* that following DNA

damage fluorescently labelled RecA recombinase assembles into bundles[29, 30, 31]. Consistent

with the literature, in unstressed cells mNG-RecA is observed evenly distributed over the

nucleoid (Supplementary Figure 3A)[32]. Following expression of nCas9 for 90 minutes during

exponential growth, mNG-RecA bundles were detected in 59% of cells when the leading

strand template was nicked and 48% of cells when the lagging strand template was nicked

(Fig. 1E-F, Supplementary Figure 3B). In contrast, expression of dCas9 did not significantly

promote mNG-RecA bundling (Fig. 1E-F, Supplementary Figure 3B), validating that mNG-

RecA is a functional reporter for DNA damage. These results are consistent with the model

that the bacterial replisome generates DNA damage upon encountering of a nick in either

strand of the parental DNA duplex.

Single-strand discontinuities in a bacterial chromosome inactivate the replisome

MFA indicated that following expression of nCas9, DNA synthesis does not proceed beyond

the nick. We wondered whether the replisome is arrested at a single-strand discontinuity or

whether it is inactivated. If the latter event occurs, then the bacterial replication restart

system would be required to reload helicase and promote replisome assembly[33, 34]. In *B.*

subtilis there is only one known replication restart system composed of the essential proteins

PriA and the helicase loaders DnaD, DnaB, and DnaI (note that in *B. subtilis* the replicative

helicase is DnaC)[35, 36, 37, 38, 39].

153   Chromatin immunoprecipitation (ChIP) followed by quantitative PCR (qPCR) was

used to determine the enrichment of helicase and replication restart proteins surrounding a

nick in either the leading or lagging strand template. Following induction of nCas9 for one

hour in exponentially growing cells, samples were harvested for ChIP. Relative protein

enrichment proximal to the nick was normalised to a control locus located on the opposite

chromosome arm equidistant from *oriC* (+90, Fig. 2A). ChIP showed that helicase is

specifically enriched upstream of nicks located in either template strand (Fig. 2B,

Supplementary Figure 4). This upstream helicase enrichment was not detected when dCas9

was expressed, indicating that the Cas9-sgRNA nucleoprotein complex alone does not

perturb replication fork progression (Fig. 2B, Supplementary Figure 4).

163   Interestingly, significant enrichment of helicase was observed downstream of the nick

site when cleavage occurred in the template used for leading strand synthesis (Fig. 2B,

Supplementary Figure 4). The bacterial replicative helicase translocates in the $5' \rightarrow 3'$

direction along the template used for lagging strand synthesis[16]. Therefore, ChIP results

suggest that when the lagging strand template is nicked helicase runs-off the DNA substrate,

whereas when the leading strand template is nicked helicase remains associated with the

template DNA and progresses past the single-strand discontinuity (Fig. 2B). Consistent with

the observed downstream helicase enrichment (Fig. 2A-B) and replisome inactivation (Fig.

1D), reconstituted helicase assays *in vitro* have shown that when the bacterial enzyme

encounters a leading strand nick it transitions to encircling double-stranded DNA (dsDNA)

and slides along the double-helix without further unwinding activity[40].

The enrichment of helicase upstream of nick sites is consistent with either helicase

stalling or helicase reloading via the replication restart pathway. To further investigate these

models, ChIP was used to probe the presence of replication restart proteins surrounding the

nicks. Following nCas9 induction, enrichment of PriA, DnaD, DnaB and DnaI was detected

upstream (but not downstream) of the sgRNA target sequences, while these proteins were

absent from this region when dCas9 was expressed (Fig. 2C-F). These results suggest that

nicks in either the leading or lagging strand template elicit the replication restart pathway to

reload the replicative helicase, and they are consistent with models where helicase either

runs off the template (lagging strand template nick) or translocates downstream (leading

strand template nick) in an inactive state (Fig. 2G).

A weakened nCas9 system permits DNA replication restart

While replication restart proteins were enriched upstream of nick sites (Fig. 2B-F), viability

assays and MFA both indicated that downstream DNA synthesis was inhibited (Fig. 1C-D).

We speculated that persistent nCas9 cleavage activity competes with productive DNA

replication restart. Therefore, to allow interrogation of processes occurring downstream of

DNA cleavage and replisome inactivation, a weaker nCas9 system was developed. An *ssrA*

degradation tag[41] was directly fused to *cas9*^{D10A} to promote nCas9-ssrA proteolysis, and

expression of both *nCas9-ssrA* and *sgRNAs* were placed under the control of xylose-

inducible promoters (Supplementary Figure 5A). In this background, induction of nCas9 did

not significantly inhibit cell viability or alter chromosome content (Supplementary Figure 5B-

C), and MFA showed that DNA synthesis continued after a one-hour xylose induction,

consistent with low tolerable levels of nickase activity (Fig. 2H, note a reduction in read

depth upstream of nickase target sites, likely reflecting recombinational repair). This

dampened nicking system was employed to determine genes essential for replication restart.

PriA is required to restart DNA replication at single-strand DNA discontinuities

The master replication restart factor *priA* is essential for growth in *B. subtilis.* To ascertain

whether PriA-dependent replication restart is required following chromosome nicking, an

IPTG-dependent titratable PriA complementation system was developed (Supplementary

Figure 6A). Here conditions were established to express PriA-ssrA at a low level, sufficient

to maintain viability while limiting the amount of PriA activity within a cell (Supplementary

Figure 6B-C). When nCas9-ssrA and sgRNA were induced under conditions with limiting

PriA-ssrA, spot-titre analyses showed that cell growth is inhibited (Fig. 2I-J). As expected,

higher expression of PriA-ssrA rescued viability (Supplementary Figure 6D). Following

induction of nCas9 and sgRNA for one hour during exponential growth, MFA showed that

limiting levels of PriA-ssrA could not sustain chromosome synthesis downstream of the nick

site (Fig. 2K). These data indicate that PriA is essential to restart DNA replication at a single-

strand DNA discontinuity, consistent with the observed enrichment of helicase loading

proteins (Fig. 2D-F) and with the model that replisomes encountering a nick are inactivated.

A core set of recombinational repair proteins are necessary for DNA replication past a

single-strand discontinuity

Genetic and cytological data suggest that when a replisome encounters a nick, it generates

a substrate with ssDNA that is sensed by RecA (Fig. 1C-E). Evidence from several

experimental systems indicate that under these conditions a single-end DSB is formed

(seDSB)[28, 42]. In bacteria, it has been proposed that a replication-dependent DSB is repaired

through homologous recombination with the intact sister chromosome, thereby repairing the

lesion and recreating a replication fork to resume DNA synthesis[43, 44, 45, 46]. To determine the

factors required for chromosome replication following nCas9 nicking, a targeted reverse

genetic analysis was employed, focused on genes previously implicated in homologous

recombination and other DNA repair processes (Supplementary Data 1). Using this

approach, spot-titre analyses revealed that eight non-essential genes were required for

normal growth upon nick induction: DNA end-processing genes *addA*/*addB* and

recombinational repair genes *recF*/*recO*/*recR*/*recA*/*recG*/*recU* (Fig. 3A and Supplementary

Figure 7A). Functional complementation of each synthetic lethal deletion confirmed that

these gene products are necessary to support growth in cells expressing nCas9

(Supplementary Figure 7B). Interestingly, the screen revealed that several factors implicated

in DNA repair are not essential for viability under these conditions, such as those classed for

end-processing (RecQ, RecS, RecJ)[28, 47], maintaining chromosome structure ("SMC-like"

RecN, SbcCD, SbcEF)[48, 49], and recombination mediation (RecD2, RecX)[50, 51, 52]. Gene

products involved in Holliday junction migration (RuvA, RuvB)[47] displayed a small colony

phenotype but not a loss of CFUs (Supplementary Figure 8D), suggesting they play a

notable yet less critical role in the repair pathway.

Focusing on the eight mutants that produced the most severe growth defects, MFA

was used to analyse DNA replication following induction of nCas9 and sgRNA in

exponentially growing cells (Supplementary Figure 9, note several of the deletion mutants

display altered replication profiles near the origin or terminus, consistent with previous

studies)[53, 54, 55]. The results showed that DNA end-processing (*addA*/*addB*) and

recombination mutants (*recF*/*recO*/*recR*/*recA*/*recG*) are defective in chromosome replication

downstream of the nick site, with the only exception being *recU* (Fig. 3B, Supplementary

Figures 10, 11, note MFA profiles reflect distinct mutant backgrounds). Together, these

observations support the recombinational repair model for rescuing a collapsed replication

fork[56]. Moreover, the requirement for the AddAB helicase/nuclease complex, which engages

linear DNA substrates at a nearly blunt end[57], strongly suggests that bacterial replisomes

encountering a nick generate a seDSB[58, 59].

Based on the proposed activities of these proteins, the following model was

developed (Fig. 3C). AddAB helicase/nuclease processes a broken replication fork at a

seDSB to generate a single-stranded substrate with a 3′-end required for recombination (Fig.

3C(I-III))[23, 58, 59]. The single-stranded DNA (ssDNA) is recognised and bound by the single-

stranded binding protein (SSB), which must be removed by the RecFOR system to allow

RecA filament formation (Fig. 3C(IV-V))[23, 60, 61, 62]. RecA is the master bacterial recombinase

that binds to ssDNA with a free 3′-end, identifies homologous DNA on the intact sister

chromosome, and performs strand invasion and exchange (Fig. 3C(V-VI))[63]. RecG is a

helicase that remodels branched DNA substrates, likely acting at the Holliday junction or at

the replication fork (these are not mutually exclusive) (Fig. 3C(VII))[64, 65]. RecU is a structure

specific nuclease involved in cleaving the Holliday junction generated by RecA (Fig.

3C(VII))[66].

AddAB helicase activity is necessary for recombinational repair at a nick

The genetic analysis above indicated that the AddAB helicase/nuclease was responsible of

processing the seDSB (Fig. 3). While both AddA and AddB subunits harbour nuclease

activity, the two enzymes cleave distinct strands of the DNA duplex following unwinding by

the AddA helicase (Fig. 4A)[59]. However, whether all AddAB activities are required for

recombinational repair in live cells remained unclear.

To determine whether AddAB was acting directly on broken replication forks

generated by replisome inactivation at a nick, enrichment of AddAB was investigated using

ChIP. AddA and AddB were independently fused to the fluorescent protein mNeonGreen

(mNG), which here was used as an epitope tag. The results showed that mNG-AddA and

mNG-AddB are maximally enriched 2 kb upstream of nicks, consistent with a direct role in

end-processing under these conditions (Fig. 4B). The enzymatic activities of AddA and AddB

were then reduced through substitution of catalytic residues (Fig. 4A)[59]. Spot-titre analyses

showed that the helicase defective AddA^{K36A} (AddA^{ΔHEL}) is unable to sustain bacterial growth

upon induction of nicks, whereas nuclease defective variants of either AddA^{D1172A} (AddA^{ΔNUC})

or AddB^{D961A} (AddB^{ΔNUC}) grew comparably to a strain encoding wild-type AddAB (Fig. 4C).

MFA confirmed that DNA synthesis is blocked downstream of a nick in the strain expressing

AddA$^{\Delta HEL}$, whereas the strains expressing AddA$^{\Delta NUC}$, AddB$^{\Delta NUC}$ or AddA$^{\Delta NUC}$AddB$^{\Delta NUC}$ were

comparable to wild-type (Fig. 4D, Supplementary Figure 12A-B). An immunoblot showed

that the AddA$^{\Delta HEL}$ variant is stably expressed (Supplementary Figure 12C), indicating that

AddAB helicase activity alone is necessary for the enzyme to process the seDSB.

Since AddAB nuclease activity was dispensable for recombinational repair, we

wondered whether another nuclease might fulfil this role. RecJ is a single-strand DNA-

specific exonuclease that has been observed to co-localise with the replisome in *B.*

subtilis[67]. Deletion of *recJ* in conjunction with either *addA$^{\Delta NUC}$*, *addB$^{\Delta NUC}$*, or

addA$^{\Delta NUC}$addB$^{\Delta NUC}$ did not reduce the number of CFU following nicking (Supplementary

Figure 12D), indicating that RecJ nuclease activity is not required. Based on these

observations, it appears that under these conditions AddAB helicase activity and an

unspecified nuclease promote end-processing of the seDSB to generate a 3′-tail.

SSB recruitment requires AddAB helicase activity

The requirement of the RecFOR system for recombinational repair (Fig. 2) implied that SSB

is bound to ssDNA at the seDSB. To interrogate this hypothesis, recruitment of SSB was

probed using ChIP following induction of nCas9 and sgRNA in exponentially growing cells.

The results showed that SSB is enriched upstream of a leading or a lagging strand nick (Fig.

5A), and that this enrichment was dependent upon AddAB (Fig. 5B). Further analysis of

AddAB showed that only helicase activity was required to promote SSB enrichment

upstream of a nick (Fig. 5C). Together these results support the role of AddAB to unwind a

seDSB and provide a ssDNA substrate for SSB.

AddAB helicase/nuclease activity is thought to be modulated by interaction of the

enzyme with ssDNA binding motifs termed *chi* sites scattered throughout the genome[59, 68, 69].

The locations of *B. subtilis chi* sites are shown below the qPCR primer pairs used to probe

SSB enrichment (Fig. 5A). The cluster of *chi* sites located 1-2 kb upstream of the nick site

correlates well with the observed peak enrichment of SSB in nicking strains (Fig. 5A). To

further investigate the connection between *chi* site location and SSB enrichment, the

endogenous *addA/addB* genes were deleted and functionally complemented by the

homologous genes from *Staphylococcus aureus* (*rexA/rexB*, Supplementary Figure 12E).

While these enzymes are related, *B. subtilis* and *S. aureus chi* sequences are distinct (*chi^Bs*

is 5′-AGCGG-3′ and *chi^Sa* is 5′-GAAGCGG-3′)[59, 70, 71] and accordingly the locations of the

respective *chi* sites upstream of the targeted nick at -90° are different (Fig. 5A). Revaluation

of SSB enrichment by ChIP in the strain expressing RexAB showed that the SSB profile is

shifted upstream to a position proximal to the first *chi^Sa* sites (Fig. 5A). Taken together, the

data suggest that end-processing is necessary to deposit SSB onto ssDNA generated from

a seDSB, and that AddAB nuclease activity can be abolished without compromising

recombinational repair.

The SSB C-terminal tail is required for recombinational repair at a nick

SSB forms a homotetramer that binds ssDNA non-specifically via its N-terminal domain[72].

Each SSB monomer harbours a long disordered C-terminal domain that contains a protein

interaction sequence at the terminus (the C-terminal tail or CTT) (Fig. 6A-B, Supplementary

Figure 13A)[73]. While the *B. subtilis* SSB-CTT is dispensable for cell viability, it is required for

efficient repair of DNA lesions created by either UV irradiation or mitomycin C treatment[49, 74].

To determine whether the SSB-CTT is required for recombinational repair following

replisome inactivation at a nick, an IPTG-inducible SSB complementation system was

generated in a Δ*ssbA* strain, allowing exogenous expression of either the wild-type protein or

a variant lacking the last six amino acids (SSB$^{\Delta CTT}$, Supplementary Figure 13B). Spot-titre

analyses confirmed that SSB$^{\Delta CTT}$ can complement a Δ*ssbA* mutant similar to wild-type SSB

and that viability was dependent upon IPTG (Supplementary Figure 13C), while

immunoblots showed that the inducible SSB proteins were being expressed at levels

comparable to the endogenous protein (Supplementary Figure 13D).

The SSB complementation system was then moved into strains capable of

expressing nCas9 and sgRNA (Supplementary Figure 13E). Spot-titre analyses showed that

wild-type SSB supports viability following induction of nicking, whereas the SSB$^{\Delta CTT}$ variant

could not (Fig. 6C). MFA confirmed that chromosome replication from the nick site to the

terminus is arrested when SSB lacks its C-terminal tail (Fig. 6D). Therefore, the results

indicate that the SSB-CTT is necessary for DNA repair following replisome inactivation at a

nick.

The SSB-CTT recruits RecO during recombinational repair

The SSB-CTT acts as a protein interaction hub[49] that directly interacts with several factors

shown above to be necessary for recombinational repair, including RecO, RecG, and PriA

(Figs. 2I-K, 3A-B, Supplementary Figure 13A). ChIP showed that the SSB$^{\Delta CTT}$ variant was

enriched upstream of a nick at similar levels to the wild-type protein (Fig. 6E), indicating that

the growth defect of the SSB$^{\Delta CTT}$ variant following nicking was not due to a general loss of

SSB activity. To investigate the essential role of the SSB-CTT, ChIP was employed to probe

the recruitment of His-RecO upstream of nick sites. Enrichment of His-RecO was abolished

in cells harbouring SSB$^{\Delta CTT}$ (Fig. 6E), indicating that the SSB:RecO interaction depends on

the SSB-CTT (Supplementary Figure 13F).

The RecFOR system is thought to promote RecA binding ssDNA by stimulating the

dissociation of SSB[23, 75]. ChIP confirmed that enrichment of His-RecA was dependent upon

both AddAB and RecFOR systems (Fig. 6F). Critically, enrichment of His-RecA was also

dependent upon the SSB-CTT (Fig. 6E). Taken together, these results are consistent with

the model that AddAB processes a seDSB to generate ssDNA for SSB, followed by RecFOR

mediating displacement of SSB to allow RecA loading on ssDNA[23, 75, 76, 77]. Further, they

suggest that an essential role of the SSB-CTT is to recruit RecO for this process.

RecG is recruited to sites of recombinational repair

ChIP enrichment of RecA was independent of RecG (Fig. 6F), suggesting that RecG acts

downstream of RecA-mediated strand exchange, consistent with proposed activities in DNA

remodelling. RecG has been implicated in a myriad of DNA transactions including reversal of

replication forks, migration of Holliday junctions, regulation of RecA strand exchange,

processing of flaps generated when DNA replication forks converge, and stabilisation of D-

loops[65, 78, 79, 80, 81]. To determine whether RecG acts directly during recombinational repair,

the endogenous *recG* allele was fused to a FLAG epitope tag and recruitment of FLAG-

RecG was probed using ChIP following induction of nCas9 and sgRNA in exponentially

growing cells. The results showed that FLAG-RecG is maximally enriched ~3 kb upstream of

a nick in either leading or lagging strand (Fig. 7A), consistent with RecG playing a direct role

in DNA remodelling necessary to resume DNA replication (Fig. 3B)[49].

RecU is recruited to sites of recombinational repair

Of the recombination proteins found to be important for supporting growth following nick

induction, the Holliday junction resolvase RecU was conspicuous[28]. First, deletion of *recU*

impaired growth less than the other mutants (Fig. 3A). Second, MFA indicated that DNA

replication past a nick was independent of *recU* (Fig. 3B, Supplementary Figure 11B).

Therefore, we considered the possibility that growth defects associated with Δ*recU* following

chromosome nicking were indirect, as the deletion mutant is known to elicit diverse

chromosome segregation abnormalities (Fig. 7B, Supplementary Figure 14)[82]. To address

this hypothesis, the endogenous *recU* allele was fused to a FLAG epitope tag and

enrichment of RecU-FLAG was determined using ChIP. The results show that RecU-FLAG

was specifically recruited to DNA upstream of a nick site (Fig. 7E), consistent with a direct

role in Holliday junction resolution. However, due to the pleiotropic phenotype of a Δ*recU*

mutant, its precise importance during recombinational repair remains unclear.

DISCUSSION

In this study we have characterised the fate of the bacterial replisome *in vivo* after it encounters a single-strand discontinuity in template DNA (Fig. 1A). The results indicate that a nick in the template for either the leading or lagging strand arrests DNA synthesis, likely generates a seDSB that elicits repair by homologous recombination, followed by PriA-dependent replication restart (Figs. 1-3). These findings are consistent with the observation that a nick in a replicating bacteriophage chromosome generates DSBs[5] and with recent reports of the impact of nicks in eukaryotic replication systems[14, 15, 22]. Importantly, while replisomes are thought to be generally tolerant to many types of DNA damage[83], it appears that the bacterial replication machinery cannot readily bypass single-strand interruptions as these lesions affect helicase progression (Figs. 2, 8(I)).

Replication beyond a nick requires recombinational repair (Fig. 3). Genetic analysis combined with ChIP indicates that a single pathway is utilised under these experimental conditions: AddAB unwinds the DNA (Fig. 8(II)); SSB binds the ssDNA produced by AddAB (Fig. 8(III)); RecFOR counteracts SSB and allows RecA to bind ssDNA (Fig. 8(IV-V)); RecA promotes strand invasion of the homologous chromosome and generates a D-loop (Fig. 8(VI)); RecG remodels a DNA substrate formed during recombination (Fig. 8(VII)); PriA binds to the repaired replication fork (Fig. 8(VIII)) and reloads helicase to promote replisome assembly and DNA replication restart (Fig. 8(IX)). The observed lack of redundancy between proteins involved in recombinational DNA repair suggests that the pathway identified here is the major route for fixing broken replication forks in *B. subtilis*[48, 84].

Building upon these findings, we investigated specific protein activities to help elucidate their roles during recombinational repair of a replisome inactivated at a single-strand discontinuity. Below we discuss these findings in what we propose is chronological order.

The replicative helicase is inactivated after it encounters a nick

In bacteria the replicative helicase encircles and tracks along the template for lagging strand

synthesis. ChIP data suggests that there are distinct fates of the helicase depending on the

location of a nick (Fig. 2A-B, Supplementary Figure 4). As expected for a nick in the lagging

strand template, helicase enrichment was only observed at the cleavage site and upstream,

consistent with the model that helicase runs-off the template when it encounters the single-

strand discontinuity (Figs. 2G, 8(I)). In contrast, when the nick was in the leading strand

template, helicase was also enriched downstream of the cleavage site, suggesting that the

enzyme continued translocation. We propose that upon encountering a nick in the leading

strand template, helicase slides onto the dsDNA template in an inactive conformation that

cannot perform further DNA unwinding (Figs. 2G, 8(I))[40]. This model is consistent with *in*

vitro experiments using *Thermus aquaticus* helicase, where after unwinding up to a leading

strand nick, the enzyme can continue to translocate with the two strands of DNA passing

through the central channel with no resultant unwinding[40]. Whether the bacterial helicase is

actively removed from dsDNA under these conditions, akin to what has been observed for

the eukaryotic CMG helicase in *Xenopus* extracts[14], requires further investigation.

AddAB helicase activity is necessary and sufficient for recombinational repair

AddAB and related enzymes have been shown to bind blunt end dsDNA or dsDNA with

short overhangs (4-20 nucleotides with either a 5′- or a 3′-end)[59, 85]. Moreover, AddAB is inert

on DNA substrates lacking a blunt end[57]. The requirement for AddAB to efficiently process

DNA past a nick (Fig. 3), specifically AddAB helicase activity to recruit SSB (Figs. 4C-D, 5),

is consistent with the model that a seDSB is generated when a bacterial replisome

encounters a single-strand discontinuity.

When the template for leading strand synthesis is nicked, the DNA polymerase likely

runs-off the end of the substrate to produce a blunt or nearly blunt end, suitable for

recognition by AddAB. However, when the template for lagging strand synthesis is nicked, a

substrate with a 3′-tail would likely be produced. Because the reverse genetic screen

showed that no other major nucleases are required for this recombinational repair pathway

(Fig. 3, Supplementary Figures 8, 12D and Supplementary Data 1), we speculate that the

high local concentration of replication factors could facilitate priming near the end of the 3′-

tail, thereby filling the ssDNA with an Okazaki fragment to create a suitable substrate for

AddAB (Fig. 8(I-II)). Alternatively, unidentified nucleases may degrade the ssDNA tail[86].

The master recombination factor RecA requires ssDNA to promote homologous

pairing. A long-standing question is how this ssDNA is generated *in vivo*. Most current

models suggest that enzymes resect a DSB to generate a single-strand terminating with a

3′-end[28, 34]. Here we tested this model by mutating the nuclease activities of AddA and AddB.

The results showed that neither nuclease activity is required for recombinational repair under

the experimental conditions tested (Fig. 4C-D). Moreover, deletion of the major exonuclease

RecJ in a strain expressing the nuclease-dead AddAB variant did not display a synthetic

phenotype (Supplementary Figure 12D). In contrast, AddAB helicase activity was required

for recombinational repair (Fig. 4D), consistent with a previous study of the ancestral AdnAB

resection machinery in *Mycobacteria tuberculosis*[87] and supported by c*hi*-dependence to

facilitate SSB loading (Fig. 5A, Supplementary Figure 12E). The local concentration of SSB

is high near the site of DNA replication[88, 89], which together with phase separation[90, 91] may

facilitate its rapid association with ssDNA emerging from AddAB. In fact, *in vitro* the

AddA^{NUC}AddB^{NUC} nuclease-dead enzyme does not unwind dsDNA in the absence of SSB[92],

suggesting that AddAB unwinding and SSB binding ssDNA may be coupled. Together these

results raise the unexpected possibility that during recombinational repair at a nick, the key

activity during end-processing is to separate the strands and create a ssDNA substrate for

SSB (Figs. 5C, 8(II-III)).

Interaction of the SSB C-terminal tail with RecO is necessary for recombinational

repair

The SSB-CTT is an established protein interaction hub that has been shown to directly bind

several of the essential replication and repair factors required at nicks, including RecO[93, 94,]

[95, 96], RecG[97, 98], and PriA[74, 99] (Supplementary Figure 13A). However, the critical activities

performed by the SSB-CTT *in vivo* had not been established[49]. Based on the results above,

we conclude that the SSB-CTT interaction with RecO is necessary for recombinational repair

(Fig. 6). RecO is part of the RecFOR system required for RecA loading onto ssDNA *in vitro*[76]

and RecA-GFP foci formation *in vivo*[23]. Because RecO activity precedes the activities of

RecG and PriA (i.e. RecA binding ssDNA is blocked, Fig. 6F), further conclusions regarding

the importance of these protein:protein interactions cannot be drawn.

Historically the RecFOR system has been proposed to mediate ssDNA gap-filling,

rather than recombination at a DSB[61]. While it is possible that multiple types of DNA lesions

are associated with nicks generated by nCas9 in *B. subtilis*, we note that recruitment of SSB

and RecA was dependent on AddAB (Figs. 5B, 6F). Because AddAB-dependent end-

processing is not associated with models for recombinational repair of ssDNA gaps, the data

presented here is most consistent with RecFOR acting within the proposed seDSB repair

pathway (Fig. 8). Supporting this viewpoint, AddAB has been shown to be epistatic to

RecFOR in *Thermus thermophilus*[100] and RecFOR is essential for DSB repair in

Deinococcus radiodurans[101].

In vitro the RecO:RecR complex facilitates loading of RecA onto SSB-coated

ssDNA[102, 103]. Because RecO, RecR, and RecF were all required for recombinational repair

following nicking (Fig. 3), we propose that the essential role of RecF here is to recruit RecR

and generate the RecO:RecR complex. This model is consistent with a hand-off mechanism

for RecR observed *in vitro*[102]. Since SSB and RecO are enriched at a broken replication fork

(Fig. 6E), we predict that there is a mechanism (potentially facilitated via RecF) to regulate

RecO:RecR complex formation until it is required, thus safeguarding the SSB-coated DNA

from promiscuous RecA loading (Figs. 5B, 6F, 8(III-V)).

RecG is necessary to complete recombinational repair prior to replication restart

The biological role of the DNA translocase RecG has been a long-standing question[64, 65].

MFA shows that a *ΔrecG* mutant cannot efficiently replicate past a nick (Fig. 3B) and ChIP

indicates that RecG is enriched upstream of the lesion (Fig. 7A)[54]. We propose that RecG

acts to remodel a DNA substrate formed after RecA-mediated strand invasion, either the

Holliday junction or the replication fork (Fig. 8(VI-VII)). Branch migration of the Holliday

junction away from the replication fork could be required for downstream steps necessary for

PriA-dependent helicase loading[104]. Alternatively, RecG may process the replication fork

itself, preparing it for PriA loading on the correct substrate (Supplementary Figure 15)[54, 55].

Holliday junction migration and resolution are not necessary for replication restart

While *recU* was identified as important in our genetic screen following nicking (Fig. 3A) and

RecU was specifically enriched upstream of a nick (Fig. 7C), it was found that RecU is not

necessary for replication restart *per se* (Fig. 3B). Single cell analyses revealed significant

chromosome segregation defects in all *ΔrecU* mutants (Fig. 7B, Supplementary Figure 14)[82],

likely explaining the observed loss in viability (Fig. 3A, Supplementary Figure 7A). Together

these results support the model that recombinational repair is mechanistically distinct from

resolution of conjoined chromosomes[56].

Potentiating ROS-mediated antibiotic toxicity by perturbing DNA repair

In this work, the nicks generated by Cas9 enzymes were used as proxies for single-strand

discontinuities formed during removal of damaged DNA from a bacterial genome. We find

that when a bacterial replication fork encounters a single strand discontinuity it becomes

inactivated. By identifying the essential factors and activities required to repair the DSB

formed after a replisome encounters a nick, we have provided a set of molecular targets

whose disruption could potentiate the activity of antimicrobial compounds that damage DNA.

METHODS

Experimental model and growth

The following bacterial organisms were used: *E. coli* and *B. subtilis.* Unless otherwise stated

in method details, strains were grown at 37°C using nutrient agar (NA) or lysogeny broth

(LB: 10 g/L tryptone, 10 g/L NaCl, 5 g/L yeast extract) for routine selection and maintenance

of bacterial strains. Supplements were added as required: ampicillin (100 μg/ml),

erythromycin (1 μg/ml) in conjunction with lincomycin (25 μg/ml), chloramphenicol (5 μg/ml),

kanamycin (5 μg/ml), spectinomycin (50 μg/ml), zeocin (10 μg/ml), xylose (1% w/v) and

IPTG (0.05 mM, 0.1 mM or 1 mM).

Bacterial strains

Bacterial strains used in this study are listed in Supplementary Data 2.

Oligonucleotides

Oligonucleotides were purchased from Eurogentec and Sigma to assemble recombinant

DNA (sequences are listed in Supplementary Data 3) or perform quantitative PCR

(Supplementary Data 4). Design of sgRNA targets was performed using the online portal

CRISPick with default parameters[105] and recombinant DNA assembly was achieved via *in*

vitro mutagenesis (see below).

E. coli plasmid construction

E. coli transformations for constructs harbouring *priA* were performed in CW198 via heat-

shock following the Hanahan method[106] and propagated in LB with appropriate antibiotics at

37°C. DH5α [*F- Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17(rk-, mk+) phoA*

supE44 thi-1 gyrA96 relA1 λ-] was used for other plasmids construction. Plasmids were

purified using QIAprep spin miniprep kits and recombinant DNA confirmed by sequencing

(Supplementary Data 3). Descriptions, where necessary, are provided below*.*

pCW367, pCW368, pCW478, pCW479, pCW510, pCW717, pCW734, pCW736 and

pCW850 were generated by Quickchange mutagenesis using mutagenic primer pairs

containing an overlap of 10-15 bp. PCR products were treated with DpnI for at least 3 hours

to digest template DNA and 10 µl were used for *E. coli* transformation.

pCW20, pCW300, pCW301, pCW327, pCW460, pCW471, pCW472, pCW575,

pCW734, pCW736 and pCW766 were generated by Gibson assembly using the NEB Hi-Fi

cloning kit according to manufacturer instructions following purification of individual PCR

fragments using QIAquick PCR purification kits.

Linear recombinant DNA assembly

For recombinant DNA assembly without *E. coli* propagation, pCW573, pCW574, pCW577,

pCW578, pCW605, pCW640, pCW672, pCW702, pCW720, pCW732, pCW758, pCW759,

pCW760, pCW777, pCW779, pCW781, pCW783, pCW785, pCW802, pCW803, pCW804,

pCW809, pCW834, pCW848, pCW854, pCW855, pCW857 and pCW859 were assembled

linearly using a multistep *in vitro* assembly process followed by nested PCR amplification

prior to *B. subtilis* transformation. Briefly, individual PCR fragments corresponding to

homology regions and insert DNA were amplified, purified and DNA ends ligated using the

NEB Hi-Fi cloning kit. Raw Gibson assembly products were then amplified using nested

primer pairs annealing towards the end of homology regions and PCR products were used

for *B. subtilis* transformation. Following isolation of *B. subtilis* colonies resistant to the

appropriate antibiotic markers, genomic DNA was extracted using the DNeasy blood and

tissue kit, regions corresponding to recombinant DNA were amplified via PCR using

oligonucleotides annealing outside homology fragments to ensure double crossover had

occurred and results were confirmed via PCR product sequencing. Note that CW2045 and

derivatives harbour the additional *addA^{D682E}* mutation necessary to propagate *addB^{D961A}*-

addA^{D1172A} in live cells, which is surface exposed and structurally unlikely to impact AddAB

end-processing activities (Supplementary Figure 12F).

B. subtilis strain construction

Transformation of competent *B. subtilis* cells was performed using an optimized two-step

starvation procedure[107]. Briefly, recipient strains were grown overnight at 37°C in

transformation medium (Spizizen salts: 0.2% w/v ammonium sulphate, 1.4% w/v

dipotassium phosphate, 0.6% w/v potassium phosphate, 0.1% w/v tri-sodium citrate, 0.02%

w/v magnesium sulphate supplemented with 1 µg/ml ammonium iron(III) citrate, 6 mM

MgSO$_4$, 0.5% w/v glucose, 0.02 mg/ml tryptophan and 0.02% w/v casein peptone)

supplemented with IPTG where required. Overnight cultures were diluted 1:20 into fresh

transformation medium supplemented with IPTG where required and grown at 37°C for 2.5-3

hours with continual shaking. An equal volume of prewarmed starvation medium (Spizizen

salts supplemented with 6 mM MgSO$_4$ and 0.5% w/v glucose) was added and the culture

was incubated at 37°C for two hours with continual shaking. DNA was added to 350 µl cells

and the mixture was incubated at 37°C for one hour with continual shaking. 20-200 µl of

each transformation was plated onto selective media supplemented with IPTG where

required and incubated at 37°C for 24-48 hours. Results were validated by propagating cells

on relevant antibiotic markers and sequencing. Sanger sequencing identified that the strong

nCas9/sgRNA nicking system is prone to accumulation of suppressor mutations either

disrupting protospacer or its promoter sequence (Supplementary Figure 1F), therefore it is

recommended to rebuild these strains periodically to validate their identity.

For strain construction of essential gene knockouts (e.g., *ssbA* and *priA*), an ectopic

copy of the gene of interest was placed under the control of an IPTG-inducible promoter and

recombinant products were transformed in wild-type *B. subtilis*. Note that *priA* was C-

terminally fused to a degradation tag (*priA-ssrA*) to promote proteolysis and reduce

background expression in the absence of IPTG. Cells harbouring an ectopic expression

cassette were transformed with recombinant DNA to knockout the corresponding

endogenous gene copy and colonies were obtained in the presence of IPTG to avoid

accumulation of suppressor mutations. Thereafter, these strains were always propagated in

the presence of IPTG to enable genetic complementation via the inducible essential gene

copy.

For the construction of protein fusions, *ssrA* tags (AANDENYSENYALAA) were

directly fused to C-terminal ends of target proteins (e.g. Cas9-ssrA variants and PriA-ssrA),

whereas the fluorescent mNG-RecA DNA break reporter was constructed by N-terminal

fusion via a short serine/glycine linker (SG$_3$).

Spot-titre assays

Cells were grown in LB overnight in the presence of 0.5% w/v glucose. Overnight cultures

were diluted in fresh LB medium using 10-fold serial dilutions and 5 µL aliquots were spotted

onto NA plates with or without xylose. NA plates were supplemented with IPTG when

required as indicated on individual figure panels. All plates were incubated for 24 hours or up

to 48 hours when indicated. Experiments were performed independently at least three times

and representative data are shown.

Antibody purification

SSB anti-serum was used as delivered from Eurogentec. Other primary antibodies received

from Eurogentec (anti-DnaI, anti-DnaC, anti-PriA) were affinity purified prior to use. Anti-sera

were diluted 1:1 with PBS and applied to respective antibody/NHS-activated agarose

columns. Purified antibodies were eluted with glycine, dialysed with PBS and concentrated.

Affinity purified antibodies were stored at -20°C in 1X PBS with 50 % v/v glycerol. The

antibody validation report is available as a Source Data file.

Immunoblot analyses

Cultures were grown overnight in LB and diluted 1:100 in fresh medium the next morning.

Diluted cells were grown to an absorbance at 600 nm (A$_{600}$) of 0.2-0.3, induced with 1% w/v

xylose or 1 mM IPTG where required and further incubated for one hour. Cells were

harvested using centrifugation, resuspended in 1x PBS supplemented with one tenth of a

peptidase inhibitor tablet and sonicated (40 amp) twice for 12 seconds with 3 second pulses

at 4°C. Samples were adjusted to 1X LDS buffer and 125 mM DTT, fixed for 5 minutes at

95°C and centrifuged for 2 minutes prior to loading the supernatant on polyacrylamide gels.

Proteins were separated by electrophoresis using NuPAGE 4-12% Bis-Tris gradient gels run

in MES SDS buffer and transferred to methanol-activated Hybond-P PVDF membranes

using Wypall X60 cloths for blotting in a semi-dry apparatus (Bio-rad Trans-Blot Turbo,

transfer buffer: 300 mM Tris-HCl, 300 mM glycine, 140 mM tricine, 0.05% w/v SDS, 2.5 mM

EDTA). Membranes were washed twice using PBST (1X PBS supplemented with 0.1% v/v

Tween-20) prior to blocking for 90 minutes at room temperature (blocking buffer: PBST

supplemented with 7.5% w/v milk). Individual polypeptides were probed using the following

primary antibodies diluted in blocking buffer following overnight incubation at 4°C: anti-Cas9

(1:1000, Merck SAB4200701), anti-FtsZ[108] (1:5000), anti-PriA (1:1000, Eurogentec), anti-

SSB (1:1000, Eurogentec) and anti-mNeonGreen (1:1000, ProteinTech 32F6). Detection

was performed with anti-rabbit (1:10000 for PriA/SSB, ProteinTech SA00001-2), anti-sheep

(1:5000 for FtsZ, Merck A3415) or anti-mouse (1:5000 for Cas9/mNeonGreen, ProteinTech

SA00001-1-A) horseradish peroxidase-linked secondary antibodies in combination with the

Clarity ECL substrate using an ImageQuant LAS 4000 mini digital imaging system. Detection

of Cas9, FtsZ, PriA, SSB and mNeonGreen was within a linear range. Experiments were

independently performed at least twice and representative data are shown.

Whole genome sequencing

Cells were grown in rich chemically defined medium (RCDM: Spizizen salts supplemented

with 1 μg/ml ammonium iron(III) citrate, 0.1 mM $CaCl_2$, 0.13 mM $MnSO_4$, 6 mM $MgSO_4$,

0.5% w/v glucose, 0.1% w/v glutamate, 0.02 mg/ml tryptophan and 0.02% w/v casein

peptone) overnight and diluted 1:100 in fresh RCDM the next morning. Diluted cells were

grown to an absorbance at 600 nm (A_{600}) of 0.2-0.3, induced with 1% w/v xylose and further

incubated for one hour. Growth was arrested by adding 0.05% w/v sodium azide to cultures

followed by vigorous mixing, cells were collected by centrifugation, the supernatant

discarded, and pellets were flash frozen in liquid nitrogen before genomic DNA extraction

using the DNeasy blood and tissue kit and a final elution volume of 50 µl in ultrapure water.

Individual gDNA libraries were prepared using the DNA Prep tagmentation kit and pre-paired

i5/i7 indices prior to pooling and DNA quantification with the 1X dsDNA high sensitivity kit

using a Qubit instrument. Pooled libraries were diluted to 100 pM in resuspension buffer and

20 µl were loaded onto iSeq 100 i1 Reagent v2 sequencing chips for whole-genome

sequencing using an iSeq100 sequencer.

Following whole-genome sequencing and demultiplexing embedded within the

iSeq100 indexed sequencing workflow, .FASTQ files were extracted and loaded onto the

CLC Genomics Workbench (version 23) for initial data processing. Paired .FASTQ files were

mapped onto the *B. subtilis* Δ6 reference genome using the 'Map Reads to Reference'

function and a .BAM file was created from mapped sequencing reads for each sequenced

genome. An in-house script written in R (version 4.5) using the Rsamtools library was

employed to deconvolute .BAM files to validate even read distribution for sense/antisense

strands prior to calculating density coverage and extracting cumulated densities per

chromosome position.

Fluorescence microscopy

Strains were grown overnight at 37°C in RCDM and the following day, cultures were diluted

1:100 into fresh imaging medium (RCDM where glucose is replaced by 0.5% v/v glycerol) in

the presence of 0.1 mM IPTG for induction of mNeonGreen-RecA expression where

required. Cells were allowed to grow to an A_{600} of 0.3, induced with 1% w/v xylose and

further incubated for 90 minutes. DAPI (5 µg/ml) and Nile red (1 µg/ml) stains were used to

visualise the nucleoid and cell membrane following a 10 minute incubation on a benchtop

incubator prior to imaging.

Cells were mounted on ~1.25% agar pads (in sterile ultrapure water) and a 0.13- to

0.17-mm glass coverslip was placed on top. Microscopy was performed on an inverted

epifluorescence microscope (Nikon Ti) fitted with a Plan Apochromat Objective 100x/1.40

NA Oil Ph3. Light was transmitted from a CoolLED pE-300 white light source through a

Sutter Instruments liquid light guide and images were collected using a Photometrics Prime

camera. Chroma fluorescence filter sets were used with 100 ms GFP, 250 ms DAPI and 50

ms mCherry exposure times at 100% LED power. Digital images were acquired using NIS

Elements (version 5) and analysed via the Fiji software (version 1.53)[109]. Quantification of

mNeonGreen-RecA features was manually curated from the count of 100 individual cells. All

experiments were independently performed at least twice and representative data are

shown.

Chromatin immunoprecipitation

Chromatin immunoprecipitation (ChIP) was performed by following an established protocol

with minor modifications[35]. Strains were grown overnight at 37°C in RCDM and the following

day, cultures were diluted 1:100 into fresh medium and allowed to grow to an A_{600} of 0.2-0.3.

Cells were induced with 1% w/v xylose and further incubated for one hour prior to ChIP.

Samples were adjusted to 1X PBS and crosslinked with 1% v/v formaldehyde for 10 minutes

on a roller at room temperature, then quenched with 0.1 M glycine. Cells were pelleted at

4°C, washed three times with ice-cold 1X PBS (pH 7.3), frozen in liquid nitrogen and stored

at -80°C. Frozen cell pellets were resuspended in 500 µl of lysis buffer (50 mM NaCl, 10 mM

Tris-HCl pH 8.0, 20% w/v sucrose, 10 mM EDTA, 100 µg/ml RNase A, one quarter of a

peptidase inhibitor tablet, 4 mg/ml lysozyme) and incubated at 37°C for 30 min to degrade

the cell wall. Protoplasts were supplemented with 500 µl of immunoprecipitation buffer (300

mM NaCl, 100 mM Tris-HCl pH 7.0, 2% v/v Triton X-100, one quarter of a peptidase inhibitor

tablet) to lyse cells and the mixture was incubated at 37°C for a further 10 minutes before

cooling on ice for 5 minutes. Lysis and immunoprecipitation buffer volumes were multiplied

by the number of antibodies to probe per sample. DNA samples were sonicated (40 amp)

three times for 2 minutes at 4°C to obtain an average fragment size of 500 to 1000 base

pairs. Cell debris were removed by centrifugation at 4°C and the supernatant transferred to a

fresh Eppendorf tube. To determine the relative amount of DNA immunoprecipitated

compared to the total amount of DNA, 100 µl of supernatant was removed, treated with 0.5

mg/ml pronase at 37°C for four hours then stored on ice. To immunoprecipate protein-DNA

complexes, 800 µl of the remaining supernatant was incubated with individual antibodies (2

µl anti-SSB, 2 µl anti-PriA, 2 µl anti-DnaD[36], 2 µl anti-DnaB[36], 2 µl anti-DnaI, 2 µl anti-DnaC,

2 µl anti-mNeonGreen (ProteinTech 32F6), 4 µl anti-His (Qiagen 34660) or 25 µl anti-FLAG

M2 beads (Merck M8823) for 90 minutes at room temperature (or 120 min for FLAG-tagged

strains). 750 µg of protein G Dynabeads (Invitrogen 10009D) were equilibrated by washing

with bead buffer (1X PBS, 0.01% v/v Tween 20), resuspended in 25 µl of bead buffer and

incubated with the sample supernatant for 1 hour at room temperature. Immunoprecipitated

complexes were collected by applying the mixture to a magnet and washed with the

following buffers in the respective order: 0.5X immunoprecipitation buffer for 15 min, 0.5X

immunoprecipitation buffer supplemented with 500 mM NaCl for 15 min, stringent wash

buffer (250 mM LiCl, 10 mM Tris-HCl pH 8.0, 0.5% v/v Igepal, 0.5% w/v sodium

deoxycholate, 10 mM EDTA) for 20 min. Finally, protein-DNA complexes were washed a

further three times with TET buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 0.01% v/v Tween

20) and resuspended in 100 µl of TEN buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 300 mM

NaCl). Formaldehyde crosslinks of both the immunoprecipitate and total DNA were reversed

by incubation at 65°C for 18 hours in the presence of 1 mg/ml proteinase K. DNA was then

removed from the magnetic beads, cleaned using QIAquick PCR purification columns and

used for quantitative PCR analyses.

Quantitative PCR

Quantitative PCR (qPCR) was performed using the Luna qPCR master mix to measure

relative amounts of DNA bound by SSB, PriA, DnaD, DnaB, DnaI, DnaC, mNeonGreen-

AddA, mNeonGreen-AddB, His-RecO, His-RecA, FLAG-RecG and RecU-FLAG at specific

genomic locations (e.g. near the nick site located at -90° on the chromosome or the origin

oriC compared to a control site located at +90° on the chromosome). All PCR reactions were

assembled using the QIAgility robotic workstation in 20 µl reaction volumes in a Rotor-Disc

100 and qPCRs were run on a Rotor-Gene Q instruments. Standard curves were obtained

using the Rotor-Gene Q software (version 2) to calculate the efficiency of each primer pair,

which varied ~5% between sets. Oligonucleotides were designed to amplify specific

genomic regions using the Primer3Plus tool[110], were typically 20-25 bases in length

(Supplementary Data 4) and amplified a ~100 bp PCR product. Individual fold enrichment

ratios were obtained as follows: first, every Ct value was converted to $1/2^{Ct}$ and technical

triplicates were averaged to generate a single enrichment value; second, genomic location

specific enrichment was normalised by corresponding values obtained at +90° on the

chromosome. Error bars indicate the standard error of the mean for three biological

replicates. Unpaired, two-sided Student t-tests were employed to evaluate enrichment

significance where relevant for n = 3 biological replicates (each averaged from three

technical replicates) with control/test groups corresponding to parent/mutant genetic

backgrounds, respectively.

Protein structure prediction and representation

Protein models for SSB:ssDNA and SSB:RecO were generated using AlphaFold 3 and

polypeptides corresponding to individual protein entities as input[111]. Main and alternative

models were manually examined and Model 0 was chosen to highlight key features using

the Pymol Molecular Graphics 2.1 software (version 3.1)[112]. Structural highlights of the

AddAB complex were derived from the crystal structure PDB: 4CEH[68].

DATA AVAILABILITY

Source data are provided with this paper. The SSB:ssDNA and SSB:RecO models

generated by AlphaFold 3 in this study have been deposited in the Zenodo database under

accession code DOI: 10.5281/zenodo.14793919. All data presented in this study are

provided in the Supplementary Information/Source Data file and are available under open

access (CC BY). For additional information and requests, please contact the lead contact:

Charles Winterhalter.

CODE AVAILABILITY

The R code used to process next-generation sequencing data used in this study is available

under open access (CC BY) in the Zenodo database under accession code DOI:

10.5281/zenodo.14793122.

REFERENCES

1. Dwyer DJ, Kohanski MA, Collins JJ. Role of reactive oxygen species in antibiotic action and resistance. *Curr Opin Microbiol* **12**, 482-489 (2009).

2. Charbon G, Bjorn L, Mendoza-Chamizo B, Frimodt-Moller J, Lobner-Olesen A. Oxidative DNA damage is instrumental in hyperreplication stress-induced inviability of. *Nucleic Acids Research* **42**, 13228-13241 (2014).

3. Lindahl T, Wood RD. Quality control by DNA repair. *Science* **286**, 1897-1905 (1999).

4. Michel B, Sinha AK, Leach DRF. Replication Fork Breakage and Restart in Escherichia coli. *Microbiol Mol Biol Rev* **82**, (2018).

5. Kuzminov A. Single-strand interruptions in replicating chromosomes cause double-strand breaks. *Proc Natl Acad Sci U S A* **98**, 8241-8246 (2001).

6. Kornberg A, Baker TA. *DNA replication*, 2 edn. W.H. Freeman and Co (1992).

7. O'Donnell M. Replisome architecture and dynamics in Escherichia coli. *J Biol Chem* **281**, 10653-10656 (2006).

8. Kim DE, Narayan M, Patel SS. T7 DNA helicase: a molecular motor that processively and unidirectionally translocates along single-stranded DNA. *J Mol Biol* **321**, 807-819 (2002).

9. Xu ZQ, Dixon NE. Bacterial replisomes. *Curr Opin Struct Biol* **53**, 159-168 (2018).

10. Fernandez AJ, Berger JM. Mechanisms of hexameric helicases. *Crit Rev Biochem Mol Biol* **56**, 621-639 (2021).

11. Beattie TR, *et al.* Frequent exchange of the DNA polymerase during bacterial chromosome replication. *Elife* **6**, (2017).

12. Graham JE, Marians KJ, Kowalczykowski SC. Independent and Stochastic Action of DNA Polymerases in the Replisome. *Cell* **169**, 1201-1213 e1217 (2017).

13. Yeeles JT, Marians KJ. The Escherichia coli replisome is inherently DNA damage tolerant. *Science* **334**, 235-238 (2011).

14. Vrtis KB, Dewar JM, Chistol G, Wu RA, Graham TGW, Walter JC. Single-strand DNA breaks cause replisome disassembly. *Mol Cell* **81**, 1309-1318 e1306 (2021).

15. Pavani R, *et al.* Structure and repair of replication-coupled DNA breaks. *Science* **385**, eado3867 (2024).

817   16.   O'Donnell M, Langston L, Stillman B. Principles and concepts of DNA replication in
818   bacteria, archaea, and eukarya. *Cold Spring Harb Perspect Biol* **5**, (2013).
819

820   17.   Pelliciari S, Dong MJ, Gao F, Murray H. Evidence for a chromosome origin unwinding
821   system broadly conserved in bacteria. *Nucleic Acids Res* **49**, 7525-7536 (2021).
822

823   18.   Kuempel PL, Duerr SA, Seeley NR. Terminus region of the chromosome in Escherichia
824   coli inhibits replication forks. *Proc Natl Acad Sci U S A* **74**, 3927-3931 (1977).
825

826   19.   Louarn J, Patte J, Louarn JM. Evidence for a fixed termination site of chromosome
827   replication in Escherichia coli K12. *J Mol Biol* **115**, 295-314 (1977).
828

829   20.   Adams RT, Wake RG. Highly specific labeling of the *Bacillus subtilis* chromosome
830   terminus. *Journal of Bacteriology* **143**, 1036-1038 (1980).
831

832   21.   Sargent MG. Specific labeling of the *Bacillus subtilis* chromosome terminus. *Journal*
833   *of Bacteriology* **143**, 1033-1035 (1980).
834

835   22.   Xu Y, *et al.* DNA nicks in both leading and lagging strand templates can trigger break-
836   induced replication. *Mol Cell* **85**, 91-106 e105 (2025).
837

838   23.   Lenhart JS, Brandes ER, Schroeder JW, Sorenson RJ, Showalter HD, Simmons LA.
839   RecO and RecR are necessary for RecA loading in response to DNA damage and
840   replication fork stress. *J Bacteriol* **196**, 2851-2860 (2014).
841

842   24.   Jinek M, East A, Cheng A, Lin S, Ma EB, Doudna J. RNA-programmed genome editing
843   in human cells. *Elife* **2**, (2013).
844

845   25.   Lee J, *et al.* Prime editing with genuine Cas9 nickases minimizes unwanted indels.
846   *Nature Communications* **14**, (2023).
847

848   26.   Peters JM, *et al.* A Comprehensive, CRISPR-based Functional Analysis of Essential
849   Genes in Bacteria. *Cell* **165**, 1493-1506 (2016).
850

851   27.   Westers H, *et al.* Genome engineering reveals large dispensable regions in Bacillus
852   subtilis. *Mol Biol Evol* **20**, 2076-2090 (2003).
853

854   28.   Lenhart JS, Schroeder JW, Walsh BW, Simmons LA. DNA repair and genome
855   maintenance in *Bacillus subtilis*. *Microbiol Mol Biol Rev* **76**, 530-564 (2012).
856

857   29.   Simmons LA, Grossman AD, Walker GC. Replication is required for the RecA
858   localization response to DNA damage in Bacillus subtilis. *Proc Natl Acad Sci U S A*
859   **104**, 1360-1365 (2007).
860

861   30.   Lesterlin C, Ball G, Schermelleh L, Sherratt DJ. RecA bundles mediate homology
862   pairing between distant sisters during DNA break repair. *Nature* **506**, 249-253 (2014).
863

864   31.   Kidane D, Graumann PL. Dynamic formation of RecA filaments at DNA double strand
865   break repair centers in live cells. *J Cell Biol* **170**, 357-366 (2005).
866

867   32.   Simmons LA*, et al.* Comparison of responses to double-strand breaks between
868   Escherichia coli and Bacillus subtilis reveals different requirements for SOS induction.
869   *J Bacteriol* **191**, 1152-1161 (2009).
870

871   33.   Smits WK, Goranov AI, Grossman AD. Ordered association of helicase loader proteins
872   with the Bacillus subtilis origin of replication in vivo. *Mol Microbiol* **75**, 452-461
873   (2010).
874

875   34.   Michel B, Sandler SJ. Replication Restart in Bacteria. *J Bacteriol* **199**, (2017).
876

877   35.   Winterhalter C*, et al.* The DNA replication initiation protein DnaD recognises a
878   specific strand of the Bacillus subtilis chromosome origin. *Nucleic Acids Res* **51**, 4322-
879   4340 (2023).
880

881   36.   Winterhalter C*, et al.* SirA inhibits the essential DnaA:DnaD interaction to block
882   helicase recruitment during Bacillus subtilis sporulation. *Nucleic Acids Res*, (2022).
883

884   37.   Matthews LA, Simmons LA. Regulation of DNA Binding and High-Order
885   Oligomerization of the DnaB Helicase Loader. *J Bacteriol* **202**, (2020).
886

887   38.   Bruand C, Farache M, McGovern S, Ehrlich SD, Polard P. DnaB, DnaD and DnaI
888   proteins are components of the Bacillus subtilis replication restart primosome.
889   *Molecular Microbiology* **42**, 245-255 (2001).
890

891   39.   Kaguni JM. Replication initiation at the Escherichia coli chromosomal origin. *Curr*
892   *Opin Chem Biol* **15**, 606-613 (2011).
893

894   40.   Kaplan DL. The 3'-tail of a forked-duplex sterically determines whether one or two
895   DNA strands pass through the central channel of a replication-fork helicase. *J Mol*
896   *Biol* **301**, 285-299 (2000).
897

898   41.   Griffith KL, Grossman AD. Inducible protein degradation in Bacillus subtilis using
899   heterologous peptide tags and adaptor proteins to target substrates to the protease
900   ClpXP. *Mol Microbiol* **70**, 1012-1025 (2008).
901

902   42.   Kowalczykowski SC. An Overview of the Molecular Mechanisms of Recombinational
903   DNA Repair. *Cold Spring Harb Perspect Biol* **7**, (2015).
904

905   43.   Kuzminov A. Instability of inhibited replication forks in E. coli. *Bioessays* **17**, 733-741
906   (1995).
907

908   44.   Kuzminov A. Collapse and repair of replication forks in Escherichia coli. *Mol Microbiol*
909   **16**, 373-384 (1995).
910

911   45.   Kuzminov A. DNA replication meets genetic exchange: chromosomal damage and its
912   repair by homologous recombination. *Proc Natl Acad Sci U S A* **98**, 8461-8468 (2001).
913

914   46.   Kuzminov A, Schabtach E, Stahl FW. Chi sites in combination with RecA protein
915   increase the survival of linear DNA in Escherichia coli by inactivating exoV activity of
916   RecBCD nuclease. *EMBO J* **13**, 2764-2776 (1994).
917

918   47.   Yasmin T, Azeroglu B, Cockram CA, Leach DRF. Distribution of Holliday junctions and
919   repair forks during Escherichia coli DNA double-strand break repair. *PLoS Genet* **17**,
920   e1009717 (2021).
921

922   48.   Sanchez H, Kidane D, Castillo Cozar M, Graumann PL, Alonso JC. Recruitment of
923   Bacillus subtilis RecN to DNA double-strand breaks in the absence of DNA end
924   processing. *J Bacteriol* **188**, 353-360 (2006).
925

926   49.   Costes A, Lecointe F, McGovern S, Quevillon-Cheruel S, Polard P. The C-terminal
927   domain of the bacterial SSB protein acts as a DNA maintenance hub at active
928   chromosome replication forks. *PLoS Genet* **6**, e1001238 (2010).
929

930   50.   Ramos C*, et al.* The RecD2 helicase balances RecA activities. *Nucleic Acids Research*
931   **50**, 3432-3444 (2022).
932

933   51.   Cárdenas PP*, et al.* RecX Facilitates Homologous Recombination by Modulating RecA
934   Activities. *Plos Genetics* **8**, (2012).
935

936   52.   Mortier-Barrière I*, et al.* A key presynaptic role in transformation for a widespread
937   bacterial protein:: DprA conveys incoming ssDNA to RecA. *Cell* **130**, 824-836 (2007).
938

939   53.   Narula J, Kuchina A, Lee DD, Fujita M, Suel GM, Igoshin OA. Chromosomal
940   Arrangement of Phosphorelay Genes Couples Sporulation and DNA Replication. *Cell*
941   **162**, 328-337 (2015).
942

943   54.   Azeroglu B, Leach DRF. RecG controls DNA amplification at double-strand breaks and
944   arrested replication forks. *FEBS Lett* **591**, 1101-1113 (2017).
945

946   55.   Azeroglu B*, et al.* RecG Directs DNA Synthesis during Double-Strand Break Repair.
947   *PLoS Genet* **12**, e1005799 (2016).
948

949   56.   Kuzminov A. Recombinational repair of DNA damage in Escherichia coli and
950   bacteriophage lambda. *Microbiol Mol Biol Rev* **63**, 751-813, table of contents (1999).
951

952   57.   Chedin F, Ehrlich SD, Kowalczykowski SC. The Bacillus subtilis AddAB
953   helicase/nuclease is regulated by its cognate Chi sequence in vitro. *J Mol Biol* **298**, 7-
954   20 (2000).
955

956   58.   Symington LS. End resection at double-strand breaks: mechanism and regulation.
957   *Cold Spring Harb Perspect Biol* **6**, (2014).

959   59.   Yeeles JT, Dillingham MS. A dual-nuclease mechanism for DNA break processing by
960   AddAB-type helicase-nucleases. *J Mol Biol* **371**, 66-78 (2007).
961

962   60.   Kidane D, Sanchez H, Alonso JC, Graumann PL. Visualization of DNA double-strand
963   break repair in live bacteria reveals dynamic recruitment of Bacillus subtilis RecF,
964   RecO and RecN proteins to distinct sites on the nucleoids. *Mol Microbiol* **52**, 1627-
965   1639 (2004).
966

967   61.   Cox MM, Goodman MF, Keck JL, van Oijen A, Lovett ST, Robinson A. Generation and
968   Repair of Postreplication Gaps in Escherichia coli. *Microbiol Mol Biol Rev* **87**,
969   e0007822 (2023).
970

971   62.   Nirwal S*, et al.* Mechanism of RecF-RecO-RecR cooperation in bacterial homologous
972   recombination. *Nat Struct Mol Biol* **30**, 650-660 (2023).
973

974   63.   Bell JC, Kowalczykowski SC. RecA: Regulation and Mechanism of a Molecular Search
975   Engine: (Trends in Biochemical Sciences, June 2016, Vol. 41, No. 6, 491--507). *Trends*
976   *Biochem Sci* **41**, 646 (2016).
977

978   64.   Bianco PR. I came to a fork in the DNA and there was RecG. *Prog Biophys Mol Biol*
979   **117**, 166-173 (2015).
980

981   65.   Lloyd RG, Rudolph CJ. 25 years on and no end in sight: a perspective on the role of
982   RecG protein. *Curr Genet* **62**, 827-840 (2016).
983

984   66.   West SC. Processing of recombination intermediates by the RuvABC proteins. *Annu*
985   *Rev Genet* **31**, 213-244 (1997).
986

987   67.   Rosch TC*, et al.* Single molecule tracking reveals spatio-temporal dynamics of
988   bacterial DNA repair centres. *Sci Rep* **8**, 16450 (2018).
989

990   68.   Krajewski WW, Fu X, Wilkinson M, Cronin NB, Dillingham MS, Wigley DB. Structural
991   basis for translocation by AddAB helicase-nuclease and its arrest at chi sites. *Nature*
992   **508**, 416-419 (2014).
993

994   69.   El Karoui M, Biaudet V, Schbath S, Gruss A. Characteristics of Chi distribution on
995   different bacterial genomes. *Res Microbiol* **150**, 579-587 (1999).
996

997   70.   Chedin F, Noirot P, Biaudet V, Ehrlich SD. A five-nucleotide sequence protects DNA
998   from exonucleolytic degradation by AddAB, the RecBCD analogue of Bacillus subtilis.
999   *Mol Microbiol* **29**, 1369-1377 (1998).
1000

1001   71.   Halpern D*, et al.* Identification of DNA motifs implicated in maintenance of bacterial
1002   core genomes by predictive modeling. *PLoS Genet* **3**, 1614-1621 (2007).
1003

1004 72. Dubiel K, *et al*. Structural Mechanisms of Cooperative DNA Binding by Bacterial
1005 Single-Stranded DNA-Binding Proteins. *J Mol Biol* **431**, 178-195 (2019).
1006

1007 73. Shereda RD, Kozlov AG, Lohman TM, Cox MM, Keck JL. SSB as an organizer/mobilizer
1008 of genome maintenance complexes. *Crit Rev Biochem Mol Biol* **43**, 289-318 (2008).
1009

1010 74. Lecointe F, *et al*. Anticipating chromosomal replication fork arrest: SSB targets repair
1011 DNA helicases to active forks. *EMBO J* **26**, 4239-4251 (2007).
1012

1013 75. Sakai A, Cox MM. RecFOR and RecOR as distinct RecA loading pathways. *J Biol Chem*
1014 **284**, 3264-3272 (2009).
1015

1016 76. Morimatsu K, Kowalczykowski SC. RecFOR proteins load RecA protein onto gapped
1017 DNA to accelerate DNA strand exchange: a universal step of recombinational repair.
1018 *Mol Cell* **11**, 1337-1347 (2003).
1019

1020 77. Raychaudhury P, Marians KJ. The recombination mediator proteins RecFOR maintain
1021 RecA* levels for maximal DNA polymerase V Mut activity. *J Biol Chem* **294**, 852-860
1022 (2019).
1023

1024 78. Carrasco B, Torres R, Moreno-Del Alamo M, Ramos C, Ayora S, Alonso JC. Processing
1025 of stalled replication forks in Bacillus subtilis. *FEMS Microbiol Rev* **48**, (2024).
1026

1027 79. Sanchez H, Carrasco B, Cozar MC, Alonso JC. Bacillus subtilis RecG branch migration
1028 translocase is required for DNA repair and chromosomal segregation. *Mol Microbiol*
1029 **65**, 920-935 (2007).
1030

1031 80. Torres R, Gandara C, Carrasco B, Baquedano I, Ayora S, Alonso JC. DisA Limits RecG
1032 Activities at Stalled or Reversed Replication Forks. *Cells* **10**, (2021).
1033

1034 81. Torres R, Romero H, Rodriguez-Cerrato V, Alonso JC. Interplay between Bacillus
1035 subtilis RecD2 and the RecG or RuvAB helicase in recombinational repair. *DNA Repair*
1036 *(Amst)* **55**, 40-46 (2017).
1037

1038 82. Ayora S, Carrasco B, Doncel-Perez E, Lurz R, Alonso JC. Bacillus subtilis RecU protein
1039 cleaves Holliday junctions and anneals single-stranded DNA. *Proc Natl Acad Sci U S A*
1040 **101**, 452-457 (2004).
1041

1042 83. Sun B, *et al*. T7 replisome directly overcomes DNA damage. *Nat Commun* **6**, 10260
1043 (2015).
1044

1045 84. Fernandez S, Ayora S, Alonso JC. Bacillus subtilis homologous recombination: genes
1046 and products. *Res Microbiol* **151**, 481-486 (2000).
1047

1048 85. Unciuleac MC, Shuman S. Double strand break unwinding and resection by the
1049 mycobacterial helicase-nuclease AdnAB in the presence of single strand DNA-binding
1050 protein (SSB). *J Biol Chem* **285**, 34319-34329 (2010).

86. Lovett ST. The DNA Exonucleases of Escherichia coli. *EcoSal Plus* **4**, (2011).

87. Gupta R, Unciuleac MC, Shuman S, Glickman MS. Homologous recombination mediated by the mycobacterial AdnAB helicase without end resection by the AdnAB nucleases. *Nucleic Acids Res* **45**, 762-774 (2017).

88. Su'etsugu M, Errington J. The replicase sliding clamp dynamically accumulates behind progressing replication forks in *Bacillus subtilis* cells. *Mol Cell* **41**, 720-732 (2011).

89. Reyes-Lamothe R, Sherratt DJ, Leake MC. Stoichiometry and architecture of active DNA replication machinery in Escherichia coli. *Science* **328**, 498-501 (2010).

90. Mascarenhas J, Weber MH, Graumann PL. Specific polar localization of ribosomes in *Bacillus subtilis* depends on active transcription. *EMBO Rep* **2**, 685-689. (2001).

91. Lewis PJ, Thaker SD, Errington J. Compartmentalization of transcription and translation in *Bacillus subtilis*. *EMBO J* **19**, 710-718 (2000).

92. Yeeles JT, van Aelst K, Dillingham MS, Moreno-Herrero F. Recombination hotspots and single-stranded DNA binding proteins couple DNA translocation to DNA unwinding by the AddAB helicase-nuclease. *Mol Cell* **42**, 806-816 (2011).

93. Umezu K, Kolodner RD. Protein interactions in genetic recombination in Escherichia coli. Interactions involving RecO and RecR overcome the inhibition of RecA by single-stranded DNA-binding protein. *J Biol Chem* **269**, 30005-30013 (1994).

94. Hobbs MD, Sakai A, Cox MM. SSB protein limits RecOR binding onto single-stranded DNA. *J Biol Chem* **282**, 11058-11067 (2007).

95. Inoue J, Honda M, Ikawa S, Shibata T, Mikawa T. The process of displacing the single-stranded DNA-binding protein from single-stranded DNA by RecO and RecR proteins. *Nucleic Acids Res* **36**, 94-109 (2008).

96. Manfredi C, Carrasco B, Ayora S, Alonso JC. Bacillus subtilis RecO nucleates RecA onto SsbA-coated single-stranded DNA. *J Biol Chem* **283**, 24837-24847 (2008).

97. Bonde NJ, Henry C, Wood EA, Cox MM, Keck JL. Interaction with the carboxy-terminal tip of SSB is critical for RecG function in E. coli. *Nucleic Acids Res* **51**, 3735-3753 (2023).

98. Buss JA, Kimura Y, Bianco PR. RecG interacts directly with SSB: implications for stalled replication fork regression. *Nucleic Acids Research* **36**, 7029-7042 (2008).

99. Kozlov AG, Jezewska MJ, Bujalowski W, Lohman TM. Binding Specificity of

1097   Single-Stranded DNA Binding Protein for the χ Subunit of DNA pol III Holoenzyme and PriA
1098   Helicase. *Biochemistry-Us* **49**, 3555-3566 (2010).

1099

1100   100.   Gomez-Campo CL*, et al.* Differential requirement for RecFOR pathway components
1101   in Thermus thermophilus. *Environ Microbiol Rep* **16**, e13269 (2024).

1102

1103   101.   Bentchikou E, Servant P, Coste G, Sommer S. A major role of the RecFOR pathway in
1104   DNA double-strand-break repair through ESDSA in Deinococcus radiodurans. *PLoS*
1105   *Genet* **6**, e1000774 (2010).

1106

1107   102.   Henry C, Mbele N, Cox MM. RecF protein targeting to postreplication (daughter
1108   strand) gaps I: DNA binding by RecF and RecFR. *Nucleic Acids Res* **51**, 5699-5713
1109   (2023).

1110

1111   103.   Henry C*, et al.* RecF protein targeting to post-replication (daughter strand) gaps II:
1112   RecF interaction with replisomes. *Nucleic Acids Res* **51**, 5714-5742 (2023).

1113

1114   104.   McGlynn P, Al-Deib AA, Liu J, Marians KJ, Lloyd RG. The DNA replication protein PriA
1115   and the recombination protein RecG bind D-loops. *J Mol Biol* **270**, 212-221 (1997).

1116

1117   105.   Doench JG*, et al.* Optimized sgRNA design to maximize activity and minimize off-
1118   target effects of CRISPR-Cas9. *Nat Biotechnol* **34**, 184-191 (2016).

1119

1120   106.   Hanahan D, Jessee J, Bloom FR. Plasmid transformation of Escherichia coli and other
1121   bacteria. *Methods Enzymol* **204**, 63-113 (1991).

1122

1123   107.   Anagnostopoulos C, Spizizen J. Requirements for transformation in *Bacillus subtilis. J*
1124   *Bacteriol* **81**, 741-746 (1961).

1125

1126   108.   Lucet I, Feucht A, Yudkin MD, Errington J. Direct interaction between the cell division
1127   protein FtsZ and the cell differentiation protein SpoIIE. *EMBO J* **19**, 1467-1475
1128   (2000).

1129

1130   109.   Schindelin J*, et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
1131   *Methods* **9**, 676-682 (2012).

1132

1133   110.   Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JA. Primer3Plus, an
1134   enhanced web interface to Primer3. *Nucleic Acids Res* **35**, W71-74 (2007).

1135

1136   111.   Abramson J*, et al.* Accurate structure prediction of biomolecular interactions with
1137   AlphaFold 3. *Nature* **630**, 493-500 (2024).

1138

1139   112.   Delano WL. The PyMOL Molecular Graphics System.) (2002).

1140

1141

1142

ACKNOWLEDGEMENTS

This work was supported by a Wellcome Trust Early-Career Award [226338/Z/22/Z] to CW and Wellcome Trust Discovery Award [225811/Z/22/Z] to HM. We would also like to express our gratitude to Frances Davison and Amandeep Kaur for technical assistance, James Grimshaw for microscope maintenance and Frederic Schramm for insightful discussions.

AUTHOR CONTRIBUTIONS STATEMENT

CW and HM contributed to the conception/design of the work. CW, KJS and SF generated results presented in the manuscript. CW created figures. CW and HM wrote/edited the manuscript.

COMPETING INTERESTS STATEMENT

The authors declare no competing interests.

FIGURE LEGENDS/CAPTIONS

Figure 1. Single-strand discontinuities arrest DNA synthesis and generate DNA damage.

(A) Diagram illustrating the impact of leading and lagging strand discontinuities on a replication fork, with the bacterial replicative helicase translocating along the lagging strand template.

(B) Genetic system employed to create a site-specific single-strand discontinuity in the *B. subtilis* chromosome. Black boxes correspond to antibiotic resistance markers, pink to Cas9 variants and green to sgRNA. Chromosome annotations refer to the replication origin (0°), terminus (180°) and sgRNA target sequence (yellow box).

(C) Spot-titre analysis of strains engineered to express (+xylose) either nCas9 or dCas9. Individual sgRNAs were designed to target the leading or lagging strand template at loci located at either -90° (top panel) or +90° (bottom panel).

(D) MFA using whole genome sequencing of strains engineered to express (+xylose) either nCas9 or dCas9. Individual sgRNAs were designed to target the leading or lagging strand template at loci located at either -90° (left panel) or +90° (right panel). The frequency of sequencing reads was plotted against genome position.

(E) Fluorescence microscopy images of the DNA damage reporter mNG-RecA in strains engineered to express (+xylose) either nCas9 or dCas9 (sgRNAs target -90°). Fluorescent RecA organises into bundles in the presence of DNA damage. Green signal corresponds to mNG-RecA fluorescence, grey scale images show corresponding phase contrast images. Scale bar indicates 2.5 µm.

(F) Quantification of fluorescence microscopy images of the DNA damage reporter mNG-RecA in strains engineered to express (+xylose) either nCas9 or dCas9 (sgRNA target -90°). Detection of spots and bundles corresponds to distinct foci or filament-like structures annotated in panel (E). Error bars indicate standard deviation and circles overlaid on bars correspond to two biological replicates.

Source data are provided as a Source Data file.

1186

Figure 2. Replication restart is necessary to resume DNA synthesis following replisome inactivation at a single-strand discontinuity.

(A) Diagram illustrating genomic loci probed via ChIP-qPCR at either -90° or +90°. The sgRNA targeted Cas9 protein variants to the locus at -90°. Lagging strand and leading strand templates are represented in black and grey lines, respectively (numbers refer to strand polarity). The blue arrow indicates leading strand synthesis and red arrows indicate lagging strand synthesis.

(B-F) ChIP-qPCR analyses of helicase (DnaC) and replication restart proteins (PriA, DnaD, DnaB, DnaI) in strains engineered to express either nCas9 or dCas9. The sgRNAs targeted Cas9 proteins to a locus located at -90° and protein enrichment was normalised to the control locus at +90°. Error bars indicate the standard error of the mean and circles overlaid on bars correspond to three biological replicates.

(G) Diagram illustrating the fate of helicase (red circle) upon encountering a single-strand discontinuity in either the leading or lagging strand template.

(H) MFA using whole genome sequencing in strains engineered to express low levels of sgRNAs and Cas9 proteins (weak nicking system). The frequency of sequencing reads was plotted against genome position.

(I) Genetic system employed to express limiting levels of PriA in strains harbouring the weak nicking system.

(J) Spot-titre analysis of strains engineered to express nCas9 (+xylose) with either high or limiting PriA. sgRNAs were targeted to a locus at -90°.

(K) MFA using whole genome sequencing in strains engineered to express (+xylose) nCas9 with either high or limiting PriA. The frequency of sequencing reads was plotted against genome position.

Source data are provided as a Source Data file.

Figure 3. The repair of single-strand discontinuities requires recombination.

(A) Spot-titre analysis of recombinational repair mutants in strains engineered to express

(+xylose) nCas9. sgRNAs were targeted to a locus at -90°. The parental nicking strain

control is shown above.

(B) MFA using whole genome sequencing of recombinational repair mutants in the presence

or absence of the weak nCas9 system. The frequency of sequencing reads was plotted

against genome position.

(C) Diagram illustrating proposed molecular events required for recombinational repair to

repair a seDSB created after a replication fork encounters a single-strand discontinuity in the

leading strand template.

Source data are provided as a Source Data file.

1224

Figure 4. The helicase activity of AddAB is necessary and sufficient to support DNA

replication past a nick.

(A) Diagram illustrating differential resection of a linear DNA in the presence of either the

AddA or AddB helicase/nuclease activities. Dotted lines indicate nuclease cleavage.

(B) ChIP-qPCR analyses of mNG-AddA and mNG-AddB in strains engineered to express

nCas9. Left panel shows the genetic system employed in this experiment. The sgRNAs

targeted nCas9 to a locus located at -90° and protein enrichment was normalised to the

control locus at +90°. Error bars indicate the standard error of the mean and circles overlaid

on bars correspond to three biological replicates.

(C) Spot-titre analysis in strains with AddAB variants (AddA$^{\Delta HEL}$, AddA$^{\Delta NUC}$, AddB$^{\Delta NUC}$

AddA$^{\Delta NUC}$AddB$^{\Delta NUC}$) engineered to express (+xylose) nCas9. sgRNAs were targeted to a

locus at -90°. The parental nicking strain control is shown above.

(D) MFA using whole genome sequencing of strains with either AddA$^{\Delta HEL}$ or

AddA$^{\Delta NUC}$AddB$^{\Delta NUC}$ engineered to express nCas9. Top panel shows the genetic system

employed in this experiment. The frequency of sequencing reads was plotted against

genome position.

Source data are provided as a Source Data file.

1242

Figure 5. SSB requires AddA helicase activity for loading upstream of a nick.

(A) ChIP-qPCR analyses of SSB in strains engineered to express nCas9 in the presence of either endogenous AddAB (top) or heterologous RexAB (bottom) end-processing machinery. Locations for respective *chi* sites are indicated. The sgRNAs targeted nCas9 to a locus located at -90° and protein enrichment was normalised to the control locus at +90°.

(B) ChIP-qPCR analyses of SSB in strains with recombinational repair mutants engineered to express nCas9. qPCR primers amplified either a site located -3 kb from the sgRNA target (-90°) or a control locus (+90°) used for normalisation. Note a significant increase in SSB enrichment in some of mutants tested under these conditions. Two-sided Student t-test results: * indicates p-value = 0.0143 for lagging strand Δ*recF*, p-value = 0.1666 and p-value = 0.0223 for leading/strand Δ*recG* backgrounds, respectively (n = 3, see Methods/Source Data for details).

(C) ChIP-qPCR analyses of SSB in strains with AddA and AddB variants engineered to express nCas9. Alleles refer to wild-type (WT), knockouts (Δ), and point mutations inactivating nuclease (ΔNUC) or helicase (ΔHEL) activities. qPCR primers amplified either a site located -3 kb from the sgRNA target (-90°) or a control locus (+90°) used for normalisation.

(A-C) Error bars indicate the standard error of the mean and circles overlaid on bars correspond to three biological replicates.

Source data are provided as a Source Data file.

Figure 6. The SSB-CTT is essential to recruit RecO following replisome inactivation at a single-strand discontinuity.

(A) Schematic of SSB primary structure with corresponding functions. Numbers indicate amino acid residue boundaries between the N-terminal domain (NTD), C-terminal domain (CTD) and C-terminal tail (CTT).

(B) Structural model *of B. subtilis* SSB homotetramer in the presence of single-strand DNA

(AlphaFold 3). A sixty nucleotide homopolymer of thymidine bases (dT_{60}) is shown in black

and SSB-CTT residues are highlighted in red.

(C) Spot-titre analysis of SSB variants in strains engineered to express (+xylose) nCas9.

Top panel shows the genetic system employed in this experiment. sgRNAs were targeted to

a locus at -90°.

(D) MFA using whole genome sequencing in strains with SSB variants engineered to

express nCas9. The frequency of sequencing reads was plotted against genome position.

(E) ChIP-qPCR analyses of SSB, His-RecO and His-RecA proteins in strains engineered to

express nCas9. qPCR primers amplified either a site located -3 kb from the sgRNA target (-

90°) or a control locus (+90°) used for normalisation. Error bars indicate the standard error of

the mean and circles overlaid on bars correspond to three biological replicates.

(F) ChIP-qPCR analyses of His-RecA in strains with recombinational repair mutants

engineered to express nCas9. Top panel shows the genetic system employed in this

experiment. qPCR primers amplified either a site located -3 kb from the sgRNA target (-90°)

or a control locus (+90°) used for normalisation. Error bars indicate the standard error of the

mean and circles overlaid on bars correspond to three biological replicates.

Source data are provided as a Source Data file.

1287

Figure 7. RecG and RecU are recruited to sites of recombinational repair.

(A) ChIP-qPCR analyses of FLAG-RecG in strains engineered to express nCas9. Top panel

shows the genetic system employed in this experiment. The sgRNAs targeted nCas9 to a

locus located at -90° and protein enrichment was normalised to the control locus at +90°.

Error bars indicate the standard error of the mean and circles overlaid on bars correspond to

three biological replicates.

(B) Fluorescence microscopy images of Δ*recU* mutant strains engineered to express nCas9

(sgRNAs targeted to -90°). Grey scale images correspond to phase contrast. Blue signal

corresponds to DAPI staining (DNA) and red signal corresponds to Nile red staining

(membrane) in composite fluorescence images. Cas9 proteins were induced (+xylose)

during the exponential growth phase for 90 minutes before cells were collected for imaging.

Scale bar indicates 5 μm.

(C) ChIP-qPCR analyses of RecU-FLAG in a strain engineered to express nCas9. Top panel

shows the genetic system employed in this experiment. The sgRNAs targeted nCas9 to a

locus located at -90° and protein enrichment was normalised to the control locus at +90°.

Error bars indicate the standard error of the mean and circles overlaid on bars correspond to

three biological replicates.

Source data are provided as a Source Data file.

1306

1307   **Figure 8. Model for seDSB repair and helicase reloading following replisome**

1308   **inactivation at a single-strand discontinuity.**

1309   Model illustrating molecular events that are required or dispensable for recombinational

1310   repair of a seDSB created after a replication fork encounters a single-strand discontinuity

1311   located on either the leading or lagging strand template.

1312

 SUPPLEMENTARY FIGURE LEGENDS/CAPTIONS

Supplementary Figure 1. Impact of Cas9 variants on *B. subtilis* growth.

(A) Spot-titre analysis of strains expressing (+xylose) either Cas9^{D10A} nickase, Cas9^{H840A}

nickase, or Cas9$^{D10A/H840A}$ catalytically inactive nuclease in the presence of the same sgRNA.

Left panel shows the genetic system employed in this experiment. sgRNAs were targeted to

a locus at -90°.

(B) Immunoblot of Cas9 variants in the presence of a sgRNA targeted to a locus at -90°.

Cas9 proteins were induced (+xylose) during the exponential growth phase for 1 hour before

cells were harvested for protein extraction. The tubulin homolog FtsZ was used as a loading

control and data is representative from at least two biological repeats.

(C) Spot-titre analysis of a strains expressing (+xylose) the Cas9^{D10A} nickase with different

sgRNAs targeting locations along the left chromosome arm.

(D) Spot-titre analysis of strains expressing (+xylose) the Cas9$^{D10A-H840A}$ catalytically inactive

nuclease. Left panel shows the genetic system employed in this experiment for sgRNAs

targeted to a locus at either -90° or +90°.

(E) Strains were induced to express (+xylose) nCas9 or dCas9 during exponential or

stationary phase for 4 hours, followed by spot-titre analysis in the absence of xylose to

determine CFUs remaining after treatment. sgRNAs were targeted to a locus at -90°.

(F) Spot-titre analysis of suppressor colonies isolated from strains expressing (+xylose)

nCas9. sgRNAs were targeted to a locus at -90°. The parental nicking strain control is

shown above. Identity of suppressors was identified via sequencing.

Source data are provided as a Source Data file.

1335

Supplementary Figure 2. MFA using whole genome sequencing of *B. subtilis* Δ6.

The frequency of sequencing reads was plotted against genome position. Note that arbitrary

chromosome boundaries indicated on x-axis may not represent termination events occurring

in the cell (*terminus* region highlighted in yellow). Redundant ribosomal DNA sequences

result in peaks throughout the genome (rDNA sequences highlighted in brown).

Source data are provided as a Source Data file.

1342

Supplementary Figure 3. Live cell fluorescence microscopy using the mNG-RecA DNA

damage reporter.

(A) Phase contrast image and fluorescence signal from the mNG-RecA reporter in wild-type

B. subtilis under standard laboratory growth conditions. Scale bar indicates 2.5 µm.

(B) Fluorescence microscopy images of mNG-RecA in strains engineered to express either

nCas9 or dCas9 (sgRNA target -90°). Fluorescent RecA organises into bundles in the

presence of DNA damage. Green signal corresponds to mNG-RecA fluorescence, grey

scale images show corresponding phase contrast images. Cas9 proteins were induced

(+xylose) during the exponential growth phase for 1 hour before cells were collected for

imaging. Scale bar indicates 5 µm.

Source data are provided as a Source Data file.

1354

Supplementary Figure 4. The replicative helicase DnaC is recruited near a single-

strand discontinuity.

ChIP-qPCR analyses of helicase (DnaC) in strains engineered to express either nCas9 or

dCas9. The sgRNAs targeted Cas9 proteins to a locus located at -90°. The replication origin

(*oriC*) was used as positive control and protein enrichment was normalised to the control

locus at +90°. Error bars indicate the standard error of the mean and circles overlaid on bars

correspond to three biological replicates.

Source data are provided as a Source Data file.

1363

Supplementary Figure 5. An attenuated nCas9 system does not arrest cell growth.

(A) Schematic showing differences between the strong and weak nicking systems used to

introduce a single-strand discontinuity.

(B) Spot-titre analysis of strains expressing (+xylose) the strong or weak nicking system in

the presence of the same sgRNA. Control indicates a strain encoding the Cas9[D10A/H840A]

catalytically inactive nuclease. sgRNAs targeted to a locus located at -90°.

(C) Live cell fluorescence microscopy in strains expressing either the strong or the weak

nCas9 system. Grey scale images correspond to phase contrast, blue signal corresponds to

DAPI staining (nucleoid), red signal corresponds to Nile red staining (membrane) and merge

corresponds to DAPI/Nile red composite images. Cas9 proteins were induced (+xylose)

during the exponential growth phase for 90 minutes before cells were collected for imaging.

Source data are provided as a Source Data file.

1376

Supplementary Figure 6. System employed to limit PriA expression during replisome

inactivation at single-strand discontinuities.

(A) Genetic system employed for *priA* complementation. The IPTG inducible *priA-ssrA*

cassette was placed at an ectopic locus.

(B) Spot-titre analysis of strains expressing different levels of PriA-ssrA (IPTG-dependent) in

the presence or absence of the endogenous *priA*.

(C) Immunoblot of PriA variants expressed at a range of IPTG concentrations. The tubulin

homolog FtsZ was used as a loading control and data is representative from at least two

biological repeats.

(D) Spot-titre analysis of strains with a low or high level of PriA-ssrA (uninduced or 1 mM

IPTG) engineered to express nCas9 (+xylose). sgRNAs were targeted to a locus at -90°.

Source data are provided as a Source Data file.

1389

Supplementary Figure 7. Recombinational repair is necessary for viability following

replisome inactivation at a single-strand discontinuity.

(A) Spot-titre analysis of recombinational repair mutants in strains engineered to express

(+xylose) nCas9. sgRNAs were targeted to a locus at -90°. The parental nicking strain

control is shown above. Plates were the same as shown in Fig. 3A imaged after 48 hours.

(B) Spot-titre analysis of recombinational repair mutants complemented with ectopic epitope-

tagged copies in strains engineered to express (+xylose) nCas9. sgRNAs were targeted to a

locus at -90°. The parental weak nicking strain is shown above.

(C) Diagram illustrating proposed molecular events required for recombinational repair to

repair the seDSB created after a replication fork encounters a single-strand discontinuity in

the lagging strand template.

1401

1402 **Supplementary Figure 8. Viability following replisome inactivation at a single-strand**

1403 **discontinuity relies on a minimal set of recombinational repair genes.**

1404 **(A-D)** Spot-titre analysis of DNA repair mutants grouped by known/proposed functions in

1405 strains engineered to express (+xylose) nCas9. sgRNAs were targeted to a locus at -90°.

1406 The parental nicking strain control is shown above.

1407

Supplementary Figure 9. MFA using whole genome sequencing of recombinational

repair mutants.

(A-C) MFA using whole genome sequencing of (A) Δ*addA*, Δ*addB,* Δ*recF*, (B) Δ*recO*,

Δ*recR,* Δ*recA*, (C) Δ*recU*, and (C-D) Δ*recG* mutant strains. The frequency of sequencing

reads was plotted against genome position (same data as in Fig. 3B for background). A

region flanking the replication origin is enlarged.

Source data are provided as a Source Data file.

1415

Supplementary Figure 10. MFA using whole genome sequencing of recombinational

repair mutants in strains expressing nCas9: highlighting the *oriC* region.

MFA using whole genome sequencing of recombinational repair mutant strains engineered

to express nCas9. The frequency of sequencing reads was plotted against genome position

(same data as in Fig. 3B knockout panels). A region flanking the replication origin is

enlarged.

Source data are provided as a Source Data file.

1423

Supplementary Figure 11. MFA using whole genome sequencing of post-synapsis recombinational repair mutants in strains expressing nCas9.

(A-B) MFA using whole genome sequencing in either (A) Δ*recG* or (A) Δ*recU* mutant strains engineered to express nCas9. The frequency of sequencing reads was plotted against genome position (same data as in Fig. 3B post-synapsis panels). Genomic regions of interest are enlarged.

Source data are provided as a Source Data file.

1431

Supplementary Figure 12. End-processing activities required following replisome inactivation at a single-strand discontinuity.

(A-B) MFA using whole genome sequencing of strains with either AddA$^{\Delta HEL}$, AddA$^{\Delta NUC}$ or AddB$^{\Delta NUC}$ engineered to express nCas9. The frequency of sequencing reads was plotted against genome position. AddA$^{\Delta HEL}$ samples were the same as shown in Fig. 4D.

(C) Immunoblot of mNG-AddA variants. The tubulin homolog FtsZ was used as a loading control and data is representative from at least two biological repeats.

(D) Spot-titre analysis of Δ*recJ* mutants combined with AddAB nuclease deficient variants in strains engineered to express (+xylose) nCas9. sgRNAs were targeted to a locus at -90°. The parental nicking strain control is shown above.

(E) Spot-titre analysis of Δ*addAB* mutants complemented with *rexAB* in strains engineered to express (+xylose) nCas9. sgRNAs were targeted to a locus at -90°. The parental nicking strain control is shown above.

(F) Crystal structure of the *B. subtilis* AddAB complex with locations of catalytic sites and the AddA^{D682} residue indicated (PDB template: 4CEH). The black backbone corresponds to DNA, AddA is displayed in orange and AddB in beige.

Source data are provided as a Source Data file.

1450

Supplementary Figure 13. System employed to investigate the function of the SSB-

CTT following replisome inactivation at single-strand discontinuities.

(A) Diagram illustrating potential essential interactions of the SSB-CTT.

(B) Genetic system employed for *ssbA* complementation. An IPTG inducible copy of *ssbA*

was placed at an ectopic locus.

(C) Spot-titre analysis of strains expressing (+IPTG) *ssbA* variants with and without the

endogenous *ssbA*.

(D) Immunoblot of SSB variants expressed using the *ssbA* complementation system. The

tubulin homolog FtsZ was used as a loading control and data is representative from at least

two biological repeats.

(E) Genetic system employed to express SSB variants in a background encoding the weak

nicking system.

(F) Structural model of a complex between a SSB homotetramer and RecO from *B. subtilis*

(AlphaFold 3). Cyan corresponds to SSB with CTT residues coloured in red and RecO is

displayed in beige.

Source data are provided as a Source Data file.

1467

Supplementary Figure 14. A Δ*recU* mutant displays chromosome segregation defects.

(A-B) Live cell fluorescence microscopy of Δ*recU* mutants using nucleoid and membrane

stains in (A) wild-type or (B) strains expressing nCas9. Grey scale images correspond to

phase contrast, blue signal corresponds to DAPI staining (nucleoid), red signal corresponds

to Nile red staining (membrane) and merge corresponds to DAPI/Nile red composite images.

Cas9 proteins were induced (+xylose) during the exponential growth phase for 90 minutes

before cells were collected for imaging.

Source data are provided as a Source Data file.

1476

1477 **Supplementary Figure 15. Potential role of RecG in replication fork remodelling.**

1478   Diagram illustrating the proposed function of RecG to ensure productive helicase reloading

1479   following replisome inactivation at a single-strand discontinuity.

1480

SUPPLEMENTARY DATA

File name: 20251019_Supplementary-Data.xlsx

Description. Excel file containing all Supplementary Data spreadsheets.

Supplementary Data 1. Supplementary Table too large to fit in a PDF containing the list of

repair gene knockouts tested in the genetic screen (related to Fig. 3 and Supplementary

Figure 8).

Supplementary Data 2. Supplementary Table too large to fit in a PDF containing genotypes

from strains generated in this study.

Supplementary Data 3. Supplementary Table too large to fit in a PDF containing genotypes

and sequences of recombinant DNA used to construct strains in this study, related to

Supplementary Data 2.

Supplementary Data 4. Supplementary Table too large to fit in a PDF containing

oligonucleotide sequences used for qPCR in this study.

GRAPHICAL ABSTRACT

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

A

B

C

FIGURE 5

FIGURE 6

A

B

C

FIGURE 7

(I) Fork collapse:
Generation of a DSB and detection via AddAB

(II) End processing:
No strict requirement for DNA end resection and *chi* dependence

(III) SSB loading:
AddAB helicase switch promotes SSB enrichment

(IV) Pre-synapsis:
Dependent on AddAB and RecO recruitment via SSB-CTT

(V) RecA recruitment:
Dependent on AddAB, RecFOR and SSB-CTT:RecO interaction

(VI) Strand exchange:
Alignment with sister chromosome and D-loop formation

(VII) Fork regression:
RecG remodels the lagging strand, holliday junction cleavage dispensable

(VIII) PriA loading:
Recruitment of restart proteins for helicase reloading

(IX) Replication restart:
Helicase loaded on lagging strand and DNA synthesis resumes

AddA-AddB end-processing machinery

SSB tetramer

RecOR:RecF

RecA loading

Synapsis

RecG   RecU resolvase

PriA   Restart pathway

Replicative helicase

Leading strand discontinuity

Lagging strand discontinuity

FIGURE 8

SUPPLEMENTARY FIGURE 1

rDNA sequences (clusters throughout genome)

terminus region

SUPPLEMENTARY FIGURE 2

A

Phase contrast   mNG-RecA

Scale bar = 2.5 µm

B

	Leading strand		Lagging strand	
	dCas9	nCas9	dCas9	nCas9

Phase contrast

mNG-RecA

bundle   bundle   Scale bar = 5 µm

SUPPLEMENTARY FIGURE 3

SUPPLEMENTARY FIGURE 4

SUPPLEMENTARY FIGURE 5

A

PriA complementation

B

C

D

SUPPLEMENTARY FIGURE 6

A

DNA repair mutants

Leading strand | Lagging strand
- xylose (48 h) | + xylose (48 h) | - xylose (48 h) | + xylose (48 h)

Parent
ΔaddA — End processing (orange)
ΔaddB
ΔrecF — Early HR
ΔrecO
ΔrecR
ΔrecA — Synapsis
ΔrecG — Late HR
ΔrecU

Homologous recombination

B

DNA repair mutants

Leading strand | Lagging strand
- xylose | + xylose | - xylose | + xylose

Parent
ΔaddA mNG-addA
ΔaddB mNG-addB
ΔrecF his-recF
ΔrecO his-recO
ΔrecR his-recR
ΔrecA his-recA
ΔrecG his-recG

C

Lagging strand discontinuity recombinational repair

Legend: AddA-AddB end-processing machinery; SSB tetramer; RecOR:RecF complex; RecA filament; RecU resolvase; RecG

Collapsed fork → Leading strand / Lagging strand

End processing

SSB loading — End resection

Pre-synapsis

RecA recruitment

Strand exchange

Post-synapsis — RecU

SUPPLEMENTARY FIGURE 7

SUPPLEMENTARY FIGURE 8

A

B

C

D

SUPPLEMENTARY FIGURE 9

Whole genome sequencing

Over-replication near the origin

End processing

Pre-synapsis

Strand exchange

Chromosome position (°)

Distance from *oriC* (kb)

SUPPLEMENTARY FIGURE 10

A

B

SUPPLEMENTARY FIGURE 11

SUPPLEMENTARY FIGURE 12

A

B

SSB complementation

C

D

E

SSB complementation

F

RecO

SSB-RecO interface

SSB tetramer

SUPPLEMENTARY FIGURE 13

A

	Phase contrast	Nucleoid	Membrane	Merge
Wild-type				
ΔrecU				

Scale bar = 5 µm

B

		Phase contrast	Nucleoid	Membrane	Merge
Leading strand	Parent				
	ΔrecU				
Lagging strand	Parent				
	ΔrecU				

Scale bar = 5 µm

SUPPLEMENTARY FIGURE 14

Presence of RecG **Absence of RecG**

Fork regression

No fork regression

↓   ↓

PriA loading at the 3'-OH group
abutting D-loop junction

Incorrect PriA loading

Helicase unwinding
towards terminus

Helicase unwinding towards origin

PriA

PriA

↓   ↓

Helicase loading on correct
lagging strand template:
productive replication restart

Incorrect helicase loading on
opposite polarity strand:
reverse-restart

SUPPLEMENTARY FIGURE 15