The basic helix-loop-helix transcription factor TCF4 recruits the Mediator Complex to activate gonadal genes and drive ovarian development

O'Neil EV, Dupont SM, Singh R, & Capel B*.

Department of Cell Biology, Duke University School of Medicine, Durham NC 27710

*Correspondence: Department of Cell Biology, Duke University School of Medicine. E-mail: blanche.capel@duke.edu

Key words: gonad, development, TCF4

Abstract

The bipotential gonad is the precursor organ to both the ovary and testis and develops as part of the embryonic urogenital system. In mice, gonadogenesis initiates around embryonic day 9.5 (E9.5), when coelomic epithelial (CE) cells overlaying the mesonephric ducts proliferate and acquire competence to differentiate into the somatic cell types of the embryonic gonad, the pre-supporting cell (Sertoli cells in the testis and granulosa cells in the ovary) and the interstitial cell lineages. While some transcription factors that drive gonadal cell fate are known, we found that basic helix-loop-helix (bHLH) binding sites are highly represented upstream of granulosa and interstitial genes. We investigated the bHLH Transcription Factor 4 (TCF4), which is expressed in GATA4+ somatic cells in both sexes prior to sex determination, maintained in ovarian pre-supporting cells and interstitial cells of both sexes, but silenced specifically in male pre-supporting cells. In a *Tcf4*$^{STOP/STOP}$ mutant mouse model that lacks the TCF4 DNA-binding domain, we found that the marker of Sertoli fate, SOX9, was higher in Sertoli cells, while markers of granulosa and interstitial fate (FOXL2 and NR2F2) were reduced and the supporting to interstitial cell ratio was altered in XX *Tcf4*$^{STOP/STOP}$ ovaries. We found that TCF4 binds the Mediator complex and chromatin remodelers to regulate expression of *Jun* and other genes in early somatic cells. Collectively, these results support the working hypothesis that TCF4 regulates a gonadal program that primes the gonad towards a female fate but is specifically silenced in male supporting cells as the testis pathway diverges.

Significant Findings

- bHLH binding sites are more abundant in genes involved in sex-determination than SOX9 or FOXL2 sites, and are 1.5 times more common upstream of granulosa than Sertoli genes.
- TCF4 is a Class I bHLH factor that is expressed in early gonadal progenitors, but silenced in Sertoli cells.
- Loss of the TCF4 binding domain results in higher levels of SOX9, suggesting that TCF4 antagonizes the Sertoli pathway.
- Loss of TCF4 binding domain reduces levels of FOXL2 and NR2F2 and leads to defects in ovary development.
- TCF4 binds with all proteins in the Mediator complex to alter the transcriptional/epigenetic landscape which may explain how it functions on target genes such as Jun.
- Our findings identify *Tcf4* as a potential upstream factor that skews the bipotential supporting cell lineage towards a granulosa cell fate.

Introduction

Differentiation of the bipotential gonad into either a testis or an ovary is a crucial process with implications for adult reproductive function. In mice, the gonad first forms from epithelial cells on the coelomic surface of the mesonephros around embryonic day 9.5 (E9.5) [1-4]. At this stage, the somatic cells of the XX and XY gonad are bipotential, with virtually identical gene expression and high expression of a common set of gonadal genes including *Wt1, Sf1, Gata4,* and *Lhx9* [5-14].

Sex determination is the process through which the gonad commits to a testis or ovary fate. The testis pathway is initiated by the expression of the Y-chromosome gene *Sry* in a subset of gonadal precursor cells between E10.5 and E12.5 [15, 16]. *Sry* activates the transcription factor *Sox9,* and SOX9 up-regulates a gene expression network that drives differentiation of these cells into Sertoli cells, the testicular supporting cell lineage [17-25]. Differentiation of XX somatic cells into the ovarian granulosa supporting cell fate is associated with expression of an alternate gene regulatory network that includes *Wnt4, Foxl2,* and *Rspo1* [26-30]. In contrast to the male pathway, the upstream factors that drive initial granulosa cell commitment are unclear [31]. This remains a substantial gap in the sex determination field.

An analysis of transcription factor (TF) binding motifs in gonadal cells showed enrichment of basic helix-loop-helix (bHLH) binding sites upstream of granulosa genes (Garcia-Moreno, unpublished data). bHLH proteins are a broad class of TFs that regulate differentiation of cells in many different tissues, including brain and muscle [32-34]. On a molecular level, bHLH TFs typically form homodimers or heterodimers and recognize E-box binding domains (CANNTG) [35, 36]. Of note, the class I bHLH factors, such as MyoD, Twist, and TCF4, are broadly expressed across tissues and are generally associated with transcriptional activation of target genes [37]. In *C. elegans,* the expression of different combinations of class I bHLH TF underly the specification of the sexually dimorphic somatic cell types in the gonad [38]. Whether bHLH factors have a role in sex determination in mammals has not been explored.

In the present study, we investigated whether the class I bHLH factor TCF4 plays a biological role in granulosa cell specification. Our results suggest that TCF4 recruits the Mediator complex to activate gonadal genes in both sexes during early gonadogenesis. While the expression of TCF4 is maintained in interstitial lineages in both XX and XY gonads, TCF4 expression is specifically silenced in XY Sertoli cells, whereas TCF4 is maintained in granulosa cells where it binds with FOXL2 and promotes the female pathway. We suggest a working model in which TCF4 binds in a large activation complex to prime the bipotential pre-supporting cell towards a granulosa cell fate in early gonad development.

Results

bHLH binding sites are ubiquitous in up- and down-regulated genes in XY and XX gonads.

To determine whether bHLH TFs might play a role in sex determination, we identified genes that were significantly (P < 0.05) up- or down-regulated (≥ 1.15-fold) between E11.5 and E13.5 in the supporting and interstitial cell lineages of XY and XX gonads using a transcriptome dataset from the lab (Supplemental Table 1) [5]. In supporting cells, 43% of the genes that were up-regulated in XX gonads contained HLH binding sites, compared to 29% of the genes that increased in XY supporting cells (Figure 1A). Meanwhile, HLH binding sites were present in 47% of the genes that were down-regulated in XX supporting cells, and in 50% of the genes that decreased in XY supporting cells. Surprisingly, we found that bHLH binding sites were more frequent in the promoter region of the top one hundred most up- or down-regulated genes in both XX and XY supporting cells compared to SOX and FOX TF binding sites (Figure 1A).

Within the interstitial cell population, HLH binding sites were present in the promoter region of 53% and 46% of up-regulated genes in XY and XX cells. Additionally, HLH binding sites were present upstream of 57% and 64% of down-regulated interstitial genes in the XY and XX gonad, respectively (Figure 1B).

Many HLH transcription factors are expressed within the early gonad (Figure 1C lists representative examples). The Class I bHLH factors *Tcf4* and *Tcf12,* which are associated with gene activation, are highly expressed in both supporting and interstitial cells (Figure 1C). Unlike *Tcf12*, *Tcf4* shows a strong down-regulation in the XY supporting cell lineage. The Class II bHLH TFs are also associated with gene activation but are either specific to the interstitial lineage (*Tcf21*) or generally expressed at lower levels than the Class I bHLH factors. Class V bHLH factors, including *Id1, Id2, Id3,* and *Bhlhe41,* are also highly expressed in the gonad but are associated with gene repression. Given the high expression levels and the activating role of TCF4, along with its significant sexually dimorphic expression pattern, we focused on investigating the role of this bHLH factor in early gonad development.

The bHLH transcription factor 4, TCF4, is highly expressed in the early gonad and down-regulated in SOX9-expressing cells.

As a role for TCF4 in gonad development has not been previously investigated, we sought to first characterize TCF4's localization in the gonad across gonadogenesis. Using immunostaining (IF), we found that TCF4 was present in the coelomic epithelium and in the broad domain of GATA4+ expressing gonadal cells in XY gonads at E10.75, just prior to sex determination (Figure 2A). At E11.5, expression of TCF4 was excluded from SOX9-expressing cells, but was maintained in XY interstitial cells that were negative for SOX9 (Figure 2B). By E12.5, TCF4 expression became restricted to the interstitial

compartment of the testis (Figure 2C), where it co-localized with the interstitial markers, VCAM and NR2F2 (Supplemental Figure 1A). This expression pattern held through E18.5, near the time of birth (Supplemental Figure 1B).

In XX gonads, TCF4 was co-expressed in GATA4+ gonadal cells at E10.75, as in the XY gonads (Figure 2D). However, while TCF4 expression was excluded from Sertoli progenitor cells in the testis, its expression was maintained in both supporting (RUNX1+) and interstitial (NR2F2+) cell types in the developing ovary (Figure 2E). By E18.5, TCF4 was highly enriched in the interstitial lineage, as in the testis, but TCF4 maintained colocalization with granulosa cell markers (Supplemental Figure 1C). Collectively, TCF4 is rapidly down-regulated in Sertoli cell progenitors, but retained in granulosa progenitors and enriched in the interstitial populations of both ovaries and testes.

Inactivation of the *Tcf4* HLH-DNA binding domain leads to dimorphic effects on the sex determination transcription factors SOX9 and FOXL2.

To understand how loss of T*cf4* affects gonad differentiation *in vivo*, we acquired a knock-out-first mutant mouse line carrying a loxP-P2A-GFP-STOP-loxP cassette in intron 17 of the *Tcf4* gene that produces a truncated form of TCF4-GFP lacking the DNA binding domain (Supplemental Figure 2A) [39]. We confirmed the presence of truncated TCF4 by western blot in protein extracts from gonads and limbs of *Tcf4$^{STOP/+}$* and *Tcf4$^{+/+}$* fetuses (Supplemental Figure 2B). Offspring homozygous for this disruption of *Tcf4* die a few days post-birth, which restricted our analysis of mutants to fetal stages. Mutant *Tcf4$^{STOP/+}$* studs were crossed with *Tcf4$^{STOP/+}$* dams, and *Tcf4$^{+/+}$* and *Tcf4$^{STOP/STOP}$* littermates were collected for downstream analysis.

First, we performed immunostaining to determine if general gonad development was disrupted by *Tcf4* inactivation. XY wildtype *Tcf4$^{+/+}$* (n = 3) and mutant *Tcf4$^{STOP/STOP}$* (n = 6) littermates showed relatively normal development of the testis, with organization of germ cells into sex cords surrounded by SOX9+ Sertoli cells (Figure 3A). Additionally, HSD3B1+ interstitial cells were present by E15.5, suggesting that interstitial cell types differentiated normally. In XX *Tcf4$^{+/+}$* (n = 5) and *Tcf4$^{STOP/STOP}$* (n = 5) fetuses, ovary development also progressed relatively normally, with the appearance and organization of FOXL2+ lined ovigerous cords (Figure 3B). However, some *Tcf4$^{STOP/STOP}$* mutant ovaries had pockets of FOXL2+ granulosa cells lacking germ cells, suggesting that there could be disruption to ovarian organization in these mutants (Supplemental Figure 3A).

Since bHLH factors specify gonadal cell types and regulate sex determination in invertebrate species [38], we used FACS analysis to test whether loss of TCF4 alters the proportions of supporting and interstitial cells in E15.5 testes and ovaries. Gonad pairs from single fetuses were dissociated, fixed, and stained for markers for supporting cells (SOX9 in XY or FOXL2 in XX) or interstitial cells (NR2F2).

Use of FACS analysis allowed us to measure cell numbers and protein abundance using fluorescent
readings on a single-cell level. The proportions, ratios, and geometric fluorescence values of *Tcf4*[+/+]
and *Tcf4*[STOP/STOP] gonads are listed in Supplemental Table 2.

XY *Tcf4*[STOP/STOP] mutant gonads (n = 3 XY gonad pairs over 3 litters) had similar proportions of
supporting and interstitial cells relative to wildtype *Tcf4*[+/+] XY gonads (n = 2 XY over 2 litters). However,
there was higher expression of SOX9 protein on a per-cell basis in Sertoli cells of *Tcf4*[STOP/STOP] mutants
(n = 10,000 cells per gonad pair, 3 gonad pairs over 3 litters) compared to *Tcf4*[+/+] littermates (n = 10,000
cells per gonad pair, 2 gonad pairs over 2 litters) (P < 0.05). There was no difference in levels of the
interstitial transcription factor, NR2F2, between wildtype and mutant gonads (Figure 3C).

Closer analysis of cell types in the ovary on the BD LSRFortessa™ Analyzer showed that mutant
Tcf4[STOP/STOP] ovaries (n = 10,000 cells per gonad pair, 4 gonad pairs over 3 litters) had a greater
supporting to interstitial cell ratio relative to XX *Tcf4*[+/+] wildtype littermates (n = 10,000 cells per gonad
pair, 2 gonad pairs over 2 litters). Further, we observed decreases in the average level of both FOXL2
and NR2F2 (P < 0.05) protein on a per-cell basis relative to wildtype *Tcf4*[+/+] ovaries (Figure 3D).

Across both sexes, TCF4 recruits the Mediator Complex and other core transcriptional
machinery to regulate gene expression.

Given the effects of TCF4 inactivation on ovarian development, we sought to further investigate TCF4's
function at E12.5 by characterizing binding partners and gene targets around the time of pre-supporting
cell commitment to the granulosa cell fate. Using the Nr5a1-GFP line, which labels both supporting and
interstitial cells at E12.5 [5, 40], we used FACS to isolate cells, immunoprecipitated TCF4, and
performed mass spectrometry (MS) (Figure 4A) (Supplemental Table 3A, B, and C). Binding partners
of TCF4 were defined as proteins that showed at least a 1.5-fold difference between the TCF4 IP and
IgG IP in each sex. In total, 2,215 proteins co-precipitated with TCF4 in dissociated XX gonadal cells,
and 3,130 proteins immunoprecipitated in dissociated XY cells as defined by a significant enrichment
(P = 0.05, rounded to two decimal places) relative to the IgG control IP. Of these, 1,774 proteins co-
precipitated with TCF4 in both XX and XY gonads (Figure 4B). Notably, TCF4 itself was enriched in the
TCF4 IP of both XX (P = 0.05) and XY cells (P = 0.04). The majority of TCF4 binding partners common
to both XX and XY gonads were not differentially abundant by sex (96%) and the HLH factor TCF21,
an interstitial-specific bHLH TF [41], immunoprecipitated with TCF4 in both XX and XY cells. Shared
TCF4 binding partners in both XX and XY cells were enriched in factors involved in positive regulation
of transcription (Figure 4C). Many key regulators of basal transcriptional processes co-precipitated with
TCF4, including all 24 of the Mediator complex proteins, as well as several classes of epigenetic
regulators, co-activators, transcription factors, and the RNA polymerase holoenzyme (Figure 4D). We

independently confirmed the presence of one core Mediator protein, Mediator 8, via western blot analysis of a TCF4 pulldown (Figure 4E). While most proteins precipitated with TCF4 in both XY and XX cells, there were also sex-specific TCF4 binders (Figure 4F). Most notably, the core transcription factor, FOXL2, which has well established roles as a crucial regulator of granulosa cell identity, co-precipitated with TCF4 in XX cells.

TCF4 regulates gonadal genes and the transcription factor *Jun* in the ovary.

Finally, ChIP-sequencing was performed using chromatin from either XX or XY *Nr5a*1-GFP cells at E12.5 to identify TCF4 targets and determine if TCF4 directly regulates genes in the granulosa network. Reads were aligned to the mm10 genome assembly, and peaks were called using MACS2 software (Supplemental Table 4). Next, stringent filtration steps were imposed on the data; peaks were filtered out if they had a Q-value < 7 or did not overlap within 1kb across all three sexed replicates (Figure 5A). While this significantly reduced the number of TCF4 targets, it enhanced the robustness of the results and minimized potential biases. In total, 96 peaks and 92 genes were identified across all three XX samples and 16 peaks and 9 genes were identified in XY samples (Figure 5B) (Supplemental Table 4).

Given that TCF4 inactivation negatively impacted ovary development, we focused on XX target genes for downstream validation. TCF4 target genes were enriched for processes in regulation of transcription and RNA metabolism (Figure 5C). TCF4 targets unique to XX cells included *Cwc22,* a splicing protein, and *Nfyc* and *Jun,* two transcription factors (Figure 5D). We measured the expression of the XX ChIP target genes *Cwc22, Nfyc*, and *Jun* in the *Tcf4$^{STOP/STOP}$* mutant gonads relative to *Tcf4$^{+/+}$* wildtype gonads. While the TCF4 putative targets *Cwc22* and *Nfyc* were not affected in mutants, we observed that expression of *Jun* as well as the granulosa-specific transcription factor *Runx1* were significantly reduced in *Tcf4$^{STOP/STOP}$* mutant ovaries compared to wildtype *Tcf4$^{+/+}$* ovaries ($P < 0.05$) (Figure 5E). In contrast, the expression of *Jun* - which was not identified as a TCF4 target in XY cells - and *Runx1* were not dysregulated in *Tcf4$^{+/+}$* mutant testes relative to *Tcf4$^{+/+}$* wildtype testes.

Although TCF4 regulates *Jun* expression in the ovary, it is unclear whether *Jun* plays a role in ovary development. Transcriptome data and single-cell RNA sequencing show that *Jun* expression becomes higher in the granulosa lineage than in the Sertoli cell lineage shortly after sex determination [5, 7]. Consistent with this, *Jun* was more highly expressed in *Tcf4$^{+/+}$* ovaries than *Tcf4$^{+/+}$* testes at E15.5 (Supplemental Figure 3B). However, JUN must be activated by phosphorylation (P-JUN) to work as a transcription factor – a process that occurs downstream of signaling cascades such as WNT signaling, which plays a pivotal role in ovary development. Thus, we measured P-JUN in whole embryonic ovaries and testes at E12.5 and in XX FACS-sorted *Nr5a1-GFP* gonadal cells across gonad development. We observed that activated P-JUN is more abundant in whole XX gonads compared to XY gonads at E12.5,

205 and that P-JUN is present in Nr5a1-GFP XX somatic gonadal cells across sex determination (from
206 E11.5 to E13.5) (Figure 5F). These results suggest that TCF4 promotes *Jun* expression, and that JUN
207 activation is elevated in XX gonads during ovary development.

Discussion

Commitment of the somatic gonad cells to the testicular Sertoli or ovarian granulosa lineage is the first
essential step in gonad differentiation. In XY fetuses, *Sry* expression activates transcriptional changes
that lead to Sertoli cell fate commitment and development of a testis [5, 7, 20, 42]. In contrast, the
upstream regulators driving the granulosa gene regulatory network and *Foxl2* expression in XX cells
remain unclear.

Across various studies, a group of genes essential for the formation of the bipotential gonad has
emerged, including *Nr5a1, Gata4, Wt1, Lhx9,* and *Emx2 [8-10, 12-14].* To search for a common element
in the regulation of these genes, we performed an investigation of TF binding sites found upstream of
early regulators of gonad differentiation. Binding sites for bHLH TFs were enriched upstream of core
gonadal genes, and were 1.5 times more common in genes associated with the ovary pathway than in
genes associated with the testis pathway [9, 10, 12, 14].

Of the many bHLH factors expressed in gonadal cells, we chose to investigate whether TCF4, which
has a known role in gene activation [43-45], plays a role in ovary development. TCF4 is highly
expressed in gonadal somatic cells before and during sex determination. However, TCF4 expression
is lost in SOX9-expressing cells, a pattern we would predict for a driver of ovarian fate.

To test whether TCF4 is involved in directing gonadal fate, we obtained a mutant mouse model for *Tcf4*
in which the binding domain is disrupted, but the rest of the protein, including the AD domains that
recruit p300 and transcriptional machinery, remain intact [16, 46]. In XY mutant gonads, the loss of
TCF4 did not lead to any obvious disruptions in testis development. Instead, we observed higher
expression of SOX9 in Sertoli cells. The fact that SOX9 binds upstream of *Tcf4* [47] and that loss of
TCF4 led to elevation of SOX9 expression, suggests an antagonistic relationship between SOX9 and
TCF4. In future studies, over-expression of TCF4 in SRY/SOX9-expressing Sertoli cells could
determine whether TCF4 is inhibitory to Sertoli cell fate.

In XX wildtype gonads, TCF4 expression is maintained in granulosa cells in the ovary with highest
abundance in the ovarian stroma. In XX *Tcf4* mutant gonads, we saw decreased expression of both the
granulosa marker FOXL2 and the ovarian stromal marker, NR2F2 at the single-cell level. Surprisingly,
the ratio of granulosa cells to stromal cells was increased in mutants, despite lower levels of FOXL2.

Further, defects in patterning and germ cell population were evident in some regions of mutant ovaries. These stronger effects are consistent with the hypothesis that TCF4 positively regulates the ovarian pathway.

To better understand how TCF4 functions in the gonad, we performed a MS analysis at E12.5 to identify TCF binding partners in XX and XY gonads. Based on mass spec analysis, binding partners identified in both XX and XY gonads included proteins associated with the Mediator Complex, the RNA polymerase complexes for initiation and elongation, epigenetic regulators, and transcription factors, such as FOXL2. These TCF4 binding partners were also reported in a paper showing that TCF4 binds with and recruits the Mediator complex to super enhancers to regulate neural stem cell differentiation [45]. If TCF4 plays a similar role in the gonad, it may bind to and open genes that establish gonadal fate and set the gonad on an ovary developmental pathway. Consistent with this, a key regulator of the ovarian pathway, FOXL2, was one of a handful of sex-specific binding partners in XX gonads. We speculate that TCF4 works in conjunction with other TF binding partners such as FOXL2 and TCF21, to specify granulosa and interstitial cell identity, respectively [37, 41, 48].

If TCF4 acts as a master regulator of gonadal genes, it is unclear why the phenotype in both XX and XY gonads is not more severe. This could be because of the complexity of the Mediator Complex in which loss of a single factor has a small impact, or, alternatively, because the *Tcf4* mutation leaves some functions of the TCF4 protein intact. We favor the latter interpretation, which needs further study.

If an effort to identify other upstream regulators of granulosa cell identity previously not recognized, we performed a TCF4 ChIP-sequencing analysis on E12.5 XX and XY gonads. Using stringent methods to confirm binding sites in XX gonads, we identified *Jun*, a transcription factor that is dysregulated in *Tcf4$^{STOP/STOP}$* mutant ovaries and known to be activated downstream of WNT signaling through the planar cell polarity pathway [49-51]. Our data show that TCF4 promotes *Jun* expression, which is preferentially activated in the embryonic ovary relative to the testis at E12.5. These findings suggest that TCF4 may enhance the availability of JUN as a key mediator of WNT-driven signaling during ovary development.

In conclusion, we hypothesize that *Tcf4* is silenced in Sertoli cells to permit their differentiation in response to *Sry* expression. Our data suggest that TCF4 recruits the Mediator complex to specific chromatin regions to maintain the progenitor state of other somatic gonadal cells and partners with proteins such as FOXL2 and TCF21 to regulate expression of granulosa and stromal factors and promote ovary differentiation (Figure 6). Many other bHLH proteins are expressed in the gonad, and further investigation of this interesting class of TFs is warranted.

Materials & Methods

Mouse strains and lines. All mice were housed in accordance with the National Institutes of Health guidelines, and experimental procedures were approved by the Duke University Medical Center Institutional Animal Care and Use Committee. For wildtype matings, CD-1 mice were obtained from Jackson Laboratories. The Tg(*Nr5a1-GFP*) (*Sf1-eGFP)* mice have been previously described [40]. The Tg(*Runx1-EGFP)*E1184Gsat/mmucd (*Runx1-eGFP)* mice were a gift from Humphrey Yao (MMRRC 010771-UCD) [52]. The *Tcf4STOP/+* mutant mouse line was a gift from Ben Philpot at UNC where they have been previously characterized [39, 53], and was maintained on a C57BL/6J (B6) background in our colony.

Timed matings. To obtain embryos at specific embryonic stages, females were placed in timed mating cages with studs. Female mice were checked daily for the presence of a vaginal plug, which was considered embryonic day 0.5. Female CD-1 mice were used for all experiments except those involving *Tcf4STOP/+ mice*. For *Tcf4STOP/+* matings, *Tcf4STOP/+* B6 studs were crossed with *Tcf4STOP/+* B6 females.

Organ collection and genotyping. Gonad-mesonephric complexes were microdissected into PBS and used for downstream applications. Tail somites were counted to ensure mice were at the appropriate developmental stage. Tails or limbs were recovered from each fetus and digested in 150 mM NaOH at 95ºC for 15 minutes and neutralized with 1M Tris HCl (pH 7.4) for genotyping. Fetuses were sexed by genotyping for the *Utx* and *Uty* genes using the following three primers (5'→3'): UTX F1: TCATGTCCATCAGGTGATGG, UTY F2: CAATGTGGACCATGACATTG, UTX/Y R: ATGGACACAGACATTGATGG. Mice from the *Tcf4STOP/+* breeding scheme were genotyped for the wildtype allele (primer sequences 5' → 3' F: GCACTTCAGGGATCGCTTA, R: CCGCCCTAATTGTTCAAAGAG) or transgenic knock-in allele (primer sequences 5' → 3': F: GCTGATCCGGAACCCTTAAGC, with the TCF4 WT R primer). PCR conditions have been previously reported [53, 54]. In the case of *Nr5a1-GFP* mice, fluorescent signal was strong enough to observe with the NIGHTSEA fluorescence detection system on the dissection microscope (NIGHTSEA, Lexington, MA), bypassing the need for genotyping.

Whole mount Immunofluorescence. Embryonic gonads were fixed for 1 hour at room temperature (RT) or overnight (O/N) at 4ºC in 4% paraformaldehyde while rocking. Samples were rinsed 3 times in PBS and then passed through an increasing methanol gradient in PBS (25%, 50%, 75%, 100%). Gonads were stored in 100% methanol O/N or longer, as needed. Samples were rehydrated stepwise into PBS and rinsed 3 times in PBS. Late-stage gonads (E15.5) were incubated in 2% Triton X-100 in PBS for 1 hour at RT to increase permeabilization. Early and late-stage gonads were then incubated in blocking

buffer (0.1% Triton X-100 for E10.5-E13.5 or 1% Triton X-100 for E15.5 and later, with 3% BSA, and 10% fetal bovine serum or horse serum [HS] in PBS) for 1 hour. Tissue was incubated with primary antibody in blocking solution O/N. Antibody details and concentrations are listed in Table 1. The following day gonads were rinsed in either 0.1% (E10.5-E14.5) or 1% Triton X-100 (E15.5 and later) in PBS and incubated with secondary antibodies (1:500) in blocking buffer O/N. Gonads were rinsed 3 times for 10 minutes in PBS and then mounted in DABCO mounting media. Images were captured with a Leica TCS SP8 confocal microscope using the associated Leica software (Leica, Wetzlar Germany). Images were processed using FIJI (ImageJ).

Mapping HLH, SOX, FOX, and COUP binding sites. HLH motif file was generated for the E-box motif (CANNTG) using the "seq2profile" command from HOMER. HLH sites genome wide were identified using the "scanMotifGenomeWide" tool against the mm9 genome [55]. The resulting bed file from this motif scan, as well as bed files from SOX9 and FOXL2 ChIP-seq studies [47, 56], were processed using the tool "annotatePeaks" to identify nearest gene transcription start sites to each genomic location present in the bed files. Next, gene lists unique to XX and XY supporting and interstitial cells were filtered by their fold-change between E11.5 and E13.5. Genes that showed a 1.15-fold change and that were statistically significant (P < 0.05) were defined as 'Increasing' or 'Decreasing' in the supporting and interstitial genes and output as a csv file (Supplemental Table 1). Annotated bed files were converted to csv and uploaded to R studio alongside the gene lists. Gene lists and motif/ChIP sites were compared for overlap using the function "inner_join" based on Gene Symbol. HLH binding sites were searched for within 1 kilobase of the gene of interest.

Fluorescence Activated Cell Sorting (FACS) Analysis. For cell analysis of E15.5 *Tcf4*$^{+/+}$ or *Tcf4*$^{STOP/STOP}$ gonads, gonad pairs from single fetuses were digested in TrypLE (Gibco, Thermo Fisher, Waltham MA) with 0.1% collagenase for 10 minutes at 37ºC. TrypLE was removed from samples and samples were then resuspended in 3% BSA in PBS and filtered through a 50-um filter. Samples were spun down at 600 g for 10 minutes at 4ºC and then fixed in 4% PFA with 0.1% Triton X-100. After a 10-minute incubation, samples were spun down again at 600 g for 10 minutes at 4ºC. Samples were left in 3% BSA in PBS at 4ºC for up to ten days before primary antibody staining. Cells were recovered, spun down at 600g for 10 minutes at 4ºC, and then incubated in blocking buffer (10% HS with 0.1% Triton X-100 in PBS) O/N with an antibody to mark interstitial cells (NR2F2) or supporting cells (SOX9/FOXL2 antibody depending on the sex of the sample) (Antibody details in Table 1). The following day samples were spun down at 600 g for 10 minutes at 4ºC and placed in blocking buffer with secondary antibody (Donkey anti-Mouse Cy3, donkey anti-rabbit AF647, or donkey anti-goat AF647) for 1 hour at RT. Samples were spun down and rinsed in 3% BSA in PBS twice before analysis. For FACS analysis, wildtype and mutant gonad suspensions were run on an BD LSRFortessa™ Analyzer in a single

session. The area of the forward side scatter (FSC-A) and side scatter area (SSC-A) were plotted to isolate cells and remove debris, and the FSC-A and FSC-H were plotted to obtain single cells. Gating conditions were established using unstained gonad cells, and single fluorophore controls (AF657 and Cy3). Voltage settings were as follows: SSC 246, FSC 360, APC 571, PE 505. Next, 5,000 cells from gonad pairs of individual, genotyped $Tcf4^{WT/WT}$ and $Tcf4^{STOP/STOP}$ fetuses were recorded. Gating was set at a signal of > $10^{2.5}$ for the 647 fluorophore and > $10^{3.1}$ for the Cy3 fluorophore. Cells with fluorescence signal greater than these gating values were identified as 'supporting' or 'interstitial' cells. The ratio of supporting: interstitial cells was then calculated and compared between wildtype and mutant conditions using a two-tailed t-test. The geometric mean, which gives the average fluorescence on a single-cell level in heterogenous populations, was also determined and compared between wildtype and mutant conditions using a two-tailed t-test.

Cell sorting. For cell sorting of *Nr5a1-GFP* gonads at E12.5, fetuses were sexed by PCR, and the gonads were dissected, pooled by sex, and dissociated for 10 minutes in TrypLE at 37ºC. After digestion, TryPLE was carefully removed, and gonads were resuspended in 10% HS in PBS. The cell suspensions were filtered through a 50-micron filter and sorted on either a BD FACSAria or Astrios Cell Sorter. GFP expression was used to sort somatic cell types, with gating set using a GFP-negative control sample by a Duke Flow Cytometry Core specialist. Cells were recovered and prepared in one of two ways for downstream analysis. For downstream IP-MS and western blot analysis, samples were spun down at 3500 g for 10 minutes at 4ºC, snap-frozen, and stored at -80ºC. For ChIP-sequencing, cells were first centrifuged at 500 g for 10 minutes at 4ºC, rinsed twice in PBS +/+, and fixed for 10 minutes at RT in freshly prepared 1% formaldehyde. Cross-linking was quenched with 2.5 M glycine and incubated for 5 minutes at 4ºC. Cells were then rinsed twice in ice cold PBS and lysed in cell lysis buffer (10 mM Tris HCl pH 7.5, 10 mM NaCl, 3 mM MgCl$_2$, 0.5% NP-40, 1 mM DTT, 1 mM PMSF, 1x Protease inhibitor) for 10 minutes at 4ºC. Nuclear pellets were spun down at 500 g for 5 minutes at 4ºC, the supernatant was removed, and samples were stored at -80ºC for later pooling

IP-MS. Frozen sorted somatic gonadal cells were pooled into three replicates of 4 million cells (± 10%). Cell pellets were lysed in a Pierce™ IP Lysis Buffer (Thermo Fisher) containing 1 mM PMSF and 1x Halt™ Protease Inhibitor Cocktail (Thermo Fisher), incubated on ice for 30 minutes, and centrifuged at 14,000 g for 10 minutes at 4ºC. Meanwhile, 50 µl of Protein G Dynabeads (Invitrogen, Waltham MA) were rinsed 3 times in PBS with 0.05% Tween-20 using a magnet bar. Beads were suspended in 237.5 µl of Conjugation buffer (20 m HEPES in PBS), and 5 µg of IgG or TCF4 antibody (Abcam, NCI-R159-6) was crosslinked to the beads using 12.5 µl of 100 mM BS3. Beads were gently vortexed and incubated for 30 minutes in crosslinking buffer. After incubation, beads were resuspended in 250 µl of Conjugation buffer and quenched with 12.5 µl of 1 M Tris HCl (pH 7.5). Beads were incubated for 15

minutes at RT and rinsed 3 times in PBS with 0.05% Tween-20. The extracted protein was then split between IgG or TCF4-conjugated beads and incubated O/N on an end-over-end rotator at 4ºC. The following day beads were rinsed in dilution buffer (10 mM Tris HCl pH 7.5 with 150 mM NaCl and 0.5 mM EDTA), dilution buffer with 1% Triton X-100, dilution buffer with 1% Triton X-100 and 360 mM NaCl, and finally in dilution buffer. Proteins were eluted at 80ºC for 10 minutes in 25 mM Tris, 50 mM NaCl, and 1% SDS and submitted to the Duke Proteomics Core for MS analysis.

Quantitative LC-MS. Samples were spiked with 1 or 2 pmol bovine casein as an internal quality control standard. The samples were brought to a final concentration of 5% SDS and reduced for 15 minutes at 80ºC. After reduction, the samples were alkylated with 20 mM iodoacetamide for 30 minutes at RT, then supplemented with 1.2% phosphoric acid and 416 µl of S-Trap (Protifi, Firport NY) binding buffer (90% MeOH/100 mM TEAB). Proteins were trapped on the S-Trap microcartridge, digested with 20 ng/µl of trypsin (Promega, Madison WI) for 1 hour at 47 ºC, and eluted sequentially with 50 mM TEAB, followed by 0.2% formic acid (FA) and 50% aceconitrile/0.2% FA. All samples were lyophilized and resuspended in 1% TFA/2% acetonitrile with 12.5 fmol/µl of yeast alcohol dehydrogenase. A study pool sample was created by combining equal volumes of each sample. Quantitative LC/MS was performed on an EvoSep One UPLC coupled to a Thermo Orbitrap Astral high-resolution, accurate-mass tandem mass spectrometer (Thermo Fisher). Each sample was eluted from the EvoTip onto a 1.5 µm EvoSep 150 µm ID x 1 5cm performance EvoSep column using the SPD30 gradient at 55ºC. Data collection on the Orbitrap Astral mass spectrophotometer was performed in a data-independent acquisition (DIA) mode of acquisition with a resolution of 240,000 for full MS scans in the m/z range of 380-980. An HCD collision energy setting of 27% was used for all MS2 scans.

Quantitative Proteomics Data Analysis. Raw data were imported into Spectronaut (Biognosis, Zurich Switzerland) and individual LC-MS data files were aligned based on the accurate mass and retention time of detected precursor and fragment ions. Relative peptide abundance was measured based on MS2 fragment ions of selected ion chromatograms of the aligned features across all runs. The MS/MS data was searched against a SwissProt M. musculus database and an equal number of reversed-sequence "decoys" for false discovery rate determination. A library free Direct DIA+ approach within Spectronaut was used to perform the database searches. Database search parameters included a fixed modification on Cys (carbamidomethyl), with variable modification on Met (oxidation) and Lys (biotin). Full trypsin enzyme rules were applied, along with 10ppm mass tolerances on precursor ions and 20ppm on product ion. Spectral annotation was set at a maximum 1% peptide false discovery rate, based on q-value calculations. Peptide homology was addressed using razor rules, where a peptide that matches to multiple different proteins is exclusively assigned to the protein with the most identified peptides. Protein homology was handled by grouping proteins that had the same set of peptides to

account for their identification. Fold-changes were calculated between the IgG and TCF4 group using a two-tailed heteroscedastic t-test on log2-transformed data. The P-Value was rounded to two decimals points. Proteins were enriched TCF4 binding partners if they met the criteria of being at least 1.5-fold higher than the IgG value with a P-value $\leq$ 0.05.

Western blot analysis. Protein was extracted from *Nr5a1-GFP* cells using Pierce™ RIPA Lysis Buffer (Thermo Fisher) with 1 mM of PMSF and 1x Halt Protease and Phosphatase inhibitor cocktail (Thermo Fisher). Protein isolates from *Nr5a1-Gfp* cells were quantified using a BCA assay. For the western blot, equivalents amount of protein [200,000 sorted cells worth of protein extract for the P-JUN western blot in *Nr5a1-GFP* cells, 20 μg of measured protein extract for the western blot measuring P-JUN and MED8 in whole gonads, or 30 μl of eluant from the IgG or TCF4 IP (approximately half the total eluant from 4 million cell IP reaction split between IgG or TCF4 antibody)] was reduced with 5x Laemmli buffer and 1 mM DTT at 95ºC for 5 minutes. Samples were loaded onto a mini-PROTEAN TGX precast gel (Bio-Rad, Hercules, CA) and proteins were separated via SDS-PAGE at a constant voltage of 100 V for 90 minutes in running buffer (25 mM Tris, 192 mM glycine, 0.1% SDS). Protein was transferred using a Trans-Blot Turbo Transfer System (Bio-Rad), following the Manufacturer's Instructions. Membrane was blocked for 1 hour at 5% milk in 0.1% Tris-Buffered saline with 0.1% Tween-20 (TBST) and incubated O/N with a polyclonal rabbit MED8, monoclonal rabbit P-JUN, or monoclonal rabbit TCF4 antibody. Blot was rinsed three times in 0.1% TBST and then put in blocking buffer with goat anti-rabbit horseradish peroxidase (HRP) conjugated secondary antibody at a 1:10,000 dilution for 1 hour at RT. Blot was rinsed three times in 0.1% TBST and developed in Super Signal West Pico Chemiluminescent Substrate (Thermo Fisher) for 5 minutes before visualization on an Amersham Imager 600. For non-IP western blots, blots were stripped in Restore PLUS Western Blot Stripping buffer for 15 minutes at RT. Blots were blocked for 1 hour in 5% milk in 0.1% TBST again and then either stained with Colloidal Gold Total Protein Stain (Bio-Rad) or incubated with a polyclonal goat anti-GAPDH antibody (1:5,000) O/N. The following day, the blot was rinsed three times in 0.1% TBST and incubated in a donkey anti-goat horseradish peroxidase (HRP) conjugated secondary antibody. Development and imaging were repeated as previously described.

ChIP Sequencing. Nuclear pellets from approximately 1.75 million (± 10%) sorted *Nr5a*1-GFP cells were pooled and placed in a MNase reaction buffer (10mM Tris HCl pH 7.5, 10mM NaCl, 3 mM MgCl2, 1 mM Cacl2, 0.2% NP-40, 1 mM DTT, 1mM PMSF, 1x protease inhibitor). Nuclei were centrifuged at 500 g for 5 minutes at 4ºC and placed in 200 μl of fresh MNAse reaction buffer. Next, 55 units of MNase was added to each tube and nuclear pellets were incubated for 15 minutes at 37ºC with vortexing every five minutes. Enzymatic DNA fragmentation was stopped by adding 20 μl of 0.5 M EDTA. Samples were centrifuged at 9400 g for 10 minutes at 4ºC and placed in 75 μl of ChIP sonication buffer (50 mM

Tris HCl pH 7.9, 10 mM EDTA, 1% SDS, 1mM PMSF, 1x protease inhibitor). Samples were left on ice for 10 minutes and 75 µl of ChIP dilution buffer (16.7 mM Tris HCl pH 7.9, 1.1% Triton X-100, 167 mM NaCl, 1.2 mM EDTA, 1 mM PMSF). DNA was sonicated four times for 20 seconds with 30 seconds of rest. Samples were centrifuged at 13000g for 10 minutes at 4ºC. Next, 600 µl of ChIP dilution buffer was added to samples, 2% of the sample volume was collected for input, and 5 µg of TCF4 antibody was added for overnight incubation at 4ºC on an end-over-end rotator. The following day, 15 µl of Dynabeads Protein G magnetic beads were added to each tube. Samples were incubated for two hours on an end-over-end rotator. Using a magnetic bar, the supernatant was removed and beads were sequentially rinsed for 10 minutes in low salt buffer (20 mM Tris HCl pH 7.9, 100 mM NaCl, 2 mM EDTA, 0.1% SDS, 1% Triton X-100), high salt buffer (20 mM Tris HCl pH 7.9, 500 mM NaCl, 2 mM EDTA, 0.1% SDS, 1% Triton X-100), LiCl Buffer (10 mM Tris HCl pH 7.9, 250 mM LiCl, 1% NP-40, 1% deoxycholate acid, 1 mM EDTA) and TE buffer. Next, 200 µl of elution buffer (1% SDS, 0.1 M NaHCO$_3$) was added and samples were left at 65ºC for 40 minutes to elute DNA. At this point, input DNA was thawed, brought to 200 µl with elution buffer, and processed in parallel with samples. Samples were recovered, 8 µl of 5 M NaCl (per 200 µl of elution buffer) was added to samples, and samples were left at 55ºC overnight to de-crosslink the DNA. The next day, 14 µl of Proteinase K solution was added to each sample (4 µl of 0.5 M EDTA, 8 µl of 1 M Tris HCl pH 6.8, 2 µl of 20 mg/ml Proteinase K) before a 2-hour incubation at 55ºC, and DNA was eluted using the Qiagen MinElute on column PCR Purification kit (Qiagen, Hilden, Germany). DNA was eluted in 10 ul of TE buffer.

ChIP sequencing data analysis. Samples were submitted to BGI Genomics for sequencing and analysis. Briefly, immunoprecipitated DNA was purified and checked for quality using a Fragment Analyzer. Sequencing libraries were prepared by adding ligating sequencing adaptors and sequenced on the DNBSEQ platform to generate 50 bp paired-end reads. After adapter trimming and removal of low-quality reads, an average of approximately 25 million high-quality reads were obtained per sample. Clean reads were aligned to the reference genome mm10 using SOAPaligner/SOAP2 software with two mismatches. Only uniquely mapped reads with a mapping quality score (MAPQ) greater than 20 were retained. Peak calling was performed using MACS1.4.2 with a Q-value cutoff of < 0.05 relative to the input control. Individual peak lists were generated for the six replicates (three sexed replicates). We then selected high-confidence peaks (Q-value < 10^-7) and designed a replicate-matching algorithm to only select peaks that were consistently observed across the three sexed replicates. Specifically, we computed clusters of peaks such that each cluster a) contains at least one peak from each replicate, and b) each peak in the cluster is within 1 kb of at least one peak from another replicate. This was achieved by constructing a tri-partite graph across three replicates where nodes are peaks, edges connect cross-replicate peaks within 1 kb and then identifying connected graph sub-components that

span all replicates. Peaks not assigned to any cluster were discarded. The Python implementation of the matching algorithm will be made available to the research community as an open-source package on Github. The peaks called after stringent filtration as well as the peak lists for each individual replicate are listed in Supplementary Table 4. Individual bigWig files are available for download at DOI 10.17605/OSF.IO/H6D2M. Raw FASTQ files will be deposited in GEO and made publicly available upon publication.

qPCR analysis. Tissue was dissected and run through a QiaShredder column in Buffer RLT (Qiagen). RNA was isolated using the MinElute RNA Kit (Qiagen), including an on-column DNAse digestion. The RNA was quantified using a Nanodrop and the purity was assessed using A260/280 values. Next, 50 ng of RNA was reverse-transcribed into cDNA using the Verso cDNA Synthesis kit. Parallel reactions were run without enzyme to assess genomic contamination. Reverse transcription was performed following manufacturers' instructions. Real-time PCR was then performed using a QuantStudio 6 Flex Real-Time System (ThermoFisher) with SsoAdvanced Universal SyberGreen Supermix (BioRad). PCR conditions were: activation, 95°C for 15 min; 40 cycles of 95°C for 15 s; 60°C for 60 s. Primers were synthesized by Integrated DNA Technologies (Primer sequences in Table 2). Technical triplicates were used to generate the cycle threshold for each gene in each sample. Melt curve analysis was performed to verify PCR product specificity. ΔCT values were calculated as the difference between the cycle threshold (CT) of the gene of interest and the housekeeping gene (*Actb)*. The housekeeping gene *Actb* is highly and stably expressed during early gonad development [5, 7]. Statistics were calculated using the ΔCT values. For visualization, fold change was calculated relative to the reference gene. Standard error was calculated by propagating the standard deviations of the gene of interest and housekeeping gene using the formula $SE_{\Delta\Delta CT} = \sqrt{(SD\_GOI/\sqrt{n})^2)+(SD\_HK/\sqrt{n})^2}$. Fold change was plotted as $2^{-\Delta\Delta CT} \pm$ propagated error.

Table 1. Key Resources.

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Gene (*M. musculus*)	Transcription factor 4 (Tcf4)	GenBank	MGI:98506	
Strain, strain background (*Mus musculus*)	Crl:CD1(ICR)	Charles River	Strain code: 022	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (*M. musculus*)	C57BL/6J	Jackson Laboratory	Stock #:000664	
Genetic reagent (*M. musculus*)	Tg(Nr5a1/EGFP)1Klp	PMID:12351700	MGI:5493455	
Genetic reagent (*M. musculus*)	Tg(Tcf4$^{STOP/+}$)	PMID: 32765228		
Antibody	Smooth muscle alpha actin (aSMA) (FITC-conjugated mouse monoclonal)	Sigma-Aldrich	F3777	1:500
Antibody	Endostatin/Collagen 18 (goat polyclonal)	R&D Systems	AF570	1:500
Antibody	FOXL2 (rabbit polyclonal)	Gift from Dagmar Wilhelm		1:250
Antibody	GATA4 (mouse monoclonal IgG$_{2\alpha}$)	Santa Cruz Biotechnology	Sc-25310	1:250
Antibody	GFP (chicken polyclonal)	Abcam	Ab13970	1:1000
Antibody	Phospho-c-Jun (Ser73) (rabbit monoclonal)	CellSignaling Technology	(D47G9)	1:1000
Antibody	Phospho-JUN (Ser73)	Proteintech	4A18	1:1000
Antibody	MED8 (rabbit polyclonal)	Proteintech	12182-1-AP	1:1000
Antibody	NR2F2 (mouse monoclonal IgG$_{2\alpha}$)	Abcam	Ab41859	1:200
Antibody	NR5A2 (goat polyclonal)	Santa Cruz Biotechnology	Sc-21132	1:500
Antibody	SOX9 (goat polyclonal)	R&D Systems	AF3075	1:500
Antibody	TCF4 (rabbit monoclonal)	Abcam	Ab217668	1:500

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Antibody	TRA98/GCNA1 (rat monoclonal)	Abcam	Ab82527	1:500
Antibody	AF488 Anti-Mouse (donkey polyclonal)	Jackson ImmunoResearch	715-545-150	1:500
Antibody	AF488 anti-Rat (donkey polyclonal)	Life Technologies	A-21208	1:500
Antibody	AF488 anti-Chicken (donkey polyclonal)	Jackson ImmunoResearch	703-545-155	1:500
Antibody	Cy3 anti-Goat (donkey polyclonal)	Jackson ImmunoResearch	705-165-147	1:500
Antibody	Cy3 anti-Mouse (donkey polyclonal)	Jackson ImmunoResearch	715-165-150	1:500
Antibody	AF647 anti-Rabbit (donkey polyclonal)	Jackson ImmunoResearch	711-605-152	1:500

499 **Table 2. qPCR Primer Sequences.**

Gene	Forward/Reverse Sequences	Amplicon Size	Origin of Primers
Cwc22	5' – CCAGAGACAACCCACGCA ATA C - 3' 5' – TGGTTCTGGGCTACGATGACC – 3'	127 nt	Origene SKU MP202029
Nfyc	5' – CCAGACACTTGCTACCAATGCC – 3' 5' – AGGCATGGT GAC TTG CTG GAT C -3'	118 nt	Origene SKU MP220268
Jun	5' – TGA TCA TCC AGT CCA GCA ATG -3' 5' – TGC TCG TCG GTC ACG TTC T -3'	85 nt	Galán-Cobo et al. 2015 [57]
Runx1	5' – CACCGTCTTTACAAATCCGCCAC– 3' **5' – CGCTCGGAAAAGGACAAACTCC - 3'**	141 nt	Lambert et al. 2024 [58]

Acknowledgments

The authors would like to thank Ben Philpot at UNC Chapel Hill for generously providing the

Tcf4$^{STOP/+}$mouse line. The authors thank the past and present members of the Capel Lab for their

support, especially Megan Harward for providing animal care and maintenance and Talia Hatkevich for

assistance in FACS analysis. Finally, the authors would like to thank Erik Soderblom, Tricia Ho, and

Greg Waitt for their technical assistance with mass spec sample analysis and interpretation. The

authors also appreciate the Staff at the Duke Light Microscopy Core Facility and the Duke Cancer
Institute Flow Cytometry Shared Resource for their excellent technical support.

Funding Sources

511 EO, SD, and B.C. are currently supported by a grant from the National Institutes of Health
512 (1R01HD090050-0 to BC).

Author contributions

515 EVO, SMD, and BC conceptualized and designed research; EVO and BC performed research; EVO,
516 SMD, RH, and BC analyzed data; and EVO and BC wrote the manuscript. All authors edited and
517 approved the final manuscript.

Competing Interests

520 None declared.

REFERENCES

1. Karl, J. and B. Capel, *Three-dimensional structure of the developing mouse genital ridge.* Philos Trans R Soc Lond B Biol Sci, 1995. **350**(1333): p. 235-42.
2. Karl, J. and B. Capel, *Sertoli cells of the mouse testis originate from the coelomic epithelium.* Dev Biol, 1998. **203**(2): p. 323-33.
3. Upadhyay S, L.J., Zombini L, *The role of the mesonephros in the development of indifferent gonads and ovaries of the mouse.* Annales de biologie animale, biochimie, biophysique, 1979. **19**(4 B): p. 1179-1196.
4. Wartenberg, H., *Human testicular development and the role of the mesonephros in the origin of a dual Sertoli cell system.* Andrologia, 1978. **10**(1): p. 1-21.
5. Jameson, S.A., et al., *Temporal transcriptional profiling of somatic and germ cells reveals biased lineage priming of sexual fate in the fetal mouse gonad.* PLoS Genet, 2012. **8**(3): p. e1002575.
6. Munger, S.C. and B. Capel, *Sex and the circuitry: progress toward a systems-level understanding of vertebrate sex determination.* Wiley Interdiscip Rev Syst Biol Med, 2012. **4**(4): p. 401-12.
7. Stevant, I., et al., *Dissecting Cell Lineage Specification and Sex Fate Determination in Gonadal Somatic Cells Using Single-Cell Transcriptomics.* Cell Rep, 2019. **26**(12): p. 3272-3283 e3.
8. Sasaki, K., et al., *The embryonic ontogeny of the gonadal somatic cells in mice and monkeys.* Cell Rep, 2021. **35**(5): p. 109075.
9. Hu, Y.C., L.M. Okumura, and D.C. Page, *Gata4 is required for formation of the genital ridge in mice.* PLoS Genet, 2013. **9**(7): p. e1003629.
10. Kreidberg, J.A., et al., *WT-1 is required for early kidney development.* Cell, 1993. **74**(4): p. 679-91.
11. Luo, X., Y. Ikeda, and K.L. Parker, *A cell-specific nuclear receptor is essential for adrenal and gonadal development and sexual differentiation.* Cell, 1994. **77**(4): p. 481-90.
12. Miyamoto, N., et al., *Defects of urogenital development in mice lacking Emx2.* Development, 1997. **124**(9): p. 1653-64.
13. Birk, O.S., et al., *The LIM homeobox gene Lhx9 is essential for mouse gonad formation.* Nature, 2000. **403**(6772): p. 909-13.
14. Luo, X., et al., *Steroidogenic factor 1 is the essential transcript of the mouse Ftz-F1 gene.* Mol Endocrinol, 1995. **9**(9): p. 1233-9.
15. Gubbay, J., et al., *A gene mapping to the sex-determining region of the mouse Y chromosome is a member of a novel family of embryonically expressed genes.* Nature, 1990. **346**(6281): p. 245-50.
16. Hiramatsu, R., et al., *A critical time window of Sry action in gonadal sex determination in mice.* Development, 2009. **136**(1): p. 129-38.
17. Colvin, J.S., et al., *Male-to-female sex reversal in mice lacking fibroblast growth factor 9.* Cell, 2001. **104**(6): p. 875-89.
18. Li, Y., M. Zheng, and Y.F. Lau, *The sex-determining factors SRY and SOX9 regulate similar target genes and promote testis cord formation during testicular differentiation.* Cell Rep, 2014. **8**(3): p. 723-33.
19. Morais da Silva, S., et al., *Sox9 expression during gonadal development implies a conserved role for the gene in testis differentiation in mammals and birds.* Nat Genet, 1996. **14**(1): p. 62-8.
20. Sekido, R. and R. Lovell-Badge, *Sex determination involves synergistic action of SRY and SF1 on a specific Sox9 enhancer.* Nature, 2008. **453**(7197): p. 930-4.
21. Bullejos, M. and P. Koopman, *Spatially dynamic expression of Sry in mouse genital ridges.* Dev Dyn, 2001. **221**(2): p. 201-5.
22. Hacker, A., et al., *Expression of Sry, the mouse sex determining gene.* Development, 1995. **121**(6): p. 1603-14.
23. Koopman, P., et al., *Male development of chromosomally female mice transgenic for Sry.* Nature, 1991. **351**(6322): p. 117-21.
24. Sekido, R., et al., *SOX9 is up-regulated by the transient expression of SRY specifically in Sertoli cell precursors.* Dev Biol, 2004. **274**(2): p. 271-9.
25. Moniot, B., et al., *The PGD2 pathway, independently of FGF9, amplifies SOX9 activity in Sertoli cells during male sexual differentiation.* Development, 2009. **136**(11): p. 1813-21.
26. Kim, Y., et al., *Fgf9 and Wnt4 act as antagonistic signals to regulate mammalian sex determination.* PLoS Biol, 2006. **4**(6): p. e187.

27. Ottolenghi, C., et al., *Loss of Wnt4 and Foxl2 leads to female-to-male sex reversal extending to germ cells.* Hum
Mol Genet, 2007. **16**(23): p. 2795-804.
28. Tomizuka, K., et al., *R-spondin1 plays an essential role in ovarian development through positively regulating Wnt-*
4 signaling. Hum Mol Genet, 2008. **17**(9): p. 1278-91.
29. Parma, P., et al., *R-spondin1 is essential in sex determination, skin differentiation and malignancy.* Nat Genet, 2006.
38(11): p. 1304-9.
30. Nicol, B., et al., *RUNX1 maintains the identity of the fetal ovary through an interplay with FOXL2.* Nat Commun,
2019. **10**(1): p. 5116.
31. Yao, H.H., *The pathway to femaleness: current knowledge on embryonic development of the ovary.* Mol Cell
Endocrinol, 2005. **230**(1-2): p. 87-93.
32. Dennis, D.J., S. Han, and C. Schuurmans, *bHLH transcription factors in neural development, disease, and*
reprogramming. Brain Res, 2019. **1705**: p. 48-65.
33. Rudnicki, M.A., et al., *The molecular regulation of muscle stem cell function.* Cold Spring Harb Symp Quant Biol,
2008. **73**: p. 323-31.
34. Schlaeger, T.M., et al., *Decoding hematopoietic specificity in the helix-loop-helix domain of the transcription factor*
SCL/Tal-1. Mol Cell Biol, 2004. **24**(17): p. 7491-502.
35. de Martin, X., R. Sodaei, and G. Santpere, *Mechanisms of Binding Specificity among bHLH Transcription Factors.*
Int J Mol Sci, 2021. **22**(17).
36. Torres-Machorro, A.L., *Homodimeric and Heterodimeric Interactions among Vertebrate Basic Helix-Loop-Helix*
Transcription Factors. Int J Mol Sci, 2021. **22**(23).
37. Wang, L.H. and N.E. Baker, *E Proteins and ID Proteins: Helix-Loop-Helix Partners in Development and Disease.* Dev
Cell, 2015. **35**(3): p. 269-80.
38. Sallee, M.D., H.E. Littleford, and I. Greenwald, *A bHLH Code for Sexually Dimorphic Form and Function of the C.*
elegans Somatic Gonad. Curr Biol, 2017. **27**(12): p. 1853-1860 e5.
39. Kim, H., et al., *Rescue of behavioral and electrophysiological phenotypes in a Pitt-Hopkins syndrome mouse model*
by genetic restoration of Tcf4 expression. Elife, 2022. **11**.
40. Stallings, N.R., et al., *Development of a transgenic green fluorescent protein lineage marker for steroidogenic factor*
1. Mol Endocrinol, 2002. **16**(10): p. 2360-70.
41. Cui, S., et al., *Disrupted gonadogenesis and male-to-female sex reversal in Pod1 knockout mice.* Development,
2004. **131**(16): p. 4095-105.
42. Neirijnck, Y., et al., *Single-cell transcriptomic profiling redefines the origin and specification of early adrenogonadal*
progenitors. Cell Rep, 2023. **42**(3): p. 112191.
43. Henthorn, P., M. Kiledjian, and T. Kadesch, *Two distinct transcription factors that bind the immunoglobulin*
enhancer microE5/kappa 2 motif. Science, 1990. **247**(4941): p. 467-70.
44. Pscherer, A., et al., *The helix-loop-helix transcription factor SEF-2 regulates the activity of a novel initiator element*
in the promoter of the human somatostatin receptor II gene. EMBO J, 1996. **15**(23): p. 6680-90.
45. Quevedo, M., et al., *Mediator complex interaction partners organize the transcriptional network that defines*
neural stem cells. Nat Commun, 2019. **10**(1): p. 2669.
46. Bullejos, M. and P. Koopman, *Delayed Sry and Sox9 expression in developing mouse gonads underlies B6-Y(DOM)*
sex reversal. Dev Biol, 2005. **278**(2): p. 473-81.
47. Rahmoun, M., et al., *In mammalian foetal testes, SOX9 regulates expression of its target genes by binding to*
genomic regions with conserved signatures. Nucleic Acids Res, 2017. **45**(12): p. 7191-7211.
48. Murre, C., et al., *Interactions between heterologous helix-loop-helix proteins generate complexes that bind*
specifically to a common DNA sequence. Cell, 1989. **58**(3): p. 537-44.
49. Park, M. and R.T. Moon, *The planar cell-polarity gene stbm regulates cell behaviour and cell fate in vertebrate*
embryos. Nat Cell Biol, 2002. **4**(1): p. 20-5.
50. Waudby, C.A., et al., *An intrinsic temporal order of c-JUN N-terminal phosphorylation regulates its activity by*
orchestrating co-factor recruitment. Nat Commun, 2022. **13**(1): p. 6133.
51. Smeal, T., et al., *Oncogenic and transcriptional cooperation with Ha-Ras requires phosphorylation of c-Jun on*
serines 63 and 73. Nature, 1991. **354**(6353): p. 494-6.
52. Gong, S., et al., *A gene expression atlas of the central nervous system based on bacterial artificial chromosomes.*
Nature, 2003. **425**(6961): p. 917-25.

53.   Kim, H., et al., *Region and Cell Type Distribution of TCF4 in the Postnatal Mouse Brain.* Front Neuroanat, 2020. **14**:
626   p. 42.
627   54.   Shpargel, K.B., et al., *KDM6 demethylase independent loss of histone H3 lysine 27 trimethylation during early*
628   *embryonic development.* PLoS Genet, 2014. **10**(8): p. e1004507.
629   55.   Heinz, S., et al., *Simple combinations of lineage-determining transcription factors prime cis-regulatory elements*
630   *required for macrophage and B cell identities.* Mol Cell, 2010. **38**(4): p. 576-89.
631   56.   Nicol, B., et al., *Genome-wide identification of FOXL2 binding and characterization of FOXL2 feminizing action in*
632   *the fetal gonads.* Hum Mol Genet, 2018. **27**(24): p. 4273-4287.
633   57.   Galan-Cobo, A., et al., *Overexpression of AQP3 Modifies the Cell Cycle and the Proliferation Rate of Mammalian*
634   *Cells in Culture.* PLoS One, 2015. **10**(9): p. e0137692.
635   58.   Lambert, J., et al., *Network-based prioritization and validation of regulators of vascular smooth muscle cell*
636   *proliferation in disease.* Nat Cardiovasc Res, 2024. **3**(6): p. 714-733.

637
638

Figure Legends.

Figure 1. Helix-loop-helix transcription factor binding sites are ubiquitous in dynamically regulated gonadal genes. A) In supporting cells, HLH binding motifs are present in the promoter region of 29% and 43% of genes that increase in XY and XX cells, respectively, and 50% and 47% of genes that decrease in XY and XX cells. Among the top 100 most up- and down-regulated genes in these sets, HLH binding sites are more frequent than SOX or FOX binding sites. B) In interstitial cells, HLH binding sites are present in 53% and 46% of genes that increase in XY and XX cells, respectively. HLH binding sites are enriched in the top 100 most up-regulated genes and occur more frequently than SOX or FOX binding sites. Additionally, HLH binding sites are upstream of 57% and 64% of genes that decrease in XY and XX interstitial cells and are present in nearly 75% of the top 100 most down-regulated interstitial genes. C) Basic helix-loop-helix transcription factors show sex- and cell-type specificity in the gonad (Jameson et al. 2012).

Figure 2. Expression of the HLH transcription factor TCF4 is lost in Sertoli cells but maintained in granulosa and interstitial cell types. A) At E10.75, TCF4 co-localizes with GATA4-expressing gonadal cells in XY gonads. B) At E11.5, TCF4 is lost in SOX9-expressing Sertoli cells in the XY testis, (shown with arrows in B' and B'') but is maintained in cells throughout the gonad which are forming sex cords (dotted circle in B' and B''). C) In the XY gonad, TCF4 is restricted to the interstitial cell population (shown with dotted lines in C' and C''). D) in the XX gonad, TCF4 is co-expressed in GATA4-expressing cells. E) By E12.5, TCF4 is expressed in both RUNX1 expressing granulosa cells and NR2F2+ interstitial cells. Dashed line highlights the boundary between ovarian supporting cells and interstitial cells. Scale bar represents 100 μm. Microscopy images are stitched together from individual panels.

Figure 3. TCF4 does not disrupt gonad differentiation but does affect expression of master regulators of sex determination and cell type ratios in the ovary. A) Immunofluorescence imaging of wildtype *Tcf4+/+* and mutant *Tcf4STOP/STOP* XY gonads. Images were taken using identical microscope settings. Representative images shown. B) Immunofluorescence imaging of wildtype *Tcf4+/+* and mutant *Tcf4STOP/STOP* XX gonads. Images were taken using identical microscope settings. Representative images shown. Microscopy images are stitched together from individual panels. C) Gonads were digested and run on a cell analyzer. While ratios of supporting: interstitial cell types did not change, SOX9 was more abundant (P < 0.05) on a per-cell bases in *Tcf4STOP/STOP* XY gonads (n=3 gonad pairs) relative to *Tcf4+/+* gonads (n=2 gonad pairs). Meanwhile, NR2F2 was not different between samples. D) Mutant ovaries had a higher supporting: interstitial cell ratio (P < 0.05). Additionally, FOXL2 and NR2F2 were less abundant (P < 0.05) on a per-cell bases in *Tcf4STOP/STOP* XX gonads (n = 4 gonad pairs) relative to *Tcf4+/+* gonads (n = 2 gonad pairs).

Figure 4. TCF4 binds core transcriptional regulators including the Mediator complex in both XX and XY sorted cells. A) *Nr5a1-Gfp* cells were sorted, immunoprecipitated for TCF4, and sent for mass spectrometry analysis to identify TCF4 binding partners. B) In total, 1,774 proteins were identified as TCF4 co-binders common to XY and XX cells. C) Proteins bound by TCF4 were enriched for processes such as positive transcription initiation and regulation of transcription, suggesting that TCF4 is a pro-transcriptional regulator in the gonad. D) Several protein complexes were identified in our mass spectrometry dataset in both XY and XX cells. Notably, the Mediator complex, as well as members of transcriptional elongation, the RNA polymerase holoenzyme, the MLL complex, and transcriptional initiation machinery. Proteins organized in a circle are known complex proteins, and lines indicate interactions. E) XY and XX *Nr5a1-Gfp* cells were sorted and western blot was performed to show the presence of Mediator 8 in somatic gonadal cells (20 μg input). Presence of Med8 was also shown in the western blot of IP'd TCF4 from XY cells (2 million cells / IP). **F**) Additionally, TCF4 co-precipitated with sex-specific proteins, including FOXL2, STAT1, and NFYB in XX cells and STAT3, LMX1B, and KDM4A in XY cells. These proteins were associated with translational machinery in XY cells, and metabolic processing in XX cells.

Figure 5. Gene targets of TCF4 include *Jun* and other transcriptional regulators. A) *Nr5a1-Gfp* cells were sorted and a ChIP-sequencing experiment was performed to identify TCF4 target genes in XX and XY chromatin. B) In total, 92 genes were identified as TCF4 targets in XX cells, while only 9 genes were identified in XY cells. C) TCF4 targets were enriched for processes including negative and positive regulation of gene transcription and RNA metabolism. D) TCF4 directly binds to the first exon of the splicing factor, *Cwc22,* the transcription factor, *Nfyc,* and across the gene body of the transcription factor *Jun.* E) *Jun,* a TCF4 ChIP target identified in XX cells, and *Runx1* expression is decreased in *Tcf4^STOP/STOP^* ovaries relative to *Tcf4^+/+^* ovaries from wildtype littermates. In contrast, *Jun* was not identified as a TCF4 ChIP target in XY cells and was not dysregulated in *Tcf4^STOP/STOP^* mutant testes. Gene expression was normalized to the housekeeping gene *Actb.* Expression is displayed as $\log_2$-fold change; statistics were calculated based on ΔCT value comparing wildtype and mutant gonads of the same sex. F) The active, phosphorylated form of JUN (P-JUN) is more abundant in E12.5 ovaries compared to E12.5 testes. By E15.5, P-JUN is less abundant in the embryonic ovary than testis. Additionally, P-JUN is present in sorted XX gonadal *Nr5a1-Gfp* cells across early gonad development (E10.5-E13.5) (approximately 200,000 cells per lane). Total protein stain showed similar loading of protein across the different samples.

Figure 6. Working model for the role of TCF4 in gonad differentiation. TCF4 drives a gonadal program, likely through recruitment of the Mediator complex and epigenetic factors, that advances

ovarian fate and is specifically silenced in Sertoli supporting cells as these pathways diverge. Figure created with Biorender.

Figure S1. Expression of TCF4 is maintained in gonadal granulosa and interstitial cells through late gestation. A) At E12.5, TCF4 co-localizes with VCAM- and NR2F2- expressing interstitial cells in XY gonads. B) At E18.5, TCF4 remains restricted to the testicular interstitial cells and absent in SOX9-expressing Sertoli cells. C) At E18.5, TCF4 is broadly expressed across both granulosa and interstitial cell lineages in the oary. Scale bar represents 100 μm. Microscopy images are stitched together from individual panels.

Figure S2. The Tcf^{STOP} allele is unable to bind DNA. A) The Tcf^{STOP} allele introduces a premature STOP codon upstream of the bHLH domain in exon 18. B) Western blot analysis shows that $Tcf^{STOP/+}$ mice express a truncated form of TCF4 without the bHLH domain that is absent in $Tcf^{+/+}$ wildtype littermates.

Figure S3. Minor defects in $Tcf^{STOP/STOP}$ mutant ovaries. A) One $Tcf^{STOP/STOP}$ mutant ovary displayed defects in patterning, as germ cells were present in the posterior portion of the ovary (white arrowheads) but were absent in the anterior portion. B) The transcription factor *Jun* is expressed in $Tcf^{+/+}$ wildtype testes at about half the level of $Tcf^{+/+}$ wildtype ovaries.

Fig1

A **Supporting cells**

Increasing ⬆

Genes with HLH binding motif

XY | 22 | XX

Pdia5... | *Osr2, Lgr5...*

130 / 447 genes
29%

135 / 312 genes
43%

TF binding sites in 100 most-upregulated genes

■ XY ■ XX

SOX FOX HLH

Decreasing ⬇

Genes with HLH binding motif

XY | 44 | XX

Emx2, Meis1... | *Gpc6...*

155 / 308 genes
50%

93 / 198 genes
47%

TF binding sites in 100 most-downregulated genes

■ XY ■ XX

SOX FOX HLH

B **Interstitial cells**

Increasing ⬆

Genes with HLH binding motif

XY | 70 | XX

Ptch1, Sorcs2... | *Lgr5...*

115 / 216 genes
53%

64 / 140 genes
46%

TF binding sites in 100 most-upregulated genes

■ XY ■ XX

SOX FOX HLH

Decreasing ⬇

Genes with HLH binding motif

XY | 42 | XX

Wt1, Emx2... | *Fli1...*

70 / 122 genes
57%

59 / 92 genes
64%

TF binding sites in 100 most-downregulated genes

■ XY ■ XX

SOX FOX HLH

C

Class I HLH

Tcf4 *Tcf12* *Twist1* *Tcf3*

Class II HLH

Tcf21 *Myc* *Srebf2* *Neurod2*

Class V HLH

Id1 *Id2* *Id3* *Bhlhe41*

Log intensity value / Stage (11.5, 12.5, 13.5)

— ♂ Supporting
— ♀ Supporting
-- ♂ Interstitial
-- ♀ Stromal

Fig2

Fig3

C

Sertoli : Interstitial Cell Ratio

Ratio — WT, Mut
n = 2 fetuses n = 3 fetuses

Geometric mean (Fluorescence Signal)

Relative Units — SOX9, NR2F2
*
WT, Mut
n = 2 fetuses n = 3 fetuses

D

Granulosa : Interstitial Cell Ratio

Ratio — WT, Mut
*
n = 2 fetuses n = 4 fetuses

Geometric mean (Fluorescence Signal)

Relative Units — FOXL2, NR2F2
* **
XX WT, XX Mut
n = 2 fetuses n = 4 fetuses

A WT Testes — SOX9 / HSD3B1 / Nuclei
Mutant Testes — SOX9 / HSD3B1 / Nuclei

B WT Ovaries — FOXL2 / NR2F2 / Nuclei
Mutant Ovaries — FOXL2 / NR2F2 / Nuclei

Fig4

A E12.5 XY or XX gonads → 1.75 million cells / replicate → Chromatin shearing, IP → Input vs TCF4-bound

Data Analysis Workflow
Quality Control →
Peak Calling →
Data filtering:
- Peaks have a –log10 Q-Value ≥ 7
- Peak identified (within 1 kb) across three biological XX or XY replicates

Fig5

B Genes nearby TCF4 peaks
XY cells: 16 peaks, 9 genes
XX cells: 96 peaks, 92 genes
Overlap: 3 genes

C Target Genes in XX Cells
Negative reg. of transcription by RNA polymerase II
Positive reg. of transcription DNA-templated
Positive reg. of nucleic acid-templated transcription
Positive reg. of transcription by RNA polymerase II
Positive reg. of RNA biosynthetic proc.
Positive reg. of nucleobase-containing compound metabolic proc.
Reg. of transcription by RNA polymerase II
Reg. of transcription DNA-templated
Reg. of nucleic acid-templated transcription
Reg. of RNA metabolic proc.
Reg. of RNA biosynthetic proc.
Transcription DNA-templated
Nucleic acid-templated transcription
Reg. of nucleobase-containing compound metabolic proc.
RNA metabolic proc.

N. of Genes: 20, 25, 30, 35, 40
-log10(FDR): 6.5, 7.0, 7.5, 8.0, 8.5
Fold Enrichment: 0, 2, 4

D
10 kb, 84, XX, *Cwc22*
10 kb, 54, XX, *Nfyc*
10 kb, 43, XX, *Jun*
TCF4 binding

E Gene expression of TCF4 ChIP targets in Tcf4+/+ and Tcf4STOP/STOP gonads at E15.5
Fold Change Relative to *Actb*
Cwc22 (NS), *Nfyc* (NS), *Jun* (P-Value 0.003), *Runx1* (P-Value 0.013)
TCF4 ChIP Targets in XX cells only
Tcf4+/+ ovaries, Tcf4STOP/STOP ovaries, Tcf4+/+ testes, Tcf4STOP/STOP testes

F Whole gonads at E12.5 — Nr5a1-GFP+ XX cells
kDa: 70, 55, 35
XX, XY, E11.5, E12.5, E13.5
P-JUN
Total protein

Fig6