**Color Desaturation in the Periphery is Explained by General Mechanisms of**

**Contrast Sensitivity and Constancy**

Ana Rozman[1,2] and Jasna Martinovic[2,3*]

[1] a.rozman@sussex.ac.uk, School of Psychology, University of Sussex, Brighton UK

[2] School of Psychology, University of Aberdeen, Aberdeen UK

[3] jmartino@ed.ac.uk, School of Philosophy, Psychology and Language Science, University of

Edinburgh, Edinburgh UK

[*]Corresponding author: Department of Psychology, School of Philosophy, Psychology and

Linguistics, University of Edinburgh, 7 George Square, EH8 9JZ, Edinburgh, Scotland, UK

jmartino@ed.ac.uk

**Author Contributions**

18 **Funding**

19 AR was funded by a BBSRS EASTBIO DTP PhD studentship (BB/M010996/1) to conduct

20 her research under the supervision of JM.

**Color Desaturation in the Periphery is Explained by General Mechanisms of**

**Contrast Sensitivity and Constancy**

**Abstract**

Colors appear less saturated in the visual periphery than in the fovea. We revisit this well-known phenomenon by characterizing parafoveal perceived contrast as a function of size. Observers (n = 20) matched perceived contrast of a parafoveally presented comparison disc (2° - 0.33°) to a standard 2° disc. For chromatic stimuli, desaturation increased with decreasing size. Unexpectedly, a similar amount of desaturation occurred for luminance-defined discs, once their perceived contrast was adjusted to match the standard chromatic discs. Desaturation was reduced as standard stimulus contrast increased, in line with contrast constancy theory, which predicts constant appearance for stimuli that are sufficiently distant from threshold. Since chromatic contrast sensitivity is reduced away from the fovea, contrast constancy is unachievable within the monitor gamut. In conclusion and somewhat counter-intuitively, the appearance of color and luminance in the periphery is affected similarly, governed by general laws of contrast sensitivity and constancy.

**Keywords**

Color vision, saturation, perceived contrast, contrast constancy, luminance contrast

**Public Significance Statement**

In the early stages of visual signal processing, color and luminance (achromatic) information are separated. If we overgeneralize on the basis of such early separation, we may conclude that the desaturated appearance of small and peripherally presented color stimuli is specific to color processing mechanisms. Here we show that this is not the case and that

45    chromatic and achromatic stimuli are processed similarly when presented in the periphery,

46    implying that such processing is governed by more general contrast-processing mechanisms.

47    Our work provides the basis for a better understanding of how the visual system builds a

48    unified experience of color appearance across chromatic and achromatic dimensions from the

49    information on contrast present in the image.

**Introduction**

Saturation is a perceptual property of color that describes the vividness of its hue. Saturation depends not only on a stimulus' colorfulness (i.e., chromatic content) but also on its brightness (i.e., achromatic content; see Fairchild, 2013). For example, adding white light to a single-wavelength light would desaturate it. Thus, saturation increases with an increase in colorfulness but also decreases with an increase in brightness/lightness. An important distinction between brightness and lightness is that lightness is relational – by normalizing brightness relative to a surface that would be perceived as white, it can account for changes in illumination. However, saturation is not relational – it can also be experienced from isolated stimuli (e.g., a traffic light seen at night; Fairchild, 2013).

A more general concept than saturation is perceived contrast – referring to the overall amount of contrast perceived within a stimulus, irrespective of whether it is defined by color or luminance. Perceived contrast varies depending on the stimulus' spatial properties and location on the retina. Chromatic patches, in particular, appear desaturated when presented parafoveally (McKeefry et al., 2007). Furthermore, chromatic patches of smaller sizes appear desaturated compared to patches of larger sizes (Abramov et al., 1991; Knau & Werner, 2002). These well documented phenomena can be linked to clear demonstrations of saturation constancy in the periphery - desaturation of peripheral stimuli is not experienced as long as their size is increased to account for the cortical magnification factor (Tyler, 2015). However, loss of perceived contrast (i.e., desaturation) in periphery does not need to be color-specific - if stimulus appearance is accounted for by general contrast constancy models, changes in the periphery should be similar across color and luminance, once they are matched in perceived contrast.

Suprathreshold appearance depends on contrast sensitivity. Contrast sensitivity

function (CSF) captures how detection thresholds relate to spatial properties of the stimulus.

Sensitivity is the inverse of threshold, i.e., the lower the amount of contrast required at a

given spatial frequency for the stimulus to be detected, the higher the sensitivity for this

frequency. For luminance-defined stimuli, this follows a bandpass curve (Campbell &

Robson, 1968). Contrast sensitivity reduces due to multiple factors, for example with

eccentricity (Rovamo et al., 1978), while higher spatial frequencies are specifically affected

by ageing (Ashraf et al., 2020; Ross et al., 1985). Contrast sensitivity function can also be

measured for chromatic patches. Chromatic CSF follows a low-pass shape for the yellow-

blue and red-green contrast alike, with sensitivity dropping off faster with an increase in

spatial frequency for yellow-blue contrast, compared to red-green (Mullen, 1985a).

Chromatic contrast sensitivity is particularly reduced in the periphery (Gordon & Abramov,

1977; Johnson, 1986; Mullen, 1991). In fact, sensitivity for chromatic stimuli shows a notable

drop at the upper end of the curve (i.e. at higher frequencies) with increasing parafoveal

presentation (Ashraf et al., 2024). This is in line with Tyler's (2015) demonstration of how

retinal eccentricity can be compensated for by an increase in stimulus size, as reductions in

stimulus size limit the range of spatial frequency detectors involved in processing the

stimulus.

The specificities of vision at or near detection threshold cannot be fully interpolated to

suprathreshold vision (Cannon, 1985; Vanston & Crognale, 2018). Still, the domains of

threshold and suprathreshold vision are linked through the notion of contrast constancy

(Georgeson & Sullivan, 1975), which describes perceived contrast of a suprathreshold

stimulus as a function of contrast detection threshold for the same stimulus. This theory was

initially defined for achromatic, luminance defined stimuli, but could also explain peripheral

color desaturation. The higher the detection threshold for a certain spatial pattern, the higher

suprathreshold contrast required to make it equal in perceived contrast to a different spatial

pattern associated with a lower detection threshold. Since detection thresholds increase with

eccentricity from the fovea for achromatic and chromatic stimuli alike, the decrease in

saturation could be described as a consequence of contrast constancy– i.e., higher absolute

contrast is required to rectify the loss of sensitivity in the periphery. Translated to stimulus

size, increasing the stimulus as it moves towards the periphery rectifies the loss of sensitivity

in higher spatial frequency channels. Therefore, rather than being reflective of color-specific

low-level processing constraints, peripheral desaturation phenomena could rather be a

governed by the same general mechanisms of contrast constancy, which have been well-

described in relation to achromatic contrast processing.

In this study, we characterize desaturation as a function of size away from the fovea

for colors that isolate the two cone-opponent mechanisms and compare it between chromatic

and luminance isolating stimuli matched in perceived contrast. We also evaluate desaturation

following reduction in stimulus size for chromatic stimuli presented on pedestals of varying

levels of luminance contrast, using stimuli with a sharp and a blurred edge.  Combined, our

experiments will answer the following question: is desaturation in the periphery specific to

color or is it governed by more general contrast constancy principles that apply equally to

luminance-defined stimuli? If contrast constancy generalizes across chromatic and luminance

contrast, we would expect to observe a similar desaturation with size across chromatic and

luminance-defined stimuli.  On the contrary, if desaturation with size in the periphery is a

color-specific phenomenon, we should fail to observe a reduction in perceived contrast for

luminance-defined stimuli. In our experiments, participants match perceived contrast of a

comparison stimulus to that of a standard, fixed-size stimulus. If desaturation away from the

fovea with a reduction of stimulus size is fully explained by contrast constancy, we should

observe a gradual reduction of desaturation for both chromatic and achromatic stimuli as

standard stimulus contrast increases and moves away from threshold. Additionally, if

desaturation in the parafovea always occurs for color, addition of luminance contrast

pedestals should not markedly reduce it, regardless of the luminance contrast added. On the

contrary, if desaturation is a product of general principles that apply equally to luminance and

chromatic vision, then it should follow a pattern predicted from these models when

luminance pedestals are added. As a final test for this, we directly manipulate the edge of the

standard disc to evaluate the dependence of perceived contrast on information carried by

lower and higher spatial frequency channels.

## Methods

### *Participants*

A total of 20 participants (15 female, aged $20 - 32$ years, $M_{age} = 26$) took part in the

experiments. No participants were excluded. Sample size was verified with an a-priori power

calculation in G*Power (Faul et al., 2009), based on data from comparable work (Abramov et

al., 1991). We extracted their data in condition most comparable to our stimuli (4 color

directions at 5 degree eccentricity and $0.25 - 2$ degrees of visual angle in size) to calculate

their observed power of the desaturation with size effect for color stimuli. This was the

weakest for their 'blue' condition (Cohen's $d$ estimate $= 1.10$, ~16% desaturation) with their

sample size of 6. In order to replicate this with power of 0.8, we would require a sample of 15

participants. For their strongest effects ('green' and 'yellow' conditions, ~54/58%

desaturation) we would require a sample size of 5. Sufficiency of our sample size has been

confirmed with post-hoc power simulation-based analyses (Kumle et al., 2021),

recommended for linear mixed effect models, where it is more difficult to estimate all the

relevant values a-priori (see Supplementary Materials for a detailed overview). This analysis

verified that at least 0.8 power for all effects in all experiments would be achieved with a

sample of 8 participants.

All observers had normal color vision as checked by City University Colour Vision

Test (Fletcher, 1984) and normal or corrected to normal visual acuity according to an EDTRS

chart check for viewing at 68 cm distance. Written informed consent was obtained from

participants. The study was conducted in line with the Declaration of Helsinki (1964) and

received ethical approval from University of Aberdeen School of Psychology Ethics

Committee.

***Apparatus***

Stimulus presentation was controlled by a VISaGe visual stimulus generator

(Cambridge Research Systems, UK) and presented on a Display++ monitor (CRS, UK), using

CRS toolbox and CRS color toolbox (Westland, Ripamonti, & Cheung, 2012) for MATLAB

(Mathworks, USA). Responses were collected using a Cedrus RB-540 button box (Cedrus,

USA). Colors were calculated on the basis of monitor spectra measurements (SpectroCAL,

CRS, UK) and 2° cone fundamentals (Stockman & Sharpe, 2000). Participants were seated in

a dark room at a 70 cm distance from the display, which was the only source of light.

***Stimuli and Procedure***

Colors used in the experiment were defined to isolate the cardinal directions of

chromatic mechanisms as represented in the DKL color space (Derrington, Krauskopf, &

Lennie, 1984). Background was set to 0.3127, 0.3290, 22.9298 cd/m2 in CIE 1931 xyY

coordinates, which is metameric to the neutral illuminant D65.

**Baseline Measurements**

To ensure the absence of luminance contrast from color-only stimuli, we measured the point of isoluminance for both red-green and yellow-blue colors for each individual observer (see Figure 1 for an outline of experimental steps). We used a heterochromatic flicker spectrometry (HCFP, Walsh, 1958) task for this purpose. Observers were presented with two 2° circles flickering parafoveally, 5.5° to the left and right from central fixation. These corresponded to the standard stimulus size and location in the main experimental contrast adjustment task. The flicker was presented at the frequency of 20 Hz. Each of the presented patches flickered between two colors, red-green or blue-yellow, corresponding to L-M and S-(L+M) mechanisms. Observers were asked to minimize the perceived flicker between the two colors whilst fixating on the center of the screen. This was done by adjusting their luminances using a button box. Observers' responses manipulated the angles of elevation in DKL space (Figure 1) in opposite directions – a left button press would increase the angle for color 1 and decrease it for color 2, and vice versa. The step size was set at a 4° luminance elevation for the RG and 1° for BY trials, corresponding to ±0.02 and ±0.12 RMS contrast units for the RG and BY mechanisms, respectively. The initial difference between the two colors in a trial was set between 1 and 5 multiples of the step size for that mechanism. Once the flicker was perceptually minimized, i.e., the point of isoluminance for the two colors was reached, observers pressed the top button to finalize their setting and proceeded to the subsequent trial. A total of 8 trials per color direction was performed. The individual isoluminance was obtained by calculating the mean luminance adjustment value for each of the color pairs based on 6 measurements, following the exclusion of the highest and lowest adjustment as outliers. The task took about 10 minutes to complete.

**Figure 1**. *Schematic Representation of the Experimental Paradigm.* Both experiments started

with participants completing a HCFP task for establishing individual isoluminance (Panel 1).

A subgroup of 3 participants completed an asymmetric matching task between cardinal colors

of DKL color space (Panel 2). The standard contrast levels obtain with this baseline

measurement were then used to match them to achromatic stimuli by a subgroup of 4

participants (Experiment 1), and to obtain JNDs in luminance contrast for all participants for

Experiment 2 (Panel 3). The main experimental task was the same saturation/contrast

adjustment task in both experiments (Panel 4). Standard stimulus, always 2° in size was

presented parafoveally at 5.5° distance from fixation. The edge of standard stimulus varied

between conditions and was sharp of blurred. The adjustable stimulus varied in size between

conditions and was presented at 4° eccentricity. In the adjustment task, the standard and

adjustable stimulus were always presented in the same color or luminance polarity and at the

same level of contrast. The position of the stimuli (left or right of the fixation cross) was

counterbalanced between conditions for all experiments.

To ensure that standard stimuli defined along different chromatic and luminance axes

were equally salient to each other, perceived contrast was matched between 2° parafoveal

discs in two baseline adjustment tasks (Switkes & Crognale, 1999). Firstly, three observers

(one of the authors) completed an asymmetric matching task comparing red and green

standard discs to yellow and blue comparison discs (Figure 1). This was done to ensure equal

perceived contrast of standard stimuli between the cardinal colors of the two cone-opponent

mechanisms. In this task, discs with a 2° diameter were presented at 4° to the left and right of

the central fixation cross. The standard was either red or green, set at 0.02 RMS contrast. The

comparison stimulus was either yellow or blue. Observers were asked to adjust the contrast of

the comparison stimulus to match that of the standard stimulus whilst fixating on the central

cross (i.e., using their peripheral vision). Stimuli were set to individual isoluminance, as

measured with HCFP. Each standard and comparison stimulus pair were shown 8 times. The

location of the stimuli in relation to the central fixation cross was counterbalanced. Results

are visualized in Figure 2a.

Secondly, four observers (two of them authors) completed an asymmetric adjustment task

between achromatic and chromatic stimuli. Two discs 2° in diameter were presented 4° to the

left or right of the central fixation cross. The standard stimulus was always presented in one

of the four cardinal colors in the cone-opponent color space (red, green, blue or yellow). The

standard was set to individual observer isoluminance measured with HCFP. The contrast of

the standard was set to the value established in the chromatic perceived contrast baseline

measurement (see description of asymmetric chromatic contrast matching above). The

comparison stimulus was of either positive or negative luminance polarity, set in the DKL

space by fixing the angle of elevation to 90° (positive) or –90° (negative). Whilst fixating the

central cross, observers were asked to adjust the contrast of the achromatic comparison so

that it appears equally salient as the chromatic standard (Figure 1). Each measurement was

performed 8 times, with the location of the standard counterbalanced. The grand-mean match

was 0.036 RMS contrast (Figure 2b). There was no time limit for making a match. Observers

completed 64 trials (4 colors, 2 polarities, 8 trials per condition) in about 20 minutes.

**Figure 2.**

*Results of Asymmetric Contrast Matching Baseline Tasks.* Adjustments in baseline

asymmetric matching task for (a) chromatic and (b) luminance stimuli. (a) Standard L-M

stimuli were set to 0.02 RMS contrast. Error bars denote +/- 1 standard deviation between

observers. The horizontal line corresponds to 0.13 RMS contrast which was used for standard

stimulus contrast for the blue-yellow mechanism in the main experiment. Matches made by

one author participant are marked with an orange line. On the x axis R, G, B and Y stand for

reddish, greenish, bluish and yellowish, e.g. 'B to R' denotes bluish matched to a reddish

standard. (b) Mean contrast adjustment is provided for positive (denoted by + on the x axis)

and negative (denoted by – on the x axis) luminance polarity stimuli matched to chromatic

standards in cardinal colors of the cone-opponent DKL color space (R, G, B and Y are used

to denote these in the same way as in panel a). Error bars show the standard deviation

between observers. A horizontal dashed line denotes the matched RMS contrast value used

for luminance contrast stimuli (0.036). Matches made by authors are marked in orange.

An additional baseline task was completed for Experiment 2, to create individually

meaningful luminance contrast pedestals for the saturation adjustment task. A just noticeable

difference (JND) in luminance contrast was measured for each observer (Figure 1, panel 3,

right). This was presented as a 2AFC task staircase, asking observers to indicate which of the

two presented stimuli is lighter. The two circles appeared on the screen briefly for 200 ms to

prevent eye movements towards the stimuli, since a brief stimulus with luminance contrast

could be expected to attract eye movements in an exogenous fashion (for literature on the

timing of saccadic eye-movements see Hutton, 2008; Pierce et al., 2019; Salthouse & Ellis,

1980). Furthermore, observers were instructed to fixate the cross and avoid eye movements

throughout the task.

Conditions were blocked by stimulus size and blocks were presented in a random

order. The stimuli were either 2°, 0.5° or 0.33° discs and were presented at 4° eccentricity.

Each block contained 8 interleaved staircases - two for each of the four cardinal colours,

measuring either a positive or a negative luminance contrast JND. Each block ran until the

termination criterion of 10 reversals was met for all of the 8 staircase procedures with

Weibull function, implemented using the Palamedes toolbox (Prins & Kingdom, 2018).

Approximately 50 matches on average were required for convergence of a single staircase.

Observers completed the task in approximately 60-90 minutes. A screen encouraging

observers to take a break was presented after each set of 80 completed trials. Before

commencing each size block of the task, observers completed a short training block to get

them adjusted to the task and stimulus size. For the first trial of each block, the luminance

difference between the stimuli was set to a relatively high value of 1 cd/m$^2$ to ensure the

observers could locate the lighter stimulus comfortably and allocate their attention to the

locations in which the stimuli would appear. This trial was excluded from the analysis.

***Experiment 1: Saturation at progressive levels of suprathreshold chromatic and***

***luminance contrast.***

To compare the effect of desaturation in the periphery, we collected contrast

adjustment data for isoluminant chromatic stimuli and luminance-only stimuli at multiple

values of standard contrast. The task was completed by 10 of the total 20 observers (8 female,

aged 20 – 32 years, $M_{age} = 26$).

Luminance standards were presented at three levels of contrast. This was either

perceptually matched to contrast of chromatic isoluminant standard stimuli (as determined in

the baseline measurements described above) or set at 3- or 9-fold multiples of that standard

contrast. Color standards could only be presented at a 3-fold increase in contrast, as it was the

maximum that could be achieved within the gamut of the monitor. The standard and

comparison stimuli always appeared in the same color or polarity. The stimuli were presented

at 4° eccentricity from the central fixation cross, either to the left or to the right, with the

location counterbalanced. The standard was 2° while the size of comparison was either 2°,

0.5° or 0.33° of visual angle. Participants adjusted the contrast of the comparison stimulus

using a button box, with the left button increasing the contrast and the right button decreasing

the contrast, until a perceptual match was achieved. Participants were asked to complete the

task whilst fixating a centrally positioned cross. Each of the 36 conditions (3 levels of

contrast, 2 polarities, 3 sizes and 2 standard edge conditions) for the achromatic matching

was presented 8 times, for a total of 288 trials. There was a total of 256 matches made for the

chromatic matching task (32 conditions - 2 levels of contrast, 4 colors, 3 sizes; presented 8

times each).

### *Experiment 2: Saturation of lighter and darker colored discs with sharp and blurred edges.*

In the second part of the study, all 20 observers completed the same adjustment task

but with chromatic contrast presented either at isoluminance or upon progressive luminance

pedestals (i.e., lighter or darker than the background).

First, to match the amount of contrast between observers, Just Noticeable Differences

(JNDs) for discriminating positive and negative luminance polarity contrasts combined with

chromatic contrast (red, green, blue and yellow) were measured for each participant (details

in Baseline Measurements section).

Colors to be used for each observer were based on these baseline matches. In addition

to an isoluminant chromatic setting (identical to the one used in the first experiment), the

same chromatic information was also presented with an addition of a positive or negative

luminance pedestal, equal to a value of 3 or 6 multiples of JNDs for a given color direction.

In the main experimental task, participants were again asked to adjust saturation of a

comparison stimulus to match that of a standard stimulus whilst fixating the central cross.

The two circular stimuli were presented at 4° eccentricity left or right of the foveal fixation

cross and always appeared in the same color (reddish, greenish, bluish or yellowish) and

luminance pedestal (isoluminant, 3 or 6 JNDs positive or negative luminance contrast

pedestal). The location of the standard and comparison stimulus was counterbalanced, and the

presentation of colors randomized. All conditions with the same luminance contrast pedestal

were blocked (split into two parts to allow for breaks) and the order of these pedestal blocks

was randomized.

The standard stimulus was always a 2° disc whilst the size of the comparison stimulus

was either 2° (baseline comparison), 0.5° or 0.33° of visual angle. Observers were asked to

adjust the amount of chromatic contrast of the comparison stimulus to match the saturation of

the standard stimulus using a button box (left button increasing, right button decreasing).

Adjustments were made 8 times per condition (4 colors, 3 comparison stimulus sizes, 5

luminance contrast pedestal conditions and two standard stimulus edge definitions; see Fig.

1), with a total of 960 trials. Trials for two standard stimulus edge conditions were split into

two testing sessions, each lasting up to 2 hours for participants to complete, depending on

chosen duration of breaks.

Finally, this experiment also manipulated stimulus spatial properties by blurring the edge of

the standard disc to reduce the information carried by the higher spatial frequency channels.

The stimulus edge was blurred by adding an outer ramp encompassing 15% of the stimulus

area, within which contrast was reduced linearly (Figure 3a). This resulted in 7.5% difference

in overall contrast between the sharp and blurred edge standard stimulus. The standard 2°

stimuli (sharp edge and blurred edge) still have comparable amount of information in the

lower spatial frequency range, but the blurred edge standard stimulus is better matched in

energy to smaller, sharp edge comparison stimuli in the higher frequency domains (Figure

3b).

**Figure 3**. *Schematic representation of stimulus edge blurring (a) and stimulus energy as a*

*function of spatial frequency (b) for blurred and sharp edge stimuli of different sizes.* The

contrast of the central region (labelled "inner section" in panel A) is set to identical contrast

levels for all discs depicted in panel B. Note that the higher frequency content of a 2° disc

stimulus depends overwhelmingly on its edge – once the edge is blurred by adding a linear

contrast ramp over its outer area, the disc maintains a highly similar amount of energy in the

lower spatial frequencies (< 0.2 cycles per degree) but its higher spatial frequency content

(>2 cycles per degree) has dropped off significantly and is now within the range of the

smaller 0.5° and 0.33° discs.

### *Statistical Data Analysis*

We used linear mixed effect models on log-transformed contrast adjustment data and therefore report effect sizes as odds ratios (ORs). Analyses were implemented in R 4.3.1 (R Core Team, 2024) using the following packages: lmerTest 3.1-3 (Kuznetsova et al., 2017) emmeans 1.10 (Lenth et al., 2025), lmeresampler 0.2.4 (Loy et al., 2023), DHARMa 0.4.6 (Hartig et al., 2024) and simr (Green & MacLeod, 2016). e report further details on statistical analyses in Supplementary Materials where we also present statistical sensitivity analyses for our models, based on Kumle et al.'s (2021) recommendations of evaluating sensitivity for LMEMs by reducing fixed effects by 15% and simulating the analyses for different sample sizes re-drawn from the original sample. Supplementary Materials also contain the full properties of best-fitting models and pairwise post-hoc tests for our effects.

To further confirm whether the increase in adjustment with decreasing stimulus size differed between chromatic and achromatic stimuli, we conducted a Bayesian model comparison using the anovaBF function, part of the BayesFactor (Morey et al., 2024) package in R. The Bayesian factor estimates for each model were based on 10000 iterations. Our interpretations of Bayes factors are based on Kass & Raftery (1995) and we refer to values $BF_{10} < 1$ as evidence against hypothesis, 1-3 as weak, 3-20 as moderate and > 20 as strong evidence for the hypothesis.

### *Transparency and Openness*

We have reported how we determined out sample size, and we also report all data exclusions, all manipulations and all measures in the study. Data collection took part between December 2021 and March 2022. All data and code required to replicate the experiment and analysis have been made available on OSF (osf.io/jmght). This study was not preregistered.

## Results

### *1. Comparing Desaturation with Size Between Chromatic and Luminance Stimuli*

We analyzed the chromatic and luminance data using linear mixed effect models. For color, we could only achieve a threefold increase in standard contrast, whilst for luminance we also tested standards that were nine times higher, so we fitted two separate models to these datasets.

First, a linear mixed effects model was fitted to chromatic and luminance data, predicting adjustment as a function of size, color and contrast level (baseline level and 3x higher contrast). The best fitting model with random by-participant intercepts explained $r^2 = 0.589$ variance, with fixed effects accounting for $r^2 = .432$. Figure 4a shows the prominent reduction of perceived contrast for smaller stimuli (0.5° vs 2°: OR 1.132, 95% CI 1.107-1.157, $t(491) = 9.740$, $p < .001$; 0.33° vs. 0.5°: OR 1.081, 95% CI 1.054-1.108, $t(491) = 5.779$, $p < .001$). A significant interaction between size and contrast level was present, with desaturation significantly reduced for higher contrast stimuli at 0.33° (OR 1.073, 95% CI 1.031-1.115, $t(491) = 3.596$, $p = .003$). Desaturation with size also significantly differed across colors, with a higher loss of perceived contrast for positive luminance polarity at 0.5° against all the hues except blue – a loss which at the smallest, 0.33° size had become more uniform across colors (see Supplementary Materials for all the post-hoc tests).

A linear mixed effects model was also fitted to the achromatic data (Figure 4b), predicting contrast adjustment as a function of size, contrast level, luminance polarity and edge definition. The best-fitting model explained $r^2 = 0.633$ through the fixed effects, rising to $r^2 = 0.745$ with the inclusion of random by-participant intercepts. The three-way interaction between size, contrast level and polarity was significant - as the standard contrast

was tripled, desaturation of 0.33° discs was lowered for negative luminance polarity only (3x

contrast vs. baseline: OR 1.124, 95% CI 1.048-1.200, $t(369)$ = 3.403, $p$ = .028). With a

further tripling of luminance contrast, perceived contrast was even less affected by stimulus

size for negative luminance polarity discs (9x vs. 3x contrast: OR 1.162, 95% CI 1.084-1.240,

$t(369)$ = 4.353, $p$ < .001). In fact, at 9 times higher contrast a flat profile is observed for

negative polarity stimuli, with a lack of statistically robust differences (2° vs 0.5°: OR 1.024,

95% CI 0.955-1.093, $t(369)$ = 0.676, $p$ = 1.00; 0.5° vs. 0.33°: OR 0.972, 95% CI 0.906-1.038,

$t(369)$ = -0.814, $p$=1.00; see Supplementary Materials for further details of the post-hoc

tests).

a)

b)

**Figure 4.** *Perceived contrast for chromatic and luminance-defined stimuli of different sizes.*

(a) Isoluminant color stimuli and luminance-defined stimuli matched to suprathreshold

standards of equal salience (i.e., at a relatively well-saturated level within the monitor gamut)

as well as to standards 3 times higher in contrast. (b) Luminance-defined stimuli as a function

of stimulus size, depicting matches for a suprathreshold contrast level (left) and for three

(middle) and nine (right) times higher contrast for sharp (full line) and blurred edge (dashed

line) stimuli. Note that the left and middle panels for sharp edge stimuli are identical to the

luminance-defined conditions in (a) and are replotted to facilitate comparisons. To compare

the extent of desaturation between chromatic and luminance-defined stimuli, we represent

saturation in units of standard stimulus contrast. Therefore, an adjustment value of 200%

would indicate twice the amount of contrast in the standard stimulus was required for the

comparison stimulus to achieve the same perceived saturation. The dotted line at 100%

indicates the contrast of the central region of the standard stimulus, which in the case of

sharp-edge standards is identical to their full contrast. Dashed blue line indicates absolute

contrast value for blurred edge stimuli (92.5%). Error bars denote 2 standard errors of the

between-subject mean.

As can be seen from Figure 4b, contrast adjustments were consistently higher when

luminance-defined stimuli were matched to a sharp edge standard than to a standard with a

blurred edge. The effect of blurring surpasses the difference in overall contrast between the

standard sharp-edged and blurred-edge stimulus (*t*-test against contrast of 92.5% $t(9)$ = -

10.848, $p < .001$, $M = 83.778\%$ 95% CI 81.354`% 86.201%, $d = -3.43$), demonstrating the

importance of edges in driving our perceptual experience of contrast (see also Figure 5 for

convergent findings from our 2nd experiment).

As outlined in the methods section, to confirm whether the increase in adjustment

with decreasing stimulus size differed between chromatic and achromatic stimuli, we

conducted a Bayesian model comparison. Firstly, as expected, we found evidence against a

model predicting adjustment performance based on the colour or luminance mechanism alone

($BF_{10} = 0.12$). Furthermore, whilst a model based on size alone provided extremely strong

evidence for desaturation ($BF_{10} = 6.22 \times 10^{43}$), this was slightly reduced with the addition of

the mechanism factor ($BF_{10} = 9.06 \times 10^{42}$), thus suggesting the desaturation with size is

comparable between mechanisms. Addition of the interaction between the factors of size and

mechanism increased the evidence compared to a non-interactive model, but this increase

was weak compared to the size only model ($BF = 1.36$). Taken together, the evidence

suggests that the effect of desaturation with a decrease in size is largely comparable between

color and luminance  mechanisms.

*2.  Desaturation with a Reduction of Stimulus Size for Light and Dark Colored Discs with*

*Blurred or Sharp Edge*

A linear mixed-effects model was fitted to the data from the 2nd experiment,

predicting perceived contrast as a function of size, color, luminance contrast pedestal and

edge definition (for full details, see Supplementary Materials). There was a series of two-way

interactions between the factors, with the best-fitting model explaining about half of the

variance in the dataset (Marginal $R^2 = 0.537$, Conditional $R^2 = 0.663$). First, an interaction

between color and size replicated our previous findings: as shown in Figure 5, reduction of

stimulus size led to desaturation: the amount of contrast required to achieve equal saturation

of the comparison stimulus increased as stimulus size decreased (2° vs. 0.5°: OR 0.848, 95%

CI 0.840-0.856, $t(2297) = -34.688$, $p < .001$; 0.5° vs. 0.33°: OR 0.944, 95% CI 0.935-0.953,

$t(2297) = -11.854$, $p < .001$). Desaturation was highest for blue (vs. yellow: OR 1.053, 95%

CI 1.041-1.065, $t(2297) = 9.145$, $p < .001$), followed by yellow (vs green: OR 1.048, 95% CI

1.037-1.059, $t(2297) = 8.806$, $p < .001$; for further post-hoc tests, see Supplementary

Materials).

**Figure 5.** *Perceived contrast for colored discs of different size, presented with and without luminance pedestal, matched to either a sharp-edged or blurred-edged standard stimulus.* Perceived contrast is presented with and without luminance contrast (i.e.., lighter: 3 JND+ and 6 JND+ and Darker: 3 JND- and 6JND-, with + standing for positive and – for negative luminance polarity), adjustment is expressed as percentage of standard stimulus contrast, as in Figure 4. The dotted line at 100% indicates the contrast of the central region of the standard stimulus, which in the case of sharp-edge standards is identical to their full contrast.

In the lower panel, an additional grey line indicates the actual contrast of the blurred edge

stimulus (92.5%). Error bars represent 2 standard errors of the between-subject mean.

Luminance pedestals interacted with color, as well as with size. Addition of

luminance pedestals had a very small effect on desaturation of red or green. However, yellow

and blue were both significantly and systematically affected. For yellow, all of the luminance

pedestals led to less desaturation compared to isoluminance, without significant differences

between the different pedestals. For blue, desaturation was reduced only by negative

luminance pedestals (see Supplementary Materials for all post-hoc tests of this interaction).

Interaction of luminance pedestal with size was such that pedestals led to no significant

changes in perceived contrast for the 2° stimulus, but produced significantly less desaturation

compared to isoluminance for the 0.33° stimulus (again, see Suppl. Materials for all the

details of the post-hoc tests).

Together with our predictions of less desaturation with the addition of luminance

pedestals, we also predicted blurring of the standard stimulus edge to decrease perceived

contrast. This would mean that less contrast would be required to find a match with a blurred-

edge standard, compared to a sharp-edge standard. There was also a significant interaction

between edge definition and luminance pedestal: for blurred-edge discs, the isoluminant

condition required more contrast to achieve equal salience in relation to all conditions with

luminance pedestals except for the positive pedestal of 3 JNDs, which similarly to

isoluminance required significantly more contrast than the negative pedestal conditions

(again, see Suppl. Materials for all the details of the post-hoc tests). This is in line with our

prediction that blurring the edge would particularly affect perceived contrast once color is

combined with luminance information.

**Discussion**

We demonstrate that parafoveal desaturation with a reduction in stimulus size is characteristic of general contrast processing by the visual system, rather than specific to color. This may appear counterintuitive, as it is generally accepted that color perception is a specialization of foveal vision, despite extensive evidence that this is not the case (e.g. Tyler, 2015). Here, we show that once achromatic stimuli are perceptually matched in contrast with chromatic stimuli, they exhibit the same changes in appearance with a reduction in stimulus size as chromatic stimuli. We demonstrate that this effect can be reduced when overall contrast of chromatic stimuli is sufficiently increased or shifted towards higher spatial frequency channels through the addition of luminance pedestals. In fact, at contrast levels tested in this study, reduction in perceived contrast with a decrease in stimulus size is eliminated for negative but not for positive luminance polarity stimuli. It is not possible to achieve sufficient contrast levels within a monitor's gamut to alleviate the desaturation with size for purely chromatic stimuli. We also show that the extent of desaturation differs between the two chromatic mechanisms and that it can be reduced for the most impacted blue-yellow mechanism when presented in combination with luminance contrast pedestals. These findings highlight that general contrast mechanisms can explain loss of perceived contrast as experienced in the parafovea with a reduction in stimulus size.

Contrast constancy theory (Kulikowski, 1976) account for our findings fairly well. For positive luminance contrasts detection threshold is higher (Lu & Sperling, 2012), which would make low contrast stimuli closer to threshold and thus reduced in perceived contrast, especially when parafoveally presented. Contrast constancy can be achieved when physical contrast is sufficiently higher than threshold. For low to medium contrast gratings or Gabors, the point of constancy depends on spatial frequency (Georgeson & Sullivan, 1975) but for disc stimuli it depends on the length of an edge, with the decrease in sensitivity as size

reduces well explained by probability summation over multiple edge detectors, following the envelope of the CSF (Ashraf et al., 2023). When contrast is at threshold, only the most sensitive channel processes the information (Kulikowski, 2003). When contrast is sufficiently increased, it breaches the contrast sensitivity limitations of other channels and they begin to contribute to contrast perception. Our investigation of perceived contrast with blurred edge standards (i.e., reduced high-frequency information) confirms this: matches of full-edge comparison discs to blurred edge standards are affected across all sizes, with the effect on appearance far exceeding the overall change in standard stimulus contrast (see Fig. 4).

Similar findings on perceived contrast across chromatic and achromatic stimuli in the periphery are reported by Jiang and colleagues (2022). Their observers compared perceived contrast of a peripheral stimulus to the foveally presented standard at varying suprathreshold contrast levels until the point of objective equality (PSE) was achieved. Whilst constancy was achieved at higher contrasts, the upper range of spatial frequencies appeared less contrasty than predicted by the contrast constancy model when the standard was presented at low or middle contrast (for consistent findings of deviations from contrast constancy, see Ashraf et al., 2022). This suggests a 'hybrid model' of contrast constancy, where constancy is not entirely achievable for higher-frequency stimuli at lower or mid-ranges of contrast. Such a hybrid model is in line with our own observations here. Put together, these findings point towards desaturation as a consequence of low-level contrast bottlenecks. Our smaller stimuli appear desaturated as the general contrast sensitivity for high spatial frequencies is low for chromatic information (Mullen, 1985) and decreases further with eccentricity (Anderson et al., 1991). Due to their small size (0.33° and smaller), their energy distribution cannot be effectively processed in the parafovea. Saturation for larger discs depends on summed information from a larger number of channels and is therefore preserved at these eccentricities.

Luminance contrast pedestals play an important role in preserving saturation for blue

and yellow stimuli. For yellow, change is observed even with lower positive and negative

luminance pedestal magnitudes, whereas for blue, only negative polarity pedestals increase

perceived contrast and thus reduce desaturation with size. The blue-yellow mechanism CSF

has a lower spatial frequency cut off than the red-green mechanism (Wuerger et al., 2020).

The actual number of spatial channels within each of the mechanisms has not yet been

definitely established (Peterzell & Teller, 2000), but previous research suggests three

channels for red-green (Losada & Mullen, 1994) and two channels for blue-yellow, matching

the sensitivity of the two lowest spatial frequency luminance channels (Humanski & Wilson,

1992). Inclusion of additional luminance contrast energy exciting the range of higher spatial

frequency channels within the luminance CSF envelope would be expected to decrease

detection threshold and hence increase perceived contrast. Different impact of luminance

contrast on yellow compared to blue may be due to differently spatially tuned Short-

wavelength-ON (responding to blue) and Short-wavelength-OFF mechanisms (responding to

yellow; Hardman & Martinovic, 2021).

Finally, we need to consider if our results are influenced by eye movements. This is

unlikely for several reasons. Our experienced psychophysical observers were instructed to

only perform the task when steadily fixating, and produced patterns of desaturation consistent

with low-level influences on perception, as described above. Jiang et al. (2022) obtained

highly convergent patterns of results with a very similar adjustment task. In fact, if

participants were foveating the stimuli consecutively, they would have produced flat lines in

both our studies, without any effects of eccentricity.

In conclusion, perceptual experience of color saturation is comparable to the well-explored

contrast perception of luminance stimuli. This means that further insights into low-level

mechanisms of contrast processing, across both luminance and color, can better inform our

understanding of high-level color properties such as saturation and how these can be

accurately represented in models of color appearance. Similar research programs have

already led to a collection of large-scale threshold and suprathreshold datasets whose

modelling not only has practical applications in the domain of display technologies (e.g.,

Ashraf et al., 2024) but can also lead to interesting predictions in relation to human

perception mechanisms. For example, they would accurately predict our current findings,

dispelling the often-accepted myth that it is only color whose appearance is affected when

presented in the periphery.

**References**

Abramov, I., Gordon, J., & Chan, H. (1991). Color appearance in the peripheral retina: Effects of stimulus size. *Journal of the Optical Society of America. A, Optics and Image Science*, *8*(2), 404–414. https://doi.org/10.1364/josaa.8.000404

Anderson, S. J., Mullen, K. T., & Hess, R. F. (1991). Human peripheral spatial resolution for achromatic and chromatic stimuli: Limits imposed by optical and retinal factors. *The Journal of Physiology*, *442*, 47–64. https://doi.org/10.1113/jphysiol.1991.sp018781

Ashraf, M., Mantiuk, R. K., Chapiro, A., & Wuerger, S. (2024). castleCSF — A contrast sensitivity function of color, area, spatiotemporal frequency, luminance and eccentricity. *Journal of Vision*, *24*(4), 5. https://doi.org/10.1167/jov.24.4.5

Ashraf, M., Wuerger, S., Kim, M., Martinovic, J., Mantiuk, R. K., Wuerger, S., Kim, M., Martinovic, J., & Mantiuk, R. K. (2020). Spatio-chromatic contrast sensitivity across the lifespan: Interactions between age and light level in high dynamic range. *Color and Imaging Conference*, *28*, 65–69. https://doi.org/10.2352/issn.2169-2629.2020.28.10

Cannon, M. W. (1985). Perceived contrast in the fovea and periphery. *Journal of the Optical Society of America. A, Optics and Image Science*, *2*(10), 1760–1768. https://doi.org/10.1364/josaa.2.001760

Curcio, C. A., Allen, K. A., Sloan, K. R., Lerea, C. L., Hurley, J. B., Klock, I. B., & Milam, A. H. (1991). Distribution and morphology of human cone photoreceptors stained with anti-blue opsin. *The Journal of Comparative Neurology*, *312*(4), 610–624. https://doi.org/10.1002/cne.903120411

Dacey, D. M. (1993). The mosaic of midget ganglion cells in the human retina. *Journal of*

*Neuroscience*, *13*(12), 5334–5355. https://doi.org/10.1523/JNEUROSCI.13-12-

05334.1993

Derrington, A. M., Krauskopf, J., & Lennie, P. (1984). Chromatic mechanisms in lateral

geniculate nucleus of macaque. *The Journal of Physiology*, *357*, 241–265.

https://doi.org/10.1113/jphysiol.1984.sp015499

Fairchild, M. D. (2013). Color Appearance Terminology. In *Color Appearance Models* (pp.

85–96). John Wiley & Sons, Ltd.

https://onlinelibrary.wiley.com/doi/abs/10.1002/9781118653128.ch4

Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using

G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research*

*Methods*, *41*, 1149–1960. https://doi.org/10.3758/BRM.41.4.1149

Fletcher, R. (1984). THE CITY UNIVERSITY COLOUR VISION TEST. *Ophthalmic and*

*Physiological Optics*, *4*(3), 279–279. https://doi.org/10.1111/j.1475-

1313.1984.tb00367.x

Georgeson, M. A., & Sullivan, G. D. (1975). Contrast constancy: Deblurring in human vision

by spatial frequency channels. *The Journal of Physiology*, *252*(3), 627–656.

https://doi.org/10.1113/jphysiol.1975.sp011162

Green, P., & MacLeod, C. J. (2016). SIMR: An R package for power analysis of generalized

linear mixed models by simulation. *Methods in Ecology and Evolution*, *7*(4), 493–

498. https://doi.org/10.1111/2041-210X.12504

Hansen, T., Pracejus, L., & Gegenfurtner, K. R. (2009). Color perception in the intermediate

periphery of the visual field. *Journal of Vision*, *9*(4), 26.

https://doi.org/10.1167/9.4.26

Hartig, F., Lohse, L., & leite, M. de S. (2024). *DHARMa: Residual Diagnostics for*

*Hierarchical (Multi-Level / Mixed) Regression Models* (Version 0.4.7) [Computer

software]. https://cran.r-project.org/web/packages/DHARMa/index.html

Humanski, R. A., & Wilson, H. R. (1992). Spatial frequency mechanisms with short-

wavelength-sensitive cone inputs. *Vision Research*, *32*(3), 549–560.

https://doi.org/10.1016/0042-6989(92)90247-G

Hutton, S. B. (2008). Cognitive control of saccadic eye movements. *Brain and Cognition*,

*68*(3), 327–340. https://doi.org/10.1016/j.bandc.2008.08.021

Jiang, Z., Shooner, C., & Mullen, K. T. (2022). Achromatic and chromatic perceived contrast

are reduced in the visual periphery. *Journal of Vision*, *22*(12), 3.

https://doi.org/10.1167/jov.22.12.3

Kass, R. E., & Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical*

*Association*, *90*(430), 773–795. https://doi.org/10.1080/01621459.1995.10476572

Knau, H., & Werner, J. S. (2002). Senescent changes in parafoveal color appearance:

Saturation as a function of stimulus area. *Journal of the Optical Society of America.*

*A, Optics, Image Science, and Vision*, *19*(1), 208–214.

https://doi.org/10.1364/josaa.19.000208

Kulikowski, J. J. (1976). Effective contrast constancy and linearity of contrast sensation.

*Vision Research*, *16*(12), 1419–1431. https://doi.org/10.1016/0042-6989(76)90161-9

Kulikowski, J. J. (2003). Neural basis of fundamental filters in vision. In G. T. Buracas, O.

Ruksenas, G. M. Boynton, & T. D. Albright, *Modulation of neuronal signalling:*

*Implications for active vision* (Vol. 334, pp. 3–68). NATO Science Series, Life

Sciences.

Kumle, L., Võ, M. L.-H., & Draschkow, D. (2021). Estimating power in (generalized) linear

mixed models: An open introduction and tutorial in R. *Behavior Research Methods*,

*53*(6), 2528–2543. https://doi.org/10.3758/s13428-021-01546-0

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26.

https://doi.org/10.18637/jss.v082.i13

Lenth, R. V., Banfai, B., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M.,

Love, J., Miguez, F., Piaskowski, J., Riebl, H., & Singmann, H. (2025). *emmeans:*

*Estimated Marginal Means, aka Least-Squares Means* (Version 1.10.7) [Computer

software]. https://cran.r-project.org/web/packages/emmeans/index.html

Losada, M. A., & Mullen, K. T. (1994). The spatial tuning of chromatic mechanisms

identified by simultaneous masking. *Vision Research*, *34*(3), 331–341.

https://doi.org/10.1016/0042-6989(94)90091-4

Loy, A., Steele, S., & Korobova, J. (2023). *lmeresampler: Bootstrap Methods for Nested*

*Linear Mixed-Effects Models* (Version 0.2.4) [Computer software]. https://cran.r-

project.org/web/packages/lmeresampler/index.html

Lu, Z.-L., & Sperling, G. (2012). Black–white asymmetry in visual perception. *Journal of*

*Vision*, *12*(10), 8. https://doi.org/10.1167/12.10.8

McKeefry, D. J., Murray, I. J., & Parry, N. R. A. (2007). Perceived shifts in saturation and

hue of chromatic stimuli in the near peripheral retina. *Journal of the Optical Society*

*of America. A, Optics, Image Science, and Vision*, *24*(10), 3168–3179.

https://doi.org/10.1364/josaa.24.003168

Morey, R. D., Rouder, J. N., Jamil, T., Urbanek, S., Forner, K., & Ly, A. (2024).

*BayesFactor: Computation of Bayes Factors for Common Designs* (Version 0.9.12-

4.7) [Computer software]. https://cran.r-

project.org/web/packages/BayesFactor/index.html

Mullen, K. T. (1985a). The contrast sensitivity of human colour vision to red-green and blue-

yellow chromatic gratings. *The Journal of Physiology*, *359*(1), 381–400.

https://doi.org/10.1113/jphysiol.1985.sp015591

Mullen, K. T. (1985b). The contrast sensitivity of human colour vision to red-green and blue-

yellow chromatic gratings. *The Journal of Physiology*, *359*(1), 381–400.

https://doi.org/10.1113/jphysiol.1985.sp015591

Mullen, K. T., & Kingdom, F. A. A. (2002). Differential distributions of red-green and blue-

yellow cone opponency across the visual field. *Visual Neuroscience*, *19*(1), 109–118.

https://doi.org/10.1017/s0952523802191103

Peterzell, D. H., & Teller, D. Y. (2000). Spatial frequency tuned covariance channels for red–

green and luminance-modulated gratings: Psychophysical data from human adults.

*Vision Research*, *40*(4), 417–430. https://doi.org/10.1016/S0042-6989(99)00187-X

Pierce, J. E., Clementz, B. A., & McDowell, J. E. (2019). Saccades: Fundamentals and

Neural Mechanisms. In C. Klein & U. Ettinger (Eds.), *Eye Movement Research: An*

*Introduction to its Scientific Foundations and Applications* (pp. 11–71). Springer

International Publishing. https://doi.org/10.1007/978-3-030-20085-5_2

Prins, N., & Kingdom, F. A. A. (2018). Applying the Model-Comparison Approach to Test

Specific Research Hypotheses in Psychophysical Research Using the Palamedes

Toolbox. *Frontiers in Psychology*, *9*.

https://www.frontiersin.org/articles/10.3389/fpsyg.2018.01250

R Core Team. (2024). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing. https://www.R-project.org/

Ross, J. E., Clarke, D. D., & Bron, A. J. (1985). Effect of age on contrast sensitivity function:

Uniocular and binocular findings. *The British Journal of Ophthalmology*, *69*(1), 51.

https://doi.org/10.1136/bjo.69.1.51

Rovamo, J., Virsu, V., & Näsänen, R. (1978). Cortical magnification factor predicts the

photopic contrast sensitivity of peripheral vision. *Nature*, *271*(5640), 54–56.

https://doi.org/10.1038/271054a0

Salthouse, T. A., & Ellis, C. L. (1980). Determinants of Eye-Fixation Duration. *The*

*American Journal of Psychology*, *93*(2), 207–234. https://doi.org/10.2307/1422228

Switkes, E., & Crognale, M. A. (1999). Comparison of color and luminance contrast: Apples

versus oranges? *Vision Research*, *39*(10), 1823–1831. https://doi.org/10.1016/s0042-

6989(98)00219-3

Tyler, C. W. (2015). Peripheral Color Demo. *I-Perception*, *6*(6), 2041669515613671.

https://doi.org/10.1177/2041669515613671

Vanston, J. E., & Crognale, M. A. (2018). Effects of eccentricity on color contrast. *Journal of*

*the Optical Society of America. A, Optics, Image Science, and Vision*, *35*(4), B122–

B129. https://doi.org/10.1364/JOSAA.35.00B122

Walsh, J. W. T. (1958). *Photometry*. Constable.

Wuerger, S., Ashraf, M., Kim, M., Martinovic, J., Pérez-Ortiz, M., & Mantiuk, R. K. (2020).

Spatio-chromatic contrast sensitivity under mesopic and photopic light levels. *Journal*

*of Vision*, *20*(4), 23. https://doi.org/10.1167/jov.20.4.23

713