Network-based population analysis of 31,595 gonococcal genomes reveals phase variability,

genetic diversity and mobile element dynamics drive antimicrobial resistance and phenotypic

diversity

Duncan Carruthers-Lay[1,2], Jessica Lam [1], Emil Jurga[1,2], Scott D. Gray-Owen[1] and John

Parkinson[1,2,3*]

[1] Department of Molecular Genetics, University of Toronto, Toronto ON, Canada

[2] Program in Molecular Medicine, Hospital for Sick Children, Toronto ON, Canada

[3] Department of Biochemistry, University of Toronto, Toronto ON, Canada.

* To whom correspondence should be addressed: jparkin@sickkids.ca

Keywords: *Neisseria gonorrhoeae*, population genomics, antimicrobial resistance, phase

variation

ABSTRACT

With a highly plastic genome allowing the accumulation of antimicrobial resistance (AMR)

determinants, *Neisseria gonorrhoeae* (*Ngo*) is an urgent public health threat. To better

understand the spread of AMR in *Ngo* and its ability to persist in humans, we performed a

systematic analysis of its global population structure. From 31,595 publicly available genomes,

we curated a maximally diverse representative set of 485 gonococcal genomes, which were

found to subdivide into 12 distinct clades on the basis of shared patterns of single nucleotide

polymorphisms. Mapping AMR determinants and mobile elements onto these clades revealed

significant correlation with gonococcal population structure. Notably, several variants associated

with AMR were identified in genes not previously linked with resistance, suggesting roles as

new determinants of resistance or compensatory mutations in resistant strains. Analysis of phase

variable motifs (a significant mechanism driving regulation of expression in *Ngo*) identified 57

genes not previously identified as phase variable; *in vivo* and *in vitro* passaging validated our

sequence-based definition of phase variability and revealed a strong association of phase

variation of *opa* and *lgtG* with improved colonization during infection. Together these findings

highlight the ability of systematic comparative genomic analyses to shed new light on the drivers

of *Ngo* population structure and identify new AMR determinants with implications for the

development of novel therapeutics.

INTRODUCTION

The gram-negative bacterial pathogen *Neisseria gonorrhoeae* (*Ngo*) is the causative agent of the sexually transmitted infection gonorrhea. The most common site of gonococcal infection is the urogenital tract, where it is typically asymptomatic in females[1]. Consequently, without treatment, infection can result in severe morbidities such as pelvic inflammatory disease or infertility[1]. Gonococcal infection also increases the risk of coinfections with chlamydia and human immunodeficiency virus[2,3]. With rapidly rising rates of infection and widespread development of antimicrobial resistance (AMR), *Ngo* is now considered a high priority pathogen by the World Health Organization, requiring the development of novel therapeutics and additional treatment options[4]. Since the advent of antimicrobials, *Ngo* has demonstrated the ability to rapidly acquire AMR mechanisms, threatening our ability to treat infections and raising the prospect of untreatable gonorrhea[5].

Supporting its ability to acquire AMR, *Ngo* is naturally competent with many DNA uptake sequences throughout genome which enables easy incorporation of genetic matter between gonococcal strains and from other pathogenic and commensal *Neisseria* species[6,7]. Driving this "promiscuity" are mobile elements, including plasmids which contain AMR encoding genes and the Gonococcal Genetic Island (GGI) which encodes a type IV secretion system, together with several proteins of unknown function[8,9]. Further, many gonococcal genes exhibit phase variability in which changes in the number of tandem repeat sequences can alter gene expression[10]. This genetic promiscuity, phase variability and high rates of recombination support *Ngo's* ability to rapidly adapt to novel environmental perturbations and result in a highly diverse population structure not observed in the closely related *Neisseria meningitidis*[7]. This population heterogeneity has confounded traditional efforts to define clonal lineages, requiring instead the use of alternative approaches such as multilocus sequence typing (MLST). While early attempts at classifying gonococci based on MLST relied on only limited sets of housekeeping genes[11], more recent scheme's adopting larger sets of core loci, such as the core genome MLST scheme, have been more successful in defining relatively stable gonococcal core genome groups[12]. Complementing such efforts are tools such as PopNet which take a more agnostic view to selection of loci for determining population structure[13]. Further, PopNet's use of network graphs help define genetic

relationships across multiple strains that are otherwise lost in the rigid structure of the more
traditional bifurcating phylogenetic trees.

With the recent availability of genome data from over 30,000 *Ngo* isolates, we sought to apply
PopNet to systematically define the global population structure of *Ngo*, investigate its pangenome
and assess the incidence of AMR mechanisms and the genetic elements helping support their
dissemination. Through this analysis we identified and subsequently validated novel phase
variable genes and AMR associated loci.

RESULTS

A representative set of 485 genomes captures temporal and geographic diversity

To begin our analyses of the global population structure of *Ngo*, we first defined a representative
set of 485 isolates from an initial set of genome sequence reads from 31,595 isolates annotated by
year and location of isolation hosted by the NCBI sequence read archive (September 26[th] 2023;
Supplementary Table S1). 1,051 isolates were removed after filtering for contaminated, poor
quality or mislabelled sequences. Of the remaining isolates, 24,431 (80.0%) were derived from
North America or Europe, 3,318 (10.9%) were derived from Oceania, primarily in Australia, while
only 2,795 (9.1%) were derived from Asia, South America, or Africa. To maximize geographic
diversity, we prioritized isolates based on their continent and year of isolation (see Methods;
Figure 1A). In brief, following alignment of genome sequence reads to the reference genome (*Ngo*
strain TUM19854), representative isolates were selected by first grouping isolates by their country
and year of isolation. Isolates within each group were then clustered based on average nucleotide
identity estimated by the Jaccard Index, resulting in an initial set of 1,307 isolates. A subsequent
round of clustering using the same method with isolates grouped by continent and year of isolation
reduced this set to 451 isolates, to which an additional 20 isolates from rare body sites together
with a set of 14 isolates representing the WHO reference collection[14,15] were added. The resulting
reference collection of 485 isolates represent 36 countries, with 38% of the isolates recovered from
Canada (72 isolates), the United States (58), and the United Kingdom (54) (**Figure 1B;**
Supplementary Table S2). The reference set captures isolates recovered between 1972-2023,
with the majority (308, 63.5%) recovered between 2010-2020. Isolates from Asia appear to be the
most evenly temporally spread, while in Africa isolates were exclusively recovered after 2013

(**Figure 1C; Supplementary Table S2**). Notably, our prioritization strategy resulting in a

substantially higher proportion (148, 30.4%) of isolates from Asia, South America, and Africa

compared to the initial set of 31,595 isolates (**Figure 1D**).

PopNet defined clades show limited enrichment in specific locales

Having defined our set of representative isolates, we next defined the population structure of these

isolates based on shared patterns of single nucleotide polymorphisms (SNPs), relative to the

reference genome (*Ngo* strain TUM19854), using PopNet. We defined 12 clades (designated A-L;

Figure 2A) ranging in size from 203 isolates (clade A) to 4 isolates (clades K & L). Within these

clades we found an enrichment of isolates from Africa (p = 1.77×10^{-13}, Fisher's Exact) and Asia

(p=0.00047, Fisher's Exact) in clade B (**Figure 2Ci**). Clades K and L were exclusively comprised

of isolates from Canada, while 4 of 5 isolates from clade J were isolated from Oceania. No other

significant associations between continent of isolation and PopNet clades were observed at a

significance level of p = 0.05. Together these findings suggest that geography has only a limited

impact on *Ngo* diversity, highlighting its ability to rapidly disseminate on a global scale (**Figure**

2B).

Comparison of PopNet clades with a maximum likelihood tree constructed using

ClonalFrameML[16] showed good concordance, with clades largely remaining intact between the

two methods (**Figure 2B**). We additionally investigated how the PopNet defined clades compared

to those obtained using two other population level clustering methods, cgMLST and PopPunk[12,17].

There was a large discrepancy between the size and number of clades obtained from PopNet (12

clades, median size 10, mean size 37.46) compared to cgMLST (99 clusters, median size 1, mean

size 4.87) and PopPunk (161 clusters, median size 1, mean size 3.03). The number and size of

clades obtained by PopNet is determined by the inflation and pre-inflation values selected based

on the silhouette index within the given set of genomes. Both cgMLST and PopPunk use pre-

determined databases in their clustering approaches, which, given the intentionally diverse set of

genomes chosen in out representative set, may have contributed to the large number of small

clusters observed with these methods. Visual inspection of the clusters (**Supplementary Figure**

S1) showed that many cgMLST or PopPunk clusters with more than 5 isolates were located within

a larger PopNet cluster. In the context of a smaller, more diverse set of genomes, the agnostic

clustering approach and 2-dimensional network generated by PopNet is better suited to highlight

similarities and differences between closely related isolates.

Pangenome analysis reveals PopNet clades reflect patterns of mobile elements

To compare gene complements across the 485 isolates, we generated genome assemblies *de novo*

using unicycler v0.5.0[18]. Genomes were subsequently annotated using Prokka[19,20] to identify genes

and orthologs defined using Panaroo[19,20] (**Supplementary Table S3**). Across all genomes we

identified a total of 2460 unique genes (i.e. orthogroups), with 1,800 "Core" genes appearing in

greater than 95% of isolates, 367 "Shell" genes appearing in between 15% and 95% of isolates and

283 "Cloud" genes appearing in less than 15% of isolates (**Figure 3A**). Accessory genes (non-core

genes) were clustered according to their presence or absence across genomes using the Jaccard

index to calculate similarity and a Markov Clustering with a granularity of 1.10 (**Figure 3C**).

Several large, distinct clusters emerged, which we theorized represented mobile elements. These

mobile elements were identified through sequence similarity searches against a database of

reference sequences generated from the GGI, the *Neisseria* cryptic plasmid, the β-lactamase

plasmid, and 3 variations of the conjugative plasmid (pConj; the pEP5050 American tetM plasmid,

the pEP5233 Dutch plasmid, and the pEP5289 Dutch tetM plasmid). Accessory genes were

assigned to mobile elements based on a nucleotide BLAST[21] search against this database with a

minimum sequence identity of 85% and an e-value less than $1x10^{-6}$. The GGI was identified in

315 (64.7%) isolates, the cryptic plasmid was identified in 429 (90.1%) isolates, the β-lactamase

plasmid was identified in 70 (14.5%) isolates, and the conjugative plasmid was identified in 165

(33.9%) isolates, of which 97 (19.9% of total isolates; 58.8% of pConj containing isolates) were a

tetM-containing plasmid.

Interestingly, clustering of isolates based on their gene complements was largely congruent with

PopNet defined clades (**Figure 3B**). This gene-based clustering appears to have been largely

driven by the presence or absence of mobile elements, specifically the GGI and the conjugative

plasmid (**Figures 3A and 3B**). Further investigation of PopNet defined clades revealed a spectrum

of associations with different mobile elements (**Figure 2Cii**). For example, clade A was

significantly depleted in both the GGI ($p=1.6x10^{-8}$, Fisher's exact) and the pConj (without tetM;

p=0.00021, Fisher's exact); clade B was significantly enriched with isolates carrying the β-

lactamase plasmid (p=0.0023, Fisher's exact) as well as the pConj-TET plasmid (p=0.00028,

Fisher's exact); clade C was defined by the absence of the pConj-TET plasmid in all 25 isolates

(p= 0.0087, Fisher's exact); while all 17 Clade D isolates possessed the GGI (p=0.0021, Fisher's

exact) and lacked the cryptic plasmid (p=3.64×10^{-19}, Fisher's exact). Of note, isolates from Africa

were found to be significantly enriched in pConj-TET (34/42 isolates; p=5.59×10^{-19}) and the β-

lactamase plasmid (21/42; p=1.34×10^{-8}), independent of the year of isolation and likely associated

with the continued use of penicillin and tetracycline to treat *Ngo* infections in these countries.

Several accessory genes clusters remained that did not match our set of known mobile elements.

Using the MCL clustering tool implemented in Graphia, we identified several clusters of genes

based on their presence or absence in our set of representative genomes[22]. While some of these

clusters lined up with the mobile elements described above, others did not. We sought to further

characterize several of these clusters that were present in <50% of our genomes and included at

least 5 genes (**Supplementary Table S4**). We identified 5 clusters meeting these criteria, with one

of them (MCL cluster 3) representing the previously identified VirB type IV secretion system

(**Figure 3C**). MCL cluster 11 was composed of 5 genes which appear to represent a fully intact

Microviridae genome present in 7 genomes. All the cluster 11 genes are found on a single contig,

and do not appear to be integrated into the gonococcal genome. MCL cluster 12 was also composed

of 5 genes present in 31 genomes, 2 of these genes were annotated as putative phage proteins

however unlike cluster 11 we were unable to identify a fully intact phage genome. The largest

cluster we identified was cluster 5, composed of 17 genes present in 17 genomes. Interestingly,

these 17 genomes all fall within PopNet clade D, and include isolates from 3 BioProjects from the

United Kingdom, Japan and the United States. Many of the genes present in cluster 5 are annotated

as putative phage protein, however, unlike the other elements described so far, these genes do not

all appear on the same contig and instead are split between 2 contigs flanked by core genes (**Figure**

3D). The cluster present in the largest number of genomes was MCL cluster 13, also composed of

5 genes present in roughly half (217/485, 44.7%) of our set of representative genomes. Unlike the

previously described clusters, cluster 13 does not appear to be phage related, with the only 2

annotated genes listed as colicin V and hemolysin D.

PopNet defined clades exhibit distinct signatures of AMR mechanisms that are often

mutually exclusive

With increased concern over the propagation of AMR within *Ngo* populations, we next predicted

resistance to six antimicrobials – penicillin (PEN), tetracycline (TET), ciprofloxacin (CIP),

azithromycin (AZM), cefixime (CFM) and ceftriaxone (CRO) - for each of the 485 reference

isolates using the online platform, PathogenWatch[23] (**Supplementary Figure S2**). The degree of

resistance (resistant, intermediate, susceptible) was classified based on the EUCAST clinical

breakpoints v9.0[24]. Incidence of resistance to the three older antimicrobials (PEN, TET & CIP)

were higher than rates of resistance to the other three and were significantly enriched in isolates

derived from Africa (p=0.029, 0.00011 and 0.016 for PEN, TET & CIP respectively, Fisher's

exact).

We found several PopNet clades were also significantly associated with resistance to specific

antimicrobials (**Figure 2B**). For example, clade A was enriched in isolates resistant to AZM

(61/202 isolates; $p=1.15 \times 10^{-17}$, Fisher's exact) and isolates with intermediate resistance to TET

(108/202 isolates; $p=8.44 \times 10^{-6}$, Fisher's exact); clade B isolates were significantly more resistant

to CFM (44/176 isolates; $p=2.31 \times 10^{-5}$, Fisher's exact) and PEN (71/176 isolates; $p=4.97 \times 10^{-5}$,

Fisher's exact); while all 13 isolates in clade F, exhibited intermediate resistance to PEN (p=0.036,

Fisher's exact), resistance to CIP (p=0.00093, Fisher's exact) and resistance to TET (p=0.00027,

Fisher's exact). Interestingly, clade C was enriched in isolates resistant to TET (23/25 isolates;

$p=1.63 \times 10^{-5}$, Fisher's exact) despite the absence of the pConj-TET plasmid. Instead, resistance

appears a consequence of a double mutation (*mtrR* promoter -5delA deletion & *rpsJ* V57M). Clade

C was also enriched in isolates with resistance to CIP (20/25 isolates; p=0.021, Fisher's exact).

The association of AMR patterns with clades suggests that AMR helps shape the global population

structure of *Ngo.*

When evaluating susceptibility to older antimicrobials (PEN, TET & CIP), isolates demonstrating

resistant to one were significantly more likely to be resistant to another. When comparing isolates

resistant to the newer antimicrobials (AZM, CFM & CRO) no significant association was observed

between AZM resistance and resistance to either CFM or CRO. Isolates resistant to cefixime were

significantly more likely to be resistant to ceftriaxone ($p=8.64 \times 10^{-6}$, Fisher's exact).

Analysis of repeat expansion distributions identifies 57 potentially novel phase variable genes

Previous studies have shown that many *Ngo* genes contain repeats that can result in slipped strand mispairing or site-specific recombination altering their expression and providing a reversible process to adapt to rapidly changing environments. Within our collection of 485 reference isolates we searched for candidate phase variable (PV) genes using PERF[25] to detect simple sequence repeats. In total, we identified 1554 genes with at least one simple sequence repeat. From this set we identified 85 genes with repeats that varied in length, with each repeat length appearing in at least 1% of available sequences for that gene (**Supplementary Table S5**). These genes were compared to those identified by Zelewska et. al in their 2016 screen based on locus tags from the reference strains FA1090 and NCCP11945, with 28 appearing in both sets. This overlap included 11 *opa* genes, 6 genes characterized as *pilC2* or Pilin cassette, and other well characterized phase variable genes such as *lgtG*, *fetA* and *modA/B* (**Supplementary Table S5**).

The remaining 57 genes represent putative phase variable genes, with the 10 genes with the greatest repeat length variation shown in **Table 1**. Among these 10, group_1317 is annotated as cytosine-specific methyltransferase, consistent with other gonococcal methyltransferases (e.g. *modA* and *modB*) known to be phase variable. Another 2 predicted genes (*argS* and *glyS*) are annotated as arginyl- and glycyl-tRNA synthetases, a system which was previously identified in a computational screen for phase variable genes[26]. We identified 2 genes (group_1584 and *prn*) which were annotated as pertactin or autotransporters, with *prn* appearing to encode a mostly intact autotransporter while group_1584 appears to only encode the N-terminal domain. Another autotransporter, *autA* is known to be phase variable, however the identified repeat for *autA* is AAGC, while we identified poly-A tracts in group_1584 and *prn*[27]. While all the genes so far described are chromosomally encoded "Core" genes, we also identified a gene (*kfrB*) carried on the conjugative plasmid. This gene possesses a poly-G repeat tract immediately after the start codon, and plays a role in plasmid partitioning[28].

Passaging gonococci *in vivo* improves colonization in a mouse model

Passaging of gonococci *in vitro* has previously been used to identify phase variable genes by

observing repeat length changes in a single strain over time[10,29]. Consequently, we wanted to

evaluate the potential impact that phase variable genes would have on enabling gonococci to adapt

to and survival in a more natural environment. To accomplish this, we utilized a lower genital tract

colonization model with female FVB mice expressing human CEACAM3, -5, and -6[30]. Prior to

infection, mice in the diestrus stage of the estrous cycle began a hormone and antibiotic regimen

to suppress commensal microflora A schematic of the mouse passaging work is shown in **Figure**

4A.

For initial infection (Passage 1; P1), 12 mice were each inoculated with 10^7 bacteria of N2009

(laboratory-passaged MS11 derivative; denoted parental strain), and vaginal washes were

performed every two days to recover viable bacteria. On Day 2, only 2 of the 12 mice remained

colonized by the bacteria; these were denoted as M1 and M2 (**Figure 4A**). Mice infected from the

M1 lineage demonstrated progressively higher rates of colonization, 75% and 88% in Passage 2

(P2) and Passage 3 (P3), respectively (**Supplementary Figure S3**). Moreover, we also observed

an increase in bacterial burden recovered from the P3 isolates compared to previous rounds of

infection (**Figure 4B**). These findings suggest that gonococcal isolates passaged *in vivo* are better

adapted for survival during the initial stages of infection **(Figure 4B)**.

Phase variation of *opa* and *lgtG* genes contributes to enhanced colonization of passaged

isolates

To delineate mutations solely conferring a fitness advantage *in vivo*, we performed complimentary

in vitro passaging using the parental strain to control for random variations introduced during the

bacterial replication process. Bacteria was sub-cultured in nutrient-rich media in 5 separate tubes

every 10-12 hours for 5 consecutive days to generate 5 independent in vitro passaged lineages.

The parental isolate, *in vivo* P2 isolate (M1.05), 7 isolates from the *in vivo* P3 (P3_580, P3_521,

P3_581, P3_676, P3_680, P3_691 & P3_698), and 5 *in vitro* passaged isolates (T1D5-T5D5) were

sequenced and analyzed for the presence of PV genes as described previously. From this

complementary screen, we identified 9 genes with signatures of phase variability including the

known phase variable genes *pilC2*, pilin, *opa* (4), *fetA,*, *lgtG* and *modA* (**Table 2**).

We examined the repeat length for each gene to identify any shifts corresponding to the improved

survival of the P3 isolates. We observed that the predicted number of repeats in group_1283

dropped from 11 to 10 in P2, P3_580 and P3_698. No shift was observed in any of the *in vitro*

strains, leading us to hypothesize that this shift may be contributing to the improved colonization

of *in vivo* passage 3 isolates. Due to the inherent population heterogeneity in *Ngo*, especially when

it comes to phase variable genes, we used capillary gel electrophoresis to determine the bacterial

population proportion containing 11 or 12 repeats based on peak density. We found that the

parental strain and the 3 tested *in vitro* passage strains showed no variation in repeat length,

remaining at 12 repeats the entire time, while the passage 2 and 3 strains were more heterogeneous

(**Figure 4C**). Approximately 32.5% of the recovered passage 2 isolates had 11 repeats, with the

remaining 67.5% staying at 12. Passage 3 isolate P3_691 also did not vary, while P3_521 (9.1%

11, 90.9% 12), and P3_581 (37.3% 11, 62.7% 12) remained majority at 12 repeats. However,

passage 3 isolates P3_676 and P3_698 were found to be 100% at 11 repeats, and the majority of

P3_580 (54.2% 11, 45.8% 12) was found to be at 11 repeats. We were unable to recover sufficient

genetic material from P3_680 for analysis. These findings indicate that the isolate populations

containing a majority of the 11 repeat *opa* gene group_1283 are turning on expression of an Opa

protein which may contribute to improved colonization in a mouse model, consistent with its key

role in adhesion to host CEACAMs[30].

We also identified *lgtG* (group_1111) as a promising candidate, as the parental and all the *in vitro*

strains contained a 10bp long poly-C repeat, while the P2 and all the P3 isolates shifted to an 11bp

long repeat. This shift enables complete transcription of *lgtG*, resulting in the attachment of the β-

chain to the gonococcal lipooligosaccharide and production of the 2C7 epitope (**Figure 4D**)[31]. We

confirmed the presence of the 2C7 epitope exclusively in the *in vivo* passaged strains through an

immunoblot assay. Due to the challenges of differentiating single nucleotide length repeats with

capillary electrophoresis, we did not perform this analysis for *lgtG*. We show here the ability of

our method to identify phase variable genes in an *in vivo* setting. We also identified phase variation

in 2 genes, *opa* and *lgtG* which may play a substantial role in improving *Ngo*'s ability to colonize

and infect a humanized mouse model.

Analysis of sequence variants reveal novel sites of genes involved in AMR as being under positive selection

In the previous sections our genomic surveys identified several phase variable genes that may help *Ngo* adapt to environmental perturbations. Another signature of genes supporting adaptation is whether they are under positive selection. To identify such genes we used the BUSTED algorithm as implemented in HyPhy[32,33]. Of the 1,853 core genes, 1,724 were sufficiently variable to test for selection. We identified 355 genes with significant evidence for positive selection (likelihood ratio test statistic > 0, p-value < 0.05). As expected, given their direct contact with the environment, positively selected genes were significantly enriched in outer membrane proteins (Benjamini-Hochberg (BH) corrected p-value 0.00615) (**Figure 5A**). In addition, we also identified several AMR-associated genes as being under positive selection, including *porB*, *penA*, *parC*, *mtrD*, and *gyrA*. To identify specific sites in AMR genes under positive selection we applied the FUBAR algorithm as implemented in HyPhy[34]. Sites with a posterior probability of positive selection > 0.9 were considered as sites with strong evidence for selection. Focusing on the gyrase, *gyrA*, our analyses identified residues S91 and D95, previously characterized as conferring resistance to ciprofloxacin[35], as being under positive selection (**Supplementary Table S6**). Ciprofloxacin resistance is also associated with mutations in residues D86, S87, S88, and E91 in the type IV topoisomerase, *parC*. In addition to sites D86, S87 and E91 which interact directly with ciprofloxacin[36,37], we also identified residues T386, G481, and S598 as being under positive selection. Residues T386 and G481 reside in the α-helices which connect the catalytic and dimerization domains, while residue S598 is located in the C-terminal domain (**Figure 5B**)[38]. We identified 8 residues (310, 642, 742, 922, 926, 1052, 1065, 1067) with signatures indicating strong positive selection in *mtrD*, likely reflecting the acquisition of mosaic alleles from other *Neisseria* sp. associated with azithromycin resistance[39]. For the known penicillin resistance gene, *penA*, we identified 6 residues with strong signatures of positive selection (**Figure 5C**). We identified residues 550, 553, 556, and 567 within the α11 helix which includes the known P551 residue previously linked to penicillin resistance through inhibition of acylation by β-lactamases[40]. We also identified a residue (A502) in the region between the β3-β4 loops which has been linked to ceftriaxone resistance[41]. The final residue under positive selection was Y201, which is located between the α3n helix and β6n loop in the N-terminal domain of penA, which to date has not been linked to AMR[40]. Previous work has used signatures of positive selection to identify genes

contributing to antimicrobial resistance in *Escherichia coli*, here we identified signatures of strong

positive selection in individual residues of AMR-associated genes to identify new or

complementary sites that may contribute to resistance[42].

Genome wide association study

Beyond signatures of positive selection, genome wide association studies offer a powerful

approach to identify SNPs associated with specific traits such as AMR[43]. Here we used the

software tool, pyseer, with the linear mixed model mode to identify SNPs significantly associated

with predicted resistance to antimicrobials (**Supplementary Table S7**)[44].

We identified several well-known SNPs together with several less characterized mutations

significantly associated with AMR. For ciprofloxacin resistance, the former include the well

studied S91F and D95A/N mutations in *gyrA*, as well as an S87R mutation in *parC* (**Figure 5Di)**

However, we also identified two mutations in (G9S & I10V) in *lrgB*, a cytoplasmic membrane

associated murein hydrolase transporter, as being associated with ciprofloxacin resistance. Of the

four variants with the greatest predicted effect size associated with penicillin resistance, two were

identified in *penA* (1719C>T & 1713A>C), while the other two were found in *murE* (216C>T,

71G>A), which lies 25bp upstream of *penA* and plays a role in cell wall biogenesis

(**Supplementary Figure 4A**). A highly significant missense variant (Arg89Glu) was also

identified in *ychF*, a ribosome binding ATPase. For tetracycline resistance, our analysis identified

a variant in the known resistance gene, *rpsJ* (Met57Val), as well as multiple mutations in *rsmB*, a

methylase which interacts with the target of tetracycline, the 16S portion of the small ribosomal

subunit (**Supplementary Figure 4B**). For both ceftriaxone and cefixime resistance, we identified

multiple variants in *penA*, consistent with known resistance mechanisms (**Supplementary Figure**

4Ci & 4Cii). Finally, we identified multiple significant variants associated with azithromycin

resistance in genes associated with the MTR multi-drug efflux pump (*mtrC, mtrD,* and *mtrE*)

(**Figure 5Dii**). Interestingly, we also identified multiple significant variants in *mqo*, a malate-

quinone oxidoreductase involved in the TCA cycle. Azithromycin is not known to target *mqo* or

the TCA cycle, and pairwise structural alignments from AlphaFold predicted structures using TM-

align show *mqo* shares minimal similarity with *rplV* (TM-score 0.11, 7% identity) or *rplD* (TM-

score 0.17, 6% identity), known ribosomal targets of azithromycin[45–48] (**Figure 5E**).

DISCUSSION

With the emergence of strains resistant to all known antibiotics, *Ngo* is a threat to global health.

To better understand how AMR mechanisms arise and disseminate on a global scale, we conducted

a systematic analysis of *Ngo* population genomics. Our set of 485 representative genomes

successfully captured the genetic diversity of the larger set of 31,595 isolates, increasing the

relative contribution of isolates collected from outside North America and Europe. Ensuring

representation across the globe is critical if we are to understand the impact of geopolitical

considerations regarding surveillance and treatment options, factors that are acutely elevated for a

pathogen that is restricted to a single host[5]. Applying the population genomics tool - PopNet, we

define 12 clades, which, with the exceptions of clades B, K and L, contain isolates that are

geographically dispersed, suggesting another mechanism my be influencing gonococcal

population structure. We observed a high degree of overlap between clades defined by other tools

based on analysis of more limited sets of loci (e.g. cgMLST), supporting the utility of PopNet to

define genetically distinct sub-populations.

Given the lack of overlap between the place of isolation and clade membership, we investigated

how global population structure may be influenced by other factors, specifically mobile elements

and AMR mechanisms. In keeping with previous findings, we observed PopNet defined clades

correlate with the presence of specific mobile elements and AMR determinants[49,50]. By contrast,

most identified clades were not significantly associated with other available metadata, including

sex of patient, site of isolation, year of isolation and country or continent of isolation. A noteworthy

exception was clade B, which was significantly associated with isolates collected from Africa and

Asia. This association can be explained by the elevated presence of the β-lactamase plasmid and

the TET-containing conjugative plasmid in group B. This is in line with previous studies that have

suggested that isolates from lower income countries are more likely to carry plasmids encoding

resistance to penicillin or tetracycline[51]. Further, penicillin resistance in African isolates and nearly

all of the observed tetracycline resistance (94.2%) could be attributed to the presence of the β-

lactamase and TET-containing conjugative plasmids, respectively, In keeping with previous

findings, we observed PopNet defined clades correlate with a strong connection between the

presence of specific mobile elements and AMR determinants with PopNet clades[49,50]. By contrast,

most identified clades were not significantly associated with other available metadata, including

sex of patient, site of isolation, year of isolation and country or continent of isolation. A noteworthy

exception was that of clade B, which was significantly associated with isolates collected from

collection in Africa and or Asia. This association can be explained by the elevated presence of the

β-lactamase plasmid and the TET-containing conjugative plasmid in group B, both of which

isolates from Africa were significantly more likely to carry. This is in line with previous studies

that have suggested that disparity has been previously noted, with isolates from lower income

countries are significantly more likely to carry plasmids encoding resistance to penicillin or

tetracycline[51]. Going further, all the penicillin resistance in African isolates and nearly all of the

observed tetracycline resistance (94.2%) could be attributed to the presence of the β-lactamase and

TET-containing conjugative plasmids, respectively. For both penicillin and tetracycline, resistance

in non-African isolates was driven by a range of chromosomal mutations, particularly in *penA*,

mtrR and *rpsJ*. It is likely that this difference is driven by the continued usage of penicillin and

tetracycline to treat *Ngo* infections in lower income countries in Africa, and may indicate some

additional fitness cost to plasmid carriage in gonococci in the absence of this pressure[52].

Phase variability is a key mechanism used by *Ngo* to evade the immune system through antigenic

variation and increasing genetic variance either by directly affecting gene transcription and

translation or by impacting downstream processes like methylation. Several well characterized

phase variable genes have been identified in *Ngo*, including those encoding for outer membrane

proteins such as *opa* and *pilC*, metabolic genes such as *lgtA*, and methyltransferases like *modA*[29].

While extensive work has been done to characterize these genes in reference strains, as well as by

identifying homologs in the closely related species, *N. meningitidis*, recent evidence has suggested

that the phase variable repertoire of *Ngo* could be much larger than we previously thought[26,29]. We

identified 57 genes containing tandem repeats of varying lengths which represent putative phase

variable genes. To complement these analyses we performed a passaging experiment in an attempt

to drive the emergence of phase variability. *In vitro* passaging experiments have previously been

successful in the identification of novel phase variable genes[29]. Here we used a tandem *in vitro &*

in vivo passaging strategy to identify phase variable genes that help support *Ngo*'s ability to infect

and propagate. During the passaging experiment, we observed improved colonization of

gonococcal isolates with each successive passage, indicating some level of adaptation to the mouse

host. Through this tandem approach, we identified several previously characterized phase variable

genes (*pilC, opa, modA, lgtG & fetA*) in our passaged isolates, of which two (group_1283, *opa* &

group_1111, *lgtG*) varied in repeat number only in *in vivo* passaged isolates. Through capillary gel

electrophoresis, we were able to confirm that through phase variation the opa protein encoded by

group_1283 varied in passage 2 and 3 isolates, likely contributing to the observed improved

colonization. We also observed a shift from 10 to 11 bp repeats in all *in vivo* passage 2 and 3

isolates, resulting in transcription of *lgtG*, with no shift observed in the *in vitro* passaged strains.

Expression of lgtG leads to the production of the 2C7 epitope, which is present in a high proportion

of clinical isolates, indicative of a role in gonococcal pathogenesis and colonization[53]. Our results

support this conclusion, and demonstrate the utility of passaging and sequencing for identifying

genomic variation linked to improved gonococcal pathogenicity.

Given the genomic plasticity of *Ngo* and selection constraints imposed by the host immune system,

we hypothesized many *Ngo* genes may be subject to positive, diversifying selective pressure. We

identified 355 of 1,853 core genes to be under positive selection. This number dramatically

expands on the previous estimate of only 10 *Ngo* genes under positive selection, likely reflecting

differences in methodology as well as the larger number of genetically diverse isolates used here[54].

A similar approach focusing on 2 isolates of *Burkholderia glumae* revealed a comparable number

of genes under positive selection (259/3,777 tested genes)[55], although we note the proportion of

gonococcal genes under positive selection is substantially higher. We found no functional

enrichment for positively selected genes, though we did observe a significant enrichment in

predicted outer membrane proteins, as would be expected given immune pressure exercised by the

host. Beyond identification of novel genes responding to host pressure, we also identified several

novel sites in known AMR genes associated with positive selection. These novel sites likely

represent complimentary variants or emerging resistance-conferring variants and require further

exploration.

Beyond using selection pressure to identify genes and variants associated with host adaptation and

environmental perturbations, we also sought to identify variants associated with the carriage of

mobile elements and AMR through a genome-wide association analysis. We successfully

identified known variants in multiple AMR-associated genes, including *gyrA* and *parC* for

ciprofloxacin, *penA* for penicillin, ceftriaxone, and cefixime, *rpsJ* for tetracycline, and multiple

mtr genes for azithromycin. We additionally identified several highly significant variants

associated with AMR in genes with no known links to AMR. These include a variant in *lrgB*

associated with ciprofloxacin, which has previously been indirectly linked to ciprofloxacin

resistance in *Ngo* and may play a role in the oxidative stress response[56]. We also found multiple

non-synonymous and missense mutations in *rsmB* associated with tetracyline resistance. This

finding is of particular interest, as *rsmB* is a methyltransferase which interacts directly with the

16S rRNA portion of the small ribosomal subunit, which is targeted by tetracycline[57]. Little is

known about *rsmB* in *Ngo*, and while it is unlikely that it represents a novel resistance mechanism,

it may complement additional tetracycline resistance-conferring mutations. Excitingly, we also

found multiple significant variants in *mqo*, a malate-quinone oxidoreductase, associated with

azithromycin resistance. Despite appearing to have minimal interaction with known azithromycin

targets, studies testing the efficacy of azithromycin against the protozoan parasite *Plasmodium*

falciparum found reduced levels of TCA associated metabolites, including malate, upon

administration of the drug[58]. It is possible a similar effect is produced in gonococci and the

mutations in *mqo* may play a role in gonococcal survival during azithromycin treatment.

Additional work has been conducted in *P. falciparum* to validate *mqo* as a target for drug

development efforts, and our finding suggest similar efforts in *Ngo* may be worth pursuing[59].

We have curated a representative set of 485 genetically and globally diverse gonococcal genomes

from the 52,874 deposited gonococcal sequences on the NCBI SRA. Using PopNet, we assessed

the population structure of our representative set, and uncovered significant relationships between

population structure, geography, mobile elements, and AMR. This analysis suggests global

gonococcal genetic diversity is driven by the pressure applied by antimicrobials and links the

carriage of mobile elements with chromosomal variants. We used our representative set of

genomes to identify putative novel phase variable genes and were able to identify phase variation

of *opa* potentially contributing to improved survival in the context of infection. Using a combined

approach of GWAS and tests for selective pressure, we identified new sites and variants potentially

associated with AMR in *Ngo*, demonstrating the utility of these large-scale analyses for studying

AMR in pathogen populations.

MATERIALS & METHODS

Selection of sequences

An initial set of 52,874 accessions of sequencing runs were obtained from the NCBI Sequence Read Archive on September 26th, 2023 based on a keyword search for "*Neisseria gonorrhoeae*". This set was reduced to 31,595 by keeping only those accessions containing paired-end Illumina sequences with associated metadata for the country and year of isolation. Additional metadata recorded when available included the site of isolation, associated BioProject, and sex of the patient. All sequence data was obtained using the SRA Toolkit v3.0.0 using the default options for the prefetch and fasterq-dump commands. Further metadata, such as minimum inhibitory concentration values were obtained from pubMLST or publications associated with the relevant BioProjects[60]. Kraken2 v2.1.3 was run to identify erroneously annotated or contaminated sequences[61]. Any sequence datasets with less than 75% of its reads annotated by Kraken as being derived from *Ngo* were discarded. Any sequences with between 75 and 85% gonococcal reads were kept if over 95% of the reads were assigned to the genus *Neisseria.*

Variant discovery for raw sequence data

The reduced set of 31,595 isolates were aligned to the chromosomal genome of *Ngo* strain TUM19854, the current recommended reference genome for gonococci from the NCBI (obtained on October 13th, 2023). Alignment was performed using the Genome Analysis Toolkit (GATK) v4.2.5.0[62]. Quality control and trimming was performed using fastp v0.23.4 with minimum qualified phred quality of 20, a 40 percent limit on unqualified bases, a mean quality requirement of 20, a cut window size of 5, both front and tail cutting, and with correction enabled[63]. Trimmed reads were then passed to GATK for alignment using bwa v0.7.17 with default settings. The GATK Haplotype caller was then run with default settings to identify variants. Dependencies for GATK included samtools v1.17 and picard v2.26.3. Variants were then filtered using bcftools v1.11 and htslib v1.11 with the following parameters: allelic depth 8, allelic depth forward/reverse 3, minimum base quality 50, minimum mapping quality score 20 and percent of reads containing variant 80[64].

Generation of a representative set of sequences

Isolates were grouped by their country of isolation and year of isolation and the Jaccard statistic
was calculated for each pair of sequences using the bedtools v2.31.0 jaccard function with default
settings[65]. An initial clustering step was performed using the k-medoids fastPAM algorithm
implemented in the R packages cluster v2.1.5 and factoextra v1.0.7[66–68]. For each group, a value
for k was determined using the silhouette method, with the maximum value of k capped at 15. The
identified medoids of the clusters were selected as representative isolates. This resulted in an initial
set of 1,307 isolates. Isolates from this set were then grouped by their continent and year of
isolation, and k-medoids clustering was performed as before without a cap on the value of k.
Strains identified as disseminated or ocular as well as the WHO global reference collection were
added to the final set, bringing the total number of isolates to 485[14,15].

PopNet analysis and phylogenetic tree construction
PopNet was used to cluster the reference set of genomes using the variants identified from the
previous analysis[13]. A variant table was generated using the GATK VariantsToTable function from
a merged vcf file of identified variants. The parameters used for the PopNet analysis were
section_length= 5000, S1_iVal = 8, S1_piVal = 19, S2_iVal= 4.5, and S2_piVal= 2.5. These
parameters were determined using the silhouette index readout provided by the tool. Initial
phylogenetic tree reconstruction was performed with IQTree v2.2.2.7 from a sequence alignment
created from the previously identified variants[69]. To begin, Model Finder Plus was run which
identified TPM2u+F+I+R7 as the best substitution model[70]. IQTree was then run using this model
with 1000 ultrafast bootstrap replicates[71]. All other parameters were left as default. This initial tree
was used as a basis for ClonalFrameML v 1.12, which was run with the standard model with 100
pseudo-bootstrap replicates[16]. Assembled genomes were uploaded to pubMLST and automatically
assigned to a cgMLST Ng_300 clade.

Pangenome analysis
Trimmed reads were assembled *de novo* using unicycler v0.5.0 with default settings[18,72]. Gene
prediction and initial annotation was performed using Prokka v1.14.5 with recommended settings
and an added protein database containing annotated proteins from *Ngo* strains MS11, FA19,
FA1090 obtained from the NCBI on March 7th, 2022[20]. The pangenome was calculated using
panaroo v1.3.4 on the strict setting with otherwise default options[19]. Representative gene

sequences from panaroo were supplied to eggNOG-mapper v2.19 with no restrictions on orthology

identification and permitting electronic GO term annotation[73]. AMR genes were predicted from

sequences uploaded to PathogenWatch on January 22nd, 2024[23]. Mobile elements were predicted

by BLAST+ v2.0.13 against reference sequences AY803022.1, NC_001377, NC_002098,

GU479464.1, GU479465, and GU479466 which were obtained from the NCBI on December 16th,

2023[21]. The Graphia MCL clustering tool was used to cluster accessory genes based on shared

presence or absence in our representative collection with a granularity of 1.5[22].

Prediction of phase variable genes

Simple sequence repeats were identified from the *de novo* assemblies and the GFF files generated

by Prokka using PERF v0.4.6[25]. Genes with repeats that varied in the number of tandem repeats in

more than 2 isolates where each repeat length was present in at least 5% of genomes were

considered as potentially phase variable. Mutations classified as intergenic were manually verified

to ensure they impacted the associated gene.

Lower genital tract colonization for in vivo passaging

<u>Bacterial growth conditions:</u> *Ngo* MS11, parental strain, was grown on GC agar (Becton,

Dickinson and Company, Sparks, USA) plates supplemented with 1% Isovitalex (ISO, Becton,

Dickinson and Company, Sparks, USA) at 37˚C with 5% CO_2 for 16-20 hours. For *Ngo* isolates

used for in vivo passaging, agar plates included a vancomycin, colistin, nystatin, and

trimethoprim antibiotic mix (VCNT, Becton, Dickinson and Company, Sparks, USA) for *Ngo*

selection.

<u>Mouse strains:</u> 6-8 week old female FVB mice containing a human chromosome-derived BAC

expressing CEACAM3, -5 and -6, were used for all infection experiments[30]. Animals were housed

in cages with sterile rodent chow, sterile water, enrichment material, and 12-hour light-dark cycles.

<u>Gonococcal *in vivo* passaging:</u> Mice were staged by cytological vaginal smears observed under

light microscopy. Those in the diestrus stage of the estrous cycle were started on an antibiotic and

hormone regimen to control impact of commensal microflora on infection. Antibiotics consisted

of 0.6 mg vancomycin hydrochloride (Bioshop, Burlington, ON, Canada) and 2.4 mg streptomycin

sulfate (Sigma-Aldrich, Oakville, Canada) resuspended in 200 µL of PBS administered by
intraperitoneal injection daily from 2 days pre-infection (-2) to 5 days post infection (+5), where
an dose was provided on Day -1. Trimethoprim (Sigma-Aldrich, Oakville, Canada) was dissolved
in the drinking water at 40 mg/mL for ad libitum consumption. Water soluble β-estradiol
(SigmaAldrich, Oakville, Canada) was constituted in PBS (0.5 mg/200 µL), filter-sterilized and
administered via subcutaneous injection on Days -2, 0 and +2.

On Day 0, bacteria were resuspended in Dulbecco's phosphate-buffered saline (DPBS) with
calcium and magnesium (PBS++ , Life Technologies, Burlington, Canada)and mice were
inoculated with 10^7 bacteria in 2.5 µL Vaginal lavages were performed every two days by washing
the genital tract with 10 µL of PBS++ and plated for bacterial recovery and enumeration. After
~20 hrs bacterial isolates were frozen in 50% brain heart infusion (BHI, Becton, Dickinson and
Company, Sparks, USA) and 25% glycerol (EMD Millipore, Darmstadt, Germany) and stored at
-80˚C. Clearance of infection was characterized by 2 consecutive vaginal lavages that tested
negative for *Ngo*.

Gonococcal *in vitro* passaging: Five culture tubes containing 5 mL of BHI supplemented with 1%
ISO was inoculated with the parental strain with a starting optical density at 550 nm (OD550) of
0.05. Bacterial cultures were placed in a 37˚C shaker at 180 rpm. Every 12 hrs, bacterial stocks
were generated and sub-cultured for 5 consecutive days.

Ngo whole genome sequencing: *Ngo* isolates were resuspended in 200 µL of PBS++ and genomic
DNA was extracted by a QIAGEN DNeasy Blood and Tissue Kit (QIAGEN, Hilden, Germany)
following manufacturer instructions. DNA quality and quantity was checked by NanoDrop ND-
1000 Spectrophotometer (NanoDrop Technologies, Wilmington, USA) and Quant-iT$^{\text{TM}}$
PicoGreen® dsDNA Assay Kit (Life Technologies, Oregon, USA) was used to quantify amount
of dsDNA.Genomic DNA was prepared for whole genome sequencing by a Nextera XT library
preparation kit (Illumina, San Diego, USA). Libraries were run on a 1% agarose gel to ensure
libraries were of the correct length prior to poolingand submission to the Donnelly Sequencing
Centre, University of Toronto for sequencing using the Illumina MiSeq platform.

Capillary electrophoresis for confirmation of *opa* repeat lengths

Capillary electrophoresis was performed as described previously for *Neisseria meningitidis*[74].
Briefly, strains were cultured overnight on GC agar as described above, and 3 distinct colonies
per plate were selected and added to 20µl of Quick Extract buffer. Suspended colonies were then
heated at 66°C for 6 minutes, then 98°C for 2 minutes to complete the extraction. Extracted
DNA was amplified with PCR using the primers in **Table 3** listed below, with the forward
primer conjugated to a 6-FAM dye. Samples were then diluted to approximately 1-2ng/µl before
being sent for capillary electrophoresis. The results were analyzed with the ThermoFisher
MicroSatellite Analysis tool.

Identification of genes under positive selection

Aligned gene sequences from panaroo for core genes with 5 or more unique sequences were used
as input for selection analyses after removing identical sequences and sequences containing
premature stop codons using Hyphy rmv[32]. First, a tree for each gene was built using IQTree
v2.2.2.7 with 1000 ultrafast bootstraps and otherwise default settings[69]. The BUSTED algorithm
was then run as implemented in HyPhy with default settings for each gene using the generated tree
and panaroo alignment[33]. Genes identified as being under positive selection were then analyzed
with FUBAR as implemented in HyPhy with default settings using the same inputs to identify
specific sites under positive selection[34]. To assess the location of predicted residues under positive
selection, structural models were generated with AlphaFold3 using the AlphaFold server based on
the consensus gene sequences from panaroo[19,75]. Predicted structures were visualized with UCSF
ChimeraX v1.9[76].

Genome wide association study

The genome wide association study was performed using pyseer with the linear mixed model
setting using the VCF file generated as described above[44]. The distance matrix was calculated
using the ClonalFrameML generated tree described above. All other pyseer settings were kept at
default. The count_patterns script was used to establish a significance threshold for each tested
phenotype. The impact of each variant was predicted with SnpEff using the *Ngo* TUM19854
reference genome obtained as previously described[77]. Structural alignments and comparisons
were performed using TM-align with default settings[45].

ACKNOWLEDGEMENTS

We are grateful to Odile Harrison, University of Oxford, for helpful advice and technical support, and the Division of Comparative Medicine & Donnelly Sequencing Centre at the University of Toronto for technical support. This study was supported by operating grants from the Canadian Institutes for Health Research (PJT-159677) and National Institutes of Health (1R01AI146941-01A1). S.D.G. is supported by the Canada Research Chairs Program. D.C. is a recipient of the EPIC Doctoral Award and the DSI Doctoral Fellowship.

ETHICS STATEMENT

All animal procedures conducted in this study were approved by the Local Animal Care Committee (LACC) at the University of Toronto (Protocol #20011775), which is in compliance with ethical and legal requirements under Ontario's Animals for Research Act and the federal Canadian Council on Animal Care.

DATA & CODE AVAILABILITY

The data sets generated in this study are provided in the Supplementary Information and Source Data files. Codes for analysis and data are available on Github [https://github.com/ParkinsonLab/Neisseria_work.git].

AUTHOR CONTRIBUTIONS

J.P. and S.D.G conceived the idea of the research and supervised it. D.C and E.J. performed the computational and data analysis. D.C. and J.M. conducted the mouse passaging and phase variation experiments and analyzed the data. D.C. wrote the manuscript with contributions from J.P., S.G.D. & J.L.

REFERENCES

1.   Mayor, M. T., Roett, M. A. & Uduhiri, K. A. Diagnosis and Management of Gonococcal Infections. *Am. Fam. Physician* **86**, 931–938 (2012).

2.   Bernstein, K. T., Marcus, J. L., Nieri, G., Philip, S. S. & Klausner, J. D. Rectal Gonorrhea and Chlamydia Reinfection Is Associated With Increased Risk of HIV Seroconversion. *JAIDS J. Acquir. Immune Defic. Syndr.* **53**, 537–543 (2010).

3.   Zhang, J. *et al. Neisseria gonorrhoeae* Enhances Infection of Dendritic Cells by HIV
Type 1. *J. Immunol.* **174**, 7995–8002 (2005).

4.   *WHO Bacterial Priority Pathogens List, 2024: Bacterial Pathogens of Public Health*
Importance to Guide Research, Development and Strategies to Prevent and Control
Antimicrobial Resistance. https://iris.who.int/bitstream/handle/10665/376776/9789240093461-
eng.pdf?sequence=1 (2024).

5.   Quillin, S. J. & Seifert, H. S. *Neisseria gonorrhoeae* host adaptation and pathogenesis.
Nat. Rev. Microbiol. **16**, 226–240 (2018).

6.   Ohnishi, M. *et al.* Spread of a chromosomal cefixime-resistant penA gene among
different *Neisseria gonorrhoeae* lineages. *Antimicrob. Agents Chemother.* **54**, 1060–1067 (2010).

7.   Harrison, O. B. *et al. Neisseria* genomics: current status and future perspectives. *Pathog.*
Dis. **75**, 60–60 (2017).

8.   Cehovin, A., Lewis, S. B. & William, S. Mobile genetic elements in *Neisseria*
gonorrhoeae: movement for change. **75**, 71–71 (2017).

9.   Ramsey, M. E., Woodhams, K. L. & Dillard, J. P. The gonococcal genetic island and type
IV secretion in the pathogenic *Neisseria. Front. Microbiol.* **2**, 61–61 (2011).

10.   Siena, E. *et al.* In-silico prediction and deep-DNA sequencing validation indicate phase
variation in 115 *Neisseria meningitidis* genes. *BMC Genomics* **17**, 843 (2016).

11.   Jolley, K. A. Multi-locus sequence typing. *Methods Mol. Med.* **67**, 173–186 (2001).

12.   Harrison, O. B. *et al. Neisseria gonorrhoeae* Population Genomics: Use of the
Gonococcal Core Genome to Improve Surveillance of Antimicrobial Resistance. *J. Infect. Dis.*
222, 1816–1825 (2020).

13.   Zhang, J., Khan, A., Kennard, A., Grigg, M. E. & Parkinson, J. PopNet: A Markov
Clustering Approach to Study Population Genetic Structure. *Mol. Biol. Evol.* **34**, 1799–1811
(2017).

14.   Unemo, M. *et al.* The novel 2016 WHO *Neisseria gonorrhoeae* reference strains for
global quality assurance of laboratory investigations: phenotypic, genetic and reference genome
characterization. *J. Antimicrob. Chemother.* **71**, 3096–3096 (2016).

15.   Phenotypic and genetic characterization of the 2008 WHO *Neisseria gonorrhoeae*
reference strain panel intended for global quality assurance and quality control of gonococcal
antimicrobial resistance surveillance for public health purposes | Journal of Antimicrobial
Chemotherapy | Oxford Academic. https://academic.oup.com/jac/article/63/6/1142/747767.

16.   Didelot, X. & Wilson, D. J. ClonalFrameML: Efficient Inference of Recombination in
Whole Bacterial Genomes. *PLOS Comput. Biol.* **11**, e1004041–e1004041 (2015).

17.   Lees, J. A. *et al.* Fast and flexible bacterial genomic epidemiology with PopPUNK.
Genome Res. **29**, 304–316 (2019).

18.   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. *PLOS Comput. Biol.* **13**, e1005595
(2017).

19.   Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo
pipeline. *Genome Biol.* **21**, 180 (2020).

20.   Prokka: rapid prokaryotic genome annotation | Bioinformatics | Oxford Academic.
https://academic.oup.com/bioinformatics/article/30/14/2068/2390517.

21.   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
(2009).

22.   Freeman, T. C. *et al.* Graphia: A platform for the graph-based visualisation and analysis
of high dimensional data. *PLoS Comput. Biol.* **18**, e1010310 (2022).

23.   Sánchez-Busó, L. *et al.* A community-driven resource for genomic epidemiology and
antimicrobial resistance prediction of *Neisseria gonorrhoeae* at Pathogenwatch. *Genome Med.*
13, 61 (2021).

24.   European Committee on Antimicrobial Susceptibility Testing (EUCAST). Breakpoint
tables for interpretation of MICs and zone diameters. Version 9.0.

25.   Avvaru, A. K., Sowpati, D. T. & Mishra, R. K. PERF: an exhaustive algorithm for ultra-
fast and efficient identification of microsatellites from large DNA sequences. *Bioinformatics* **34**,
943–948 (2018).

26.   Wanford, J. J., Green, L. R., Aidley, J. & Bayliss, C. D. Phasome analysis of pathogenic
and commensal *Neisseria* species expands the known repertoire of phase variable genes, and
highlights common adaptive strategies. *PLoS ONE* **13**, e0196675 (2018).

27.   The meningococcal autotransporter AutA is implicated in autoaggregation and biofilm
formation - PubMed. https://pubmed.ncbi.nlm.nih.gov/25059714/.

28.   Pachulec, E. & van der Does, C. Conjugative Plasmids of *Neisseria gonorrhoeae*. *PLoS*
ONE **5**, e9962 (2010).

29.   Zelewska, M. A. *et al.* Phase variable DNA repeats in *Neisseria gonorrhoeae* influence
transcription, translation, and protein sequence variation. *Microb. Genomics* **2**, e000078 (2016).

30.   Islam, E. A. *et al.* Specific binding to differentially expressed human carcinoembryonic
antigenrelated cell adhesion molecules determines the outcome of *Neisseria gonorrhoeae*
infections along the female reproductive tract. *Infect. Immun.* **86**, (2018).

31.   Erwin, A. L., Haynes, P. A., Rice, P. A. & Gotschlich, E. C. Conservation of the
lipooligosaccharide synthesis locus lgt among strains of *Neisseria gonorrhoeae*: requirement for
lgtE in synthesis of the 2C7 epitope and of the beta chain of strain 15253. *J. Exp. Med.* **184**,
1233–1241 (1996).

32. Pond, S. L. K., Frost, S. D. W. & Muse, S. V. HyPhy: hypothesis testing using
phylogenies. *Bioinformatics* **21**, 676–679 (2005).

33. Gene-Wide Identification of Episodic Selection | Molecular Biology and Evolution |
Oxford Academic. https://academic.oup.com/mbe/article/32/5/1365/1134918.

34. Murrell, B. *et al.* FUBAR: A Fast, Unconstrained Bayesian AppRoximation for Inferring
Selection. *Mol. Biol. Evol.* **30**, 1196–1205 (2013).

35. Belland, R. J., Morrison, S. G., Ison, C. & Huang, W. M. *Neisseria gonorrhoeae* acquires
mutations in analogous regions of gyrA and parC in fluoroquinolone-resistant isolates. *Mol.*
Microbiol. **14**, 371–380 (1994).

36. Collins, J. A., Oviatt, A. A., Chan, P. F. & Osheroff, N. Target-Mediated
Fluoroquinolone Resistance in *Neisseria gonorrhoeae*: Actions of Ciprofloxacin against Gyrase
and Topoisomerase IV. *ACS Infect. Dis.* **10**, 1351–1360 (2024).

37. The InterPro database, an integrated documentation resource for protein families,
domains and functional sites - PMC. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC29841/.

38. Corbett, K. D., Schoeffler, A. J., Thomsen, N. D. & Berger, J. M. The Structural Basis for
Substrate Specificity in DNA Topoisomerase IV. *J. Mol. Biol.* **351**, 545–561 (2005).

39. Wadsworth, C. B., Arnold, B. J., Sater, M. R. A. & Grad, Y. H. Azithromycin resistance
through interspecific acquisition of an epistasis-dependent efflux pump component and
transcriptional regulator in *Neisseria gonorrhoeae*. *mBio* **9**, (2018).

40. Powell, A. J., Tomberg, J., Deacon, A. M., Nicholas, R. A. & Davies, C. Crystal
Structures of Penicillin-binding Protein 2 from Penicillin-susceptible and -resistant Strains of
Neisseria gonorrhoeae Reveal an Unexpectedly Subtle Mechanism for Antibiotic Resistance. *J.*
Biol. Chem. **284**, 1202–1212 (2009).

41. Fenton, B. A. *et al.* Mutations in PBP2 from ceftriaxone-resistant *Neisseria gonorrhoeae*
alter the dynamics of the β3–β4 loop to favor a low-affinity drug-binding state. *J. Biol. Chem.*
297, 101188 (2021).

42. Malekian, N., Agrawal, A. A., Berendonk, T. U., Al-Fatlawi, A. & Schroeder, M. A
genome-wide scan of wastewater *E. coli* for genes under positive selection: focusing on
mechanisms of antibiotic resistance. *Sci. Rep.* **12**, 8037 (2022).

43. A genome-wide association analysis reveals a potential role for recombination in the
evolution of antimicrobial resistance in *Burkholderia multivorans* | PLOS Pathogens.
https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1007453.

44. Lees, J. A., Galardini, M., Bentley, S. D., Weiser, J. N. & Corander, J. pyseer: a
comprehensive tool for microbial pangenome-wide association studies. *Bioinformatics* **34**, 4310–
4312 (2018).

45. Bittrich, S., Segura, J., Duarte, J. M., Burley, S. K. & Rose, Y. RCSB protein Data Bank:
exploring protein 3D similarities via comprehensive structural alignments. *Bioinformatics* **40**,
btae370 (2024).

46. Varadi, M. *et al.* AlphaFold Protein Structure Database in 2024: providing structure
coverage for over 214 million protein sequences. *Nucleic Acids Res.* **52**, D368–D375 (2024).

47. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583–589 (2021).

48. Ng, L.-K., Martin, I., Liu, G. & Bryden, L. Mutation in 23S rRNA Associated with
Macrolide Resistance in *Neisseria gonorrhoeae*. *Antimicrob. Agents Chemother.* **46**, 3020–3025
(2002).

49. de Korne-Elenbaas, J., Bruisten, S. M., van Dam, A. P., Maiden, M. C. J. & Harrison, O.
B. The *Neisseria gonorrhoeae* Accessory Genome and Its Association with the Core Genome
and Antimicrobial Resistance. *Microbiol. Spectr.* **10**, e02654-21.

50. Pinto, M. *et al. Neisseria gonorrhoeae* clustering to reveal major European whole-
genome-sequencing-based genogroups in association with antimicrobial resistance. *Microb.*
Genomics **7**, 000481 (2020).

51. Cehovin, A., Jolley, K. A., Maiden, M. C. J., Harrison, O. B. & Tang, C. M. Association
of *Neisseria gonorrhoeae* Plasmids With Distinct Lineages and The Economic Status of Their
Country of Origin. *J. Infect. Dis.* **222**, 1826–1836 (2020).

52. San Millan, A. & MacLean, R. C. Fitness Costs of Plasmids: a Limit to Plasmid
Transmission. *Microbiol. Spectr.* **5**, (2017).

53. Gulati, S., McQuillen, D. P., Mandrell, R. E., Jani, D. B. & Rice, P. A. Immunogenicity
of *Neisseria gonorrhoeae* Lipooligosaccharide Epitope 2C7, Widely Expressed In Vivo with No
Immunochemical Similarity to Human Glycosphingolipids. *J. Infect. Dis.* **174**, 1223–1237
(1996).

54. Yu, D. *et al.* A Genome-Wide Identification of Genes Undergoing Recombination and
Positive Selection in *Neisseria*. *BioMed Res. Int.* **2014**, 815672 (2014).

55. Wang, S. *et al.* Next Generation Sequencing and Comparative Genomic Analysis Reveal
Extreme Plasticity of Two *Burkholderia glumae* Strains HN1 and HN2. *Pathogens* **11**, 1265
(2022).

56. Schubert, B., Maddamsetti, R., Nyman, J., Farhat, M. R. & Marks, D. S. Genome-wide
discovery of epistatic loci affecting antibiotic resistance in *Neisseria gonorrhoeae* using
evolutionary couplings. *Nat. Microbiol.* **4**, 328–338 (2019).

57. Chopra, I. & Roberts, M. Tetracycline Antibiotics: Mode of Action, Applications,
Molecular Biology, and Epidemiology of Bacterial Resistance. *Microbiol. Mol. Biol. Rev.* **65**,
232 (2001).

58.   Burns, A. L. *et al.* Retargeting azithromycin analogues to have dual-modality antimalarial
activity. *BMC Biol.* **18**, 133 (2020).

59.   Wang, X. *et al.* Identification of Plasmodium falciparum Mitochondrial Malate: Quinone
Oxidoreductase Inhibitors from the Pathogen Box. *Genes* **10**, 471 (2019).

60.   Jolley, K. A., Bray, J. E. & Maiden, M. C. J. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res.* **3**, 124
(2018).

61.   Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol. **20**, 257 (2019).

62.   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1297 (2010).

63.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

64.   Li, H. A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987–
2993 (2011).

65.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841–842 (2010).

66.   Schubert, E. & Rousseeuw, P. J. Fast and eager k-medoids clustering: O (k) runtime
improvement of the PAM, CLARA, and CLARANS algorithms. *Inf. Syst.* **101**, 101804 (2021).

67.   Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M. & Hornik, K. cluster: Cluster
Analysis Basics and Extensions. (2023).

68.   Kassambara, A. & Mundt, F. factoextra: Extract and Visualize the Results of Multivariate
Data Analyses. (2020).

69.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

70.   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–
589 (2017).

71.   Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

72.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications
to Single-Cell Sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

73.   eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale | Molecular Biology and Evolution | Oxford Academic.
https://academic.oup.com/mbe/article/38/12/5825/6379734.

74.   Jen, F. E.-C., Seib, K. L. & Tan, A. Screening DNA Repeat Tracts of Phase Variable
Genes by Fragment Analysis. in *Neisseria meningitidis: Methods and Protocols* (eds. Seib, K. L.
& Peak, I. R.) 93–104 (Springer, New York, NY, 2019). doi:10.1007/978-1-4939-9202-7_6.

75.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500 (2024).

76.   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

77.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff. *Fly (Austin)* **6**, 80–92 (2012).

TABLES

Table 1: Predicted new phase variable genes.

Gene	Repeat	No. of repeats	Annotation	NGK locus	NGO locus
group_1584	A	33	Pertactin	NGK_2016	NGO_1158
group_280	AACGC	24	Hypothetical protein	-	NGO_0366
argS	AAGC	18	Arginyl-tRNA synthetase	NGK_0832	NGO_0963
group_1317	AACGC	17	C-5 cytosine-specific DNA methylase	NGK_0522	NGO_0365
group_1449	AT	11	Spermidine synthase	NGK_2156	NGO_1752
group_1206	C	11	ISXO2-like transposase domain	NGK_1147	NGO_1157
kfrB	C	10	-	-	-
glyS	C	9	Glycyl-tRNA synthetase beta	NGK_2629	NGO_2154
prn	A	8	Autotransporter	NGK_1206	NGO_0694
yigL	C	7	Hydrolase	NGK_1967	-

874 **Table 2: Predicted phase variable genes from mouse passaging experiment.**

Gene	Repeat	No. of repeats	Annotation	NGK locus	NGO locus
group_1059	C	3	Pilus assembly protein	NGK_0074	NGO_0055
group_1287	C	2	Pilus assembly protein	NGK_0074	NGO_0055
group_399	CCTCT	2	Opacity protein	NGK_0693	NGO_07725
group_579	CCTCT	2	Opacity protein	NGK_0693	NGO_1513
group_784	C	3	Enterobactin receptor *fetA*	NGK_2557	NGO_2093
group_1283	CCTCT	3	Opacity-associated protein	NGK_0749	NGO_08230
group_973	AGCC	2	Adenine specific DNA methylase	NGK_1273	NGO_0641
group_1471	CCTCT	2	Opacity-associated protein	NGK_2410	NGO_1513
group_1111	C	2	Glycosyl transferase family 1	NGK_2534	NGO_2072

877   **Table 3: Primers used for capillary electrophoresis.**

Primer	Modification	Sequence
Opa-For	5' 6-FAM	CAACATCAGTGAAAATCGTTCTTTT
Opa-Rev	None	TCGTGGGTAATGTGTTCGTAG

<u>FIGURES</u>

Figure 1: Overview of representative set of 485 *N. gonorrhoeae* genomes.

a Schematic of selection of representative genomes. **b** Geographic locations of isolates in the

representative set. **c** Geographic and temporal distribution of isolates in the representative set. **d**

Comparison of the geographic distribution of isolates between the initial set of 52,784 SRA runs

and the representative set of 485.

Figure 2: Population structure of *N. gonorrhoeae* is influenced by AMR and the presence of

mobile elements.

a PopNet network view of 485 representative *Ngo* genomes. Nodes represent genomes, coloured

by their assigned clades. Annuli show the chromosome painting generated for each genome, with

regions sharing a colour showing overlap between genomes. A selection of genomes from clades

B, E, I & K are shown to illustrate the chromosome painting visualization. **b** ClonalFrameML

generate phylogenetic tree of representative *Ngo* genomes. Outer rings show the assigned

PopNet clade, MDR (multi-drug-resistant) status and continent of isolation for each genome.

Top: MDR status overlaid on the PopNet network view. Bottom: Continent of isolation overlaid

on the PopNet network view. AZM: azithromycin, CFM: cefixime, XDR: extreme drug resistant.

c Association between antimicrobial resistance (left, i) or mobile elements (right, ii) and PopNet

clades. Only significant associations with a p-value < 0.05 as determined by Fisher's exact test

with Bonferonni multiple testing correction are shown.

Figure 3: The pangenome of *N. gonorrhoeae* is dominated by large mobile elements, which are associated with PopNet clades determined by chromosomal variants.

a Pangenome of *Ngo* is composed of 2460 genes, with 1800 Core (present in > 95% of genomes), 367 Shell (present in between 15% and 95% of genomes) & 283 Cloud (present in <15% of genomes) genes. Dendrogram and ordering of genomes on the y-isolates calculated using complete linkage from a Euclidian distance matrix generated based on shared gene content. **b** Top: Network view of isolates clustered by shared gene content coloured by the presence or absence of 2 large mobile elements: the GGI (gonococcal genetic island) and the pConj (conjugative plasmid). Bottom: Genomes coloured by PopNet clade overlaid on the network view of isolates clustered by shared gene content. **c** Clustering of genes based on their co-occurrence in genomes. Clusters defined by MCL clustering as implemented in Graphia which are present in <50% of genomes and composed of 5 or more genes are coloured. **d** Organization of genes from MCL cluster 11 associated with clade D.

Figure 4: Passaging of *N. gonorrhoeae* improves colonization in a mouse model of infection

which may be associated with phase variation of *opa*.

a Schematic overview of *in vivo* and *in vitro* passaging experiment. Coloured mice represent

mice successfully colonized by *Ngo* 2 days post infection. Coloured gonococci indicate isolates

sequenced by capillary electrophoresis to confirm the repeat lengths of *opa*. **b** Recovered CFU

(colony-forming units) of *Ngo* recovered 2 days post infection. Black bars represent the mean

CFUs recovered for each passage. **c** Ratio of *opa* repeats in each isolate sequenced by capillary

electrophoresis. **d** Schematic of *Ngo* lipooligosaccharide with (bottom) and without (top) lgtG

expression, resulting in the production of the 2C7 epitope. Isolates passaged *in vivo* express lgtG,

while no expression is observed in isolates passaged *in vitro.* Immunoblot assay for the 2C7

isotope confirms the expression of lgtG in a passage 3 isolate, which is absent from the parental

isolate. KDO: 3-Deoxy-D-manno-oct-2-ulosonic. Hep: Heptose. Glc: Glucose. Gal: Galactose.

GlcNAc: N-acetylglucosamine. GalNAc: N-Acetylgalactosamine.

Figure 5: GWAS and tests for positive selection identify sites linked to AMR in *N. gonorrhoeae*.

a Boxplot of LRT (likelihood ratio test statistic) of positive selection for *Ngo* genes for predicted cellular localization. **b** Predicted residues with strong signatures of positive selection in *parC*. Residues highlighted in green are under positive selection and known to contribute to ciprofloxacin resistance. Residues in blue are under positive selection with no known links to AMR. Structural prediction from AlphaFold3 visualized with USCF ChimeraX. **c** Predicted residues with strong signatures of positive selection in *penA*. Residues in blue are under positive selection with no known links to AMR. Structural prediction from Alphafold3 visualized with USCF ChimeraX. **d** Manhattan plots of variants with significant associations with ciprofloxacin (top) and azithromycin (bottom) resistance. Only variants with an LRT p-value <0.05 are shown. Red lines indicate significance thresholds determined by Bonferroni correction with the number of unique variant patterns as the number of multiple tests. **e** Pairwise alignment of AlphaFold predicted structures for Mqo (yellow) with RplV (left, blue) and RplD (right, blue) using RSCPD TM-align. Darker residues indicate regions of structural overlap.