Viral lifecycle dynamics and spatial structure explain why pathogenicity is prophage-encoded

Sultan A. Nazir[1,*] and Bram van Dijk[1]

[1]Theoretical Biology, Utrecht University, The Netherlands
*Corresponding author: s.a.nazir@uu.nl

May 13, 2025

Abstract

Genes encoding for virulence factors are frequently found on prophages, yet the evolutionary forces driving this association remain unclear. The evolutionary association of mobile genetic elements (MGEs) with host-beneficial genes is known to be hindered by the stability of chromosomal loci and competition with MGEs lacking accessory genes. Using a mathematical model that incorporates these constraints, we identify two key mechanisms that help overcome these barriers, resulting in the evolutionary linkage of accessory genes, like virulence genes, to prophages. First, we show that migration across bacterial populations favours the association of genes with prophages when the gene is beneficial in certain environments (e.g. virulence genes in the gut) that also trigger higher rates of prophage induction. Second, we show that within-population spatial dynamics also promotes the association of weakly selected genes and phages. Here, virion dispersal allows phage-encoded genes to spread into patches of bacteria lacking the gene, giving them a selective advantage over immobile chromosomal genes. We argue that these mechanisms are less applicable for plasmids and other MGEs, highlighting a potentially unique role for phages in shaping bacterial adaptation. By demonstrating how phage lifecycle dynamics and spatial heterogeneity drive MGE-gene associations, our work provides new insights into the evolution of phage-encoded virulence.

Keywords: prophages, virulence, mobile genetic elements, spatial structure, induction, dispersal

Background

Mobile genetic elements (MGEs) are genome segments that can move within or between genomes, often replicating independently of the chromosome. Although MGEs such as bacteriophages, plasmids, and transposons have been the focus of extensive research for several decades, their eco-evolutionary functions are still being elucidated [1, 2]. Increasingly, studies reveal the crucial roles that MGEs play in shaping the ecological and clinical functions of microbial communities [3, 4]. MGEs are especially relevant because they can carry accessory genes, which are non-essential for the MGE itself, but may substantially alter the bacterial phenotype. Different types of MGEs appear to vary in their likelihood of carrying specific accessory genes [3], but the mechanisms behind these associations remain poorly understood. For example, a recent analysis shows that phages are more likely to encode virulence factors than plasmids [5]. This raises the question: what makes phages uniquely predisposed to turn their hosts into pathogens?

Different hypotheses have been proposed to explain why non-essential and cooperative traits can be found on MGEs rather than on chromosomes (or immobile parts of the genome) [3]. For phage-encoded pathogenicity in particular, Abedon and LeJeune have written an excellent review of different hypothesised and empirically supported mechanisms [6]. Models have been a useful tool to unpack the conditions under which MGEs carry certain accessory genes. For example, Bergstrom et al. proposed a local adaptation hypothesis, wherein plasmid-encoded genes persist by transferring locally adapted genes to newly arriving strains [7]. Later, Smith proposed the infectivity hypothesis, which stated that MGEs may evolve to carry genes for public goods in order to force defectors to cooperate [8]. More recently, van Dijk et al. have shown that HGT can rescue a beneficial gene (therewith increasing population fitness) when the benefit is not strong enough to maintain the gene against mutational loss [9]. From these studies, we can generalise a *rescue hypothesis*[10]: genes are MGE-encoded when horizontal gene transfer (HGT) can rescue the gene from being lost through negative selection, mutational loss or the emergence of cheaters.

While the above-mentioned models and the rescue hypothesis are insightful, they do not fully capture the biology of how MGE-driven mobilisation comes about. For an accessory gene to become mobilised by residing on an MGE, its mobility must not only be beneficial, it must also persist under competition against MGEs that do not encode the accessory gene. Competition with such "non-carrier" MGEs has indeed been shown to hinder the association of MGEs with accessory genes [11, 12]. In addition, most mechanisms of gene loss will act on the mobile locus as much as they do on the chromosomal locus, and the gene may fail to persist by mobility alone.

More recent work by Lehtinen et al. has shown that an accessory gene that is initially more abundant on the mobile (MGE) locus than on the immobile (chromosomal) locus will, under weak selection, stay on the mobile locus [13]. This "priority effect" may emerge when a species is more likely to acquire the gene via an MGE than by mutational innovation on the chromosome. While this mechanism is of likely importance to plasmids which have host ranges spanning broad taxa [14], temperate phages are generally thought to have more narrow host ranges, often limited to strains of a single species, due to

their specificity during integration [15–17]. Moreover, there is evidence that genes associated with prophages have a chromosomal origin due to their localisation close to the attachment site of the prophage DNA [18]. In other words, while much research on MGEs was done with plasmids as a model MGE – implicitly assuming generalisability to other MGEs – few models have been developed to enhance our understanding of phage-encoded accessory traits.

In this article, we focus on understanding why a beneficial chromosomal gene can evolve to be associated with bacteriophages by means of mathematical modelling. While our model could be applied to accessory genes more generally, we built our model especially in the context of exotoxin production, a very common virulence factor encoded by prophages [19, 20]. The benefits provided by these extracellular virulence factors may be partially enjoyed by other bacteria that also gain access to resources of the plant or animal through tissue damage. Therefore, we model the expression of the gene as a cooperative trait with certain degree of privatisation of benefits in order to maintain a positive selection pressure. As a preliminary analysis, our mathematical model verifies that the "rescue" of certain beneficial genes by lysogenic phages is highly sensitive to competition with non-carrier phages. In competition with non-carrier phages, we find that phage-encoded genes cannot invade when the gene is already chromosomally encoded. However, we find that the gene-phage association can invade under certain conditions within two distinct implementations of spatial structure. Firstly, by extending our ODE model to include a host-environment metapopulation structure, we find that when prophage induction is upregulated in the host environment where the benefits of the gene are limited to, natural selection favours the beneficial gene to be phage-encoded. Secondly, using an individual-based model, we propose virion dispersal within a spatially structured population as a mechanism that catalyses the evolution of phage-encoded cargo genes under weak selection. Here, since natural selection occurs at a local scale, phage particles help the gene disperse to pockets lacking the gene, reinforce selection and replace the chromosomal gene at a global scale. Taken together, our work shines light on the exogenous factors that contribute to the evolution of phage-encoded accessory traits, explaining the observed association of virulence genes and prophages.

Methods

Infection cycle and genotype space: We implemented an infection model, as shown in Figure 1(A), where uninfected prokaryotic cells (denoted by U) are susceptible to infection by virion particles (denoted by V) such that the viral DNA integrates into the cell as a prophage and converts each cell into a lysogen (denoted by L). Then, the prophage is induced at a rate α which causes cell lysis and the release virion particles equal to its burst size r_V. Assuming they occur simultaneously, we ignore the lag in time that occurs between infection and integration and between induction and lysis. Virion particles in the external environment can etiher decay at a rate δ or infect a susceptible cell at a rate β. In this framework, we assume superinfection immunity such that a cell cannot be infected by a virion particle if it already harbours a prophage.

The viral DNA may or may not carry the gene of interest. If the virion particle is a carrier of the gene, it is marked by a

subscript τ. When a non-carrier virion V infects a susceptible cell, it creates a non-carrier prophage, but if a carrier virion V_τ does, it creates a carrier prophage, and a lysogen with a carrier prophage is indicated by a subscript p to denote that the gene is phage encoded. Every cell, regardless of whether it is infected by a prophage or not, has a chromosomal locus where the gene may or may not be present. If the cell is a carrier of the gene at the chromosomal locus, it is indicated by a subscript c. Hence, as shown in Figure 1(B), there are six possible genotypes of prokaryotic cells (U, U_c, L, L_c, L_p, and L_{cp}) and two kinds of virion particles (V and V_τ).

Cooperation and privatisation: All of the cells grow at a fixed rate and compete for resources along with an intrinsic death rate of d. The effect of carrying the gene of interest on the growth dynamics is modelled as follows. We define "minimum carrying capacity" as the carrying capacity in the absence of the gene and set its value to one for non-dimensionalisation, i.e. all population sizes are expressed relative to the minimum carrying capacity. Let ϕ represent the maximum additional biomass that can be generated due to gene expression. If f_τ denotes the fraction of carriers in the population, by the definition of ϕ, the net carrying capacity becomes $1 + \phi f_\tau$. If the benefits produced by the carriers are equally shared among all cells, each cell receives a benefit of ϕf_τ. However, positive assortment and other mechanisms enable cooperators to have disproportionate access to these benefits. To model this, as shown in Figure 1(C), we introduce the parameter π, the fraction of the benefits produced by a carrier that is privatised, i.e. retained exclusively by the producer. Consequently, all cells (including carriers themselves) receive a shared benefit of $(1-\pi)\phi f_\tau$, while carriers gain an additional private benefit of $\pi\phi$. This framework allows π to function as a "privatisation parameter", or more generally, for tuning the rate of selection acting on the gene.

Mutations: In the model, we consider two kinds of deletion mutations, as shown in Figure 1(D). Firstly, every "carrier" (henceforth used to refer to a cell that has the gene of interest) can mutate into a "non-carrier" by gene loss at a rate μ. Note that this mutation occurs at the prophage locus and the chromosomal locus, at the same rate. Secondly, every lysogen can lose its prophage at a rate λ. For simplicity, we assume that this loss coincides with the loss of superimmunity, i.e. the cell will immediately be available for future phage infections. If a cargo gene resided on the prophage, this gene is lost along with the phage.

The above processes are first analysed within a framework of differential equations (Appendix A), where we focus on the ability of a phage-encoded trait to replace a chromosomally-encoded trait. In addition, this analytical model is compared and contrasted with an individual-based framework, described in the next section. The parameters and their values are summarised in Appendix B.

Spatial model

For the spatially structured version of the model, we implemented an Individual based model (IBM) of a finite population of bacteria on a square grid. The model captures the same processes described above using the ODEs while introducing spatial structure and finiteness to the cell and virus populations with the following elements.

Figure 1: **An illustration of all of the model components:** (A) The lifecycle of the phage consists of a prophage stage that begins upon infecting a susceptible bacterium and a virion state which is produced upon induction of the prophage. As a virion, the phage can be lost to environmental degradation. (B) Based on whether the infecting phage carries the gene of interest or not, and whether the susceptible cell carries the gene at the chromosomal locus, we get six cell genotypes and four of them carry the gene. The subscript denotes whether the gene is present on the chromosomal locus (c) or the phage locus (p). (C) Out of the ϕ benefits produced by a carrier, π fraction is privatised by the producer and the remaining is shared with the whole cell population. (D) Both the gene and the phage can be lost from a microbial genome due to mutations.

Birth and Death: A cell can copy itself within a neighbouring empty site with a probability b as shown in Figure 2(A). We assume the rate of birth is equal for all cells at all times. The effect of cooperative interations is expressed in terms of varying death rates. We chose to vary the death rates instead of the birth rates, because birth is inherently density-dependent due to our assumption that reproduction requires empty sites. In the absence of any benefit from carriers, a cell dies with probability $d_s + \Delta d$ where d_s is the intrinsic death rate in the spatial model and Δd is the rate of death that can be avoided with benefits from the gene. The maximum population size is limited by the size of the grid (total number of grid points). Therefore, to simulate dynamics comparable to the ODE, we adjust the death rate parameters such that the grid size (the maximum carrying capacity) is $1 + \phi$ times the average steady population size in the absence of the gene (the minimum carrying capacity). Analogously to the ODE model, the benefits of the cooperative gene are partially privatised. A carrier privatise π fraction of its benefits to itself (lowering its own chance of

death), with the remaining fraction $(1 - \pi)$ of benefits shared with others. Therefore, the probability of death is

$$\mathbb{P}_{\text{death}}(\pi) = d_s + \Delta d\Big(1 - \pi\mathbb{1}_{\text{carrier}} - (1 - \pi)f_\tau\Big)$$

where $\mathbb{1}_{\text{carrier}}$ is the indicator variable

$$\mathbb{1}_{\text{carrier}} = \begin{cases} 1, & \text{cell is a carrier} \\ 0, & \text{cell is not a carrier} \end{cases}$$

and f_τ is the fraction of carriers on the grid.

To ensure that the ODE is the a good approximation of the spatial model when well-mixed, all the rate parameters in the IBM are scaled by a factor of $\frac{b}{1+\phi}$ (for example, $d_s = d\frac{b}{1+\phi}$) and we set $\Delta d = \phi\frac{b}{1+\phi}$. See Appendix C for verification of the well-mixed approximation.

Phage lifecycle: Extracellular virions are also modelled as discrete entities. Therefore, at every grid point, we have an integer measure of the number of extracellular virion particles

Figure 2: **Spatial model:** (A) Individual cells are placed on a square grid and the occupied grid points are coloured according to the cell genotype (identical to the colours in Figure 1(B). On the grid, the cells can reproduce into an unoccupied grid point in its Moore neighbourhood. The cells can also die, leaving an unoccupied grid point. The propensity of death reduces with the beneifts received from the gene. If a cell is a carrier, it privatises π fraction of benefits and shares the remaining $1 - \pi$ fraction with other cells on the grid. (B) Apart from the 'Cell grid', there is a 'Virion grid' where more than one virion particle can occupy a single grid point. They infect cells and undergo part of their lifecycle as prophages on the Cell grid. Upon induction and lysis, virions are released onto the Virion grid where it can diffuse and ultilmately infect a susceptible cell or undergo decay.

of each type - non-carrier and carrier. Upon induction and lysis, the cell releases the virion particles into the 'Virion grid' at the same grid coordinate as shown in Figure 2(B). The released virions are carriers if the induced prophage was a carrier. Irrespective of whether it is a carrier or not, at every time point, each virion particle can decay or migrate to a neighbouring grid point with fixed probabilities for each. The migration probabilty V_{diff} controls the rate of diffusion of these particles. When a susceptible cell is located at a specific coordinate on the 'Cell grid', all virion particles present at the same coordinate on the Virion grid compete to infect the cell. Each virion particle has a fixed probability of infecting the cell. Therefore, the probability that the cell becomes infected increases linearly with the local virion abundance.

Results

Before exploring the conditions under which the gene evolves to be phage-associated, we first identified conditions under

which the phage and/or the gene can persist in the population at all. As both the gene and the phage are subject to loss mutations, they will naturally disappear from the system in the absence of self-replication. They must make sufficient copies of themselves, at a rate higher than at which they are lost, in order to persist.

Ignoring the effects of the phage, the gene can persist if selection is sufficiently high:

$$\pi > \frac{\mu}{\phi}$$

Similarly, ignoring the effects of the gene, the phage can persist if the induction rate is sufficiently high:

$$\alpha > \frac{\lambda(\beta + \delta)}{\beta(r_V - 1) - \delta}$$

See Appendix D for the derivations.

When both the gene and the phage (or mobile element in general) are present in the population, studies have shown that one can rescue the other by compensating for weak selection with high transmission rate [9] or vice versa [21], shifting the thresholds we identified above. However, we verify that such rescue mechanisms are not successful in our model. This is because the gene is lost due to mutation on the chromosomal locus at the same rate as it is lost on the phage locus. Hence, the phage cannot rescue the gene by transmission alone. This observation is in agreement with the findings of Mc Ginty et al. [12]. Similarly, if the phage is lost due to mutation, it cannot be rescued by linking with a beneficial gene when the gene can instead be on a more stable chromosomal locus [7]. Upon numerically simulating the model for different values of the parameters π and α, we observed that the thresholds for gene and phage persistence do rigidly exist, roughly partitioning the $\pi - \alpha$ (selection-transmission) parameter space into four quadrants (see Figure 3(A)). Although there is a quadrant where both the gene and the phage persist, this does not necessarily imply that they are evolutionarily associated. For example, a population with a chromosomally encoded gene (U_c) could simply be chronically infected by a non-carrier phage (L_c).

Gene association with prophage does not evolve in a well-mixed population

To investigate eco-evolutionary outcomes where phages carry the beneficial gene as cargo from a chromosomal origin, we performed numerical simulations across the $\pi - \alpha$ parameter space. Since our model does not include gene translocation from the chromosome to the prophage, we simulate an invasion attempt of a mutant that encodes the gene on a prophage instead of its chromosome, by setting the initial state to $L = 0.500, L_c = 0.495, L_p = 0.005$, and the remaining genotype abundances to zero. In other words: all cells have a prophage, almost all carrier cells encode the gene chromosomally, but 1% of cells have the genes encoded on a prophage instead. Hence, the question becomes whether the rare mutant encoding the gene on a prophage invades the population. Within the persistence quadrant (see Figure 3(A)) — where both the phage and gene can coexist — we examined the association between the gene and the phage in terms of the percentage of prophages that carry the gene and the percentage of gene copies that occur on the prophage (shown with a 2D colour gradient).

We observed that these conditions are not sufficient for the gene-phage association, as the mutant does not establish in the population (Figure 3(A) remains fully dark purple). This result indicates that, assuming genes with chromosomal origins, it is difficult for the gene to be genetically linked to phages as cargo. Hence, in agreement with previous models [12, 13], prophage-encoded (more generally MGE-encoded) accessory genes that are typically traced back to a chromosomal origin cannot evolve in a well-mixed population.

Why is it so difficult for a beneficial gene to become stably associated with a prophage? When the gene is initially located on the bacterial chromosome, infection by phages carrying the gene does not confer an additional fitness advantage to already gene-bearing cells, and thus these phages are not selectively favoured. Although susceptible cells lacking the gene can gain a fitness benefit upon infection by carrier phages, the overall propagation of the gene is limited by the presence of a competing population of non-carrier phages which are always present due to gene loss mutations. Additionally, the gene can be lost from the prophage due to phage loss mutations. As a result, the net growth rate of the phage-associated gene cannot surpass that of the chromosomally encoded gene.

1. Spatial structure between populations: differential induction promotes gene-prophage association

In their review, Abedon and LeJeune have explored endogenous factors such as epistasis that promote the association of beneficial genes with prophages [6]. However, beyond within-cell biochemical interactions, the co-regulation of prophage induction and gene expression can also emerge from *exogenous* factors such as environmental heterogeneity. The gene may not be advantageous in all environments — for example, a pathogenicity trait is only beneficial within a host environment. Similarly, prophage induction is often influenced by extracellular conditions [22]. Henrot and Petit identified multiple signalling molecules within the mammalian gut microbiota that can trigger prophage induction [23]. Additionally, expression of benefits from the gene itself may alter the cellular environment in ways that promote induction. For example, Pattenden et al. propose that virulent bacterial strains, due to their faster growth and increased exposure to stress, induce their prophages more frequently [24]. Furthermore, immune responses targeting virulent cells may inadvertently trigger prophage activation.

Motivated by these observations, we next consider a scenario in which the gene is beneficial only in a specific environment, and this same environment promotes prophage induction. To model this, we introduce a second compartment representing the "outside world" where toxin production confers no fitness benefit and prophage induction signals are weak (see Appendix E for modified equations). In the absence of migration, the system in this outside world will converge towards a population of non-carrier susceptible (U) cells. We connect this compartment with the "host environment" by introducing migration between them. We assume that the migration is symmetric and that both compartments have equal maximum carrying capacities.

With this modification, we see a large portion of the parameter space permitting the invasion of phage-borne genes within the host environment (bright red colouring in Figure 3(B)). This is because the cells in the outside world are inclined to lose both the gene and the phage, and the influx of these non-carrier susceptible cells increases the rate of phage-infections that successfully convert non-carriers into carriers within the host environment, hence promoting selection for mobility at the gene-level. This is only true if this rate of conversion by phage transmission can exceed the rate of selection replacing incoming non-carriers with carriers. Hence, the bright red region appears only for weak selection and a high rate of prophage induction. It is interesting to note that gene-phage association is not a stable attractor when either of the compartments is considered to be a closed system. However, upon permitting migration between them, phage-association becomes the stable evolutionary attractor.

To highlight the requirement of weaker rate of prophage induc-

Figure 3: **ODE model:** (A) In an unstructured environment, i.e. one well-mixed compartment of cells, gene-phage association does not evolve. The degree of association of gene and phage (represented by the 2D colour map on the right) is plotted after simulation time 10^6 for different values of α and π. (B) When a new compartment simulating the "outside world", where the gene is not beneficial, is introduced, gene-phage association evolves but it requires that $\alpha > \alpha_{\text{out}}$.

tion in the outside world (α_{out}), we repeated the simulations for different values of α_{out} (shown in red dashed line). We saw

that the gene-phage association evolves only when the rate of prophage induction in the host environment exceeds α_{out}.

Figure 4: Differential dispersal of virions in a spatially structured cell population leads to the evolution of gene-phage association for intermediate selection. The plots show the distribution of final fraction of prophage-association in the gene population, excluding runs where the gene went extinct. For each value of π and V_{diff}, the distribution is plotted over 20 replicates of the stochastic simulation. (A) Cells are stationary on the grid. (B) Cells are well-mixed, i.e. the grid points on the 'Cell grid' are shuffled every time-step.

Therefore, the higher proportion of uninfected cells in incoming cells, due to the weaker transmission rate of the phage in the outside world, is a requirement for the emergent selection pressure on mobility of the gene within the host environment.

We demonstrate that when prophage induction and gene expression become environmentally co-regulated, stable gene-phage associations can evolve *de novo*. More generally, we postulate the following hypothesis from our metapopulation framework.

> *When there is a positive correlation between the benefits offered by a gene and the transmission rate of a mobile element across patches, the gene evolves linkage with the mobile element.*

The analysis so far has been performed using a set of 16 differential equations (8 equations for each compartment), which assumes that the population is homogeneous within each compartment. However, natural microbial populations are also spatially structured *within* these patches, exhibiting local variations in bacterial density and gene/phage distribution. In the next section, we explore whether spatial structure within a patch alone can drive gene-phage associations without requir-

ing additional regulatory mechanisms.

2. Spatial structure within a population: virion dispersal promotes gene-prophage association

In the individual-based spatial model (described in Methods), due to discrete population sizes, a gene can be lost entirely and not re-emerge. To investigate whether phage-encoded cargo genes can establish, we conducted an invasion experiment. We initialized a population of L_c cells on a 120×120 grid and allowed mutant types (U, U_c, and L) to arise and stabilize their abundances. We then introduced a spontaneous translocation mutation, converting 1% of L_c cells into L_p cells, effectively moving the gene from the chromosome to the prophage. If the phage-associated gene went extinct, we reintroduced the mutation in the same manner. The simulation ran until either the gene was lost from both loci or the maximum time limit (5×10^6 steps) was reached. A "successful invasion" was defined as the complete replacement of the chromosomal gene by the phage-associated gene.

Running 20 replicates of the experiment across different values of privatisation (π) and virion diffusion rates (V_{diff}) revealed two key findings (Figure 4). First, phage-associated genes successfully invaded in some replicates under intermediate pri-

Figure 5: **Snapshots of the grid model at different time-points of a single run of the simulation.** The three cases shown here give a side-by-side comparison to illustrate the limitations of lowering virion dispersal and increasing selection coefficient in the successful invasion of the phage-encoded gene.

vatisation. While the ODE model never allowed full takeover of the phage locus, spatial simulations frequently did. Control experiments with well-mixed populations confirmed that this effect was driven by spatial structure rather than finite population size (see Figure 4(B)). Second, increasing virion dispersal significantly enhanced phage-associated gene invasion. While a change from 0.1 to 0.001 in virion diffusion rate had no noticeable effect, well-mixed virions led to successful invasion in all replicates with intermediate privatisation. However, when both cells and virions were well-mixed, virion diffusion no longer influenced gene-phage association, highlighting the role of differential dispersal in moving the phage-associated gene away from local competition with the chromosomal gene.

To better understand the conditions under which the phage-encoded gene can invade, we analysed spatial snapshots and temporal dynamics across different parameter regimes (Figure 5). In spatially structured environments, local reproduction leads to clustering of genotypes, forming spatial niches where the success of an invading genotype depends on local context. Carrier genotypes have a higher growth rate and are able to invade patches occupied by non-carrier cells (visible in orange and yellow in panels a and c). However, gene loss mutations

continually regenerate non-carrier genotypes, allowing gene-free clusters to persist under weak selection.

While both chromosomal and phage-encoded versions of the gene can invade these clusters, the phage locus has a unique advantage: virion dispersal allows it to reach distant, gene-free regions that the chromosomal gene cannot access as easily. This effect is evident under weak selection and high virion diffusion (Figure 5(a)), where phage-mediated transmission facilitates successful invasion. However, when virion diffusion is limited (Figure 5 (c,d)), the advantage of the phage locus is lost and virions remain localized, unsuccessful in reaching susceptible cells in distant gene-free patches.

In contrast, strong selection (Figure 5(b)) eliminates gene-free regions through selective sweeps. Under these conditions, the chromosomal locus dominates due to its mutational robustness. Interestingly, under strong selection with high virion diffusion (Figure 5(b)), the system stabilizes into a persistent coexistence of the chromosomal and phage-encoded gene, as neither is able to fully outcompete the other.

In summary, the success of the phage-encoded gene depends

critically on two factors: effective virion dispersal and the continued presence of gene-free populations. These conditions allow the phage to exploit spatial heterogeneity and temporarily outcompete the chromosomal gene.

Based on these findings, we hypothesize the following.

> *Accessory genes in bacteria evolved to be associated with phages because of its ability to disperse these genes to favourable niches through virion particles.*

See Figure 6 for an illustration of the hypothesis.

The occurrence of these favourable niches can be attributed to the weak selection pressure on the gene in a spatially structured cell population. This mechanism sets apart phage-encoded genes from genes associated with more locally transmitted MGEs such as plasmids and could potentially explain why phages differ from other MGEs in terms of the composition of their cargo.

Discussion

This study presents a suite of theoretical models to investigate the evolution of phage-associated genes originating from chromosomal loci. We verify that chromosomal mutational robustness and competition with non-carrier phages create significant barriers to the evolution of phage-encoded accessory genes. However, we find that spatial structure – both between and within populations – can overcome these barriers. Previous studies have highlighted co-regulation between virulence and prophage genes as a mechanism for their association [6]. Extending this idea, we propose that environmental heterogeneity can drive phage-gene associations, while spatial structure within bacterial populations can promote association via virion-mediated gene dispersal, without any regulatory coupling. In other words, we identified exogenous factors that can explain the observed association between prophages and virulence genes in contrast to previously hypothesised endogenous mechanisms.

Our results suggest that when prophage induction is upregulated within host environments, genes that provide fitness advantages in these environments readily evolve to associate with these prophages. This process is driven by the influx of non-carrier susceptible cells from the external environment. In our analysis, we assumed symmetric cell densities and migration rates between the two environments, but in reality, these factors may vary. Exploring how different metapopulation structures influence this evolutionary process would be an interesting direction for future research. Nonetheless, our findings help explain the empirical tendency of phages to encode virulence factors [5, 6, 20, 25]. Environmental conditions at sites of bacterial infection, such as the gut [23] or urinary tract [26], have been shown to elevate prophage induction rates. In light of our findings, it would be valuable to experimentally test whether such prophages exhibit an increased propensity to link with virulence genes. More broadly, our model applies to any weakly selected beneficial trait. Virulence genes fit this criterion due to their known tendency to share the benefits of pathogenicity with surrounding bacteria [27]. The requirement for weak selection also aligns with a scenario where increasing host immunity decreases the selective pressure on virulence genes.

Phages offer distinct advantages over plasmids in associating with virulence genes due to their lifecycle. Unlike plasmids, which rely on contact-dependent conjugation, phages achieve horizontal transfer through virion dispersal, and induction allows phages to synchronise gene transfer with adaptive stress responses. While cross-species transfer has been proposed to facilitate plasmid-associated gene evolution [13], temperate phages may be more host-specific due to receptor-mediated infection and site-specific integration [17, 28]. While plasmids carrying antimicrobial resistance genes may have host ranges spanning different genera or beyond [14], broad host range temperate phages are typically limited to strains within the same species or genus [29–31]. Furthermore, evidence suggests that phage-associated virulence genes originate from chromosomal loci via erroneous excision of prophages [18]. While priority effect has been shown to drive plasmid-gene associations [13], our models suggest that localised priority effects in spatially structured populations may instead facilitate the evolution of phage-encoded genes.

Weak selection emerges as a key condition for MGE-borne gene evolution [9, 13, 32]. Strong selection, typically associated with essential genes, favours chromosomal association due to stability and reliability. With weak selection, frequent loss of the gene creates opportunities for lineages of non-carriers to coexist with carriers, making mobility advantageous as it enables the gene to spread independent from their host reproductive cycle. Due to weak selection, our simulations show that fixation of mobile genes occurs over long timescales, amidst significant demographic noise (see Appendix F).

In a spatially structured population, cooperation can persist through spatial privatisation – benefits being shared primarily within local neighbourhoods. When we restrict sharing of benefits from carrier cells to a radial neighbourhood, we found that phage-gene association evolved more easily with increasing virion dispersal as well as increasing the range of cooperation (see Appendix G). This is because increasing the radius of sharing benefits increases the chance of non-carrier cells receiving the benefit and lowers the selection coefficient acting on the gene. Therefore, we highlight the generalisability of our dispersal hypothesis to any weakly selected gene in a spatially structured bacterial population.

For illustrative purposes, our models rely on several simplifying assumptions, deviating from the realities of natural systems. One notable assumption is superinfection immunity. In natural bacterial populations, particularly among pathogens, multiple prophages often coexist within a single host cell [18, 28]. These prophages can undergo high order interactions and genetic exchange that reflect in dynamics at higher levels of biological systems [17, 33]. While embedding these interactions in ODE models inflates the number of equations, it will be interesting to relax superinfection immunity in individual-based models in the future. These models may also include other processes we have thus far neglected, such as retention of partial immunity after phage loss [34]. While these unmodelled complexities are potentially significant, they do not invalidate the core insights gained from our current model.

In summary, our findings highlight spatial structure as a critical mechanism for the evolution of phage-encoded accessory genes. By explicitly modelling phage-specific dynamics,

Figure 6: **Dispersal hypothesis illustrated for phage-encoded beneficial traits.** When a chromosomal gene can locally outcompete its phage-associated counterpart, but both types can locally outcompete non-carriers, the phage-associated gene gains a unique dispersal advantage. By diffusing through virion particles, it increases the mixing of carrier and non-carrier genotypes and thereby has a higher effective rate of selection over the chromosomal gene. This enables the phage-associated gene to outcompete the chromosomal gene on a larger spatial scale.

rather than MGEs in the abstract, we provide a framework to understand why phages are more likely to associate with virulence genes [5]. Our work provides mechanistic insights into the unique evolutionary role of temperate phages in bacterial adaptability, and their role in the persistence of virulence.

Authors' contributions: S.A.N. conceived the model with input from B.D. and designed and programmed the model, ran simulations, generated figures from the output data and drafted the initial manuscript. All authors gave final approval for publication and agree to be held accountable for the work performed therein.

Acknowledgements: We thank Paulien Hogeweg, Otto Cordero, Nobuto Takeuchi and Hilje Doekes for valuable discussions and feedback on earlier versions of this work.

Competing interests: We declare that we have no competing interests.

Software: Simulations of the analytical model were run using Jupyter notebook [35], and the packages Matplotlib [36], NumPy [37], SciPy [38], and Pandas [39]. The spatial model is implemented using Cacatoo [40], and an interactive version is available at `https://sultannazir.github.io/bacteria_prophage_codes/`.

Data availability: All code and data for the model are available online at `https://github.com/sultannazir/bacteria_prophage_codes`.

References

[1] A. Mark Osborn and D. Böltner, *Plasmid*, 2002, **48**, 202–212.

[2] J. P. J. Hall, E. Harrison and D. A. Baltrus, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 2021, **377**, Publisher: Royal Society, 20200460.

[3] D. J. Rankin, E. P. C. Rocha and S. P. Brown, *Heredity*, 2011, **106**, Publisher: Nature Publishing Group, 1–10.

[4] A. J. Weisberg and J. H. Chang, *Annual Review of Microbiology*, 2023, **77**, Publisher: Annual Reviews, 603–624.

[5] N. Takeuchi, S. Hamada-Zhu and H. Suzuki, *Proceedings of the Royal Society B: Biological Sciences*, 2023, **290**, Publisher: Royal Society, 20231088.

[6] S. T. Abedon and J. T. LeJeune, *Evolutionary Bioinformatics Online*, 2007, **1**, 97–110.

[7] C. T. Bergstrom, M. Lipsitch and B. R. Levin, *Genetics*, 2000, **155**, 1505–1519.

[8] J. Smith, *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 2001, **268**, Publisher: Royal Society, 61–69.

[9] B. van Dijk et al., *eLife*, 2020, **9**, ed. S. Mitri, P. J. Wittkopp and P. G. Higgs, Publisher: eLife Sciences Publications, Ltd, e56801.

[10] B. van Dijk, *Current genetics*, 2020, **66**, 1069–1071.

[11] F. Geoffroy and H. Uecker, *Theoretical Population Biology*, 2023, **154**, 102–117.

[12] S. E. Mc Ginty, D. J. Rankin and S. P. Brown, *Evolution*, 2011, **65**, 21–32.

[13] S. Lehtinen, J. S. Huisman and S. Bonhoeffer, *Evolution Letters*, 2021, **5**, 290–301.

[14] C. Coluzzi and E. P. Rocha, *Molecular Biology and Evolution*, 2025, msaf060.

[15] L.-M. Bobay, E. P. Rocha and M. Touchon, *Molecular Biology and Evolution*, 2013, **30**, 737–751.

[16] L.-M. Bobay and H. Ochman, *Proceedings of the National Academy of Sciences*, 2018, **115**, Publisher: Proceedings of the National Academy of Sciences, 6040–6045.

[17] J. A. Moura de Sousa et al., *Molecular Biology and Evolution*, 2021, **38**, 2497–2512.

[18] H. Brüssow, C. Canchaya and W.-D. Hardt, *Microbiology and Molecular Biology Reviews*, 2004, **68**, Publisher: American Society for Microbiology, 560–602.

[19] V. S. Gummalla et al., *Microorganisms*, 2023, **11**, Number: 3 Publisher: Multidisciplinary Digital Publishing Institute, 541.

[20] E. F. Boyd and H. Brüssow, *Trends in Microbiology*, 2002, **10**, 521–529.

[21] A. S. Millan et al., *Nature Communications*, 2014, **5**, Publisher: Nature Publishing Group, 5208.

[22] J. S. Huisman, A. Bernhard and C. Igler, *Trends in Microbiology*, 2025, **0**, Publisher: Elsevier, DOI: 10.1016/j.tim.2025.01.007.

[23] C. Henrot and M.-A. Petit, *Molecular Microbiology*, 2022, **118**, 494–502.

[24] T. Pattenden, C. Eagles and L. M. Wahl, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 2021, **377**, Publisher: Royal Society, 20200465.

[25] K. Schroven, A. Aertsen and R. Lavigne, *FEMS Microbiology Reviews*, 2021, **45**, fuaa041.

[26] T. Miller-Ensminger et al., *PeerJ*, 2020, **8**, Publisher: PeerJ Inc., e9718.

[27] T. Nogueira et al., *Current Biology*, 2009, **19**, Publisher: Elsevier, 1683–1691.

[28] G. López-Leal et al., *mSystems*, 2022, **7**, Publisher: American Society for Microbiology, e00326–22.

[29] S. Ford et al., *Frontiers in Microbiology*, 2021, **12**, Publisher: Frontiers, DOI: 10.3389/fmicb.2021.765271.

[30] J. A. Hammerl et al., *International Journal of Molecular Sciences*, 2022, **23**, Number: 12 Publisher: Multidisciplinary Digital Publishing Institute, 6779.

[31] S. Z. Ilyas et al., *Frontiers in Microbiology*, 2022, **12**, Publisher: Frontiers, DOI: 10.3389/fmicb.2021.768931.

[32] B. van Dijk and P. Hogeweg, *Genome biology and evolution*, 2016, **8**, 176–188.

[33] C. C. Wendling, *Current Opinion in Systems Biology*, 2023, **35**, 100469.

[34] C. F. Arias et al., *Nature Communications*, 2022, **13**, Publisher: Nature Publishing Group, 7412.

[35] T. Kluyver et al., Positioning and Power in Academic Publishing: Players, Agents and Agendas, ed. F. Loizides and B. Schmidt, 2016, pp. 87–90.

[36] J. D. Hunter, *Computing in Science & Engineering*, 2007, **9**, 90–95.

[37] C. R. Harris et al., *Nature*, 2020, **585**, 357–362.

[38] P. Virtanen et al., *Nature Methods*, 2020, **17**, 261–272.

[39] W. McKinney, Proceedings of the 9th Python in Science Conference, ed. S. van der Walt and J. Millman, 2010, pp. 56–61.

[40] B. van Dijk, *Journal of Open Source Software*, 2022, **7**, 3948.

A   Ordinary Differential Equations

The described model can be expressed in the form of differential equations. We use the notations used for the cell genotypes as non-negative real-valued variables to measure their corresponding abundances in the population. Therefore, the total cell population size (N) and the fractional abundance of carriers (f_τ) can be expressed as

$$N = U + U_c + L + L_c + L_p + L_{cp}$$
$$f_\tau = (U_c + L_c + L_p + L_{cp})/N$$

Similarly, we use V and V_τ to denote the abundances of non-carrier and carrier virion particles. Hence, we describe the temporal dynamics of the cell and phage abundances uding the following system of differential equation:

$$\frac{dU}{dt} = U\left(1 - d - N + (1-\pi)\phi f_\tau\right) \qquad - \beta U(V + V_\tau) \qquad + \lambda(L + L_p) + \mu U_c$$

$$\frac{dU_c}{dt} = U_c\left(1 - d - N + \pi\phi + (1-\pi)\phi f_\tau\right) \qquad - \beta U_c(V + V_\tau) \qquad + \lambda(L_c + L_{cp}) - \mu U_c$$

$$\frac{dL}{dt} = L\left(1 - d - N + (1-\pi)\phi f_\tau\right) \qquad + \beta UV \qquad - \lambda L + \mu(L_c + L_p) \qquad - \alpha L$$

$$\frac{dL_c}{dt} = L_c\left(1 - d - N + \pi\phi + (1-\pi)\phi f_\tau\right) \qquad + \beta U_c V \qquad - (\lambda + \mu)L_c + \mu L_{cp} \qquad - \alpha L_c$$

$$\frac{dL_p}{dt} = L_p\left(1 - d - N + \pi\phi + (1-\pi)\phi f_\tau\right) \qquad + \beta U V_\tau \qquad - (\lambda + \mu)L_p + \mu L_{cp} \qquad - \alpha L_p$$

$$\frac{dL_{cp}}{dt} = \underbrace{L_{cp}\left(1 - d - N + \pi\phi + (1-\pi)\phi f_\tau\right)}_{\text{Growth}} \qquad + \underbrace{\beta U_c V_\tau}_{\text{Infection}} \qquad - \underbrace{(\lambda + 2\mu)L_{cp}}_{\text{Mutations}} \qquad - \underbrace{\alpha L_{cp}}_{\text{Induction}}$$

$$\frac{dV}{dt} = r_V \alpha(L + L_c) \qquad - \beta(U + U_c)V \qquad - \delta V$$

$$\frac{dV_\tau}{dt} = \underbrace{r_V \alpha(L_p + L_{cp})}_{\text{Lytic release}} \qquad - \underbrace{\beta(U + U_c)V_\tau}_{\text{Infection}} \qquad - \underbrace{\delta V_\tau}_{\text{Decay}}$$

B Model parameters

Variable	Meaning	Value (ODE/IBM)
U	Abundance of non-carrier susceptible cells	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
U_c	Abundance of suceptible cells carrying gene on the chromosomal locus	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
L	Abundance of non-carrier lysogens	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
L_c	Abundance of lysogens carrying the gene on the chromosomal locus	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
L_p	Abundance of lysogens carrying the gene on the phae locus	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
L_{cp}	Abundance of lysogens carrying the gene on both loci	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
V	Abundance of extracellular non-carrier virions	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$
V_τ	Abundance of extracellular carrier virions	$[0, \infty) \; / \; \{0, 1, 2, 3...\}$

Parameter	Meaning	Value (ODE/IBM)
ϕ	Ratio of maximum increase in carrying capacity to minimum carrying capacity	1 / 1
b	Intrisic birth rate	1 / 0.5
d	Intrinsic death rate	$0.02 \; / \; 5 \times 10^{-3}$
π	Fraction of benefits privatised by carriers of gene	$[10^{-5}, 1] \; / \; [10^{-5}, 1]$
β	Rate of phage infection	$0.02 \; / \; 5 \times 10^{-3}$
α	Rate of prophage induction and lysis	$[10^{-5}, 10^{-1}] \; / \; 5 \times 10^{-4}$
r_V	Burst size	30 / 30
δ	Rate of virion decay	$0.02 \; / \; 5 \times 10^{-3}$
μ	Rate of gene loss	$2 \times 10^{-3} \; / \; 5 \times 10^{-4}$
λ	Rate of phage loss	$2 \times 10^{-3} \; / \; 5 \times 10^{-4}$
m	Rate of migration	0 (one compt.), 0.01 (two compt.) / n.a. for IBM
V_{diff}	Rate of virion diffusion	$\{10^{-3}, 10^{-1}, \infty\}$ (IBM)

Table 1: Variables and parameters of the model, their meaning and the values used. The variables are non-dimensionalised in the ODEs and the rate parameters are scaled when switching between the ODE and IBM. The details of these scales are given in Appendix C

C   Well-mixed approximation of the spatial model

Let x be the number of cells on the grid of a certain genotype and k be the minimum carrying capacity, i.e. the grid size is $k(1 + \phi)$. Then, the probability of finding an empty grid point is

$$\mathbb{P}(\text{empty grid point}) = 1 - \frac{n}{k(1 + \phi)}$$

The average death rate of a genotype will be

$$\frac{\sum_{\text{genotype}} \mathbb{P}_{\text{death}}(\pi)}{\sum_{\text{genotype}} 1} = \begin{cases} d_s + \Delta d\Big(1 - \pi - (1 - \pi)f_\tau\Big), & \text{genotype is a carrier} \\ d_s + \Delta d\Big(1 - (1 - \pi)f_\tau\Big), & \text{genotype is not a carrier} \end{cases}$$

The the well-mixed growth dynamics of a carrier genotype is given by

$$\frac{dx}{dt'} = \text{Avg. } \#\text{births into empty grid points} - \text{Avg. } \# \text{ deaths}$$

$$\frac{dx}{dt'} = bx\Big(1 - \frac{n}{k(1 + \phi)}\Big) - x\Big(d + \Delta d\Big(1 - \pi - (1 - \pi)f_\tau\Big)\Big)$$

$$\frac{dx}{dt'} = \frac{b}{1 + \phi}x\Big(1 + \phi - \frac{n}{k} - d_s\frac{1 + \phi}{b} - \Delta d\frac{1 + \phi}{b}\Big(1 - \pi - (1 - \pi)f_\tau\Big)\Big)$$

We apply the following non-dimensionalisation

$$X = \frac{x}{k}, \quad N = \frac{n}{k}, \quad t = t'\frac{b}{1 + \phi}$$

and the following substitutions

$$d_s = d\frac{b}{1 + \phi}, \quad \Delta d = \frac{b\phi}{1 + \phi}$$

to obtain

$$\frac{dX}{dt} = X(1 - N + \pi\phi + (1 - \pi)\phi f_\tau)$$

which is the same growth dynamics we implemented in the ODEs. The same derivation can be done for the growth dynamics of a non-carrier genotype. Notice that time is scaled by a factor of $\frac{b}{1+\phi}$ in the ODE, i.e. one time-step in the IBM is equal to $\frac{1+\phi}{b}$ (which is equal to 4 for our choice of parameters) many time-steps of the ODE. Therefore, all the rate parameters, just like the intrinsic death rate, need to be adjusted to the time scale by this factor when switching between the models.

D Selection- and transmission-mutation thresholds

To derive the persistence condition of the gene in the face of mutation, we can consider a simpler version of the model consisting only of two kinds of cells: carrier (U) and non-carrier (U_c).

$$\frac{dU}{dt} = U\left(1 - U - U_c + (1-\pi)\phi\frac{U_c}{U+U_c}\right) + \mu U_c$$

$$\frac{dU_c}{dt} = U_c\left(1 - U - U_c + \pi\phi + (1-\pi)\phi\frac{U_c}{U+U_c}\right) - \mu U_c$$

We obtain the condition that for a positive abundance of the carrier type at steady-state ($U_c(t=\infty) > 0$) requires the rate of selection to exceed the rate of mutation:

$$\pi\phi > \mu$$

Proof. Consider the non-dimensionalised version of the model such that $u = U/N$ and $u_c = U_c/N$, where $N = U + U_c$. The equations then become

$$\frac{du}{dt} = -\pi\phi u u_c + \mu u_c$$

$$\frac{du_c}{dt} = \pi\phi u u_c - \mu u_c$$

Solving for steady state, the non-trivial solutions are $u^\infty = \frac{\mu}{\pi\phi}, u_c^\infty = 1 - \frac{\mu}{\pi\phi}$, and $u^\infty = 1, u_c^\infty = 0$.

We can infer that a stable positive solution for u_c^∞ exists only when $\pi\phi > \mu$. $\qquad\square$

Now to derive the persistence condition of the MGE, let us look at a simpler version of the model consisting only of two kinds of bacteria: uninfected (U) and infected (L). First, let us consider a plasmid.

$$\frac{dU}{dt} = U(1 - U - L) - \beta U L + \lambda L$$

$$\frac{dL}{dt} = L(1 - c - U - L) + \beta U L - \lambda L$$

We obtain the condition that the plasmid will persist in the population ($L(t=\infty) > 0$) if the conjugation rate is higher than the sum of rate of segregational loss and cost of carrying the plasmid:

$$\beta > \lambda + c$$

Proof. The plasmid will persist if it has a positive growth rate when it is rare in the population, i.e. the susceptible cells are maximally abundant ($U = 1$) and the infected cell abundance is close to zero ($L^2 \simeq 0$). Then,

$$\frac{dL}{dt} > 0 \implies \beta - \lambda - c > 0$$

$\qquad\square$

When the MGE is a phage, we will consider an eqaution describing the virion abundance (V) as well.

$$\frac{dU}{dt} = U(1 - U - L) - \beta U V + \lambda L$$

$$\frac{dL}{dt} = L(1 - U - L) + \beta U V - \lambda L - \alpha L$$

$$\frac{dV}{dt} = r_V \alpha L - \beta U V - \delta V$$

We obtain the condition that the phage will persist in the population ($L(t=\infty) > 0$) if the lysis rate is above a certain threhsold:

$$\alpha > \frac{\lambda(\beta + \delta)}{\beta(r_V - 1) - \delta}$$

Proof. By explicitly considering the extracellular virions, we have modelled an infection cycle with two stages: the prophage stage and the virion stage. For a two-stage infection cycle, the net basic reproductive number of the infection, R_0 is given by the product of R_0 of each step. Therefore, the phage will presist if the net R_0 is greater than 1.

$$R_0 = R_0(\text{prophage}) \times R_0(\text{virion})$$

$$= \frac{\text{max. infection rate}}{\text{phage loss rate} + \text{lysis rate}} \times \frac{\text{burst size} \times \text{lysis rate}}{\text{max. infection rate} + \text{decay rate}}$$

$$= \frac{r_V \alpha \beta}{(\lambda + \alpha)(\beta + \delta)}$$

Therefore,

$$R_0 > 1 \implies \alpha > \frac{\lambda(\beta + \delta)}{\beta(r_V - 1) - \delta}$$

□

E   Modified ODEs

Let m be the rate of migration between the 'host' and the 'outside world' compartments. Then the ODEs describing the dynamics in the outside world (variables denoted with a superscript o represent abundances in the outside world).

$$\frac{dU^o}{dt} = U^o(1 + \phi - d - N^o) - \beta U^o(V^o + V^o_\tau) + \lambda(L^o + L^o_p) + \mu U^o_c + m(U - U^o)$$

$$\frac{dU^o_c}{dt} = U^o_c(1 + \phi - d - N^o) - \beta U^o_c(V^o + V^o_\tau) + \lambda(L^o_c + L^o_{cp}) - \mu U^o_c + m(U_c - U^o_c)$$

$$\frac{dL^o}{dt} = L^o(1 + \phi - d - N^o) + \beta U^o V^o - (\alpha_{\text{out}} + \lambda)L^o + \mu(L^o_c + L^o_p) + m(L - L^o)$$

$$\frac{dL^o_c}{dt} = L^o_c(1 + \phi - d - N^o) + \beta U^o_c V^o - (\alpha_{\text{out}} + \lambda + \mu)L^o_c + \mu L^o_{cp} + m(L_c - L^o_c)$$

$$\frac{dL^o_p}{dt} = L^o_p(1 + \phi - d - N^o) + \beta U^o V^o_\tau - (\alpha_{\text{out}} + \lambda + \mu)L^o_p + \mu L^o_{cp} + m(L_p - L^o_p)$$

$$\frac{dL^o_{cp}}{dt} = L^o_{cp}(1 + \phi - d - N^o) + \beta U^o_c V^o_\tau - (\alpha_{\text{out}} + \lambda + 2\mu)L^o_p + m(L_{cp} - L^o_{cp})$$

$$\frac{dV^o}{dt} = r_V \alpha_{\text{out}}(L^o + L^o_c) - \beta(U^o + U^o_c)V^o - \delta V^o$$

$$\frac{dV^o_\tau}{dt} = r_V \alpha_{\text{out}}(L^o_p + L^o_{cp}) - \beta(U^o + U^o_c)V^o_\tau - \delta V^o_\tau$$

where $N^o = U^o + U^o_c + L^o + L^o_c + L^o_p + L^o_{cp}$. The only modification to the system of ODEs for the host comparmtent is the addition of a migration term $+m(X^o - X)$ to the ODE for the corresponding X genotype abundance variable.

F Timeseries evolution prophage-association in space

(a) Cells are stationary on the grid

(b) Cells are well-mixed on the grid

Supplementary Figure. 1: Timeseries plots of the fraction of phage-associated genes for 20 replicates per value of privatisation (rows, $p(\pi) = -\log_{10}\pi$) and virion diffusion rate (columns). The red triangles mark points in the simulation runs where the gene went extinct.

G   Spatial privatization of benefits

Supplementary Figure. 2: (Left) In a spatial model, privatization can be modelled mechanistically through spatial privatization, i.e. the benefits of expression of a cooperative trait is only experienced by cells in an immediate neighbourhood. Therefore, we introduce a parameter "Range of cooperation" which measures the maximum distance of a cell from a carrier cell in order to obtain benefits from the latter. Therefore, decreasing this radius increases the privatization. Furthermore, since a cell is likely surrounded by similar genotype, decreasing the radius also increases the selection pressure acting on the gene. (Right) With the mechanistic implementation of privatization, we yet again observe that the chance of prophage-association increases with increasing virion dispersal and decreasing selection coefficient (increasing range of cooperation). In comparison with the case where the cells are well-mixed, we can clearly see that spatial structure is required for this trend. We also see that gene extinction happens sooner with spatial structure and that the effect is stochasticity is much stronger. This is because at the local level, the carrier genotypes have the same growth rate as the non-carrier (cheater) genotypes and hence evolves towards gene extinction. Selection in this model occurs at a higher (group-) level. This sufficiently reduces the effective population size on which selection is acting - leading to increased genetic drift. To increase the effective population size, we lowered the parameter ϕ to 0.1 and adjusted other parameter values accordingly.