PHIROS: Integrated microfluidic platform for multi-day high-resolution imaging of organotypic slices

Jana B. Petr[1], Meng-Syuan Lin[2], Catarina M. Gomes[1], Dina Hochuli[2], Andreas Hierlemann[1], Martin Baumgartner[2*], Mario M. Modena[1*]

[1]ETH Zurich, Department of Biosystems Science and Engineering, Basel, Switzerland
[2]University Children's Hospital, Children's Research Centre, Division of Oncology, Zurich, Switzerland
*Corresponding authors: martin.baumgartner@kispi.uzh.ch, mario.modena@bsse.ethz.ch

Abstract

Replicating the cytoarchitecture and cell heterogeneity of the brain *in vitro* remains challenging. Although *ex vivo* organotypic slices preserve native tissue complexity, current culturing methods limit long-term, high-resolution imaging and restrict temporally controlled perturbations. Here, we present PHIROS, a microfluidic platform for high-resolution imaging of organotypic slices that supports static culturing of excised tissue at the air–liquid interface and provides continuous optical access for extended imaging at subcellular resolution. Controlled perfusion with oxygenated medium preserves tissue viability over several days, enabling, e.g., the monitoring of spontaneous and pharmacologically modulated astrocytic calcium activity. Using PHIROS we characterized medulloblastoma (MB) cell behavior in a physiological tumor microenvironment and observed dynamic F-actin-driven interactions with tissue-resident astrocytes, as well as leading edge localization of the immune-checkpoint marker B7-H3 in invading tumor cells. Quantitative assessment of mitochondria transfer across heterotypic actin-rich connections evidences the potential of PHIROS as a versatile system for mechanistic studies in a tissue context, enabling controlled compound exposure and high-resolution imaging in physiologically relevant tissue settings.

Main

Replicating the complexity of intact tissue *in vitro* remains a major challenge in biology. Intact tissues integrate diverse cell types within a highly organized cytoarchitecture, enabling complex cell signaling and physiological functions. Conventional *in vitro* systems, such as 2D co-cultures, spheroids and organoids, provide valuable insights into cell-to-cell interactions and cellular mechanisms, but fall short of reproducing complex tissue environments, ultimately resulting in limited physiological relevance[1,2]. Consequently, multicellular processes, such as neural network activity, tumor-host interactions, and drug responses, cannot be investigated under conditions that reflect native tissue organization by using these models.

Organotypic tissue slices offer a physiologically relevant alternative to overcome these limitations and preserve cytoarchitecture, cell heterogeneity, cytokine signaling and functional connectivity *ex vivo*[3,4]. These features enable studies of cell motility and signaling within a native tissue context, e.g., in the brain, where neuronal and glial interactions have shown to remain intact[3,5-7]. Organotypic slice models are commonly used in neuroscience and have recently increasingly been applied to cancer neuroscience research[8-12]. The preservation of a neural tumor microenvironment (TME) enables to investigate the interplay between tumor and cells of the TME, where connectivity and heterogeneity strongly influence tumor initiation[13], progression[14-17] and metastasis[18]. By engrafting tumor cells on healthy organotypic slices, tumor-host tissue interactions can be studied in a controlled and accessible environment over multiple days[9,19]. While these models can overcome many limitations of conventional *in vitro* systems, the currently used culturing methods introduce restrictions on imaging and experimental flexibility.

Current slice culturing techniques rely on air-liquid interfaces (ALIs)[5] or alternating exposure
to medium and air[20], which limit continuous optical access to the slice and confine imaging to
short time windows or endpoint analysis. Orthotopic tumor models offer restricted imaging and
typically require adult animals, large numbers of implanted cells, and long preparation times,
constraints that hinder approaches of personalized medicine and raise ethical concerns[10,11]. In
contrast, slice-based engraftment requires fewer cells, leverages tissue from young mice that
better mimic pediatric tumor biology, and enables short experimental timelines. These
advantages make organotypic slices particularly suited for modelling brain tumors, including
pediatric brain tumors. such as medulloblastoma (MB), the most common malignant brain
tumor in children[21].

To address slice-culturing limitations and unlock the full potential of slice-based disease
modeling, we developed a microfluidics-based **P**latform for **Hi**gh-**R**esolution imaging of
Organotypic **S**lices (PHIROS), which features long-term tissue viability, along with continuous,
unhindered optical access, and enables precise temporal control of compound delivery.
PHIROS enables real-time monitoring of subcellular events, including calcium dynamics,
mitochondrial transfer, and actin remodeling over multiple days in a tissue context at
unprecedented resolution. To demonstrate potential uses of PHIROS in neuroscience and
cancer research, we showcase here two different applications: First, we monitored the calcium
(Ca^{2+}) dynamics of astrocytic populations in organotypic cerebellar slices over 3 days, and we
perturbed network activity by blocking gap-junction intercellular communication. Secondly, we
implanted different fluorescently labelled medulloblastoma cells into cerebellar slices and
monitored tissue invasion, tumor-tumor and tumor-TME interactions, and the dynamic
localization of B7-homologoues 3 (B7-H3), a clinically relevant checkpoint inhibitor, during
tissue invasion.

Results

Platform for long-term high-resolution imaging of organotypic brain slices

PHIROS integrates four key functions: (i) static culturing at the air-liquid interface (ALI) for
tissue recovery and reorganization after slicing, (ii) perfused culturing in a closed chamber for
continuous, long-term high-resolution imaging under aseptic conditions, (iii) controlled change
of medium conditions for, e.g., drug delivery, and (iv) tissue fixation and recovery for
downstream analysis. Once placed on the platform, the brain slices remain undisturbed during
culturing and imaging, which prevents mechanical damage through transfers.

The platform comprises four main components (**Supplementary Figure 1A-B**): i) a
polymethylmethacrylate (PMMA)-based slice holder, featuring a microfluidic channel for the
perfusion culturing; ii) a hydrophilic 0.4-μm-pore-diameter polytetrafluoroethylene (PTFE)
membrane bonded to the slice holder; iii) a PMMA-based support part; iv) an adhesive foil for
sealing the assembly and enabling perfusion. The assembly of i) and ii) is further referred to
as the "culture chip". For static culturing, the culture chip is placed on a standard tissue-culture
insert, so that medium exchange can be performed following standard culturing procedures by
simply transferring the inserts into a well with fresh medium. The chip can then be removed
from the insert and sealed before transfer to a microscope stage at the beginning of the
perfusion protocol to enable long-term, high-resolution imaging (**Figure 1A-B**).

We cultured murine cerebellar slices for 14-18 days at the ALI to enable tissue recovery and
Purkinje cell-layer reorganization. ALI incubation enabled experimental interventions, such as
adeno-associated-virus (AAV)-based labeling of specific cell subpopulations and
medulloblastoma spheroid engrafting onto the cerebellar slices. Imaging was initiated after
sealing the slice compartment under sterile conditions (**Supplementary Figure 1B**). Up to 3
chips, accommodating a total of up to 9 cerebellar slices, were perfused and imaged in parallel

on a spinning-disk confocal microscope (**Supplementary Figure 1C**). To support brain-slice
viability, the medium was oxygenated using a custom-made gas exchanger. We found a
dissolved oxygen concentration of ~30 mg/L being sufficient to support tissue health without
bubble formation in the tubing. Oxygenation was achieved by using the custom-made gas
exchanger between medium reservoir and chip (**Figure 1C, Supplementary Figure 2**). By
combining static and perfusion culturing, PHIROS enables continuous, multi-day monitoring of
tissue and single-cell dynamics, including subcellular events, such as actin remodeling and
mitochondrial transfer (**Figure 1D**).

PHIROS preserves cerebellar slice cytoarchitecture and sustains tissue viability

We characterized cerebellar-slice cultures on chip and compared tissue organization with
those of standard ALI cultures and *in vivo* metrics, and we assessed tissue health under
perfusion. Our main questions were: i) does chip geometry and dual-membrane ALI culture
support tissue reorganization, and ii) is viability under perfusion with oxygenated medium
maintained.

We cultured P10-P12 murine cerebellar slices either on standard tissue-culture inserts or on
the chip placed on an insert and assessed Purkinje cell layer reorganization[8] after 16 days *in*
vitro (DIV 16, **Figure 2A,** dashed line at DIV 16). Under both conditions, slices appeared
translucent, indicative of good tissue health and maturation[3] (**Supplementary Figure 3**).
Immunofluorescence staining revealed a continuous Purkinje cell layer along the tissue
convolutions, a dense granule cell population, and astrocyte networks in the slices (**Figure 2B,**
Extended Data 1). The Purkinje cell layer was flanked by organized granular and molecular
layers, indicating a physiologically relevant reorganization of the tissue slice during the static
culturing.

We next quantified Purkinje cell-soma size and cell density and compared these values to
those of standard ALI cultures and published *in vivo* data. Early culture chip prototypes with
suboptimal culture conditions yielded sparse, disorganized, and disrupted Purkinje cell layers
with enlarged cell somas (**Supplementary Figure 4**). In contrast, the Calbindin staining of the
final chip design revealed a mean Purkinje cell-soma area of 2028 μm^3 and a cell density of
0.0014 cells/μm^2, matching data of standard ALI cultures and previously reported values for
Purkinje cell density *in vivo*[22] (**Figure 2C-D**). These results confirmed that chip geometry and
dual-membrane culture supported the reorganization of the cerebellar architecture in
organotypic slices without compromising viability.

To perform high-resolution imaging, the tissue slices need to be submerged in medium.
However, submerged slices develop hypoxia in the absence of perfusion, and imaging is
restricted to short time periods of a few hours only. To overcome this limitation, we implemented
a pulsatile-perfusion protocol with oxygenated medium, designed to minimize disruption of
endogenous gradients in the culture chamber while ensuring sufficient oxygen supply. We
established a flow protocol consisting of hourly cycles of partial medium exchange with
intermittent 1 h no-flow intervals. This strategy resulted in a daily medium consumption of ~1.4
mL, a volume comparable to that used under standard culturing conditions (1 mL) (**Figure 2A**).

To assess cell viability under perfusion, we used a cleaved caspase-3/7 live-cell stain, which
visualizes apoptotic cells in real time. Cell viability was assessed at the start of the perfusion
(day 1), after 24 (day 2) and 48 h (day 3). As a positive control, slices were treated with 40 µM
doxorubicin after 48 h of perfusion to induce apoptosis (**Figure 2E, Extended Data 2A**). The
low number of caspase-3/7-positive cells prior to doxorubicin exposure indicated good tissue
health in the sealed chip throughout the 48 h of perfusion (**Figure 2F**). No significant increase
in apoptotic cells was observed during the perfusion period, which further confirmed that
medium oxygenation and pulsatile perfusion supported cell viability.

To evaluate hypoxic stress during perfusion culturing, we determined the expression levels of
the hypoxia transcription factor HIF1-α in the tissue slices by immunofluorescence analysis.
We compared the cerebellar slices cultured using the pulsatile flow protocol with slices either
cultured at the ALI or submerged for 24 or 48 hours under 2 ml of medium (~2 mm of medium
column above the slices), to mimic the conditions of continuous, long-term imaging using
immersion objectives. Nuclear localization of HIF1-α, indicative of hypoxia, was quantified and
normalized against control ALI conditions. Submerged slices under static conditions exhibited
a 16.8- and 18.8-fold increase in hypoxia levels after 24 and 48 hours, respectively. In contrast,
hypoxia levels under pulsatile perfusion with oxygenated medium did not show any significant
difference with respect to ALI control condition (**Figure 2G, Extended Data 2B**). These findings
indicate that slice submersion, which would be required for imaging using high numerical-
aperture objectives, would rapidly compromise slice conditions, while pulsatile flow with
oxygenated medium in a closed environment preserves slice viability, thereby enabling long-
term imaging investigations.

Multi-day imaging of subcellular and network-wide astrocytic calcium events

To demonstrate PHIROS' ability to perform functional imaging in an intact tissue context, we
monitored spontaneous astrocytic Ca^{2+} dynamics in cerebellar slices. Astrocytes form a
network that is highly interconnected through gap junctions, enabling the transfer of ions and
small molecules between cells. In response to neuronal excitation, astrocytes exhibit complex
intracellular Ca^{2+} fluctuation and global Ca^{2+} dynamics. These calcium changes can be
visualized using genetically encoded calcium indicators[23,24] and detected using fluorescence
microscopy.

We transduced cerebellar slices from P8-P10 mice with AAV-hGFAP-GCaMP6f and
maintained the cultures at the ALI for 10 days (**Figure 3A**). During calcium imaging, we
transiently replaced the culture medium with artificial cerebrospinal fluid (aCSF) using the flow
system. To support tissue viability, standard culture-medium conditions were reinstalled
between imaging sessions. PHIROS enabled repeated imaging of the same region of interest
(ROI) over three consecutive days, allowing long-term monitoring of astrocytic activity in a
physiologically relevant tissue context (**Figure 3B-D**, **Supplementary Video 1**).

We detected spontaneous intracellular Ca^{2+} signal fluctuations with consistent intensity
changes (~0.2 ΔF/F) in astrocytic processes across the three days (**Figure 3E**). Ca^{2+} event
characteristics were analyzed using AQuA2[25], including event duration, area, and intensity,
and remained stable throughout the 48 hours of perfusion culture (**Figure 3F**), confirming the
platform's ability to support tissue functionality and facilitate functional imaging over extended
periods.

Next, we investigated the difference in Ca^{2+} events between Bergmann glia (**Figure 3D**) and
velate astrocytes (**Figure 3G**), two cerebellar astrocyte subtypes that display distinct
morphological and molecular signatures and local connectivity[26]. Bergmann glia cells extend
radial processes across the Purkinje cell layer to the pial surface, while velate astrocytes are
distributed within the granule cell layer[27]. We observed Ca^{2+} fluctuations in the cell body and
in processes in both astrocyte subtypes. Notably, we observed Ca^{2+} waves spreading within
local astrocyte networks predominantly in regions rich in velate astrocytes (**Figure 3G,**
Supplementary Video 2). While further investigating the Ca^{2+} dynamics of these two astrocyte
subtypes, we found that Ca^{2+} events in Bergmann glia were smaller in size, showed a smaller
amplitude and were shorter than observed events in velate astrocytes. Furthermore, the
number of events detected simultaneously was higher in velate astrocyte regions, indicating
higher cellular connectivity (**Figure 3H**). We next assessed whether velate network activity
could be modulated acutely using the gap-junction inhibitor carbenoxolone (CBX). Following

exposure, spontaneous Ca^{2+} fluctuations persisted, but event areas became smaller, likely
including single cells or small clusters of neighboring cells (**Figure 3I**). Quantification revealed
a decrease in the size and duration of events covering an area larger than 150 μm^2, as well as
a reduction in simultaneously occurring events, indicating carbenoxolone-induced disruption
of gap-junction-mediated network communication. Together, our findings underline PHIROS'
ability to support multiscale Ca^{2+} event visualization at sub-cellular resolution and over multiple
days, while enabling precise control over medium conditions at high temporal resolution.

Actin cytoskeleton dynamics and tissue-invasion behavior of medulloblastoma cells

Tumor cell invasion is strongly influenced by tissue stiffness and composition, features that
differ between *in vivo* and classical *in vitro* migration models. Tissue composition forces cells
to dynamically adapt motility modes during microenvironmental exploration[28]. We used
PHIROS to capture time-lapse recordings of MB spheroids, co-cultured with cerebellar slices,
and evaluated TME-dependent changes in MB cell morphology and motility by comparing them
to tumours grown on dissociated neural cells. During the static culturing period of the cerebellar
tissue slices, we transduced the slices with hGFAP-driven mCherry, enabling the visualization
of astrocyte somata and processes. To simultaneously monitor F-actin dynamics, we used
ONS-76 cells expressing LifeAct-eGFP. Tumor spheroids were placed on slices 72 h before
imaging (DIV 16–19) to allow engraftment, and sealed chips were imaged for 48 h under
periodic perfusion (**Figure 4A**).

We found that the tumor cells stopped exploring the cerebellar tissue microenvironment under
continuous perfusion, indicating that the flow might interfere with microenvironmental factor
gradients. This observation highlights the immediate impact of perturbation and underscores
the importance of the microenvironment for invasion. To minimize the disruption of endogenous
gradients, we limited the flow to only 7 minutes every hour, while image acquisitions were
performed outside the flow intervals. Under these flow conditions, we could observe different
patterns of motile cell behavior, ranging from random motility in the spheroid center (**Figure
4B i**) to directional streams of collectively invading cells (**Figure 4B ii**). The invading cells
exhibited directional movement trajectories only in certain regions of the tumor cell mass on
the tissue slice. In contrast, ONS-76 cells migrated radially on dissociated cultures of neural
cells without constraining tissue barriers (**Supplementary Figure 5**). Particle image
velocimetry (PIV) showed spatial heterogeneity in tumor cell dynamics when comparing the
2D with the 3D TME (**Figure 4C, Supplementary Figure 6**). The velocity magnitude analysis
revealed uniform radial movements in the neural cell-tumor co-culture, whereas corresponding
directional movements in the tissue slices were spatially restricted (**Figure 4C**). Speed analysis
revealed higher speeds of cells seeded on neural co-culture compared to speeds on slice co-
cultures, the latter of which were in good accordance with *in vivo* values[29,30], suggesting that
the microenvironment architecture served as an important factor controlling tumor motility
(Figure 4D).

In the slice co-culture models, the largest velocities were recorded in cluster *iii*, where cells
displayed actin-rich structures that were associated with altered local velocity patterns of
mCherry-positive astrocytes (**Supplementary Figure 6**), suggesting a potential physical
interaction of these two cell populations. Cluster *i* cells display marked F-actin dynamics with
little discernible directional movement of the mostly rounded cells (**Figure 4E**). In contrast,
cluster *ii* cells displayed flat and elongated morphologies with prominent lamellipodia and
filopodia, reminiscent of mesenchymal cells[31,32] at the invasion front (**Figure 4E,
Supplementary Video 4**). These cells exhibited F-actin-rich lamellipodia that displayed
explorative contacts with surrounding cells via filopodia-like protrusions (**Figure 4F**).
Importantly, while tumor cell-cell contacts in clusters *i* were mostly short lived (0.5 – 6 h), the
tumor cell-cell contacts in cluster *ii* persisted for more than 12 h and were an indication of
collective invasive behavior. We also observed fewer contacts between rapidly invading tumor

cells of this cluster with mCherry-positive astrocytes, suggesting an avoidance mechanism that
allows tumor cells to navigate through glial structures during invasion. (**Extended Data 3**).

To assess whether acute treatment effects could be monitored using PHIROS, we monitored
the effects of actin-polymerization inhibition by exposing both the slice and neural co-cultures
to cytochalasin D (CytoD) for 1 hour. In both models, subcellular actin distribution changed
after exposure, with a marked loss of F-actin and increased punctate actin distribution (**Figure**
4G). Furthermore, in neural co-cultures, CytoD induced an acute inhibition of cell dynamics,
which was not displayed in slice settings. Conversely, CytoD-treated tumor cells on cerebellar
slices elongated markedly within 24 h of exposure, possibly resulting from tissue-associated
responses to CytoD (**Figure 4H**).

B7-H3 localized to leading edge of explorative and invasive tumor cells

We next investigated whether PHIROS could be utilized to study the behavior of tumor-
associated surface molecules within a tissue context. To this end, we engineered ONS-76 cells
to express the transmembrane protein B7-H3/CD276, fused with mNeon-Green (ONS-76-B7-
H3-mNG, **Figure 5A**). B7-H3 is highly expressed in malignant pediatric brain tumors, including
MB, and is a validated target for CAR-T cell therapies[33,34]. In tissue-invading tumor cells, B7-
H3-mNG was prominently localized at the leading edge and along the plasma membrane
(**Figure 5A-B**). We observed both dynamic and static cell-cell contacts. Dynamic contacts were
defined as cell-cell interactions that formed during an observation period of 30 min, whereas
static contacts were those that persisted throughout the 30-min period. A rapid increase in B7-
H3 intensity was detected at the contact sites of dynamic cell-cell interactions, whereas the
B7-H3-mNG signal at static contacts remained relatively stable (**Figure 5C**). Quantification of
relative mNG fluorescence across the lamellipodia of invading cells confirmed a pronounced
enrichment of B7-H3 at the leading edge (**Figure 5D**). This polar distribution and accumulation
were maintained in individual cells for at least 24 hours (**Figure 5E**).
The persistent accumulation of B7-H3 at the leading edge of invading cells and at the tip of
TME-exploring lamellipodia indicates a critical role of B7-H3 in lamellipodial guidance and
tumor invasion (**Figure 5G, Extended Data 4**).

Long-term, high-resolution imaging with PHIROS to monitor cell-cell interactions and organelle trafficking in tissue slices

With the established methodology, we were able to repeatedly image the same MB-astrocyte
clusters over three consecutive days without any loss of registration or resolution (**Figure 6A**).
This stability enabled morphological quantification of astrocyte features, including branching
parameters and astrocyte area (**Figure 6B-C**). Our analysis revealed an initial decrease in
astrocyte area, followed by an increase during which astrocytes reached similar area values
as at the assessment on the first day of the imaging assay. Notably, the number of branches
decreased over the three-day observation period, while the branch lengths increased. These
morphological changes could indicate an initial, transient state of astrocyte reactivity, triggered
by the proximity of the tumor cells, as has been observed in neurological injuries[35].
Furthermore, we observed astrocytic death in proximity to the tumor cells, likely caused by
tumor-astrocyte interactions (**Extended Data 5**).The ability to observe these morphological
alterations underlines PHIROS' capability to provide robust structural analysis in a complex
tissue environment.

In our acquisitions, we regularly observed microtubular connections between tumor cells and
cells of the cerebellar slices (**Figure 6D**). Some of these connections displayed moving
mitochondria, indicating mitochondrial transfer activities across F-actin-positive cell-cell
connections (**Figure 6G, Extended Data 5**). In pediatric and adult glial tumors, these
connections, referred to as tumor microtubes, generate interconnected networks of cells
implicated in therapy resistance[36,37]. Recently, they were also identified in patient-derived

xenografts of Li-Fraumeni SHH human medulloblastoma[38]. We observed that tumor-tumor
connections were shorter and thinner compared to the microtubes linking tumor cells and cells
of the microenvironment (**Figure 6E-F**). While the connections remained stable for several
hours, we observed a few events where cells disconnected, or tubes extended towards other
cells (**Extended Data 5**). We observed mitochondria moving through these microtubes
between tumor cells, a phenomenon not yet addressed in MB, with displacements of 22.8±16.1
μm and a mean speed of 0.012±0.01 μm/s (**Figure 6H-I**).

Discussion

Organotypic slices retain native cytoarchitecture and multicellular interactions for weeks ex
vivo, offering an advantageous model for studying complex cellular dynamics that dissociated
cultures or organoids cannot faithfully reproduce. Yet the fact that slices require ALI culturing
has prevented continuous high-resolution imaging. PHIROS directly addresses this limitation
by enabling multiday, subcellular-resolution imaging under controlled perfusion while
maintaining tissue viability. The platform integrates ALI culturing for tissue recovery and
extends experimental options, including AAV transduction and spheroid engraftment, while
pulsatile-perfusion culturing can be performed in a transparent, sealed chamber that is
compatible with high-NA objectives. The platform design ensures stable registration, aseptic
conditions, and, by relying on inert, biocompatible plastics, it eliminates adsorption issues
associated with soft polymers[39] .

Cerebellar slices retained an intact Purkinje cell layer and preserved laminar organization,
confirming tissue health. Although validated for the cerebellum, the developed protocols are
applicable to other brain regions, such as hippocampus or thalamus, which are challenging to
image dynamically *in vivo*[40].

PHIROS enabled long-term visualization of spontaneous Ca^{2+} activity at subcellular resolution,
revealing astrocytic Ca^{2+} heterogenic dynamics. Pulse delivery of a gap junction inhibitor
rapidly disrupted network activity, demonstrating the platform's capacity to probe acute and
sustained responses to targeted perturbations under physiological tissue conditions *in vitro*.

We further used PHIROS to dissect MB invasion in a tissue context. Tumor cells on organotypic
slices recapitulated patient and *in vivo*–like phenotypes more faithfully than in neuronal co-
cultures. The platform uncovered heterogeneous invasion behavior and helped to resolve
complex F-actin polymerization dynamics in invading and non-invading cells, illustrating how
MB cells sense and respond to the TME. Single-cell analyses further identified behaviorally
distinct MB clusters shaped by local cues. Tracking the transmembrane receptor B7-H3
revealed its persistent accumulation in tumor cell lamellipodia during invasion, suggesting an
additional role of B7-H3 in TME interaction and motility beyond its known immunomodulatory
functions[41].

Finally, PHIROS revealed dynamic remodeling of the surrounding TME, including astrocytic
processes, and enabled real-time visualization of actin-rich microtubes connecting MB cells,
which mediates organelle exchange, a process which has been linked to therapy resistance of
gliomas. The ability to track mitochondria trafficking demonstrates that PHIROS provides the
resolution and stability that are required to capture subcellular organelle dynamics over
multiple days, which is pivotal for studying intercellular communication in brain tumors and
neural diseases.

Taken together, our results establish PHIROS as a robust and broadly applicable platform for
long-term, high-resolution interrogation of cellular dynamics in intact physiological tissues,
enabling mechanistic studies that are not feasible with conventional *in vitro* or *in vivo*
approaches.

Acknowledgements

This work was financially supported by Innosuisse – the Swiss Innovation Agency- under grant 120.665 IP-LS. The authors acknowledge the Single Cell Facility (SCF) at D-BSSE of ETH Zürich for help and support, as well as Dr. Sreedhar Kumar and Dr. Michele Nava, also at D-BSSE, and Dr. Bernard Ciraulo, University Children's Hospital Zürich, for support with data interpretation and visualization.

Author contributions

J.B.P. designed and fabricated the platform, designed and performed all experiments, analyzed and interpreted tissue reorganization, tissue viability, hypoxia, calcium imaging, tumor microtube and mitochondria trafficking data, pre-processed the tumor invasion data, and wrote the manuscript. M.S.L designed and performed experiments, interpreted tissue reorganization, analyzed tumor invasion data, and edited the manuscript. C.M.G. analyzed and interpreted astrocyte morphology. D.H. generated and validated the fluorescent cell lines. M.B. conceived the project, acquired funding, provided experimental advice and supervision of the experiments, analyzed tumor invasion and B7H3 accumulation data, and edited the manuscript. A.H. supervised the project, provided experimental advice, and edited the manuscript. M.M.M. conceptualized the project, acquired funding, supervised the project, designed the platform, supported with experimental design and experiments, analyzed tissue viability and tumor invasion data, and edited the manuscript. All authors read and approved the final version of the manuscript.

Main figures

Figure 1: Platform design for long-term high-resolution imaging of organotypic slices

Figure 1: **(A)** Experimental set-up of the platform and workflow using mouse-derived cerebellar slices to investigate cellular dynamics, such as brain tumor motility or intracellular calcium dynamics. An exploded view of the platform design for both static and perfused configuration can be seen in **Supplementary 1.** (**B**) Photo of the developed platform placed on top of a tissue culture insert with two cerebellar slices. **(C)** Schematic of the developed fluidic system to maintain organotypic-slice health on a spinning-disc confocal microscope. A peristaltic pump was used to drive the medium from the main reservoir to the oxygenator (gas-exchanger), buffering the medium to a pH of 7.4, and oxygenating the medium before it reaches the culture chip in the environmental box of the microscope. A detailed view of the oxygenator and the measured dissolved oxygen concentrations can be seen in **Supplementary 2**. **(D)** The developed platform enables to study events across spatial and temporal scales, ranging from 4X or 10X magnification to single-cell dynamics and cell-cell interactions at 60X magnification over multiple days. Images show maximum intensity projections at the respective magnification.

Figure 2: Characterisation of chip-based cerebellar slice cultures and perfusion strategy for long-term imaging

A

Chip loading → Static culture → Perfused culture and live-cell imaging → Fixation

DIV 1   DIV 16   DIV 18

Fixation static charcterization

Tissue slice, Culture chip, 6-well plate, Tissue culture insert

Medium exchange, Tissue slice

Culture chip, Microfluidic channel, Sealing layers, PS foil

Oxygenated medium

Staining solution

4 % PFA in PBS

B

Molecular layer, Purkinje cell layer, Granule cell layer
500 µm   50 µm
Calbindin, GFAP, NeuN, DAPI, 50 µm

C

Purkinje cell soma volume [µm³]

PHIROS   Insert   ns

D

Purkinje cells per area [#/µm²]

PHIROS   Insert   ns

E

Perfusion Start - T 0h, DIV 16
Cleaved Caspase-3/7 (Apoptotic cells), Hoechst (Nuclei)

Day 2 perfusion culture - T 24h, DIV 17

Day 3 perfusion culture - T 48h, DIV 18

Day 3 perfusion culture - T 50h, Treated with 40µM Doxorubicin

100 µm

F

Normalized number of caspase-positive cells

Day 1, Day 2, Day 3, Doxorubicin controls

, ns, *, ***, ns, ns

G

Normalized increase in HIF1α-positive cells

ALI culture, 24h submerged static culture, 48h submerged static culture, 48h perfusion culture

ns, ***, ****, ns, ****, ****, ***

Figure 2: **(A)** Experimental design to assess tissue reorganization during double-membrane culturing
(dotted line endpoint at DIV 16) and to determine tissue viability under perfusion at DIV 18. **(B)** Stitched,
maximum-intensity projection image of a representative cerebellar slice. The integrity of the Purkinje cell
layer marked in yellow using anti-Calbindin is visualized. Composite image of the close-up in **B** with split
channels showing markers for anti-Calbindin (yellow), anti-GFAP (cyan), anti-NeuN (magenta) and DAPI
(blue). **(C)** Quantitative comparison of Purkinje cell volumes between cerebellar slices cultured on
culture chips (n= 6 chips/9 slices, 1658 cells) and on inserts (n=3 inserts, 754 cells). **(D)** Quantitative
comparison of Purkinje cell density per area between cerebellar slices cultured on chips (n= 6 chips)
and on inserts (n= 3 inserts). Statistics for C and D were derived using an unpaired t-test with Welch's
correction. **(E)** Composite, maximum-intensity projection images of the same slice stained for nuclei
(cyan) and apoptotic cells (cleaved caspase 3-7, magenta) after perfusion start, 24h into the perfusion
culture, 48h into the perfusion culture, and 2h post exposure to 40 µM doxorubicin. **(F)** Quantitative
assessment of caspase-positive cells (n = 6 chips/12 slices). **(G)** Quantification of hypoxic cells by using
immunostaining for anti-HIF1-alpha and nuclear co-localization. Results are normalized with respect to
hypoxic-cell counts found in standard ALI cultures. N = 4 slices per condition. Statistics for F and G were
derived using 1-way ANOVA.

Figure 3: Multi-day monitoring of spontaneous calcium dynamics in cerebellar tissue

Figure 3: (A) Experimental protocol to investigate intracellular calcium changes in cerebellar astrocytes;
the static culturing period was used to label astrocytes via AAV transduction. **(B-D)** Images of the same
ROI during the calcium imaging assay on day 1, day 2 and day 3 of the perfusion culture. Close-ups
show a calcium event indicated by the yellow arrow in the astrocytic process of likely the same cell
across the three time points. Scale bar of the close-ups is 20 µm. **(E)** Fluorescence intensity changes
of the events depicted in **B-D**. **(F)** Detected changes in calcium event area, duration, and changes in
fluorescence intensity from day 1 to day 3 of the imaging assay (n = 4 slices from 2 independent
experiments, 3-4 ROIs per slice), visualized as probability density functions. **(G)** Representative
example of calcium dynamics of velate astrocytes. The insets show a calcium event likely traveling
across multiple cells in the astrocyte network. **(H)** Comparison of different calcium-event metrics
between velate and Bergmann glia cells (n= 4 slices/ 2 chips each). **(I)** Calcium event in the same ROI
as in G after carbenoxolone exposure. **(J)** Comparison of different metrics of calcium events in velate
astrocytes before and after carbenoxolone exposure.

Figure 4: Dynamics of cell flow and single-cell trajectories in ONS76-LAeGFP spheroids.

Figure 4: **(A)** Experimental workflow to establish the medulloblastoma co-culture model for high-resolution imaging assays. **(B)** Representative image showing the ONS-76-LA-eGFP cells and astrocytes during 12-hour time-lapse imaging. Green: LA-eGFP; orange: GFAP-driven mCherry. Areas *i* and *iii* exhibit increased cell motility, while area *ii* shows confined cellular movement. **(C)** Heat maps display the mean velocity magnitude across the spheroid invading a slice (left) or migrating on dissociated neural cells (right) over 12 hours. Arrows indicate the mean motion direction. **(D)** Fitted mean velocity magnitudes from **(C)** plotted as a density distribution (n = 4 spheroids from slice co-cultures and n=3 spheroids from dissociated neural cell co-cultures) **(E)** Representative images of tumor cells invading slices displaying rounded and mesenchymal-like invasion phenotypes **(F)** Representative image of the actin-cytoskeleton of an invading medulloblastoma cell in the cerebellar slice. Arrows point to filopodia extending from the cell body. Quantifications show the normalized number of filopodia and filopodia lengths of MB invading the tissue (n= 12 cells from 4 slices/ 2 chips) **(G)** Representative images of untreated and 1 µM cytochalasin-D treated ONS-76 cells co-cultured with cerebellar tissue slices or neural dissociated cells. Arrowheads indicate exemplary TME-exploring lamellipodia, arrows filopodia in untreated cells. **(H)** ONS-76-LA-eGFP cell area, cell area variability, and cell elongation were assessed at day 2 under standard medium conditions and during CytoD exposure (D2 Ctrl and D2 CytoD, respectively), and 24h after exposure (D3 Ctrl and D3 CytoD). N = 4 control slices, n = 5 CytoD slices, n = 5 ROIs control neural co-cultures, n= 3 ROIs CytoD treated neural co-cultures). Nonparametric one-way ANOVA was used to derive statistical differences.

Figure 5: Persistent accumulation of B7-H3 at the leading edge of tissue-invading tumor cells

Figure 5: (A) Grayscale image of a polarized ONS-76-B7-H3-mNG cell on a tissue slice during perfusion culturing, two days after chip closure. The plot above the cell shows fluorescence values along the longitudinal profile (orange, horizontal line), while the plot on the right shows values of the cross-sectional profile (blue, vertical line). The schematic at the top right illustrates the B7-H3 fusion construct. **(B)** Maximum intensity projection of a confocal image stack after 48 hours of perfusion culture. Green: B7-H3-mNG; orange: GFAP-driven mCherry. The inset highlights the green channel, showing membrane-localized B7-H3 (arrowheads) and the leading-edge lamellipodium (arrow). **(C)** Dynamics of B7-H3-mNG at dynamic (top) versus static (bottom) cell-cell contacts. Heatmap kymographs show normalized fluorescence values along a 15-µm line (yellow dotted line) at cell-cell contacts, with arrows indicating B7-H3 accumulation. **(D)** Representative B7-H3-mNG fluorescence in tissue-invading tumor cells. Below: Normalized values of fluorescence intensity along a 30-µm line. **(E)** Normalized fluorescence values across leading-edge lamellipodia in cells invading during 24 hours (n=17) and 48 hours (n=21). **(F)** Left: Maximum intensity projection of confocal imaging after 48 hours of perfusion culture (green: B7-H3-mNG, orange: GFAP-driven mCherry). Right: Inverted grayscale of B7-H3-mNG fluorescence intensity shows lamellipodia dynamics over 30 minutes, with arrowheads indicating accumulation at the leading edge, whereas the initial position of the leading edge is visualized with a blue dotted line **(G)** Top: Green channel of image in B, with yellow-dotted arrows marking quantified areas of leading-edge lamellipodia. Bottom: Heat map kymographs show relative fluorescence values over 30 minutes along a 40–48 µm line across leading-edge lamellipodia of representative invading cells.

 Figure 6: Cell-cell interactions and mitochondria trafficking in a tissue context

Figure 6: (A): Interactions of the same cluster of ONS-76-La-eGFP medulloblastoma cells (green: LA-eGFP) with astrocytes (orange: GFAP-driven mCherry) over three days of perfusion culturing and imaging. Images are maximum intensity projections. **(B)** Number of branches and average branch length of astrocytes in close proximity to the tumor. **(C)** Cell areas of astrocytes in close proximity to the tumor cells. Data in (B) and (C) was derived from astrocytes of three different chips (n = 955 cells for D1, 702 cells for D2 and 1334 cells for D3 from 4 slices). Statistical differences between days were derived using a one-way ANOVA. **(D)** Representative maximum intensity projection image of a region featuring actin-rich tumor microtubular connections (green: LA-eGFP) and mitochondria (magenta: MitoTracker deep red). Insets show 3D renderings of a tumor-tumor connection (framed in green) and a tumor-TME connection (framed in blue). **(E)** Length and **(F)** diameter of the tumor-tumor connections (n = 124 TMs from 6 chips/14 slices) and tumor-TME microtubes (connections between medulloblastoma and cells of the TME (blue triangles, n = 78 connections from 6 chips). **(G)** 3D renderings of different timepoints showing a mitochondria (magenta: MitoTracker deep red) transfer between two connected medulloblastoma cells (green: La-eGFP) and in a tumor-TME microtube (blue: LA-eGFP) protrusion connecting to the TME (in blue: LA-eGFP). Colored arrows indicate the movement of the same mitochondria between frames. **(H)** Displacement and **(I)** speed of mitochondria in tumor-tumor connections (n=61 from 6 chips/14 slices) and tumor-TME connections (n=31 from 6 chips/14 slices. Data in G, H, J, K, L was first tested for normal distribution before applying either a Welch's t-test for lognormal distributed data or a Mann-Whitney t-test.

Extended Data

Extended Data 1: **(A)** 10X maximum intensity projections of the astrocytic networks labeled with anti-S100β in magenta and anti-GFAP in cyan. Close-ups show 40X maximum intensity projections of the same, representative cerebellar slices cultured on PHIROS. **(B-C)** 10X maximum intensity projections of the Purkinje cell layer of slices cultured on PHIROS and stained with anti-Calbindin after fixation at DIV 14 of two additional experiments. Close-ups show 40X images of the Purkinje cell reorganization. The red line in close-up **C** indicates how the boundaries of the Purkinje cell layer were defined to assess the Purkinje cell density. **(D)** Purkinje cell reorganization after 14 days of culturing during which slices were cultured on standard tissue-culture inserts. The close-ups at 40X show the presence of different neuronal cell types, identified by the anti-Calbindin and NeuN staining, as well as glial cells, as indicated by the anti-GFAP staining.

Extended Data 2: ROIs of different slice experiments of **Figure 2**. **(A-C)** Viability assessment using cleaved Caspase-3/7 stain labeling apoptotic nuclei (magenta) co-stained with NucBlue (cyan). The images show the same tissue slice during 48 h of perfusion culturing. **(D-F)** Representative immunofluorescent staining for transcription factor HIF1-α (red) and nuclei (DAPI, in blue) of ALI cultured slices (**D**), slices cultured for 24h (**E**) or 48h (**F**) submerged under 2 ml of medium, or for 48h under perfusion on PHIROS (**G**).

Extended Data 3: (A) Representative image showing tumor-astrocyte interactions during 12-hour time-lapse imaging. Area *ii* highlights a region with low astrocyte density (orange: GFAP-driven mCherry), where pronounced ONS-76-La-eGFP cell movement (green: LA-eGFP) was observed (arrows indicate movement direction). Area *i* shows higher astrocyte density, where cell movement was minimal. Time refers to time after platform sealing; the scale bar of both sets of insets is 20 μm; **(B)** Example of repulsive and attractive interactions between tumor cells (green) and astrocytes (orange) over 14 hours. Movements of cells 1, 2, and 3 are shown with arrows representing their respective trajectories. **(C)** Individual cell trajectories on a background image show only astrocytes at the imaging starting point, illustrating movement patterns in relation to astrocyte density.

Extended Data 4: (A) Left: Green channel of the image in **Figure 5B**, with yellow-dotted arrows indicating additionally quantified areas of leading-edge lamellipodia. Right: Heat map kymographs show relative gray fluorescence values of the green fluorescence channel over 30 minutes along a 48 µm line across leading-edge lamellipodia in representative invading cells, in addition to the cells shown in **Figure 5G**. **(B-C)** Inverted grayscale of B7-H3-mNG fluorescence intensity of two cells from **(A)**, indicated by the dotted bounding box. The images show lamellipodia dynamics over 30 minutes of acquisition, with arrowheads indicating accumulation at the leading edge. The initial position of the leading edge is visualized with the blue dotted line.

Extended Data 5: (A) Example of an observed disrupting tumor-tumor cell connection. Green: LA-eGFP; Magenta: MitoTracker deep red. **(B)** Still frames of a microtube extending from an ONS-76-LA-eGFP cell towards the TME. The arrows indicate the tip of the microtube. Green: LA-eGFP; orange: GFAP-driven mCherry; magenta: MitoTracker deep red. **(C)** Still frames of an event where tumor-cell contact likely elicited astrocyte death. **(D-E)** Representative images showing live observations of tumor cells interconnected through long, actin-rich protrusions in two additional independent experiments. Images show maximum intensity projections of still images. All scale bars in the close-ups are 20 μm.

References

Baker, B. M. & Chen, C. S. Deconstructing the third dimension – how 3D culture microenvironments alter cellular cues. *Journal of Cell Science* **125**, 3015-3024 (2012). https://doi.org:10.1242/jcs.079509

Jensen, C. & Teng, Y. Is It Time to Start Transitioning From 2D to 3D Cell Culture? *Frontiers in Molecular Biosciences* **7** (2020). https://doi.org:10.3389/fmolb.2020.00033

Humpel, C. Organotypic brain slice cultures: A review. *Neuroscience* **305**, 86-98 (2015). https://doi.org:10.1016/j.neuroscience.2015.07.086

Croft, C. L., Futch, H. S., Moore, B. D. & Golde, T. E. Organotypic brain slice cultures to model neurodegenerative proteinopathies. *Mol Neurodegener* **14**, 45 (2019). https://doi.org:10.1186/s13024-019-0346-0

Stoppini, L., Buchs, P. A. & Muller, D. A simple method for organotypic cultures of nervous tissue. *J Neurosci Methods* **37**, 173-182 (1991). https://doi.org:10.1016/0165-0270(91)90128-m

Eisemann, T. *et al.* An advanced glioma cell invasion assay based on organotypic brain slice cultures. *BMC Cancer* **18**, 103 (2018). https://doi.org:10.1186/s12885-018-4007-4

Santhana Kumar, K. *et al.* TGF-β Determines the Pro-migratory Potential of bFGF Signaling in Medulloblastoma. *Cell Reports* **23**, 3798-3812.e3798 (2018). https://doi.org:https://doi.org/10.1016/j.celrep.2018.05.083

Neve, A., Santhana Kumar, K., Tripolitsioti, D., Grotzer, M. A. & Baumgartner, M. Investigation of brain tissue infiltration by medulloblastoma cells in an ex vivo model. *Sci Rep* **7**, 5297 (2017). https://doi.org:10.1038/s41598-017-05573-w

Steindl, A. & Valiente, M. Potential of ex vivo organotypic slice cultures in neuro-oncology. *Neuro-Oncology* (2024). https://doi.org:10.1093/neuonc/noae195

Comba, A. *et al.* Spatiotemporal analysis of glioma heterogeneity reveals COL1A1 as an actionable target to disrupt tumor progression. *Nature Communications* **13**, 3606 (2022). https://doi.org:10.1038/s41467-022-31340-1

Mangukiya, H. B. *et al.* Reconstructing the Single-Cell Spatiotemporal Dynamics of Glioblastoma Invasion. *bioRxiv*, 2025.2003.2020.644331 (2025). https://doi.org:10.1101/2025.03.20.644331

Ravi, V. M. *et al.* Human organotypic brain slice culture: a novel framework for environmental research in neuro-oncology. *Life Science Alliance* **2**, e201900305 (2019). https://doi.org:10.26508/lsa.201900305

Pan, Y. *et al.* NF1 mutation drives neuronal activity-dependent initiation of optic glioma. *Nature* **594**, 277-282 (2021). https://doi.org:10.1038/s41586-021-03580-6

Quail, D. F. & Joyce, J. A. The Microenvironmental Landscape of Brain Tumors. *Cancer Cell* **31**, 326-341 (2017). https://doi.org:10.1016/j.ccell.2017.02.009

Winkler, F. *et al.* Cancer neuroscience: State of the field, emerging directions. *Cell* **186**, 1689-1707 (2023). https://doi.org:10.1016/j.cell.2023.02.002

16 Charles, N. A., Holland, E. C., Gilbertson, R., Glass, R. & Kettenmann, H. The brain
tumor microenvironment. *Glia* **60**, 502-514 (2012).

17 Hanahan, D. & Monje, M. Cancer hallmarks intersect with neuroscience in the tumor
microenvironment. *Cancer Cell* **41**, 573-580 (2023).
https://doi.org:10.1016/j.ccell.2023.02.012

18 Chen, Q. *et al.* Carcinoma–astrocyte gap junctions promote brain metastasis by
cGAMP transfer. *Nature* **533**, 493-498 (2016). https://doi.org:10.1038/nature18268

19 Parker, J. J., Lizarraga, M., Waziri, A. & Foshay, K. M. A Human Glioblastoma
Organotypic Slice Culture Model for Study of Tumor Cell Migration and Patient-
specific Effects of Anti-Invasive Drugs. *J Vis Exp* (2017).
https://doi.org:10.3791/53557

20 Gahwiler, B. H. Organotypic monolayer cultures of nervous tissue. *J Neurosci
Methods* **4**, 329-342 (1981). https://doi.org:10.1016/0165-0270(81)90003-0

21 Northcott, P. A. *et al.* Medulloblastoma. *Nat Rev Dis Primers* **5**, 11 (2019).
https://doi.org:10.1038/s41572-019-0063-6

22 Keller, D., Erö, C. & Markram, H. Cell Densities in the Mouse Brain: A Systematic
Review. *Frontiers in Neuroanatomy* **12** (2018).
https://doi.org:10.3389/fnana.2018.00083

23 Semyanov, A., Henneberger, C. & Agarwal, A. Making sense of astrocytic calcium
signals — from acquisition to interpretation. *Nature Reviews Neuroscience* **21**, 551-
564 (2020). https://doi.org:10.1038/s41583-020-0361-8

24 Bazargani, N. & Attwell, D. Astrocyte calcium signaling: the third wave. *Nature
Neuroscience* **19**, 182-189 (2016). https://doi.org:10.1038/nn.4201

25 Mi, X. *et al.* Fast, accurate, and versatile data analysis platform for the quantification
of molecular spatiotemporal signals. *Cell* **188**, 2794-2809.e2721 (2025).
https://doi.org:https://doi.org/10.1016/j.cell.2025.03.012

26 Farmer, W. T. *et al.* Neurons diversify astrocytes in the adult brain through sonic
hedgehog signaling. *Science* **351**, 849-854 (2016).
https://doi.org:doi:10.1126/science.aab3103

27 Hoogland, T. M. & Kuhn, B. Recent Developments in the Understanding of Astrocyte
Function in the Cerebellum In Vivo. *The Cerebellum* **9**, 264-271 (2010).
https://doi.org:10.1007/s12311-009-0139-z

28 Almagro, J., Messal, H. A., Elosegui-Artola, A., van Rheenen, J. & Behrens, A. Tissue
architecture in tumor initiation and progression. *Trends in Cancer* **8**, 494-505 (2022).
https://doi.org:10.1016/j.trecan.2022.02.007

29 Alieva, M. *et al.* Intravital imaging of glioma border morphology reveals distinctive
cellular dynamics and contribution to tumor cell invasion. *Scientific Reports* **9**, 2054
(2019). https://doi.org:10.1038/s41598-019-38625-4

30 Winkler, F. *et al.* Imaging glioma cell invasion in vivo reveals mechanisms of
dissemination and peritumoral angiogenesis. *Glia* **57**, 1306-1315 (2009).
https://doi.org:https://doi.org/10.1002/glia.20850

31 Vilchez Mercedes, S. A. *et al.* Decoding leader cells in collective cancer invasion.
Nature Reviews Cancer **21**, 592-604 (2021). https://doi.org:10.1038/s41568-021-
00376-8

32 Mayor, R. & Etienne-Manneville, S. The front and rear of collective cell migration.
Nature Reviews Molecular Cell Biology **17**, 97-109 (2016).
https://doi.org:10.1038/nrm.2015.14

33 Haydar, D. *et al.* Cell-surface antigen profiling of pediatric brain tumors: B7-H3 is
consistently expressed and can be targeted via local or systemic CAR T-cell delivery.
Neuro Oncol **23**, 999-1011 (2021). https://doi.org:10.1093/neuonc/noaa278

34 Epperly, R., Gottschalk, S. & DeRenzo, C. CAR T cells redirected to B7-H3 for
pediatric solid tumors: Current status and future perspectives. *EJC Paediatr Oncol* **3**
(2024). https://doi.org:10.1016/j.ejcped.2024.100160

35 Zhou, B., Zuo, Y.-X. & Jiang, R.-T. Astrocyte morphology: Diversity, plasticity, and role
in neurological diseases. *CNS Neuroscience & Therapeutics* **25**, 665-673 (2019).
https://doi.org:https://doi.org/10.1111/cns.13123

36 Hausmann, D. *et al.* Autonomous rhythmic activity in glioma networks drives brain
tumour growth. *Nature* **613**, 179-186 (2023). https://doi.org:10.1038/s41586-022-
05520-4

37 Venkataramani, V. *et al.* Disconnecting multicellular networks in brain tumours. *Nature*
Reviews Cancer **22**, 481-491 (2022). https://doi.org:10.1038/s41568-022-00475-0

38 Kats, I. *et al.* Spatio-temporal transcriptomics of chromothriptic SHH-medulloblastoma
identifies multiple genetic clones that resist treatment and drive relapse. *Nature*
Communications **15**, 10370 (2024). https://doi.org:10.1038/s41467-024-54709-w

39 Toepke, M. W. & Beebe, D. J. PDMS absorption of small molecules and
consequences in microfluidic applications. *Lab on a Chip* **6**, 1484-1486 (2006).
https://doi.org:10.1039/B612140C

40 Grienberger, C., Giovannucci, A., Zeiger, W. & Portera-Cailliau, C. Two-photon
calcium imaging of neuronal activity. *Nature Reviews Methods Primers* **2**, 67 (2022).
https://doi.org:10.1038/s43586-022-00147-1

41 Zhao, B. *et al.* Immune checkpoint of B7-H3 in cancer: from immunology to clinical
immunotherapy. *J Hematol Oncol* **15**, 153 (2022). https://doi.org:10.1186/s13045-022-
01364-7

568