Family-based Selection: An Efficient Method for Increasing Phenotypic Variability

Shraddha Lall[1], Chris Milton[1], and Benjamin de Bivort[1,2,*]

[1]Department of Organismic and Evolutionary Biology, Harvard University

[2]Center for Brain Science, Harvard University

*Corresponding author: debivort@oeb.harvard.edu

Abstract

Persistent idiosyncrasies in behavioral phenotypes have been documented across animal taxa. These individual differences among organisms from the same genotype and reared in identical environments can result in phenotypic variability in the absence of genetic variation. While there is strong evidence to suggest that variability of traits can be heritable and determined by the genotype of an organism, little is known about how selection can specifically shape this heritable variance. Here, we describe a Python-based model of directional artificial selection for increasing the variability of a polygenic trait of interest. Specifically, our model focuses on variability in left-vs-right turn bias in *D. melanogaster*. While the mean value of turn bias for a genotype is non-heritable and constant across genotypes, the variability of turn bias is a heritable and polygenic trait, varying dramatically among different genetic lines. Using our model, we compare different selection regimes and predict selection dynamics at population and genetic levels. We find that introducing population structure via a family-based selection regime can significantly affect selection response. When selection for increased variability is implemented on the basis of independently measured traits of individuals, the response is slower, but leads to a population with a greater genetic diversity. In contrast, when selection is implemented by measuring traits of families with half or full siblings, the response is faster, albeit with a reduced final genetic diversity in the population. Our model provides a useful starting point to study the effect of different selection regimes on any polygenic

trait of interest. We can use this model to predict responses of laboratory-based selection experiments and implement feasible experiments for selection of complex polygenic traits in the lab.

Introduction

Individuals exhibit idiosyncratic behavioral phenotypes, even in the absence of genetic variation [10, 34]. These behavioral differences persist over long periods of time and have been found in behaviors such as activity, aggression [4, 7, 22] and exploration [13, 26, 42], in species such as fruit flies [20], Amazon mollies [3], and mice [16]. At the mechanistic level, such differences may be an unavoidable consequence of the stochastic nature of molecular and cellular function, particularly during development [37, 44]. Clonal single-celled organisms exhibit a significant amount of phenotypic variability, attributed to stochasticity in gene expression [15]. In fruit flies, 23% of genes were differentially expressed among genetically identical individuals reared under essentially identical laboratory conditions [23]. Stochasticity in gene expression may affect behavior through its impact on neuronal physiology, and specific neurons have been implicated in determining specific idiosyncratic behaviors. For example, the degree of left-right asymmetry in the wiring of fruit fly dorsal cluster neurons (DCN) has been shown to predict individual differences in visual object orientation in freely walking flies [24]. Individual differences in calcium responses [19] of specific glomeruli in the antennal lobe of fruit flies predict individual odor preferences [9]. In zebrafish, short-term habituation of the acoustic startle response has been found to be modulated by serotonergic neurons of the dorsal raphe nucleus (DRN) at the individual level [29].

At first glance, the stochastic, noisy nature of molecular processes suggests that intragenotypic behavioral variability may be non-adaptive. But theoretical models [12, 25, 47, 28], and empirical studies from microbes [36, 2] and plants [30, 36, 35, 8] suggest that phenotypic variability can, in some circumstances, be adaptive, allowing organisms to "hedge their bets" against fluctuations in the environment. If variability is adaptive, it may be the object of evolution by natural selection, provided there is heritable variation for the level of variability. Evidence that variability of a trait is under genetic control (and therefore heritable) is mounting, with empirical studies of morphology [11, 14, 41] and behavior [1, 40] supporting this. These studies suggest that selection can act to increase or decrease variability in trait

values. Consequently bet-hedging strategies can potentially be fine-tuned to the ecological and environmental fluctuations that favor them.

In *Drosophila melanogaster*, locomotor handedness is a behavioral measure that is readily estimated [5] and suitable for experimental evolutionary genetics. Flies have idiosyncratic tendencies to turn left or right in Y-shaped mazes or open arenas [6]. The distribution of average turn biases across flies is unimodal and broad. That is, the most common bias is symmetrical left-vs-right, but some individuals have strong left- or right-biases. Notably, different wild-derived inbred fruit fly lines exhibit differing degrees of intragenotypic variability in this behavior. This variation has a complex genetic architecture with many sites of small effect correlated with the extent of variability [1]. Further, the mean value of turn-bias, regardless of genotypes, is ~0.5 [1, 6]. The mean value of turn bias therefore has very low broad sense heritability and is unlikely to respond to selection. In contrast, variability of turn bias does have substantial heritability and could potentially respond to selection. Demonstrating that variability of a behavioral trait can indeed evolve under artificial selection would be an important step in establishing whether it can evolve in natural circumstances as an adaptive strategy. Locomotor handedness, measured as turn bias, offers several advantages for such an effort, including little risk that selection on trait variability has confounding interactions with the trait mean.

Different approaches can be used over the course of artificial selection to choose which individuals are allowed to mate and contribute to the next generation. In mass selection, also known as individual-based selection, an individual is chosen based on its own score on the trait being selected. Alternatively, selection decisions can be made considering the trait scores of the individual and its relatives (an approach termed family-based or family selection) [45]. The latter approach may enrich the statistical signal in selected individuals, as related individuals may have correlated phenotypes. Additionally, family-based selection can be more likely to maintain allelic combinations that contribute to the selected phenotype non-additively. The families chosen in each generation may be composed of half siblings, often with the same mother but different potential fathers, or of full siblings, generated from one breeding pair. Family-based selection is often employed in plant and animal breeding [39], and is sometimes employed in lab-based selection studies [17, 49].

Laboratory-based selection experiments are long and labor-intensive. Further, factors such as the population size used in the experiment and the number of generations for which

selection should be implemented are difficult to predict in advance, and can have significant impacts on the success of the experiment [33]. Theoretical and computational models can be helpful in this regard, allowing one to predict the dynamics of selection at a population and genetic level prior to embarking on a selection experiment. In this paper, we describe an individual-based model of directional artificial selection on a polygenic trait of interest, specifically behavioral variability in locomotor handedness in *Drosophila melanogaster*. Informed by empirical data [1], this model compares the phenotypic and genotypic responses of mass versus family-based selection regimes (both half-sib and full-sib). This study is novel in focusing on a selected trait that is measured using many individuals, the variance of individual behavior. Using this model, we predict how different selection regimes change the underlying genetic architecture of the evolving population, with focus on how epistatic interactions can modulate this change. While our model is motivated by an interest in behavior, it is generalizable and can be readily extended to a diversity of traits with differing genetic architectures. Importantly, it provides specific predictions about responses to selection on a phenotypic and genetic level, helping biologists optimize the parameters of their selection regime before committing to large-scale experiments.

Methods

Our model assumes a polygenic trait with additive and epistatic components. All individuals are diploid, and reproduction is sexual with a fixed recombination rate. There are no *de novo* mutations. Generations are non-overlapping, with a fixed, finite population size. These assumptions are implemented in the model through variables whose default values are given in Table 1. We chose values appropriate to behavioral experiments in *Drosophila*, but these variables can be readily changed for traits with different architectures or to model populations of various sizes and family compositions.

The model was implemented entirely in Python 3. Initial code optimization was done on a desktop-PC, and the full model was run on the FAS Research Computing Cluster at Harvard University. Statistical analyses were also performed in Python 3, using built-in numpy [18] and scipy [43] functions. Annotated code can be found at: lab.debivort.org/family-selection-for-variability

Table 1: Model variables and their default values

Variable	Variable Symbol	Default Value
Population Size	N	980
Number of Sites	L	500
Generations	G	100
Per-base Recombination Rate	ρ	0.002
Percent Strength of Selection	s	20%
Number of Trials per Individual	t	500
For Mass Selection		
Number of Individuals Screened	N_{sc}	$\frac{N}{2}$
Number of Individuals Selected	N_{sel}	$\frac{s}{100} \times N$
For family-based selection		
Number of Families	f	35
Number of Individuals per Family	n	$\frac{N}{f}$
Individuals Screened per Family	n_{sc}	$\frac{n}{2}$
Number of Families Selected	f_{sel}	$\frac{s}{100} \times f$

We assumed that variability as a trait is determined by L sites, each of which has two
possible alleles: one that has no effect and one that contributes a direct additive effect and
contributes to epistatic effects. Specifically, we assumed that additive effects (A_i) come from
a beta prime distribution (Fig. S1), which closely matches the empirical distribution of effect
sizes for SNPs associated with variability in locomotor handedness (Ayroles et al., 2015). We
used beta prime parameters of $a = 0.976$, $b = 5.789$, loc $= 2.79 \times 10^{-7}$ and scale $= 0.272$. To
model epistatic effects, we defined an L x L symmetrical interaction matrix with each entry
$E_{i,j}$ drawn from an exponential distribution with parameter λ and sign randomly assigned
as positive or negative with equal probability (i.e., a Laplace distribution with mean 0 and
variance $2\lambda^{-2}$). The value of λ was determined based on the desired relative contribution of

epistasis to the overall effect size, further detailed below. The values of site effect sizes and the epistatis interaction matrices were drawn once and fixed for all runs and all regimes, thus avoiding any effects introduced by random variation in effect sizes. All individuals in our simulations were diploid, so each site has two loci — individuals could have 0, 1 or 2 alleles at each site with an effect on variability. Thus, the variability trait value for individual a was given by

$$V_a = \sum_{i=1}^{L_a} n_{a,i} A_i + \sum_{i=1}^{L_a} \sum_{j=i+1}^{L_a} E_{i,j}$$

where L_a is the number of sites in individual a for which at least one allele has an effect on variability, i and j index those sites, and $n_{a,i}$ is, for individual a, the number of alleles at site i which have an effect on variability (1 or 2 depending whether that site is heterozygous or homozygous). Three levels of epistasis, defined as the ratio of the variance in epistatic effects to the overall variance of genotypic values across the population, were simulated. Specifically, for high epistasis, $\frac{Var(E)}{Var(V)} = 0.5$, $\lambda = 0.0021$; for medium epistasis, $\frac{Var(E)}{Var(V)} = 0.25$, $\lambda = 0.0012$; for low epistasis, $\frac{Var(E)}{Var(V)} = 0.125$, $\lambda = 0.00079$.

In our model, an individual's genotype determines their potential to have a phenotype far from the mode [1]. In other words, if an individual comes from a high variability genotype, it will have a phenotype that is drawn from a wider distribution of potential trait values. The particular behavioral trait inspiring this analysis, locomotor handedness, has a mean value very close to 0.5 across all assessed genotypes [5, 6, 1]. Therefore, we used each individual's V_a value to compute a true locomotor handedness phenotype P_a from a normal distribution with mean 0.5. In order to make the standard deviation for these distributions match empirical data, V_a values were scaled and matched via linear interpolation (using the `interp` function in numpy) to a distribution of empirical standard deviations of locomotor handedness for 167 genotypes (Fig. S2). Accordingly, for each individual, the phenotype was determined as: $P_a \sim Norm(0.5, (d(V_a))^2)$, where $d(V_a)$ is the genotypic value after mapping. If the sampled value was less than 0 or greater than 1, we set it to 0 or 1, respectively. Once the true phenotype of an individual was assigned, we could determine its phenotype as would be measured in an actual selection experiment (π_a). Locomotor handedness is measured as the fraction of t turns that an animal makes to the right. Thus, $\pi_a \sim Binom(t, P_a)/t$.

New offspring individuals in our simulations were generated via sexual reproduction. The sex ratio at birth was fixed at 50:50, but the distributions of fitness varied by sex. All

females had equal probability of being a parent. However, fruit fly males do not have equal likelihood of successfully mating and fathering offspring. Accordingly, the probability of each male being a parent was drawn from the empirical distribution of male fitness in Fig. 1 of Pischedda and Rice, 2011 [32]. For each new offspring, each of their diploid parents produced two gametes, with recombination. Each site had an independent probability ρ of being a site of recombination. For our default values of $L = 500$ sites and recombination rate $\rho = 0.002$, there was on average one recombination event per haploid genome per generation. Of the two gametes generated by this process, one was chosen at random as the haploid contribution to the offspring. Note, in *Drosophila* there is no recombination in males [27], so in this respect our simulation was more typical of other species.

In every generation of both family-based and mass selection simulations, we implemented a strength of selection $s = 0.2$ or 20%. For simulations of family-based selection, the initial population was seeded with f families, each of which have n individuals, for a total population size of $f * n = N$. The default values of $f = 35$ and $N = 980$ were chosen as plausible numbers for a potential real implementation of this experiment. Measured phenotypes (π_a) were computed for all of the individuals from each family. The measured variability for each family m, in generation 0, was calculated as the standard deviation of individual phenotypes: $Var_m(0) = \sigma(\pi_a)$. s% of f families with the highest variability were selected to seed the next generation (Fig. 1A). The average population variability was estimated as the average of family variabilities for all f families: $Var_0 = \mu(Var_m(0))$

For half-sib selection, individuals from all selected families were pooled, and f families were seeded for the next generation by randomly picking f multiply-mated females from the pool. For full-sib selection, male and female individuals were randomly paired from among the selected families to create f monogamous pairs, that were used to generate f families for the next generation. The variability for all families in the population ($Var_m(g)$) was estimated at every generation g, and selection was carried out as before.

For simulations of mass selection, the initial population was seeded with N individuals. Measured phenotypes (π_a) were computed for all of the individuals in the population. The variability of this population was calculated as the standard deviation of the measured phenotypes: $Var_0 = \sigma(\pi_a)$. s% of N individuals were chosen to seed the next generation, such that the selected individuals formed a subset with the highest possible variability (Fig. 1B). This was implemented by sampling across the distribution of individuals, selecting those that

189 have the highest probability of being from a target high variability distribution. The next
190 generation of N individuals was created via reproduction among all selected individuals.
191 The variability of the population Var_g was estimated at each generation, and selection was
192 carried out as before.

Figure 1: **Variability selection regimes**

(A) Schematic of family-based selection regime used in the model, showing half-sib and
full-sib variants; (B) Mass selection regime; (C) Change in turn bias variability ($\Delta Var_g =
Var_g - Var_0$) over 100 generations of mass, half-sib and full-sib selection. Lines are the
average ΔVar_g at each generation g, averaged across $k = 10000$ replicate runs of model.
Shaded regions indicate ± 1 standard deviation due to run-to-run variation $(\sigma_r)_g$ estimated
as the standard deviation of ΔVar_g across k replicates; dotted line indicates the variability
value of a population where all individuals have the maximum variability trait value V_a
possible with this genetic architecture.

To track the genotypic changes associated with selection, each variability site was evaluated for fixation at the end of selection. All selection regimes with various combinations of parameters were simulated k times under different random seeds to assess variability in selection response trajectories.

Results

Family-based selection for variability is faster than mass selection

Starting with a simple additive model (no epistasis), our simulations showed robust responses to selection for higher turn bias variability (Fig. 1C). Family-based selection (half-sib or full-sib) yielded a faster response than mass selection. Half-sib family-based selection yielded 75% of its total response to selection by generation 6, whereas mass selection attained 75% of its final response by generation 11. Full-sib selection was slightly faster than half-sib selection, attaining 75% of its final response approximately 2 generations sooner. We additionally implemented a particularly aggressive form of mass selection, by selecting 20% of the individuals with the most extreme turn-bias phenotypes at every generation. The response with this method was faster than the default mass selection regime, but was still slower than both half and full-sib family-based selection through at least 5 generations (Fig. S3). However, a downside of such a strategy is the possibility of evolving an undesirable bimodal phenotype distribution if the mean of the trait is heritable.

The change in variability over 100 generations of selection was consistent across all regimes. The final variability attained was, on average, almost as high as the theoretical maximum possible value, calculated as the variability of a population where all the individuals have all the sites fixed at the high variability allele (Fig. 1C).

Family-based selection for variability fixes more sites than mass selection

Many more variability-determining sites were fixed in 100 generations of family-based selection compared to mass selection, with full-sib selection fixing 94.9% of all sites on average (Fig. 2A). This was an expected consequence of the lower effective population size, and therefore higher inbreeding in family-based selection [31]. Sites with large effects on turn

bias variability had higher fixation probabilities in all regimes (Fig. 2B-D), as expected. This relationship was strongest in mass selection ($r^2 = 0.81$ using a second-degree polynomial fit), weaker in half-sib selection ($r^2 = 0.20$), and weakest in full-sib selection ($r^2 = 0.07$), where a large number of sites were likely to fix regardless of their effect size (Fig. 2D).

Figure 2: **Fixation of variability determining sites after 100 generations of selection** (A) Mean fixation percentage of sites at generation 100 for the three selection regimes, averaged over 1000 runs. (B-D) Site fixation probability versus effect size in mass, half-sib and full-sib regimes, respectively. Lines are second-degree polynomial fits. Tick marks at bottom indicate the values of site effect sizes, which were fixed across all runs.

Family structure modulates selection response

To determine the effects of family and population size on selection response, we systematically varied these parameters (f and N) and measured peak variability change $\Delta Var_{100} =$

$Var_{100} - Var_0$. Selection response was smaller at both extremes of f (Fig. 3A). The smallest response to selection was found with populations with few families of many individuals each (low f, high n). Slightly higher variability was attained with populations made of a large number of families with few individuals in each family (high f, low n). The greatest selection responses were found in populations with families of intermediate size. Regardless of family composition, increasing population size boosted the selection response.

Run-to-run noise (σ_r) can affect the reliability of selection, an important consideration for artificial selection experiments that might actually be implemented in the lab. With fewer, larger, families (low f, high n), selection responses were more variable (Fig. 3B). Populations with more, small families (high f, low n) were generally less subject to noisy fluctuations. Unsurprisingly, increasing total population size reduced run-to-run variability. Overall, our simulations indicated that intermediate number of moderate sized families can optimize selection for variability at various population sizes, allowing for reliable, consistently high responses.

Figure 3: **Effects of population structure in family-based selection**
Changes in (A) peak variability (ΔVar_{100}) and (B) run-to-run variation (σ_r) in selection response over 100 generations of half-sib selection. Each row has a fixed total population size, with population structure varied by changing the distribution of animals within families. The leftmost column corresponds to a many families with few individuals in each, while the rightmost column corresponds to fewer, larger families.

Selection Outcomes are Robust to Change in Model Parameters

Changing model parameters (in Table 1) altered the selection responses quantitatively, but did not alter the overall trends and differences between the selection regimes. For example, changing the number of sites that contribute to phenotypic variability from 63 to 1000 had little discernible effect on the selection response in any of the regimes (Fig. S4). Likewise, changing the distribution of site effects had no qualitative effect on selection outcomes (Fig. S5). This suggest that results from our model apply to traits with a variety of effect size statistics.

When selection was weaker (i.e., more individuals were chosen to found subsequent generations) the selection response was slower, as expected. This effect was most pronounced under mass selection (Fig. S6). Modulating population size had similar effects across all regimes, with larger population sizes showing faster and stronger responses to selection. This effect was more pronounced in family-based selection, where the effective population size is reduced due to inbreeding (Fig. S7).

Epistasis Attenuates Selection Responses

Our model accommodates pairwise epistatic interactions between sites that additively contribute to variability. These epistatic effects can be positive or negative (Fig. 5B), increasing or decreasing an individual's latent variability (V_a). When epistasis was added to the model, the response to selection was slower across all selection regimes. However, over 100 generations, a similar value of variability was attained (Fig. 4A). The effect of epistasis was quantitative: increasing the magnitude of epistatic interactions resulted in slower selection responses (Fig. 4B, Supplementary Figure 9).

Sites Are Differentially Fixed With and Without Epistasis

We examined the effect of epistasis on the probability of site fixation after selection (Fig. 5A). In our model, a site's additive effect and its average epistatic effect were uncorrelated(Fig. 5B). How these two effects determined a site's fixation probability depended on the selection regime. In general, fixation probabilities were higher when both additive and epistatic effects were included (Fig. 5C-H). This difference was more pronounced in family-selection regimes, where the population structure could preserve combinations of linked sites with

Figure 4: **Effect of epistasis on selection response**

(A) Average change in variability versus generations of selection, with high level of epistasis $(\frac{Var(E)}{Var(V)} = 0.5)$ and without epistasis, in different selection regimes; (B) Average change in variability versus generations of mass selection, with varying magnitudes of epistatic interactions.

strong epistatic interactions. Unsurprisingly, the fixation probability of a site depended on its additive effect, with larger effect sites being more likely to fix (Fig. 2C-E). This relationship was strongest when epistasis was not included in the model.

Conversely, sites that had very positive average epistatic effects were more likely to fix when the model included epistatis (Fig. 5F-H). This was true even for sites with low additive effects, on account of their large positive epistatic contribution to variability. Similarly, sites that had highly negative epistatic effect sizes fixed with a higher probability with epistasis, potentially due to their strong negative effects on variability driving them to extinction (Fig. 5F-H).

Family-based Selection is Particularly Efficient for Variance- but not Mean-Based Traits

In our model, family-based selection produced a faster response than mass selection when the trait under selection was variance-based, such as turn bias variability (Fig. 1C, Fig. 6A). We hypothesized that the efficiency of family-based selection resulted from the concordance of the unit of selection (families) and a trait that can only be measured from multiple individuals (variance). To test this hypothesis, we used the same model (genetic architecture, selection

Figure 5: **Effect of epistasis on fixation of variability-influencing sites**
(A) Schematic for interpreting fixation probability plots in (C to H); (B) Additive effect size versus average epistatic effect size for each site (purple point). (C-E) Site fixation probability with epistasis versus without epistasis, under mass, half-sib and full-sib selection regimes after 100 generations. Points (sites) are colored by their additive effect sizes. Diagonal line indicates the unity line and X marks average fixation probabilities across sites. (F-H) as in C-E, but with points colored by their average epistatic effect.

regimes, etc.) to instead select for increased mean value of the trait. When the trait under selection was mean-based, and not variance-based, all selection regimes were equally rapid (Fig. 6B). Both mass and half-sib selection yielded close to their maximum selective response within 8 generations (Fig. 6D & F). This was in contrast to what we saw for a variance-based trait, in which mass selection did not yield nearly half of its selective response within 8 generations (Fig. 6C & E), while half-sib selection reached close to maximum variability. Therefore, the increased efficiency of family-based designs appears specific to selection that acts on the variability of a trait, and not its mean.

Discussion

We examined the efficiency of different artificial selection regimes at evolving variability of a trait. We modeled a trait with a polygenic architecture, including the possibility of epistatic interactions, as is typical of behaviors. Our simulations demonstrated that selection can indeed increase the variability of a polygenic trait (Fig. 1C), without impacting the mean (Fig. 6A,C,E). Family-based selection allowed for an efficient response of increasing variability in the focal trait. The parameters of our trait were based on locomotor handedness in *Drosophila melanogaster*, but should generalize to many traits, and our major findings are robust to large parameter changes (Fig. S4, Fig. S5, Fig. S6, Fig. S7). Mass selection did result in a strong selection response, but this response was slower, a difference with practical consequences for actual selection experiments. For instance, laboratory artificial selection in fruit flies over a year would result in 20 to 25 generations [38]; at this point family-based selection is still likely to have yielded a larger response than mass selection. An added practical advantage of family-based selection is that family-wise variability estimates can be obtained by screening a subset of the individuals in each family [45]. Therefore, a comparable strength of selection can be achieved by screening fewer animals than in mass selection.

Since estimating variance requires the measurement of multiple individuals, it makes sense that family-based selection will be more efficient at selecting for variability. Family-based selection measures the phenotypes of multiple, related individuals that are enriched for common variants, and propagates families as the unit of selection. Consistent with this reasoning, we found that family-based selection was faster than mass selection on our variance-based

Figure 6: **Efficacy of selection regimes for variance-based and mean-based traits**
Change in selected phenotype versus generations of selection for increased (A) variability of
the trait (same as Fig. 1C) and (B) mean of the trait. (C) Kernel density estimates of the
distributions of trait values at generations 0, 8 and 100 of mass selection for increased trait
variability. The generation 8 distribution is not as wide as the generation 100 distribution.
(D) As in C, but for selection for increased trait mean. Note that the generation 8 distribution
has a similar mean to that of generation 100. (E, F) As in C and D, but for half-sib selection.
In both panels, the generation 8 and 100 distributions are very similar.

trait (Fig. 6). When the same model was used to select for an increase in the mean value of the phenotype, all three regimes performed essentially identically (Fig. 6B,D,E). This indicated that family-based selection might be specifically effective when the selected trait reflects variability among individuals.

Population size plays a key role in selection, with larger populations being more resistant to drift, resulting in stronger selection response [21, 48]. Accordingly, we found larger population sizes resulted in faster, more reliable responses across all selection regimes (Fig. S7). Family-based selection schemes have smaller effective populations than mass selection [31, 46], and we found that their maximum selection responses were correspondingly sensitive to decreased total population size(Fig. S7). The relative size of families within the population also mattered for selective responses. Given a constant total population size, more, smaller families improved the reliability of selection across independent runs (Fig. 3B). Reliability is important when planning selection experiments in organisms with substantial generation times. Even fruit flies, with 10 day generations, require months or years to complete an artificial selection experiment. Unfortunately, while increasing the number of families improves reliability, it tends to diminish the magnitude of selection responses (Fig. 3A). Therefore, a population that has an intermediate number of moderate sized families represented a good compromise between maximizing selection response and minimizing run-to-run noise. Our model thus is helpful for planning experiments and predicting the population structure most likely to maximize the chances of seeing a selection response within experimental constraints.

Site fixation probabilities varied considerably across regimes (Fig. 2) and with the presence or absence of epistasis (Fig. 5). Even with a large total population size, family-based selection's reduced effective population size led to substantially lower heterozygosity after selection in our model (Fig. 2A). The effects of inbreeding were most prominent in full-sib selection. The absence of inbreeding in mass selection resulted in far fewer sites being fixed after 100 generations (Fig. 2A). Also, in mass selection the effect sizes of the sites were more strongly correlated with fixation probability. Sites of large effect had higher chances of fixing in all regimes, but sites of low effect were much less likely to fix in mass selection (Fig. 2), consistent with drift being a powerful dynamic in family-based selection. Therefore, while the population structure in family-based selection can accelerate the speed of selection, the loss of genetic diversity could potentially lead to inbreeding depression ([31]).

When epistatis was factored into the model, fixation probability rose (Fig. 5). Since

inbreeding and family structure can preserve larger regions of linked sites, they make it more likely that sites with neutral or mild direct effects but more positive or negative epistatic effects will fix (Fig. 5F-H). Conversely, sites of high direct effect can be prevented from fixing if they are epistatically coupled with sites that negatively impact the trait under selection. This was evident in our model, with sites of high additive effect size having a lower likelihood of fixation when epistasis was included, across all selection regimes (Fig. 5C-E). When planning actual selection experiments, the true level of epistasis may be unknown. Fortunately, our model suggests that the extent of epistasis does not change the relative efficacy of the three regimes (nor did it impact the final variability values attained after many generations of selection; Fig. 4).

Our model is flexible, accommodating populations of varying structures and traits having different underlying genetic architectures. We observed a robust response to selection for variability across a wide range of configurations with our model (Fig. S4 - Fig. S7). Our model highlights that selection can change the variance of traits, and that family-based regimes are an effective way of implementing directional selection for increased variability. Intragenotypic behavioral variability can be a target of selection, in the lab and potentially in nature. Responsiveness to selection is a precondition for bet-hedging to evolve as an adaptive phenotypic strategy. Experimental studies that implement selection on variability in the lab can are a promising path forward to understand the evolutionary role of variability and its genetic basis.

Acknowledgments

We thank Ian Dworkin for the conceptual understanding of family-based selection, Danylo Lavrentovich for assistance in using the FAS Research Computing Cluster, and Vikram Ravindranath for help with LaTeX typesetting.

References

[1] Julien F Ayroles et al. "Behavioral idiosyncrasy reveals genetic control of phenotypic variability". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 112.21 (May 2015), pp. 6706–6711.

[2] Hubertus J E Beaumont et al. "Experimental evolution of bet hedging". en. In: *Nature* 462.7269 (Nov. 2009), pp. 90–93.

[3] David Bierbach, Kate L Laskowski, and Max Wolf. "Behavioural individuality in clonal fish arises despite near-identical rearing conditions". en. In: *Nat. Commun.* 8 (May 2017), p. 15361.

[4] Peter A Biro and Judy A Stamps. "Are animal personality traits linked to life-history productivity?" en. In: *Trends Ecol. Evol.* 23.7 (July 2008), pp. 361–368.

[5] Benjamin de Bivort et al. "Precise quantification of behavioral individuality from 80 million decisions across 183,000 flies". en. In: *Front. Behav. Neurosci.* 16 (May 2022), p. 836626.

[6] Sean M Buchanan, Jamey S Kain, and Benjamin L de Bivort. "Neuronal control of locomotor handedness in Drosophila". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 112.21 (May 2015), pp. 6700–6705.

[7] Ben B Chapman et al. "Behavioural syndromes at multiple scales in Myrmica ants". In: *Anim. Behav.* 82.2 (Aug. 2011), pp. 391–397.

[8] Dylan Z Childs, C J E Metcalf, and Mark Rees. "Evolutionary bet-hedging in the real world: empirical evidence and challenges revealed by plants". en. In: *Proc. Biol. Sci.* 277.1697 (Oct. 2010), pp. 3055–3064.

[9] Matthew A Churgin et al. "Neural correlates of individual odor preference in Drosophila". en. In: *eLife* (Oct. 2023).

[10] Sasha R X Dall, Alasdair I Houston, and John M McNamara. "The behavioural ecology of personality: consistent individual differences from an adaptive perspective". en. In: *Ecol. Lett.* 7.8 (Aug. 2004), pp. 734–739.

[11] Vincent Debat, Allan Debelle, and Ian Dworkin. "Plasticity, canalization, and developmental stability of the Drosophila wing: joint effects of mutations and developmental temperature". en. In: *Evolution* 63.11 (Nov. 2009), pp. 2864–2876.

[12] Niels J Dingemanse and Max Wolf. "Recent models for adaptive personality differences: a review". en. In: *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365.1560 (Dec. 2010), pp. 3947–3958.

[13] Niels J Dingemanse et al. "Variation in personality and behavioural plasticity across four populations of the great tit Parus major". en. In: *J. Anim. Ecol.* 81.1 (Jan. 2012), pp. 116–126.

[14] Ian Dworkin and Greg Gibson. "Epidermal growth factor receptor and transforming growth factor-beta signaling contributes to variation for wing shape in Drosophila melanogaster". en. In: *Genetics* 173.3 (July 2006), pp. 1417–1431.

[15] Michael B Elowitz et al. "Stochastic gene expression in a single cell". en. In: *Science* 297.5584 (Aug. 2002), pp. 1183–1186.

[16] Julia Freund et al. "Emergence of individuality in genetically identical mice". en. In: *Science* 340.6133 (May 2013), pp. 756–759.

[17] Araceli Gallego and Carlos López-Fanjul. "The number of loci affecting a quantitative trait in Drosophila melanogaster revealed by artificial selection". In: *Genet. Res.* 42.2 (Oct. 1983), pp. 137–149.

[18] Charles R Harris et al. "Array programming with NumPy". en. In: *Nature* 585.7825 (Sept. 2020), pp. 357–362.

[19] Kyle S Honegger et al. "Idiosyncratic neural coding and neuromodulation of olfactory individuality in Drosophila". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 117.38 (Sept. 2020), pp. 23292–23297.

[20] Jamey S Kain, Chris Stokes, and Benjamin L de Bivort. "Phototactic personality in fruit flies and its suppression by serotonin and white". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 109.48 (Nov. 2012), pp. 19834–19839.

[21] M Kimura, T Maruyama, and J F Crow. "The mutation load in small populations". en. In: *Genetics* 48.10 (Oct. 1963), pp. 1303–1312.

[22] Raine Kortet and Ann Hedrick. "A behavioural syndrome in the field cricket Gryllus integer: intrasexual aggression is correlated with activity in a novel environment". en. In: *Biol. J. Linn. Soc. Lond.* 91.3 (July 2007), pp. 475–482.

[23]   Yanzhu Lin et al. "Microenvironmental Gene Expression Plasticity Among Individual Drosophila melanogaster". en. In: *G3* 6.12 (Dec. 2016), pp. 4197–4210.

[24]   Gerit Arne Linneweber et al. "A neurodevelopmental origin of behavioral individuality in the Drosophila visual system". en. In: *Science* 367.6482 (Mar. 2020), pp. 1112–1119.

[25]   John M McNamara, Zoltan Barta, and Alasdair I Houston. "Variation in behaviour promotes cooperation in the Prisoner's Dilemma game". en. In: *Nature* 428.6984 (Apr. 2004), pp. 745–748.

[26]   Jeroen Minderman et al. "Personality traits in wild starlings: exploration behavior and environmental sensitivity". In: *Behav. Ecol.* 20.4 (July 2009), pp. 830–837.

[27]   T Morgan. "Complete linkage in the second chromosome of the male of Drosophila". en. In: *Science* 36.934 (Nov. 1912), pp. 719–720.

[28]   Helen Olofsson, Jörgen Ripa, and Niclas Jonzén. "Bet-hedging as an evolutionary game: the trade-off between egg size and number". en. In: *Proc. Biol. Sci.* 276.1669 (Aug. 2009), pp. 2963–2969.

[29]   Carlos Pantoja et al. "Neuromodulatory Regulation of Behavioral Individuality in Zebrafish". en. In: *Neuron* 91.3 (Aug. 2016), pp. 587–601.

[30]   Juli G Pausas et al. "Bet-hedging and best-bet strategies shape seed dormancy". en. In: *New Phytol.* 236.4 (Nov. 2022), pp. 1232–1236.

[31]   Nina Pekkala et al. "The effect of inbreeding rate on fitness, inbreeding depression and heterosis over a range of inbreeding coefficients". en. In: *Evol. Appl.* 7.9 (Nov. 2014), pp. 1107–1119.

[32]   Alison Pischedda and William R Rice. "Partitioning sexual selection into its mating success and fertilization success components". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 109.6 (Feb. 2012), pp. 2049–2053.

[33]   Derek Roff and Daphne Fairbairn. "Modeling Experimental Evolution Using Individual-Based Variance-Componenets Models". In: *Experimental evolution: Concepts, methods, and applications of selection experiments.* Ed. by Theodore Garland and Michael R Rose. Berkeley, CA: University of California Press, Dec. 2009, pp. 31–64.

[34] Andrew Sih et al. "Behavioral syndromes: an intergrative overiew". en. In: *Q. Rev. Biol.* 79.3 (Sept. 2004), pp. 241–277.

[35] Andrew M Simons. "Fluctuating natural selection accounts for the evolution of diversification bet hedging". en. In: *Proc. Biol. Sci.* 276.1664 (June 2009), pp. 1987–1992.

[36] Ana Solopova et al. "Bet-hedging during bacterial diauxic shift". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 111.20 (May 2014), pp. 7427–7432.

[37] Judy A Stamps and Ton G G Groothuis. "Developmental perspectives on personality: implications for ecological and evolutionary studies of individual differences". en. In: *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365.1560 (Dec. 2010), pp. 4029–4041.

[38] Hugo Stocker and Peter Gallant. "Getting Started". In: *Drosophila: Methods and Protocols.* Ed. by Christian Dahmann. Totowa, NJ: Humana Press, 2008, pp. 27–44.

[39] M A Toro et al. "Inbreeding and family index selection for prolificacy in pigs". In: *Anim. Sci.* 46.1 (Feb. 1988), pp. 79–85.

[40] Thomas L Turner, Paige M Miller, and Veronica A Cochrane. "Combining genome-wide methods to investigate the genetic complexity of courtship song variation in Drosophila melanogaster". en. In: *Mol. Biol. Evol.* 30.9 (Sept. 2013), pp. 2113–2120.

[41] Thomas L Turner et al. "Population-based resequencing of experimentally evolved populations reveals the genetic basis of body size variation in Drosophila melanogaster". en. In: *PLoS Genet.* 7.3 (Mar. 2011), e1001336.

[42] Monica E M Verbeek, Piet J Drent, and Piet R Wiepkema. "Consistent individual differences in early exploratory behaviour of male great tits". In: *Anim. Behav.* 48.5 (Nov. 1994), pp. 1113–1121.

[43] Pauli Virtanen et al. "SciPy 1.0: fundamental algorithms for scientific computing in Python". en. In: *Nat. Methods* 17.3 (Mar. 2020), pp. 261–272.

[44] Günter Vogt et al. "Production of different phenotypes from the same genotype in the same environment by developmental variation". en. In: *J. Exp. Biol.* 211.Pt 4 (Feb. 2008), pp. 510–523.

[45] Bruce Walsh and Michael Lynch. "Family-Based Selection". In: *Evolution and Selection of Quantitative Traits*. Ed. by Bruce Walsh and Michael Lynch. Oxford: Oxford University Press, 2018, pp. 719–768.

[46] Bruce Walsh and Michael Lynch. "Long-term Response: 2. Finite Population Size and Mutation". In: *Evolution and Selection of Quantitative Traits*. Oxford University Press, 2018.

[47] Max Wolf, G Sander van Doorn, and Franz J Weissing. "Evolutionary emergence of responsive and unresponsive personalities". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 105.41 (Oct. 2008), pp. 15825–15830.

[48] S Wright. "Evolution in Mendelian Populations". en. In: *Genetics* 16.2 (Mar. 1931), pp. 97–159.

[49] Bas Zwaan, R Bijlsma, and R F Hoekstra. "Direct selection on life span in Drosophila melanogaster". en. In: *Evolution* 49.4 (Aug. 1995), pp. 649–659.

 # Supplementary Figures

Figure S1: **Distribution of Site Effect Sizes**

Probability density function of betaprime distribution [1] from which site effect sizes are drawn (black line); example values of 500 sites with effect sizes drawn from the distribution from one particular run of our model.

Figure S2: **Empirical turn bias variability scores**

Histogram of turn bias variability as measured across 167 inbred fruit fly lines [1]. We mapped the summed variant effect sizes (and epistatic effects, if they were included in a model run) of individuals in our simulation to this distribution to determine the latent turn bias variability from which their individual turn bias phenotypes were drawn.

Figure S3: **Comparison of mass selection methods**

(A) Average change in variability as a function of the number of generations of selection for different selection regimes, including mass selection implemented by keeping individuals with most extreme phenotypic values; (B) Average change in variability for the first 10 generations of selection

Figure S4: **Effect of number of sites on selection response**

Average change in variability as a function of the number of generations of selection with varying number of sites that contribute to the variability phenotype in (A) Mass Selection; (B) Half-Sib Selection and (C) Full-Sib Selection

Figure S5: **Effect of site effect size distribution on selection response**

Probability density functions (PDFs) of distributions and corresponding example values of effect sizes for sites that contribute to variability for (A) Betaprime distribution, (C) Exponential distribution, (E) Normal distribution and (G) Skewed-Normal distribution; (B,D,F,H) Corresponding change in variability (i.e., standard deviation) as a function of number of generations of selection for various selection regimes. $[(f(G_a) = G_a)$ for all cases]

Figure S6: **Effect of selection strength on selection response**

Average change in variability as a function of the number of generations of selection with varying percentage of individuals kept every generation to sire offspring in (A) Mass Selection; (B) Half-Sib Selection and (C) Full-Sib Selection

Figure S7: **Effect of population size on selection response**

Average change in variability as a function of the number of generations of selection with varying total population size in (A) Mass Selection; (B) Half-Sib Selection and (C) Full-Sib Selection

Figure S8: **Effect of epistasis on family-based selection response**

Average change in variability as a function of the number of generations of selection with varying contributions of epistatic interactions on variability phenotype in (A) Half-Sib Selection and (B) Full-Sib Selection