Predicting Biological Age Using an Accumulated Neurotoxicity Biomarker for Amyloid Beta Oligomers

Andrey V. Kuznetsov

Department of Mechanical and Aerospace Engineering, North Carolina State University,

Raleigh, NC 27695-7910, USA; e-mail: avkuznet@ncsu.edu

Abstract

This study proposes using accumulated neurotoxicity, defined as the time integral of Aβ oligomer concentration, as a biomarker for neuronal aging. A relationship between biological age and accumulated neurotoxicity is proposed. Numerical analysis guided the development of a new analytical solution linking the biological and calendar ages of neurons. The effects of Aβ monomer and oligomer half-lives—key indicators of proteolytic efficiency—on biological age are examined. Both constant and age-dependent (exponentially increasing) half-life scenarios are considered. The findings indicate that increasing the half-life of Aβ monomers and oligomers with age accelerates biological aging. Reducing Aβ monomer production is shown to slow biological aging, with a linear relationship established between these two quantities. Additionally, biological age is found to depend linearly on the half-deposition time of Aβ oligomers into senile plaques.

Keywords: biomarker of aging; neuron; Alzheimer's disease; Finke−Watzky model; senile plaques

Abbreviations

Aβ   amyloid beta

CV   control volume

F-W   Finke−Watzky

IDP   intrinsically disordered protein

1. Introduction

Aging is a universal process in complex multicellular organisms (hydra, which seemingly can renew its stem cells indefinitely, is a very simple organism). Understanding aging is, therefore, critically important. It seems logical to associate aging with irreversible changes that occur in an organism as its chronological age increases. However, the term "irreversible" should be used cautiously, as in thermodynamics it is used for a process in which total entropy generation, including the system and its surroundings, is positive. Although nearly all biological processes are thermodynamically irreversible, this does not imply a continuous increase in an organism's internal entropy. As open systems, living organisms maintain their low-entropy state by exchanging energy and increasing the entropy of their surroundings.

(Reed et al., 2024) proposed using the term resilience, which is defined as a system's ability to return to its original state after a perturbation. Here, resilience is interpreted as the capacity to recover from damage. However, the ability of a biological system to restore itself is not unlimited. Many types of damage cannot be fully repaired and accumulate over time, including epigenetic changes (Wang et al., 2022), errors in transcription and translation (Vermulst et al., 2015), damage caused by reactive oxygen species (Labunskyy and Gladyshev, 2013), (Giorgi et al., 2018), and various other forms of deterioration. Aging can therefore be understood as a consequence of the biological system's limited resilience.

Aging arises from the accumulation of damage across multiple biological levels, from molecules to entire organ systems, which the body cannot fully repair. Here, "damage" refers to detrimental changes associated with aging. However, it is important to note that some changes may aid in adaptation to aging, while others may be neutral and have no harmful effects on the organism (Gladyshev et al., 2021).

Studying interventions that may slow aging requires an assessment of how rapidly an individual is aging, which is determined by their biological age. Unlike chronological age, biological age varies among individuals due to differences in aging rates. Various aging clocks, which are computational models based on biomarkers of aging, are used to estimate biological age (Min et al., 2024). Examples include epigenetic clocks that analyze DNA methylation patterns, telomere clocks that measure telomere length, and epitranscriptomic clocks that track RNA methylation (Wang et al., 2023), (Han, 2024), (Crimmins et al., 2024).

The abundance of aging clocks reflects the complexity of the aging process, which involves numerous biological mechanisms. (Dormann and Lemke, 2024) proposed developing a novel aging clock based on the aggregation of intrinsically disordered proteins (IDPs) such as amyloid beta (Aβ), tau, α-synuclein, and TDP-43. IDPs have a high tendency to aggregate due to the lower free energy of their aggregated

state compared to their native, disordered form (Vetri and Foderà, 2015). The significance of an IDP-based aging clock lies in the fact that aging is the primary risk factor for neurodegenerative diseases, including Alzheimer's and Parkinson's, which are closely linked to IDP aggregation.

This paper proposes using the accumulated neurotoxicity of Aβ oligomers as a biomarker for the biological age of neural cells (hereafter referred to as biological age). Numerical simulations are conducted to explore the relationship between the proposed biomarker, calendar age, and other model parameters. A hypothetical 70-year timespan is considered.

A new analytical solution has been derived. Unlike the results in (Kuznetsov, 2024a), which provided analytical solutions only for the extreme cases of infinitely slow ($\theta_{1/2,B} \to \infty$) and infinitely fast ($\theta_{1/2,B} \to 0$) deposition rates of Aβ aggregates into senile plaques, this work presents an analytical solution for a finite, physiologically relevant half-deposition times of Aβ oligomers into senile plaques, $\theta_{1/2,B}$. Due to the non-linearity of the governing equations, the behavior of the analytical solution presented in this paper differs significantly from previously reported solutions. The sensitivity of the predicted biological age to model parameters characterizing the efficiency of protein degradation machinery and the Aβ monomer production rate is investigated.

2. Materials and models

2.1. Mathematical framework for Aβ peptide aggregation

As in (Kuznetsov, 2024a), (Kuznetsov, 2025) the aggregation of Aβ peptides is described using the Finke-Watzky (F-W) model, which simplifies the process into two pseudo-elementary steps: nucleation and autocatalytic growth. During nucleation, initial aggregates form at a continuous rate, while in the autocatalysis phase, pre-existing aggregates catalyze the conversion of monomers into additional aggregates (Morris et al., 2008), (Iashchishyn et al., 2017). These two reaction mechanisms are mathematically represented as follows:

$$A \xrightarrow{k_1} B \tag{1}$$

$$A + B \xrightarrow{k_2} 2B \tag{2}$$

In Eqs. (1) and (2), A represents a monomeric peptide, while B denotes an amyloid-converted peptide that remains free and has not yet been incorporated into a plaque. The kinetic constants k_1 and k_2 correspond to the nucleation and autocatalytic growth rates, respectively (Morris et al., 2008). Primary nucleation,

simulated by Eq. (1), involves only monomers, whereas secondary nucleation, described by Eq. (2), requires interactions between monomers and pre-existing free aggregates of the same peptide (Thacker et al., 2023).

When Aβ monomers diffuse rapidly, their concentration, C_A, can be considered nearly uniform within the control volume (CV) illustrated in Fig. 1. Under this assumption, time t serves as the only independent variable in the model. The dependent variables used in the analysis are summarized in Table 1, while the model parameters are summarized in Table 2.

By scaling governing equations, (Kuznetsov, 2024b) introduced the following dimensionless parameter:

$$\xi = \frac{q_{A,0}L}{D_A} t_f k_2 \tag{3}$$

Parameter ξ represents the ratio of the variation in Aβ monomer concentration across the CV to the average concentration of Aβ monomers within the CV at time t_f. As demonstrated in (Kuznetsov, 2024b), the lumped capacitance approximation, which assumes that Aβ concentrations depend only on time and not on location within the CV, is valid when the condition $\xi \ll 1$ holds. In Eq. (3), the diffusivity of Aβ monomers is denoted by D_A. Using a diffusivity value of 62.3 μm²/s (Waters, 2010) and the other parameter values given in Table 2, ξ is estimated to be 0.026. This confirms that the criterion $\xi \ll 1$ is satisfied, and the spatial variation of Aβ monomer concentration in the CV can be neglected.

The model proposed in (Kuznetsov, 2024a) is adopted. By applying the conservation of Aβ monomers within the CV depicted in Fig. 1, the following equation is obtained:

$$\frac{dC_A}{dt} = -k_1 C_A - k_2 C_A C_B - C_A \frac{\ln(2)}{T_{1/2,A}} + \frac{q_{A,0}}{L} \tag{4}$$

In Eq. (4), t is the time, which is also synonymous with calendar age. The first term on the right-hand side of Eq. (4) represents the conversion rate of Aβ monomers into aggregates driven by nucleation. The second term describes the conversion rate through autocatalytic growth. The third term accounts for the degradation of Aβ monomers, while the fourth term models the production of Aβ monomers on lipid membranes. It should be noted that $\frac{q_{A,0}}{L}$ represents the average volumetric generation rate of Aβ monomers within the CV.

By applying the conservation of free Aβ aggregates within the CV, the following equation is derived:

$$\frac{dC_B}{dt} = k_1 C_A + k_2 C_A C_B - C_B \frac{\ln(2)}{T_{1/2,B}} - C_B \frac{\ln(2)}{\theta_{1/2,B}} \tag{5}$$

In Eq. (5), the first two terms on the right-hand side mirror the first two terms of Eq. (4), but with opposite signs. The third term represents the degradation rate of free Aβ aggregates, while the fourth term simulates the deposition rate of these aggregates into senile plaques. The formation of Aβ plaques from adhesive Aβ fibrils is modeled similarly to colloidal suspension coagulation (Boltachev and Ivanov, 2020). In the model, free Aβ aggregates (B) are assumed to have a half-life of $\theta_{1/2,B}$. These aggregates deposit into Aβ plaques, and the concentration of the deposited aggregates is denoted as C_D.

$$\frac{dC_D}{dt} = C_B \frac{\ln(2)}{\theta_{1/2,B}} - C_D \frac{\ln(2)}{T_{1/2,D}} \tag{6}$$

In Eq. (6), the first term on the right-hand side corresponds to the fourth term in Eq. (5) with the opposite sign, while the second term represents the half-life of Aβ aggregates within the senile plaques.

The initial conditions are specified as follows:

At $t = 0$: $C_A = 0$, $C_B = 0$, $C_D = 0$ (7a,b,c)

The time-dependent behaviors of Aβ monomers, aggregates, and plaque half-lives were modeled using two scenarios. In the first scenario, the half-lives remain constant, independent of calendar age:

$$T_{1/2,A} = T_{1/2,A,0} \tag{8}$$

$$T_{1/2,B} = T_{1/2,B,0} \tag{9}$$

$$T_{1/2,D} = T_{1/2,D,0} \tag{10}$$

In the second scenario, the half-lives increase exponentially with calendar age:

$$T_{1/2,A} = T_{1/2,A,0} \exp(bt) \tag{11}$$

$$T_{1/2,B} = T_{1/2,B,0} \exp(bt) \tag{12}$$

$$T_{1/2,D} = T_{1/2,D,0} \exp(bt) \tag{13}$$

$b = 0.0183$ hour^{-1} represents a situation in which the half-lives increase 2.5-fold over a 50-year period (Patterson et al., 2015).

2.2. Analytical solution for the scenario with $T_{1/2,A} \to \infty$, $T_{1/2,B} \to \infty$, and $T_{1/2,D} \to \infty$

For the scenario with $T_{1/2,A} \to \infty$ and $T_{1/2,B} \to \infty$, the numerical solutions presented in Figs. 2 and 3 indicate that the concentrations of Aβ monomers and oligomers, C_A and C_B, respectively, approach asymptotic steady-state values over time. This section determines these asymptotic values. The solution derived here is labeled as "analytical" in the figures. The steady-state values of C_A and C_B are obtained from the following equations, derived from Eqs. (4) and (5):

$$-k_1 C_A - k_2 C_A C_B + \frac{q_{A,0}}{L} = 0 \tag{14}$$

$$k_1 C_A + k_2 C_A C_B - C_B \frac{\ln(2)}{\theta_{1/2,B}} = 0 \tag{15}$$

Adding Eqs. (14) and (15) yields the following equation:

$$\frac{q_{A,0}}{L} - C_B \frac{\ln(2)}{\theta_{1/2,B}} = 0 \tag{16}$$

Solving Eq. (16) for C_B results in the following:

$$C_B = \frac{q_{A,0} \theta_{1/2,B}}{L \ln(2)} \tag{17}$$

Substituting Eq. (17) into Eq. (14) and solving for C_A yields:

$$C_A = \frac{1}{L\left(\dfrac{k_1}{q_{A,0}} + k_2 \dfrac{\theta_{1/2,B}}{L\ln(2)}\right)} \tag{18}$$

In agreement with Eq. (17), Figs. 2b and 3b demonstrate that the asymptotic value of C_B obtained for $t \to \infty$ is affected by both $q_{A,0}$ and $\theta_{1/2,B}$. While the asymptotic value of C_A remains unaffected by $q_{A,0}$ for small values of k_1, it does depend on the value of $\theta_{1/2,B}$ (Figs. 2a and 3a). This result is consistent with Eq. (18). The numerical and analytical solutions show excellent agreement when time exceeds five years (Figs. 2 and 3).

For $T_{1/2,D} \to \infty$, Eq. (6) can be rewritten as:

$$\frac{dC_D}{dt} = C_B \frac{\ln(2)}{\theta_{1/2,B}}$$ (19)

Substituting Eq. (17) into Eq. (19) yields:

$$\frac{dC_D}{dt} = \frac{q_{A,0}}{L}$$ (20)

Integrating Eq. (20) with the initial condition (7c) gives:

$$C_D = \frac{q_{A,0}}{L}t$$ (21)

According to Eq. (21), C_D increases linearly with time, depending on $q_{A,0}$ (Fig. S1a) but remaining independent of $\theta_{1/2,B}$ (Fig. S2a). The numerical and analytical solutions are nearly identical (Figs. S1a and S2a).

An equation for the radius of senile plaques as a function of C_D is derived in Section S1 of the Supplemental Materials, using the approach developed in (Watzky et al., 2008). By substituting Eq. (21) into Eq. (S5), an analytical solution is obtained for the scenario with $T_{1/2,A} \to \infty$, $T_{1/2,B} \to \infty$, and $T_{1/2,D} \to \infty$:

$$r_{ABP} = \left(\frac{3}{2\pi} \frac{MW}{\rho} \frac{L^2}{a_{21}} q_{A,0}t \right)^{1/3}$$ (22)

According to Eq. (22), the plaque radius grows in direct proportion to the cube root of time. It depends on $q_{A,0}$ (Fig. S1b) but is independent of $\theta_{1/2,B}$ (Fig. S1b). The numerical and analytical solutions for the plaque radius show excellent agreement (Figs. S1b and S2b).

2.2. Analytical solution for the scenario with constant finite half-lives $T_{1/2,A}$, $T_{1/2,B}$, and $T_{1/2,D}$

Numerical solutions of Eqs. (4) and (5) with initial conditions (7a, b) for finite, time-independent values of $T_{1/2,A}$ and $T_{1/2,B}$, as shown in Figs. S3 and S4, indicate that C_A and C_B reach steady-state

concentrations over time. These asymptotic concentrations C_A and C_B can be determined by solving the following steady-state equations, derived from Eqs. (4) and (5):

$$-k_1 C_A - k_2 C_A C_B - C_A \frac{\ln(2)}{T_{1/2,A}} + \frac{q_{A,0}}{L} = 0 \tag{23}$$

$$k_1 C_A + k_2 C_A C_B - C_B \frac{\ln(2)}{T_{1/2,B}} - C_B \frac{\ln(2)}{\theta_{1/2,B}} = 0 \tag{24}$$

The analytical solution for this case, labeled as "full analytical" in the figures, is derived in Section S2 of the Supplemental Materials.

Table 1. List of dependent variables used in the model.

Symbol	Definition	Units
C_A	Concentration of Aβ monomers	μM
C_B	Concentration of free Aβ aggregates (such as oligomers and protofibrils)	μM
C_D	Concentration of Aβ aggregates incorporated into amyloid plaques	μM

Table 2. List of model parameters.

Symbol	Definition	Units	Value or range	Reference or estimation method	Value(s) used in computations
a_{21}	Factor that converts from $\frac{\text{mol}}{\mu\text{m}^3}$ to μM	$\frac{\mu\text{M } \mu\text{m}^3}{\text{mol}}$	10^{21}		10^{21}
b	The rate constant describing the exponential increase in the half-lives of	hour^{-1}		(Patterson et al., 2015)	0.0183 [a]

	Aβ monomers, aggregates, and plaques			
c	Proportionality coefficient in Eq. (28) that defines the relationship between biological age and the accumulated neurotoxicity of Aβ oligomers	$\dfrac{\text{year}}{\mu\text{M s}}$			1.44×10^{-8} [b]
k_1	Rate constant for the nucleation step of the F-W model for Aβ aggregation	s^{-1}	2.22×10^{-9} [c]	(Morris et al., 2008), (Bieschke et al., 2005)	2.22×10^{-9}
k_2	Rate constant for the autocatalytic growth step of the F-W model for Aβ aggregation	$\mu\text{M}^{-1}\ \text{s}^{-1}$	9.44×10^{-6} [c]	(Morris et al., 2008), (Bieschke et al., 2005)	9.44×10^{-6}
L	One-half of the inter-plaque spacing in neural tissue	µm	50	Estimated based on data presented in Fig. 1a of (Querol-Vilaseca et al., 2019)	50
MW	Molecular weight of the Aβ monomer, calculated as the total mass of atoms that compose the monomer	g mol^{-1}	4.51×10^{3} [d]	(Raskatov, 2019)	4.51×10^{3}

$q_{A,0}$	Aβ monomer flux per unit area at the left face of the CV. The CV, represented by the gray-shaded square in Fig. 1, has a volume of L^3. In the lumped capacitance approximation, $q_{A,0}/L$ gives the volumetric rate of monomer production per unit volume.	μM μm s⁻¹	1.1×10^{-5} [e]	(Raskatov, 2019)	1.1×10^{-5}
t_f	Representative lifespan	s	2.21×10^{9} [f]	(Hampel et al., 2021), (Dormann and Lemke, 2024)	2.21×10^{9}
$T_{1/2,A,0}$	Half-life of Aβ monomers at the beginning of the process, $t = 0$	s	4.61×10^{3} [g]	(Waters, 2010)	4.61×10^{3}, 10^{20} (to represent $T_{1/2,A,A\beta} \to \infty$)
$T_{1/2,B,0}$	Half-life of free Aβ aggregates at the beginning of the process, $t = 0$	s	4.61×10^{4} [h]		4.61×10^{4}, 10^{20} (to represent $T_{1/2,B,A\beta} \to \infty$)
$T_{1/2,D,0}$	Half-life of Aβ aggregates after their incorporation into senile plaques at $t = 0$	s	4.61×10^{5} [i]		4.61×10^{5}, 10^{20} (to represent $T_{1/2,D,A\beta} \to \infty$)

ρ	Density of the Aβ plaque	g/μm^3	1.35×10^{-12} [j]	(Cavicchi et al., 2018)	1.35×10^{-12}
$\theta_{1/2,B}$	Half-deposition time of free Aβ aggregates into senile plaques, defined as the time required for half of the free Aβ aggregates within the CV (Fig. 1) to be deposited into the senile plaque	s	8.64×10^{4} - 1.73×10^{5} [k]	(Han et al., 2017), (Meyer-Luehmann et al., 2008)	10^{7}

[a] $b=\ln(2.5)/50=0.0183$ hour^{-1} represents the case where $T_{1/2,A}$, $T_{1/2,B}$, and $T_{1/2,D}$ increase by a factor of 2.5 over the span of 50 years, as reported in (Patterson et al., 2015) and illustrated in Fig. S6.

[b] The value of c is determined from the assumption that the highest accumulated neurotoxicity value in Fig. 2a, 6.95×10^{9} μM·s, corresponds to a biological age of 100 years.

[c] The values reported in (Morris et al., 2008) were obtained by curve-fitting experimental data from (Bieschke et al., 2005). The specific parameter values are $k_1 =8\times10^{-6}$ h^{-1}, $k_2 =3.4\times10^{-2}$, as detailed in (Morris et al., 2008).

[d] The molecular weight of Aβ42, as is reported in (Raskatov, 2019).

[e] As noted in (Raskatov, 2019), the human brain synthesizes Aβ42 monomers at a rate of 0.33×10^{-6} μmol s^{-1}. Given the brain's estimated volume of 1.5×10^{15} μm³, the production rate per unit volume of Aβ42 is approximately 2.2×10^{-22} μmol μm^{-3} s^{-1}. In a small cube with side length L (representing the gray-shaded CV in Fig. 1), the total production rate of Aβ42 peptide is calculated to be 2.75×10^{-17} μmol s^{-1}. Aβ monomers are produced at lipid membranes (Dawkins et al., 2023). Assuming that all Aβ42 monomers are released from the left face of the cube, which has an area of L^2 (as shown in Fig. 1), the release rate per unit area is 1.1×10^{-20} μmol μm^{-2} s^{-1}, or equivalently 1.1×10^{-5} μM μm s^{-1}.

[f] Pathological processes involving Aβ may begin decades before clinical symptoms appear (Hampel et al., 2021). The formation of Aβ aggregates likely starts early in life (Dormann and Lemke, 2024), initially kept in check by proteolytic degradation (Saido and Leissring, 2012). This study hypothesizes that the Aβ aggregation process continues throughout an individual's lifetime (assumed to be 70 years in the model) and can be used as one of the biomarkers of biological aging (Dormann and Lemke, 2024).

[g] The half-life of Aβ monomers is estimated to be 1.28 hours (Waters, 2010).

[h] Free Aβ aggregates are expected to have a longer half-life than Aβ monomers, although they can still be degraded by microglia (Nakanishi, 2003). In this study, their half-life is assumed to be ten times that of Aβ monomers.

[i] In the absence of therapeutic intervention, Aβ plaques degrade at a significantly slower rate than Aβ monomers. (Madrasi et al., 2021) indicates that the turnover of senile plaques occurs over several years. To account for this, two values for $T_{1/2,D}$ are considered: one representing an infinite half-life of Aβ plaques, 10^{20} s, and another set to 4.61×10^5 s, reflecting a scenario where the half-life of Aβ plaques is ten times that of free Aβ aggregates.

[k] Research indicates that individual Aβ plaques can develop rapidly, with formation occurring within a span of 24 hours (Meyer-Luehmann et al., 2008) to 48 hours (Han et al., 2017).

2.2. Criterion of accumulated neurotoxicity of Aβ oligomers and proposed biomarker for biological aging

The precise mechanism underlying the neurotoxicity of Aβ oligomers remains unclear. Some Aβ oligomers may be inherently more toxic than others, or neurotoxicity could result from a heterogeneous mixture of oligomers with varying structures, stability, and concentrations. The mixture of oligomers may induce nonspecific neurotoxicity by interacting with membrane proteins, disrupting membrane lipids, triggering oxidative stress, and altering membrane dielectric properties and ion permeability (Benilova et al., 2012). If neurotoxicity results from the combined effects of various oligomers, a parameter quantifying the accumulated time-dependent damage caused by Aβ oligomers can be defined, following (Kuznetsov, 2024a), (Kuznetsov, 2024c), as:

$$\Xi(t) = \int_0^t C_B(\hat{t}) d\hat{t} \tag{25}$$

While a few drugs, such as aducanumab and lecanemab, have received FDA approval, numerous clinical trials of Aβ-clearing drugs for AD treatment have failed (Zhang et al., 2023), (Xiao and Zhang, 2024). This aligns with Eq. (25), which suggests that removing Aβ plaques does not undo the cumulative damage caused by Aβ oligomers.

By substituting the analytical solution for C_B from Eq. (17), which holds for $T_{1/2,A} \rightarrow \infty$ and $T_{1/2,B} \rightarrow \infty$, into Eq. (25), the following expression is derived:

$$\Xi(t) = \frac{q_{A,0}\theta_{1/2,B}}{L\ln(2)}t \tag{26}$$

The biological age of neurons (hereafter referred to as biological age) is defined by the following equation:

$$\text{biological age} = c\Xi \tag{27}$$

Substituting the analytical solution from Eq. (26) into Eq. (27) results in:

$$\text{biological age} = c\frac{q_{A,0}\theta_{1/2,B}}{L\ln(2)}t \tag{28}$$

2.3. Analysis of the sensitivity of biological age to various parameters

To assess the sensitivity of biological age to model parameters, local sensitivity coefficients were calculated. These coefficients represent the first-order partial derivatives of biological age with respect to the model parameters (Beck and Arnold, 1977), (Zadeh and Montas, 2010), (Zi, 2011), (Kuznetsov and Kuznetsov, 2019). The dimensionless relative sensitivity coefficients were determined following the methodology outlined in (Zadeh and Montas, 2010) and (Kacser et al., 1995). For instance, the dimensionless sensitivity of biological age to the flux of Aβ monomers into the CV is given by:

$$S_{q_{A,0}}^{\text{biological age}} = \frac{q_{A,0}}{\text{biological age}}\frac{\partial \text{biological age}}{\partial q_{A,0}} \tag{29}$$

Utilizing the analytical solution given by Eq. (28) yields the following expression for the sensitivity of biological age to calendar age t:

$$S_t^{\text{biological age}} = 1 \tag{30}$$

Eq. (30) is consistent with the fact that biological age increases in direct proportion to calendar age t; see Eq. (28). Also,

$$S_{q_{A,0}}^{\text{biological age}} = 1 \tag{31}$$

$$S_{\theta_{1/2,B}}^{\text{biological age}} = 1 \tag{32}$$

$S_{T_{1/2,A}}^{\text{biological age}}$ is calculated using the same parameter values as those used for one of the curves in Fig. 4a: $T_{1/2,A} = 4.61 \times 10^3$ s (the biologically relevant value) and $T_{1/2,B} = 10^{20}$ s. Utilizing Eq. (S16) yields the following result:

$$S_{T_{1/2,A}}^{\text{biological age}} = 1.25 \tag{33}$$

$S_{T_{1/2,B}}^{\text{biological age}}$ is calculated using the same parameter values as those used for one of the curves in Fig. 6a: $T_{1/2,A} = 10^{20}$ s and $T_{1/2,B} = 4.61 \times 10^4$ s (the biologically relevant value). Using Eq. (S17) yields the following result:

$$S_{T_{1/2,B}}^{\text{biological age}} = 0.995 \tag{34}$$

3. Results

Details of the numerical solution are outlined in Section S3 of the Supplementary Materials. Unless otherwise specified in a figure or its caption, parameter values are taken from Table 2.

Accumulated neurotoxicity, Ξ, quantifies the neuronal damage caused by Aβ oligomers. Therefore, it is proposed as a biomarker for neuronal biological age. Biological age is assumed to be directly proportional to accumulated neurotoxicity. The maximum accumulated neurotoxicity value, 6.95×10^9 μM·s in Fig. S5a, is assumed equivalent to a biological age of 100 years, as defined by Eq. (27).

If the proteolytic machinery responsible for Aβ monomer degradation remains functional ($T_{1/2,A} = 4.61 \times 10^3$ s, a physiologically relevant value), neuronal aging due to Aβ oligomer neurotoxicity remains minimal (Fig. 4a). This does not imply that neurons will not age, but rather that Aβ oligomer neurotoxicity will not be a significant contributing factor in aging. In contrast, if Aβ monomer degradation is entirely impaired ($T_{1/2,A} = 10^{20}$ s), the resulting neurotoxicity will drive neurons to a biological age of 100 years by a calendar age of 70 years. If the half-life of Aβ monomers is increased

tenfold beyond the physiologically relevant value ($T_{1/2,A} = 4.61 \times 10^4$ s), neurons will reach a biological age of approximately 47 years by the time the calendar age reaches 70 years (Fig. 4a). The observed increase in biological age with the half-life of Aβ monomers (Fig. 4a) is consistent with a positive value of $S_{T_{1/2,A}}^{\text{biological age}}$; see Eq. (33).

For $T_{1/2,A} = 4.61 \times 10^3$ s and $T_{1/2,A} = 10^{20}$ s, the numerical and full analytical solutions for the biological age (given by Eq. (S15)) show excellent agreement. However, for $T_{1/2,A} = 4.61 \times 10^4$ s, a systematic discrepancy between the two solutions persists over time (Fig. 4a). To understand the origin of this discrepancy, Fig. 5 plots the concentration of Aβ oligomers as a function of time for the scenario depicted in Fig. 4a. As seen in Fig. 5, the discrepancy arises from the difference between the numerical and full analytical solutions (given by Eq. (S10)) for C_B during the first 10 years. Although the agreement between the two solutions improves after this period, the initial discrepancy leads to a persistent offset in the accumulated neurotoxicity (Fig. S5a). This occurs because accumulated neurotoxicity is computed as the integral of the oligomer concentration over time (Eq. (25)), and during the first 10 years, the numerically predicted oligomer concentration is lower than the analytically predicted one (Fig. 5).

Fig. 4b depicts the scenario in which the half-life of Aβ monomers, $T_{1/2,A}$, increases exponentially with calendar age from its initial value, $T_{1/2,A,0}$, at t=0. The exponential growth of $T_{1/2,A} / T_{1/2,A,0}$, represented by $\exp(bt)$, is shown in Fig. S6, where the parameter b is estimated based on data from (Patterson et al., 2015). This model is plausible, as the Aβ monomer degradation machinery is expected to function efficiently in youth but gradually decline with age, leading to an age-dependent increase in the half-life of Aβ monomers. Under this scenario, if $T_{1/2,A,0}$ is set to 4.61×10^4 s, the biological age at a calendar age of 70 years reaches approximately 70 years (Fig. 4b), compared to about 47 years for the same $T_{1/2,A,0}$ without exponential growth over time (Fig. 4a).

When the half-life of Aβ oligomers is at a physiologically relevant value of 4.61×10^4 s, the degradation machinery functions efficiently, resulting in a very insignificant increase in biological age with calendar age (Fig. 6a). However, increasing the half-life to 4.61×10^6 s accelerates aging, raising the biological age to approximately 32 years by the time the calendar age reaches 70 years. If the half-life becomes infinitely large, the biological age reaches 100 years at a calendar age of 70 years (Fig. 6a). The increase of the biological age with $T_{1/2,B}$ observed in Fig. 6a agrees with the positive sensitivity $S_{T_{1/2,B}}^{\text{biological age}}$; see Eq. (34).

If the half-life of Aβ oligomers increases exponentially with age, an initial value of 4.61×10^6 s results in a biological age of approximately 47 years at a calendar age of 70 years (Fig. 6b), compared to 32 years for the same value of $T_{1/2,B,0}$ in Fig. 6a. The trends in accumulated neurotoxicity (Fig. S7) closely follow those in Fig. 6 for biological age, as the model assumes a direct proportionality between biological age and accumulated neurotoxicity; see Eq. (27).

Variations in the half-life of Aβ aggregates within senile plaques do not affect biological age (Fig. S8a,b). This is because the model assumes that accumulated neurotoxicity is driven by Aβ oligomers, whereas senile plaques, considered non-toxic, serve as relatively benign reservoirs for these oligomers—a hypothesis consistent with (Gouras et al., 2015), (Cline et al., 2018).

Fig. 7a illustrates the effect of Aβ monomer flux into the CV, $q_{A,0}$, representing the production rate of Aβ monomers, on biological age. Aβ monomers are generated at lipid membranes, and an increase in their flux significantly accelerates biological aging. When the flux is reduced to half of its physiologically relevant value ($0.5 \times 1.1 \times 10^{-5}$ μM μm s^{-1}), the biological age at a calendar age of 70 years is approximately 50 years (Fig. 7a). At a physiologically relevant value of the flux (1.1×10^{-5} μM μm s^{-1}), the biological age at 70 years reaches 100 years. If the flux is doubled from its physiologically relevant value ($2 \times 1.1 \times 10^{-5}$ μM μm s^{-1}), the biological age reaches 100 years as early as a calendar age of 35 years (Fig. 7a).

Fig. 7b illustrates the effect of the half-deposition time of free Aβ aggregates into senile plaques, $\theta_{1/2,B}$, on biological age. Interestingly, its effect is analogous to that of the Aβ monomer flux: when $\theta_{1/2,B}$ is reduced to half of its physiologically relevant value, the biological age at a calendar age of 70 years is approximately 50 years (Fig. 7b). At the physiologically relevant value of $\theta_{1/2,B}$, the biological age at 70 years reaches 100 years. When $\theta_{1/2,B}$ is doubled from its physiologically relevant value, the biological age reaches 100 years as early as a calendar age of 35 years (Fig. 7b).

Figs. 7a and 7b demonstrate excellent agreement between the numerical and analytical solutions; hence, the analytical solution given by Eq. (28) can be used to explain this similarity. Eq. (28) shows that biological age is proportional to the product of $q_{A,0}$ and $\theta_{1/2,B}$. Consequently, variations in these parameters produce identical effects on biological age. The effects of $q_{A,0}$ and $\theta_{1/2,B}$ on accumulated neurotoxicity follow the same trends, as biological age is assumed to be directly proportional to accumulated neurotoxicity (Figs. S9a,b).

Biological age exhibits minimal dependence on the nucleation rate constant, k_1 (Fig. 8a). However, it increases with an increase in the autocatalytic growth rate constant, k_2, indicating that Aβ oligomer production is primarily driven by the autocatalytic process (Fig. 8b). For biologically relevant values of k_2 (represented by the line with red crosses), the relationship between biological age and calendar age is initially nonlinear for the first 30 years. Beyond this period, it transitions into a linear trend, consistent with Eq. (28). The curves depicting accumulated neurotoxicity vs time for different values of k_1 and k_2 exhibit similar trends (Fig. S10a and S10b).

4. Discussion, limitations of the model, and future directions

The biological age of neurons is assumed to be characterized by the accumulated neurotoxicity of Aβ oligomers. The half-lives of Aβ monomers, oligomers, and aggregates serve as indicators of the efficiency of the Aβ degradation machinery. When these half-lives are assumed to be constant, biological age exhibits a linear dependence on calendar age (Figs. 4a, 6a, 7a, and 7b). This aligns with the analytical solution for biological age given by Eq. (28), which predicts a linear relationship with calendar age t. An exception is observed in Fig. 8b, which depicts biological age as a function of time for different values of the autocatalytic constant k_2. For biologically relevant values of k_2, the curve exhibits a parabolic shape at small t but transitions to a linear trend at larger times. This remains consistent with Eq. (28), as the analytical solution applies to sufficiently large time values where linearity emerges.

If the half-lives of monomers and oligomers are assumed to increase exponentially with age, simulating the decline of proteolytic machinery over time, the relationship between biological age and calendar age becomes nonlinear for intermediate initial half-life values (Figs. 4b and 6b).

Numerical analysis revealed a similarity between the effects of Aβ monomer production rate (represented by Aβ monomer flux into the CV, $q_{A,0}$) and the half-deposition time of free Aβ aggregates into senile plaques, $\theta_{1/2,B}$ (Fig. 7a and 7b). This similarity is explained by the analytical solution for biological age, which depends on the product $q_{A,0}\theta_{1/2,B}$.

A potential limitation of the proposed biomarker is the assumption that accumulated neurotoxicity begins to rise immediately after birth, whereas it is more likely to increase only after a certain age. Future research should focus on experimental evaluation of the proposed biomarker, which represents accumulated neurotoxicity, for assessing neuronal aging. Additionally, the potential applicability of

accumulated toxicity (defined similarly) as a biomarker for aging in non-neural tissues should be explored.

Data accessibility. This article has no additional data.

Authors' contributions. AVK is the sole author of this paper.

Competing interests. The author declares no competing interests.

Funding statement. The author acknowledges the support provided by the National Science Foundation (grant CBET-2042834) and the Alexander von Humboldt Foundation through the Humboldt Research Award.

References

Beck, J.V., Arnold, K.J., 1977. Parameter Estimation in Science and Engineering. Wiley, New York.

Benilova, I., Karran, E., Strooper, B.D., 2012. The toxic Aβ oligomer and Alzheimer's disease: an emperor in need of clothes. Nature Neuroscience 15, 349–357. https://doi.org/10.1038/nn.3028

Bieschke, J., Zhang, Q., Powers, E.T., Lerner, R.A., Kelly, J.W., 2005. Oxidative metabolites accelerate Alzheimer's amyloidogenesis by a two-step mechanism, eliminating the requirement for nucleation. Biochemistry 44, 4977–4983. https://doi.org/10.1021/bi0501030

Boltachev, G.S., Ivanov, M.G., 2020. Effect of nanoparticle concentration on coagulation rate of colloidal suspensions. Heliyon 6, e03295. https://doi.org/10.1016/j.heliyon.2020.e03295

Cavicchi, R.E., King, J., Ripple, D.C., 2018. Measurement of Average Aggregate Density by Sedimentation and Brownian Motion Analysis. Journal of Pharmaceutical Sciences 107, 1304–1312. https://doi.org/10.1016/j.xphs.2018.01.013

Cline, E.N., Bicca, M.A., Viola, K.L., Klein, W.L., 2018. The Amyloid-β Oligomer Hypothesis: Beginning of the Third Decade. J Alzheimer's Dis 64, S567–S610. https://doi.org/10.3233/JAD-179941

Crimmins, E.M., Klopack, E.T., Kim, J.K., 2024. Generations of epigenetic clocks and their links to socioeconomic status in the Health and Retirement Study. Epigenomics 16, 1031–1042. https://doi.org/10.1080/17501911.2024.2373682

Dawkins, E., Derks, R.J.E., Schifferer, M., Trambauer, J., Winkler, E., Simons, M., Paquet, D., Giera, M., Kamp, F., Steiner, H., 2023. Membrane lipid remodeling modulates γ-secretase processivity. Journal of Biological Chemistry 299, 103027. https://doi.org/10.1016/j.jbc.2023.103027

Dormann, D., Lemke, E.A., 2024. Adding intrinsically disordered proteins to biological ageing clocks. Nature Cell Biology 26, 851–858. https://doi.org/10.1038/s41556-024-01423-w

Giorgi, C., Marchi, S., Simoes, I.C.M., Ren, Z., Morciano, G., Perrone, M., Patalas-Krawczyk, P., Borchard, S., Jędrak, P., Pierzynowska, K., Szymański, J., Wang, D.Q., Portincasa, P., Węgrzyn, G., Zischka, H., Dobrzyn, P., Bonora, M., Duszynski, J., Rimessi, A., Karkucinska-Wieckowska, A., Dobrzyn, A., Szabadkai, G., Zavan, B., Oliveira, P.J., Sardao, V.A., Pinton, P., Wieckowski, M.R., 2018. Mitochondria and Reactive Oxygen Species in Aging and Age-Related Diseases. Int Rev Cell Mol Biol 340, 209–344. https://doi.org/10.1016/bs.ircmb.2018.05.006

Gladyshev, V.N., Kritchevsky, S.B., Clarke, S.G., Cuervo, A.M., Fiehn, O., de Magalhães, J.P., Mau, T., Maes, M., Moritz, R.L., Niederhofer, L.J., Van Schaftingen, E., Tranah, G.J., Walsh, K., Yura, Y., Zhang, B., Cummings, S.R., 2021. Molecular damage in aging. Nature Aging 1, 1096–1106. https://doi.org/10.1038/s43587-021-00150-3

Gouras, G.K., Olsson, T.T., Hansson, O., 2015. β-Amyloid peptides and amyloid plaques in Alzheimer's disease. Neurotherapeutics 12, 3–11. https://doi.org/10.1007/s13311-014-0313-y

Hampel, H., Hardy, J., Blennow, K., Chen, C., Perry, G., Kim, S.H., Villemagne, V.L., Aisen, P., Vendruscolo, M., Iwatsubo, T., Masters, C.L., Cho, M., Lannfelt, L., Cummings, J.L., Vergallo, A., 2021. The Amyloid-β Pathway in Alzheimer's Disease. Molecular Psychiatry 26, 5481–5503. https://doi.org/10.1038/s41380-021-01249-0

Han, J.-D.J., 2024. The ticking of aging clocks. Trends in Endocrinology & Metabolism 35, 11–22. https://doi.org/10.1016/j.tem.2023.09.007

Han, S., Kollmer, M., Markx, D., Claus, S., Walther, P., Fändrich, M., 2017. Amyloid plaque structure and cell surface interactions of β-amyloid fibrils revealed by electron tomography. Scientific Reports 7, 43577. https://doi.org/10.1038/srep43577

Iashchishyn, I.A., Sulskis, D., Ngoc, M.N., Smirnovas, V., Morozova-Roche, L.A., 2017. Finke-Watzky Two-Step Nucleation-Autocatalysis Model of S100A9 Amyloid Formation: Protein Misfolding as "Nucleation" Event. ACS Chemical Neuroscience 8, 2152–2158. https://doi.org/10.1021/acschemneuro.7b00251

Ikonomovic, M.D., Klunk, W.E., Abrahamson, E.E., Mathis, C.A., Price, J.C., Tsopelas, N.D., Lopresti, B.J., Ziolko, S., Bi, W., Paljug, W.R., Debnath, M.L., Hope, C.E., Isanski, B.A., Hamilton, R.L., DeKosky, S.T., 2008. Post-mortem correlates of in vivo PiB-PET amyloid imaging in a typical case of Alzheimer's disease. Brain 131, 1630–1645. https://doi.org/10.1093/brain/awn016

Kacser, H., Burns, J.A., Fell, D.A., 1995. The Control of Flux. Biochemical Society transactions 23, 341–366. https://doi.org/10.1042/bst0230341

Kuznetsov, A.V., 2025. Evaluating the Combined Neurotoxicity of Amyloid Beta and Tau Oligomers in Alzheimer's Disease: A Novel Cellular-Level Criterion. ASME Journal of Biomechanical Engineering 147, 041003. https://doi.org/10.1115/1.4067701

Kuznetsov, A.V., 2024a. A criterion characterizing accumulated neurotoxicity of Aβ oligomers in Alzheimer's disease. bioRxiv 2024.08.19.608707. https://doi.org/10.1101/2024.08.19.608707

Kuznetsov, A.V., 2024b. Lewy body radius growth: The hypothesis of the cube root of time dependency. J Theor Biol. 581, 111734. https://doi.org/10.1016/j.jtbi.2024.111734

Kuznetsov, A.V., 2024c. Criterion for assessing accumulated neurotoxicity of alpha-synuclein oligomers in Parkinson's disease. bioRxiv 2024.09.23.614584. https://doi.org/10.1101/2024.09.23.614584

Kuznetsov, I.A., Kuznetsov, A.V., 2019. Investigating sensitivity coefficients characterizing the response of a model of tau protein transport in an axon to model parameters. Computer Methods in Biomechanics and Biomedical Engineering 22, 71–83. https://doi.org/10.1080/10255842.2018.1534233

Labunskyy, V.M., Gladyshev, V.N., 2013. Role of reactive oxygen species-mediated signaling in aging. Antioxid Redox Signal 19, 1362–1372. https://doi.org/10.1089/ars.2012.4891

Liu, P., Reichl, J.H., Rao, E.R., McNellis, B.M., Huang, E.S., Hemmy, L.S., Forster, C.L., Kuskowski, M.A., Borchelt, D.R., Vassar, R., Ashe, K.H., Zahs, K.R., 2017. Quantitative Comparison of Dense-Core Amyloid Plaque Accumulation in Amyloid-β Protein Precursor Transgenic Mice. J Alzheimers Dis 56, 743–761. https://doi.org/10.3233/JAD-161027

Madrasi, K., Das, R., Mohmmadabdul, H., Lin, L., Hyman, B.T., Lauffenburger, D.A., Albers, M.W., Rissman, R.A., Burke, J.M., Apgar, J.F., Wille, L., Gruenbaum, L., Hua, F., 2021. Systematic in silico analysis of clinically tested drugs for reducing amyloid-beta plaque accumulation in Alzheimer's disease. Alzheimer's & Dementia 17, 1487–1498. https://doi.org/10.1002/alz.12312

Meyer-Luehmann, M., Spires-Jones, T.L., Prada, C., Garcia-Alloza, M., Calignon, A. de, Rozkalne, A., Koenigsknecht-Talboo, J., Holtzman, D.M., Bacskai, B.J., Hyman, B.T., 2008. Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease. Nature 451, 720–724. https://doi.org/10.1038/nature06616

Min, M., Egli, C., Dulai, A.S., Sivamani, R.K., 2024. Critical review of aging clocks and factors that may influence the pace of aging. Front Aging 5, 1487260. https://doi.org/10.3389/fragi.2024.1487260

Morris, A.M., Watzky, M.A., Agar, J.N., Finke, R.G., 2008. Fitting neurological protein aggregation kinetic data via a 2-step, Minimal/"Ockham's Razor" model: The Finke-Watzky mechanism of nucleation followed by autocatalytic surface growth. Biochemistry 47, 2413–2427. https://doi.org/10.1021/bi701899y

Nakanishi, H., 2003. Microglial functions and proteases. Molecular Neurobiology 27, 163–176. https://doi.org/10.1385/MN:27:2:163

Patterson, B.W., Elbert, D.L., Mawuenyega, K.G., Kasten, T., Ovod, V., Ma, S., Xiong, C., Chott, R., Yarasheski, K., Sigurdson, W., Zhang, L., Goate, A., Benzinger, T., Morris, J.C., Holtzman, D., Bateman, R.J., 2015. Age and amyloid effects on human central nervous system amyloid-beta kinetics. Ann Neurol 78, 439–453. https://doi.org/10.1002/ana.24454

Querol-Vilaseca, M., Colom-Cadena, M., Pegueroles, J., Nuñez-Llaves, R., Luque-Cabecerans, J., Muñoz-Llahuna, L., Andilla, J., Belbin, O., Spires-Jones, T., Gelpi, E., Clarimon, J., Loza-Alvarez, P., Fortea, J., Lleó, A., 2019. Nanoscale structure of amyloid-β plaques in Alzheimer's disease. Scientific Reports 9, 5181. https://doi.org/10.1038/s41598-019-41443-3

Raskatov, J.A., 2019. What Is the "Relevant" Amyloid β42 Concentration? ChemBioChem 20, 1725–1726. https://doi.org/10.1002/cbic.201900097

Reed, J.M., Wolfe, B.E., Romero, L.M., 2024. Is resilience a unifying concept for the biological sciences? iScience 27, 109478. https://doi.org/10.1016/j.isci.2024.109478

Saido, T., Leissring, M.A., 2012. Proteolytic degradation of amyloid β-protein. Cold Spring Harbor Perspectives in Medicine 2, a006379. https://doi.org/10.1101/cshperspect.a006379

Thacker, D., Barghouth, M., Bless, M., Zhang, E., Linse, S., 2023. Direct observation of secondary nucleation along the fibril surface of the amyloid β 42 peptide. Proceedings of the National Academy of Sciences 120, e2220664120. https://doi.org/10.1073/pnas.2220664120

Vermulst, M., Denney, A.S., Lang, M.J., Hung, C.-W., Moore, S., Moseley, M.A., Thompson, J.W., Madden, V., Gauer, J., Wolfe, K.J., Summers, D.W., Schleit, J., Sutphin, G.L., Haroon, S., Holczbauer, A., Caine, J., Jorgenson, J., Cyr, D., Kaeberlein, M., Strathern, J.N., Duncan, M.C., Erie, D.A., 2015. Transcription errors induce proteotoxic stress and shorten cellular lifespan. Nat Commun 6, 8065. https://doi.org/10.1038/ncomms9065

Vetri, V., Foderà, V., 2015. The route to protein aggregate superstructures: Particulates and amyloid-like spherulites. FEBS Letters 589, 2448–2463. https://doi.org/10.1016/j.febslet.2015.07.006

Wang, K., Liu, H., Hu, Q., Wang, L., Liu, J., Zheng, Z., Zhang, W., Ren, J., Zhu, F., Liu, G.-H., 2022. Epigenetic regulation of aging: implications for interventions of aging and diseases. Signal Transduction and Targeted Therapy 7, 374. https://doi.org/10.1038/s41392-022-01211-8

Wang, Qingyi, Hou, T., Wang, Qiwen, He, J., Wang, L., Si, J., Chen, S., 2023. An evaluation of aging measures: from biomarkers to clocks. Biogerontology 24, 303–328. https://doi.org/10.1007/s10522-022-09997-4

Waters, J., 2010. The Concentration of Soluble Extracellular Amyloid-beta Protein in Acute Brain Slices from CRND8 Mice. PLoS One 5, e15709. https://doi.org/10.1371/journal.pone.0015709

Watzky, M.A., Finney, E.E., Finke, R.G., 2008. Transition-Metal Nanocluster Size vs Formation Time and the Catalytically Effective Nucleus Number: A Mechanism-Based Treatment. Journal of the American Chemical Society 130, 11959–11969. https://doi.org/10.1021/ja8017412

Xiao, D., Zhang, C., 2024. Current therapeutics for Alzheimer's disease and clinical trials. Exploration of Neuroscience 3, 255–271. https://doi.org/10.37349/en.2024.00048

Zadeh, K.S., Montas, H.J., 2010. A class of exact solutions for biomacromolecule diffusion-reaction in live cells. Journal of Theoretical Biology 264, 914–933. https://doi.org/10.1016/j.jtbi.2010.03.028

Zhang, Y., Chen, H., Li, R., Sterling, K., Song, W., 2023. Amyloid β-based therapy for Alzheimer's disease: challenges, successes and future. Signal Transduction and Targeted Therapy 8, 248. https://doi.org/10.1038/s41392-023-01484-7

Zi, Z., 2011. Sensitivity analysis approaches applied to systems biology models. IET Systems Biology 5, 336–346. https://doi.org/10.1049/iet-syb.2011.0015

Figure captions

Fig. 1. A 2D view of two cubic control volumes (CVs) with side length L used to model the aggregation of Aβ monomers into free aggregates and their eventual deposition into senile plaques (Liu et al., 2017), (Ikonomovic et al., 2008). Aβ monomers are assumed to be generated at lipid membranes. In the shaded CV, the production occurs at the left face ($x=0$), which has an area of L^2. The right boundary of the shaded CV, $x=L$, is treated as symmetric, preventing Aβ monomer flux across it. A senile plaque is assumed to form at the interface between two adjacent CVs at $x=L$. A lumped capacitance approximation is utilized, simplifying the governing equations by assuming that all Aβ concentrations depend only on time.

Fig. 2. Molar concentration of (a) Aβ monomers, C_A, and (b) Aβ oligomers, C_B, as a function of time for different values of Aβ monomer flux into the CV, $q_{A,0}$. Parameters: $T_{1/2,A}=10^{20}$ s, $T_{1/2,B}=10^{20}$ s. The plotted analytical solutions are derived from Eqs. (17) and (18). Note that in each plotted case C_A and C_B reach constant asymptotic values as time increases. This is essential for deriving the analytical solution.

Fig. 3. Molar concentration of (a) Aβ monomers, C_A, and (b) Aβ oligomers, C_B, as a function of time for different values of half-deposition times of free Aβ aggregates into senile plaques, $\theta_{1/2,B}$. Parameters: $T_{1/2,A}=10^{20}$ s, $T_{1/2,B}=10^{20}$ s. The plotted analytical solutions are derived from Eqs. (17) and (18). Note that in each plotted case C_A and C_B reach constant asymptotic values as time increases. This is essential for deriving the analytical solution.

Fig. 4. Biological age as a function of calendar age for different half-lives of Aβ monomers, $T_{1/2,A}$. (a) The half-life of Aβ monomers, $T_{1/2,A}$, remains constant and independent of calendar age. (b) The half-life of Aβ monomers, $T_{1/2,A}$, increases exponentially with calendar age. Parameters: $T_{1/2,B}=10^{20}$ s.

Fig. 5. Molar concentration of Aβ oligomers, C_B, as a function of time for different half-lives of Aβ monomers, $T_{1/2,A}$. This figure corresponds to the scenario shown in Fig. 4 and uses the same parameter values. The full analytical solution, obtained from Eq. (S10), is plotted. Notably, a significant discrepancy between the numerical and full analytical solutions is observed during the first 10 years for the case where

$T_{1/2,A,0} = 4.61 \times 10^4$ s. However, in all cases, C_B eventually reaches a constant asymptotic value over time. Parameters: $T_{1/2,A} = 10^{20}$ s.

Fig. 6. Biological age as a function of calendar age. (a) The half-life of free Aβ aggregates, $T_{1/2,B}$, remains constant over time. (b) The half-life of free Aβ aggregates, $T_{1/2,B}$, increases exponentially with calendar age. Parameters: $T_{1/2,A} = 10^{20}$ s.

Fig. 7. (a) Biological age as a function of calendar age for different values of the flux, $q_{A,0}$, of Aβ monomers per unit area at the left-hand boundary of the CV shown in Fig. 1. (b) Biological age as a function of calendar age for different values of the half-deposition time, $\theta_{1/2,B}$, of free Aβ aggregates into senile plaques. Parameters: $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s.

Fig. 8. (a) Biological age as a function of calendar age for different values of the nucleation rate constant, k_1, in the first pseudo-elementary reaction step of the F-W model for Aβ peptide. Parameters: $k_2 = 9.44 \times 10^{-6}$ μM⁻¹ s⁻¹, $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s, $T_{1/2,D} = 10^{20}$ s. (b) Biological age as a function of calendar age for different values of the autocatalytic growth rate constant, k_2, in the second pseudo-elementary reaction step of the F-W model for Aβ peptide. Parameters: $k_1 = 2.22 \times 10^{-9}$ s⁻¹, $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s, $T_{1/2,D} = 10^{20}$ s.

Figure 1

(a)

(b)

Figure 2a,b

Figure 3a,b

Figure 4a,b

Figure 5

Figure 6a,b

Figure 7a,b

Figure 8a,b

Predicting Biological Age Using an Accumulated Neurotoxicity Biomarker for Amyloid Beta Oligomers

Andrey V Kuznetsov

Department of Mechanical and Aerospace Engineering, North Carolina State University,

Raleigh, NC 27695-7910, USA; e-mail: avkuznet@ncsu.edu

Supplemental Materials

S1. Modeling senile plaque growth

This study follows the derivation developed in (Kuznetsov, 2024a), (Kuznetsov, 2024c). To simulate the growth of an Aβ plaque (illustrated in Fig. 1), the approach focuses on determining the total number of Aβ monomers, denoted as N, that accumulate within the plaque over time t. This is done by considering the average concentration of Aβ monomers deposited into the plaque, as proposed by (Watzky et al., 2008):

$$N(t) = 2N_A \frac{L^3}{a_{21}} C_D(t) \tag{S1}$$

where N_A is Avogadro's number. The factor of 2 on the right-hand side of Eq. (S1) accounts for the assumption that an Aβ plaque forms in the space between two adjacent neurons. Each neuron's membrane releases Aβ monomers at a rate of $q_{A,0}$, contributing symmetrically to plaque formation (Fig. 1).

An alternative approach, also based on (Watzky et al., 2008), involves determining $N(t)$ using the volume of an individual inclusion body, denoted as V_{ABP}:

$$N(t) = N_A \frac{V_{ABP}(t)\rho}{MW} \tag{S2}$$

where MW represents the average molecular weight of an Aβ monomer.

Equating the right-hand sides of Eqs. (S1) and (S2) and solving for the volume of the Aβ plaque yields the following expression:

$$V_{ABP}(t) = 2\frac{MW}{\rho} \frac{L^3}{a_{21}} C_D(t) \tag{S3}$$

where $a_{21} = 10^{21} \dfrac{\mu M \; \mu m^3}{mol}$ represents the conversion factor from $\dfrac{mol}{\mu m^3}$ to μM.

Assuming a spherical shape for the Aβ plaque, its volume can be expressed as:

$$V_{ABP}(t) = \frac{4\pi}{3} r_{ABP}^3(t) \tag{S4}$$

Equating the right-hand sides of Eqs. (S3) and (S4) and solving for the radius of the Aβ plaque yields the following expression:

$$r_{ABP} = \left(\frac{3}{2\pi} \frac{MW}{\rho} \frac{L^3}{a_{21}} C_D(t) \right)^{1/3} \tag{S5}$$

The size of an Aβ plaque is often quantified by its surface area, which can be measured from stained brain tissue sections examined under a microscope. Within the framework of this model, the plaque area can be calculated as follows:

$$A_{ABP} = \pi r_{ABP}^2 \tag{S6}$$

Aβ plaque load, defined as the percentage of the area occupied by amyloid plaques (Liu et al., 2017), (Ikonomovic et al., 2008), can be determined using the following calculation:

$$ABP\,load = \frac{A_{ABP}}{2L^2} 100\% \tag{S7}$$

The factor of 2 in the denominator of Eq. (S7) accounts for the assumption in the model that a single plaque occupies two control volumes (Fig. 1).

S2. Derivation of the "full analytical" solution for the scenario with constant, finite half-lives $T_{1/2,A}$, $T_{1/2,B}$, and $T_{1/2,D}$

Summing Eqs. (23) and (24) gives:

$$C_A \frac{\ln(2)}{T_{1/2,A}} + C_B \ln(2) \left(\frac{1}{T_{1/2,B}} + \frac{1}{\theta_{1/2,B}} \right) - \frac{q_{A,0}}{L} = 0 \tag{S8}$$

Solving for C_A yields:

$$C_A = -C_B T_{1/2,A}\left(\frac{1}{T_{1/2,B}} + \frac{1}{\theta_{1/2,B}}\right) + \frac{q_{A,0}}{L}\frac{T_{1/2,A}}{\ln(2)} \tag{S9}$$

Substituting Eq. (S9) into Eq. (S8) results in a quadratic equation for C_B. The physically meaningful root, as concentration cannot be negative, is given by:

$$C_{B,fa} = \frac{1}{2k_2 L T_{1/2,A}\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2}\Big[k_2 q_{A,0} T_{1/2,A} T_{1/2,B}\theta_{1/2,B} - L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right)$$

$$+ \left\{-4k_2 Lq_{A,0}T_{1/2,A}T_{1/2,B}\theta_{1/2,B}\left(T_{1/2,B} + \theta_{1/2,B}\right)(\ln 2)^2 + \left(k_2 q_{A,0}T_{1/2,A}T_{1/2,B}\theta_{1/2,B} + L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right)\right)^2\right\}^{1/2}\Big] \tag{S10}$$

The subscript "fa" stands for "full analytical," indicating that the analytical solution applies to finite, constant values of $T_{1/2,A}$, $T_{1/2,B}$, and $T_{1/2,D}$.

Substituting Eq. (S10) into Eq. (S9) and solving for C_A, yields the following expression:

$$C_{A,fa} = \frac{1}{2k_2 L T_{1/2,B}\theta_{1/2,B}\ln 2}\Big[k_2 q_{A,0} T_{1/2,A} T_{1/2,B}\theta_{1/2,B} + L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right)$$

$$- \left\{-4k_2 Lq_{A,0}T_{1/2,A}T_{1/2,B}\theta_{1/2,B}\left(T_{1/2,B} + \theta_{1/2,B}\right)(\ln 2)^2 + \left(k_2 q_{A,0}T_{1/2,A}T_{1/2,B}\theta_{1/2,B} + L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right)\right)^2\right\}^{1/2}\Big] \tag{S11}$$

It is important to note that $C_{B,fa}$ and $C_{A,fa}$, as given by Eqs. (S10) and (S11), are independent of time.

Figs. S3b and S4b illustrate that the asymptotic values of C_B depend on both $T_{1/2,A,0}$ and $T_{1/2,B,0}$. A similar dependence for C_A is observed in Figs. S3a and S4a. Notably, the numerical and analytical solutions exhibit excellent agreement once the time exceeds five years (Figs. S3 and S4).

Solving Eq. (6) for C_D for the case of constant, finite $T_{1/2,D}$ value yields:

$$C_D = \frac{\left(1 - \exp\left(-\dfrac{t\ln(2)}{T_{1/2,D}}\right)\right)C_B T_{1/2,D}}{\theta_{1/2,B}} \tag{S12}$$

The value of $C_{D,fa}$ (which is time-dependent) can be determined by substituting $C_{B,fa}$ from Eq. (S10) into Eq. (S12). The plaque radius can be obtained by substituting $C_{D,fa}$ from Eq. (S12) into Eq. (S5). Note that for small values of t, Eq. (S12) collapses to Eq. (21) when the expression from Eq. (17) is substituted for C_B.

Additionally, Eqs. (S12) and (17) can be used to predict the asymptotic value that C_D approaches as $t \to \infty$:

$$C_{D,fa} = \frac{q_{A,0} T_{1/2,D}}{L \ln(2)} \tag{S13}$$

However, this asymptotic value is excessively large and unlikely to have biological significance.

Utilizing Eq. (25) yields the following result:

$$\Xi_{fa}(t) = C_{B,fa} t \tag{S14}$$

where $C_{B,fa}$ is defined by Eq. (S10).

The biological age is calculated by substituting Eq. (S14) into Eq. (27):

$$\text{biological age}_{fa} = c C_{B,fa} t \tag{S15}$$

where the expression for $C_{B,fa}$ is again provided in Eq. (S10).

By utilizing the "full analytical" solution from Eqs. (S15) and (S10), the following expressions are obtained for the dimensionless sensitivities of biological age to the half-lives of Aβ monomers and free aggregates, which reflect the efficiency of the Aβ degradation machinery:

$$S_{T_{1/2,A}}^{\text{biological age}} = \left[L\left(T_{1/2,B} + \theta_{1/2,B}\right)(\ln 2)^2 \right]\left\{ -4k_2 Lq_{A,0} T_{1/2,A} T_{1/2,B} \theta_{1/2,B} \left(T_{1/2,B} + \theta_{1/2,B}\right)(\ln 2)^2 \right.$$

$$\left. + \left(k_2 q_{A,0} T_{1/2,A} T_{1/2,B} \theta_{1/2,B} + L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right)\right)^2 \right\}^{-1/2} \tag{S16}$$

$$S_{T_{1/2,B}}^{\text{biological age}} = \left\{ \theta_{1/2,B} \left[k_2 q_{A,0} T_{1/2,A} T_{1/2,B} \theta_{1/2,B} + L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right) \right. \right.$$

$$\left. + \left\{ -4k_2 Lq_{A,0} T_{1/2,A} T_{1/2,B} \theta_{1/2,B} \left(T_{1/2,B} + \theta_{1/2,B}\right)(\ln 2)^2 + \left(k_2 q_{A,0} T_{1/2,A} T_{1/2,B} \theta_{1/2,B} + L\left(T_{1/2,B} + \theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A} + \ln 2\right)\right)^2 \right\}^{1/2} \right] \right\}$$

$$\times \left\{ 2\left(T_{1/2,B} + \theta_{1/2,B}\right)\left[-4k_2 Lq_{A,0} T_{1/2,A} T_{1/2,B} \theta_{1/2,B} \left(T_{1/2,B} + \theta_{1/2,B}\right)(\ln 2)^2 \right. \right.$$

$$+\left(k_2 q_{A,0} T_{1/2,A} T_{1/2,B}\theta_{1/2,B} + L\left(T_{1/2,B}+\theta_{1/2,B}\right)\ln 2\left(k_1 T_{1/2,A}+\ln 2\right)\right)^2 \ ]^{1/2}\Big\}^{-1} \qquad \text{(S17)}$$

S3. Numerical Solution

The system of differential equations (Eqs. (4)–(6)), along with the initial conditions defined in Eq. (7), was solved numerically using MATLAB's ODE45 solver (MATLAB R2024a, MathWorks, Natick, MA, USA). ODE45, which uses an adaptive timestep-step Runge-Kutta method, was chosen for its accuracy and robustness. To ensure high numerical precision, both the relative tolerance (RelTol) and absolute tolerance (AbsTol) were set to 1e-10.

S4. Supplementary figures

Fig. S1. (a) Molar concentration of Aβ aggregates deposited in plaques, C_D, as a function of time and (b) radius of a growing Aβ plaque, r_{ABP}, as a function of time. The plots are presented for different values of Aβ monomer flux into the CV, $q_{A,0}$. Parameters: $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s, $T_{1/2,D} = 10^{20}$ s. The plotted analytical solutions are obtained from Eq. (21) and (22).

Fig. S2. (a) Molar concentration of Aβ aggregates deposited in plaques, C_D, as a function of time and (b) radius of a growing Aβ plaque, r_{ABP}, as a function of time. The plots are presented for different values of

half-deposition times of free Aβ aggregates into senile plaques, $\theta_{1/2,B}$. Parameters: $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s, $T_{1/2,D} = 10^{20}$ s. The plotted analytical solutions are obtained from Eq. (21) and (22).

Fig. S3. Molar concentration of (a) Aβ monomers, C_A, and (b) Aβ oligomers, C_B, as a function of time for different values of the half-life of Aβ monomers, $T_{1/2,A}$. Parameters: $T_{1/2,B} = 10^{20}$ s. The plotted "full analytical" solutions are derived from Eqs. (S11) and (S10). Note that in each plotted case C_A and C_B reach constant asymptotic values as time increases. This is essential for deriving the full analytical solution.

Fig. S4. Molar concentration of (a) Aβ monomers, C_A, and (b) Aβ oligomers, C_B, as a function of time

for different values of the half-life of Aβ oligomers, $T_{1/2,B}$. Parameters: $T_{1/2,A}=10^{20}$ s. The plotted "full

analytical" solutions are derived from Eqs. (S11) and (S10). Note that in each plotted case C_A and C_B reach constant asymptotic values as time increases. This is essential for deriving the full analytical solution.

Fig. S5. Accumulated neurotoxicity of Aβ oligomers as a function of calendar age. (a) The half-life of Aβ monomers, $T_{1/2,A}$, remains constant regardless of calendar age. (b) The half-life of Aβ monomers, $T_{1/2,A}$, increases exponentially with calendar age. Parameters: $T_{1/2,B} = 10^{20}$ s.

Fig. S6. A function illustrating the exponential increase in the half-lives of Aβ monomers, $T_{1/2,A} / T_{1/2,A,0}$.

In this scenario, the half-life of Aβ monomers is assumed to increase by a factor of 2.5 by the time the calendar age reaches 50 years, as described in (Patterson et al., 2015). The same model is applied to the half-lives of free aggregates and aggregates deposited in Aβ plaques in the scenario of exponentially increasing half-life.

(a)

(b)

Fig. S7. Accumulated neurotoxicity of Aβ oligomers as a function of calendar age. (a) The half-life of free Aβ aggregates, $T_{1/2,B}$, remains constant over time. (b) The half-life of free Aβ aggregates, $T_{1/2,B}$, increases exponentially with calendar age. Parameters: $T_{1/2,A} = 10^{20}$ s.

Fig. S8. Biological age as a function of calendar age. (a) The half-life of Aβ aggregates after deposition into senile plaques, $T_{1/2,D}$, remains constant over time. (b) The half-life of Aβ aggregates after deposition into senile plaques, $T_{1/2,D}$, increases exponentially with calendar age. Parameters: $T_{1/2,A}=10^{20}$ s, $T_{1/2,B}=10^{20}$ s.

Fig. S9. (a) Accumulated neurotoxicity of Aβ oligomers as a function of calendar age for different values of the flux, $q_{A,0}$, of Aβ monomers per unit area at the left-hand face of the shaded CV shown in Fig. 1. (b)

Accumulated neurotoxicity of Aβ oligomers as a function of calendar age for different values of the half-deposition time, $\theta_{1/2,B}$, of free Aβ aggregates into senile plaques. Parameters: $T_{1/2,A}=10^{20}$ s, $T_{1/2,B}=10^{20}$ s.

Fig. S10. (a) Accumulated neurotoxicity of Aβ oligomers as a function of calendar age for different values of the nucleation rate constant, k_1, in the first pseudo-elementary reaction step of the F-W model for Aβ peptide. Parameters: $k_2 = 9.44 \times 10^{-6}$ μM⁻¹ s⁻¹, $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s, $T_{1/2,D} = 10^{20}$ s. (b) Accumulated neurotoxicity of Aβ oligomers as a function of calendar age for different values of the autocatalytic growth rate constant, k_2, in the second pseudo-elementary reaction step of the F-W model for Aβ peptide. Parameters: $k_1 = 2.22 \times 10^{-9}$ s⁻¹, $T_{1/2,A} = 10^{20}$ s, $T_{1/2,B} = 10^{20}$ s, $T_{1/2,D} = 10^{20}$ s.