Mesoderm-specific transcript reduces ciliary sphingomyelin levels to promote tendon stem/progenitor cells osteochondrogenesis in traumatic heterotopic ossification

Authors: Bowen Lai[1†], Yuan Gao[1†], Zhilong Shen[2], Jianquan Zhao[1], Heng Jiang[1*], Xuhui Zhou[1,3*]

Affiliations:

[1]Department of Orthopedics, The Second Affiliated Hospital of Naval Medical University; Shanghai, 200003, China.

[2]Department of Orthopedic Oncology, The Second Affiliated Hospital of Naval Medical University; Shanghai, 200003, China.

[3]Translational Research Centre of Orthopedics, Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine; Shanghai, 200080, China.

*Corresponding author. Email: include the email addresses of the corresponding author(s). Please use the asterisk (*) symbol for the corresponding author information.

† These authors contributed equally to this work.

One Sentence Summary: novel mechanisms and potential therapeutic strategies of traumatic heterotopic ossification

Abstract: Traumatic heterotopic ossification (tHO) is a prevalent musculoskeletal disorder with ectopic bone forms in soft tissues post-trauma. Despite its substantial impact on patients, which may range from local discomfort to severe functional impairments, the etiology and underlying mechanisms of tHO remain unknown, and no treatments have been demonstrated to be effective for tHO. Here, we found that ciliary Hedgehog (Hh) and PI3K-Akt signaling in tendon-derived stem/progenitor cells (TSPCs) is upregulated during tHO development, leading to the activation of GLI family zinc finger 2 (GLI2) transcription factor, which promotes mesoderm-specific transcript (MEST) gene expression. We further showed that MEST can change ciliary lipids composition and reduce the abundance of sphingomyelin in cilia, creating a positive feedback loop that amplifies ciliary Hh signaling, enhancing the osteogenesis and chondrogenesis of TSPCs and driving tHO. Cilia suppression by knocking down ciliary genes intraflagellar transport 88 (IFT88) or ADP ribosylation factor like GTPase 3 (ARL3), or using the SMO inhibitor vismodegib, restores normal TSPCs osteo-chondrogenic differentiation via downregulating Hh signaling. Inhibiting MEST expression or adding exogenous sphingomyelin in vivo also effectively alleviate tHO. These findings deepen our understanding of tHO pathogenesis and provide novel insights to develop novel therapeutic strategies against tHO.

Main Text:

INTRODUCTION

Traumatic heterotopic ossification (tHO) is a common musculoskeletal disorder characterized by the formation of ectopic bone in soft tissues, affecting a significant proportion of patients who experience severe injuries or orthopedic procedures(*1*). Although most cases of tHO present with localized symptoms such as pain, swelling, and restricted joint movement,

severe cases can lead to significant functional impairments, including joint ankylosis and chronic
disability, necessitating intensive surgical interventions like amputation(*2*). However, the
etiology and underlying biological mechanisms of tHO remain unknown. Notably, there are no
effective drugs available that can specifically eliminate ectopic bone formation or decelerate the
progression of tHO. Therefore, it is necessary to conduct more studies exploring the molecular
mechanisms of tHO to provide new strategies for clinical treatment that alleviate the suffering of
patients and the burden on medical resources.
Primary cilia are solitary, non-motile microtubule-based organelles that protrude from the
cell surface, acting as essential sensory antennae for cells to receive and transduce extracellular
signals(*3*). They play crucial roles in various developmental and physiological processes,
including embryonic patterning, tissue homeostasis, and cellular differentiation(*4*). Recent
studies have increasingly highlighted the involvement of primary cilia in skeletal biology. For
instance, primary cilia in osteocyte function as critical mechanosensors to maintain normal bone
formation and help to treat osteoporosis(*5*). Our previous study found that primary cilia can
positively regulate the process of osteochondral differentiation of stem cells from patients with
fibrodysplasia ossificans progressiva (FOP), a genetic form of HO caused by gene point
mutations(*6*). However, there is currently no research involving the function and regulatory
mechanisms of primary cilia in the pathogenesis of tHO.
The recruitment of stem/progenitor cells and their differentiation into osteoblasts or
chondrocytes are key steps in the formation of tHO(*7*). Tendon stem/progenitor cells (TSPCs)
are located between the long and parallel collagen fiber chains of tendon and are surrounded by
the extracellular matrix(*8*). TSPCs can differentiate into tenocyte tissue (tendon) and non-
tenocyte tissues (such as bone, cartilage, and adipose tissue) in different circumstances. Studies
have reported that the increased ectopic bone volume in Achilles tendon of rats is accompanied
with increased bone morphogenetic protein (BMP) expression and enhanced osteogenesis in
TSPCs(*9*). Moreover, the inhibition of the mTOR signaling by rapamycin treatment can suppress
the osteochondral differentiation of TSPCs and tHO(*10*). Thus, it is important to explore whether
primary cilia regulate the abnormal osteochondrogenic differentiation in TSPCs and how
primary cilia contribute to the progression of tHO via specific mechanisms.
MEST (mesoderm-specific transcript) is an imprinted gene predominantly expressed from
the paternal allele. The protein encoded by it is widely expressed in mesodermal tissues, belongs
to the α/β hydrolase superfamily, and has enzymatic activity. Through high-throughput
sequencing technology, we indicated that MEST might be a key downstream molecule in the
regulation of tHO by primary cilia. Studies have confirmed that knocking down MEST in human
periodontal ligament stem cells will inhibit the expression of their stem cell markers and
proliferation ability and weaken their multi-lineage differentiation abilities. However, no study
focuses on whether MEST is involved in regulating TSPCs osteogenic or chondrogenic
differentiation. Therefore, it is necessary to explore the role of MEST in the development of tHO
and the possibility of targeting MEST as a novel therapeutic strategy.
There is a diffusion barrier in the transition zone at the base of the primary cilia, which can
restrict the free movement of proteins and lipids in and out of the cilia, resulting in differences in
the composition of proteins and lipids on the ciliary membrane and the cytoplasmic membrane.
Under the stimulation of mechanical stress or chemical signals, the selective movement of
specific lipids in and out of the cilia can dynamically influence the function of receptors on the
ciliary surface and the activity of downstream signaling pathways, thereby regulating cell
proliferation and differentiation(*11*). The disruption of the homeostasis of ciliary lipids has been

reported to be associated with a variety of diseases. Some studies have reported that
abnormalities in genes encoding key enzymes involved in cholesterol metabolism can lead to
defects in the formation of primary cilia in osteoblasts and in the differentiation of osteoblasts,
resulting in craniofacial malformation(*12, 13*). Since MEST has been found to possess the
characteristics of lipase, which can promote the hypertrophy of adipocytes and expand the
volume of adipose tissue(*14*), the regulation of lipid changes within the cilia of TSPCs by MEST
may be an important part of the pathogenesis of tHO.
In this study, we demonstrated that the activity of ciliary Hh and PI3K-Akt signaling of
TSPCs is significantly upregulated during tHO development, leading to the activation and
nuclear translocation of the transcription factor GLI2, which promotes the expression of the
MEST gene. The upregulation of MEST expression results in a reduction of sphingomyelin
abundance within the primary cilia, which in turn amplifies the activation of the ciliary Hh
signaling. This forms a positive feedback loop that significantly enhances the osteogenic and
chondrogenic differentiation potential of TSPCs, thereby driving local ectopic bone deposition
and contributing to the development and progression of tHO. Moreover, knockdown ciliary gene
or suppressing Hh signaling restored the TSPCs differentiation. Inhibiting MEST expression or
adding exogenous sphingomyelin in vivo also had great therapeutic effects on tHO. Unraveling
these mechanisms may enhance our understanding of the pathogenesis of tHO and pave the way
for the development of innovative and effective therapeutic approaches.

RESULTS

1. Inhibition of ciliary function alleviates the formation of tHO

To induce tHO, 6- to 8-week-old male C57BL/6 mice were subjected to a burn injury using a
60°C heated aluminum block on the dorsal skin, followed by resection of the right Achilles
tendon (Figure 1a). Tendon tissues from the injured sites were harvested at 0 days (prior to
injury), and at 3, 7, 21, and 56 days after injury for subsequent histological examination (Figure
1b). HE staining demonstrated disorganized and irregular tendon fibrous tissue at 3 to 7 days
post-injury. Chondrogenic structures were detected at 21 days, and osteogenic structures were
observed at 56 days. Masson trichrome staining showed blue-stained tendon collagen fibers, with
the progression of structural disorganization and tHO being similar to that observed with HE
staining. Von Kossa staining revealed the appearance of brown calcified nodules at 21 days,
which became deeply stained by 56 days, suggesting calcium deposition due to new bone
formation.
To explore the function of cilia in tHO, we performed immunofluorescence staining on
tendon tissues collected from mice at 0, 3, 7, 21, and 56 days post-injury (Figure 1c). ARL13B, a
specific marker for primary cilia, was used to quantify ciliated cells. The percentage of ARL13B-
positive cells was 27.21% in the control group (0 days post-injury). After injury, the proportion
of ARL13B-positive cells increased to 40.14% at 3 days, 38.07% at 7 days, and 39.75% at 21
days, before declining to 33.62% at 56 days. These data indicated that ciliated cells increased
rapidly during tHO and may contribute to its development (Figure 1d). To further identify the
potential specific cell types targeted by cilia in the process of tHO, immunofluorescence double-
staining was performed on tendon tissue samples collected at various time points post-injury.
The following marker pairs were used: ARL13B with CD31 (endothelial cells marker), ARL13B
with F4/80 (macrophages marker), ARL13B with PDGFRα (stem/progenitor cells marker),
ARL13B with TNMD (tendon cells marker), and ARL13B with α-SMA (smooth muscle cells
marker). Immunofluorescence analysis revealed clear co-localization between ARL13B and

PDGFRα after tHO injury (Figure 1c). Quantitatively, the proportion of
ARL13B+PDGFRα+cells was the highest among the five double-staining groups. Specifically,
the percentage of ARL13B+PDGFRα+cells was 55.91% in the control group. After injury, this
proportion significantly increased to 95.29% at 3 days, 85.56% at 7 days, 90.17% at 21 days, and
95.49% at 56 days (Figure 1e, f). These data indicate that cilia mainly affect tHO by targeting
stem/progenitor cells.
Intraflagellar transport protein 88 homolog (IFT88) functions as a core protein of the IFT-B
complex, driving anterograde transport within cilia. Arf-like Protein 3 (ARL3), a small GTPase,
is involved in ciliogenesis and the maintenance of ciliary function. To locally inhibit ciliary
function, we administered adeno-associated virus (AAV) vectors carrying *shIft88* and *shArl3* via
intramuscular injection into the lower limbs of mice with tHO. The results showed that
knockdown of *Ift88* and *Arl3* significantly decreased the bone volume of ectopic bone formation,
whereas bone density was not significantly altered. This suggests that inhibition of ciliary
function can reduce the severity of tHO (Figure 1g).
146
2. Primary cilia in stem/progenitor cells contribute to tHO progression

Single-cell RNA sequencing (scRNA-seq) can distinguish differences among individual
cells in a high-throughput manner, thereby clarifying cellular heterogeneity and underlying
molecular mechanisms during disease progression. Therefore, in our study, we reanalyzed a
scRNA-seq dataset (GSE126060) from mouse tHO tissues. The dataset included 12 samples,
with three samples per group at four time points: pre-injury (Day 0) and post-injury (Day 3, Day
7, and Day 21). The UMAP plot showed a total of 21,880 cells detected and divided into nine
subclusters (Figure 2a) according to the previous study(*15*). Bubble plot revealed the marker
genes of each cluster, and markers for the Mesenchymal cluster were Pdgfra and Prrx1 (Figure
2b, c). Group-stratified UMAP plots revealed that, as tHO progresses, the percentage of
Macrophage cluster cells peaked at 3 days (74.43%) post-injury, later rapidly reduced to 37.10%
at 7 days and 15.02% at 21 days post-injury, indicating the evolution of the local inflammatory
phase (Figure 2d). In contrast, the Mesenchymal cluster cells exhibited a larger proportion at 7
(43.44%) and 21 days (49.43%), suggesting that the osteogenic and chondrogenic differentiation
of stem/progenitor cells contributes to tHO during this period (Figure 2d). Consequently, we
analyzed the differentially expressed genes in the Mesenchymal subcluster at 7 and 21 days post-
injury compared with the control group and revealed that 82 genes were upregulated in the
Mesenchymal subcluster at Day 7, and 53 genes were upregulated at Day 21 (Figure 2e). Since
the knockdown of ciliary genes *Ift88* and *Arl3* can alleviate tHO formation in vivo, we are
interested in the expression of ciliary genes at single-cell level. *Ift88* has low expression at single
cell level likely due to limited scRNA-seq depth, but still predominantly localizes to the
mesymenchal subpopulation, and shows a subtle upward trend during tHO (Figure S1). We then
shifted our focus to another ciliary gene IFT20 which is also a component of the IFT-B complex
and has been reported to regulate ciliogenesis and skeletal development(*16*). It can be shown that
both *Ift20* and *Arl3* are highly expressed in the Mesenchymal cluster (Figure 2f). Moreover, with
the developmnet of tHO after injury, the expression of *Ift20* and *Arl3* in Mesencymal cluster
rapidly increased in Day 7 and Day 21 group, while the trends over time were not significant in
other clusters (Figure 2g). The results indicated primary cilia and ciliary genes may mainly
functioned in stem/progenitor cell (like TSPCs as we identified above) during the progression of
tHO.
177

3. Primary cilia regulate the osteochondrogenic differentiation potential of TSPCs

Since the above results suggested that cilia on stem/progenitor cells may regulate tHO, and considering that abnormal differentiation of tendon stem/progenitor cells (TSPCs) has been reported to potentially contribute to the pathogenesis of tHO(*17, 18*), we isolated TSPCs from the tendon tissues of C57BL/6 mice. Microscopic examination showed that these TSPCs exhibited a fibroblast-like spindle-shaped morphology (Figure 3a). Flow cytometry analysis demonstrated that 88.1% of the TSPCs were double-positive for Pdgfrα and Prrx1 (consistent with the scRNA-seq results), thereby confirming the stem/progenitor cell characteristics of the isolated TSPCs (Figure 3b).

To investigate the regulatory effects of cilia on the functions of TSPCs, we transfected lentivirus vectors carrying *shIft88* and *shArl3* into mouse TSPCs to establish stable transfectants with knockdown of ciliary genes. qPCR and Western blot analyses confirmed successful knockdown of *Ift88* and *Arl3*, indicating that these transfectants were suitable for subsequent experiments (Figure S2). Immunofluorescence staining for ARL13B and α-tubulin (Acetyl Lys40) was used to observe changes of primary cilia in TSPCs following knockdown of cilia. The results showed a significant reduction in cilia counts and length (Figure 3c). Quantitative analysis revealed that the proportion of ciliated TSPCs was approximately 53.54% in the control group but decreased to 19.36% in the *shIft88* group and 34.70% in the *shArl3* group, indicating a marked decrease in cilia number in TSPCs following ciliary gene knockdown. Additionally, the average cilia length was 3.37 μm in the control group but reduced to 2.06 μm in the *shIft88* group and 2.14 μm in the *shArl3* group, demonstrating a significant shortening of cilia length in TSPCs after ciliary gene knockdown (Figure 3d, e).

Previous studies have shown that the osteogenic and chondrogenic differentiation capabilities of TSPCs are abnormally activated during the process of tHO, leading to the formation of ectopic bone in tendon tissues(*18*). Therefore, investigating whether cilia can regulate the osteogenic and chondrogenic differentiation of TSPCs is important for exploring the regulatory mechanisms of cilia in tHO. We induced osteogenic differentiation in mouse TSPCs and performed alizarin red staining after 21 days of induction. Compared with the control group, the alizarin red staining was lighter in the *shIft88* and *shArl3* groups, indicating less calcium deposition (Figure 3f, S2). This suggests that knockdown of ciliary genes attenuates the osteogenic differentiation capacity of TSPCs. qPCR analysis revealed that the expression of osteogenic differentiation-related genes *Alpl*, *Bglap*, and *Sp7* was significantly reduced in the *shIft88* and *shArl3* groups compared with the control group, while *Runx2* expression remained largely unchanged (Figure 3g). These findings further confirm a positive correlation between cilia function and the osteogenic differentiation capacity of TSPCs. Similarly, we induced chondrogenic differentiation in the stable transfectants of mouse TSPCs with ciliary gene knockdown and performed alcian blue staining after 21 days. Compared with the control group, the blue staining was lighter (Figure 3f, S3). Meanwhile, qPCR analysis revealed that the expression of chondrogenic differentiation-related genes *Col2a1* and *Acan* was significantly reduced, while *Sox9* expression retained (Figure 3g). These results indicate that primary cilia may positively modulate the osteogenic and chondrogenic differentiation of TSPCs, leading to abnormal osteogenesis and chondrogenesis in local soft tissues and ultimately resulting in tHO.

4. Primary cilia modulate MEST expression to promote tHO

To further identify key genes involved in tHO and their potential crosstalk with primary cilia,
we obtained bulk-RNA sequencing data from a public database for mouse tHO tissues
(GSE233201) (Figure 4a). We then integrated these data with RNA sequencing results from
TSPCs with ciliary gene *Ift88* knockdown and control groups to identify common differentially
expressed genes (Figure 4b). Genes upregulated in the Mesenchymal cluster at 7 and 21 days
during tHO (GSE126060), genes upregulated in injured tissue at 21 days of tHO (GSE233201),
and genes downregulated in TSPCs with ciliary gene *Ift88* knockdown were obtained
respectively (Table S1). Venn diagram analysis identified three common differentially expressed
genes: MEST, PTN, and COL2A1 among these three datasets (Figure 4c). COL2A1 is a key
gene for chondrogenic differentiation, while PTN showed insignificant differences after
validation (Figure S4). Thus, we focused on the role of the MEST gene in tHO in subsequent
studies. scRNA-seq analysis revealed that *Mest* is predominantly expressed in the Mesenchymal
cluster (Figure 4d). Both qPCR and Western blot analyses demonstrated a significant reduction
in *Mest* RNA and protein expression following knockdown of the ciliary genes *Ift88* or *Arl3*
(Figure 4e, f). Moreover, scRNA-seq analysis showed that the expression of *Mest* in the
Mesenchymal cluster exhibits an upward trend with the progression of tHO (Figure 4g).
Consistently, Immunohistochemical analysis of tendon tissues at various time points after
establishment of tHO further showed that MEST protein expression is progressively upregulated
during the tHO development (Figure 4h), suggesting that MEST may play a significant role in
the regulatory mechanisms of tHO. Notably, immunofluorescence staining showed no significant
co-localization between MEST and α-tubulin (Figure 4i), indicating that MEST likely functions
outside the cilia in TSPCs to influence their function and thereby contribute to the occurrence
and progression of tHO.
245
5. MEST facilitates the TSPCs differentiation and tHO in a ciliary localization-
independent manner

As MEST is primarily not localized or expressed on the cilia of TSPCs, we explored how
MEST can regulate tHO in a cilia localization-independent manner. We constructed two small
interfering RNAs (siRNAs): *siMest-1* and *siMest-2*, to inhibit *Mest* gene expression in vitro.
qPCR analysis revealed a significant decrease in *Mest* expression following *siMest* intervention,
confirming that *siMest* can effectively suppress MEST function (Figure 5b). Alizarin red staining
showed that the *siMest* group exhibited lighter staining and fewer calcium nodules compared
with the control group after 21 days of osteogenic differentiation induction (Figure 5a, S5).
Additionally, the expression of osteogenic differentiation-related genes *Alpl, Runx2* and *Bglap*
was significantly reduced (Figure 5b). Similarly, alcian blue staining after 21 days of
chondrogenic differentiation induction showed that, compared with the control group, the *siMest*
group exhibited lighter staining and reduced chondrogenic capacity (Figure 5a, S5). The
expression of chondrogenic differentiation-related genes *Sox9, Col2a1*, and *Acan* was also
decreased (Figure 5b). Moreover, we divided the Mesenchymal cluster to Mest-high and Mest-
low subclusters based on the expression level of Mest in tHO scRNA-seq data (Figure 5c).
Though *Alpl, Sp7,* and *Col2a1* were not significantly expressed between two subclusters
probably because of their extreme low expression in single-cell level (Figure S6), the expression
of *Runx2, Sox9,* and *Acan* were greatly higher in Mest-high subcluster, which further
demonstrated our results. These findings indicate that MEST promotes the TSPCs

osteochondrogenesis and inhibiting MEST function can diminish the osteogenic and
chondrogenic differentiation capacity of TSPCs in tHO.
We further employed immunofluorescence staining with α-tubulin to label the cilia of TSPCs
in the control and *siMest* groups (Figure 5c). Quantitative analyses showed no significant
difference in cilia number (58.26% vs 53.11%), while cilia length was significantly shorter in the
siMest group compared with the control group (2.46 μm vs 3.48 μm) (Figure 5d). In addition, we
performed in vivo experiments by locally injecting a AAV vector carrying *shMest* into mice with
tHO to knock down *Mest* expression in local tissues. Micro-CT imaging revealed that the volume
of ectopic new bone in the *shMest* group was significantly lower than that in the control group,
while bone density remained largely unchanged (Figure 5e, f). Collectively, these results suggest
that MEST are involved in regulating the ciliary function of TSPCs and their osteochondrogenic
differentiation in a ciliary localization-dependent manner. Inhibition of *Mest* expression in the
specific way may serve as a potential therapeutic target for treating tHO.
279
6. MEST targets ciliary Hh and PI3K-Akt signaling to mediate TSPCs differentiation in
tHO

To further investigate the specific molecular mechanisms and downstream targets through
which cilia regulate tHO, we conducted enrichment analysis on our previously obtained
sequencing data. Gene Ontology (GO) enrichment analysis showed that differentially expressed
genes at the site of tHO 21 days post-injury were significantly enriched in functions related to the
extracellular matrix and ossification (Figure 6a). Consistent with these findings, differentially
expressed genes in TSPCs following knockdown of cilia-related genes were also enriched in
ossification (Figure 6b). These results suggest a strong correlation between the differentially
expressed genes and tHO, indicating that gene engineering therapies targeting these genes may
represent a potential strategy for treating tHO. We further conducted Gene Set Enrichment
Analysis (GSEA) on the differentially expressed genes in TSPCs following knockdown of cilia-
related genes, and the results showed significant enrichment in the Hh signaling pathway (Figure
6c). Given that the activity of the Hedgehog (Hh) signaling pathway is dependent on the primary
cilia for regulating embryonic development and tissue differentiation, and considering that
aberrant upregulation of Hh has been implicated in the pathogenesis of tHO following injury(*19*),
the Hh signaling activity within TSPCs cilia is likely closely related to tHO. Additionally, KEGG
enrichment analysis indicated that differentially expressed genes in the Mesenchymal cluster of
tHO tissues, as well as those in TSPCs following knockdown of ciliary gene *Ift88*, were
significantly enriched in the PI3K-Akt signaling pathway (Figure 6d, e). To verify the
bioinformatics analysis results, we performed qPCR and found that the mRNA expression of key
Hh signaling molecules *Shh* and *Ptch1* was significantly downregulated in TSPCs undergoing
osteogenic and chondrogenic differentiation after *shIft88* or *shArl3* treatment (Figure 6f).
Western blot analysis also demonstrated a significant reduction in the protein expression of SHH
and PTCH1 (Figure 6g). Additionally, during the osteogenic and chondrogenic differentiation of
TSPCs, the ratios of phosphorylated PI3K to total PI3K (p-PI3K/PI3K) and phosphorylated AKT
to total AKT (p-AKT/AKT) were significantly decreased following ciliary gene knockdown
(Figure 6g), indicating reduced phosphorylation of PI3K and AKT and thus inhibition of the
PI3K-Akt signaling pathway. Smoothened (SMO) is a key seven-transmembrane receptor
involved in Hh signaling and is inhibited by PTCH1 when Hh signaling is inactive. Upon
binding of the Hh ligand to PTCH1, the inhibitory effect of PTCH on SMO is alleviated, thereby

enabling the activation of SMO. Interestingly, scRNA-seq analysis showed that *Smo* mainly
expressed and increased over time in Mesenchymal cluster (Figure 6h, i) after trauma, further
demonstrating the Hh signaling was activated in tHO development. Collectively, these results
suggest that the primary cilia of TSPCs may lead to aberrant activation of the Hh and PI3K-Akt
signaling pathways, thereby promoting osteogenic and chondrogenic differentiation of TSPCs
and exacerbating tHO.
Previous results have confirmed that MEST can influence the osteogenic and chondrogenic
differentiation of TSPCs and regulate tHO in ciliary localization-independent manner, yet the
underlying molecular mechanisms remain unclear. qPCR (Figure S7) and Western blot (Figure
7a) analyses indicated that, compared with the control group, the expression levels of key Hh
signaling molecules SHH and PTCH1 were significantly downregulated at both the mRNA and
protein levels in *siMest*-transfected TSPCs undergoing osteogenic and chondrogenic
differentiation. In addition, the phosphorylation levels of PI3K and AKT were also decreased
(Figure 7b). These findings suggest that MEST may promote the osteogenic and chondrogenic
differentiation of TSPCs and the formation of tHO by mediating the Hh and PI3K-Akt signaling
pathways. GLI2, a key transcriptional activator of the Hh signaling pathway, translocate to the
nucleus upon Hh signal activation and induces the expression of downstream genes involved in
cell proliferation and differentiation. The sequence logo of GLI2 transcription factor was shown
and the JASPAR database predicts that the GLI2 can bind to the MEST gene promoter at
positions 904-916 and 1698-1710, thereby activating MEST transcription and enhancing its
expression (Figure 7c). Dual-luciferase reporter assays using pGL4.10 as vector demonstrated
that GLI2 can bind to the MEST promoter at both predicted sites, with overexpression of GLI2
significantly increasing luciferase activity, thus confirming that GLI2 positively regulates Mest
transcription (Figure 7d). To further elucidate the regulatory role of the GLI2 transcription factor
in tHO, we performed immunohistochemistry on tendon tissues obtained from mice with tHO
and found that GLI2 expression progressively increased with the development of the ossification
(Figure 7e). Meanwhile, qPCR and Western blot analyses revealed that, compared with the
control group, the *Gli2* mRNA and protein expression levels of GLI2 were significantly
downregulated in TSPCs undergoing osteogenic and chondrogenic differentiation after
knockdown of ciliary genes (Figure 7f, g). Moreover, in vivo experiments involving local
injection of a AAV vector carrying *shMest* into mice with tHO to knockdown *Mest* expression
showed decreased protein expression levels of GLI2, PTCH1, and SMO, as shown by
immunohistochemistry (Figure 7h), indicating that targeting Mest may alleviate the occurrence
and progression of tHO by reducing the activation level of the Hh signaling pathway.
Collectively, these results suggest a self-amplifying loop of GLI2 and MEST via Hedgehog
signaling that promotes the osteogenic and chondrogenic differentiation of TSPCs, and thereby
exacerbates the tHO phenotype.
348

7.   MEST reduced ciliary sphingomyelin levels in TSPCs

Recent studies have shown that ciliary lipids play a crucial role in maintaining ciliary
structure stability and regulating ciliary mechanosensation and signal transduction. Aberrant
distribution and dynamic changes in ciliary lipids have been implicated in various diseases,
including polycystic kidney disease(*20*) and obesity(*21*). Moreover, deficiencies in key
cholesterol-metabolizing enzymes have been shown to cause ciliary defects that impact bone
development(*13*). These findings have prompted our interest in investigating the changes in
lipids within TSPCs cilia during tHO. We first isolated cilia from TSPCs in the control and

siMest groups using ultracentrifugation to prepare ciliary suspensions (for more details see
Methods). Immunofluorescence confirmed that the cilia were effectively removed from TSPCs
after treatment (Figure 8a). Then we conducted Untargeted lipidomics using liquid
chromatography-tandem mass spectrometry (LC-MS/MS) on the collected ciliary suspensions.
OPLS-DA analysis showed obvious differences in the lipid profiles of TSPCs cilia between the
control (NC) and *siMest* (SIM) groups (Figure 8b). We performed differential analysis of lipids
based on the LIPID MAPS® classification system and the differential expressed primary (Figure
8c, d) and secondary (Figure S8a) lipids were classified. Totally, we screened 159 upregulated
and 157 downregulated lipids in the SIM group with the criteria of $|log2FC| > 1$ and P value <
0.05 (Table S2, Figure 8e, S8b). Compared with the NC group, the abundance of sphingomyelin
(SM) within the cilia of TSPCs in the SIM group was significantly increased, while the content
of triglycerides (TG) was significantly decreased (Figure 8f). Heatmaps revealed consistent
differential changes in almost all subclasses of sphingomyelins and triglycerides, including 22
sphingomyelins and 53 triglycerides (Figure S8c). To further explore how the differentially
expressed lipids within the cilia of TSPCs might contribute to tHO, we conducted KEGG
pathway enrichment analysis. Consistent with our prior findings, the differentially expressed
metabolites were enriched in pathways including glycerophospholipid metabolism, sphingolipid
signaling, and sphingolipid metabolism (Figure 8g, S8d).
SGMS2 (sphingomyelin synthase 2) gene encodes sphingomyelin synthase 2, which is
mainly involved in the synthesis of sphingomyelin on the cell membrane. It has been reported
that SGMS2-mediated sphingomyelin metabolism was involved in skeletal homeostasis
regulation and its mutation could lead to a series of bone diseases like osteoporosis and skeletal
dysplasia(*22*). We found that *Sgms2* was mainly expressed in Mesenchymal cluster but rarely
expressed in Mest-high subcluster (Figure 8h). Moreover, the expression level of *Sgms2* in Mest-
high subcluster gradually reduced over time after tHO at single-cell level (Figure 8i), indicating
that Mest reduced ciliary sphingomyelin synthesis in tHO possibly by inhibiting *Sgms2*
expression in TSPCs. These results indicated that MEST may reduce the synthesis and
abundance of sphingomyelin within the cilia of TSPCs, and that silencing *Mest* expression in
vitro can reverse the decrease in sphingomyelins within the cilia of TSPCs.
386
8.   Targeting ciliary sphingomyelin in TSPCs rescue tHO development via Hh signaling

Since previous findings have demonstrated that MEST can induce the osteogenic and
chondrogenic differentiation of TSPCs in ciliary localization-independent manner, thereby
promoting the occurrence and development of tHO, it is essential to explore whether lipids
within the cilia of TSPCs also contribute to this pathogenesis. Research has shown that
sphingomyelins and cholesterol interact on the cellular membrane to regulate intracellular
signaling processes, and their dynamic balance is crucial for maintaining normal cellular
functions(*23*). Considering the parallel roles of cholesterol and triglycerides under physiological
and pathological conditions, we hypothesize that the decrease in triglycerides within the cilia of
TSPCs following *Mest* knockdown may follow the increase in sphingomyelins. Therefore,
studying how sphingomyelins regulate TSPCs function is of great significance. We induced
osteogenic and chondrogenic differentiation in TSPCs and performed alizarin red and alcian blue
staining with different treatments respectively (Figure 9a, b). Compared with the control group,
treatment with exogenous sphingomyelin significantly diminished the osteogenic and
chondrogenic differentiation capacities, indicating that sphingomyelins can suppress the
osteogenic and chondrogenic differentiation of TSPCs. Treatment of TSPCs with myriocin (a

sphingomyelin synthesis inhibitor) significantly enhanced the osteogenic and chondrogenic differentiation capacities of TSPCs. However, when sphingomyelins and myriocin were applied together, the osteogenic and chondrogenic differentiation capacities of TSPCs were not significantly different from the control group. Furthermore, downregulation of *Mest* gene expression in TSPCs using *siMest* mitigated the excessive upregulation of osteogenic and chondrogenic differentiation induced by myriocin-mediated reduction in sphingomyelins. qPCR analysis of the expression of osteogenic (*Alpl, Runx2,* and *Bglap*) and chondrogenic (*Col2a1, Sox9,* and *Acan*) differentiation-related genes in TSPCs revealed trends consistent with the staining results (Figure 9c), suggesting that inhibiting *Mest* gene expression may increase the sphingomyelin content in TSPCs cilia, thereby attenuating their osteogenic and chondrogenic differentiation capacities and alleviating tHO.

Notably, a study has demonstrated that depletion of sphingomyelins on the ciliary membrane can increase the accessibility of cholesterol, thereby amplifying the transmission of the Hh signaling pathway within the cilia[24]. Therefore, we hypothesize that the results may be associated with the activity of the Hh signaling pathway. Smoothened (SMO), a seven-transmembrane receptor protein, serves as a bridge between Hh ligands and GLI transcription factors in the Hh pathway. Direct targeting of SMO has become a trending area in drug research targeting the Hh signaling pathway, with the SMO inhibitor vismodegib being FDA-approved for the treatment of basal cell carcinoma. When vismodegib was administered to TSPCs treated with myriocin, the osteogenic and chondrogenic differentiation capacities were significantly reduced (Figure 9a, b), and the expression of osteogenic and chondrogenic differentiation-related genes was downregulated, with no significant differences compared to the control group (Figure 9c). qPCR analysis of key Hh signaling molecules (*Shh, Ptch1, and Gli2*) during osteogenic differentiation of TSPCs showed that myriocin strongly activated the Hh signaling pathway, while exogenous sphingomyelin had the opposite effect, inhibiting Hh signaling (Figure 9d). In contrast, groups treated with myriocin+sphingomyelin, myriocin+siMest, and myriocin+vismodegib exhibited Hh signaling pathway component gene expression levels that were not significantly different from the control group (Figure 9d). In vivo experiments by local injection of exogenous sphingomyelin into tHO mice injury site revealed that the volume of ectopic bone in mice treated with exogenous sphingomyelin was significantly lower than that in the PBS-injected control group, with no significant differences in bone mineral density between the two groups (Figure 9e). These findings suggest that the depletion of cilia sphingomyelins may facilitate the osteogenic and chondrogenic differentiation of TSPCs through the activation of the Hh signaling pathway, thereby exacerbating tHO. This effect can be attenuated by silencing the MEST gene or using a SMO inhibitor vismodegib, which may represent emerging candidate therapeutic targets for tHO.

DISCUSSION

The formation of traumatic heterotopic ossification (tHO) is a complex process involving the aberrant differentiation of different cells into osteogenic and chondrogenic lineages. Primary cilia are microtubule-based, non-motile organelles that act as cellular antennae, sensing and transducing extracellular signals to regulate various developmental and physiological processes. Several studies have revealed the relationship between primary cilia and skeletal diseases, including defects of skeletal development such as polydactyly[25] and craniofacial abnormalities[12]. Our previous research found that primary cilia in skeletogenic cells is crucial for BMP and ActA-mediated osteogenesis and chondrogenesis and thereby governs the development of fibrodysplasia ossificans progressiva (FOP). In this study, we demonstrate that

cilia function is significantly upregulated during tHO, with an increased prevalence of ciliated cells in the injury site, particularly among tendon stem/progenitor cells (TSPCs). This observation aligns with previous studies indicating that primary cilia are essential for tissue repair(26) and regeneration(27) but extends our understanding by linking cilia activity directly to the pathogenesis of tHO.

Our study reveals that primary cilia positively modulate the osteogenic and chondrogenic differentiation of TSPCs, contributing to the development of tHO. Knockdown of cilia-related genes (Ift88 and Arl3) significantly reduced the differentiation capacity of TSPCs, as evidenced by decreased calcium deposition and downregulation of key osteogenic and chondrogenic markers, resulting in reduced ectopic bone volume in tHO mice. This finding is consistent with the established role of primary cilia in regulating stem/progenitor cell functions in various cell types. For examples, loss of Numb (a ciliary protein) in spinal neural progenitors lead to reduced cerebellar size via ciliary Hh signaling in mice(28). The Hh signaling pathway, which is cilia-dependent, is known to be aberrantly activated in postnatal skeletogenesis(29) and heterotopic ossification in osteoblast(30). Meanwhile, inactivated PI3K-Akt signaling after ciliary gene IFT80 deficiency results in decreased odontogenic differentiation in dental pulp stem cells(31). Our study further supports this by showing that cilia gene knockdown in TSPCs downregulates Hh signaling components and reduces PI3K-Akt signaling activation, leading to reduced osteogenesis and chondrogenesis. This suggests that primary cilia serve as a crucial regulatory hub, integrating signals that drive the aberrant osteogenic and chondrogenic differentiation of TSPCs during tHO.

A significant contribution of our study is the identification of MEST as a core regulator of tHO for the first time. MEST encodes protein that is widely expressed in mesoderm. As we know, the mesoderm develops into the skin, muscle, bone, and other connective tissues in our body, which indicate the potential role of MEST in skeletogenesis. Our scRNA-seq results showed that MEST is predominantly expressed in stem/progenitor cells and is upregulated during tHO progression. A recent study reported that MEST specifically expressed in mesenchymal cells and induced bone formation during mouse digit tip regeneration(32). In our study, scRNA-seq showed that the expression levels of osteogenic and chondrogenic genes in Mest highly expressed stem/progenitor cells were higher than Mest low expressed stem/progenitor cells. Moreover, knockdown of MEST in TSPCs attenuated their osteogenic and chondrogenic differentiation capacities, while shMest significantly alleviate severity of tHO in mice, indicating a pivotal role of MEST in promoting tHO by regulating TSPCs differentiation. Interestingly, MEST functions in a cilia localization-independent manner, suggesting that it may act downstream of cilia signaling pathways. This finding is innovative as it uncovers a novel regulatory axis involving MEST and highlights its potential as a therapeutic target for tHO.

Our study elucidates the molecular mechanisms through which MEST regulates tHO. We demonstrate that MEST modulates the Hh and PI3K-Akt signaling pathways, which are critical for TSPC differentiation. Furthermore, we show that the GLI2 transcription factor, a key activator of the Hh pathway, binds to and promotes MEST transcription. This finding is supported by previous studies indicating that GLI2 is a master regulator of bone homeostasis(33) and that its activity is tightly regulated by cilia function(34). In summary, the results indicated a positive feedback loop where Hh signaling activates MEST, which in turn enhances TSPC osteogenic and chondrogenic differentiation and tHO progression via Hh and PI3K-Akt signaling.

In the presence of external stimuli, the selective influx and efflux of specific lipids in primary cilia can dynamically modulate the functions of cilia and cells(11). This process is

intricately linked to adipogenesis(*21*) and diseases such as polycystic kidney disease(*20*). Thus,
another novel aspect of our study is the investigation of ciliary lipid metabolism in TSPCs during
tHO. By untargeted lipidomic after isolation of primary cilia, we show that MEST regulates the
abundance of various important lipids within TSPC cilia, especially with sphingomyelin levels
being significantly upregulated upon MEST knockdown. This finding is intriguing as
sphingomyelins are known to interact with cholesterol to regulate ciliary signaling processes(*24*).
Our study suggests that the balance of sphingomyelins and triglycerides within cilia is crucial for
TSPCs differentiation. Specifically, we demonstrated that exogenous sphingomyelin treatment
inhibits TSPCs osteogenic and chondrogenic differentiation and reduce ecotopic bone volume in
tHO mice. This effect is mediated through the ciliary Hh signaling, as evidenced by the
activation of Hh signaling upon myriocin treatment and its inhibition by exogenous
sphingomyelin. Vismodegib, a SMO inhibitor targeting Hh signaling, can also diminish
osteogenesis and chondrogenesis of TSPCs. These findings provide a new perspective on the role
of lipid metabolism in tHO and suggest that targeting ciliary sphingomyelin levels may represent
a potential therapeutic strategy.
Several limitations should be acknowledged in the study. First, our study primarily focuses
on the role of primary cilia in TSPCs, while other cell types involved in tHO, such as
macrophages and endothelial cells, were not extensively investigated. Future studies should
explore the cilia-dependent functions of these cells and their interactions with TSPCs during tHO.
Second, our in vivo experiments were limited to local injection of AAV vectors carrying *shRNA*,
which may not fully recapitulate the complex physiological environment of tHO. Further studies
using genetically modified animal models are needed to validate our findings. Third, while we
identified MEST as a key regulator of tHO, the detailed molecular mechanisms through which
MEST modulates sphingomyelin metabolism within cilia remain unclear. Future studies should
investigate the specific interactions between MEST and lipid metabolic enzymes and their
impact on cilia function as the regulator of tHO.
The schematic diagram of the study is shown in Figure 10. Our study opens several avenues
for future research. First, further investigation into the role of primary cilia in other cell types
involved in tHO is warranted. This will provide a more comprehensive understanding of the
cellular and molecular mechanisms underlying tHO progression. Second, exploring the
therapeutic potential of targeting cilia function or MEST expression in preclinical models of tHO
is an important next step. This could involve the development of small molecules or gene
therapies aimed at modulating cilia activity or inhibiting MEST function. Third, elucidating the
detailed molecular mechanisms through which MEST regulates lipid metabolism within cilia
will provide new insights into the pathogenesis of tHO and identify potential therapeutic targets.
Finally, translating these findings into clinical applications will require rigorous preclinical and
clinical trials to assess the safety and efficacy of cilia-targeted therapies for tHO.

MATERIALS AND METHODS

1. Experimental tHO mice

Male C57/BL6 mice weighing approximately 20 g and aged 6–8 weeks bought from
Hangzhou Ziyuan Laboratory Animal Technology Co., Ltd were used for construction of
burn/tenotomy tHO mice model as previously reported(*35*). Briefly, mice were anesthetized via
intraperitoneal injection of sodium pentobarbital. Hair was removed from the left hind limb
(from the knee to the ankle) and a 2×3 cm area on the left side of the midline of the spine using
an electric clipper for surgical preparation. A longitudinal incision was made along the Achilles

tendon to expose the gastrocnemius tendon, which was then transected transversely. The skin
was sutured using 5-0 surgical thread. Subsequently, a 2×3×3 cm aluminum block preheated to
60°C in a water bath was applied to the 2×3 cm area on the left side of the midline of the spine
for approximately 17 seconds without additional pressure. The surgical site was disinfected with
an alcohol swab postoperatively, and mice were housed individually. Antibiotics and analgesics
were administered as needed. The success of the tHO model was evaluated 8–10 weeks post-
surgery.
547
2. Micro-CT

After euthanasia, the lower leg and foot of the mice were harvested and fixed overnight at
4°C in 10% formalin solution. Subsequently, the samples were subjected to high-resolution
micro-CT using a SkyScan CT reconstruction software (Bruker Corporation). Three-dimensional
modeling and additional data analysis were performed using 3D.SUITE (Bruker Corporation).
The entire ectopic bone region of the specimen was selected as the region of interest (ROI) for
quantification of bone volume (BV). Additionally, the distal tibial epiphysis of the mouse was
selected as the ROI for quantification of bone mineral density (BMD) and trabecular thickness
(Tb.Th).
557
3. Immunofluorescence

For immunofluorescence staining of tissues and cells, samples were prepared by fixing in
4% paraformaldehyde at 4°C for 24 hours (tissues) or at room temperature for 15 minutes (cells).
Tissues were subsequently deparaffinized and rehydrated through gradient ethanol, while cells
were permeabilized with ice-cold methanol or 0.3% Triton X-100 (Solarbio Life Sciences) as
needed. Both tissues and cells were blocked with 10% normal goat serum or 1% BSA to reduce
nonspecific binding. Primary antibodies, raised in different host species, were applied and
incubated overnight at 4°C. After washing, fluorophore-conjugated secondary antibodies (#A-
11034, Thermo Fisher Scientific) were added and incubated for 1 hour at room temperature in
the dark. Nuclei were stained with DAPI for 10–15 minutes, followed by washing. Sections and
cells were mounted with an anti-fade medium and imaged using Nikon ECLIPSE Ni microscopic
system. Primary antibodies used are as follows: ARL13B (1:500, #17711-1-AP, Proteintech
Group Inc.); Acetyl-α-Tubulin (Lys40) (1:800, #12152, Cell Signaling Technology) ; PDGFRα
(1:100, #sc-338, Santa Cruz); CD31 (1:500, #11265-1-AP, Proteintech Group Inc.); F4/80 (1:500,
#28463-1-AP, Proteintech Group Inc.); TNMD(1:100, #PA5-112767, Invitrogen); α-SMA(1:500,
#14395-1-AP, Proteintech Group Inc.); MEST (1:250, #11118-1-AP, Proteintech Group Inc.).
574
4. Histology and immunohistochemistry

Following euthanasia, the tHO mice tendon tissues were collected and fixed in 4%
paraformaldehyde at 4°C for 24 hours. After three washes with PBS, the samples were
decalcified in 10% EDTA at 4°C for 21 days. The tissues were then embedded in paraffin, and
sagittal sections of the Achilles tendon were obtained via paraffin microtomy. Histological
staining was performed using hematoxylin and eosin (H&E), Masson's trichrome, and Von
Kossa staining to evaluate ectopic ossification in the Achilles tendon. The sections were
dehydrated in ethanol, mounted, and imaged using Nikon ECLIPSE Ni microscopic system.
For immunohistochemistry on paraffin-embedded sections, the slides were first
deparaffinized in xylene and rehydrated through a graded ethanol series. Endogenous peroxidase

activity was quenched with 3% hydrogen peroxide in methanol. Antigen retrieval was performed
by heating the sections in citrate buffer (pH 6.0) for 10 minutes at 95–100°C. Sections were then
blocked with 10% fetal bovine serum or 1% BSA for 1 hour at room temperature. Primary
antibodies were applied and incubated either for 1 hour at room temperature or overnight at 4°C,
followed by washing with PBS. Secondary antibodies (#Ab205718, abcam) were added and
incubated for 30 minutes at room temperature. For visualization, sections were developed with
DAB substrate and counterstained with hematoxylin. Finally, the sections were dehydrated,
cleared, and mounted with neutral gum for microscopic examination using Nikon ECLIPSE Ni
microscopic system. Primary antibodies used are as follows: MEST (1:200, #11118-1-AP,
Proteintech Group Inc.), GLI2 (1:200, #28245-1-AP, Proteintech Group Inc.), PTCH1 (1:100,
#ab53715, abcam), SMO (1:1000, #20787-1-AP, Proteintech Group Inc.).
596

5. Isolation of TSPCs and cell culture

The isolation of TSPCs was conducted as previously reported(*36*). Briefly, 6- to 8-week-old
C57/BL6 mice were sacrificed, and their entire gastrocnemius tendons were aseptically harvested.
The tendons were washed three times with 1X PBS and finely minced using ophthalmic scissors.
The minced tissue was collected in a 15 ml centrifuge tube and digested with 3 ml of 3 mg/ml
collagenase type I (#SCR103, Sigma-Aldrich) at 37°C in a shaking incubator for 45 minutes.
The mixture was then centrifuged at 500 rpm for 5 minutes to remove the collagenase.
Subsequently, the pellet was resuspended in 3 ml of 2 U/ml dispase type II (#4942078001,
Roche) and digested at 37°C for another 45 minutes. After centrifugation at 500 rpm for 1
minute to remove the dispase, the cell pellet was resuspended in 3 ml of low-glucose DMEM
supplemented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin. The cell
suspension was filtered through a 70 μm cell strainer into a 10 cm culture dish and supplemented
with 6-8 ml of complete culture medium. The medium was changed for the first time after 3 days
and then every other day. When the cells reached 80% confluence (7-10 days later), they were
passaged for further experimental use.
612

6. Virus and siRNA construction and transfection

The pLVX-shRNA-puro lentivirus vector carrying shRNA was constructed by Shanghai
OBiO Technology Inc. for targeted gene knockdown in vitro. The pAAV-U6-shRNA-CMV-
MCS-WPRE adeno-associated virus (AAV) vector carrying shRNA was constructed by
Shanghai OBiO Technology Inc. for targeted gene knockdown in vivo. The shRNA sequences
used are *shARL3:* 5'-AAGCTGAATGTATGGGACATT-3'; *shIFT88*: 5'-
GCCCTCAGATAGAAAGACCAA-3'; *shMest*: 5'-CACCGGCCATTGGATCCTAT-3' ;*shNC*:
5'-CCTAAGGTTAAGTCGCCCTCG-3'.
Forty-eight hours after tHO induction in mice, 10 μL of the AAV was injected
intramuscularly into the medial side of the injured hind limb. Subsequent injections were
administered every 3–5 days to achieve sustained knockdown of the specific gene expression.
For lentivirus-mediated transduction in TSPCs, 8×10^4 cells were seeded into each well of a 6-
well plate. When the cell density reached 20%–30%, cells were transduced with lentivirus in the
presence of polybrene (8–10 hours). Subsequently, stable transduced cells were selected using 2
μg/ml puromycin for 5–10 days. The successful establishment of stable cell lines was verified by
quantitative qPCR and Western blot.

The siRNA targeting MEST gene (*siMest-1*: 5'-UGGUCAUCCAGAAUCGACACUGUGG-
3'; *siMest-2*: 5'-GAGGAUCCCAUGGGCUUCUUGAAUG-3') and nontargeting control siRNA
(*siNC*: 5'-UGGGUUCUACAGACGCUACACAUGG-3') was synthesized by Shanghai
GenePharma. For siRNA transfection, 1×10^5 cells were seeded into each well of a 6-well plate.
When the cell density reached 50%, cells were transfected with siRNA using Lipofectamine
RNAiMAX (Invitrogen). Specifically, the siRNA was mixed with Lipofectamine RNAiMAX
and incubated at room temperature for 10 minutes. The mixture was then added to cells cultured
in Opti-MEM reduced-serum medium. After incubation for 48–72 hours, cells were harvested for
subsequent experiments.
638
7.   Induction of TSPCs osteogenic and chondrogenic differentiation

To induce osteogenic differentiation of TSPCs, gelatin-coated 6-well plates were used to
promote cell attachment and growth. TSPCs were seeded at a density of 1×10^4 cells/cm^2 and
cultured in osteogenic induction medium containing 10 mM β-glycerophosphate, 50 μg/ml
ascorbic acid, and 100 nM dexamethasone. The medium was changed every 2-3 days to maintain
optimal conditions. Osteogenic differentiation was assessed by alizarin red staining after 2-4
weeks to visualize calcium deposits. Additionally, qPCR was performed to evaluate the
expression of osteogenesis related genes.
To induce chondrogenic differentiation of TSPCs, cells were seeded at a density of 1×10^4
cells/cm^2 on gelatin-coated 6-well plates and cultured in chondrogenic induction medium
containing 10 ng/ml transforming growth factor-β3 (TGF-β3), 100 nM dexamethasone, 50 μg/ml
ascorbic acid, and 10 mM β-glycerophosphate. The medium was refreshed every 2-3 days to
ensure optimal conditions for differentiation. Chondrogenic differentiation was assessed by
histological staining with alcian blue to detect glycosaminoglycans at 2-4 weeks. qPCR was also
performed to evaluate the expression of chondrogenic markers.
654
8.   Quantitative polymerase chain reaction (qPCR)

To assess the mRNA levels of the target genes, total RNA was purified from TSPCs using
an RNA Easy Fast Tissue/Cell Kit (#DP451, TIANGEN). The extracted mRNA was reverse-
transcribed into cDNA using FastKing gDNA Dispelling RT SuperMix (#KR118, TIANGEN).
Quantitative PCR was performed using the ABI Step One Plus system (Applied Biosystems).
Each reaction mixture (final volume of 20 μL) contained 1 μL of total cDNA, Power SYBR
Green PCR Master Mix (#A57155, Invitrogen), and gene-specific primers. The PCR protocol
included an initial denaturation at 95°C for 10 minutes, followed by 45 cycles of 95°C for 20
seconds, 72°C for 30 seconds, and a final dissociation curve analysis from 60°C to 95°C at a
heating rate of 0.1°C per second. GAPDH was used as an internal control. Relative gene
expression was calculated using the $2^{-\Delta\Delta Ct}$ method. Primers used in the study are listed as follows:
Shh (F: 5'-CTGGCTCGCCTGGCTGTG-3'; R: 5'-CCACGGAGTTCTCTGCTTTCAC-3');
Ptch1 (F: 5'-GGCTGGCTGGCGTCCTG-3'; R: 5'-CATCATCCACACCAACACCAAGAG-3');
Alpl (F: 5'-CACGGCGTCCATGAGCAGAAC-3'; R: 5'-CAGGCACAGTGGTCAAGGTTGG-
3'); *Runx2* (F: 5'-TCCCAGGCAGGCACAGTCTTC-3'; R: 5'-AGCGGCGTGGTGGAGTGG-
3'); *Bglap* (F: 5'-AGGGCAGCGAGGTAGTGAAGAG-3'; R: 5'-
GCCGATGTGGTCAGCCAACTC-3'); *Sp7* (F: 5'-CCTCTGCGGGACTCAACAAC-3'; R: 5'-
AGCCCATTAGTGCTTGTAAAGG-3'); *Col2a1* (F: 5'-GGTCCTCCTGGTCCTGGCATC-3';
R: 5'-CGTGCTGTCTCAAGGTACTGTCTG-3'); *Sox9* (F: 5'-

ATGACCGACGAGCAGGAGAAGG-3'; R: 5'-CCGAGGGACAGGGCGAACC-3'); *Acan* (F:
5'-GGAGACCCAGACAGCAGAAACAAC-3'; R: 5'-GCAGGTGGCTCCATTCAGACAAG-
3'); *Ift88* (F: 5'-TTCAGCAAGCAGTGAGAACCAGTC-3'; R: 5'-
TGTCCCTGTCATCGGTCTTCCC-3'); *Arl3* (F: 5'-TGTGTGCCCGTGCTCATCTTTG-3'; R:
5'-TGGACGCCCTCGCCTGTG-3'); *Gli2* (F: 5'-GCTCCACACACCCGCAACAC-3'; R: 5'-
CTCAGCATCGTCACTTCGGTCAG-3'); *Mest* (F: 5'-CTTAGGCTTTGGCTTCAGTGACAA-
3'; R: 5'-AGGTTGATTCTGCGGTTCTGGAG-3');
681
9. Western blot

To perform western blot analysis, proteins were extracted from cells by washing with PBS,
lysing in RIPA buffer (#R0278, Sigma-Aldrich) containing Protease & Phosphatase Inhibitor
Cocktail (#78441, Thermo Fisher Scientific), and sonicating on ice. The lysates were centrifuged
at 4°C to collect the supernatant, and protein concentrations were determined using a BCA
Protein Assay Kit (#P0011, Beyotime). Equal amounts of protein were denatured by heating in
SDS loading buffer, separated by SDS-PAGE, and transferred to PVDF or nitrocellulose
membranes. Membranes were blocked with 5% skim milk in TBST to reduce nonspecific
binding. Primary antibodies were applied overnight at 4°C, followed by washing and incubation
with secondary antibodies conjugated to horseradish peroxidase (#SA00001-2, Proteintech Inc.)
for 1 hour at room temperature. After additional washing, the membranes were developed using
Super ECL Detection Reagent (#36208ES60, YEASEN) and imaged with ChemiDoc XRS+ Gel
Imaging System (Bio-Rad). Band intensities were quantified using ImageJ software, and target
protein expression levels were normalized to those of the internal control. Primary antibodies
used in western blot are as follows: MEST (1:1000, #11118-1-AP, Proteintech Group Inc.), GLI2
(1:2000, #28245-1-AP, Proteintech Group Inc.), PTCH1 (1:1000, #ab53715, abcam), SHH
(1:2000, #20697-1-AP, Proteintech Group Inc.), IFT88 (1:6000, #13967-1-AP, Proteintech
Group Inc.), ARL3 (1:1000, #10961-1-AP, Proteintech Group Inc.), PI3K (1:1000, #ab191606,
abcam), p-PI3K (1:2000, #ab278545, abcam), AKT (1:2000, #ab8805, abcam), p-AKT (1:1000,
#ab38449, abcam).
702
10. RNA-seq

Total RNA was isolated from samples using RNA Easy Fast Tissue/Cell Kit (#DP451,
TIANGEN). The purity and concentration of the extracted RNA were assessed using the
NanoDrop One spectrophotometer (Thermo Fisher Scientific). RNA integrity was further
evaluated using the Agilent 2100 Bioanalyzer (Agilent Technologies). Subsequently, RNA
sequencing libraries were prepared using the VAHTS Universal V6 RNA-seq Library Prep Kit
(Vazyme). The transcriptome sequencing and subsequent bioinformatics analysis were
performed by Shanghai OE Biotech. Sequencing libraries were sequenced on the Illumina
Novaseq 6000 platform to generate 150-bp paired-end reads. The raw sequencing reads in
FASTQ format were processed using the fastp software to remove low-quality reads, yielding
clean reads for subsequent data analysis. The clean reads were aligned to the reference genome
using the HISAT2 software, and gene expression levels were calculated in terms of fragments
per kilobase of transcript per million mapped reads (FPKM). Additionally, read counts for each
gene were obtained using HTSeq-count. Subsequent analyses, including principal component
analysis (PCA), differential gene expression analysis, and enrichment analysis, were performed
using the R software.

11. Untargeted lipidomics using liquid chromatography-tandem mass spectrometry (LC-MS/MS)

TSPCs were cultured in serum-free medium to promote cilium growth until confluence. The cells were washed with PBS and subjected to shear force by rotating at 360 rpm in a shaking incubator at 37°C for 4 minutes to detach cilia as previously reported(*37*). The cell debris was removed by centrifugation at 1000×g for 10 minutes at 4°C, followed by ultracentrifugation at 40,000×g for 30 minutes at 4°C to isolate the ciliary suspension from the supernatant. The ciliary suspension was then used for untargeted lipidomics sequencing.

For lipid extraction, 200 µL of pre-chilled methanol/water (3:1, v/v) was added to the ciliary suspension, followed by sonication on ice for 15 minutes. Then, 1 mL of pre-chilled methyl tert-butyl ether (MTBE) was added and vortexed thoroughly. The mixture was rotated at 4°C for 1 hour, followed by sonication on ice for another 30 minutes. After adding 100 µL of water and vortexing for 1 minute, the mixture was left at room temperature for 10 minutes. The sample was then centrifuged at 14,000×g for 15 minutes at 4°C. The supernatant was collected, and the pellet was dried under a stream of nitrogen. The dried pellet was re-dissolved in 200 µL of SDT buffer. Protein concentration was measured using the BCA assay (Table 4-2). Based on the protein content, the cell quantity in each sample was estimated, and equal cell amounts were taken from each supernatant. The supernatant was dried using a high-speed vacuum concentrator, re-dissolved in 60 µL of pre-chilled isopropanol/methanol (1:1, v/v), and centrifuged at 20,000×g for 20 minutes at 4°C. The supernatant was vortexed and transferred to an injection vial for analysis. The entire analysis was performed at 4°C using an automatic sampler. The samples were analyzed using a SHIMADZU-LC30 ultra-high-performance liquid chromatography (UHPLC) system equipped with an ACQUITY UPLC® HSS C18 column (2.1×100 mm, 1.9 µm) (Waters, Milford, MA, USA). To avoid signal drift, samples were analyzed in a random order. A quality control (QC) sample was inserted every 5-10 experimental samples to monitor system stability and data reliability. The samples were analyzed in both positive and negative ion modes using electrospray ionization (ESI). After separation by UHPLC, the samples were analyzed using a QExactive Plus mass spectrometer (Thermo Fisher Scientific). Lipid identification and quantification were performed using the MSDIAL software (Version 5.2.240218.2-net472) with the following parameters: precursor mass tolerance of 10 ppm, product mass tolerance of 10 ppm, retention time alignment of 0.1 min, and peak area filtering to reduce false positives. Lipids with relative standard deviations (RSD) exceeding 30% in QC samples were excluded. The data were normalized to the total peak area, preprocessed with UV-scaling, and subjected to multivariate statistical analyses, including unsupervised principal component analysis (PCA), supervised partial least squares discriminant analysis (PLS-DA), and orthogonal partial least squares discriminant analysis (OPLS-DA). Univariate statistical analyses included Student's t-test and fold-change analysis, and volcano plots and hierarchical clustering were generated using R software.

12. Dual-luciferase reporter assay

The Dual-Luciferase Reporter Assay was performed using the Dual-Luciferase Reporter Assay System (Promega). The firefly luciferase reporter gene was cloned downstream of the wild-type (WT) or mutant (MUT) pGL4.10-Mest promoter regions and co-transfected into TSPCs with pcDNA3.1-GLI2 and Renilla expression vector pRL-TK. Following a 48-hour

incubation, cell lysates were prepared, and firefly luciferase activity was quantified and normalized to Renilla luciferase activity as an internal control.

13. Bioinformatics analysis

The scRNA sequencing dataset pertaining to tHO in mice was retrieved from the GEO database (GSE126060). Dimensionality reduction, clustering, and cell-type identification of the scRNA sequencing data were performed using the Seurat package in R software. Additionally, bulk RNA sequencing data from the heterotopic ossification sites of mice three weeks post-trauma was also obtained from the GEO database (GSE233201). Differential gene expression analysis was conducted using the DESeq2 package, while enrichment analysis was performed with the clusterProfiler package. Transcription factor binding sites were predicted using the JASPAR database (https://jaspar.elixir.no/).

14. Statistical analysis

All statistical analyses were performed using SPSS software and R Studio. Data are presented as mean ± standard deviation (SD) of three independent assays. Statistical analyses were performed by Student's t-test for two-group comparison and one-way ANOVA analyses with Tukey's post hoc test for multiple comparisons. Statistical significance was set at $P < 0.05$.

List of Supplementary Materials

Fig. S1 to S8 for multiple supplementary figures

Table S1 to S2 for multiple supplementary tables (Excel files)

References and Notes

1. D. Dey *et al.*, The traumatic bone: trauma-induced heterotopic ossification. *Transl Res* **186**, (2017).
2. K. Ranganathan *et al.*, Heterotopic Ossification: Basic-Science Principles and Clinical Correlates. *The Journal of Bone and Joint Surgery. American Volume* **97**, 1101-1111 (2015).
3. P. Mill, S. T. Christensen, L. B. Pedersen, Primary cilia as dynamic and diverse signalling hubs in development and disease. *Nature Reviews. Genetics* **24**, 421-441 (2023).
4. Z. Anvarian, K. Mykytyn, S. Mukhopadhyay, L. B. Pedersen, S. T. Christensen, Cellular signalling by primary cilia in development, organ function and disease. *Nature Reviews. Nephrology* **15**, 199-219 (2019).
5. M. Spasic, M. P. Duffy, C. R. Jacobs, Fenoldopam Sensitizes Primary Cilia-Mediated Mechanosensing to Promote Osteogenic Intercellular Signaling and Whole Bone Adaptation. *J Bone Miner Res* **37**, 972-982 (2022).
6. K. He *et al.*, Primary cilia mediate skeletogenic BMP and Hedgehog signaling in heterotopic ossification. *Sci Transl Med* **16**, eabn3486 (2024).
7. C. D. Hwang *et al.*, Contemporary perspectives on heterotopic ossification. *JCI Insight* **7**, (2022).
8. Y. Bi *et al.*, Identification of tendon stem/progenitor cells and the role of the extracellular matrix in their niche. *Nat Med* **13**, 1219-1227 (2007).
9. G. Dai *et al.*, Higher BMP Expression in Tendon Stem/Progenitor Cells Contributes to the Increased Heterotopic Ossification in Achilles Tendon With Aging. *Front Cell Dev Biol* **8**, 570605 (2020).
10. Y. Chen *et al.*, Targeted pathological collagen delivery of sustained-release rapamycin to prevent heterotopic ossification. *Sci Adv* **6**, eaay9526 (2020).
11. P. Dutta, K. Ray, Ciliary membrane, localised lipid modification and cilia function. *Journal of Cellular Physiology* **237**, 2613-2631 (2022).
12. C. Iwaya, A. Suzuki, J. Iwata, Loss of Sc5d results in micrognathia due to a failure in osteoblast differentiation. *J Adv Res* **65**, 153-165 (2024).
13. A. Suzuki *et al.*, Disruption of Dhcr7 and Insig1/2 in cholesterol metabolism causes defects in bone formation and homeostasis through primary cilium formation. *Bone Research* **8**, 1 (2020).

14. L. Nikonova *et al.*, Mesoderm-specific transcript is associated with fat mass expansion in response to a
positive energy balance. *FASEB Journal : Official Publication of the Federation of American Societies For*
Experimental Biology **22**, 3925-3937 (2008).
15. A. K. Huber *et al.*, Immobilization after injury alters extracellular matrix and stem cell fate. *The Journal of*
Clinical Investigation **130**, 5444-5460 (2020).
16. K. Noda, M. Kitami, K. Kitami, M. Kaku, Y. Komatsu, Canonical and noncanonical intraflagellar transport
regulates craniofacial skeletal development. *Proc Natl Acad Sci U S A* **113**, E2589-2597 (2016).
17. H. Feng *et al.*, Tendon-derived cathepsin K-expressing progenitor cells activate Hedgehog signaling to
drive heterotopic ossification. *The Journal of Clinical Investigation* **130**, 6354-6365 (2020).
18. S. Chen *et al.*, A CD26+ tendon stem progenitor cell population contributes to tendon repair and
heterotopic ossification. *Nat Commun* **16**, 749 (2025).
19. J. B. Regard *et al.*, Activation of Hedgehog signaling by loss of GNAS causes heterotopic ossification. *Nat*
Med **19**, 1505-1512 (2013).
20. K. Ha *et al.*, Cilia-enriched oxysterol 7β,27-DHC is required for polycystin ion channel activation. *Nat*
Commun **15**, 6468 (2024).
21. D. Yamakawa *et al.*, Primary cilia-dependent lipid raft/caveolin dynamics regulate adipogenesis. *Cell*
Reports **34**, 108817 (2021).
22. M. Pekkinen *et al.*, Osteoporosis and skeletal dysplasia caused by pathogenic variants in SGMS2. *JCI*
Insight **4**, (2019).
23. G. Luchetti *et al.*, Cholesterol activates the G-protein coupled receptor Smoothened to promote Hedgehog
signaling. *ELife* **5**, (2016).
24. M. Kinnebrew *et al.*, Cholesterol accessibility at the ciliary membrane controls hedgehog signaling. *ELife* **8**,
(2019).
25. S. P. Taylor *et al.*, Mutations in DYNC2LI1 disrupt cilia function and cause short rib polydactyly syndrome.
Nat Commun **6**, 7092 (2015).
26. H. Chen *et al.*, The effects of primary cilia-mediated mechanical stimulation on nestin+-BMSCs during
bone-tendon healing. *J Adv Res*, (2024).
27. D. Tao *et al.*, Primary cilia support cartilage regeneration after injury. *Int J Oral Sci* **15**, 22 (2023).
28. X. Liu *et al.*, Numb positively regulates Hedgehog signaling at the ciliary pocket. *Nat Commun* **15**, 3365
(2024).
29. B. A. Allard *et al.*, Thm2 interacts with paralog, Thm1, and sensitizes to Hedgehog signaling in postnatal
skeletogenesis. *Cell Mol Life Sci* **78**, 3743-3762 (2021).
30. Q. Cong *et al.*, A self-amplifying loop of YAP and SHH drives formation and expansion of heterotopic
ossification. *Sci Transl Med* **13**, (2021).
31. X. Yuan, M. Liu, X. Cao, S. Yang, Ciliary IFT80 regulates dental pulp stem cells differentiation by
FGF/FGFR1 and Hh/BMP2 signaling. *International Journal of Biological Sciences* **15**, 2087-2099 (2019).
32. V. Jou, S. M. Peña, J. A. Lehoczky, Regeneration-specific promoter switching facilitates Mest expression
in the mouse digit tip to modulate neutrophil response. *NPJ Regen Med* **9**, 32 (2024).
33. W. Lu *et al.*, Hedgehog signaling regulates bone homeostasis through orchestrating osteoclast
differentiation and osteoclast-osteoblast coupling. *Cell Mol Life Sci* **80**, 171 (2023).
34. Y. Wang *et al.*, TMEM216 promotes primary ciliogenesis and Hedgehog signaling through the SUFU-
GLI2/GLI3 axis. *Sci Signal* **17**, eabo0465 (2024).
35. J. R. Peterson *et al.*, Direct Mouse Trauma/Burn Model of Heterotopic Ossification. *J Vis Exp*, e52880
(2015).
36. M. Chen *et al.*, Noncanonical Wnt5a signaling regulates tendon stem/progenitor cells senescence. *Stem Cell*
Res Ther **12**, 544 (2021).
37. D. R. Raleigh *et al.*, Cilia-Associated Oxysterols Activate Smoothened. *Mol Cell* **72**, (2018).
858

859 **Acknowledgments:** We thank Airu Zhu at Shanghai Bioprofile Technology CompanyLtd for
860 her technical support in mass spectroscopy.

861 **Funding:** National Natural Science Foundation of China 82302084 (HJ)

862 **Author contributions:**
863 Conceptualization: HJ, XHZ

864   Methodology: BWL, JQZ
865   Investigation: BWL, YG
866   Visualization: BWL, ZLS
867   Funding acquisition: HJ
868   Project administration: HJ, XHZ
869   Supervision: JQZ
870   Writing – original draft: BWL, YG
871   Writing – review & editing: HJ, XHZ

872 **Competing interests:** Authors declare that they have no competing interests.

877 **Ethical approval :** All animal procedures were conducted in accordance with the guidelines of
878   the Ethics Committee on Animal Experiments of the Naval Medical University.

879 **Figures**

Figure 1. Primary cilia defects rescue the ecotopic bone formation of tHO. (a) Illustration of establishment of
the tHO burn/tenotomy mouse model. (b) HE, Masson trichrome, and Von Kossa staining of tHO tendon tissues
collected from differnet time points after injury (scale bar 50 um). (c) Immunofluorescence images of PDGFRα
(green) and ARL13B (red) at differnet time points during tHO development. DAPI staining indicates nuclei (blue),
scale bar 50 um. (d) The percentage of ARL13B+ cells at differnet time points after injury are shown. Data are
presented as means ± SD (n= 3). (e) The percentage of ARL13+ cells in PDGFRα+ cells are shown (n= 3). (f) The
percentage of ARL13+ cells in five types of cells (CD31+, F4/80+, PDGFRα+, TNMD+, and α-SMA+ cells) are
shown (n= 3). (g) Micro-CT images show tHO of lower limbs among *shNC, shIFT88*, and *shARL3* groups (n=
4/group). Bone volume (BV, mm^3), bone mineral dendity (BMD, g/cc), and Tb.Th (trabecular thickness, mm) are
quantified. Data are presented as means ± SD of three independent assays. Statistical analyses were performed by
one-way ANOVA analyses with Tukey's post hoc test for multiple comparisons. **P < 0.01; ns, P > 0.05.

892
Figure 2. scRNA-seq indicated the potential role of cilia and stem/progenitor cells in tHO development. (a)
UMAP plot of nine clusters identified in 21880 cells in total of tHO site. (b) Bubble plot shows the high variable
marker genes of each cluster. (c) The expression density of Pdgfra+, Prrx1+, and Pdgfra+Prrx1+ cells in each cluster
are shown by UMAP plot. (d) UMAP plot of nine clusters with single cells grouped by different tim points (Day0 ,3,
7, 21) during tHO demonstrates the trend of changes in cell composition. (e) The volcano plot visualizes the
differential expressed genes in the Mesenchymal subcluster at Day 7 (left) or Day 21 (right) compared to the Day 0
group. (f) The dot plot visualizes the different expression of *Ift20* and *Arl3* in nine clusters. (g) UMAP plot shows
the expression of *Ift20* (upper) and *Arl3* (lower) at different time points (Day0 ,3, 7, 21) during tHO development.

901
Figure 3. TSPCs osteochondrogenic differentiation is regulated by primary cilia in tHO. (a) Fibroblast-like
TSPCs extracted from mice for subsequent experiments (scale bar 100 um). (b) Flow cytometry analysis of the
expression of Pdgfra and Prrx1 (stem/progenitor cell surface markers) on TSPCs. (c) Immunofluorescence images of
ARL13B (red) and α-tubulin (green) mark the cilia in *shNC, shIFT88,* and *shARL3* TSPCs. DAPI staining indicates
nuclei (blue), scale bar 10 um (upper) and 30 um (lower). (d) The percentage of ciliated TSPCs marked by ARL13B
show the cilia counts in the control group and ciliary gene knockdown groups. (e) Cilia length in TSPCs with or
without *shIFT88* or *shARL3* treatment (n= 25/group). (f) Representative images of alizarin red staining (upper) and
alcian blue staining (lower) in TSPCs with or without *shIFT88* or *shARL3* treatment after 3 weeks of in vitro
osteogenic or chondrogenic differentiation. (g) qPCR showed relative osteogenesis-related genes (*Alpl, Runx2,
Bglap, Sp7*) and chondrogenesis-related genes (*Col2a1, Sox9, Acan*) mRNA expression in TSPCs with or without
shIFT88 or *shARL3* treatment after 3 weeks of in vitro osteogenic or chondrogenic differentiation. Data are
presented as means ± SD of three independent assays. Statistical analyses were performed by one-way ANOVA
analyses with Tukey's post hoc test for multiple comparisons. *P < 0.05; **P < 0.01; ***P < 0.005; ****P < 0.001.
915   ns, P > 0.05.

916
Figure 4. Primary cilia contribute to tHO pathogenesis by regulating MEST expression. (a) The volcano plot
visualizes the differential expressed genes in injury site of the tHO mice 3 weeks after burn/tenotomy compared to
the control mice. (b) RNA-seq performed between *shIft88* and *shNC* TSPCs after 3-7 days of in vitro osteogenic
differentiation and differential expressed genes are visualized. (c) Venn diagram shows the common differential
expressed genes among three genesets and identified MEST, PTN, COL2A1 as the key genes. (d) The expression
density of *Mest* in each cluster are shown by violin plot and UMAP plot. (e) qPCR shows relative *Mest* mRNA
expression in TSPCs with or without *shIFT88* or *shARL3* treatment during in vitro osteogenic (left) or chondrogenic
(right) differentiation. (f) Western blot shows MEST protein expression levels in TSPCs with or without *shIFT88* or
shARL3 treatment during in vitro osteogenic (left) or chondrogenic (right) differentiation. (g) UMAP plot shows the
expression of *Mest* at different time points (Day0 ,3, 7, 21) during tHO development. (h) Immunohistochemistry
images demonstrates MEST in tHO tendon tissues at different time post burn/tenotomy injury (scale bar 100 um). (i)
Immunofluorescence images of α-tubulin (green) and MEST (red) in mouse primary TSPCs. DAPI staining
indicates nuclei (blue), scale bar 10 um. Data are presented as means ± SD of three independent assays. Statistical
analyses were performed by one-way ANOVA analyses with Tukey's post hoc test for multiple comparisons. *$P <$
0.05; **$P < 0.01$; ***$P < 0.005$; ****$P < 0.001$.

932
Figure 5. MEST modulates TSPCs cilia length and osteochondral differentiation in tHO. (a) Representative
images of alizarin red staining (upper) and alcian blue staining (lower) in TSPCs with or without *siMest* treatment
after 3 weeks of in vitro osteogenic or chondrogenic differentiation. (b) qPCR shows relative *Mest* and osteogenic
differentiation-related (*Alpl, Runx2, Bglap*) or chondrogenic differentiation-related genes (*Col2a1, Sox9, Acan*)
mRNA expression in TSPCs with or without *siMest* treatment after 3 weeks of in vitro osteogenic or chondrogenic
differentiation. (c) tSNE map shows Mest-high and Mest-low subclusters divided from Mesenchymal cluster. The
expression level of *Mest, Runx2, Sox9*, and *Acan* in Mest-high and Mest-low subclusters were visualize by violin
plots. (d) Immunofluorescence images of α-tubulin (green) mark the cilia of TSPCs. DAPI staining indicates nuclei
(blue), scale bar 10 um. Ciliated TSPCs proportion (cilia counts) marked by α-tubulin and cilia length (n= 25/group)
in the *siNC* and *siMest* group. (e) Micro-CT images show tHO of lower limbs in *shNC* and *shMest* groups (n=
4/group). Quantification of tHO degree by bone volume (BV, mm^3) and bone mineral dendity (BMD, g/cc). Data
are presented as means ± SD of three independent assays. Statistical analyses were performed by Student's t-test for

947
Figure 6. TSPCs ciliary Hh and PI3K-Akt signaling are involved in tHO progression. (a) GO enrichment
anayslis of differential expressed genes in tHO mice injury site shows top 10 enriched functions that are important
for tHO. (b) GO enrichment analysis of differential expressed genes in TSPCs treated with *shIft88* shows top 10
enriched biological processes. (c) GSEA enrichment of TSPCs treated with *shIft88* or *shNC*. The enriched pathway
with the absolute value of normalized enrichment score (NES) greater than 1 and P-value less than 0.05 was
considered significant. (d) KEGG enrichment analysis of differential expressed genes in Mesenchymal subcluster of
tHO injury site identified by scRNA-seq shows top 10 enriched pathways. (e) KEGG enrichment analysis of
differential expressed genes in TSPCs treated with *shIft88* showed top enriched pathways. (f) qPCR showed relative
Hh signaling genes (Shh, Ptch1) mRNA expression in TSPCs with or without *shIft88* or *shARL3* treatment after 3

weeks of in vitro osteogenic or chondrogenic differentiation. (g) Western blot showed Hh signaling (SHH, PTCH1)
and PI3K-Akt signaling (PI3K, p-PI3K, AKT, p-AKT) protein expression levels in TSPCs with or without *shIFT88*
or *shARL3* treatment during in vitro osteogenic or chondrogenic differentiation. (h) The expression level of *Smo* in
each cluster are shown by violin plot. (i) UMAP plot shows the expression of *Smo* at different time points (Day0 ,3,
7, 21) during tHO development. Data are presented as means ± SD of three independent assays. Statistical analyses
were performed by one-way ANOVA analyses with Tukey's post hoc test for multiple comparisons. ***$P < 0.005$;
****$P < 0.001$.

Figure 7. A self-amplifying loop of GLI2 and MEST drives TSPCs differentiation and tHO progression via Hh signaling. (a) Western blot shows Hh signaling (SHH, PTCH1) protein expression levels in TSPCs with or without *siMest* treatment during in vitro osteogenic or chondrogenic differentiation. (b) Western blot showed PI3K-Akt signaling (PI3K, p-PI3K, AKT, p-AKT) protein expression levels in TSPCs with or without *siMest* treatment

durig in vitro osteogenic or chondrogenic differentiation. (c) The sequence logo of GLI2 transcription factor and two potential binding sites of the GLI2 transcription factor on the MEST promoter region predicted by the JASPAR database. (d) The luciferase reporter assay using pGL4.10 as vector showed that GLI2 can bind to the MEST promoter at both site 1 and 2. (e) Immunohistochemistry images demonstrats GLI2 in tHO tendon tissues at different time post burn/tenotomy injury (scale bar 100 um). (f) qPCR shows relative *Gli2* mRNA expression in TSPCs with or without *shIft88* or *shARL3* treatment after 3 weeks of in vitro osteogenic or chondrogenic differentiation. (g) Western blot shows GLI2 protein expression levels in TSPCs with or without *shIFT88* or *shARL3* treatment during in vitro osteogenic or chondrogenic differentiation. (h) Immunohistochemistry images demonstrats GLI2, PTCH1, and SMO expression in tHO tendon tissues with or without shMest treatment after injury (scale bar 150 um). Data are presented as means ± SD of three independent assays. Statistical analyses were performed by one-way ANOVA analyses with Tukey's post hoc test for multiple comparisons. **P < 0.01; ****P < 0.001.

Figure 8. MEST regulates ciliary lipid metabolism in TSPCs. (a) Immunofluorescence images of α-tubulin (green) marked cilia in ciliated and non-ciliated mouse primary TSPCs after cilia isolation. DAPI staining indicates nuclei (blue). (b) OPLS-DA plot shows the differences between NC (blue, n= 7) and SIM (green, n= 6) groups. (c) Circular diagram of the classification statistics of differential lipids. (d) Bubble chart of lipid classification. The horizontal axis represents the quantity range of lipids, and the vertical axis represents the logarithmic transformation of fold change. Each circle represents a lipid substance. The larger the circle, the higher the variable importance.

Different colors represent the primary classification, and lipids of the same classification are arranged together. (e)
The volcano plot visualizes the differential expressed lipids in SIM group compared to the NC group. (f) Relative
abundance of sphingomyelin and triglyceride are significantly different in SIM and NC group. (g) KEGG
enrichment analysis of differential expressed lipids between SIM and NC group (C: cellular processes; E:
environmental information processing; H: human diseases; M: metabolism; O:organismal systems). (h) The
expression density of *Sgms2* in each cluster (left) and in Mesenchymal subcluster (right) are shown by UMAP plot.
(i) UMAP plot shows the expression of *Sgms2* in Mesenchymal subcluster at different time points (Day0 ,3, 7, 21)
during tHO development.

995
Figure 9. Targeting MEST or ciliary sphingomyelin in TSPCs may alleviate tHO via Hh signaling. (a)
Representative images of alizarin red staining (upper) and alcian blue staining (lower) in TSPCs after 3 weeks of in
vitro osteogenic (upper) and chondrogenic (lower) differentiation in control, myriocin, sphingomyelin,
myriocin+sphingomyelin, myriocin+siMest, and myriocin+vismodegib treated groups. (b) Quantification of the
percentages of (alizarin red) ARS+ and alcian blue (AB+) areas in six groups. (c) qPCR showed relative osteogenic
differentiation-related genes *(Alpl, Runx2, Bglap)* and chondrogenic differentiation-related genes (*Sox9, Col2a1,*

Acan) mRNA expression in TSPCs after 3 weeks of in vitro osteogenic (upper) or chondrogenic (lower)
differentiation in six groups. (d) qPCR showed relative Hh signaling genes (*Shh, Ptch1, Gli2*) mRNA expression in
TSPCs after 3 weeks of in vitro osteogenic differentiation in six groups. (e) Micro-CT images show tHO of lower
limbs between 1X PBS and sphingomyelin treated groups (n= 4/group). Quantification of tHO degree by bone
volume (BV, mm^3) and bone mineral dendity (BMD, g/cc). Data are presented as means ± SD of three independent
assays. Statistical analyses were performed by one-way ANOVA analyses with Tukey's post hoc test for multiple
comparisons. *P < 0.05; **P < 0.01; ***P < 0.005; ****P < 0.001. ns, P > 0.05.

1009
Figure 10. Schematic diagram of the study. After trauma, the level of SHH within the primary cilia of TSPCs is
significantly elevated. This increase in SHH activates the Hh signaling through its interaction with PTCH1 and SMO
receptors. Subsequently, the GLI2 transcription factor is activated and translocates to the nucleus, where it promotes
the expression of the MEST gene. Concurrently, the phosphorylation levels of PI3K and AKT are enhanced. The
upregulation of MEST expression leads to a reduction in sphingomyelin within the cilia, further amplifying the
activation of the ciliary Hh signaling. This positive feedback loop dramatically enhances the osteogenic and
chondrogenic differentiation potential of TSPCs, resulting in localized ectopic bone deposition and tHO. Conversely,
when the ciliary genes Ift88 and Arl3 are knocked down in TSPCs, the production of SHH ligands is markedly
reduced. Thus, SMO is inhibited by PTCH1, preventing the activation of GLI2. Additionally, the phosphorylation of
PI3K and AKT is suppressed. The reduction in MEST expression subsequently increases the sphingomyelin within
the cilia, which further inhibits the ciliary Hh signaling. As a result, the osteogenic and chondrogenic differentiation
capacity of TSPCs is significantly diminished, thereby alleviating the occurrence and development of tHO.