Title

Amyloid beta is released by lysosomal exocytosis from hiPSC-derived neurons

Authors

Adrianna R Tsang[1,2], Manoj Reddy Medapati[2], Claudia Seah[2], Jordan M Krupa[2,3],

Stephen H. Pasternak[1,2,3,4*]

[1]Department of Physiology and Pharmacology, Schulich School of Medicine and

Dentistry, Western University, London, Ontario, Canada

[2]Robarts Research Institute, Schulich School of Medicine and Dentistry, Western

University, London, Ontario, Canada

[3]Neuroscience, Schulich School of Medicine and Dentistry, Western University, London,

Ontario, Canada

[4]Department of Clinical Neurological Sciences, Schulich School of Medicine and

Dentistry, Western University, London, Ontario, Canada

[*]Corresponding Author

spasternak@robarts.ca

RRI 3243, Robarts Research Institute

Schulich School of Medicine & Dentistry

Western University

1151 Richmond St. N.

London, ON N6A 5B7

Tel. 519-931-5294

Declarations of Interest: Zywie Bio LLC has provided grant support to the institution.

The PI is a shareholder and patent holder in Zywie Bio LLC.

Abstract

The endosomal-lysosomal system has long been linked to the production of amyloid beta

(Aβ), but the specific intracellular compartments involved in Aβ secretion remain

contentious. While lysosomes are typically associated with Aβ degradation, studies have

also shown that lysosomes also serve as site of Aβ accumulation in cells, mouse models,

and human tissues. Lysosomal exocytosis is a major secretory pathway in non-neuronal

cells, but few studies have investigated this pathway in neurons. Here, we examined the

potential role and mechanism of lysosomal exocytosis in human induced pluripotent stem

cell (hiPSC)-derived neurons, and we hypothesized that lysosomal exocytosis is a

pathway for Aβ secretion from these neurons. Using total internal reflection fluorescence

(TIRF) microscopy, lysosomes filled with fluorescently labelled amyloid were seen

approaching and fusing with the plasma membrane in real-time. The number and

composition of the released particles were characterized using nanoscale flow cytometry.

Silencing two proteins, Rab27b and munc13-4, significantly reduced these events and

blocked the release of amyloid into the extracellular space. Our results provide direct

evidence for the involvement of lysosomal exocytosis in the release of Aβ from neurons

and highlight its potential as a target for therapeutic intervention in Alzheimer's disease.

Keywords: amyloid beta, lysosomes, exocytosis, Alzheimer's disease, hiPSC-derived

neurons

Introduction

Amyloid deposition is one of the pathological hallmarks of Alzheimer's disease (AD),

the most common age-related neurodegenerative disorder worldwide. Although the exact

cause of AD pathology is unknown, it is widely accepted that the accumulation of

aggregated amyloid beta (Aβ) and tau-containing neurofibrillary tangles are central to

AD. Amyloidogenic processing of Aβ by sequential cleavages from β-secretase BACE1

and γ-secretase leads to the generation of toxic Aβ peptides, $A\beta_{40}$ and $A\beta_{42}$.[1] The

amyloidogenic processing of the amyloid precursor protein (APP) is believed to occur

after APP has been internalized into the endosomal-lysosomal system. Our group has

shown that APP can be transported directly to lysosomes and processed into Aβ.[2–4]

Lysosomes are acidic compartments that are typically viewed as end state organelles

responsible for degradation.[5] Lysosomal dysfunction and the accumulation of

intralysosomal Aβ have been implicated in progression of AD.[6] Perhaps the earliest

connection between lysosomes and AD was over 60 years ago when high levels of

phosphatase-positive lysosomal dense bodies were localized within amyloid plaques of

AD brains.[7] Early electron microscopy studies also revealed swollen neurites filled with

lysosome-like organelles surrounding Aβ plaques in the brains of AD patients.[8] These

initial findings have been supported by identification of enlarged or swollen lysosomes in

neurons from post-mortem AD brain samples and mouse models of AD.[9–12]

In addition to digesting and recycling material, lysosomes are dynamic compartments that

can alter their position in the cell to sense and respond to cellular needs in order to

maintain homeostasis.[13,14] Lysosomes also serve as secretory organelles undergoing Ca^{2+}

dependent exocytosis in many cell types.[13,15–18] For example, lysosomes function as

major secretory granules in many haematopoietic cells, as well as osteoclasts and

melanocytes.[16,19–26] Cytotoxic granules in T-cells, basophil cell granules in basophils,

azurophil granules in neutrophiles, and platelet dense in platelets are also key examples

of cell-specific secretory lysosomes. The presence of secretory lysosomes in neurons and

astrocytes has also been reported.[27–33] As neuronal exocytosis of lysosomes is still an

emerging field, most of our knowledge of lysosomal exocytosis has been studied in non-

neuronal cells, including cargo, function and machinery. Exocytosis of lysosomes

requires the recruitment of specific proteins which mediate their movement, tethering,

and fusion to the plasma membrane.

Lysosomal exocytosis typically begins with a rise in intracellular Ca^{2+} that is detected by

a Ca^{2+} sensor, synaptotagmin VII (SytVII).[20,21,27,34] This triggers the mobilization of the

lysosome toward the cell surface via a series of kinesin, actin and myosin proteins[27,35–37]

At the cell surface, the lysosome tethers and fuses with the plasma membrane through

specific docking and fusion machinery, such as Soluble *N*-Ethylmaleimide-Sensitive

Factor Attachment Receptor (SNARE) complexes and small GTPases of the Ras analog

in brain (Rab) family.[15,16,26,27,35,38,39] The fusion of lysosomal and plasma membranes

results in the release of the original luminal cargo into the extracellular space. Many

proteins have been implicated in this fusion, including v-SNARE, VAMP7, t-SNAREs,

SNAP23 and syntaxin-4, and the small GTPases, Rab3 and Rab27.[15,23,35,39] While none of

these are unique to lysosomal exocytosis, Rab3 and Rab27 are the two so-called

"secretory Rabs" and are present in neurons.[40–43] Of these, Rab27 has been shown to be

more stably localized at vesicle membranes.[44,45] Rab27 exists as two isoforms, Rab27a

and Rab27b, in which Rab27b displays higher localized expression in the central nervous

system (CNS).[39,46] In neurons, Rab27b targets and recruits its effector protein unc-13

homolog D (munc13-4), which plays a key role in priming of exocytic vesicles by

interacting with SNARE complexes.[43]

As lysosomal dysfunction is linked to AD, the role of lysosomal exocytosis in the context

of AD is an important field of research that is relatively understudied. It has been

proposed that lysosomal exocytosis may serve as a prion-like transmission mechanism in

which Aβ, along with other toxic proteins, can spread between cells.[29,33,47] A study by

Annunziata et al. (2013) investigated the link between lysosomal exocytosis and AD

using knockout mice of neuraminidase 1 (NEU1), which displayed higher levels of

lysosomal exocytosis, leading to an increase in Aβ secretion.[29] Besides this study, not

much is known about the regulated secretion of Aβ from neurons and this secretory event

has never been directly imaged. Here, using human induced pluripotent stem cell

(hiPSC)-derived neurons, we sought to visualize the real time release of Aβ by lysosomal

exocytosis and investigate key machinery involved in this process, specifically focusing

on Rab27b and its effector munc13-4.

Results

Aβ is preferentially localized in neuronal lysosomes

iPSC-derived human neurons were generated and differentiated using standard

techniques adapted from Stem Cell Technologies (Supplementary Figure 1). To

determine the intracellular localization of Aβ, wild-type iNeurons were loaded with

exogenous fluorescently labelled $Aβ_{42}$ monomers or APP-Swe iNeurons were incubated

with Amytracker-520 to stain endogenous Aβ. The localization of Aβ was examined by

the colocalization with transduced fluorescently-tagged compartment markers including,

Rab5 for early endosomes, Rab9 for late endosomes, and LAMP1 for lysosomes (Figure

1). In wild-type iNeurons, only 15.61% of exogenously loaded $Aβ_{42}$ co-localized with

Rab5 positive compartments. By comparison, 74.78% of $Aβ_{42}$ co-localized with Rab9

positive compartments (p=0.001) and 87.00% of $Aβ_{42}$ co-localized with LAMP1 positive

compartments (p<0.0001). However, no significant difference was seen between Rab9

late endosomes and LAMP1 lysosomes (p=0.38). On the other hand, in APP-Swe

iNeurons, 87.92% of intracellular Aβ marked by Amytracker-520 co-localized with

LAMP1 positive compartments. This was significantly higher than either endosomal

marker, with only 5.30% Amytracker-520 co-localizing with Rab5 positive

compartments (p<0.001) and 20.16% with Rab9 positive compartments (p<0.001).

This supports the idea that accumulation of aggregated Aβ occurs within lysosomes in

AD. The non-significant differences between late endosomes and lysosomes, may be due

to the overlapping expression of Rab9 and LAMP1. Thus, dextran loading was used to

more accurately identify terminal lysosomes.[48] Wild-type iNeurons were incubated with

10 kDa dextran-TMR for 6 hours, followed by an overnight incubation at 37°C for 20

hours (Figure 2). The timing of this pulse-chase was used to ensure that the dextran had

fully traversed the endocytic pathway and was confined to lysosomes, which has been

previously demonstrated.[49] Across all compartment markers, 88.19% of all $Aβ_{42}$ spots

contained dextran and 92.39% contained LAMP-1. Importantly, when Rab5 and Rab9

compartment markers were exclusively separated from dextran signal (i.e, Rab5(+) dex(-)

and Rab9(+) and dex(-)), the percentage of $A\beta_{42}$ spots significantly fell to 3.18% and

4.93% for Rab5 (p<0.001) and Rab9 (p<0.001), respectively. This strongly supports the

idea that $A\beta_{42}$ accumulates primarily in lysosomes in AD.

TIRF microscopy captures lysosomal release of amyloid

To follow lysosomal trafficking, iNeurons were transduced with LAMP1-BFP to label

lysosomes and pHmScarlet-LAMP1 to monitor intraluminal pH. pHmScarlet exhibits

minimal fluorescence under acidic conditions but demonstrates strong fluorescence at

neutral pH. Upon fusion of lysosomes with the plasma membrane, the incoming culture

media neutralizes the acidic environment of the lysosomes. This shift to a neutral pH

triggers a marked increase in fluorescence of pHmScarlet.

The changes in fluorescence intensity of individual lysosomes were captured using total

internal reflection fluorescence (TIRF) microscopy. In brief, TIRF microscopy provides

higher resolution of events occurring near the plasma membrane by restricting the depth

of light energy able to penetrate a cell. To do this, laser light is reflected off the surface of

the glass coverslip at a critical reflection angle in which almost all of the energy is

reflected. However, an evanescent wave that can excite fluorophores penetrates 70-250

nm into the specimen surface. This eliminates any background signal outside this specific

z-plane that would be captured under normal epifluorescence microscopy and allows for

high-resolution time-lapse videos.

At rest, exogenously loaded fluorescent $A\beta_{42}$ and endogenous Amytracker-stained $A\beta_{42}$

was seen in LAMP1 compartments that had little pHmScarlet signal, indicating that they

were acidified lysosomes captured by live-cell TIRF microscopy (Figure 3). Ionomycin, a

Ca^{2+} ionophore, was used to trigger exocytosis, and live-cell videos were acquired in

wild-type cells (Supplementary Movie 1) and APP-Swe cells (Supplementary Movie

2).[21,28,35,50] In wild-type iNeurons loaded with $A\beta_{42}$, lysosomes began fusing with the

plasma membrane 10-20 seconds following addition of ionomycin, resulting in a gain in

pHmScarlet-LAMP1 intensity. Meanwhile, there was a simultaneous reduction in $A\beta_{42}$

intensity within LAMP1-BFP positive compartments. In APP-Swe cells labelled with

Amytracker-520, there was an increase in pHmScarlet-LAMP1 intensity for

approximately 30 seconds following ionomycin treatment, along with a simultaneous

decrease of LAMP1-BFP and Amytracker-520 intensities.

Silencing of Rab27b and munc13-4 impedes lysosomal exocytosis of amyloid

To study the machinery involved in lysosomal exocytosis in neurons, silencing of

Rab27b and munc13-4 was performed using shRNA adenoviruses (Supplementary Figure

2). Silencing of both proteins in wild-type and APP-Swe iNeurons completely abolished

the rise in the pHmScarlet signal captured by TIRF microscopy (Figure 4). The intensities

of LAMP1-BFP, pHmScarlet-LAMP1, $A\beta_{42}$-647 or Amytracker-520 all remained

relatively stable. This lack of intensity changes represented a failure of LAMP1-positive

vesicles to fuse with the plasma membrane and thus, prevent the release of $A\beta$ into the

extracellular space.

To verify that $A\beta$ was released during ionomycin-induced exocytosis and that this release

was blocked by silencing of Rab27b and munc13-4, media samples were collected before

and after exocytosis, and $A\beta$ levels were analyzed by ELISAs. Wild-type and APP-Swe

cells demonstrated a significant increase in $A\beta$ release following ionomycin treatment. In

wild-type cells, the levels of extracellular $A\beta_{42}$ rose from undetectable levels to 458.33

pg/mL following ionomycin treatment ($p<0.001$). This was reduced to 70.77 pg/mL by

silencing of Rab27b ($p<0.001$) and to 19.67 by silencing of munc13-4 ($p<0.001$).

Likewise, this was also seen in APP-Swe iNeurons, in levels of extracellular $A\beta_{42}$ rose

from undetectable levels to 391 pg/mL following ionomycin treatment ($p<0.001$). This

was reduced to 75.33 pg/mL from shRab27b treated cells ($p<0.001$) and to 10.67 pg/mL

from shMunc13-4 treated cells ($p<0.001$).

Lysosomes are the main secretory organelle releasing amyloid-containing particles upon

ionomycin treatment

To better examine the particles that are released in the extracellular space,

nanoscale flow cytometry (or nanoflow analysis) was performed using an Apogee A60

Microplus system, which is optimized for the detection of 80-100 nm range particles with

much higher sensitivity. We have previously used this system to identify tau and

amyloid-containing particles from AD patient plasma.[51] iNeurons were transduced with

fluorescently-tagged compartment markers and either loaded with fluorescent Aβ (for

wild-type iNeurons) or labelled with Amytracker (for APP-Swe iNeurons) (Figure 5).

The media was collected pre- and post-ionomycin treatment and fluorescent particles in

the media were sorted by nanoscale flow cytometry. Prior to ionomycin treatment, almost

no fluorescent particles were detected in the media. Following ionomycin treatment to

trigger exocytosis, majority of BFP-containing particles were detected from cells

expressing LAMP1-BFP, but few from cells transduced with Rab5-BFP or Rab9-BFP. In

wild-type cells, an average of 6981.89 events/µL LAMP1 containing particles was

detected in samples post-ionomycin treatment, which is a significant increase from 77.33

events/µL pre-ionomycin treatment (p<0.001). In contrast, no significant levels of Rab5

(688.16 events/µL, p=0.5) or Rab9 (305.91 events/µL, p=0.77) were measured in the

media following ionomycin treatment. In APP-Swe cells, an average of 66338 events/µL

LAMP1 containing particles was measured in samples post-ionomycin treatment, which

was a significant increase from 1138.22 events/µL pre-ionomycin treatment (p<0.001).

No significant changes were detected for Rab5 (1416.33 events/µL, p=0.89) or Rab9

(3583 events/µL, p=0.72).

Nanoscale flow cytometry reveals differences in Aβ-containing particles released from

wild-type and APP-Swe iNeurons

Interestingly, our nanoscale flow cytometry results also found a stark difference

between the type of Aβ-containing particles released by wild-type versus APP-Swedish

iNeurons (Figure 6). In addition to detecting LAMP1, we also examined the presence of

common exosomal marker CD9. In wild-type cells, 52.1% of smaller fluorescent Aβ-

containing particles (180-300 nm) did not contain LAMP1 or CD9 (LAMP1-/CD9-). This

was significantly greater than Aβ secreted which contained LAMP1 only (15.2%,

p<0.001), CD9 only (3.73%, p<0.001), or with LAMP1 and CD9 together (21.2%,

p<0.001). In contrast, 57.3% of larger Aβ-containing particles (300-1000 nm) was

secreted with LAMP1 and CD9 together, which was significantly greater than Aβ

secreted with LAMP1 only (10.7%, p<0.001), CD9 only (4.78%, p<0.001), or as

unbound cargo (21.1%, p<0.001).

In APP-Swedish cells, the majority of Amytracker-labelled Aβ was released in particles

that contained LAMP1 and CD9 together, regardless of size. 66.7% of smaller Aβ (180-

300 nm) was secreted in particles bearing LAMP1 and CD9, which was significantly

greater than Aβ secreted with LAMP1 only (14.2%, p<0.001), CD9 only (2.78%,

p<0.001), or as unbound cargo (11.2%, p<0.001). Likewise, 87.5% of larger Aβ (300-

1000 nm) was secreted in particles containing both LAMP1 and CD9, which was

significantly greater than Aβ secreted bearing LAMP1 alone (4.49%, p<0.001), CD9

alone (2.27%, p<0.001), or lacking both (3.89%, p<0.001).

These findings suggest that small exogenously loaded Aβ$_{42}$ aggregates are released as

unbound cargo in wild-type cells, whereas aggregated Aβ is mainly released within

membrane bound compartments in APP-Swe cells.

Discussion

Despite extensive research on Aβ, direct imaging of Aβ secretion in human neurons has

never been shown. Prior work in our lab has established the involvement of endosomes

and lysosomes in Aβ production.[2–4,52] Here, we confirmed that exogenously loaded

fluorescent-labelled Aβ and endogenously produced Aβ resides within lysosomes. Using

TIRF microscopy, we directly observed exogenously-loaded fluorescent Aβ$_{42}$ release via

lysosomal exocytosis following ionomycin treatment. Similarly, the secretion of

endogenous Aβ, identified with Amytacker in APP-Swe cells, was also imaged by TIRF

microscopy as it was secreted from the cells upon ionomycin induction. This was blocked

by shRNA knockdown of genes known to participate in lysosomal secretion (Rab27b and

Munc13-4), and confirmed by ELISA-based detection of Aβ in the media following

stimulation. Nanoscale flow cytometry revealed that small Aβ-containing aggregates

released in this process are predominantly released without the membrane/exosome

markers LAMP1 or CD9, while larger aggregated Aβ is released within LAMP1/CD9

containing structures. The findings of this study demonstrated the exocytic capabilities of

neuronal lysosomes, and that lysosomes release of Aβ and Aβ-containing aggregates into

the extracellular space.

This adds to the limited research on the presence of secretory lysosomes in cells of the

central nervous system and in neurons.[27–33] However, most of our understanding of the

machinery involved in this process comes from studies in non-neuronal cells.[16,19–26] The

involvement of Rab27b in lysosomal secretion is well-supported by research studies that

underscore its critical role in vesicular trafficking and exocytosis.[42–45] For instance,

silencing of Rab27b results in the perinuclear accumulation of CD63, a marker for late

endosomes and lysosomes.[39] In the context of elimination of pathological substrates,

Rab27b has been implicated in the exocytosis of alpha-synuclein.[53–55] In particular,

knocking down Rab27b disrupts lysosomal function, leading to the accumulation of

defects in primary neuronal cultures and in the cortex and hippocampus of Rab27b

knockout mice.[53] Future work will be needed to map out the regulatory structure of

lysosomal exocytosis in neurons, including the role of calcium sensors such as

SytVII,[27,34] and proteins of the SNARE machinery, particularly the t-SNARE syntaxin-4

and the v-SNARE VAMP-7 that are critical for the fusion of lysosomes with the plasma

membrane.[24,35,56] While Rab27b is a known secretory Rab that has been shown to be

highly abundant in the central nervous system, only a handful of papers have examined

Rab27 in AD.[40–43]

This work supports also the longstanding idea that lysosomes are important in the

aggregation and release of Aβ. Our own work suggests that APP can be rapidly

transported to lysosomes and from the plasma membrane by macropinocytosis, and

directly from the Golgi apparatus, where it can be cleaved into Aβ.[2–4] Even in the 1990s,

there was evidence that extracellular Aβ is taken up into lysosomes, where it can nucleate

endogenous Aβ aggregation.[57,58] Aβ is known to accumulate and aggregate intracellularly

in lysosomes.[11,12,59–61] The aggregation of Aβ in late endosomes and lysosomes can

disrupt synapses[62] and cause lysosomal rupture leading to cell death.[63] Aβ secretion is

known to be driven by neural activity.[64] Because lysosomal enzymes are only activated in

the lysosomes, presence of active lysosomal enzymes in amyloid plaques suggests that

they are derived from lysosomes.[65]

In summary, here we demonstrate that Aβ is directly secreted from neurons by lysosomal

secretion. This work dovetails with longstanding observations of Aβ aggregation and

accumulation in lysosomes and the observation that extracellular Aβ plaques contain

active lysosomal enzymes. The secretion of Aβ in aggregates from lysosomal secretion

suggests that this process may play a role in generating aggregates that could propagate in

the development of AD.

Methods

Cell culturing of hiPSC-derived neurons

hiPSCs were obtained from the New York Stem Cell Foundation, including cell lines

containing the APP Swedish mutation (APP-Swe; CO002-01-CS-003, Lot# E137-1D)

and its isogenic control (wild-type; CO0002-01-SV-003, Lot# E259-1H). The induction

of hiPSCs into neural progenitor cells (NPCs) was performed using STEMdiff™ SMADi

neural induction kit (Stem Cell Technologies, 08581) and STEMdiff™ neural induction

media (Stem Cell Technologies, 05835), following the manufacturer's protocols outlined

in 'Generation and Culture of Neural Progenitor Cells Using the STEMdiff™ Neural

System'. In brief, embryoid bodies were initially generated, maintained for 10 days, and

replated for neural rosette selection using STEMdiff™ neural rosette selection reagent

(Stem Cell Technologies, 05832). NPC outgrowths were then formed from selected

rosette-containing clusters.

Selected NPCs were then maintained and passaged (every 5-7 days) in STEMdiff™

neural progenitor medium (Stem Cell Technologies, 05833), or differentiated for 7 days

using STEMdiff™ forebrain neuron differentiation kit (Stem Cell Technologies, 08600).

Following differentiation, cells were matured into iNeurons for 21 days in vitro (DIV)

using Neurobasal™ medium (Gibco, 21103049) supplemented with 1x B27 (Gibco,

17504044), 0.5x GlutaMAX (Gibco, 35050061), 1x antibiotic-antimycotic (Gibco,

15240096), 0.5 mM db-cAMP (Sigma-Aldrich, D0627), 0.2 ug/mL ascorbic acid (Sigma-

Aldrich, A92902), 20 ng/mL BDNF (PeproTech, 450-02), and 20 ng/mL GDNF

(PeproTech, 450-10). For all experiments, cells were maintained or plated on cultureware

coated with 15 ug/mL poly-L-ornithine (Sigma-Aldrich, P4957) and 10 ug/mL laminin

(Corning, 354232).

Viral vectors

All constructs were designed, packaged into adenovirus serotype 5, and purchased from

VectorBuilder. iNeurons were transduced at 18 DIV with the listed multiplicity of

infections (MOIs) and performed following recommendations set by VectorBuilder.

Virus	MOI	Vector Builder ID	General Purpose
CD9-SNAP	50	VB221202-1061typ	exosome marker
LAMP1-BFP	50	VB191029-14394qax	lysosome marker
LAMP1-miRFP	50	VB211121-1082udt	lysosome marker
pHmScarlet-LAMP1	30	VB220418-1238mbc	pH sensitive lysosomal maker (only at neutral pH)
Rab5-BFP	50	VB200411-2007gzu	early endosome marker
Rab5-miRFP	50	VB211125-1183cef	early endosome marker
Rab9-BFP	50	VB200410-1338nsa	late endosome marker
Rab9-miRFP	50	VB211125-1182dhn	late endosome marker
shMUNC13-4	100	VB200421-1904fgb	target sequence: GAGTCCTGGCCCTGGATAATT
shRab27b	100	VB191029-4720pgh	target sequence: GACGCCATGGGCTTCTTATTA
shScramble	100	VB240723-1508pnw	silencing control

Aβ monomer preparation and loading & Amytracker detection

$A\beta_{42}$ monomers were generated from recombinant $A\beta_{42}$ tagged with HiLyte™ Fluor 647

(Anaspec, AS-64161), following previously established protocols.[66] In brief, lyophilized

peptides were solubilized in 1,1,1,3,3,3-hexafluoro-2-propanol (Sigma-Aldrich, 105288).

Vacuum centrifugation was used to evaporate the HFIP for subsequent resuspension in

dimethyl sulfoxide (DMSO). Following a 10-minute period of sonication, $A\beta_{42}$

monomers were diluted in ice-cold PBS. Matured wild-type iNeurons (20 DIV) were

loaded with 250 nM of $A\beta_{42}$ overnight prior to experimentation for total internal

fluorescence (TIRF) microscopy.

Instead of $A\beta_{42}$ loading, APP-Swe iNeurons were incubated with Amytracker-520 (1:250,

Ebba Biotech) for 2 hours to mark endogenous aggregated $A\beta$ for microscopy

experiments. For nanoscale flow cytometry experiments, the media of APP-Swe

iNeurons was incubated with Amytracker-680 (1:100, Ebba Biotech) for 10 minutes prior

to analysis.

Ionomycin treatment

Following $A\beta_{42}$ loading overnight or Amytracker incubation for 2 hours, iNeurons were

washed and treated with 5 or 10 µM of ionomycin (Sigma-Aldrich, C7522) in Hank's

Balanced Salt Solution (HBSS, Gibco, 14025092) to induce lysosomal exocytosis. 5 µM

of ionomycin was used for microscopy experiments, whereas 10 µM of ionomycin was

used for enzyme-linked immunosorbent assays (ELISAs) and nanoscale flow cytometry.

This difference is mainly due to the fact that for microscopy experiments, ionomycin was

directly injected over the confocal dish for live-cell capture of events, and thus a lower

concentration was used, whereas HBSS containing ionomycin was replaced across the

entire 6-well for ELISAs and nanoscale flow cytometry.

Immunofluorescence staining

To verify the neural cell yield, immunofluorescence staining was performed for nestin

and MAP2 (Supplementary Figure 1). In brief, iNeurons were fixed for 15 minutes with

4% paraformaldehyde. Cells were permeabilized for 10 minutes with 0.1% Triton X-100

in HBSS and blocked with 2% bovine serum albumin (BSA) for 30 mins. Cells were

incubated with primary antibodies for MAP2 (1:1000; Sigma-Aldrich, M9942) and nestin

(1:200; Invitrogen, PA5-11887) for 1 hour at RT, washed three times with HBSS, and

incubated with species-appropriate secondary antibodies conjugated to Alexa Fluor 488

(1:1000; Invitrogen, A21202) or Alexa Fluor 647 (1:1000; Invitrogen, A31573) for 1

hour at RT. Cells were then counterstained with DAPI (Sigma-Aldrich, D9542).

Confocal microscopy and image analysis

Images were acquired on a Leica TCS SP8 confocal microscope (Leica, Germany) using

an HC PL APO CS2 63x/1.40 oil objective, with excitation lasers of 405 nm, 488 nm,

552 nm, and 638 nm. Emission filters were consistent for each set of experiments.

For immunofluorescence experiments with fixed cells, a minimum of 3 regions were

acquired per confocal dish, with each region consisting of a stitched merge of 5x5 tiles.

To quantify the neural yield of 3 experimental replicates, the relative percentage of nestin

and MAP2 positive cells was calculated using the filament tracer function on Imaris 10

Software (Oxford Instruments).

For all co-localization experiments, Z-stack images of the entire cell were acquired from

a minimum of 3 live-cells per confocal dish, per replicate. Imaris was used for semi-

automated quantification across 3 experimental replicates. In Aβ and compartmental

marker experiments, object-object based colocalization was measured by calculating the

percentage of co-localization between spots (semi-automatically generated for Aβ) and

surfaces (semi-automatically generated for compartment markers). In experiments using

10 kDa dextran-tetramethylrhodamine (TMR; Invitrogen, D1868), spots were created to

represent $A\beta_{42}$, and intensity filters were then used to classify spots based on the presence

of dextran-TMR and/or compartment maker signals.

TIRF microscopy

To capture live cell videos of individual lysosomes collapsing at the plasma membrane,

total internal reflection fluorescence (TIRF) microscopy was performed using a Leica

DMI 6000B TIRF microscope (Leica, Germany) with an HCX PL APO 63X oil

immersion objective with numerical aperture of 1.47. Fluorescence emitted from

excitation lasers at 405 nm, 488 nm, 561 nm, and 635 nm was captured after passing

through a Quad-Band filter cube. The Quad-Band filter cube setup allowed for a very

short switching time and thus, an ultra-high synchronized frame rate that is required for

the analysis of TIRF videos.

Following $A\beta_{42}$ loading overnight or Amytracker incubation for 2 hours, iNeurons were

washed with pre-warmed HBSS. Cells were then transported to the microscope stage in

the incubation chamber and treated directly with 5 µM of ionomycin. To capture rapid

exocytic events, the imaging time for each wavelength was synchronized with the

hardware. For every video, images were captured approximately every 0.3-0.5 seconds

(across 3 channels) for 150-300 time points.

A minimum of 10 live cells were captured using TIRF microscopy and analyzed with

Imaris 10 Software (Oxford Instruments). Using the semi-automated spot detection

feature on Imaris, a spot was created to represent each lysosome (based on the

pHmScarlet-LAMP1 channel), and tracks were generated to follow the fate of each

lysosome over time. At each single time point, the mean intensity of each channel was

measured from that spot and averaged across all spots within a single cell. Tracks were

normalized to its maximum intensity, and the change in normalized intensity was plotted

against time to compare across conditions.

Western blotting

Western blotting for Rab27b and munc13-4 was performed to verify their silencing

(Supplementary Figure 2). iNeurons were washed on ice with ice-cold HBSS and then

lysed in N-PER lysis buffer (Thermo Scientific, 87792) with 1X protease and

phosphatase inhibitor cocktail (Thermo Scientific, 78442). Protein concentrations were

measured using the Pierce BCA Protein Assay (Thermo Scientific, 23225).

Equal amounts of total protein were mixed with 2X Laemmli buffer (Bio-Rad, 1610737),

denatured at 95°C for 5 minutes, and loaded onto 4-20% Mini-PROTEAN® TGX™

precast protein gels (Bio-Rad, 4561096). Gels were run at 125V for 75 minutes and

subsequently, transferred onto methanol-activated Immun-Blot® PVDF membranes (Bio-

Rad, 1620177) at 90 V for 75 minutes. Following Ponceau verification of transfer,

membranes were blocked with 5% BSA for 1 hour. Membranes were incubated in

primary antibodies (α-Rab27b: 1:500, Proteintech, 13412-1-AP, α-munc13-4: 1:500,

Proteintech, 16905-1-AP, and α-alpha-tubulin: 1:100,000, Sigma-Aldrich, T5168) in

blocking solution overnight at 4°C. Membranes were incubated with HRP-conjugated

goat anti-rabbit (1:5000; Bio-Rad, 1706515) for 1 hour at RT, followed by visualization

using Amersham ECL prime western blotting detection reagent (Cytvia, RPN2236) on a

Bio-Rad ChemiDoc MP. Densitometric analysis was conducted using ImageLab

Software (Bio-Rad).

Enzyme-linked immunosorbent assays (ELISAs)

For ELISAs, matured iNeurons grown on coated 6-well plates were washed with pre-

warmed HBSS and incubated with 10 µM ionomycin in HBSS at 37°C for 5 minutes

prior to supernatant collection. Samples were centrifuged at 500 x g for 5 minutes to

remove any debris and the supernatant was then collected and frozen at -80°C. For

measurement of $A\beta_{42}$ levels, supernatants were diluted 10-fold and detected using an

ELISA for human $A\beta_{42}$ (Invitrogen, KHB3544). Triplicates of each sample from 3

replicates were loaded per plate.

Nanoscale flow cytometry

Supernatant samples containing exocytosed contents were collected similarly to the

ELISA samples, as described above. For detection of CD9-SNAP in supernatants by

nanoscale flow cytometry, labelling was performed using SNAP-Cell™ Oregon-Green

(New England Biolabs, S9104S) according to the manufacturer's instructions.

Undiluted samples were analyzed on the Apogee A60 Microplus Nanoflow Cytometer

(Apogee Flow Systems Inc.). Prior published work from our lab has already established

technical settings to achieve data linearity using this system.[51,67] Size gating was

determined by calibration beads to observe smaller (180-300 nm) and larger particles

(300-1000 nm), as previously published.[67] Instrument settings included sheath pressure at

200 mbar, flow rate of 1.50 µL/min for 130 µL and lasers at 70 mW (405 nm), 70 mW

(488 nm), and 70 mW (638 nm). Light scatter of events was produced using the 405 nm

laser, with thresholds to eliminate background noise of 37 a.u. for small angle light

scatter (SALS) and 40 a.u. for large angle light scatter (LALS). Photomultiplier tube

(PMT) voltages were set at 260 V at LALS, 340 V at SALS, 525 V at L405-Blu, 525 V at

L488-Grn, and 575 V at L638-Red. For each nanoflow experiment, samples were run in

triplicate from 3 replicates.

Statistical analysis

Statistical analysis was conducted using GraphPad Prism 10 (GraphPad Software, USA).

All data is represented as mean ± SEM. Unpaired t-tests and one-way or two-way

ANOVAs with Tukey's multiple comparisons test were used for statistical analysis.

Statistical significance was defined as $*p < 0.05$, $**p < 0.01$, or $***p < 0.001$.

Figure 1 Live cell imaging of amyloid with compartment markers in wild-type and APP-Swe iNeurons

A) Representative images of wild-type cells (top) loaded with Aβ$_{42}$-647 (red) or APP-Swe cells (bottom) incubated with Amytracker-520 (red). Cells were transduced with adenoviral vectors containing fluorescently tagged Rab5, Rab9, LAMP1 to mark early endosomes, late endosomes and lysosomes, respectively (green). B) Object-object based colocalization was calculated using the spot and surfaces functions on Imaris. All results are presented as mean ± SEM. Asterisks indicate significant changes according to one-way ANOVA and post hoc Tukey test. **p<0.01 ***p<0.001

Figure 2 Live cell imaging of amyloid with dextran in wild-type iNeurons

A) Representative images of wild-type cells loaded with Aβ$_{42}$-647 (red). Cells were transduced with adenoviral vectors containing fluorescently tagged Rab5, Rab9, LAMP1 to mark early endosomes, late endosomes and lysosomes, respectively (green). A 6-hour pulse with dextran and 20 hour chase was used to ensure that the probe was confined to lysosomes (blue). Arrowheads represent Aβ$_{42}$ with dextran positive signal but lacking Rab5 or Rab9 signal. B) Classification and quantification of Aβ$_{42}$ spots containing dextran and/or compartment markers signals was calculated by Imaris. All results are presented as mean ± SEM. Asterisks indicate significant changes according to one-way ANOVA and post hoc Tukey test. ***p<0.001

Figure 3 TIRF microscopy of amyloid released from lysosomes in wild-type and APP-Swe iNeurons

A) Representative images from a live cell video captured through a TIRF microscope demonstrating lysosomal exocytosis in a wild-type and APP-Swe cell after ionomycin treatment over 60 seconds. Cells were transduced with adenoviral vectors containing fluorescently tagged LAMP1-BFP (blue) and pHmScarlet-LAMP1 (green) and incubated with Aβ (red). B) Change in normalized mean intensity of each channel was tracked after ionomycin treatment (at time = 0s). A total of 538 lysosomes were analyzed in wild-type cells (n=10), and 785 lysosomes were tracked in APP-Swe cells (n=10).

Figure 4 Silencing of Rab27b and munc13-4 halts lysosomal exocytosis

A) Representative images from a live cell video captured through a TIRF microscope

demonstrating the lack of lysosomal exocytosis in a wild-type cell upon shRab27b

silencing. B) Silencing of Rab27b and munc-13-4 results in intensities across all 3

channels to remain relatively stable, demonstrating a lack of lysosomal secretion. C)

Significant reduction of Aβ is detected in the media of iNeurons following silencing, as

measured by ELISAs. All results are presented as mean ± SEM. Asterisks indicate

significant changes according to two-way ANOVA with post hoc Tukey test. *p<0.05

***p<0.001

Figure 5 Nanoscale flow cytometry of secreted vesicles triggered by ionomycin

A) Total BFP events/μL were analyzed by nanoscale flow cytometry on media of wild-type and APP-Swe iNeurons following ionomycin treatment. B) Representative cytograms of LAMP1-BFP, Rab5-BFP and Rab9-BFP collected from cell culture media prior and following ionomycin treatment. All results are presented as mean ± SEM. Asterisks indicate significant changes according to two-way ANOVA and post hoc Tukey test. ***p<0.001

Figure 6 Nanoscale flow cytometry of particles containing Aβ

A) Percentage of unbound Aβ (LAMP1-/CD9-), Aβ with LAMP1, CD9 or both
(LAMP1+/CD9+) was measured by nanoscale flow cytometry on media from wild-type
and APP-Swe iNeurons following ionomycin treatment. Particles were sorted by size
(established with calibration beads). B) Representative cytograms of Aβ collected from
cell culture media following ionomycin treatment. Blue boxes indicate particles
containing Aβ only, red boxes indicate particles of Aβ with LAMP1, green boxes indicate
particles of Aβ with CD9, and purple boxes indicate particles of Aβ with LAMP1 and
CD9. All results are presented as mean ± SEM. Asterisks indicate significant changes
according to one-way ANOVA and post hoc Tukey test. *p<0.05, **p<0.01. ***p<0.001

References

1.   Haass, C., Kaether, C., Thinakaran, G. & Sisodia, S. Trafficking and proteolytic
processing of APP. *Cold Spring Harb. Perspect. Med.* **2**, (2012).

2.   Tam, J. H. K., Seah, C. & Pasternak, S. H. The amyloid precursor protein is
rapidly transported from the golgi apparatus to the lysosome and where it is
processed into beta-Amyloid. *Mol. Brain* **7**, 1–18 (2014).

3.   Tang, W. *et al.* Arf6 controls beta-amyloid production by regulating
macropinocytosis of the Amyloid Precursor Protein to lysosomes. *Mol. Brain* **8**,
(2015).

4.   Chiu, J., Krupa, J. M., Seah, C. & Pasternak, S. H. Small GTPases control
macropinocytosis of amyloid precursor protein and cleavage to amyloid-β. *Heliyon*
10, e31077 (2024).

5.   de Duve, C. & Wattiaux, R. Functions of Lysosomes. *Annu. Rev. Physiol.* **28**, 435–
492 (1966).

6.   Nixon, R. A. & Cataldo, A. M. Lysosomal system pathways: Genes to
neurodegeneration in Alzheimer's disease. *J. Alzheimer's Dis.* **9**, 277–289 (2006).

7.   Suzuki, K. & Terry, R. D. Fine structural localization of acid phosphatase in senile
plaques in Alzheimer's presenile dementia. *Acta Neuropathol.* **8**, 276–284 (1967).

8.   Tsukita, S. & Ishikawa, H. The movement of membranous organelles in axons.
Electron microscopic identification of anterogradely and retrogradely transported
organelles. *J. Cell Biol.* **84**, 513–530 (1980).

9.   Cataldo, A. M. *et al.* Aβ localization in abnormal endosomes: association with
earliest Aβ elevations in AD and Down syndrome. *Neurobiol. Aging* **25**, 1263–
1272 (2004).

10.   Margallo-Lana, M. *et al.* Influence of the amyloid precursor protein locus on

dementia in Down syndrome. *Neurology* **62**, 1996–1998 (2004).

11.   Cataldo, A. M. *et al.* Presenilin Mutations in Familial Alzheimer Disease and

Transgenic Mouse Models Accelerate Neuronal Lysosomal Pathology. *J.*

Neuropathol. Exp. Neurol. **63**, 821–830 (2004).

12.   Langui, D. *et al.* Subcellular Topography of Neuronal Aβ Peptide in APPxPS1

Transgenic Mice. *Am. J. Pathol.* **165**, 1465–1477 (2004).

13.   Settembre, C., Fraldi, A., Medina, D. L. & Ballabio, A. Signals from the lysosome:

A control centre for cellular clearance and energy metabolism. *Nat. Rev. Mol. Cell*

Biol. **14**, 283–296 (2013).

14.   Settembre, C. *et al.* A lysosome-to-nucleus signalling mechanism senses and

regulates the lysosome via mTOR and TFEB. *EMBO J.* **31**, 1095–1108 (2012).

15.   Blott, E. J. & Griffiths, G. M. Secretory lysosomes. *Nat. Rev. Mol. Cell Biol.* **3**,

122–131 (2002).

16.   Chieregatti, E. & Meldolesi, J. Regulated exocytosis: New organelles for non-

secretory purposes. *Nat. Rev. Mol. Cell Biol.* **6**, 181–187 (2005).

17.   Buratta, S. *et al.* Lysosomal exocytosis, exosome release and secretory autophagy.

Int. J. Mol. Sci **21**, 1–20 (2020).

18.   Tancini, B. *et al.* Lysosomal exocytosis: The extracellular role of an intracellular

organelle. *Membranes (Basel).* **10**, 1–21 (2020).

19.   Chavez, R. A., Miller, S. G. & Moore, H. P. H. A biosynthetic regulated secretory

pathway in constitutive secretory cells. *J. Cell Biol.* **133**, 1177–1191 (1996).

20.   Coorssen, J. R., Schmitt, H. & Almers, W. Ca2+ triggers massive exocytosis in

Chinese hamster ovary cells. *EMBO J.* **15**, 3787–3791 (1996).

21.   Rodríguez, A., Webster, P., Ortego, J. & Andrews, N. W. Lysosomes Behave as

Ca2+-regulated Exocytic Vesicles in Fibroblasts and Epithelial Cells. *J. Cell Biol.*

137, 93–104 (1997).

22. Stinchcombe, J. C. & Griffiths, G. M. Regulated secretion from hemopoietic cells.
J. Cell Biol. **147**, 1–5 (1999).

23. Andrews, N. W. Regulated secretion of conventional lysosomes. *Trends Cell Biol.*
10, 316–321 (2000).

24. Jaiswal, J. K., Andrews, N. W. & Simon, S. M. Membrane proximal lysosomes are
the major vesicles responsible for calcium-dependent exocytosis in nonsecretory
cells. *J. Cell Biol.* **159**, 625–635 (2002).

25. Logan, M. R., Odemuyiwa, S. O. & Moqbel, R. Understanding exocytosis in
immune and inflammatory cells: The molecular basis of mediator secretion. *J.*
Allergy Clin. Immunol. **111**, 923–932 (2003).

26. Verhage, M. & Toonen, R. F. Regulated exocytosis: merging ideas on fusing
membranes. *Curr. Opin. Cell Biol.* **19**, 402–408 (2007).

27. Arantes, R. M. E. A Role for Synaptotagmin VII-Regulated Exocytosis of
Lysosomes in Neurite Outgrowth from Primary Sympathetic Neurons. *J. Neurosci.*
26, 4630–4637 (2006).

28. Bowser, D. N. & Khakh, B. S. Two forms of single-vesicle astrocyte exocytosis
imaged with total internal reflection fluorescence microscopy. *Proc. Natl. Acad.*
Sci. **104**, 4212–4217 (2007).

29. Annunziata, I. *et al.* Lysosomal NEU1 deficiency affects amyloid precursor
protein levels and amyloid-β secretion via deregulated lysosomal exocytosis. *Nat.*
Commun. **4**, 2734 (2013).

30. Padamsey, Z. *et al.* Activity-Dependent Exocytosis of Lysosomes Regulates the
Structural Plasticity of Dendritic Spines. *Neuron* **93**, 132–146 (2017).

31. Zhang, Z. *et al.* Regulated ATP release from astrocytes through lysosome

exocytosis. *Nat. Cell Biol.* **9**, 945–953 (2007).

32.   Tsunemi, T. *et al.* Increased Lysosomal Exocytosis Induced by Lysosomal Ca2+
Channel Agonists Protects Human Dopaminergic Neurons from α-Synuclein
Toxicity. *J. Neurosci.* **39**, 5760–5772 (2019).

33.   Xie, Y. X. *et al.* Lysosomal exocytosis releases pathogenic α-synuclein species
from neurons in synucleinopathy models. *Nat. Commun.* **13**, (2022).

34.   Reddy, A., Caler, E. V., Andrews, N. W. & Haven, N. Plasma Membrane Repair Is
Mediated by Ca2+-Regulated Exocytosis of Lysosomes. *Cell* **106**, 157–169
(2001).

35.   Rao, S. K., Huynh, C., Proux-Gillardeaux, V., Galli, T. & Andrews, N. W.
Identification of SNAREs Involved in Synaptotagmin VII-regulated Lysosomal
Exocytosis. *J. Biol. Chem.* **279**, 20471–20479 (2004).

36.   Burkhardt, J. K., McIlvain, J. M., Sheetz, M. P. & Argon, Y. Lytic granules from
cytotoxic T cells exhibit kinesin-dependent motility on microtubules in vitro. *J.
Cell Sci.* **104**, 151–162 (1993).

37.   Langford, G. M. Actin- and microtubule-dependent organelle motors:
interrelationships between the two motility systems. *Curr. Opin. Cell Biol.* **7**, 82–
88 (1995).

38.   D'Adamo, P. *et al.* RAB GTPases and RAB-interacting proteins and their role in
the control of cognitive functions. *Neurosci. Biobehav. Rev.* **46**, 302–314 (2014).

39.   Ostrowski, M. *et al.* Rab27a and Rab27b control different steps of the exosome
secretion pathway. *Nat. Cell Biol.* **12**, 19–30 (2010).

40.   Fukuda, M. Membrane traffic in the secretory pathway. *Cell. Mol. Life Sci.* **65**,
2801–2813 (2008).

41.   Bustos, M. A., Lucchesi, O., Ruete, M. C., Mayorga, L. S. & Tomes, C. N. Rab27

and Rab3 sequentially regulate human sperm dense-core granule exocytosis. *Proc. Natl. Acad. Sci.* **109**, (2012).

42. Gomi, H., Mori, K., Itohara, S. & Izumi, T. Rab27b Is Expressed in a Wide Range of Exocytic Cells and Involved in the Delivery of Secretory Granules Near the Plasma Membrane. *Mol. Biol. Cell* **18**, 4377–4386 (2007).

43. Fukuda, M. Rab27 Effectors, Pleiotropic Regulators in Secretory Pathways. *Traffic* **14**, 949–963 (2013).

44. Handley, M. T. W., Haynes, L. P. & Burgoyne, R. D. Differential dynamics of Rab3A and Rab27A on secretory granules. *J. Cell Sci.* **120**, 973–984 (2007).

45. Izumi, T. In vivo Roles of Rab27 and Its Effectors in Exocytosis. *Cell Struct. Funct.* **46**, 79–94 (2021).

46. Sekine, Y. *et al.* Functional Genome-wide Screen Identifies Pathways Restricting Central Nervous System Axonal Regeneration. *Cell Rep.* **23**, 415–428 (2018).

47. Guo, J. L. & Lee, V. M. Y. Seeding of normal tau by pathological tau conformers drives pathogenesis of Alzheimer-like tangles. *J. Biol. Chem.* **286**, 15317–15331 (2011).

48. Barral, D. C. *et al.* Current methods to analyze lysosome morphology, positioning, motility and function. *Traffic* **23**, 238–269 (2022).

49. Johnson, D. E., Ostrowski, P., Jaumouillé, V. & Grinstein, S. The position of lysosomes within the cell determines their luminal pH. *J. Cell Biol.* **212**, 677–92 (2016).

50. Liu, C. & Hermann, T. E. Characterization of ionomycin as a calcium ionophore. *J. Biol. Chem.* **253**, 5892–5894 (1978).

51. Dayarathna, T. *et al.* Nanoscale flow cytometry-based quantification of blood-based extracellular vesicle biomarkers distinguishes MCI and Alzheimer's disease.

Alzheimer's Dement. (2024) doi:10.1002/alz.14087.

52. Pasternak, S. H. *et al.* Presenilin-1, nicastrin, amyloid precursor protein, and γ-
secretase activity are co-localized in the lysosomal membrane. *J. Biol. Chem.* **278**,
26687–26694 (2003).

53. Underwood, R. *et al.* The GTPase Rab27b regulates the release, autophagic
clearance, and toxicity of α-synuclein. *J. Biol. Chem.* **295**, 8005–8016 (2020).

54. Pattanayak, R. *et al.* Deletion in chromosome 6 spanning alpha-synuclein and
multimerin1 loci in the Rab27a/b double knockout mouse. *Sci. Rep.* **12**, 9837
(2022).

55. Bellucci, A., Longhena, F. & Spillantini, M. G. The Role of Rab Proteins in
Parkinson's Disease Synaptopathy. *Biomedicines* **10**, 1941 (2022).

56. Luzio, J. P., Parkinson, M. D. J., Gray, S. R. & Bright, N. A. The delivery of
endocytosed cargo to lysosomes. *Biochem. Soc. Trans.* **37**, 1019–1021 (2009).

57. Yang, A. J., Knauer, M., Burdick, D. A. & Glabe, C. Intracellular Aβ1-42
Aggregates Stimulate the Accumulation of Stable, Insoluble Amyloidogenic
Fragments of the Amyloid Precursor Protein in Transfected Cells. *J. Biol. Chem.*
270, 14786–14792 (1995).

58. Burdick, D., Kosmoski, J., Knauer, M. F. & Glabe, C. G. Preferential adsorption,
internalization and resistance to degradation of the major isoform of the
Alzheimer's amyloid peptide, Aβ1–42, in differentiated PC12 cells. *Brain Res.*
746, 275–284 (1997).

59. Hu, X. *et al.* Amyloid seeds formed by cellular uptake, concentration, and
aggregation of the amyloid-beta peptide. *Proc. Natl. Acad. Sci. U. S. A.* **106**,
20324–20329 (2009).

60. Gouras, G. K. *et al.* Intraneuronal Aβ42 Accumulation in Human Brain. *Am. J.*
Pathol. **156**, 15–20 (2000).

61.   D'Andrea, M. R., Nagele, R. G., Wang, H. Y., Peterson, P. A. & Lee, D. H. S.
Evidence that neurones accumulating amyloid can undergo lysis to form amyloid
plaques in Alzheimer's disease. *Histopathology* **38**, 120–134 (2001).

62.   Takahashi, R. H. *et al.* Intraneuronal Alzheimer abeta42 accumulates in
multivesicular bodies and is associated with synaptic pathology. *Am. J. Pathol.*
161, 1869–79 (2002).

63.   Yang, A. J., Chandswangbhuvana, D., Margol, L. & Glabe, C. G. Loss of
endosomal/lysosomal membrane impermeability is an early event in amyloid A?1-
42 pathogenesis. *J. Neurosci. Res.* **52**, 691–698 (1998).

64.   Cirrito, J. R. *et al.* Endocytosis Is Required for Synaptic Activity-Dependent
Release of Amyloid-β In Vivo. *Neuron* **58**, 42–51 (2008).

65.   Cataldo, A. M. & Nixon, R. A. Enzymatically active lysosomal proteases are
associated with amyloid deposits in Alzheimer brain. *Proc. Natl. Acad. Sci.* **87**,
3861–3865 (1990).

66.   Stine, W. B., Dahlgren, K. N., Krafft, G. A. & LaDu, M. J. In vitro
characterization of conditions for amyloid-β peptide oligomerization and
fibrillogenesis. *J. Biol. Chem.* **278**, 11612–11622 (2003).

67.   Gomes, J. *et al.* Analytical Considerations in Nanoscale Flow Cytometry of
Extracellular Vesicles to Achieve Data Linearity. *Thromb. Haemost.* **118**, 1612–
1624 (2018).

705