Identification of Genes Required for Spatial Control and Mechanical Resilience of Cytokinesis during *Caenorhabditis elegans* Embryogenesis

-Author names and Affiliations

Chelsey Lynn Gough[1,2,3], Yuxuan Rain Xiong[1,2,3], Aoi Hiroyasu[1,2,3], Christina Rou Hsu[1,2],

Viktorija Juciute[1,2], MinJee Kim[1,2], Kalen Dofher[1,2], and Kenji Sugioka[1,2*]

[1]Life Sciences Institute, The University of British Columbia, 2350 Health Sciences Mall, Vancouver, BC V6T1Z3, Canada

[2]Department of Zoology, The University of British Columbia, 2350 Health Sciences Mall, Vancouver, BC V6T1Z3, Canada

[3]These authors contributed equally to this work.

*Corresponding author

Kenji Sugioka

Life Sciences Institute, Department of Zoology, The University of British Columbia, 2350 Health Sciences Mall, Vancouver, BC V6T1Z3, Canada

kenji.sugioka@ubc.ca

Phone: +1-604-822-4628

1 Abstract

Cytokinesis is the final step of cell division, in which the dividing cell is physically

separated into two daughter cells by the contractile ring. The contractile ring is a highly

resilient molecular machine that can function   properly under mechanical stress.

Additionally, its function, position, and orientation are spatially modulated in developing

animals to regulate morphogenesis. Although essential regulators of cytokinesis have

been identified through previous genetic screens, the molecular mechanisms underlying

these spatial controls and the mechanical resilience of cytokinesis remain elusive. To

identify cytokinesis regulators involved in these processes, we performed a high-

throughput RNAi screen using a gain-of-function mutant of actin that exhibits ectopic

cortical contraction and abnormal spatial control of cytokinesis in *Caenorhabditis*

elegans embryos. We obtained a total of 483 early embryonic genes that suppress

embryonic lethality in an *act-2* mutant background. Two parallel secondary screens of

candidate genes were conducted. The first secondary screen in a wild-type background

identified 71 candidate genes regulating spatial cytokinesis control—asymmetric ring

closure, positioning, and rotation—during early embryogenesis. The second secondary

screen in the *act-2(or295)* background identified four genes required for cytokinesis in

this background, including microtubule regulators, *evl-20/ARL2*, and *lpin-1/Lipin1*. This

study will serve as a useful resource for the development of future hypotheses and

provide insights into the precise regulation of cytokinesis in tissues.

Short title: Spatial and Robust Cytokinesis Control in *C. elegans*

Keywords: Cytokinesis, C. elegans, Contractile ring, Cell division

Introduction

Cytokinesis is the final step of cell division, in which the contractile ring physically separates the dividing cell into two daughter cells (Pollard and O'Shaughnessy 2019). The contractile ring is a mechanically resilient molecular machine capable of withstanding mechanical stress and repairing physical damage (Silva *et al.* 2016; Singh *et al.* 2019). Additionally, its orientation and position are spatially regulated in three distinct ways during morphogenesis (Sugioka 2021). First, asymmetric contractile ring closure, often found in zygotes and epithelial tissue, is critical for maintaining cytokinesis resilience, epithelial integrity, and polarity (Maddox *et al.* 2007; Herszterg *et al.* 2013; Taulet *et al.* 2017). Second, cell rotation during cytokinesis is required for dorsal-ventral axis establishment in *C. elegans* (Priess and Thomson 1987; Schierenberg and Junkersdorf 1992) and is implicated in the establishment of animal left-right asymmetry in pond snails and *C. elegans* (Meshcheryakov and Beloussov 1975; Wood 1991; Kuroda *et al.* 2009; Naganathan *et al.* 2014). Finally, asymmetric positioning of the contractile ring during asymmetric cell division is critical for generating unequal daughter cell sizes, which is required for subsequent cell fate decisions (Jankele *et al.* 2021). Despite their significance, the regulators of mechanical resilience and the spatial control of cytokinesis remain largely unknown.

Decades of studies have revealed the essential components of the contractile ring, which are evolutionarily conserved in eukaryotes (Pollard 2017). First, the small GTPase RhoA is activated by the centralspindlin complex, thereby initiating contractile ring assembly via stimulation of downstream effectors such as formin and Rho-associated

kinase (Basant and Glotzer 2018). Contractile ring function also requires structural components such as anillin (Piekny and Maddox 2010). As the contractile ring constricts through actin-myosin sliding, membrane trafficking pathways contribute to cytokinesis by delivering additional membrane to the ingressing furrow to accommodate the growing cell surface area (Frémont and Echard 2018). Finally, the daughter cells separate via ESCRT-III-mediated abscission at the midbody (Mierzwa and Gerlich 2014). Previous forward genetic and RNAi-based screens in yeast and various animal systems were typically performed in wild-type backgrounds and focused on identifying essential regulators of cytokinesis (Eggert et al. 2004; Echard et al. 2004; Sönnichsen et al. 2005). A recent study in *C. elegans* performed a small-scale candidate screen in zygotes under uniaxial compression and identified several regulators required for the mechanical resilience of cytokinesis (Singh *et al.* 2019). However, a large-scale genetic screen focusing on the physical and developmental control of cytokinesis is still needed.

In this study, we performed an RNAi screen targeting 4,218 early embryonic genes in an actin gain-of-function mutant of *C. elegans*. The *act-2(or295)* mutation lies near the ATP-binding pocket, leading to filament stabilization and hyperactive cortical contractility (Willis *et al.* 2006). We found that the spatial control of cytokinesis, such as asymmetric ring closure in the zygote, rotational cytokinesis in the 2-cell stage, and asymmetric ring positioning in the 6-cell stage, is also abnormal in this mutant. Therefore, we reasoned that RNAi knockdown that either suppresses or enhances *act-2(or295)* embryonic lethality may reveal regulators of both the spatial control and mechanical resilience of cytokinesis. Among the 483 genes that suppress *act-2(or295)* embryonic lethality, we performed two secondary screens: (i) a screen in a wild-type background for

regulators of spatial cytokinesis control; and (ii) a screen in the *act-2(or295)* background to uncover genes essential for furrow completion under heightened cortical contractility. We identified 71 candidate regulators of the spatial control of cytokinesis and 4 genes required for cytokinesis completion in an *act-2(or295)* background.

Results

act-2(or295) exhibits defects in three distinct modes of spatial control of cytokinesis

The *C. elegans* embryo is a suitable model system for studying the mechanisms underlying the spatial control of cytokinesis, as it exhibits an invariant cell lineage and three distinct modes of spatially controlled cytokinesis within an hour after fertilization (Figure 1A). In zygotes, the cleavage furrow forms asymmetrically, leading to asymmetric contractile ring closure (Figure 1A, left; Figure 1B, wild-type)(Maddox *et al.* 2007). At the 2-cell stage, the somatic blastomere AB undergoes rotation during cytokinesis (Figure 1A, middle; Figure 1C, wild-type) (Schierenberg and Junkersdorf 1992). This rotation positions the posterior daughter cell ABp on the dorsal side and is critical for the establishment of the dorsal-ventral axis (Priess and Thomson 1987; Schierenberg and Junkersdorf 1992). At the 6-cell stage, the endomesodermal precursor EMS undergoes asymmetric cell division (Goldstein 1992). We found that the contractile ring is initially formed in the equatorial region but becomes displaced posteriorly during cytokinesis (Figure 1A, right; Figure 1D, wild-type).

We used a temperature-sensitive and semi-dominant actin gain-of-function mutant, *act-2(or295),* which exhibits hypercontractility and stabilized actin filaments (Willis *et al.* 2006), to investigate the role of actin dynamics in the spatial control of cytokinesis. In

zygotes, *act-2(or295)* exhibited symmetric furrowing more often than the wild-type (Figure 1B and 1E). Additionally, cytokinesis often resulted in extra furrow formation (Figure 1B, bottom right), suggesting that the normal cytokinesis process is also compromised by this mutation. During AB cell division at the 2-cell stage, *act-2(or295)* embryos exhibited a reduced degree of rotation during cytokinesis (Figure 1C and 1F). During EMS cell division at the 6-cell stage, *act-2(or295)* displayed an abnormal shift of the contractile ring; some embryos showed no shift in position, whereas others showed an anterior shift (Figure 1D and 1G). In summary, *act-2(or295)* mutants display defects in all three modes of spatially controlled cytokinesis and also exhibit defects in proper contractile ring formation.

High-throughput RNAi screen in *act-2(or295)* mutant background

We reasoned that any RNAi knockdown that either suppresses or enhances *act-2(or295)* embryonic lethality might include regulators of the spatial control and mechanical resilience of cytokinesis. According to single-cell RNA-seq data, 5,928 genes are expressed in early *C. elegans* embryos (Tintori et al., 2016). Of these, 4,445 genes are included in the Ahringer feeding RNAi library (Kamath and Ahringer, 2003), and we performed a feeding RNAi screen targeting them (Figure 2A). Both wild-type and *act-2(or295)* mutants expressing an intestinal GFP reporter (*vha-6*p::GFP) were fed dsRNA-expressing bacteria at 20°C, a temperature at which *act-2(or295)* exhibits 80.2% embryonic lethality (n > 100). Instead of performing standard feeding RNAi on Nematode Growth Media (NGM) plates, we incubated the L1-stage larvae in liquid NGM supplemented with dsRNA-expressing bacteria (hereafter "liquid-based RNAi") in 96-

well plates (Figure 2A). We then counted the number of hatched worms in each well by imaging the intestinal GFP signal using high-content screening microscopy (Figure 2B-C). The effects of RNAi on embryonic viability in both wild-type and *act-2(or295)* strains were measured by computing Z-scores using the mean and standard deviation of worm counts within each 96-well plate, rather than using the global mean and standard deviation of the entire screen, to mitigate technical variation. For each of the 4,218 analyzed genes, we obtained three parameters: (1) Z-score in the control background, (2) Z-score in the *act-2(or295)* background, and (3) the difference between the two (Figure 3A).

We found that liquid-based feeding RNAi was consistently less effective than standard plate-based feeding RNAi, possibly due to the lower density of bacteria or differences in worm feeding behavior. Because of this weaker knockdown effect, we identified the essential cytokinesis regulator *nmy-2* (non-muscle myosin II) as a suppressor of *act-2(or295)* lethality in our screen (Figure 2C). Furthermore, we identified components of the RhoA signaling pathway (*ect-2/RhoGEF*, *rga-1/RhoGAP, cyk-1/formin*), structural components of the contractile ring (*act-1/-2/-3*, *ani-2/anillin*), and a regulator of branched actin assembly (*wsp-1/N-WASP*) as suppressors (Figure 3B).

In this primary screen, we set a less conservative threshold and designated any RNAi with a mutant Z-score ≥ 1.5 or a Z-score difference ≥ 1.5 as a suppressor of *act-2* lethality (Figure 3C). Conversely, RNAi knockdown resulting in failed hatching or significantly reduced hatching in the *act-2(or295)* background was classified as an enhancer. Hereafter, we focus only on *act-2(or295)* suppressors, as enhancers may include genes whose knockdown additively increased the lethality of *act-2(or295)*

without indicating a specific genetic interaction. We found that 310 genes displayed Z-

scores ≥ 1.5 in *act-2(or295)*, and 547 genes exhibited Z-score differences ≥ 1.5 (Figure

3C). In pursuit of genes directly regulating cytoskeletal dynamics or cell signaling, we

excluded those encoding DNA-binding, RNA-binding, and ribosomal proteins (which

likely affect general viability through transcription or translation). After this filtering, 388

genes had Z-score differences ≥ 1.5, and 210 genes showed Z-scores ≥ 1.5 in *act-*

2(or295), leading to a total of 483 *act-2* suppressor genes (Figure 3C).

Identification of regulators of spatial control of cytokinesis

Next, we performed a secondary screen to identify regulators of the spatial control

of cytokinesis among 483 genes characterized as *act-2(or295)* suppressors (Figure 4A).

For each RNAi knockdown, we collected three live-imaging movies capturing

embryogenesis from the 1-cell to the 7-cell stage.

For asymmetric contractile ring closure in zygote, we visually classified an

embryo as defective if the cleavage furrow ingressed asymmetrically. Because

asymmetric ring closure can occur in any orientation, furrowing asymmetry is not

detectable in our 2D imaging setup when the ring closure occurs along the imaging axis.

Indeed, about 23% of wild-type zygotes exhibited symmetric furrowing patterns (n = 22,

Figure 1E). Thus, we aimed to identify extreme cases that displayed a high penetrance of

defects. We scored phenotypes for each collected movie (score = 0 if defective, score = 1

if normal) and calculated a Z-score for each gene using the mean and standard deviation

across all *act-2* suppressors. We identified 36 genes exhibiting symmetric contractile ring

closure in all three movies (Z-score ≥ 1.96; Figure 4B). These candidates include *dhhc-*

9/ZDHHC16, which encodes S-palmitoyltransferase, and *cyn-6*/PPIB, which encodes

peptidylprolyl isomerase B (Figure 5A).

For the rotational cytokinesis during AB cell division, we measured the angle of

AB division axis rotation. We identified 13 genes with Z-scores > 1.96 (indicating

reduced rotation) and 3 gene with Z-score ≤ -1.96 (indicating excess rotation) (Figure 4C).

We found that *sem-5*/Grb2, involved in epidermal growth factor signaling, causes strong

defects in AB cell rotation (Figure 5B).

For the asymmetric contractile ring displacement during the EMS cell cytokinesis,

we calculated the shift of the cleavage furrow along the anterior-posterior axis before and

after cytokinesis for each movie. We identified 11 genes with Z-score ≥ 1.96 (indicating

less furrow displacement) and 11 genes with Z-score ≤ -1.96 (indicating excessive

displacement). These candidates include *pfd-1*/PFDN1, which encodes a prefoldin

subunit known to be involved in actin folding.

Finally, we performed a gene network analysis using the 71 identified genes and

found that 16 candidate genes—either genetically or physically, and directly or

indirectly— interact with known cytokinesis regulators (Figure 5D).

Identification of regulators of mechanical resilience of cytokinesis

In *act-2(or295)* embryos, which exhibit hypercontractility, cells are under constant

mechanical stress yet display only subtle defects in contractile ring formation (e.g., extra

furrow shown in Figure 1B) and show no defects in cytokinesis completion in zygotes

(Figure 6B). Thus, we reasoned that the genes required for cytokinesis in *act-2(or295)*

may include regulators of the mechanical resilience of cytokinesis. We selected 139

genes from the 483 *act-2* suppressors based on the following criteria: for genes on

Chromosome I and II, RNAi knockdown caused > 67% defects in asymmetric furrowing

in the wild-type background; for genes on the remaining chromosomes, RNAi

knockdown caused 100% embryonic lethality in *act-2(or295)* (Figure 6A). We then

performed an RNAi screen targeting these 139 genes in an *act-2* mutant background

using a standard plate-based feeding RNAi (Figure 13A). Of the 139 candidates

examined, RNAi knockdown of 12 genes led to mild to severe cytokinesis defects.

Among these, five genes are known cytokinesis regulators of cytokinesis: the actin genes

(*act-1, act-2, act-3*), *ect-2/RhoGEF* (which activates RhoA), and *nmy-2* (non-muscle

myosin II). Another three genes (*egg-6, cks-1, ubc-12*) are involved in eggshell formation,

cell cycle regulation, and ubiquitination, respectively, and are known to cause cytokinesis

defects when disrupted in wild-type background. Two additional genes (*tba-1*/α-tubulin

and the centrosomal/centriolar component *spd-2*) function in microtubule assembly and

organization.

The final two genes are relatively less understood. First, *evl-20/ARL2* encodes

ADP-ribosylation factor-like GTPase that localizes to the periphery of astral microtubules

and the plasma membrane (Antoshechkin and Han 2002). A previous study reported that

evl-20 knockdown or mutation reduces microtubule levels during embryogenesis and

produces multinucleated cells later in development, such as in vulval precursor cells and

the somatic gonad (Antoshechkin and Han 2002). However, the early embryonic

phenotype was not described, and cytokinesis has not been analyzed in detail. *lpin-*

1/Lipin 1 encodes phosphatidic acid phosphatase involved in polyunsaturated fatty acids

(PUFAs) production and longevity (Jung *et al.* 2020). Previous studies showed that *lpin-1*

is required for proper endoplasmic reticulum organization and timely nuclear envelope breakdown during embryogenesis (Golden *et al.* 2009; Gorjánácz and Mattaj 2009). In *Toxoplasma gondii*, the loss of Lipin results in multinucleation, possibly due to cytokinesis failure (Dass *et al.* 2021).

To identify genes specifically required for cytokinesis in *act-2(or295)*, we compared RNAi phenotypes between wild-type and *act-2* backgrounds and found that *evl-20*, *lpin-1*, *spd-2*, and *tba-1* induced more severe cytokinesis defects in *act-2(or295)* (Figure 6B). The phenotypes of *spd-2(RNAi);act-2(or295)* and *tba-1(RNAi); act-2(or295)* were relatively mild, as these embryos still formed a furrow that sometimes regressed later (Figure 6C, bottom). By contrast, *evl-20(RNAi);act-2(or295)* and *lpin-1(RNAi); act-2(or295)* failed to form a fully developed furrow, suggesting that these genes are involved in contractile ring formation (Figure 6C).

Discussion

In this study, we found that the gain-of-function actin mutant *act-2(or295)* exhibits defects in the spatial control of cytokinesis. By leveraging this genetic background, we conducted a high-throughput RNAi screen and identified 483 suppressors of *act-2(or295)* embryonic lethality. In a secondary, live-imaging-based screen in a wild-type background, we identified 36 genes involved in asymmetric contractile ring closure in zygotes, 16 genes involved in rotational cytokinesis in the 2-cell stage AB cell, and 22 genes involved in asymmetric contractile ring positioning in the 6-cell stage EMS cell. Furthermore, we conducted another secondary RNAi screen in *act-2(or295)* background

to search for genes enabling successful cytokinesis under mechanical stress. We

identified four genes that exhibit cytokinesis defects in *act-2(or295).*

Although the position and function of the contractile ring are spatially controlled

in tissues, the mechanisms underlying this spatial control remain unclear. Through our

RNAi screen, we identified potential regulators for each of the three modes of spatial

cytokinesis control. Among the 71 candidate genes, only three genes were common to at

least two modes of spatial control. Some of the identified genes are known to be involved

in cytokinesis. For example, we identified *dhhc-9*/ZDHHC16, which encodes S-

palmitoyltransferase, as a regulator of asymmetric contractile ring closure (Figure 5A). A

recent study in mammalian cultured cells shows that ZDHHC5 regulates the

palmitoylation of protocadherin 7 (PCDH7) during mitosis, thereby targeting its cleavage

furrow localization and activating RhoA (Ozkan *et al.* 2023). Disruption of PCDH7

increases cytokinesis failure, but its role in asymmetric contractile ring closure remains

unexplored. We also identified *sem-5*/Grb2 as a regulator of rotational cytokinesis

(Figure 5B). Grb2 is a protein containing Src homology domain 2 (SH2) and SH3

domains, acting as an adaptor in various cell signaling pathways, such as FGF and EGF

signaling (Giubellino *et al.* 2008). Grb2 interacts with Wiskott-Aldrich Syndrome protein

(WASp) and with cortactin, both of which regulate the branched actin nucleator Arp2/3

complex (She *et al.* 1997; Martinez-Quiles *et al.* 2004). Interestingly, our candidates for

rotational cytokinesis regulators also include the N-WASp homolog *wsp-1*. The list of

candidate genes generated in this study should inspire novel hypotheses by integrating

this existing molecular knowledge.

The contractile ring is a molecular machine known for its remarkable resilience, yet the mechanisms underlying this resilience of cytokinesis are not yet fully understood. Our secondary screen in *act-2(or295)* mutants identified four genes that contribute to cytokinesis completion in this background. Genes, such as *tba-1*, *spd-2*, and *evl-20* are involved in microtubule assembly. The mitotic spindle plays a critical role in activating RhoA through the central spindle and astral microtubules (Pollard and O'Shaughnessy 2019). A previous small-scale RNAi screen in *C. elegans* also identified microtubule regulators as genes required for the mechanical resilience of cytokinesis (Singh *et al.* 2019). That experimental setup was more specific, subjecting embryos to uniaxial compression using small-diameter beads and a coverslip. Thus, our identification of these microtubule regulators confirms their requirement in the mechanical resilience of cytokinesis. We also identified the Lipin homolog *lpin-1* as a regulator of mechanical resilience of cytokinesis. The specific role of Lipin in contractile ring formation and function has yet to be determined and will be an interesting direction for future research.

It is worth noting that this study has three main limitations. First, the primary RNAi screen, which analyzed 4,218 early embryonic genes, was performed only once. Consequently, some pseudonegatives may have gone undetected due to technical errors. Second, both secondary RNAi screens used feeding RNAi with relatively small sample sizes, limiting our ability to fully elucidate the roles of candidate genes in cytokinesis. Finally, *act-2(or295)* enhancers likely include other interesting candidates, but we did not analyze them in order to streamline the study.

Overall, this study identified a number of potentially novel regulators of cytokinesis, with a particular focus on the relatively underexplored areas of cytokinesis

research, mechanical resilience and spatial control. The identified list of genes will facilitate the generation of future hypotheses aimed at understanding the multicellular-specific regulation of cytokinesis during animal development.

Acknowledgements

We thank Caenorhabditis Genetics Center (funded by the NIH Office of Research Infrastructure Programs; P40 OD010440) for sharing worm strains and the Sugioka lab members for general discussions. We also thank Life Sciences Institute Imaging Core Facility, Michele Roberge, Aruna Balgi for sharing high-content screening microscopy, Don Moerman and Mark Edgley for sharing liquid handling machine. This work was supported by the Canadian Institutes of Health Research (Project Grant; PJT-169145), Government of Canada's New Frontiers in Research Fund (NFRFE-2019-00310), and the Michael Smith Health Research BC (Scholar Award; SCH-2020-0406) to K.S.

Author contributions

C.G.: Formal analysis, Investigation, Data curation, Writing-Original Draft, Validation, Visualization. **Y.R.X.**: Formal analysis, Investigation, Data Curation, Project administration. **A. H.**: Formal analysis, Investigation, Methodology, Data Curation, Project administration. **C.R.H.**: Investigation. **V. J.**: Data Curation. **M.J.K.** Investigation. **K.D.** Investigation. **K.S.**: Conceptualization, Formal analysis, Investigation, Methodology, Validation, Writing-Original Draft, Writing-Review & Editing, Supervision, Funding acquisition.

Declaration of interests

We declare no competing interests.

Figure legends

Figure 1. *act-2(or295)* mutant exhibits abnormal spatial control of

cytokinesis.

(**A**) Three modes of spatial control of cytokinesis during animal development. (**B**)

Asymmetric contractile ring closure in zygotes. The contractile ring closes

asymmetrically in wild-type embryos but is defective in *act-2(or295)*. *act-2(or295)*

mutants also exhibit extra furrow phenotypes at later stage of cytokinesis. (**C**) Rotational

cytokinesis in the AB cell. At the two-cell stage, the AB cell normally rotates during

cytokinesis, but this rotation is defective in *act-2(or295)*. Dotted white lines indicate

mitotic spindle position. (**D**) Asymmetric positioning of the contractile ring in the EMS

cell. At the 6-cell stage, the EMS forms a cleavage furrow in the equatorial region, which

then shifts towards the posterior. In *act-2(or295),* this posterior furrow shift is often

defective. (**E**-**G**) Quantification of zygote furrow asymmetry (E), cellular rotation during

AB cell division (F), and posterior furrow displacement during EMS cell division (G).

Yellow arrowheads indicate the furrow position. Scale bars represent 10 µm. Error bars

represent 95% confidence intervals. Fisher's exact test was performed in panel E and

Welch's t-test was used for panels F and G. P-values: "****", "**", "*"and "ns" indicate

$p < 0.0001$, $p < 0.01$, $p < 0.05$, and $p > 0.05$, respectively.

Figure 2. High-throughput RNAi screen using *act-2(or295)* mutants.

(**A**) Design of the RNAi screen. Wild-type and *act-2(or295)* mutants expressing an

intestinal GFP reporter were bleached, and purified embryos were hatched without food

to induce starvation-induced L1 larval arrest. The synchronized L1 larvae were cultured

in liquid NGM containing dsRNA expressing feeding RNAi bacteria. Hatched F1 larvae

were then imaged using high-content screening microscopy. (**B**) Intestinal GFP reporter

signals for the control and one suppressor, *nmy-2(RNAi)*. Segmented images were used to

count the number of hatched worms. (**C**) The number of hatched worms in a 96-well

plate. *nmy-2(RNAi)* resulted in a significantly higher number of hatched worms,

identified it as a suppressor of *act-2(or295)*.

Figure 3. Summary of the primary RNAi screen.

(**A**) Scatter plot of Z-scores for 4,218 analyzed genes. Z-scores in the control strain and

act-2 mutants are shown on the x- and y-axes, respectively. The difference in Z-scores

between control and *act-2* is indicated by the color code. (**B**) Scatter plot of Z-scores

highlighting known regulators of cytokinesis. As indicated by red, many of them were

identified as suppressors. (**C**) Summary of identified actin suppressors. Among the non-

DNA/RNA-binding and non-ribosomal genes, 388 and 210 suppressors were identified

based on the Z-score difference and the Z-score in *act-2* mutants, respectively. A total of

483 genes were used for the secondary screens.

Figure 4. Secondary screen for spatial cytokinesis regulators.

(**A**) Design of the secondary screen. A total of 483 *act-2* suppressor genes were targeted

by standard plate-based feeding RNAi in a wild-type background. For asymmetric

furrowing in zygotes, we performed a qualitative assessment, whereas for AB cell

rotation and EMS cell furrow shift, we measured the degree of rotation and the extent of

furrow shift, respectively. (**B-D**) Candidates regulators of asymmetric contractile ring

closure (B), rotational cytokinesis (C), and asymmetric furrow shift (D). Orange bars

indicate extreme samples with Z-scores $\geq$ 1.96 or $\leq$ -1.96.

Figure 5. Identification of spatial cytokinesis regulators.

(**A**) Defects in asymmetric furrowing in *dhhc-9(RNAi)*. (**B**) Defects in rotational

cytokinesis in *sem-5(RNAi)*. (**C**) Defects in asymmetric furrow displacement in *pfd-*

1(RNAi). (**D**) Gene interaction network among identified candidate genes and known

cytokinesis regulators. Protein-protein interactions (PPI), genetic interactions, shared

domain information are shown. Candidates without known interactions are not displayed.

Scale bar indicates 10 µm.

Figure 6. Identification of genes required for cytokinesis in the *act-2(or295)*

background.

(**A**) Design of the secondary screen. A total of 139 genes were selected based on defects

in asymmetric furrowing in zygotes or 100% embryonic lethality in the *act-2* mutant

background from 483 *act-2* suppressor genes. These genes were knocked-down in *act-*

2(or295) mutants and tested for their requirement in cytokinesis completion. (**B**) Genes

identified as required for cytokinesis completion in *act-2(or295)* mutants. P-values were

calculated by chi-square test, followed by Holm-Sidak adjustment for multiple

comparisons. P-values: "**", "*", and "ns" indicate $p < 0.01$, $p < 0.05$, and $p > 0.05$,

respectively. (**C**) Cytokinesis phenotype after candidate gene knockdown. White dotted

lines indicate the mitotic spindle position, and yellow arrowheads indicate the cleavage

furrow. Scale bars, 10 µm.

7
Materials and Methods

Maintenance of worm strains

All *C. elegans* strains were cultured and maintained using standard methods

(Brenner 1974). The transgene, *oxTi872*[*vha-6p*::GFP::*tbb-2* 3'UTR + Cbr-*unc-119*(+)],

was used to monitor intestinal GFP expression. Additionally, *act-2(or295)* mutants were

used as our sensitized genetic background. Both the *act-2(or295)* mutants and wild-type

strains carrying *oxTi872* were maintained on NGM agar plates at 15˚C. During the liquid

culture RNAi screen, these strains were incubated at 20˚C, the semi-permissive

temperature for *act-2(or295)* mutant strain. For both secondary screens, these strains

were incubated at 25˚C, the restrictive temperature for *act-2(or295)*.

RNAi

Feeding RNAi was performed using the Ahringer RNAi library (Fraser *et al.*

2000; Ahringer 2006). The expression of dsRNA in the *Escherichia coli* strain HT115

was induced by overnight incubation at 37˚C in Nematode Growth Medium (NGM)

containing 10 µg/mL Ampicillin and 1 mM isopropyl ß-D-1-thiogalactopyranoside

(IPTG).

For the high-throughput RNAi screen, gravid adults were bleached with

hypochlorite solution, and the resulting purified embryos were incubated in M9 buffer

overnight to obtain developmentally arrested L1 stage worms. These worms were

suspended in liquid NGM and approximately 4-5 L1 worms were transferred into each

well of 96-well plates containing various dsRNA-expressing bacteria. After 5 days of

incubation at 20˚C, the 96-well plates were imaged using a high-content screening system

(Cellomics Arrayscan; Thermo Fisher).

For the two secondary RNAi screens, dsRNA-expressing bacteria were growin on

solid NGM agar plates. Developmentally arrested L1 worms were transferred onto these

plates and incubated for 3 days at 25˚C. For genes required for fertility, we used L4 stage

worms were used and incubated overnight before live imaging. Wild-type worms and

act-2(or295) mutants were used for the spatial cytokinesis regulator screen and the

cytokinesis resilience regulator screen, respectively. A bacterial strain containing the

empty vector L4440 served as a negative control in both secondary screens.

Live-imaging

To obtain embryos for live imaging by brightfield microscopy, we dissected

gravid adults on a coverslip in a 5 µl droplet of M9 buffer. We then placed the coverslip

gently onto a 2% agarose pad mounted on a glass slide and sealed the edges with

petroleum jelly (Vaseline) to prevent dehydration.

For characterization of the strains used in the high-throughput RNAi screen and

secondary screens, we performed live imaging at room temperature (~22.5ºC) using

brightfield compound microscopes (B290-TB; OPTIKA), equipped with a 100X oil

objective lens (1.2NA) and controlled by OPTIKA Vision Lite software. For the

secondary screen of spatial cytokinesis regulators, images were taken every 2 seconds for

60 minutes, beginning at the 1-cell stage and continuing until after EMS cell division. For

the secondary screen of cytokinesis resilience regulators, images were taken every 2

seconds for 10 minutes, beginning at the1-cell stage and ending at the 2-cell stage. In

most cases, filming began during late pronuclear migration stage but before nuclear

envelope breakdown (NEBD).

Image analysis for the high-throughput RNAi screen

To quantify the number of hatched worms, we processed images obtained from

the high-content screening system using Fiji (Schindelin *et al.* 2012). We applied a

Laplacian filter to extract edge information while removing background signals. Next, we

used Otsu's threshold to identify relevant signals and then applied morphological filters

to select rod-like worm bodies while excluding circular objects, such as embryos and

debris. The resulting segmentation data was used to count the number of hatched F1 and

F2 worms. We then standardized the worm counts for each 96-well plate by calculating

Z-scores, using the mean and standard deviation of worm counts within each plate to

minimize batch effects. We computed three different Z-scores for each candidate gene:

one in the control background, one in the *act-2(or295)* mutant background, and the

difference between the two. Genes with Z-scores greater than 1.5 in the *act-2(or295)*

mutant background were considered actin suppressors. Additionally, genes with a Z-score

difference greater than 1.5 between act-2 and control strains were also designated as actin

suppressors.

Selection of candidate genes for secondary screens

For the secondary screen of spatial regulators of cytokinesis, all actin suppressor

genes were assessed using plate-based RNAi in a wild-type strain (EG8829). At least

three embryos were imaged for each gene, and phenotypes were manually annotated in

Fiji using the collected live-imaging movies.

For the secondary screen of regulators of cytokinesis resilience, we first selected

actin suppressor genes using two different criteria. Initially, we chose genes based on

defects in asymmetric ring closure in 1-cell stage zygotes, as identified in the spatial

cytokinesis regulator screen. However, we found that most of these genes were not

required for cytokinesis in the *act-2* mutant background. Therefore, we selected genes

whose RNAi knockdown cased 100% embryonic lethality in *act-2* mutants at 25˚C for

genes on chromosomes III-X.

Bioinformatics

19 Scatter plots for the high-throughput screening data were generated using

20 Matplotlib. To analyze and visualize gene networks among the identified candidates, we

21 used GeneMANIA in the Cytoscape software (Shannon *et al.* 2003).

Statistics

Statistical analyses were performed using Prism9 (GraphPad Software) and

custom Python scripts. For Figure 1E, we performed Fisher's exact test on raw data

(rather than percentages). For Figure 1F-G, we used Welch's t-test. For Figure 6B, we

used Fisher's exact test on raw data. The resulting P-values were corrected for multiple

comparisons using a custom Python script (Holm-Sidak method). The symbols such as

****, ***, **, *, and ns indicate $p < 0.0001$, $p < 0.001$, $p < 0.01$, $p < 0.05$, and $p > 0.05$,

respectively.

References:

Ahringer   J.,   2006   Reverse   genetics.   WormBook.

12   https://doi.org/10.1895/wormbook.1.47.1

13   Antoshechkin I., and M. Han, 2002 The C. elegans evl-20 gene is a homolog of the small

14   GTPase ARL2 and regulates cytoskeleton dynamics during cytokinesis and

15   morphogenesis. Dev. Cell 2: 579–591.

16   Basant A., and M. Glotzer, 2018 Spatiotemporal Regulation of RhoA during Cytokinesis.

17   Curr. Biol. 28: R570–R580.

18   Brenner S., 1974 The genetics of Caenorhabditis elegans. Genetics 77: 71–94.

Dass S., S. Shunmugam, L. Berry, C.-S. Arnold, N. J. Katris, *et al.*, 2021 Toxoplasma
LIPIN is essential in channeling host lipid fluxes through membrane biogenesis
and lipid storage. Nat. Commun. 12: 2813.

Fraser A. G., R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann, *et al.*,
2000 Functional genomic analysis of C. elegans chromosome I by systematic
RNA interference. Nature 408: 325–330.

Frémont S., and A. Echard, 2018 Membrane traffic in the late steps of cytokinesis. Curr.
Biol. 28: R458–R470.

Giubellino A., T. R. Burke Jr, and D. P. Bottaro, 2008 Grb2 signaling in cell motility and
cancer. Expert Opin. Ther. Targets 12: 1021–1033.

Golden A., J. Liu, and O. Cohen-Fix, 2009 Inactivation of the C. elegans lipin homolog
leads to ER disorganization and to defects in the breakdown and reassembly of
the nuclear envelope. J. Cell Sci. 122: 1970–1978.

Goldstein B., 1992 Induction of gut in Caenorhabditis elegans embryos. Nature 357:
255–257.

Gorjánácz M., and I. W. Mattaj, 2009 Lipin is required for efficient breakdown of the
nuclear envelope in Caenorhabditis elegans. J. Cell Sci. 122: 1963–1969.

Herszterg S., A. Leibfried, F. Bosveld, C. Martin, and Y. Bellaiche, 2013 Interplay
between the Dividing Cell and Its Neighbors Regulates Adherens Junction
Formation during Cytokinesis in Epithelial Tissue. Dev. Cell 24: 256–270.

Jankele R., R. Jelier, and P. Gönczy, 2021 Physically asymmetric division of the C.

elegans zygote ensures invariably successful embryogenesis. Elife 10.

https://doi.org/10.7554/eLife.61714

Jung Y., S. Kwon, S. Ham, D. Lee, H.-E. H. Park, *et al.*, 2020 Caenorhabditis elegans

Lipin 1 moderates the lifespan-shortening effects of dietary glucose by

maintaining ω-6 polyunsaturated fatty acids. Aging Cell 19: e13150.

Kuroda R., B. Endo, M. Abe, and M. Shimizu, 2009 Chiral blastomere arrangement

dictates zygotic left-right asymmetry pathway in snails. Nature 462: 790–794.

Maddox A. S., L. Lewellyn, A. Desai, and K. Oegema, 2007 Anillin and the septins

promote asymmetric ingression of the cytokinetic furrow. Dev. Cell 12: 827–835.

Martinez-Quiles N., H.-Y. H. Ho, M. W. Kirschner, N. Ramesh, and R. S. Geha, 2004

Erk/Src phosphorylation of cortactin acts as a switch on-switch off mechanism

that controls its ability to activate N-WASP. Mol. Cell. Biol. 24: 5269–5280.

Meshcheryakov V. N., and L. V. Beloussov, 1975 Asymmetrical rotations of blastomeres

in early cleavage of gastropoda. Wilehm Roux Arch Dev Biol 177: 193–203.

Mierzwa B., and D. W. Gerlich, 2014 Cytokinetic abscission: molecular mechanisms and

temporal control. Dev. Cell 31: 525–538.

Naganathan S. R., S. Furthauer, M. Nishikawa, F. Julicher, and S. W. Grill, 2014 Active

torque generation by the actomyosin cell cortex drives left-right symmetry

breaking. Elife 3: e04165.

Ozkan N. E., B. N. Yigit, B. S. Degirmenci, M. H. Qureshi, G. N. Yapici, *et al.*, 2023 Cell cycle-dependent palmitoylation of protocadherin 7 by ZDHHC5 promotes successful cytokinesis. J. Cell Sci. 136: jcs260266.

Piekny A. J., and A. S. Maddox, 2010 The myriad roles of Anillin during cytokinesis. Semin. Cell Dev. Biol. 21: 881–891.

Pollard T. D., 2017 Nine unanswered questions about cytokinesis. J. Cell Biol. 216: 3007–3016.

Pollard T. D., and B. O'Shaughnessy, 2019 Molecular Mechanism of Cytokinesis. Annu. Rev. Biochem. 88: 661–689.

Priess J. R., and J. N. Thomson, 1987 Cellular interactions in early C. elegans embryos. Cell 48: 241–250.

Schierenberg E., and B. Junkersdorf, 1992 The role of eggshell and underlying vitelline membrane for normal pattern formation in the early C. elegans embryo. Rouxs. Arch. Dev. Biol. 202: 10–16.

Schindelin J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, *et al.*, 2012 Fiji: an open-source platform for biological-image analysis. Nat. Methods 9: 676–682.

Shannon P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, *et al.*, 2003 Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res. 13: 2498–2504.

She H. Y., S. Rockow, J. Tang, R. Nishimura, E. Y. Skolnik, *et al.*, 1997 Wiskott-Aldrich
syndrome protein is associated with the adapter protein Grb2 and the epidermal
growth factor receptor in living cells. Mol. Biol. Cell 8: 1709–1721.

Silva A. M., D. S. Osório, A. J. Pereira, H. Maiato, I. M. Pinto, *et al.*, 2016 Robust gap
repair in the contractile ring ensures timely completion of cytokinesis. J. Cell Biol.
215: 789–799.

Singh D., D. Odedra, P. Dutta, and C. Pohl, 2019 Mechanical stress induces a scalable
switch in cortical flow polarization during cytokinesis. J. Cell Sci. 132: jcs231357.

Sugioka K., 2021 Symmetry-breaking of animal cytokinesis. Semin. Cell Dev. Biol.
https://doi.org/10.1016/j.semcdb.2021.12.008

Taulet N., B. Vitre, C. Anguille, A. Douanier, M. Rocancourt, *et al.*, 2017 IFT proteins
spatially control the geometry of cleavage furrow ingression and lumen
positioning. Nat. Commun. 8: 1928.

Willis J. H., E. Munro, R. Lyczak, and B. Bowerman, 2006 Conditional dominant
mutations in the Caenorhabditis elegans gene act-2 identify cytoplasmic and
muscle roles for a redundant actin isoform. Mol. Biol. Cell 17: 1051–1064.

Wood W. B., 1991 Evidence from reversal of handedness in C. elegans embryos for early
cell interactions determining cell fates. Nature 349: 536–538.

Figure 1. Actin gain-of-function mutant exhibits abnormal spatial control of cytokinesis.

(**A**) Three modes of spatial control of cytokinesis during animal development. (**B**) Asymmetric contractile ring closure in zygotes. The contractile ring closes asymmetrically in wild-type embryos but is defective in *act-2(or295)*. *act-2(or295)* mutants also exhibit extra furrow phenotypes at later stage of cytokinesis. (**C**) Rotational cytokinesis in the AB cell. At the two-cell stage, the AB cell normally rotates during cytokinesis, but this rotation is defective in *act-2(or295)*. Dotted white lines indicate mitotic spindle position. (**D**) Asymmetric positioning of the contractile ring in the EMS cell. At the 6-cell stage, the EMS forms a cleavage furrow in the equatorial region, which then shifts towards the posterior. In *act-2(or295)*, this posterior furrow shift is often defective. (**E**-**G**) Quantification of zygote furrow asymmetry (E), cellular rotation during AB cell division (F), and posterior furrow displacement during EMS cell division (G). Yellow arrowheads indicate the furrow position. Scale bars represent 10 µm. Error bars represent 95% confidence intervals. Fisher's exact test was performed in panel E and Welch's t-test was used for panels F and G. P-values: "****", "**", "*"and "ns" indicate p < 0.0001, p < 0.01, p < 0.05, and p > 0.05, respectively.

Figure 2. High-throughput RNAi screen using *act-2(or295)* mutants.

(**A**) Design of the RNAi screen. Wild-type and *act-2(or295)* mutants expressing an intestinal GFP reporter were bleached, and purified embryos were hatched without food to induce starvation-induced L1 larval arrest. The synchronized L1 larvae were cultured in liquid NGM containing dsRNA expressing feeding RNAi bacteria. Hatched F1 larvae were then imaged using high-content screening microscopy. (**B**) Intestinal GFP reporter signals for the control and one suppressor, *nmy-2(RNAi)*. Segmented images were used to count the number of hatched worms. (**C**) The number of hatched worms in a 96-well plate. *nmy-2(RNAi)* resulted in a significantly higher number of hatched worms, identified it as a suppressor of *act-2(or295)*.

Figure 3. Gough et al.

Figure 3. Summary of the primary RNAi screening.
(**A**) Scatter plot of Z-scores for 4,218 analyzed genes. Z-scores in the control strain and act-2 mutants are shown on the x- and y-axes, respectively. The difference in Z-scores between control and act-2 is indicated by the color code.
(**B**) Scatter plot of Z-scores highlighting known regulators of cytokinesis. As indicated by red, many of them were identified as suppressors. (**C**) Summary of identified actin suppressors. Among the non-DNA/RNA-binding and non-ribosomal genes, 388 and 210 suppressors were identified based on the Z-score difference and the Z-score in act-2 mutants, respectively. A total of 483 genes were used for the secondary screens.

Figure 4. Secondary screen for spatial cytokinesis regulators.

(**A**) Scatter plot of Z-scores for 4,218 analyzed genes. Z-scores in the control strain and act-2 mutants are shown on the x- and y-axes, respectively. The difference in Z-scores between control and act-2 is indicated by the color code. (**B**) Scatter plot of Z-scores highlighting known regulators of cytokinesis. As indicated by red, many of them were identified as suppressors. (**C**) Summary of identified actin suppressors. Among the non-DNA/RNA-binding and non-ribosomal genes, 388 and 210 suppressors were identified based on the Z-score difference and the Z-score in act-2 mutants, respectively. A total of 483 genes were used for the secondary screens.

Figure 5. Gough et al.

A Cytokinesis in zygotes
dhhc-9(RNAi)

early

cyn-6(RNAi)

early

Symmetric furrowing

B Cytokinesis in AB cell
sem-5(RNAi)

anaphase

early

No rotation

C Cytokinesis in EMS cell
pfd-1(RNAi)

early

late

No furrow displacement

D

Legend:
- Known cytokinesis regulators (black)
- Identified candidates (yellow)
- Protein-protein interaction (blue)
- Genetic interaction (red)
- Shared domain (yellow)

Figure 5. Identification of spatial cytokinesis regulators.

(**A**) Defects in asymmetric furrowing in *dhhc-9(RNAi)*. (**B**) Defects in rotational cytokinesis in *sem-5(RNAi)*. (**C**) Defects in asymmetric furrow displacement in *pfd-1(RNAi)*. (**D**) Gene interaction network among identified candidate genes and known cytokinesis regulators. Protein-protein interactions (PPI), genetic interactions, shared domain information are shown. Candidates without known interactions are not displayed. Scale bar indicates 10 µm.

Figure 6. Identification of genes required for cytokinesis in *act-2(or295)* background

(**A**) Design of the secondary screen. A total of 139 genes were selected based on defects in asymmetric furrowing in zygotes or 100% embryonic lethality in the act-2 mutant background from 483 act-2 suppressor genes. These genes were knocked-down in act-2(or295) mutants and tested for their requirement in cytokinesis completion. (**B**) Genes identified as required for cytokinesis completion in act-2(or295) mutants. P-values were calculated by chi-square test, followed by Holm-Sidak adjustment for multiple comparisons. P-values: "**", "*", and "ns" indicate p < 0.01, p < 0.05, and p > 0.05, respectively. (**C**) Cytokinesis phenotype after candidate gene knockdown. White dotted lines indicate the mitotic spindle position, and yellow arrowheads indicate the cleavage furrow. Scale bars, 10 μm.