Identification of Genes Required for Spatial Control and Mechanical Resilience of Cytokinesis during *Caenorhabditis elegans* Embryogenesis

-Author names and Affiliations

Chelsey Lynn Gough[1,2,3], Yuxuan Rain Xiong[1,2,3], Aoi Hiroyasu[1,2,3], Christina Rou Hsu[1,2],

Viktorija Juciute[1,2], MinJee Kim[1,2], Kalen Dofher[1,2], and Kenji Sugioka[1,2*]

[1]Life Sciences Institute, The University of British Columbia, 2350 Health Sciences Mall,

Vancouver, BC V6T1Z3, Canada

[2]Department of Zoology, The University of British Columbia, 2350 Health Sciences Mall,

Vancouver, BC V6T1Z3, Canada

[3]These authors contributed equally to this work.

*Corresponding author

Kenji Sugioka

Life Sciences Institute, Department of Zoology, The University of British Columbia, 2350

Health Sciences Mall, Vancouver, BC V6T1Z3, Canada

kenji.sugioka@ubc.ca

Phone: +1-604-822-4628

1 Abstract

Cytokinesis is the final step of cell division, in which the dividing cell is physically

separated into two daughter cells by the contractile ring. The contractile ring is a highly

resilient molecular machine that can function properly under mechanical stress.

Additionally, its function, position, and orientation are spatially modulated in developing

animals to regulate morphogenesis. Although essential regulators of cytokinesis have been

identified through previous genetic screens, the molecular mechanisms underlying these

spatial controls and the mechanical resilience of cytokinesis remain elusive. To identify

cytokinesis regulators involved in these processes, we performed a high-throughput RNAi

screen using a gain-of-function mutant of actin that exhibits ectopic cortical contraction

and abnormal spatial control of cytokinesis in *Caenorhabditis elegans* embryos. We

obtained a list of early embryonic genes that suppress embryonic lethality in an *act-2*

mutant background. Two parallel secondary screens of candidate genes were conducted.

The first secondary screen in a wild-type background identified 69 candidate genes

regulating spatial cytokinesis control—asymmetric ring closure, positioning, and

rotation—during early embryogenesis. The second secondary screen in the *act-2(or295)*

background identified four genes required for cytokinesis in this background, including

microtubule regulators, *evl-20/ARL2*, and *lpin-1/Lipin1*. This study will serve as a useful

resource for the development of future hypotheses and provide insights into the precise

regulation of cytokinesis in tissues.

Short title: Spatial and Robust Cytokinesis Control in *C. elegans*

Keywords: Cytokinesis, C. elegans, Contractile ring, Cell division

Introduction

Cytokinesis is the final step of cell division, in which the contractile ring physically separates the dividing cell into two daughter cells (Pollard and O'Shaughnessy 2019). The contractile ring is a mechanically resilient molecular machine capable of withstanding mechanical stress and repairing physical damage (Silva *et al.* 2016; Singh *et al.* 2019). Additionally, its orientation and position are spatially regulated in three distinct ways during morphogenesis (Sugioka 2021). First, asymmetric contractile ring closure, often found in zygotes and epithelial tissue, is critical for maintaining cytokinesis resilience, epithelial integrity, and polarity (Maddox *et al.* 2007; Herszterg *et al.* 2013; Taulet *et al.* 2017). Second, cell rotation during cytokinesis is required for dorsal-ventral axis establishment in *C. elegans* (Priess and Thomson 1987; Schierenberg and Junkersdorf 1992) and is implicated in the establishment of animal left-right asymmetry in pond snails and *C. elegans* (Meshcheryakov and Beloussov 1975; Wood 1991; Kuroda *et al.* 2009; Naganathan *et al.* 2014). Finally, asymmetric positioning of the contractile ring during asymmetric cell division is critical for generating unequal daughter cell sizes, which is required for subsequent cell fate decisions (Jankele *et al.* 2021). Despite their significance, the regulators of mechanical resilience and the spatial control of cytokinesis remain largely unknown.

Decades of studies have revealed the essential components of the contractile ring, which are evolutionarily conserved in eukaryotes (Pollard 2017). First, the small GTPase RhoA is activated by the centralspindlin complex, thereby initiating contractile ring assembly via stimulation of downstream effectors such as formin and Rho-associated

kinase (Basant and Glotzer 2018). Contractile ring function also requires structural

components such as anillin (Piekny and Maddox 2010). As the contractile ring constricts

through actin-myosin sliding, membrane trafficking pathways contribute to cytokinesis by

delivering additional membrane to the ingressing furrow to accommodate the growing cell

surface area (Frémont and Echard 2018). Finally, the daughter cells separate via ESCRT-

III-mediated abscission at the midbody (Mierzwa and Gerlich 2014). Previous forward

genetic and RNAi-based screens in yeast and various animal systems were typically

performed in wild-type backgrounds and focused on identifying essential regulators of

cytokinesis (Eggert et al. 2004; Echard et al. 2004; Sönnichsen et al. 2005). A recent study

in *C. elegans* performed a small-scale candidate screen in zygotes under uniaxial

compression and identified several regulators required for the mechanical resilience of

cytokinesis (Singh *et al.* 2019). However, a large-scale genetic screen focusing on the

physical and developmental control of cytokinesis is still needed.

In this study, we performed an RNAi screen targeting 4,218 early embryonic genes

in an actin gain-of-function mutant of *C. elegans*. The *act-2(or295)* mutation lies near the

ATP-binding pocket, leading to filament stabilization and hyperactive cortical contractility

(Willis *et al.* 2006). We found that the spatial control of cytokinesis, such as asymmetric

ring closure in the zygote, rotational cytokinesis in the 2-cell stage, and asymmetric ring

positioning in the 6-cell stage, is also abnormal in this mutant. Therefore, we reasoned that

RNAi knockdown that either suppresses or enhances *act-2(or295)* embryonic lethality may

reveal regulators of both the spatial control and mechanical resilience of cytokinesis.

Among the 689 genes that suppress *act-2(or295)* embryonic lethality, we performed two

secondary screens on 385 selected candidates: (i) a screen in a wild-type background for

regulators of spatial cytokinesis control; and (ii) a screen in the *act-2(or295)* background to uncover genes essential for furrow completion under heightened cortical contractility. We identified 69 candidate regulators of the spatial control of cytokinesis and 4 genes required for cytokinesis completion in an *act-2(or295)* background.

Results

act-2(or295) exhibits defects in three distinct modes of spatial control of cytokinesis

The *C. elegans* embryo is a suitable model system for studying the mechanisms underlying the spatial control of cytokinesis, as it exhibits an invariant cell lineage and three distinct modes of spatially controlled cytokinesis within an hour after fertilization (Figure 1A). In zygotes, the cleavage furrow forms asymmetrically, leading to asymmetric contractile ring closure (Figure 1A, left; Figure 1B, wild-type)(Maddox *et al.* 2007). At the 2-cell stage, the somatic blastomere AB undergoes rotation during cytokinesis (Figure 1A, middle; Figure 1C, wild-type) (Schierenberg and Junkersdorf 1992). This rotation positions the posterior daughter cell ABp on the dorsal side and is critical for the establishment of the dorsal-ventral axis (Priess and Thomson 1987; Schierenberg and Junkersdorf 1992). At the 6-cell stage, the endomesodermal precursor EMS undergoes asymmetric cell division (Goldstein 1992). We found that the contractile ring is initially formed in the equatorial region but becomes displaced posteriorly during cytokinesis (Figure 1A, right; Figure 1D, wild-type).

We used a temperature-sensitive and semi-dominant actin gain-of-function mutant, *act-2(or295),* which exhibits hypercontractility and stabilized actin filaments (Willis *et al.* 2006), to investigate the role of actin dynamics in the spatial control of cytokinesis. In

zygotes, *act-2(or295)* exhibited symmetric furrowing more often than the wild-type

(Figure 1B and 1E). Additionally, cytokinesis often resulted in extra furrow formation

(Figure 1B, bottom right), suggesting that the normal cytokinesis process is also

compromised by this mutation. During AB cell division at the 2-cell stage, *act-2(or295)*

embryos exhibited a reduced degree of rotation during cytokinesis (Figure 1C and 1F).

During EMS cell division at the 6-cell stage, *act-2(or295)* displayed an abnormal shift of

the contractile ring; some embryos showed no shift in position, whereas others showed an

anterior shift (Figure 1D and 1G). In summary, *act-2(or295)* mutants display defects in all

three modes of spatially controlled cytokinesis and also exhibit defects in proper contractile

ring formation.

High-throughput RNAi screen in *act-2(or295)* mutant background

We reasoned that any RNAi knockdown that either suppresses or enhances *act-2(or295)*

embryonic lethality might include regulators of the spatial control and mechanical

resilience of cytokinesis. According to single-cell RNA-seq data, 5,928 genes are

expressed in early *C. elegans* embryos (Tintori et al., 2016). Of these, 4,445 genes are

included in the Ahringer feeding RNAi library (Kamath and Ahringer, 2003), and we

performed a feeding RNAi screen targeting them (Figure 2A). Both wild-type and *act-*

2(or295) mutants expressing an intestinal GFP reporter (*vha-6*p::GFP) were fed dsRNA-

expressing bacteria at 20°C, a temperature at which *act-2(or295)* exhibits 80.2%

embryonic lethality (n > 100). Instead of performing standard feeding RNAi on Nematode

Growth Media (NGM) plates, we incubated the L1-stage larvae in liquid NGM

supplemented with dsRNA-expressing bacteria (hereafter "liquid-based RNAi") in 96-well

plates (Figure 2A). We then counted the number of hatched worms in each well by imaging

the intestinal GFP signal using high-content screening microscopy (Figure 2B-C). The

effects of RNAi on embryonic viability in both wild-type and *act-2(or295)* strains were

measured by computing Z-scores using the mean and standard deviation of worm counts

within each 96-well plate, rather than using the global mean and standard deviation of the

entire screen, to mitigate technical variation. For each of the 4,218 analyzed genes, we

obtained three parameters: (1) Z-score in the control background, (2) Z-score in the *act-*

2(or295) background, and (3) the difference between the two (Figure 3A).

We found that liquid-based feeding RNAi was consistently less effective than

standard plate-based feeding RNAi, possibly due to the lower density of bacteria or

differences in worm feeding behavior. Because of this weaker knockdown effect, we

identified the essential cytokinesis regulator *nmy-2* (non-muscle myosin II) as a suppressor

of *act-2(or295)* lethality in our screen (Figure 2C). Furthermore, we identified components

of the RhoA signaling pathway (*ect-2/RhoGEF*, *rga-1/RhoGAP, cyk-1/formin*), structural

components of the contractile ring (*act-1/-2/-3*, *ani-2/anillin*), and a regulator of branched

actin assembly (*wsp-1/N-WASP*) as suppressors (Figure 3A-B).

In this primary screen, we set a less conservative threshold and designated any

RNAi with a mutant Z-score ≥ 1.5 or a Z-score difference ≥ 1.5 as a suppressor of *act-2*

lethality (Figure 3C). Conversely, RNAi knockdown resulting in failed hatching or

significantly reduced hatching in the *act-2(or295)* background was classified as an

enhancer. Hereafter, we focus only on *act-2(or295)* suppressors, as enhancers may include

genes whose knockdown additively increased the lethality of *act-2(or295)* without

indicating a specific genetic interaction. We found that 304 genes displayed Z-scores ≥ 1.5

in *act-2(or295)*, and 547 genes exhibited Z-score differences ≥ 1.5 (Figure 3C; a total of

689 genes). In pursuit of genes directly regulating cytoskeletal dynamics or cell signaling,

we excluded those encoding DNA-binding, RNA-binding, and ribosomal proteins (which

likely affect general viability through transcription or translation). After this filtering, 385

genes had Z-score differences ≥ 1.5, and 184 genes showed Z-scores ≥ 1.5 in *act-2(or295)*,

leading to a total of 454 *act-2* suppressor genes (Figure 3C). From this list, we selected

genes that have a known function, a conserved domain, or are conserved in humans,

yielding a final set of 385 candidate genes used for the secondary screens (Figure 3C).

Identification of regulators of spatial control of cytokinesis

Next, we performed a secondary screen to identify regulators of the spatial control

of cytokinesis among 385 genes characterized as *act-2(or295)* suppressors (Figure 4A).

For each RNAi knockdown, we collected three live-imaging movies capturing

embryogenesis from the 1-cell to the 7-cell stage.

For asymmetric contractile ring closure in zygote, we visually classified an embryo

as defective if the cleavage furrow ingressed asymmetrically. Because asymmetric ring

closure can occur in any orientation, furrowing asymmetry is not detectable in our 2D

imaging setup when the ring closure occurs along the imaging axis. Indeed, about 23% of

wild-type zygotes exhibited symmetric furrowing patterns (n = 22, Figure 1E). Thus, we

aimed to identify extreme cases that displayed a high penetrance of defects. We scored

phenotypes for each collected movie (score = 0 if defective, score = 1 if normal) and

calculated a Z-score for each gene using the mean and standard deviation across all *act-2*

suppressors. We identified 35 genes exhibiting symmetric contractile ring closure in all

three movies (Z-score $\geq$ 1.96; Figure 4B). These candidates include *dhhc-9*/ZDHHC16, which encodes S-palmitoyltransferase, and *cyn-6*/PPIB, which encodes peptidylprolyl isomerase B (Figure 5A).

For the rotational cytokinesis during AB cell division, we measured the angle of AB division axis rotation. We identified 13 genes with Z-scores > 1.96 (indicating reduced rotation) and 3 genes with Z-score $\leq$ -1.96 (indicating excess rotation) (Figure 4C). We found that *sem-5*/Grb2, involved in epidermal growth factor signaling, causes strong defects in AB cell rotation (Figure 5B).

For the asymmetric contractile ring displacement during the EMS cell cytokinesis, we calculated the shift of the cleavage furrow along the anterior-posterior axis before and after cytokinesis for each movie. We identified 11 genes with Z-score $\geq$ 1.96 (indicating less furrow displacement) and 10 genes with Z-score $\leq$ -1.96 (indicating excessive displacement). These candidates include *pfd-1*/PFDN1, which encodes a prefoldin subunit known to be involved in actin folding (Figure 5C).

Finally, we performed a gene network analysis using the 69 identified genes and found that 16 candidate genes—either genetically or physically, and directly or indirectly— interact with known cytokinesis regulators (Figure 5D).

Identification of regulators of mechanical resilience of cytokinesis

In *act-2(or295)* embryos, which exhibit hypercontractility, cells are under constant mechanical stress yet display only subtle defects in contractile ring formation (e.g., extra furrow shown in Figure 1B) and show no defects in cytokinesis completion in zygotes (Figure 6B). Thus, we reasoned that the genes required for cytokinesis in *act-2(or295)* may

include regulators of the mechanical resilience of cytokinesis. We selected 139 genes from the 385 *act-2* suppressors based on the following criteria: for genes on Chromosome I and II, RNAi knockdown caused > 67% defects in asymmetric furrowing in the wild-type background; for genes on the remaining chromosomes, RNAi knockdown caused 100% embryonic lethality in *act-2(or295)* (Figure 6A). We then performed an RNAi screen targeting these 139 genes in an *act-2* mutant background using a standard plate-based feeding RNAi (Figure 6A). Of the 139 candidates examined, RNAi knockdown of 12 genes led to mild to severe cytokinesis defects. Among these, five genes are known cytokinesis regulators of cytokinesis: the actin genes (*act-1, act-2, act-3*), *ect-2/RhoGEF* (which activates RhoA), and *nmy-2* (non-muscle myosin II). Another three genes (*egg-6, cks-1, ubc-12*) are involved in eggshell formation, cell cycle regulation, and ubiquitination, respectively, and are known to cause cytokinesis defects when disrupted in wild-type background. Two additional genes (*tba-1*/α-tubulin and the centrosomal/centriolar component *spd-2*) function in microtubule assembly and organization.

The final two genes are relatively less understood. First, *evl-20/ARL2* encodes ADP-ribosylation factor-like GTPase that localizes to the periphery of astral microtubules and the plasma membrane (Antoshechkin and Han 2002). A previous study reported that *evl-20* knockdown or mutation reduces microtubule levels during embryogenesis and produces multinucleated cells later in development, such as in vulval precursor cells and the somatic gonad (Antoshechkin and Han 2002). However, the early embryonic phenotype was not described, and cytokinesis has not been analyzed in detail. *lpin-1/Lipin 1* encodes phosphatidic acid phosphatase involved in polyunsaturated fatty acids (PUFAs) production and longevity (Jung *et al.* 2020). Previous studies showed that *lpin-1* is required for proper

endoplasmic reticulum organization and timely nuclear envelope breakdown during embryogenesis (Golden *et al.* 2009; Gorjánácz and Mattaj 2009). In *Toxoplasma gondii*, the loss of Lipin results in multinucleation, possibly due to cytokinesis failure (Dass *et al.* 2021).

To identify genes specifically required for cytokinesis in *act-2(or295)*, we compared RNAi phenotypes between wild-type and *act-2* backgrounds and found that *evl-20*, *lpin-1*, *spd-2*, and *tba-1* induced more severe cytokinesis defects in *act-2(or295)* (Figure 6B). The phenotypes of *spd-2(RNAi);act-2(or295)* and *tba-1(RNAi); act-2(or295)* were relatively mild, as these embryos still formed a furrow that sometimes regressed later (Figure 6C, bottom). By contrast, *evl-20(RNAi);act-2(or295)* and *lpin-1(RNAi); act-2(or295)* failed to form a fully developed furrow, suggesting that these genes are involved in contractile ring formation (Figure 6C).

Discussion

In this study, we found that the gain-of-function actin mutant *act-2(or295)* exhibits defects in the spatial control of cytokinesis. By leveraging this genetic background, we conducted a high-throughput RNAi screen and identified 689 suppressors of *act-2(or295)* embryonic lethality. In a secondary, live-imaging-based screen in a wild-type background, we identified 35 genes involved in asymmetric contractile ring closure in zygotes, 16 genes involved in rotational cytokinesis in the 2-cell stage AB cell, and 21 genes involved in asymmetric contractile ring positioning in the 6-cell stage EMS cell. Furthermore, we conducted another secondary RNAi screen in *act-2(or295)* background to search for genes

enabling successful cytokinesis under mechanical stress. We identified four genes that

exhibit cytokinesis defects in *act-2(or295)*.

Although the position and function of the contractile ring are spatially controlled in

tissues, the mechanisms underlying this spatial control remain unclear. Through our RNAi

screen, we identified potential regulators for each of the three modes of spatial cytokinesis

control. Among the 69 candidate genes, only three genes were common to at least two

modes of spatial control. Some of the identified genes are known to be involved in

cytokinesis. For example, we identified *dhhc-9*/ZDHHC16, which encodes S-

palmitoyltransferase, as a regulator of asymmetric contractile ring closure (Figure 5A). A

recent study in mammalian cultured cells shows that ZDHHC5 regulates the palmitoylation

of protocadherin 7 (PCDH7) during mitosis, thereby targeting its cleavage furrow

localization and activating RhoA (Ozkan *et al.* 2023). Disruption of PCDH7 increases

cytokinesis failure, but its role in asymmetric contractile ring closure remains unexplored.

We also identified *sem-5*/Grb2 as a regulator of rotational cytokinesis (Figure 5B). Grb2 is

a protein containing Src homology domain 2 (SH2) and SH3 domains, acting as an adaptor

in various cell signaling pathways, such as FGF and EGF signaling (Giubellino *et al.* 2008).

Grb2 interacts with Wiskott-Aldrich Syndrome protein (WASp) and with cortactin, both

of which regulate the branched actin nucleator Arp2/3 complex (She *et al.* 1997; Martinez-

Quiles *et al.* 2004). Interestingly, our candidates for rotational cytokinesis regulators also

include the N-WASp homolog *wsp-1*. The list of candidate genes generated in this study

should inspire novel hypotheses by integrating this existing molecular knowledge.

The contractile ring is a molecular machine known for its remarkable resilience, yet

the mechanisms underlying this resilience of cytokinesis are not yet fully understood. Our

secondary screen in *act-2(or295)* mutants identified four genes that contribute to

cytokinesis completion in this background. Genes, such as *tba-1*, *spd-2*, and *evl-20* are

involved in microtubule assembly. The mitotic spindle plays a critical role in activating

RhoA through the central spindle and astral microtubules (Pollard and O'Shaughnessy

2019). A previous small-scale RNAi screen in *C. elegans* also identified microtubule

regulators as genes required for the mechanical resilience of cytokinesis (Singh *et al.* 2019).

That experimental setup was more specific, subjecting embryos to uniaxial compression

using small-diameter beads and a coverslip. Thus, our identification of these microtubule

regulators confirms their requirement in the mechanical resilience of cytokinesis. We also

identified the Lipin homolog *lpin-1* as a regulator of mechanical resilience of cytokinesis.

The specific role of Lipin in contractile ring formation and function has yet to be

determined and will be an interesting direction for future research.

It is worth noting that this study has three main limitations. First, the primary RNAi

screen, which analyzed 4,218 early embryonic genes, was performed only once.

Consequently, some pseudonegatives may have gone undetected due to technical errors.

Second, both secondary RNAi screens used feeding RNAi with relatively small sample

sizes, limiting our ability to fully elucidate the roles of candidate genes in cytokinesis.

Finally, *act-2(or295)* enhancers likely include other interesting candidates, but we did not

analyze them in order to streamline the study.

Overall, this study identified a number of potentially novel regulators of cytokinesis,

with a particular focus on the relatively underexplored areas of cytokinesis research,

mechanical resilience and spatial control. The identified list of genes will facilitate the

generation of future hypotheses aimed at understanding the multicellular-specific regulation of cytokinesis during animal development.

Acknowledgements

We thank Caenorhabditis Genetics Center (funded by the NIH Office of Research Infrastructure Programs; P40 OD010440) for sharing worm strains and the Sugioka lab members for general discussions. We also thank Life Sciences Institute Imaging Core Facility, Michele Roberge, Aruna Balgi for sharing high-content screening microscopy, Don Moerman and Mark Edgley for sharing liquid handling machine. This work was supported by the Canadian Institutes of Health Research (Project Grant; PJT-169145), Government of Canada's New Frontiers in Research Fund (NFRFE-2019-00310), and the Michael Smith Health Research BC (Scholar Award; SCH-2020-0406) to K.S.

Author contributions

C.G.: Formal analysis, Investigation, Data curation, Writing-Original Draft, Validation, Visualization. **Y.R.X.**: Formal analysis, Investigation, Data Curation, Project administration. **A. H.**: Formal analysis, Investigation, Methodology, Data Curation, Project administration. **C.R.H.**: Investigation. **V. J.**: Data Curation. **M.J.K.** Investigation. **K.D.** Investigation. **K.S.**: Conceptualization, Formal analysis, Investigation, Methodology, Validation, Writing-Original Draft, Writing-Review & Editing, Supervision, Funding acquisition.

Declaration of interests

We declare no competing interests.

Figure legends

Figure 1. *act-2(or295)* mutant exhibits abnormal spatial control of

cytokinesis.

(**A**) Three modes of spatial control of cytokinesis during animal development. (**B**)

Asymmetric contractile ring closure in zygotes. The contractile ring closes

asymmetrically in wild-type embryos but is defective in *act-2(or295)*. *act-2(or295)*

mutants also exhibit extra furrow phenotypes at later stage of cytokinesis. (**C**) Rotational

cytokinesis in the AB cell. At the two-cell stage, the AB cell normally rotates during

cytokinesis, but this rotation is defective in *act-2(or295)*. Dotted white lines indicate

mitotic spindle position. (**D**) Asymmetric positioning of the contractile ring in the EMS

cell. At the 6-cell stage, the EMS forms a cleavage furrow in the equatorial region, which

then shifts towards the posterior. In *act-2(or295),* this posterior furrow shift is often

defective. (**E**-**G**) Quantification of zygote furrow asymmetry (E), cellular rotation during

AB cell division (F), and posterior furrow displacement during EMS cell division (G).

Yellow arrowheads indicate the furrow position. Scale bars represent 10 μm. Error bars

represent 95% confidence intervals. Fisher's exact test was performed in panel E and

Welch's t-test was used for panels F and G. P-values: "****", "**", "*"and "ns" indicate

$p < 0.0001$, $p < 0.01$, $p < 0.05$, and $p > 0.05$, respectively.

Figure 2. High-throughput RNAi screen using *act-2(or295)* mutants.

(**A**) Design of the RNAi screen. Wild-type and *act-2(or295)* mutants expressing an

intestinal GFP reporter were bleached, and purified embryos were hatched without food

to induce starvation-induced L1 larval arrest. The synchronized L1 larvae were cultured

in liquid NGM containing dsRNA expressing feeding RNAi bacteria. Hatched F1 larvae

were then imaged using high-content screening microscopy. (**B**) Intestinal GFP reporter

signals for the control and one suppressor, *nmy-2(RNAi)*. Segmented images were used to

count the number of hatched worms. (**C**) The number of hatched worms in a 96-well

plate. *nmy-2(RNAi)* resulted in a significantly higher number of hatched worms,

identified it as a suppressor of *act-2(or295)*.

Figure 3. Summary of the primary RNAi screen.

(**A**) Scatter plot of Z-scores for 4,218 analyzed genes. Z-scores in the control strain and

act-2 mutants are shown on the x- and y-axes, respectively. The difference in Z-scores

between control and *act-2* is indicated by the color code. (**B**) Scatter plot of Z-scores

highlighting known regulators of cytokinesis. As indicated by red, many of them were

identified as suppressors. (**C**) Summary of identified actin suppressors. Among the non-

DNA/RNA-binding and non-ribosomal genes, 385 and 184 suppressors were identified

based on the Z-score difference and the Z-score in *act-2* mutants, respectively. Of the 454

suppressor genes, worm-specific genes lacking known functions or conserved domains

were removed, leaving 385 genes for the secondary screens.

Figure 4. Secondary screen for spatial cytokinesis regulators.

(**A**) Design of the secondary screen. A total of 385 5 *act-2* suppressor genes were targeted

by standard plate-based feeding RNAi in a wild-type background. For asymmetric

furrowing in zygotes, we performed a qualitative assessment, whereas for AB cell

rotation and EMS cell furrow shift, we measured the degree of rotation and the extent of

furrow shift, respectively. (**B-D**) Candidate regulators of asymmetric contractile ring

closure (B), rotational cytokinesis (C), and asymmetric furrow shift (D). Orange bars

indicate extreme samples with Z-scores ≥ 1.96 or ≤ -1.96.

Figure 5. Identification of spatial cytokinesis regulators.

(**A**) Defects in asymmetric furrowing in *dhhc-9(RNAi)*. (**B**) Defects in rotational

cytokinesis in *sem-5(RNAi)*. (**C**) Defects in asymmetric furrow displacement in *pfd-*

1(RNAi). (**D**) Gene interaction network among identified candidate genes and known

cytokinesis regulators. Protein-protein interactions (PPI), genetic interactions, shared

domain information are shown. Candidates without known interactions are not displayed.

Scale bar indicates 10 µm.

Figure 6. Identification of genes required for cytokinesis in the *act-2(or295)*

background.

(**A**) Design of the secondary screen. A total of 139 genes were selected based on defects

in asymmetric furrowing in zygotes or 100% embryonic lethality in the *act-2* mutant

background from 385 5 *act-2* suppressor genes. These genes were knocked-down in *act-*

2(or295) mutants and tested for their requirement in cytokinesis completion. (**B**) Genes

identified as required for cytokinesis completion in *act-2(or295)* mutants. P-values were

1 calculated by chi-square test, followed by Holm-Sidak adjustment for multiple

2 comparisons. P-values: "**", "*", and "ns" indicate $p < 0.01$, $p < 0.05$, and $p > 0.05$,

3 respectively. (**C**) Cytokinesis phenotype after candidate gene knockdown. White dotted

4 lines indicate the mitotic spindle position, and yellow arrowheads indicate the cleavage

5 furrow. Scale bars, 10 µm.

7
Materials and Methods

Maintenance of worm strains

All *C. elegans* strains were cultured and maintained using standard methods

(Brenner 1974). The transgene, *oxTi872*[*vha-6p*::GFP::*tbb-2* 3'UTR + Cbr-*unc-119*(+)],

was used to monitor intestinal GFP expression. Additionally, *act-2(or295)* mutants were

used as our sensitized genetic background. Both the *act-2(or295)* mutants and wild-type

strains carrying *oxTi872* were maintained on NGM agar plates at 15˚C. During the liquid

culture RNAi screen, these strains were incubated at 20˚C, the semi-permissive

temperature for *act-2(or295)* mutant strain. For both secondary screens, these strains were

incubated at 25˚C, the restrictive temperature for *act-2(or295)*.

RNAi

20   Feeding RNAi was performed using the Ahringer RNAi library (Fraser *et al.* 2000;

21 Ahringer 2006). The expression of dsRNA in the *Escherichia coli* strain HT115 was

22 induced by overnight incubation at 37˚C in Nematode Growth Medium (NGM) containing

23 10 µg/mL Ampicillin and 1 mM isopropyl ß-D-1-thiogalactopyranoside (IPTG).

For the high-throughput RNAi screen, gravid adults were bleached with hypochlorite solution, and the resulting purified embryos were incubated in M9 buffer overnight to obtain developmentally arrested L1 stage worms. These worms were suspended in liquid NGM and approximately 4-5 L1 worms were transferred into each well of 96-well plates containing various dsRNA-expressing bacteria. After 5 days of incubation at 20˚C, the 96-well plates were imaged using a high-content screening system (Cellomics Arrayscan; Thermo Fisher).

For the two secondary RNAi screens, dsRNA-expressing bacteria were cultured on solid NGM agar plates. Developmentally arrested L1 worms were transferred onto these plates and incubated for 3 days at 25˚C. For genes required for fertility, we used L4 stage worms were used and incubated overnight before live imaging. Wild-type worms and *act-2(or295)* mutants were used for the spatial cytokinesis regulator screen and the cytokinesis resilience regulator screen, respectively. A bacterial strain containing the empty vector L4440 served as a negative control in both secondary screens.

Live-imaging

To obtain embryos for live imaging by brightfield microscopy, we dissected gravid adults on a coverslip in a 5 μl droplet of M9 buffer. We then placed the coverslip gently onto a 2% agarose pad mounted on a glass slide and sealed the edges with petroleum jelly (Vaseline) to prevent dehydration.

For characterization of the strains used in the high-throughput RNAi screen and secondary screens, we performed live imaging at room temperature (~22.5ºC) using brightfield compound microscopes (B290-TB; OPTIKA), equipped with a 100X oil

objective lens (1.2NA) and controlled by OPTIKA Vision Lite software. For the secondary

screen of spatial cytokinesis regulators, images were taken every 2 seconds for 50 minutes,

beginning at the 1-cell stage and continuing until after EMS cell division. For the secondary

screen of cytokinesis resilience regulators, images were taken every 2 seconds for 10

minutes, beginning at the 1-cell stage and ending at the 2-cell stage. In most cases, filming

began during late pronuclear migration stage but before nuclear envelope breakdown

(NEBD).

Image analysis for the high-throughput RNAi screen

To quantify the number of hatched worms, we processed images obtained from the

high-content screening system using Fiji (Schindelin *et al.* 2012). We applied a Laplacian

filter to extract edge information while removing background signals. Next, we used Otsu's

threshold to identify relevant signals and then applied morphological filters to select rod-

like worm bodies while excluding circular objects, such as embryos and debris. The

resulting segmentation data was used to count the number of hatched F1 and F2 worms.

We then standardized the worm counts for each 96-well plate by calculating Z-scores,

using the mean and standard deviation of worm counts within each plate to minimize batch

effects. We computed three different Z-scores for each candidate gene: one in the control

background, one in the *act-2(or295)* mutant background, and the difference between the

two. Genes with Z-scores greater than 1.5 in the *act-2(or295)* mutant background were

considered actin suppressors. Additionally, genes with a Z-score difference greater than

1.5 between *act-2* and control strains were also designated as actin suppressors.

Selection of candidate genes for secondary screens

For the secondary screen of spatial regulators of cytokinesis, all actin suppressor genes were assessed using plate-based RNAi in a wild-type strain (EG8829). At least three embryos were imaged for each gene, and phenotypes were manually annotated in Fiji using the collected live-imaging movies.

For the secondary screen of regulators of cytokinesis resilience, we first selected actin suppressor genes using two different criteria. Initially, we chose genes based on defects in asymmetric ring closure in 1-cell stage zygotes, as identified in the spatial cytokinesis regulator screen. However, we found that most of these genes were not required for cytokinesis in the *act-2* mutant background. Therefore, we selected genes whose RNAi knockdown cased 100% embryonic lethality in *act-2* mutants at 25˚C for genes on chromosomes III-X.

Bioinformatics

Scatter plots for the high-throughput screening data were generated using Matplotlib. To analyze and visualize gene networks among the identified candidates, we used GeneMANIA in the Cytoscape software (Shannon *et al.* 2003).

Statistics

Statistical analyses were performed using Prism9 (GraphPad Software) and custom Python scripts. For Figure 1E, we performed Fisher's exact test on raw data (rather than percentages). For Figure 1F-G, we used Welch's t-test. For Figure 6B, we used Fisher's exact test on raw data. The resulting P-values were corrected for multiple comparisons

using a custom Python script (Holm-Sidak method). The symbols such as ****, ***, **,

*, and ns indicate $p < 0.0001$, $p < 0.001$, $p < 0.01$, $p < 0.05$, and $p > 0.05$, respectively.

References:

Ahringer J., 2006 Reverse genetics. WormBook. https://doi.org/10.1895/wormbook.1.47.1

Antoshechkin I., and M. Han, 2002 The C. elegans evl-20 gene is a homolog of the small
GTPase ARL2 and regulates cytoskeleton dynamics during cytokinesis and
morphogenesis. Dev. Cell 2: 579–591.

Basant A., and M. Glotzer, 2018 Spatiotemporal Regulation of RhoA during Cytokinesis.
Curr. Biol. 28: R570–R580.

Brenner S., 1974 The genetics of Caenorhabditis elegans. Genetics 77: 71–94.

Dass S., S. Shunmugam, L. Berry, C.-S. Arnold, N. J. Katris, *et al.*, 2021 Toxoplasma
LIPIN is essential in channeling host lipid fluxes through membrane biogenesis and
lipid storage. Nat. Commun. 12: 2813.

Fraser A. G., R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann, *et al.*, 2000
Functional genomic analysis of C. elegans chromosome I by systematic RNA
interference. Nature 408: 325–330.

Frémont S., and A. Echard, 2018 Membrane traffic in the late steps of cytokinesis. Curr. Biol. 28: R458–R470.

Giubellino A., T. R. Burke Jr, and D. P. Bottaro, 2008 Grb2 signaling in cell motility and cancer. Expert Opin. Ther. Targets 12: 1021–1033.

Golden A., J. Liu, and O. Cohen-Fix, 2009 Inactivation of the C. elegans lipin homolog leads to ER disorganization and to defects in the breakdown and reassembly of the nuclear envelope. J. Cell Sci. 122: 1970–1978.

Goldstein B., 1992 Induction of gut in Caenorhabditis elegans embryos. Nature 357: 255–257.

Gorjánácz M., and I. W. Mattaj, 2009 Lipin is required for efficient breakdown of the nuclear envelope in Caenorhabditis elegans. J. Cell Sci. 122: 1963–1969.

Herszterg S., A. Leibfried, F. Bosveld, C. Martin, and Y. Bellaiche, 2013 Interplay between the Dividing Cell and Its Neighbors Regulates Adherens Junction Formation during Cytokinesis in Epithelial Tissue. Dev. Cell 24: 256–270.

Jankele R., R. Jelier, and P. Gönczy, 2021 Physically asymmetric division of the C. elegans zygote ensures invariably successful embryogenesis. Elife 10. https://doi.org/10.7554/eLife.61714

Jung Y., S. Kwon, S. Ham, D. Lee, H.-E. H. Park, *et al.*, 2020 Caenorhabditis elegans Lipin 1 moderates the lifespan-shortening effects of dietary glucose by maintaining ω-6 polyunsaturated fatty acids. Aging Cell 19: e13150.

Kuroda R., B. Endo, M. Abe, and M. Shimizu, 2009 Chiral blastomere arrangement

dictates zygotic left-right asymmetry pathway in snails. Nature 462: 790–794.

Maddox A. S., L. Lewellyn, A. Desai, and K. Oegema, 2007 Anillin and the septins

promote asymmetric ingression of the cytokinetic furrow. Dev. Cell 12: 827–835.

Martinez-Quiles N., H.-Y. H. Ho, M. W. Kirschner, N. Ramesh, and R. S. Geha, 2004

Erk/Src phosphorylation of cortactin acts as a switch on-switch off mechanism that

controls its ability to activate N-WASP. Mol. Cell. Biol. 24: 5269–5280.

Meshcheryakov V. N., and L. V. Beloussov, 1975 Asymmetrical rotations of blastomeres

in early cleavage of gastropoda. Wilehm Roux Arch Dev Biol 177: 193–203.

Mierzwa B., and D. W. Gerlich, 2014 Cytokinetic abscission: molecular mechanisms and

temporal control. Dev. Cell 31: 525–538.

Naganathan S. R., S. Furthauer, M. Nishikawa, F. Julicher, and S. W. Grill, 2014 Active

torque generation by the actomyosin cell cortex drives left-right symmetry breaking.

Elife 3: e04165.

Ozkan N. E., B. N. Yigit, B. S. Degirmenci, M. H. Qureshi, G. N. Yapici, *et al.*, 2023 Cell

cycle-dependent palmitoylation of protocadherin 7 by ZDHHC5 promotes

successful cytokinesis. J. Cell Sci. 136: jcs260266.

Piekny A. J., and A. S. Maddox, 2010 The myriad roles of Anillin during cytokinesis.

Semin. Cell Dev. Biol. 21: 881–891.

Pollard T. D., 2017 Nine unanswered questions about cytokinesis. J. Cell Biol. 216: 3007–

3016.

Pollard T. D., and B. O'Shaughnessy, 2019 Molecular Mechanism of Cytokinesis. Annu.

Rev. Biochem. 88: 661–689.

Priess J. R., and J. N. Thomson, 1987 Cellular interactions in early C. elegans embryos.

Cell 48: 241–250.

Schierenberg E., and B. Junkersdorf, 1992 The role of eggshell and underlying vitelline

membrane for normal pattern formation in the early C. elegans embryo. Rouxs.

Arch. Dev. Biol. 202: 10–16.

Schindelin J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, *et al.*, 2012 Fiji: an

open-source platform for biological-image analysis. Nat. Methods 9: 676–682.

Shannon P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, *et al.*, 2003 Cytoscape: a

software environment for integrated models of biomolecular interaction networks.

Genome Res. 13: 2498–2504.

She H. Y., S. Rockow, J. Tang, R. Nishimura, E. Y. Skolnik, *et al.*, 1997 Wiskott-Aldrich

syndrome protein is associated with the adapter protein Grb2 and the epidermal

growth factor receptor in living cells. Mol. Biol. Cell 8: 1709–1721.

Silva A. M., D. S. Osório, A. J. Pereira, H. Maiato, I. M. Pinto, *et al.*, 2016 Robust gap

repair in the contractile ring ensures timely completion of cytokinesis. J. Cell Biol.

215: 789–799.

Singh D., D. Odedra, P. Dutta, and C. Pohl, 2019 Mechanical stress induces a scalable
switch in cortical flow polarization during cytokinesis. J. Cell Sci. 132: jcs231357.

Sugioka K., 2021 Symmetry-breaking of animal cytokinesis. Semin. Cell Dev. Biol.
https://doi.org/10.1016/j.semcdb.2021.12.008

Taulet N., B. Vitre, C. Anguille, A. Douanier, M. Rocancourt, *et al.*, 2017 IFT proteins
spatially control the geometry of cleavage furrow ingression and lumen positioning.
Nat. Commun. 8: 1928.

Willis J. H., E. Munro, R. Lyczak, and B. Bowerman, 2006 Conditional dominant
mutations in the Caenorhabditis elegans gene act-2 identify cytoplasmic and muscle
roles for a redundant actin isoform. Mol. Biol. Cell 17: 1051–1064.

Wood W. B., 1991 Evidence from reversal of handedness in C. elegans embryos for early
cell interactions determining cell fates. Nature 349: 536–538.

Figure 1. Gough et al.

Figure 1. Actin gain-of-function mutant exhibits abnormal spatial control of cytokinesis.
(**A**) Three modes of spatial control of cytokinesis during animal development. (**B**) Asymmetric contractile ring closure in zygotes. The contractile ring closes asymmetrically in wild-type embryos but is defective in *act-2(or295)*. *act-2(or295)* mutants also exhibit extra furrow phenotypes at later stage of cytokinesis. (**C**) Rotational cytokinesis in the AB cell. At the two-cell stage, the AB cell normally rotates during cytokinesis, but this rotation is defective in *act-2(or295)*. Dotted white lines indicate mitotic spindle position. (**D**) Asymmetric positioning of the contractile ring in the EMS cell. At the 6-cell stage, the EMS forms a cleavage furrow in the equatorial region, which then shifts towards the posterior. In *act-2(or295)*, this posterior furrow shift is often defective. (**E**-**G**) Quantification of zygote furrow asymmetry (E), cellular rotation during AB cell division (F), and posterior furrow displacement during EMS cell division (G). Yellow arrowheads indicate the furrow position. Scale bars represent 10 μm. Error bars represent 95% confidence intervals. Fisher's exact test was performed in panel E and Welch's t-test was used for panels F and G. P-values: "****", "**", "*"and "ns" indicate $p < 0.0001$, $p < 0.01$, $p < 0.05$, and $p > 0.05$, respectively.

Figure 2. High-throughput RNAi screen using *act-2(or295)* mutants.

(**A**) Design of the RNAi screen. Wild-type and *act-2(or295)* mutants expressing an intestinal GFP reporter were bleached, and purified embryos were hatched without food to induce starvation-induced L1 larval arrest. The synchronized L1 larvae were cultured in liquid NGM containing dsRNA expressing feeding RNAi bacteria. Hatched F1 larvae were then imaged using high-content screening microscopy. (**B**) Intestinal GFP reporter signals for the control and one suppressor, *nmy-2(RNAi)*. Segmented images were used to count the number of hatched worms. (**C**) The number of hatched worms in a 96-well plate. *nmy-2(RNAi)* resulted in a significantly higher number of hatched worms, identified it as a suppressor of *act-2(or295)*.

Figure 3. Summary of the primary RNAi screening.
(**A**) Scatter plot of Z-scores for 4,218 analyzed genes. Z-scores in the control strain and *act-2* mutants are shown on the x- and y-axes, respectively. The difference in Z-scores between control and *act-2* is indicated by the color code. (**B**) Scatter plot of Z-scores highlighting known regulators of cytokinesis. As indicated by red, many of them were identified as suppressors. (**C**) Summary of identified actin suppressors. Among the non-DNA/RNA-binding and non-ribosomal genes, 385 and 184 suppressors were identified based on the Z-score difference and the Z-score in act-2 mutants, respectively. Of the 454 suppressor genes, worm-specific genes lacking known functions or conserved domains were removed, leaving 385 genes for the secondary screens.

Figure 4. Secondary screen for spatial cytokinesis regulators.

(**A**) Design of the secondary screen. A total of 385 act-2 suppressor genes were targeted by standard plate-based feeding RNAi in a wild-type background. For asymmetric furrowing in zygotes, we performed a qualitative assessment, whereas for AB cell rotation and EMS cell furrow shift, we measured the degree of rotation and the extent of furrow shift, respectively. (**B**-**D**) Candidates regulators of asymmetric contractile ring closure (B), rotational cytokinesis (C), and asymmetric furrow shift (D). Orange bars indicate extreme samples with Z-scores ≥ 1.96 or ≤ -1.96.

Figure 5. Gough et al.

Figure 5. Identification of spatial cytokinesis regulators.

(**A**) Defects in asymmetric furrowing in *dhhc-9(RNAi)*. (**B**) Defects in rotational cytokinesis in *sem-5(RNAi)*. (**C**) Defects in asymmetric furrow displacement in *pfd-1(RNAi)*. (**D**) Gene interaction network among identified candidate genes and known cytokinesis regulators. Protein-protein interactions (PPI), genetic interactions, shared domain information are shown. Candidates without known interactions are not displayed. Scale bar indicates 10 μm.

Figure 6. Identification of genes required for cytokinesis in *act-2(or295)* background

(**A**) Design of the secondary screen. A total of 139 genes were selected based on defects in asymmetric furrowing in zygotes or 100% embryonic lethality in the act-2 mutant background from 385 act-2 suppressor genes. These genes were knocked-down in act-2(or295) mutants and tested for their requirement in cytokinesis completion. (**B**) Genes identified as required for cytokinesis completion in act-2(or295) mutants. P-values were calculated by chi-square test, followed by Holm-Sidak adjustment for multiple comparisons. P-values: "**", "*", and "ns" indicate p < 0.01, p < 0.05, and p > 0.05, respectively. (**C**) Cytokinesis phenotype after candidate gene knockdown. White dotted lines indicate the mitotic spindle position, and yellow arrowheads indicate the cleavage furrow. Scale bars, 10 μm.