Signaling of trans-saccadic prediction error by foveal neurons of the monkey superior colliculus

Tong Zhang[1, 2]*, Amarender R. Bogadhi[1, 2, 3]*, and Ziad M. Hafed[1, 2]

1 Werner Reichardt Centre for Integrative Neuroscience, University of Tübingen, Tübingen, Germany

2 Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany

3 Neuroscience and Mental Health Discovery Research, Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach, Germany

* Contributed equally

Short title:

Trans-saccadic predictive coding by foveal superior colliculus neurons

Correspondence:

Ziad M. Hafed

Otfried-Müller Str. 25

Tübingen

72076

Germany

Tel: +49 7071 29 88819

E-mail: ziad.m.hafed@cin.uni-tuebingen.de

Abstract

Across saccades, neurons in retinotopically organized visual representations experience drastically different images, but visual percepts remain stable. Here we investigated whether such stability can be mediated, in part, via prediction-error signaling by neurons processing post-saccadic visual images. We specifically recorded from foveal superior colliculus (SC) neurons when a visual image only overlapped with their response fields (RF's) after foveating saccades but not pre-saccadically. When we rapidly changed the target features intra-saccadically, the foveal neurons' post-saccadic visual reafferent responses were elevated, even though the neurons did not directly sample the pre-saccadic extrafoveal target features. This effect did not occur in the absence of saccades, and it also scaled with the extent of the introduced intra-saccadic image feature discrepancies. These results suggest that foveal SC neurons may signal a trans-saccadic prediction error when the foveated image stimulating them is inconsistent with that expected from pre-saccadic extrafoveal representations, a potential perceptual stability mechanism.

Introduction

Sensory systems benefit from spatial and temporal continuity of incoming information streams. Often, such continuity is disrupted, whether through exogenous environmental factors or through self-movement. One example such disruption occurs every time we generate a rapid eye movement. Across saccades, which we typically use to foveate relevant extrafoveal image patches, retinotopic representations of foveal visual inputs can experience drastically different images. For example, when viewing a painting of a natural scene, we might generate a saccade from one foveated image patch, say containing a flower, to another one, say containing a leaf. In this case, the pre-saccadic extrafoveal image of the leaf is rapidly translated to become the post-saccadic foveal image; thus, the pre- and post-saccadic foveal visual images are decidedly different from each other. Despite that, our percept of the portrait remains seamless.

The question of perceptual stability across saccades has intrigued scientists for many decades (1, 2, 3, 4, 5, 6, 7, 8). Among the theories invoked for such stability are ones utilizing a form of prediction (meant in the phenomenological sense) of the visual consequences of eye movements (3, 9). Conceptually, such prediction may be thought of as a template matching between newly incoming foveal visual information and the expected appearance, obtained a priori, of the post-saccadic foveal image (3, 9). Such matching could arise, for example, by relaying a pre-saccadic extrafoveal preview of the eye movement's target (10) to neurons processing the newly foveated visual evidence. Here, we searched for a neural correlate of such a process in the visual responses of foveal superior colliculus (SC) neurons. We specifically asked whether these neurons' post-saccadic reafferent responses (after target foveation) are sensitive to the pre-saccadic extrafoveal image appearance, even when such extrafoveal visual input never activated the neurons' classic response fields (RF's) pre-saccadically. If so, then this would suggest a potential trans-saccadic dispatching of extrafoveal pre-saccadic visual information to the foveal SC.

The SC is particularly attractive for exploring hallmarks of extrafoveal-to-foveal trans-saccadic integration. First, SC neurons exhibit sufficiently varied feature tuning properties (11), allowing us to investigate reafferent visual responses for different image manipulations. Importantly, such feature tuning endows extrafoveal SC neurons with integrated evidence about the extrafoveal appearance of saccade targets (11); while likely not enough to process very fine details (11, 12, 13), this integrated evidence can provide a sufficient (13, 14, 15) pre-saccadic preview. Second, the SC possesses a putative mechanism for relaying exactly such an extrafoveal preview to the rest of the visual system (including foveal representations) at the time of saccades (16, 17, 18, 19). In particular, SC motor bursts, time-locked to saccade onset (20), embed within them a representation of the visual properties of the extrafoveal saccade target (16, 17, 19). Thus, whether through intra- or extra-collicular connections, SC foveal representations (21) can be "informed" about extrafoveal saccade target appearance, despite not being directly visually-stimulated by the saccade target pre-saccadically. Finally, the SC signals salience (22, 23, 24, 25); hence, trans-saccadic integration by foveal SC neurons would be functionally useful to signal failures of image stability. In other words, potential inconsistencies between experienced versus expected foveal images constitute salient events worthy of flagging, so detecting them is desirable.

Here, we show that foveal SC neurons are indeed sensitive to the pre-saccadic extrafoveal
target appearance, despite never being stimulated by the extrafoveal targets before the eye
movements. We also show that such sensitivity is contingent upon saccade generation,
depends on extrafoveal target visibility, and scales with the extent of sensory conflict
experienced between pre-saccadic extrafoveal and post-saccadic foveal image appearances.
We suggest that the SC is endowed with mechanisms for signaling salient events in the
temporal domain (26, 27, 28), such as happens trans-saccadically, and not just in the image-
based one (22, 23, 24, 25).
108
109
110

Results

111
112
113

Foveal superior colliculus neurons detect unexpected trans-saccadic stimulus
changes

116
We recorded from foveal SC neurons while monkeys generated visually-guided saccades to
extrafoveal images containing different spatial frequency textures (12, 13, 16, 29) (Methods).
In our basic condition, the saccade target was a circular patch containing one of two possible
spatial frequencies, and it appeared either to the right or left of fixation (Methods). A
saccade "go" signal was provided a few hundred milliseconds later, after which the monkeys
foveated the circular patch (Fig. 1a). The patch itself never moved on the display, but it was
rapidly translated on the retina by saccades, eventually entering the RF's of our recorded
foveal SC neurons and visually stimulating them.
125
126

Figure 1 Testing for trans-saccadic prediction error signaling by foveal SC neurons. (a) Monkeys generated a
delayed, visually-guided saccade towards an extrafoveal target. We used a delayed paradigm to make sure that
there was a stable visual image upon saccade generation. In some trials, the saccade target was unchanged
throughout the whole trial (high spatial frequency grating embedded within a circular patch for this shown
example). In other trials, we detected saccade onset and immediately flipped the saccade target to another

feature (from a low to a high spatial frequency texture in the shown example; Methods). **(b)** We only selected foveal SC neurons with response fields (RF's) not extending towards the pre-saccadic extrafoveal stimulus location. In this example, the RF was almost entirely contained within <2 deg eccentricity. Each black dot is a stimulus onset location during RF mapping, and the white circle (3 deg radius) shows the extent of the saccade target if it was perfectly foveated post-saccadically. The target covered the RF post-saccadically but not pre-saccadically. The z-axis indicates the visual response strength of the neuron at each stimulus location (Methods). **(c)** Activity of the same example neuron when the extrafoveal stimulus appeared. There was no visual response, confirming that the pre-saccadic stimulus did not drive the neuron. The faint lines show eye position traces (the monkey was still fixating the central fixation spot). **(d)** After saccades, the neuron exhibited a visual reafferent response to the image feature entering its RF. Note how the neuron completely paused intra-saccadically and only responded after the new visual evidence entered its RF (there was additionally a pre-saccadic transient elevation before the pause due to the release of fixation, which we do not focus on in this study). Note also that the neuron was visually-tuned to the low spatial frequency (12, 29). The faint lines indicate eye position traces, showing how the neural modulations were time-locked to saccade onset. Error bars denote SEM, and numbers of trials are evident from the rasters.

We employed a strict selection of neurons to include in our analyses (Methods). Specifically, we ensured that the neurons were not visually-stimulated by the saccade target except after eye movement generation. Thus, we only accepted neurons with foveal RF's that were not visually-driven (before the saccades) by the extrafoveal image patch (RF's were estimated through a mapping task run before our main paradigm; Methods). Fig. 1b shows an example such RF, with the white circle indicating the size of the saccade-target image patch (were the eye to be perfectly aligned with the patch's center). In the actual experiment, this patch was centered at an eccentricity of 8 deg before the saccade, and was thus far from the neuron's RF; however, the same patch was large enough to clearly visually drive the neuron after the eye movement has ended (i.e. after the patch was retinally translated towards the fovea by the saccade). We confirmed these two points by measuring the neuron's activity after extrafoveal stimulus onset and also upon saccade generation. In the former case, there were no visually-evoked bursts (Fig. 1c shows the neuron's activity when extrafoveal image patches containing two different spatial frequency textures appeared; Methods). If anything, there was a transient reduction in baseline activity, particularly for the high spatial frequency texture. Such transient reduction has been observed before upon extrafoveal stimulus onset (for examples, see ref. (30)), and it can aid in subsequent winner-take-all processes needed to recruit the extrafoveal saccade-related motor bursts (31). Upon saccade generation, the neuron now exhibited a visual reafferent response caused by translating the saccade-target image into its RF (Fig. 1d). Importantly, by employing two different spatial frequencies, we ensured that this response was visual in nature, because it depended on the visual appearance of the foveated saccade target (29). For example, the reafferent response of the neuron of Fig. 1b, c was earlier and stronger for the low spatial frequency texture, as might be expected (29) (Fig. 1d).

Having established that our neurons were not visually stimulated by the pre-saccadic extrafoveal stimuli (S1 Figure), we next analyzed the situation in which there was an image mismatch between the experienced post-saccadic visual stimulus and the pre-saccadic (extrafoveal) target image. For each saccade to a given image feature (low or high spatial frequency), we had two conditions: no change in the image feature across the saccade (control), or an instantaneous intra-saccadic change (to the other spatial frequency; Methods). For example, if the pre-saccadic image had a high spatial frequency texture, then, during the saccade, we flipped it to the low spatial frequency texture. Thus, the foveal

neurons experienced a visual feature different from that expected from the extrafoveal pre-
saccadic feature, but these neurons were never directly stimulated by the extrafoveal
feature pre-saccadically. We used instantaneous eye position to detect saccade onset and
updated the display accordingly (either fictively in the control condition or actually;
Methods). This allowed us to compare the visual reafferent response for the same image
feature in the fovea (e.g. landing on a low spatial frequency), but either when the
extrafoveal pre-saccadic image was the same (low spatial frequency) or different (high
spatial frequency).
191
Figure 2a shows example radial eye speed traces for rightward saccades from the same
session of the example neuron of Fig. 1. Here, we aligned all traces to the end of the
stimulus update on trials in which the saccade target flipped from containing a high to a low
spatial frequency texture during the saccade (similar results were obtained for the other
image flip condition). As can be seen, the stimulus update was intra-saccadic, and the eye
speed by the end of the stimulus change was systematically higher than ~100 deg/s. Thus,
given that the gratings were vertically oriented and the saccades were predominantly
horizontal, this ensured substantial blurring of both spatial frequencies on the retina by the
ongoing eye movements (and during the image change). In Fig. 2b, we plotted the visual
reafferent response of the same example neuron of Fig. 1, but now aligned to the time of
intra-saccadic stimulus update completion in all trials (note that control trials had fictive
updates, allowing for similar temporal alignment in the two conditions). In both shown
conditions, we were measuring the neuron's visual reafferent response for the same post-
saccadic foveal stimulus (low spatial frequency); the only difference is that, in one case
(blue), the extrafoveal pre-saccadic image was the same (control; low-to-low spatial
frequency from pre-saccadic to post-saccadic image appearances); in the other (yellow), it
was different (high-to-low spatial frequency change). The neuron's visual reafferent response
was elevated when there was a mismatch in the pre-saccadic image, even though the
foveated visual feature was the same as in the control condition. This was the case even
though the extrafoveal image location was outside of the neuron's RF before the saccade
(Fig. 1).
213
214

Figure 2 Signaling of trans-saccadic prediction error by the foveal SC. (a) Radial eye speed traces from the same session as in Fig. 1b-d when the monkey was generating foveating saccades. Here, we aligned the traces to the end of the stimulus update event (on trials in which the pre-saccadic extrafoveal texture was of high spatial frequency and the post-saccadic foveal feature was a low spatial frequency texture). Eye speed was consistently >100 deg/s during stimulus updates (also see S2a, b Figure demonstrating that we always achieved stimulus updates intra-saccadically and with large eye speeds, maximizing stimulus blurring). **(b)** Relative to control (blue; same data as in Fig. 1), the initial reafferent response of the neuron was elevated (yellow). This happened even though the neuron was not stimulated by the pre-saccadic extrafoveal stimulus (Fig. 1). **(c)** Similar observations when the foveated image feature contained a high spatial frequency; in this case, the control condition also contained a high spatial frequency pre-saccadically. Thus, the neuron's visual reafferent response was systematically elevated when the experienced post-saccadic visual feature was different from that expected based on the pre-saccadic extrafoveal target appearance; this was despite the lack of visual stimulation of the neuron pre-saccadically by the extrafoveal saccade target (Fig. 1). Note also how the reafferent response timing was similar on control and stimulus-change trials, confirming that our temporal alignment (to the stimulus update time) was similar across the different conditions; thus, the different reafferent response magnitudes across conditions were not an artifact of different temporal jitters across them. Error bars denote SEM, and numbers of trials in each condition are evident from the individual traces and rasters.

The same observation was also made when the post-saccadic foveal visual image feature was the high spatial frequency (Fig. 2c). Again, the neuron's reafferent response was elevated (relative to control) when there was a mismatch with the pre-saccadic extrafoveal saccade-target appearance (i.e. when the pre-saccadic image was a low spatial frequency). Note that there was no other change in the firing patterns of the neuron in Fig. 2b, c, for example in the reafferent response timing on stimulus change trials. This suggests that we were not observing a simple superposition of the original reafferent response and an additional sensory drive by the stimulus flip event (or other transients associated with saccade execution); rather, there was a modulation of the otherwise-normal reafferent response, as if it was more salient than expected. Thus, this example foveal SC neuron signaled a trans-saccadic change in the saccade-target appearance, even though it did not "see" the saccade target pre-saccadically.

249 Across neurons and sessions, we confirmed that all of our stimulus updates were intra-
250 saccadic and with high eye speeds maximizing the likelihood of large retinal image blur (S2a,
251 b Figure). We also confirmed that saccade speeds (including post-saccadic drift (32)) (S2c, d
252 Figure) and radial amplitudes (S2e, f Figure) were not altered by the intra-saccadic stimulus
253 changes. And, we confirmed that the neurons' RF hotspot locations were foveal (S3 Figure).
254
At the level of reafferent responses, there was a systematic elevation of their magnitudes,
for each spatial frequency, if the pre-saccadic image contained a different feature from that
experienced foveally after the saccade. This result was assessed in multiple ways. First, we
calculated the normalized population firing rate in each condition, and visualized the
different conditions in Fig. 3a, b (Methods). As can be seen, the population dynamics
behaved very similarly to those of the example neuron of Fig. 2b, c. Second, we performed
receiver operating characteristic (ROC) analysis, like we recently performed (16). Elevations
in the area under the ROC curve indicated elevated firing rates in intra-saccadic stimulus
change trials. As can be seen, whether the post-saccadic image was a low (Fig. 3c) or high
(Fig. 3d) spatial frequency, there was always an elevation of reafferent visual response
amplitudes on stimulus change trials relative to pre-saccadically (error bars denote 95%
confidence intervals). We then plotted individual neuron reafferent responses as scatter
plots (Fig. 3e, f). For each neuron, we measured the strength of the visual reafferent
response (peak firing rate in a prescribed interval after the end of the real or fictive stimulus
update; Methods). We did this twice: once when the pre-saccadic extrafoveal visual image
was the same as that stimulating the foveal RF after the saccade (control), and once when
the pre-saccadic extrafoveal image was of a different spatial frequency (intra-saccadic image
change). We then plotted the two values against each other. Figure 3e shows the condition
when the foveated stimulus had a low spatial frequency texture; the reafferent response to
this texture was elevated if it was inconsistent with the extrafoveal pre-saccadic texture's
spatial frequency ($p=4.91 \times 10^{-8}$; Wilcoxon signed-rank test). Figure 3f shows the results from
the same neurons when the foveated stimulus was a high spatial frequency and the
extrafoveal stimulus in the image flip condition was a low spatial frequency ($p=0.0031$;
Wilcoxon signed-rank test). In both cases, the reafferent response expected from a particular
visual feature in the RF (low or high spatial frequency) was systematically elevated when the
visual feature was unexpected based on the extrafoveal pre-saccadic feature. This is a
hallmark of prediction-error signaling.
282
283

284
Figure 3 Consistency of trans-saccadic error signaling by foveal SC neurons. (a, b) Similar to Fig. 2b, c but now
showing the normalized firing rate of the population. Each neuron's results were normalized before averaging
across neurons (Methods) and error bars denote SEM across neurons. The example neuron results of Fig. 2b, c
were representative of the whole population. **(c, d)** ROC analyses (Methods; also see ref. (16)) comparing intra-
saccadic change trials to control trials across the population. Elevations in the area under the ROC value indicate
higher firing rates on intra-saccadic change trials. For both low and high spatial frequency foveated targets, there
was an elevation in area under the ROC curve post-saccadically when compared to pre-saccadically (and also
surpassing the 0.5 chance level). Note that the values below 0.5 at negative times in **d** reflect an earlier transient
associated with the release of fixation, which is not the focus here. Error bars denote 95% confidence intervals.

 (e) Reafferent response strength (Methods) on trials with an intra-saccadic spatial frequency change versus
reafferent response strength on control trials. In both cases, the foveated image feature was the low spatial
frequency texture within the circular patch of the saccade target. There was a systematic elevation of reafferent
response strengths across the population when there was a visual feature inconsistency across saccades. **(f)**
Similar results for the case in which the foveated texture had a high spatial frequency. **(g, h)** Neural modulation
indices (Methods) for the results in **e**, **f**. For both foveated image features, there was a systematic elevation of
foveal neuron visual reafferent response strength when the experienced post-saccadic image was inconsistent
with that expected from the pre-saccadic target appearance. In each panel, the solid vertical line indicates the
mean of the distribution, and the dashed vertical line indicates the median. Dark histograms indicate the neurons
that were individually significant within a session (Methods).

Finally, we also created a neural modulation index for each foveated image spatial frequency
(Fig. 3g, h; the dark histograms indicate the data for the neurons that were individually
significant within a session; Methods). This index was calculated by subtracting the
reafferent response strength on the control condition from that in the image flip condition
and then dividing by the sum of response strengths (Methods). Again, there was a
systematic elevation of reafferent response strengths when the experienced post-saccadic
foveal visual image feature was not consistent with that expected from the pre-saccadic
extrafoveal visual appearance of the saccade target. Statistically, comparing population
modulation indices to zero revealed significant elevations for both the low ($p=1.47 \times 10^{-7}$;
Wilcoxon signed-rank test) and high ($p=0.0119$; Wilcoxon signed-rank test) spatial frequency
foveated targets. Thus, foveal SC neurons signaled the occurrence of intra-saccadic stimulus
changes, a kind of prediction error from the expected pre-saccadic extrafoveal target image
appearance.

We also considered the possibility of systematic differences in landing positions across pre-
saccadic target spatial frequencies and sessions. These did indeed exist (due to expected
variances in saccades), but the retinotopic positions of the foveated image patches were
always encompassing our neurons' RF hotspot locations (S3a Figure). Moreover, variations in
landing positions by the saccades across conditions (thus causing variations in foveated
retinotopic image positions relative to the foveal RF's) would not be expected to always
systematically elevate reafferent visual responses; rather, they might equally likely also
diminish responses if the eye landing position resulted in a sub-optimal visual stimulation of
the RF by the image patch. This was clearly not the case in our data (Fig. 3).

Therefore, our results so far indicate that, even though our foveal SC neurons were not
directly visually stimulated pre-saccadically by the extrafoveal saccade targets, they were
still, nonetheless, sensitive to these targets' pre-saccadic appearances: intra-saccadically
introduced conflicts in post-saccadic target appearances resulted in elevated reafferent
response magnitudes. Such elevation, under prediction error situations, is a hallmark of
predictive coding in a variety of brain areas (7, 8, 33, 34, 35, 36).

Independence of trans-saccadic error signaling from neuron preference

We next checked whether an individual neuron's preference for a particular foveated image
feature dictated whether the neuron was sensitive to a conflict with pre-saccadic image

features. We assessed neuron preference from the control conditions of Figs. 1-3: we
measured the reafferent response strength when foveating a low spatial frequency (that was
unchanged across the saccade) and compared it to the case when the saccade landed on a
high spatial frequency (again with no intra-saccadic change). In our database, 30/81 neurons
(37.04%) had numerically stronger visual reafferent responses when the foveated image
feature was a high spatial frequency, whereas 51/81 neurons (62.96%) responded more
strongly for the low spatial frequency patch. This is consistent with SC neurons generally
preferring low spatial frequencies (12, 29) (Fig. 4a). Next, we plotted the results of Fig. 3e, f
against each other. Specifically, we measured, for each neuron, the difference in reafferent
response strength between the low-to-high image flip condition and the corresponding
control scenario (high-to-high), and we plotted it against the difference between the high-to-
low image flip condition and the corresponding control condition (Fig. 4b). There was no
correlation between the two measures (r = 0.0062; p = 0.956), suggesting that we were not
merely observing a single, universal sensitivity (per neuron) for just any intra-saccadic image
update. Moreover, when we colored the neurons according to whether they preferred high
or low spatial frequencies in Fig. 4a, we found that neurons from both groups could signal a
trans-saccadic inconsistency in foveated image appearance (by elevated reafferent response
strengths; Fig. 4b), and for both types of spatial frequency changes. For example, 38/51
neurons (74.5%) preferring the low spatial frequency had an elevated reafferent response
when the pre-saccadic image was of a high spatial frequency. For the neurons preferring the
high spatial frequency, 17/30 neurons (56.7%) signaled a trans-saccadic inconsistency by
elevating their reafferent response strengths (even though this proportion was less than for
neurons preferring the low spatial frequency, the trend was still in the same direction). Thus,
the results of Fig. 2, 3 were a general property of our population, and could happen
regardless of individual neuron preferences for the foveated image features.
368
369

Figure 4 Elevations of foveal reafferent response strengths with trans-saccadic image inconsistencies for both
neurons preferring low and neurons preferring high spatial frequency textures. (a) We assessed neuron
preferences for low or high spatial frequency textures based on their reafferent responses in the control trials.
More neurons preferred low spatial frequencies, as might be expected (12, 29). The different colors are used to
help in classifying the different neuron groups in **b**. **(b)** For the same neurons, we plotted whether they elevated
or decreased their reafferent responses on image-flip trials. Neurons with both preferences in **a** could elevate
their reafferent responses with trans-saccadic image inconsistencies. Thus, there was no systematic dependence
of the results of Figs. 2-3 on neuron preferences.
379

In Fig. 4b, there were additional interesting differences in the foveal neurons' abilities to signal intra-saccadic stimulus changes in their reafferent responses. For example, when landing on a low spatial frequency image patch (x-axis), trans-saccadic error signaling seemed to be slightly more systematic than when landing on a high spatial frequency image patch, independent of neuron preferences (79.01% of all neurons elevated their reafferent response strength in the former case, as opposed to 67.9% in the latter, but this difference was not significant; $p > 0.5$; binomial distribution fit). In addition, going from a low spatial frequency pre-saccadic extrafoveal image to a high spatial frequency post-saccadic foveal image had neurons with more marked modulations in either direction (elevations or reductions relative to control; y-axis). These observations might suggest two possibilities. One, it could be that the SC might favor certain types of trans-saccadic image inconsistencies to "flag" by its foveal neurons (such as high-to-low spatial frequency image flips). Alternatively, since high spatial frequencies are significantly less visible and less salient than low spatial frequencies extrafoveally (due to visual acuity and contrast sensitivity functions of the visual system), a flip from high to low spatial frequency may elicit a more robust prediction error signal (on average) than a flip from an already highly salient extrafoveal image feature (low spatial frequency) to a less salient one. In a later experiment described below, we show evidence consistent with this latter possibility.

Trans-saccadic error signaling does not occur with simulated saccades

To check whether a saccade was needed for foveal neurons to signal a trans-saccadic inconsistency between extrafoveal and foveal visual image appearances, we also ran a simulated saccade condition. In this condition, the monkeys always maintained gaze fixation, and we rapidly translated the extrafoveal image patch (initially appearing to the right or left of fixation at the same eccentricity as in the active saccade task) towards the fovea (Methods). On some trials, the patch had a fictive image change (control), and, on others, it had a real one, just like in the saccade task. Naturally, due to our discretized (both spatially and temporally) display, we could not mimic the exact saccadic eye movement profile, but we used a similar duration and peak speed of image translations (Methods).

During maintained fixation, externally translating extrafoveal images into the neurons' foveal RF's did not elevate the neurons' visual responses to the stimuli on the image flip conditions relative to control; that is, there was no added spiking when the experienced foveal image feature was different from the pre-translation extrafoveal one. Consider, for example, Fig. 5a, b, showing the responses of the same neuron of Figs. 1, 2 but in the simulated saccade condition. First, it should be noted that the neuron's activity was not completely paused during the image translation phase of the trials (compare Fig. 5a, b to Fig. 2b, c around the time of the stimulus update event). This is expected: the image translation in this condition was no different (conceptually) from a stimulus onset event used in many visual neurophysiology experiments; thus, the neuron just maintained its baseline firing rate until new sensory evidence entered into its RF. On the other hand, during real saccades, extrafoveal neurons were bursting and, simultaneously, foveal neurons paused (30, 37, 38, 39). More importantly, the same neuron's response to the afferent sensory input itself (the

circular patch entering it's RF) was the same, for a given spatial frequency, irrespective of the image flip condition. That is, when the stimulus entering the RF was a low spatial frequency, it did not matter whether the initial extrafoveal stimulus was a low or high spatial frequency (Fig. 5a). Similarly, when the stimulus entering the RF was a high spatial frequency, it did not matter whether the extrafoveal stimulus was the same or different (Fig. 5b). We also assessed this statistically: we measured the peak firing rate in a prescribed interval (Methods) from (fictive or real) stimulus update time; whether the final foveal stimulus was of low or high spatial frequency, there was no difference in visual response strength relative to the corresponding control condition (p=0.1303; Wilcoxon rank sum test comparing low-to-low and high-to-low; and p=0.1260; Wilcoxon rank sum test comparing high-to-high and low-to-high trials). Thus, besides confirming our earlier assertion that our foveal neurons were not sensitive to extrafoveal image appearance during fixation, these results indicate that signaling of trans-saccadic extrafoveal-versus-foveal image inconsistencies in Figs. 2-4 above was contingent upon saccade generation.

Figure 5 Lack of foveal neural response elevations in the absence of real saccadic eye movements. (a, b) Responses of the same example neuron of Figs. 1, 2, but this time in the absence of saccades. The monkey maintained gaze fixation, and we rapidly translated the extrafoveal image patch to the fovea (Methods). This resulted in a visual response to the patch entering the neuron's RF, and there was no prior pause in neural activity because there was no actual saccade generation (compare to Figs. 1, 2). Importantly, whether there was an intra-translation image flip or not, the neuron simply reflected the visual appearance of the final foveal stimulus in its RF. Thus, the neuron was neither sensitive to the pre-translation extrafoveal appearance, nor was it sensitive to an intra-translation image change. Error bars denote SEM across trials.

Across the population, we reached similar conclusions. Figure 6 shows results of analyses like those in Fig. 3, but for the present simulated saccade condition. For each foveated image feature, there were no statistically significant differences between control and image flip conditions (p=0.44 in Fig. 6e and p=0.055 in Fig. 6f, Wilcoxon signed-rank test in each case; and p=0.71 in Fig. 6g and p=0.09 in Fig. 6h, Wilcoxon signed-rank test in each case), indicating that the foveal neurons did not "detect" a change from pre-translation extrafoveal

461   image appearance like we saw with real saccades. Consistent with this, there was no
462   significant correlation between the effects of real and simulated saccades in the neurons in
463   which we ran both conditions within the same session (S4 Figure; r=-0.1034, p=0.4701 for
464   panel **a**; r=-0.1703, p=0.2323 for panel **b**). Moreover, the population temporal dynamics (Fig.
465   6a-d) were consistent with the temporal dynamics shown in the example neuron of Fig. 5.
466
467   Thus, when foveal SC neurons were visually-stimulated through exogenous means, and not
468   through self-generated saccades, they expectedly only reflected the instantaneous sensory
469   evidence entering into their RF's. Signaling of trans-saccadic inconsistencies between
470   extrafoveal and foveal image appearances by foveal SC neurons was contingent upon
471   saccade generation.
472
473

474
475 **Figure 6 Lack of foveal neural response elevations in the absence of real saccadic eye movements.** Analyses
476 similar to those in Fig. 3, demonstrating a lack of foveal response elevations in the absence of saccades. Error
477 bars denote SEM in **a**, **b**, and 95% confidence intervals in **c, d**. All other conventions are like in Fig. 3 above.

Extrafoveal stimulus visibility interacts with trans-saccadic error signaling

Because extrafoveal visual acuity is not as good as that in the fovea, including in the SC's retinotopic visual field representation (12, 21, 40), one should not necessarily expect universally successful signaling of trans-saccadic prediction errors by foveal SC neurons. Specifically, if the relevant extrafoveal visual feature was only subtly changed intra-saccadically, then trans-saccadic elevations of foveal SC reafferent responses might be reduced or eliminated: when extrafoveal visibility is already limited, there is little evidence to allow signaling a trans-saccadic image mismatches.

To test this, we employed object shape changes; rather than changing the spatial frequency of the texture inside the saccade target across saccades, we changed only the outline form encompassing the texture. For either a low or high spatial frequency texture, the pre-saccadic extrafoveal target was a circle, and the post-saccadic foveal target was now a square. These trials were interleaved with all of the other trials of Figs. 1-4 (Methods). We hypothesized that, with the low spatial frequency texture, extrafoveal target visibility was dominated by the highly salient texture and not by the outline shape (especially for SC neurons preferring low spatial frequencies; ref. (12)). On the other hand, with the less salient fine texture, the form (circle versus square) could be better distinguished extrafoveally, and could, in turn, be sensed by the foveal neurons if it was changed trans-saccadically. Thus, we expected that trans-saccadic signaling of shape changes (between the pre-saccadic extrafoveal and post-saccadic foveal image scenarios) would be weakened with the low spatial frequency texture embedded within the shape.

Figure 7a, b shows the results of the same example neuron of Figs. 1, 2, 5 under these additional image manipulations. When the shape of the saccade target was changed across saccades, this change was only detected by the foveal SC neuron when the texture inside the shape was a high spatial frequency. We also confirmed this statistically. Across trials, the peak visual reafferent response in the shape-change trials was not different from that in the control trials for the low spatial frequency texture embedded within the saccade targets (p=0.7587; Wilcoxon rank sum test; Fig. 7a). However, the reafferent response was elevated relative to control in the shape-change trials when the embedded texture was of high spatial frequency (p=2.67×10^{-4}; Wilcoxon rank sum test; Fig. 7b).

518
Figure 7 Generalizability of trans-saccadic error signaling to the case of intra-saccadic outline shape changes.
(a) Visual reafferent responses of the same foveal neuron of Figs. 1, 2, 5. Here, we compared the control condition
(blue; same as in Figs. 1, 2) of landing on a circular target containing a low spatial frequency texture to the one
in which only the outline form was changed (but the foveated texture unaltered). There was no elevation of
reafferent response strength by the intra-saccadic image change. We hypothesize that the salient low spatial
frequency texture dominated the pre-saccadic visibility of the extrafoveal target, rendering the shape harder to
discern pre-saccadically. **(b)** Consistent with this, when the embedded texture was a high spatial frequency, and
thus putatively less salient, a trans-saccadic shape mismatch was successfully detected by the neuron. **(c, d)** Same
as in Fig. 3e, f, but for the shape-change manipulation. Across the population, there was a significant elevation
in both cases (p=0.0025 in **c** and p=0.0056 in **d**; Wilcoxon signed-rank test). **(e, f)** For the neural modulation
indices, there was also a positive population index in both cases (p=0.03 in **e** and p=0.0016 in **f**; Wilcoxon signed-
rank test). However, as mentioned in the text, the modulation indices in **e** were significantly weaker than those
in Fig. 3g, suggesting a weaker effect of the shape changes when the embedded textures were of low spatial
frequency. All conventions are similar to those in Figs. 2, 3. Also see S5a, b, e, f Figure for population ROC and
neural firing rate dynamic analyses, supporting the results in this figure.

534
535
536
Such an effect was also present across the population (Fig. 7c-f, using similar formatting to
earlier figures). When the pre- and post-saccadic targets contained within them a low spatial
frequency texture, the median modulation index across the population was 0.0161 (Fig. 7e).
This was lower than the median modulation index seen in Fig. 3g when landing on a low
spatial frequency texture (median 0.0541; p=0.0228; Wilcoxon signed-rank test comparing
the modulation indices in the two conditions). This suggests weakening of trans-saccadic
prediction error signaling when the trans-saccadic change was rendered more subtle (also
see below for further evidence of such scaling effects). Interestingly, when the embedded
texture was of a high spatial frequency, putatively rendering the shape change more visible
due to the lower salience of high spatial frequencies, there was no difference in modulation
indices between the shape-change trials (Fig. 7f) and the spatial-frequency change trials (Fig.
3h; p=0.6530; Wilcoxon signed-rank test comparing the modulation indices in Figs. 3h, 7f).
Once again, these results were not an outcome of variability in saccade landing positions
(S3b Figure).
551
552 Therefore, trans-saccadic signaling of inconsistencies between experienced and expected
553 foveal visual target appearances by SC neurons may fail when the limits on pre-saccadic
554 extrafoveal preview visibility are reached.
555
556

557 *Scaling of error signaling by combined image conflicts*
558
Finally, our shape-change experiments allowed us to next test combined image
manipulations. On intra-saccadic image flip trials, we now flipped both the spatial frequency
and the encompassing outline form at the same time. This way, we could ask whether
signaling of trans-saccadic prediction errors by foveal SC neurons may be viewed as just an
"event detector" of any trans-saccadic image inconsistency, or whether the signaling
strength might scale with the magnitude of the feature changes generated by the intra-
saccadic image flips. From the comparisons made between the neural modulation indices of
Figs. 3, 7, we had reason to believe that trans-saccadic prediction error signaling by foveal SC
neurons is more than just a simple event detection process.
568
At the individual neuron level, consistent with the differences between the Fig. 3 and Fig. 7
population results, we did indeed notice that the modulation of reafferent response
strengths on combined feature change trials could be significantly stronger than its
modulation on individual feature change trials. For example, Fig. 8a, b shows the results of
the same neuron as in Figs. 1, 2, 5, 7 above, but now with both a shape and spatial
frequency change across saccades. When saccades landed on a low spatial frequency texture
(plus a changed shape; Fig. 8a), there was an elevated reafferent response relative to
control, and this elevation was larger in magnitude than that observed on either the spatial-
frequency (Fig. 2b) or shape (Fig. 7a) change trials alone: the difference in peak reafferent
response from control was 26.82 spikes/s in the combined change trials, whereas it was
16.11 and 1.02 spikes/s in Fig. 2b and Fig. 7a, respectively. When the saccades landed on a
high spatial frequency target instead, and again in the combined change trials (Fig. 8b), the

reafferent response elevation was larger than that observed on the shape-only trials (Fig. 7b;
38.74 versus 19.01 spikes/s), and also more robust than that observed for the spatial
frequency trials alone (Fig. 2c; 38.74 versus 27.06 spiks/s). Thus, combining multiple intra-
saccadically altered image features strengthened the elevations of reafferent response
magnitudes caused by inconsistencies between the pre- and post-saccadic visual image
appearances.
587
588

589
Figure 8 Scaling of trans-saccadic prediction error signaling by foveal SC neurons with combined intra-saccadic
feature changes. (a, b) Results from the same example neuron of Figs. 1, 2, 5, 7. Here, relative to the same control
trials of Figs. 1, 2, we checked reafferent responses when both the outline shape and the embedded spatial
frequency texture within it were altered intra-saccadically. Both when landing on a low (**a**) or high (**b**) spatial
frequency texture, the reafferent responses of the neuron were elevated more strongly than either for only the

spatial frequency (Fig. 2) or for only the shape (Fig. 7) change trials. **(c, d)** Consistent trans-saccadic error
elevations in foveal neuron reafferent responses were observed across the population (p=0.0033 for **c** and
p=6.22x10^{-5} for **d**; Wilcoxon signed-rank test). **(e, f)** Similarly, population modulation indices were also positive
(p=0.0023 for **e** and p=6.6x10^{-5} for **f**; Wilcoxon signed-rank test). Error bars denote SEM, and all conventions are
the same as in earlier figures. Also see S5c, d, g, h Figure for population ROC and neural firing rate dynamic
analyses, supporting the results in this figure.

Next, after confirming that the combined image feature inconsistencies still caused trans-
saccadic error signaling in our population (Fig. 8c-f; S5c, d, g, h Figure), we tested to what
extent the neural modulation indices on combined (shape plus spatial frequency) image
trials approached the sum of the neural modulation indices obtained in the individual image
manipulations separately. Specifically, our control trials were the same for both types of
experiments, since these control conditions constituted only circular saccade targets and a
single spatial frequency (Methods). Thus, on trials with only spatial frequency flips or only
shape flips, the control condition that we were comparing to was the same. This allowed us
to compute neural modulation indices on the trials with combined image changes and see
whether they approached the sum of the individual modulation indices.

For each neuron, we plotted, on the x-axis, the sum of neural modulation indices obtained
from the modulation indices of the individual image manipulations (shown in Figs. 3 and 7).
Then, on the y-axis, we plotted the actual (measured) neural modulation indices obtained
from the combined image manipulations (Fig. 8e, f). There was indeed a clear positive
correlation, both when landing on a low (Fig. 9a; r=0.6632; p<0.000001) and high (Fig. 9b;
r=0.8521; p<0.000001) spatial frequency texture. Therefore, signaling of trans-saccadic
prediction error by foveal SC neurons does scale (albeit sub-additively when landing on a low
spatial frequency texture) with the amount of feature conflicts experienced by the neurons.

625
Figure 9 Scaling of trans-saccadic error signaling in the foveal SC by combined intra-saccadic image changes.
(a, b) When landing on either a low (**a**) or high (**b**) spatial frequency texture, we computed on the x-axis the sum
of modulation indices obtained from the experimental trials with only spatial frequency intra-saccadic changes
(Fig. 3) and only shape changes (Fig. 7). Then, on the y-axis, we calculated the actual measured modulation
indices from the trials containing both spatial frequency and shape intra-saccadic image updates. In both cases,
there was a clear positive correlation, suggesting a scaling of prediction error signaling effects in the foveal SC

with the magnitude of feature inconsistencies across saccades (correlation coefficients are indicated in each
panel; $p < 0.000001$ in each case).

We should also emphasize here that we additionally ran the simulated saccade experiments,
on the same neurons as in Fig. 6, with the shape-only and combined (shape plus spatial
frequency) change experiments. In both cases, we did not observe systematic elevations in
sensory responses to the afferent signals received by the translating images into the
neurons' RF's. This can be seen in S6 Figure.

Discussion

In this study, we investigated the activity of foveal SC neurons (21) as these neurons were
visually stimulated by saccade-induced translations of extrafoveal visual stimuli into their
RF's. We tested the hypothesis that foveal SC neurons can receive, trans-saccadically, an
expected appearance of the foveated saccade target (based on this target's pre-saccadic
visual features). We picked only neurons that were not visually stimulated by the extrafoveal
saccade target. Then, during the saccade, we altered the target's appearance, such that the
post-saccadic foveated image feature (now visually driving the neurons) was different from
that expected from the pre-saccadic visual property. We found that visual reafferent
responses were systematically elevated if there was a trans-saccadic mismatch in saccade
target appearance. This elevation did not happen with simulated saccades, and it also scaled
with the extent of feature mismatches that we introduced across saccades.

We interpret our results within the framework of predictive coding (7, 8, 33, 34). In this
framework, neural responses would be elevated under so-called prediction error situations.
In our case, a stable saccade target across saccades is the most expected sensory
consequence of eye movements (that is, having no prediction error). This is because the real
world is generally stable across rapid eye movements (41, 42), despite translations of its
image on the retina. Thus, experiencing a different foveal image from that expected based
on the pre-saccadic extrafoveal saccade-target appearance constitutes a trans-saccadic
prediction error; accordingly, the visual environment can be interpreted as having been
unstable across the saccade.

Consistent with our interpretation, some of the oldest ideas about perceptual stability in the
face of eye movements have revolved around concepts of "template matching": a pre-
movement template (or prediction) is formed, and post-saccadic perceptual stability is
established when a post-movement image matches the pre-emptively formed template (3,
9). Neurobiologically, changes in visual RF shapes and locations do indeed occur peri-
saccadically (43, 44, 45, 46, 47, 48, 49, 50). In sensory-motor areas, such as frontal eye field
(FEF), lateral intraparietal area (LIP), and SC, these RF changes have long been termed
"predictive remapping" (43, 44, 50). However, while the exact details of spatial updating
continue to be under scrutiny (46, 48, 49, 51, 52), such spatial updating remains significantly
and conceptually different from our focus here: how is visual sensation of a stably present

visual object in the environment handled when an eye movement brings this object inside a
given neuron's RF? In particular, how are visual feature representations, not just retinotopic
spatial representations, handled across saccades?
682
One aspect of this question was already addressed previously by Crapse and Sommer (34),
albeit again with a focus on spatial location. These authors recorded from extrafoveal FEF
neurons. They presented a visual stimulus that was stable and brought into the visual RF of
an FEF neuron by a saccade. When the stimulus remained stable across the saccade, the
visual reafferent response to the stimulus (as it entered into the RF of the recorded neuron)
was not as strong as when the same stimulus was experimentally translated intra-
saccadically (such that it still occupied the same post-saccadic RF location). Thus, the
recorded FEF neuron "signaled" (by elevating its firing rate) a deviation from the "predicted"
location of a stable stimulus, even though the stimulus was outside its RF pre-saccadically;
the neuron, therefore, detected the unexpected intra-saccadic change in stimulus position.
We do not know of such effects in the SC, and certainly not with visual features other than
spatial location. Thus, we attempted to fill this gap here, and we also used target features,
rather than location, as the manipulation of interest for us. Importantly, we additionally did
this in the foveal visual processing domain, which is, in any case, one of the most important
aspects of active visual scanning by saccadic eye movements. Our results, in this sense,
support the recent experiments demonstrating that foveal perception is influenced, right
before saccade onset, by extrafoveal saccade target appearance (53, 54). These results are
also in line with previously described peripheral preview effects; in such effects, changes in
both behavioral performance (such as reaction times) and cortical brain activity have been
observed if the post-saccadic foveated image was rendered different from the pre-saccadic
preview (4, 55, 56, 57).
704
Having said that, we do not wish to imply a complete dissociation of our effects from
predictive spatial remapping models. We believe that accounts of predictive remapping of
spatial representations that include remapping of visual feature information to the fovea
likely contribute to our observations (53, 57, 58, 59, 60, 61). In that regard, we find it
intriguing that we observed evidence of a kind of "capacity limit" to the utility of pre-
saccadic previews for our foveal SC neurons to signal trans-saccadic prediction errors. For
example, our neural modulation effects scaled with the amount of intra-saccadic feature
changes that we introduced (e.g. Figs. 8, 9) and also with the putative pre-saccadic
extrafoveal visibility of the saccade target (e.g. Fig. 7). This makes sense. In fact, spatial
resolution changes strongly with retinotopic eccentricity, and SC spatial frequency tuning
curves also exhibit a gradient of decreasing preferred spatial frequency with increasing
eccentricity (12). Thus, if the extrafoveal preview is limited in its information, then so is the
ability to detect intra-saccadic feature changes from this preview. This is not necessarily a
disadvantage for trans-saccadic integration; rather, it confers the visual system with some
tolerance to incidental retinotopic image changes. Otherwise, prediction errors might be
signaled by the brain all too often and thus disrupt visual processing as opposed to
supporting perceptual stability. Interestingly, the limits of such a tolerance window of the
visual system might have been successfully exploited by the famous Mona Lisa painting (62).
Specifically, when viewed peripherally, the Mona Lisa's mouth is perceived as smiling more
than when viewed foveally; low-pass filtering of the painting also revealed a smiling mouth,
suggesting that the peripheral percept is mediated by spatial frequency (62).

The above example suggests that it would be interesting in the future to test foveal SC
neurons with faces and other object stimuli as the saccade targets. Indeed, the (extrafoveal)
SC detects faces (13, 15, 63, 64, 65, 66), and real-life objects in general (13), in its very first
visual responses. Moreover, the (extrafoveal) SC senses object stimuli already in its saccade-
related motor bursts (16). Thus, SC neurons may already have a peripheral "object" preview
both at stimulus onset and at saccade generation. Testing foveal SC neurons with these
ecologically-relevant stimuli across saccades would then allow direct translations of cortical
peripheral preview effects (55, 56) to relevant subcortical brain areas, and to the fovea.

The concept of remapping also invites mechanistic studies of how rerouting of information
happens across saccades. It has been suggested that such rerouting might be facilitated by
oscillatory coherence between retinotopic cortical sites that would experience the pre- and
post-saccadic image (67), and it was also found that saccades might trigger cortical traveling
waves (68). These ideas are interesting to revisit from an SC perspective especially because
classic SC studies have indeed revealed wave-like spreading of SC activity across saccades
(69, 70). In fact, such SC spreading might lead cortical waves in time, and therefore be more
relevant for the predictive mechanisms suggested by our work.

Our choice to study the SC was also motivated by additional factors. For example, besides
integrating visual evidence from the entire visual system (11), the SC issues saccade motor
commands (20). Importantly, and as just mentioned, at the time of saccade generation, such
integrated visual evidence gets reactivated in the SC, exactly when predictions might be
relayed to foveal visual representations according to our framework: SC saccade-related
motor bursts embed within them a sensory representation of the saccade target's visual
properties (16, 19). Thus, our foveal neurons might receive such evidence from the
extrafoveal motor bursts of the SC. It would be interesting in the future to explore whether
such information transfer happens intra-collicularly or through loops outside of the SC,
especially because it is also known that an influence in the reverse direction (from the fovea
to the periphery) additionally occurs (71). This could be done, for example, by reversibly
inactivating the extrafoveal SC representation and checking whether trans-saccadic error
signal by foveal neurons would survive.

We also think that the SC is interesting to study because of its importance for signaling
salience. Classically, such salience has been described in the image domain. That is, when
spatiotemporal movies are played to the retina, then the SC can signal salient "locations" in
retinotopic images at specific time snapshots (22, 23, 24, 25). However, salience can also be
represented in the temporal domain. For example, our results indicate that foveal SC
neurons can signal salient post-saccadic events highlighting a "failure" of expected visual
scene stability as seen by the very same neurons. Thus, the same retinotopic location
responds differentially at different times. We believe that this idea can also extend to non-
saccadic situations. For example, if there are known temporal contingencies in the
environment with certain types of stimuli, then we can expect that SC neurons can still be
sensitive to violations in these contingencies even in the absence of saccades. This can
happen, say, if certain object motions follow strict or systematic physical rules to their
sequencing, such as in the case of biological motion. If the SC does indeed behave as a
temporal salience detector for detecting violations in temporal contiguity across saccades,

then this can be part of a larger framework for building invariant object detection in the
cortex (72, 73).
775

We now turn to potential limitations of our study and its interpretations. For one, it could be
argued that our elevated reafferent responses reflect a relief from adaptation. That is, in the
control trials, the pre-saccadic image was the same as the post-saccadic one, whereas it was
different on the image-flip trials. However, because the SC is retinotopic, adaptation in our
experiments would have been specific to the extrafoveal neurons that saw the pre-saccadic
stimulus and not to our recorded foveal neurons. It is not clear why foveal SC neurons
should reflect the adaptation state of extrafoveal ones. This possibility could potentially be
tested in the future by exploring neural trajectories in the population manifold represented
by foveal SC neurons, both at the time of extrafoveal stimulus presentation and also around
saccade onset. For example, population manifolds of extrafoveal SC neurons are orthogonal
to each other between the stimulus and saccade generation epochs (16), suggesting that
such an approach may reveal more than just potential across-SC adaptation effects. More
importantly, if adaptation was indeed present, then this in itself would constitute a trans-
saccadic transfer of information from the extrafoveal to the foveal SC. This would be directly
consistent with our interpretation that foveal SC may receive information about the state of
extrafoveal scene appearances before saccades.
792

Our simulated saccades were also not direct replicas of real saccadic trajectories. Thus, it
might be argued that our simulated saccades were not "sufficient" to cause elevated
responses in image-flip trials. However, our experience with these types of simulated
saccades from earlier experiments is that they often lead to stronger, rather than weaker,
modulations (74). Also, the fact that we did not see pauses in activity before the visual
responses (Fig. 5) supports our view that the neurons were not stimulated before the image
translations, and that they simply passively received new stimuli entering their RF's. Thus, it
is not too surprising that they might not detect a pre-translation image difference.
801

It would also be interesting in the future to have a perceptual readout from our paradigm.
That is, here, we did not query whether the monkeys experienced perceptual instability or
not. This would be important to establish, and we already have interesting hints from our
current data. Specifically, when we analyzed the timing of the first catch-up saccade after the
primary saccades in our data, we found that it was sensitive to intra-saccadic stimulus
changes, especially when landing on a high spatial frequency texture (S7 Figure); the effect
was consistent with a slight delaying of catch-up saccades on prediction error trials, which is
the expected consequence of elevated foveal SC activity. This motivates further searches of
perceptual readouts of trans-saccadic integration in our paradigm, along with causal
manipulations of SC activity, in order to establish whether prediction error signaling has
direct influences on perceptual experience. Given the role of the SC in attention (75), we
suspect that our neural modulations would be relevant for altering behavior. For example,
we could inactivate the extrafoveal SC and ask whether trans-saccadic error signaling by
foveal SC neurons would survive, and whether perceptual readouts can also be affected.
816

817　Finally, we think that SC neurons with large RF's can potentially contribute to the transfer of
818　information from the extrafoveal neurons at the time of saccade onset (16) to the foveal
819　neurons post-saccadically, exactly for realizing prediction error signaling. In our analyses, we

strictly excluded such neurons (Methods). However, we had plenty of such excluded neurons with subtle elevations in their activity at extrafoveal stimulus onset. That is, their peri-foveal RF's extended out to be marginally driven by extrafoveal stimulus onset. When we analyzed those excluded neurons like we did in this study, we found that they had trans-saccadic modulations that were not too different from those reported here (for example, like the reafferent responses of Figs. 1, 2). Thus, intermediate RF's spanning both pre- and post-saccadic visual representations could be critical for the mechanisms explaining our results, and they could exist through wide-field synaptic connections in the SC (76).

Methods

Experimental animals and ethical approvals

We collected data from two adult, male rhesus macaque (macaca mulatta) monkeys (A and M), aged 8-9 years.

The animals were prepared previously for behavioral and neurophysiological procedures. They each had a head holder for stabilization of head position during the experiments, scleral search coils for eye tracking, and a recording chamber with craniotomy for access to the superior colliculus (SC).

All experiments were approved by ethical committees at the regional governmental offices of Tübingen (Regierungspräsidium Tübingen), under licenses CIN 3/13 and CIN 4/19G, and the experiments complied with all guidelines and regulations.

Laboratory setup

The experiments were conducted using the same setup as in our earlier studies (13). Specifically, we used a custom-built modification to PLDAPS (77), which employed the PsychToolbox (78, 79, 80) for display control. The system synchronized with the OmniPlex system from Plexon for data acquisition and storage.

The use of the PsychToolbox allowed us to synchronize our experimental control system with the graphics card controlling our display. This enabled direct access to the vertical blanking signal of each frame update, which was important for achieving gaze-contingent intra-saccadic display changes in some trials (see details below). In previous studies performed during the earlier years of the laboratory's existence (32, 81, 82, 83), we performed extensive tests of this system with external photodiodes detecting every single frame update. This confirmed that we successfully synchronized our experimental control system with each frame command. Consistent with this, the reafferent responses after saccades (e.g. Figs. 1, 2, 3a, b) had very similar timing across trials, regardless of whether there was an intra-saccadic display change or not (also see Fig. 2a for display timing relative to saccades, and S2 Figure for other eye movement parameters with and without intra-saccadic display changes).

We used a linearized and calibrated CRT display monitor with a refresh rate of 85 Hz and ~34 pixels/deg resolution. The monkeys sat ~72 cm from the display, and experiments were conducted with a gray background on the display (having 26.11 cd/m^2 luminance). White stimuli (for example, for a fixation spot) had a luminance of 79.9 cd/m^2. We tracked eye movements using the magnetic induction technique (84, 85), and we recorded neural activity using 16- or 24-channel V-Probes from Plexon.

Experimental procedures

For response field (RF) mapping, we employed a fixation task. The monkeys maintained fixation on a central fixation spot (black), and we then presented a small white spot at

877 pseudorandom locations across trials (86). The fixation and target spots were each ~10.8 x
878 10.8 min arc in size. We mapped RF's to ensure that there were no visual responses to the
879 extrafoveal visual stimulus onset in the main tasks (e.g. Fig. 1b).
880
For the main saccade task, the monkeys fixated a central (white) fixation spot of the same
dimensions as above. After 300-700 ms from foveating the fixation spot, we presented the
extrafoveal stimulus. This stimulus appeared at 8 or 10 deg to the right or left of the fixation
spot in all sessions except two; for these two sessions, we placed the stimulus at +/- 5.5 deg
horizontally and -4.7 deg vertically, or +/- 4.6 deg horizontally and +4.4 deg vertically. This
placement was dictated by other unrelated experiments that we were simultaneously
performing during these two sessions in the primary visual cortex. The extrafoveal stimulus
was a circle of 3 deg radius. Inside, there could be a vertical sine wave texture of high
contrast (100%), having either 1 cycle/deg (cpd) or 4 cycles/deg (cpd) spatial frequency. We
called the former the low spatial frequency texture and the latter the high spatial frequency
texture. After a delay period of 500-1000 ms, we removed the fixation spot, instructing the
monkeys to generate a saccade towards the extrafoveal stimulus.
893
When eye position exited a virtual window surrounding the fixation spot, the computer
flagged a saccade detection event. This triggered a stimulus update on the very next
graphics display frame update. In control trials, the initial shape was always a circle (with
either low or high spatial frequency embedded within it), and the display update event was
fictive. That is, the stimulus was unchanged in this case (of course, the display itself was
always being refreshed from frame to frame, as expected from such display technology).
Such control trials constituted 50% of all trials within a session. The remaining trials were
split among the different intra-saccadic change conditions. In one case, the initial stimulus
was a circle and the final stimulus was still a circle, but containing the other spatial
frequency. That is, if the initial circle encompassed 1 cpd, then the changed stimulus had 4
cpd. These spatial-frequency change trials constituted 16.67% of all trials. In another case,
the initial shape was a circle containing either the low or spatial frequency texture, and the
final shape was a square (6x6 deg in size) with the same texture. These shape-change trials
constituted 16.67% of all trials in a session. Finally, there were trials with a circle initial
shape; the final shape was a square (6x6 deg) containing a different spatial frequency (shape
plus spatial frequency trials). These constituted 16.67% of all trials in the session. Note that
the stimulus never moved in any of the trials. It was the eye that was moving and causing
the stimulus to move towards the fovea on the retina. After the foveating saccade to the
extrafoveal stimulus, the monkeys maintained fixation for another 500 ms before being
rewarded with juice droplets. We collected 322-1440 trials of this task per session.
914
In some sessions, we also ran a simulated saccade version of the above paradigm. Here, the
monkey always maintained fixation. Instead of a go signal to generate a saccade (removal of
the fixation spot), we simply translated the extrafoveal stimulus towards the center of the
display. The translation proceeded as follows: the target at the initial 8 deg eccentricity was
translated to 5.5 deg in the first frame (i.e. 2.5 deg in 11.76 ms), and then to 1.5 deg in the
second frame (i.e. a shift of 4 deg in 11.76 ms), and finally to the final central position in the
next frame (i.e. a shift of 1.5 deg in 11.76 ms). Thus, the simulated saccades had an
accelerating and decelerating profile with a peak speed of 340 deg/s, which is similar to our
real saccades' peak speeds (e.g. Fig. 2a). Image-change trials involved changing the relevant

image feature in the first frame of the translation. The experiment was otherwise the same
as the real saccade version. We did not run the simulated saccade experiment during the
two sessions in which the saccade target was at an oblique position relative to the fixation
spot. We collected 647-841 trials of this task per session.
928
929

Data analysis

931 We detected all microsaccades and saccades using our established methods (32, 87). In the
932 saccade task, we only accepted trials in which saccades landed within a radius of <2deg from
933 the center of the saccade target. This ensured that the saccade target activated our foveal
934 neurons post-saccadically (S3 Figure).
935
We sorted neurons offline using Kilosort (88). In total, we had 249 neurons in our database.
However, we excluded neurons in which there was an elevation (however small) in firing
rates after extrafoveal stimulus onset in our main task (relative to pre-stimulus firing rates).
This is because we wanted to exclude any remote possibility that our foveal neurons were
visually-stimulated by the pre-saccadic extrafoveal stimulus. We also confirmed this with our
RF mapping task, as well as by measuring post-stimulus activity in our accepted neurons (S1
Figure). We also excluded neurons in which the average firing rate in the interval 0-100 ms
from fictive stimulus update time (i.e. in control trials) was less than 5 spikes/s, to ensure
that we had a proper post-saccadic visual response to analyze. This left us with 81 neurons
that we included in the final database.
946
947 To check for proper intra-saccadic stimulus updates, we collected all saccades and related
948 their parameters (onset, peak velocity, and end times) to the time after stimulus update
949 completion (S2a Figure). We also measured eye speed at the time of stimulus update
950 completion (S2b Figure and Fig. 2a), in order to confirm that we had as much stimulus
951 motion blur on the retina as possible at the time of intra-saccadic target changes.
952
We checked other behavioral metrics of the saccades as well. For example, we plotted radial
eye speeds with and without intra-saccadic stimulus changes (S2c, d Figure), as well as radial
eye displacements (S2e, f Figure). The former plotting allowed us to check whether post-
saccadic drifts (32) were altered by intra-saccadic stimulus changes, and the former and
latter plotting allowed us to investigate whether saccades were truncated (89) or otherwise
altered by the intra-saccadic changes. Finally, we measured the latency of the first catch-up
saccade after the primary saccade (S7 Figure). This allowed us to check whether elevations
in reafferent responses in foveal SC neurons (on intra-saccadic change trials) altered
subsequent saccade timing.
962
To measure reafferent response strength, we calculated the peak firing rate in the interval
after the end of stimulus updates. For low spatial frequency foveated targets, the interval
was 0-120 ms from real or fictive stimulus update; for high spatial frequency foveated
targets, it was 0-150 ms. The two different intervals reflected the fact that responses were
delayed for high spatial frequency foveated stimuli (12, 29). For the simulated saccade
version of the task, we used the same measurement intervals relative to the stimulus update
times. Note that in all cases, we focused only on the initial reafferent response and not also
later modulations in firing rate that could happen. This is because such later modulations

coincided with potential catch-up saccades, which (due to their small size) were expected to
significantly alter foveal SC activity. Also note that we used stimulus update time as the
alignment point for measuring our reafferent responses. Because we had a fictive update
time stamp also for the control trials, and because online saccade detection was quite
repeatable across trials (e.g. Fig. 2a), this meant that we had similar alignment of firing rates
for both the control and intra-saccadic stimulus change trials. This, in turn, allowed for a fair
comparison between the conditions. In addition, the stimulus update time was always at a
time of maximal suppression of neural activity (e.g. Figs. 2b, c, 3a, b, 7a, b, 8a, b), which was
a suitable point of temporal reference. Having said that, we also tried all analyses after
aligning to saccade end instead, and we reached the very same conclusions. Firing rate
curves looked shifted in time, confirming the repeatable nature of our saccade-contingent
stimulus updates. Our choice to use stimulus update times made it easier for us to also use
this time in the cases in which no real saccades were generated.

Our modulation indices were calculated for each neuron as follows: firing rate on the intra-
saccadic change condition of interest minus firing rate on the corresponding control
condition, divided by the sum of the firing rates.

We performed statistical analyses in two ways. First, across neurons, we measured the firing
rate on the intra-saccadic change of interest and the firing rate on the corresponding control
condition. We then performed a Wilcoxon signed-rank test across neurons. Second, we
checked whether the population modulation indices were different from zero using a
Wilcoxon signed-rank test. For all depictions of population modulation indices, we also
highlighted the neurons that were individually significant in a given analysis. Individual
significance was assessed by comparing the firing rate measures of the population of control
trials to the firing rate measures of the population of trials in a given intra-saccadic stimulus
change condition, using a permutation test across trials (with 10000 repetitions).

Finally, we also depicted the neural population dynamics in two additional ways. In one, we
plotted the average normalized population firing rate curve in a given condition. To do so, we
obtained the average firing rate curve of each neuron and each condition (across trials of the
same condition). Then, we normalized each average firing rate curve of the neuron by the
peak of the average firing rate curve of the corresponding control condition (low-to-low
spatial frequency when landing on a low spatial frequency texture, or high-to-high spatial
frequency when landing on a high spatial frequency texture). Then, we averaged across
neurons and depicted mean and SEM curves. In the second approach, we performed
receiver-operating-characteristic (ROC) analyses, like we recently did (16). Here, we were
testing the discrimination performance between a stimulus change condition and a control
condition as a function of time relative to the stimulus change. If reafferent responses were
elevated on intra-saccadic change trials, then the area under the ROC curve was expected to
elevate relative to pre-saccadic periods. All of our area under the ROC curve plots included
95% confidence intervals around them.

1013

1014

Competing interests statement

The authors declare no competing interests.

Data availability statement

Data tables are provided in Supplementary Information.

Acknowledgements

We were funded by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation) through the Special Priority Programme: "SPP 2411 Sensing LOOPS: cortico-subcortical interactions for adaptive sensing" (project numbers: 520617944 and 520283985, HA 6749/11-1). We were also funded by the DFG project BO5681/1-1. The funders did not play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Wurtz RH. Neuronal mechanisms of visual stability. Vision Res. 2008;48(20):2070-89.
2. Watson TL, Krekelberg B. The relationship between saccadic suppression and perceptual stability. Curr Biol. 2009;19(12):1040-3.
3. Bridgeman B, van der Heijden AH, Velichkovsky BM. A theory of visual stability across saccadic eye movements. Behav Brain Sci. 1994;17:247-92.
4. Melcher D. Visual stability. Philos Trans R Soc Lond B Biol Sci. 2011;366(1564):468-75.
5. Wurtz RH. Corollary Discharge Contributions to Perceptual Continuity Across Saccades. Annu Rev Vis Sci. 2018;4:215-37.
6. Ross J, Morrone MC, Goldberg ME, Burr DC. Changes in visual perception at the time of saccades. Trends Neurosci. 2001;24(2):113-21.
7. Baltaretu BR, Dunkley BT, Stevens WD, Crawford JD. Occipital cortex is modulated by transsaccadic changes in spatial frequency: an fMRI study. Scientific reports. 2021;11(1):8611.
8. Baltaretu BR, Stevens WD, Freud E, Crawford JD. Occipital and parietal cortex participate in a cortical network for transsaccadic discrimination of object shape and orientation. Scientific reports. 2023;13(1):11628.
9. McConkie GW, Currie CB. Visual stability across saccades while viewing complex pictures. J Exp Psychol Hum Percept Perform. 1996;22(3):563-81.
10. Nuthmann A, Clayden AC, Fisher RB. The effect of target salience and size in visual search within naturalistic scenes under degraded vision. J Vis. 2021;21(4):2.
11. Hafed ZM, Hoffmann KP, Chen CY, Bogadhi AR. Visual Functions of the Primate Superior Colliculus. Annu Rev Vis Sci. 2023;9(1):361-83.
12. Chen CY, Sonnenberg L, Weller S, Witschel T, Hafed ZM. Spatial frequency sensitivity in macaque midbrain. Nat Commun. 2018;9(1):2852.
13. Bogadhi AR, Hafed ZM. Express detection of visual objects by primate superior colliculus neurons. Scientific reports. 2023;13(1):21730.
14. Bogadhi AR, Buonocore A, Hafed ZM. Task-irrelevant visual forms facilitate covert and overt spatial selection. J Neurosci. 2020;40(49):9496-506.
15. Soares SC, Maior RS, Isbell LA, Tomaz C, Nishijo H. Fast Detector/First Responder: Interactions between the Superior Colliculus-Pulvinar Pathway and Stimuli Relevant to Primates. Front Neurosci. 2017;11:67.
16. Baumann MP, Bogadhi AR, Denninger AF, Hafed ZM. Sensory tuning in neuronal movement commands. Proc Natl Acad Sci U S A. 2023;120(38):e2305759120.
17. Zhang T, Malevich T, Baumann MP, Hafed ZM. Superior colliculus saccade motor bursts do not dictate movement kinematics. Commun Biol. 2022;5(1):1222.
18. Willeke KF, Tian X, Buonocore A, Bellet J, Ramirez-Cardenas A, Hafed ZM. Memory-guided microsaccades. Nat Commun. 2019;10(1):3710.
19. Hafed ZM. Superior colliculus peri-saccadic field potentials are dominated by a visual sensory preference for the upper visual field. iScience. 2025;28(3):112021.
20. Gandhi NJ, Katnani HA. Motor functions of the superior colliculus. Annual review of neuroscience. 2011;34:205-31.
21. Chen CY, Hoffmann KP, Distler C, Hafed ZM. The Foveal Visual Representation of the Primate Superior Colliculus. Curr Biol. 2019;29(13):2109-19 e7.

22. Veale R, Hafed ZM, Yoshida M. How is visual salience computed in the brain? Insights
from behaviour, neurobiology and modelling. Philos Trans R Soc Lond B Biol Sci.
2017;372(1714).
23. White BJ, Berg DJ, Kan JY, Marino RA, Itti L, Munoz DP. Superior colliculus neurons
encode a visual saliency map during free viewing of natural dynamic video. Nat Commun.
2017;8:14263.
24. White BJ, Kan JY, Levy R, Itti L, Munoz DP. Superior colliculus encodes visual saliency
before the primary visual cortex. Proc Natl Acad Sci U S A. 2017;114(35):9451-6.
25. Heeman J, White BJ, Van der Stigchel S, Theeuwes J, Itti L, Munoz DP. Saliency
Response in Superior Colliculus at the Future Saccade Goal Predicts Fixation Duration during
Free Viewing of Dynamic Scenes. J Neurosci. 2025;45(3).
26. Herman JP, Krauzlis RJ. Color-Change Detection Activity in the Primate Superior
Colliculus. eNeuro. 2017;4(2).
27. Cavanaugh J, Wurtz RH. Subcortical modulation of attention counters change
blindness. J Neurosci. 2004;24(50):11236-43.
28. Cone JJ, Mitchell AO, Parker RK, Maunsell JHR. Stimulus-dependent differences in
cortical versus subcortical contributions to visual detection in mice. Curr Biol.
2024;34(9):1940-52 e5.
29. Khademi F, Chen C-Y, Hafed ZM. Visual feature tuning of superior colliculus neural
reafferent responses after fixational microsaccades. Journal of Neurophysiology.
2020;123:2136-53.
30. Hafed ZM, Krauzlis RJ. Goal representations dominate superior colliculus activity
during extrafoveal tracking. J Neurosci. 2008;28(38):9426-39.
31. Arai K, Keller EL, Edelman JA. Two-dimensional neural network model of the primate
saccadic system. Neural Networks. 1994;7(6/7):1115-35.
32. Chen CY, Hafed ZM. Postmicrosaccadic enhancement of slow eye movements. The
Journal of neuroscience : the official journal of the Society for Neuroscience.
2013;33(12):5375-86.
33. Rao RP, Ballard DH. Predictive coding in the visual cortex: a functional interpretation
of some extra-classical receptive-field effects. Nat Neurosci. 1999;2(1):79-87.
34. Crapse TB, Sommer MA. Frontal eye field neurons assess visual stability across
saccades. J Neurosci. 2012;32(8):2835-45.
35. Summerfield C, Egner T, Greene M, Koechlin E, Mangels J, Hirsch J. Predictive codes
for forthcoming perception in the frontal cortex. Science. 2006;314(5803):1311-4.
36. O'Doherty JP, Buchanan TW, Seymour B, Dolan RJ. Predictive neural coding of reward
preference involves dissociable responses in human ventral midbrain and ventral striatum.
Neuron. 2006;49(1):157-66.
37. Munoz DP, Wurtz RH. Fixation cells in monkey superior colliculus. I. Characteristics of
cell discharge. J Neurophysiol. 1993;70(2):559-75.
38. Everling S, Pare M, Dorris MC, Munoz DP. Comparison of the discharge characteristics
of brain stem omnipause neurons and superior colliculus fixation neurons in monkey:
implications for control of fixation and saccade behavior. J Neurophysiol. 1998;79(2):511-28.
39. Hafed ZM, Krauzlis RJ. Similarity of superior colliculus involvement in microsaccade
and saccade generation. J Neurophysiol. 2012;107(7):1904-16.
40. Hafed ZM, Chen CY. Sharper, Stronger, Faster Upper Visual Field Representation in
Primate Superior Colliculus. Curr Biol. 2016;26(13):1647-58.

41.   Wexler M, Panerai F, Lamouret I, Droulez J. Self-motion and the perception of stationary objects. Nature. 2001;409(6816):85-8.

42.   Hafed ZM, Krauzlis RJ. Ongoing eye movements constrain visual perception. Nat Neurosci. 2006;9(11):1449-57.

43.   Duhamel JR, Colby CL, Goldberg ME. The updating of the representation of visual space in parietal cortex by intended eye movements. Science. 1992;255(5040):90-2.

44.   Walker MF, Fitzgibbon EJ, Goldberg ME. Neurons in the monkey superior colliculus predict the visual result of impending saccadic eye movements. Journal of neurophysiology. 1995;73(5):1988-2003.

45.   Tolias AS, Moore T, Smirnakis SM, Tehovnik EJ, Siapas AG, Schiller PH. Eye movements modulate visual receptive fields of V4 neurons. Neuron. 2001;29(3):757-67.

46.   Zirnsak M, Moore T. Saccades and shifting receptive fields: anticipating consequences or selecting targets? Trends Cogn Sci. 2014;18(12):621-8.

47.   Zirnsak M, Steinmetz NA, Noudoost B, Xu KZ, Moore T. Visual space is compressed in prefrontal cortex before eye movements. Nature. 2014;507(7493):504-7.

48.   Hartmann TS, Zirnsak M, Marquis M, Hamker FH, Moore T. Two Types of Receptive Field Dynamics in Area V4 at the Time of Eye Movements? Frontiers in systems neuroscience. 2017;11:13.

49.   Neupane S, Guitton D, Pack CC. Two distinct types of remapping in primate cortical area V4. Nat Commun. 2016;7:10402.

50.   Sommer MA, Wurtz RH. Influence of the thalamus on spatial visual processing in frontal cortex. Nature. 2006;444(7117):374-7.

51.   Neupane S, Guitton D, Pack CC. Dissociation of forward and convergent remapping in primate visual cortex. Curr Biol. 2016;26(12):R491-2.

52.   Wang X, Fung CC, Guan S, Wu S, Goldberg ME, Zhang M. Perisaccadic Receptive Field Expansion in the Lateral Intraparietal Area. Neuron. 2016;90(2):400-9.

53.   Kroell LM, Rolfs M. Foveal vision anticipates defining features of eye movement targets. eLife. 2022;11.

54.   Kroell LM, Rolfs M. The magnitude and time course of pre-saccadic foveal prediction depend on the conspicuity of the saccade target. BioRxiv. 2023.

55.   Huber-Huber C, Buonocore A, Dimigen O, Hickey C, Melcher D. The peripheral preview effect with faces: Combined EEG and eye-tracking suggests multiple stages of trans-saccadic predictive and non-predictive processing. Neuroimage. 2019;200:344-62.

56.   Buonocore A, Dimigen O, Melcher D. Post-Saccadic Face Processing Is Modulated by Pre-Saccadic Preview: Evidence from Fixation-Related Potentials. J Neurosci. 2020;40(11):2305-13.

57.   Melcher D, Colby CL. Trans-saccadic perception. Trends Cogn Sci. 2008;12(12):466-73.

58.   Knapen T, Swisher JD, Tong F, Cavanagh P. Oculomotor Remapping of Visual Information to Foveal Retinotopic Cortex. Frontiers in systems neuroscience. 2016;10:54.

59.   Rolfs M, Jonikaitis D, Deubel H, Cavanagh P. Predictive remapping of attention across eye movements. Nat Neurosci. 2011;14(2):252-6.

60.   Kämmer L, Kroell LM, Knapen T, Rolfs M, Hebart MN. Feedback of peripheral saccade targets to early foveal cortex. BioRxiv. 2025.

61.   Melcher D. Predictive remapping of visual features precedes saccadic eye movements. Nat Neurosci. 2007;10(7):903-7.

62. Livingstone MS. Is it warm? Is it real? Or just low spatial frequency? Science.
2000;290(5495):1299b.
63. Nguyen MN, Matsumoto J, Hori E, Maior RS, Tomaz C, Tran AH, et al. Neuronal
responses to face-like and facial stimuli in the monkey superior colliculus. Front Behav
Neurosci. 2014;8:85.
64. Nguyen MN, Nishimaru H, Matsumoto J, Van Le Q, Hori E, Maior RS, et al. Population
Coding of Facial Information in the Monkey Superior Colliculus and Pulvinar. Front Neurosci.
2016;10:583.
65. Le QV, Le QV, Nishimaru H, Matsumoto J, Takamura Y, Hori E, et al. A Prototypical
Template for Rapid Face Detection Is Embedded in the Monkey Superior Colliculus. Frontiers
in systems neuroscience. 2020;14:5.
66. Yu G, Katz LN, Quaia C, Messinger A, Krauzlis RJ. Short-latency preference for faces in
primate superior colliculus depends on visual cortex. Neuron. 2024;112(16):2814-22 e4.
67. Neupane S, Guitton D, Pack CC. Coherent alpha oscillations link current and future
receptive fields during saccades. Proc Natl Acad Sci U S A. 2017;114(29):E5979-E85.
68. Zanos TP, Mineault PJ, Nasiotis KT, Guitton D, Pack CC. A sensorimotor role for
traveling waves in primate visual cortex. Neuron. 2015;85(3):615-27.
69. Munoz DP, Wurtz RH. Saccade-related activity in monkey superior colliculus. II.
Spread of activity during saccades. J Neurophysiol. 1995;73(6):2334-48.
70. Nakahara H, Morita K, Wurtz RH, Optican LM. Saccade-related spread of activity
across superior colliculus may arise from asymmetry of internal connections. J Neurophysiol.
2006;96(2):765-74.
71. Zhang T, Tian X, Malevich T, Baumann MB, Hafed ZM. Foveal action for the control of
extrafoveal vision. BioRxiv. 2024.
72. Li N, DiCarlo JJ. Unsupervised natural experience rapidly alters invariant object
representation in visual cortex. Science. 2008;321(5895):1502-7.
73. Li N, DiCarlo JJ. Unsupervised natural visual experience rapidly reshapes size-
invariant object representation in inferior temporal cortex. Neuron. 2010;67(6):1062-75.
74. Idrees S, Baumann MP, Franke F, Munch TA, Hafed ZM. Perceptual saccadic
suppression starts in the retina. Nat Commun. 2020;11(1):1977.
75. Krauzlis RJ, Lovejoy LP, Zenon A. Superior colliculus and visual spatial attention. Annu
Rev Neurosci. 2013;36:165-82.
76. Gale SD, Murphy GJ. Distinct representation and distribution of visual information by
specific cell types in mouse superficial superior colliculus. J Neurosci. 2014;34(40):13458-71.
77. Eastman KM, Huk AC. PLDAPS: A Hardware Architecture and Software Toolbox for
Neurophysiology Requiring Complex Visual Stimuli and Online Behavioral Control. Front
Neuroinform. 2012;6:1.
78. Brainard DH. The Psychophysics Toolbox. Spatial vision. 1997;10(4):433-6.
79. Pelli DG. The VideoToolbox software for visual psychophysics: transforming numbers
into movies. Spatial vision. 1997;10(4):437-42.
80. Kleiner M, Brainard D, Pelli DG. What's new in Psychtoolbox-3? (Abstract).
Perception. 2007;36.
81. Hafed ZM, Ignashchenkova A. On the dissociation between microsaccade rate and
direction after peripheral cues: microsaccadic inhibition revisited. J Neurosci.
2013;33(41):16220-35.
82. Chen CY, Ignashchenkova A, Thier P, Hafed ZM. Neuronal Response Gain
Enhancement prior to Microsaccades. Curr Biol. 2015;25(16):2065-74.

83.   Tian X, Yoshida M, Hafed ZM. A Microsaccadic Account of Attentional Capture and
Inhibition of Return in Posner Cueing. Frontiers in systems neuroscience. 2016;10:23.
84.   Judge SJ, Richmond BJ, Chu FC. Implantation of magnetic search coils for
measurement of eye position: an improved method. Vision Res. 1980;20(6):535-8.
85.   Fuchs AF, Robinson DA. A method for measuring horizontal and vertical eye
movement chronically in the monkey. J Appl Physiol. 1966;21(3):1068-70.
86.   Malevich T, Zhang T, Baumann MP, Bogadhi AR, Hafed ZM. Faster Detection of
"Darks" than "Brights" by Monkey Superior Colliculus Neurons. J Neurosci.
2022;42(50):9356-71.
87.   Bellet ME, Bellet J, Nienborg H, Hafed ZM, Berens P. Human-level saccade detection
performance using deep neural networks. J Neurophysiol. 2019;121(2):646-61.
88.   Pachitariu M, Steinmetz NA, Kadir SN, Carandini M, Harris KD. Fast and accurate spike
sorting of high-channel count probes with KiloSort. Advances in Neural Information
Processing Systems (NIPS 2016). 2016;29.
89.   Buonocore A, Chen CY, Tian X, Idrees S, Munch TA, Hafed ZM. Alteration of the
microsaccadic velocity-amplitude main sequence relationship after visual transients:
implications for models of saccade control. J Neurophysiol. 2017;117(5):1894-910.
1241
1242

1243 **Supporting figure legends**

1244

1245

S1 Figure. Lack of visually-evoked bursts in the foveal neurons at the time of extrafoveal
stimulus onsets. (a) For all neurons included in our main analyses of Figs. 1-4, 7-9, we
measured neural activity 50-150 ms after extrafoveal stimulus onset (during maintained gaze
fixation), and we compared it to pre-stimulus activity. Here, the appearing stimulus was a
circular patch containing a low spatial frequency texture. There was a reduction in activity,
rather than visual bursts. **(b)** Similar observations for the case in which the extrafoveal
stimulus had a high spatial frequency texture embedded within it.

1253

1254
1255 **S2 Figure. Confirmation of intra-saccadic stimulus updates, as well as confirmation of a lack**
1256 **of truncation of primary saccades. (a)** For all trials containing a real or fictive image update
1257 (from the basic paradigm of spatial frequency changes and circular outline shapes of the

saccade targets), we measured (also across all sessions) the times of saccade onset, saccade peak velocity, and saccade end relative to when the image flip event was completed. We always had intra-saccadic image updates in our experiments. **(b)** Importantly, the eye speed was always high by the end of the image update. Here, for each session, we calculated the average eye speed by the end of the image update event, and we plotted a histogram across sessions. For the great majority of sessions, the image speeds on the retina by the end of intra-saccadic image updates were >150 deg/s. For the two sessions with speeds of ~50 deg/s, the saccade targets were placed a bit closer to the fovea (thus having smaller saccades) because the sessions involved other experiments involving the primary visual cortex (we thus tailored the target locations to the cortical neurons explored during the same sessions for other purposes). However, eye speed was still relatively large during these outlier sessions. **(c, d)** For the main conditions of Figs. 2, 3 we plotted radial eye velocity on control and intra-saccadic stimulus change trials. Each plot shows the average velocity curve across sessions (with horizontal saccade target locations, which were the majority), and the error bars show SEM across sessions. The saccadic profiles were not affected by intra-saccadic display updates, including post-saccadic drift speeds. This means that the saccades were not truncated by the stimulus changes. Similar observations were made for the few oblique sessions. **(e, f)** Similar to **c**, **d** but now plotting radial saccade amplitude across sessions. There was again no evidence of saccade truncation by the intra-saccadic stimulus changes. Similar conclusions were reached for all other experiments (e.g. with shape changes).

S3 Figure. Confirmation of post-saccadic visual stimulation of our foveal neurons by the saccade targets. **(a)** Across trials and sessions from our basic paradigm, we plotted the retinotopic position of the saccade-target patch (circles) upon saccade end. There was variability due to variability in saccade metrics. However, the foveated image patches were always encompassing our recorded foveal neurons (dark gray spots indicating the foveal RF hotspot locations). Thus, we always had robust visual reafferent responses. Note also that such jitter in saccade landing should have weakened reafferent responses in some cases, rather than strengthened them, because it could cause visual stimulation at sub-optimal RF positions. This suggests that our main results (dominated by elevated reafferent responses) are not explained by jitter in saccade landing positions. **(b)** Similar observations from the shape-change trials (Methods) of our paradigm. Here, the foveated stimulus had a square outline instead of a circular one.

1295
S4 Figure. Lack of relationship between the modulation indices associated with real versus
simulated saccades. (a) For a low spatial frequency foveated patch after either real or
simulated saccades, we plotted the modulation indices from Fig. 6g on the x-axis (for
simulated saccades) and those from Fig. 3g on the y-axis (for real saccades). The shown
neurons are those for which we ran both conditions together in the same session. In the
simulated saccade case (x-axis), the modulation indices straddled zero; however, in the real
saccade case (y-axis), the modulation indices were largely positive. Thus, there was no
correlation between the two situations (p=0.4701). **(b)** Similar observations for a high spatial
frequency foveated image (p=0.2323).

1305
1306
1307

1308
S5 Figure. Population firing rate dynamics in the shape change and shape and frequency
change conditions. (a, b) ROC analyses for the experiments of Fig. 7. Whether landing on a
low or high spatial frequency texture, there was an elevation in area under the ROC curve in
the reafferent response epoch. **(c, d)** The effects were larger for the combined shape and
spatial frequency change trials (Fig. 8). **(e, f)** Population firing rate dynamics for the
experiments of Fig. 7. **(g, h)** Population firing rate dynamics for the experiments of Fig. 8. Error
bars denote 95% confidence intervals for **a**-**d** and SEM for **e**-**h**.

1316

S6 Figure. Confirmation that shape-change and shape plus spatial frequency changes had much weaker effects on foveal SC neurons during simulated saccades than during our main experimental manipulations with real saccades. (a, b) Population results like in Fig. 6e, f, but now for the shape change trials and a low (**a**) or high (**b**) spatial frequency post-translation texture. There were weaker differences in visual response strength relative to control than with real saccades. **(c, d)** Similar observations for the shape plus spatial frequency change trials. **(e, f)** Neural modulation indices for **a**, **b** (p=0.8 and 0.049 for **e** and **f**, respectively; Wilcoxon signed-rank test). **(g, h)** Neural modulation indices for **c**, **d** (p=0.22 and 0.40 for **g** and **h**, respectively; Wilcoxon signed-rank test).

S7 Figure. Influence of the neural modulations in this study on the timing of catch-up saccades after the primary eye movements. (a) For the basic experiments of Figs. 2, 3, and when landing on a low spatial frequency texture, the distribution of catch-up saccade times (for the first saccade after foveating the target) was very slightly skewed towards longer

latencies on intra-saccadic change trials, but there were also saccades with very short latency. **(b, c)** Similar observations for the shape (**b**) and shape plus frequency change experiments (**c**). **(d)** The averages of the distributions in **a**-**c**. For the spatial frequency condition (High to low), the average latency was reduced relative to control, likely due to the few saccades with very express latencies. With shape plus frequency changes, it was increased again, but there was (overall) no significant effect of condition on catch-up saccade latency. A Kruskal-Wallis nonparametric ANOVA across all conditions showed no significance (p=0.2617). **(e-h)** Same as **a**-**d** but when landing on a high spatial frequency texture. Here, the catch-up saccade latencies were generally faster than when landing on a low spatial frequency (compare **h** to **d**). Perhaps as a result, it was easier to see the influence of intra-saccadic stimulus changes. Specifically, catch-up saccade latencies increased relative to control (High to high), and there was a significant effect across conditions in **h** (p=0.0049). Thus, elevation of foveal SC activity on intra-saccadic change trials was associated with slightly delayed corrective saccades. This is an expected consequence of prediction error signaling and also of elevated foveal SC activity. Error bars denote SEM. Note that the histograms show fewer numbers of catch-up saccades on intra-saccadic change trials when compared to control trials. This is because there were fewer intra-saccadic change trials than control trials in our experiments (Methods).