Title

Functional prediction of DNA/RNA-binding proteins using deep

learning based on gene expression correlations

Naoki Osato [1*]

[1] School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan.

E-mail: naokiosato11@gmail.com

*Corresponding author.

Abstract

Deep learning methods have been developed to predict gene expression levels

across various cell and tissue types based on DNA-binding sites of protein identified by

ChIP-seq. However, the experimental identification of DNA/RNA-binding sites and the

functional characterization of certain proteins remain challenging. Here I predicted the

regulatory target genes and functional roles of DNA/RNA-binding proteins using gene

co-expression data. Gene co-expression correlations are derived from gene expression

profiles across cells and tissues and represent interactions between a DNA/RNA-binding

protein and other genes. I used these interactions to predict gene expression levels by

partially replacing the input data in the deep learning analysis. Functional annotations

statistically overrepresented in the predicted target genes of ten DNA/RNA-binding

proteins examined in this study were consistent with their known biological roles.

Furthermore, several annotations not present in current gene function databases were

supported by findings in previously published studies.

Keywords

Computational biology, Deep learning, Gene expression, Gene regulation, Protein

function prediction

Introduction

Deep learning approaches have improved the accuracy of predicting gene

expression levels from the binding sites of DNA/RNA-binding proteins, such as

transcription factors, as well as other regulatory factors including miRNAs [1].

Transcription factors bind to promoters near transcription start sites (TSSs) and to

enhancers, which are located distally upstream or downstream of genes. The method

predicts gene expression levels in 53 human tissues based on experimentally determined

binding sites of DNA/RNA-binding proteins in promoter regions and on predicted

binding sites for miRNAs in transcribed regions. The correlation between actual and

predicted gene expression levels was 0.82. In a previous study, I modified the method by

including data on DNA-binding proteins in open chromatin regions within 1 Mbp of

TSSs—some of which likely correspond to enhancers—and by excluding RNA-binding

proteins and miRNAs. This resulted in a correlation coefficient of 0.70 [2].

The binding sites of DNA/RNA-binding proteins are identified using experimental

assays such as ChIP-seq and ChIA-PET. These assays require high-quality antibodies

against the target proteins, which are difficult to obtain for some proteins. For some

DNA/RNA-binding proteins, determining their binding sequences through experimental

analysis may be impractical due to limitations in time, cost, and labor. Large language

model (LLM)-based deep learning architectures, such as Enformer, can predict DNA

sequences that are critical for regulating gene expression [3]. Massively parallel reporter

assays (MPRAs) have been used to measure the regulatory activity of hundreds of

thousands of sequences [4]. However, it remains challenging to determine which DNA- or

RNA-binding proteins interact with these sequences.

Analysis of gene co-expression across different cell and tissue types has suggested

regulatory interactions among genes, including those encoding DNA- or RNA-binding

proteins. Gene expression data from RNA-seq experiments and gene co-expression

analysis results are publicly available in various databases [5-9]. Numerous studies have

reported gene co-expression network analyses to infer functional relationships among

genes [5,9-13]. Gene expression correlations may be influenced by both direct and indirect

molecular interactions. Some of these molecules may be elusive and difficult to detect experimentally, as well as challenging to predict using alternative computational approaches. These correlations may reflect regulatory mechanisms that have yet to be fully characterized.

In this study, to investigate the functional annotations enriched in the putative target genes of DNA- or RNA-binding proteins, I utilized gene co-expression data to identify interactions between these proteins and other genes. Based on gene co-expression data, the DNA-binding sites of approximately 300 proteins that showed low contribution scores in my previous deep learning model were replaced with those of other DNA- or RNA-binding proteins. The deep learning model learned to predict gene expression levels and assigned importance to specific interactions between DNA-binding sites and genes that contribute to the prediction. Regulatory target genes of a transcription factor often include functionally related genes [14]. Therefore, functional enrichment among the predicted target genes can be identified, even if the predictions are not always accurate enough for precise identification of regulatory relationships. I examined the overlap between the predicted regulatory target genes of DNA- or RNA-binding proteins and the binding sites identified by experimental assays such as ChIP-seq and eCLIP. I found that the predicted regulatory target genes whose binding sites were identified by experimental assays for a given DNA- or RNA-binding protein exhibited higher absolute contribution scores than those for other proteins. I then performed functional enrichment analyses on the predicted regulatory target genes of ten DNA- or RNA-binding proteins and found that the enriched functional annotations corresponded to known functions documented in the UniProt database. I also identified functional annotations that were enriched in the predicted regulatory targets but were not annotated in the UniProt database, although they have been described in prior

publications.

These analyses can facilitate the identification of functional roles and regulatory

targets of DNA/RNA-binding proteins without the need for experimental protein-binding

data. They can also reveal both direct and indirect interactions, including those difficult

to detect experimentally, thereby reducing analysis costs and enabling the discovery of

novel functions.

Results

Relationship between gene expression levels and contribution scores

To examine the effect of gene co-expression interactions on gene expression

prediction, the input data for 302 out of 1,310 DNA-binding proteins with low

contribution scores (DeepLIFT scores) were substituted with co-expression interaction

data involving mRNAs encoding DNA-binding proteins and other mRNAs not encoding

DNA- or RNA-binding proteins. DNA-binding proteins encoded by the co-expressed

genes were randomly selected from gene symbols listed in the ENPD database and other

databases used for DNA-binding motif analysis [15]. Notably, the correlation coefficient

between actual and predicted gene expression levels increased from 0.70 to 0.80 in human

foreskin fibroblast (HFF) cells when co-expression data were used (**Fig. 2a, b**). Then,

among the 1,310 DNA-binding proteins, 325 proteins with low contribution scores were

replaced with co-expression interaction data involving genes encoding RNA-binding

proteins and other mRNAs. The RNA-binding proteins were randomly selected from gene

symbols listed in the ENPD database. The correlation coefficient increased from 0.70 to

0.81 when co-expression data were used as well (**Fig. 2c**). These results suggest that gene

co-expression interactions contribute significantly to the prediction of gene expression

levels.

The contribution scores of the input features selected based on gene expression

correlations were distributed across a wide range, but both their mean and median values

were higher than those of the original input (**Fig. 2d and Supplementary Table S1**). I

examined the distribution of Spearman's correlation coefficients between the contribution

scores and gene expression levels ($\log_2$-TPMs; transcripts per million) (**Supplementary**

Fig. S2), and plotted example relationships between the contribution scores of a DNA-

or RNA-binding protein and the expression levels of its predicted regulatory target genes

across cell types (**Fig. 2e**). Based on the interactions between co-expressed genes and

DNA/RNA-binding proteins, the deep learning model adjusted its predictions of gene

expression levels in a cell type–specific manner.

Overlap between high-scoring genes and experimental targets

To investigate whether the regulatory target genes of a DNA/RNA-binding protein,

selected based on their contribution scores, are associated with the binding sites of that

protein, I used ChIP-seq and eCLIP data. Specifically, a gene was considered associated

with the protein if a ChIP-seq peak for a DNA-binding protein was located within ±5 kb

of its transcription start site (TSS), or if an eCLIP peak for an RNA-binding protein was

present in its mRNA-transcribed region. ChIP-seq data for 32 DNA-binding proteins in

HepG2 cells and eCLIP data for 29 RNA-binding proteins in K562 cells were used. Genes

with a DNA- or RNA-binding site for the protein showed high positive or low negative

contribution scores among the predicted regulatory target genes of the DNA/RNA-

binding protein derived from deep learning on gene co-expression data (Fig. 3 and

Supplementary Fig. S1). These high and low contribution scores are likely associated

with the protein's regulatory roles in activating or repressing gene expression, as

described in the documentation of DeepLIFT on its GitHub page and in previous studies

using DeepLIFT [1][16].

Functional enrichment analysis of the target genes of a DNA- or RNA-binding proteins

The transcriptional target genes of a transcription factor often share functionally

related annotations. Functional enrichment analysis of gene annotations can help predict

the function of a transcription factor and verify that the predicted target genes are not

randomly selected. To assess whether the regulatory target genes of a DNA/RNA-binding

protein, predicted by deep learning based on gene expression correlations, include

functionally related genes, I performed functional enrichment analysis using Gene

Ontology (GO) annotations for the target genes of ten DNA/RNA-binding proteins [17,18].

The regulatory target genes of DNA/RNA-binding proteins exhibited functionally related

annotations and significant enrichment in Gene Ontology (GO) terms. However, when

the same number of genes were randomly selected from the entire gene set, GO term

enrichment was negligible, indicating that the predicted target genes are functionally

coherent rather than randomly distributed (**Fig. 4**). For a more detailed analysis of

functional enrichment in the regulatory target genes of DNA/RNA-binding proteins, I

performed a statistical overrepresentation test using the gene list analysis tool provided

by PANTHER (https://pantherdb.org/) [19]. This test utilizes functional annotation

databases, including PANTHER Pathways, Protein Class, GO (Gene Ontology) Slim, GO,

and Reactome Pathways. The functional annotations enriched in the putative regulatory

target genes of a DNA/RNA-binding protein were compared with those of the protein

retrieved   from   the   UniProt   database   (e.g.,

https://www.uniprot.org/uniprotkb/Q9Y2L8/entry) [20]. Ten examples of DNA/RNA-

binding proteins are presented in **Table 1** and **Supplementary Table S4**. The predicted

functional annotations of DNA/RNA-binding proteins correspond to those in the UniProt

database and published studies. Notably, among the ten examples, AKAP8 is annotated

with cell cycle-related functions, such as "GO:0000278 mitotic cell cycle" and

"GO:0044839 cell cycle G2/M phase transition," in the UniProt database

(https://www.uniprot.org/uniprotkb/O43823/entry). In the present study, AKAP8 was

also implicated in circadian clock regulation. A search of the PubMed database identified

two studies supporting the involvement of AKAP8 in circadian rhythms: one states that

"genes such as NFIL3, AKAP8, and SALL3, which are highly correlated with known

molecular clock genes (CRY1), can be identified as candidate circadian genes," while the

other reports that "AKAP5 and AKAP8 are both involved in the biosynthesis of the

circadian hormone melatonin, which is markedly disrupted in individuals with ASDs."

[21,22]. These findings demonstrate that the methods and analyses used in this study

successfully identified both known and novel functional annotations associated with

DNA/RNA-binding proteins.

Deep learning adjusted the contribution (i.e., neural network weights) of

DNA/RNA-binding proteins to the prediction of their regulatory target gene expression

during training. To evaluate the effect of learning on functional enrichment, I analyzed

all co-expressed genes (z score > 2) of ten DNA/RNA-binding proteins using Panther DB.

I then compared the enriched annotations in these co-expressed genes with those of the

regulatory target genes identified by deep learning (**Table 2** and **Supplementary Table**

S5). Although the number of enriched functional annotations increased for all co-

expressed genes due to the larger gene set, some annotations were uniquely or more

significantly enriched in the regulatory target genes of the DNA/RNA-binding protein

after learning. Notably, circadian rhythm-related functional annotations, as seen for

AKAP8, were identified only after learning (**Table 2**; yellow highlights in the

'S2_AKAP8' tab of **Supplementary Table S4**). Similarly, for PKM, kinase-related

functional annotations were detected only after learning (**Table 2** and 'S2_PKM' in

Supplementary Table S4). PKM interacts with ALDOA, as shown in the STRING

database for both mouse (https://string-db.org/network/10090.ENSMUSP00000034834)

and human [23,24]. Previous studies support the association between the functional

annotation "aldolase (PC00044)" and PKM in glycolysis. For example, one study

reported that "Aldolase A (ALDOA) is an enzyme that plays an important role in

glycolysis and gluconeogenesis, which is closely related to tumor metabolism." [25,26].

These functional annotations were enriched after the deep learning process. Thus, they

appear to be involved in biological processes in K562 cells, where the regulatory target

genes of the RNA-binding protein were identified. Circadian clocks and PKM have been

associated with human acute myeloid leukemia (AML) cells through glycolysis [27-30].

These results suggest that combining gene co-expression data with deep learning to

predict gene expression levels can reveal the functional associations of DNA/RNA-

binding proteins in a cell-type- or tissue-specific context.

Discussion

Deep learning models have been used to predict human gene expression levels

across cell types based on the DNA-binding sites of proteins. Here, I applied this approach

to predict novel regulatory interactions between DNA/RNA-binding proteins and their

target genes based on gene expression correlations across cell types. Gene expression

correlations were statistically analyzed using RNA-seq data available from public

databases across various cell and tissue types. Experimental identification of DNA/RNA-

binding sites using assays such as ChIP-seq and eCLIP is more time-consuming, costly,

and labor-intensive than RNA-seq experiments. Experimental determination of

DNA/RNA-binding sites remains challenging for certain DNA/RNA-binding proteins.

Deep learning analysis based on gene expression correlations predicted the regulatory

target genes of DNA/RNA-binding proteins. Functional enrichment analysis of these

genes highlighted potential functional associations and roles of the DNA/RNA-binding

proteins. Many of the predicted annotations were consistent with known functional

annotations reported in public databases. In addition, some predicted annotations not

present in these databases were supported by previously published studies. Functionally

related annotations were commonly observed among the regulatory target genes.

Functional enrichment analysis revealed the potential roles and associations of

DNA/RNA-binding proteins. Many of the predicted functional annotations were

consistent with those reported in public databases. Additionally, annotations not

previously described in these databases were also identified and supported by published

studies.

When comparing DNA-binding sites predicted based on contribution scores for

gene expression prediction with those identified by ChIP-seq assays, the two sets of

binding sites may not completely overlap. Although ChIP-seq can identify tens of

thousands of DNA-binding sites for a given protein, not all of them may be directly

involved in gene expression regulation. In addition, some transcription factors are known

to participate in multiple cellular functions [31,32]. The number of co-expressed genes (z

score > 2) for each DNA/RNA-binding protein was generally less than 1,500, which is

smaller than the number of genes with ChIP-seq peaks for the same proteins. However,

most of the statistically significant co-expressed genes are likely to be involved in gene

expression regulation. In our previous analysis of insulator-associated DNA-binding

proteins, 97 transcription factors and DNA-binding proteins were identified as being

potentially associated with insulator functions that influence enhancer–promoter

interactions [2]. Some of these proteins may play important roles in the formation of

chromatin loops, transcriptional condensates, and/or phase separation—mechanisms that

influence gene expression differently from the direct regulation by transcription factors.

In our deep learning analysis of insulator-associated DNA-binding proteins, we

incorporated eQTL data to identify DNA-binding sites associated with changes in gene

expression levels. However, in this study, I did not use DNA-binding motif sequences or

open chromatin regions identified by DNase-seq or ATAC-seq assays to determine DNA-

binding sites, as the objective was to investigate DNA-binding proteins with unknown

motifs. Moreover, the associations between eSNPs and eGenes in eQTL data may reflect

indirect interactions. The contribution scores of DNA-binding sites selected based on

gene expression correlations exhibited both high and low values, suggesting that indirect

interactions between DNA-binding proteins and other genes may influence these scores.

In the functional enrichment analysis of regulatory target genes, ten DNA/RNA-

binding proteins showed functional annotations consistent with those reported for the

respective proteins, suggesting that the number of predicted target genes was sufficient to

detect statistically significant functional associations. Statistical enrichment was assessed

using Fisher's exact test with false discovery rate (FDR) correction in PANTHER DB,

which generally required more than 100 regulatory target genes to achieve statistical

significance. For some proteins, fewer than 100 predicted target genes were identified,

limiting the detection of enriched functions. A potential next step in this study is to

enhance the detection of functional enrichment in small gene sets by incorporating

additional features, such as gene expression levels and gene co-expression correlations.

Furthermore, functional enrichment analysis using high or low-ranking regulatory target

genes revealed novel functions potentially associated with specific DNA/RNA-binding

proteins. These results suggest that regulatory target genes with high absolute

contribution scores can facilitate the identification of functional roles of DNA/RNA-

binding proteins in a cell type–specific manner.

When all input data in the deep learning analysis were replaced with interactions

between DNA/RNA-binding proteins and other genes based on gene expression

correlations, the number of functionally enriched annotations in the predicted regulatory

target genes decreased, and known functional annotations from the UniProt database were

no longer included. This suggests that a high replacement ratio of input data may result

in learning error-prone interactions between DNA/RNA-binding proteins and target

genes. In a previous study, when the regulatory target genes of a DNA-binding protein

were replaced with randomly selected genes, the number of enriched functional

annotations decreased as the number of random genes increased. The number of enriched

functions, normalized to the total number of predicted genes, reflects the accuracy of the

predicted regulatory target genes [33].

DNA-binding proteins were ranked for each gene based on the contribution scores

of their DNA-binding sites. When a protein's rank was within the top 50 or bottom 50

(i.e., rank ≤ 50 or ≥ 1,260 out of 1,310), functional enrichment of its putative target genes

was typically detectable using statistical tests in PANTHER DB. Some DNA/RNA-

binding proteins did not exhibit high- or low-ranking target genes in HepG2 cells.

However, these proteins may have such rankings in other cell types, allowing functional

enrichment analysis to reveal their roles in different cellular contexts. Some transcription

factors and DNA-binding proteins form protein complexes that regulate gene expression

by binding cooperatively to the same promoters [34-36]. In such cases, not all proteins within

the complex necessarily have consistently high or low ranks. Co-localization analysis of

DNA/RNA-binding proteins may help uncover the functional roles of these complexes.

In conclusion, the deep learning model, its parameters, and analytical framework

developed in this study provide a foundation for predicting the functional roles of

DNA/RNA-binding proteins based on gene expression correlations. Although the present

analysis focused on a limited number of proteins and cell types, the approach

demonstrated its potential for identifying both known and novel functional annotations.

These findings may contribute to a deeper understanding of the biological and molecular

functions of DNA/RNA-binding proteins and their regulatory target genes, especially

those with previously uncharacterized roles.

Methods

305 **Methods**

Deep learning analysis

Deep learning analysis

I modified a previously developed deep learning approach to predict gene

expression levels across different cell and tissue types using DNA-binding site

information of transcription factors and other regulatory proteins. In this study, DNA-

binding sites located in regions far upstream and downstream of transcription start sites

(TSSs) were included in the input data [1,2] (**Fig. 1**). To reduce the effects of cellular

heterogeneity within tissues, gene expression data were collected at the cell-type level.

RNA-seq data from various human cell types were obtained from public databases:

human foreskin fibroblasts (HFFs; GSM941744), human mammary epithelial cells

(HMECs; ENCODE at UCSC Genome Browser,

wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam) [37,38], neural progenitor cells

(NPCs; GSM915326 and ENCFF529SIO), HepG2 hepatoma cells (ENCFF670LIE), and

K562 leukemia cells (ENCFF584WML). FPKM values of gene expression were

calculated using the RSeQC tool (http://rseqc.sourceforge.net/) based on GENCODE v19

gene annotations in the ENCODE portal (https://www.encodeproject.org/). FPKM values

were transformed to $\log_2$ scale with the addition of 0.25 to avoid negative infinity. $\log_2$-

fold changes were calculated relative to the median $\log_2$-FPKM across the four cell types,

following the transcriptome data processing method described by Tasaki et al 2020 [1,2].

DNA-binding sites overlapping open chromatin regions in putative enhancers

located outside of promoter regions were included as input features in the deep learning

model. To identify the DNA-binding sites of DNA-binding proteins (DBPs) in promoter

and enhancer regions, ChIP-seq data for DBPs were obtained from the GTRD database

(release v20.06; https://gtrd.biouml.org) [42]. The binding sites of DBPs regulating the

expression of each transcript were selected based on expression quantitative trait loci

(eQTL) data (GTEx_Analysis_v8_eQTL.tar) from the GTEx portal (release v8;

https://gtexportal.org). A DNA-binding site located within a 100-bp window centered on

a single nucleotide polymorphism (SNP) of an eQTL was considered to regulate the

corresponding transcript. Only cis-eQTLs were used; trans-eQTLs, where the SNP and

the transcript are located on different chromosomes, were excluded. The genomic

coordinates of the eQTLs were converted to the hg19 human genome assembly using the

UCSC LiftOver tool (http://hgdownload.soe.ucsc.edu/downloads.html#liftover) [43].

The original deep learning model used DNA-binding site data in promoter regions

as input features. Specifically, the input was defined as the length of ChIP-seq peaks of a

DNA-binding protein (DBP) within each 100-base-pair bin in the region spanning from

2 kilobases (kb) upstream to 1 kb downstream of the transcription start site (TSS) of each

gene or transcript (**Fig. 1**). To incorporate DNA-binding sites located in more distant

upstream and downstream regions and to reduce the total number of input bins, I first

computed the peak length of a DBP within each 100-bp bin across a ±1 megabase (Mb)

region surrounding the TSS. I then selected the maximum peak length within each

window of 100 consecutive bins. This procedure resulted in up to 200 input values per

transcript, representing the longest ChIP-seq peaks in the extended region. The original

model was modified to accommodate this additional data.

The modified deep learning model was trained using the same test, training, and

validation datasets as in the original method [1,2]: all 2,705 genes located on chromosome

1 were used as the test set; 22,251 genes were randomly selected from other chromosomes

for training, and 2,472 genes were selected for validation. To interpret the model,

DeepLIFT [16] was used to calculate the contribution scores of DNA-binding sites,

indicating their importance in gene expression prediction. Input data representing

promoter and putative enhancer regions were concatenated into a single input matrix

comprising 230 bins: 30 bins for promoter regions and 200 for putative enhancers, each

bin representing the length of ChIP-seq peaks for 1,310 DNA-binding proteins (DBPs).

This concatenation was necessary because the DeepLIFT configuration used

('RevealCancel rule on dense layers and Rescale rule on conv layers', recommended for

genomics) does not support models with multiple input layers. Model training and

inference were performed on a system running Ubuntu 20.04 with Python 3.8.10,

TensorFlow 2.6.0, the NVIDIA CUDA toolkit 11.4.2, a GeForce RTX 3080 GPU (10 GB

memory), and an Intel Core i7-10700 CPU with 128 GB RAM. Original scripts were

updated to support the newer software versions. The training procedure followed the

original method, and training time ranged from two to four hours.

Functional prediction of DNA/RNA-binding proteins

To predict the regulatory target genes of DNA/RNA-binding proteins without

relying on protein binding assay data such as ChIP-seq or eCLIP, I modified the deep

learning input by substituting low-contribution DNA-binding proteins with other proteins.

Among the 1,310 DNA-binding proteins originally used in the model, 302 proteins with

lower contribution scores for predicting gene expression levels were replaced with other

DNA-binding proteins. For RNA-binding protein analysis, 325 DNA-binding proteins

were replaced with RNA-binding proteins. Gene symbols of RNA-binding proteins were

obtained from the ENPD database [15]. Interactions between the substituted DNA/RNA-

binding proteins and their putative target genes were estimated based on gene co-

expression pairs across different cells and tissues, as obtained from COXPRESdb [5].

Gene co-expression refers to the similarity in expression profiles between pairs of

genes. Functionally related genes, as well as genes under common regulatory control, are

generally co-expressed across different cell and tissue types. When a gene encoding a

DNA/RNA-binding protein exhibits expression patterns similar to those of other genes,

those genes are likely to be regulated by the protein and functionally associated with it.

Co-expression interactions with a z-score > 2 between genes encoding DNA/RNA-

binding proteins and other genes (excluding DNA/RNA-binding proteins) were used to

replace those with lower contribution scores in the input data. The binding sites of the

DNA/RNA-binding proteins were hypothetically assigned to 2 kb upstream of the

transcription start sites (TSSs) and at the TSSs of their co-expressed genes, respectively.

Using this updated input, deep learning analysis and the DeepLIFT tool were applied to

calculate the contribution scores of the binding sites of DNA/RNA-binding proteins for

predicting gene expression levels.

For each DNA/RNA-binding protein, I evaluated the rank of its contribution score

by comparing the contribution scores of its binding sites with those of other DNA/RNA-

binding proteins for each gene. Based on the tendency of high or low ranks (i.e., high

positive or low negative contribution scores) across the predicted regulatory target genes,

I selected genes in which the protein's binding site ranked within the top five or bottom

five as its primary regulatory target genes. To detect functional enrichment, the rank

threshold was adjusted individually for each DNA/RNA-binding protein. In most cases,

the threshold was limited to the top five or bottom five ranks. For RPLP0, the threshold

was extended to the top eight to increase the number of selected genes.

Functional enrichment analysis of the primary regulatory target genes of each

DNA/RNA-binding protein was performed using in-house scripts and the Statistical

Overrepresentation Test provided by PANTHER (https://pantherdb.org/) [19]. To evaluate

the specificity of enriched Gene Ontology (GO) annotations, I compared the number of

significantly enriched GO terms among three gene sets: (1) the primary target genes of

each DNA/RNA-binding protein, (2) all co-expressed genes with a z-score > 2, and (3)

randomly selected genes matched in number to the co-expressed genes. GO annotations

from the three major categories—biological process, molecular function, and cellular

component—were combined for the analysis. The GO ontology file

(https://purl.obolibrary.org/obo/go.obo) and the human gene annotation file

(https://current.geneontology.org/annotations/goa_human.gaf.gz) were downloaded from

the Gene Ontology resource (https://geneontology.org/docs/download-ontology/) [17,18].

Human gene symbols were converted to Ensembl Gene IDs using the HUGO Gene

Nomenclature Committee (HGNC) database (https://www.genenames.org/). For

PANTHER analysis, Ensembl Gene IDs of the primary target genes were used as the test

gene set, and Ensembl Gene IDs from the full gene expression dataset used for deep

learning were used as the reference set. Functional enrichment was assessed using

Fisher's exact test, with multiple testing correction applied using the Benjamini–

Hochberg false discovery rate (FDR), with significance defined as FDR < 0.05.

To investigate the overlap between experimentally identified binding sites and the

predicted regulatory target genes of DNA/RNA-binding proteins derived from deep

learning on gene co-expression data, I selected DNA-binding sites located within ±5 kb

of a gene's transcription start site (TSS) based on ChIP-seq data, and RNA-binding sites

located between the TSS and transcription end site (TES) based on eCLIP data. Gene

annotations were based on the human GENCODE v19 release [40]. For this analysis, ChIP-

seq data for 32 DNA-binding proteins and eCLIP data for 29 RNA-binding proteins were

obtained from the ENCODE portal and a published study (**Supplementary Table S6**) [44].

I also evaluated the proportions of genes with high or low contribution scores, as defined

above, among those with ChIP-seq or eCLIP binding sites near the TSS or transcribed

regions, respectively.

To examine the binding sites of ChIP-seq data and eCLIP data around the

regulatory target genes of DNA/RNA-binding proteins predicted by deep learning of gene

co-expression data, I selected the DNA-binding sites within ±5 kbp of a TSS of a gene by

ChIP-seq assay or the RNA-binding sites between the transcription start and end of a gene

by eCLIP assay based on the Human GENCODE v19 gene annotation [40]. For this analysis,

ChIP-seq data of 32 DNA-binding proteins and eCLIP data of 29 RNA-binding proteins

were obtained from the ENCODE portal and the paper (**Supplementary Table S6**) [44]. I

also examined the ratios of the number of genes with high and low contribution scores of

the DNA/RNA-binding site around genes with the DNA/RNA-binding site by ChIP-seq

or eCLIP assays.

Statistical analysis

Functional enrichment was assessed using Fisher's exact test, with multiple testing

correction applied using the Benjamini–Hochberg false discovery rate (FDR) procedure.

Statistical significance was defined as FDR < 0.05. Distributions of DeepLIFT

contribution scores for DNA/RNA-binding sites derived from gene co-expression data

and ChIP-seq data were compared using the Mann-Whitney U test (also known as the

Wilcoxon rank sum test). A P value < 0.05 was considered statistically significant.

Data availability

RNA-seq data of human foreskin fibroblast (HFF) cells were obtained from the Gene

Expression Omnibus (GEO) under accession number GSM941744. RNA-seq data of

human mammary epithelial cells (HMEC) were retrieved from ENCODE at the UCSC

Genome Browser (http://genome.ucsc.edu/encode/) under the file name

wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam. RNA-seq data of human neural

progenitor cells (NPC) were obtained from GSM915326. RNA-seq data for HepG2 and

K562 cells were obtained from the ENCODE portal under accession numbers

ENCFF670LIE and ENCFF584WML, respectively. Co-expression data of mRNAs were

downloaded from COXPRESdb (https://zenodo.org/record/6861444). The list of

DNA/RNA-binding proteins was obtained from the ENPD database

(http://qinlab.sls.cuhk.edu.hk/ENPD/). ChIP-seq data of transcription factors were

downloaded from the v20.06 release of the GTRD database (https://gtrd.biouml.org).

eQTL data (GTEx_Analysis_v8_eQTL.tar) were obtained from the GTEx portal (v8

release) (https://gtexportal.org). eCLIP data of RNA-binding proteins were retrieved from

ENCSR456FVU in the ENCODE portal and from Supplementary Data 2 of Van Nostrand

et al., *Nature* (2020), https://doi.org/10.1038/s41586-020-2077-3.

Code availability

The deep learning model used in this study is available at:

https://github.com/reposit2/coexgene .

Figures

Figure 1. Overview of gene expression prediction. Gene expression levels in human

cells were predicted from the genomic locations of DNA-binding sites at promoters and

distal regions. During model training, a DNA-binding protein bound to a distal region

overlapping an expression-associated SNP (eSNP) was linked to a potentially regulated

gene (eGene) based on eQTL data. For each gene–protein association, ChIP-seq peak

locations were binned (green box) and summarized into a matrix as input for deep

learning.

Figure 2. Relationship between gene expression levels and contribution scores.

Predicted gene expression levels (log$_2$-TPM) were compared with actual values in HFF

cells using Spearman's rank correlation. **a** Input data included DNA-binding sites of 1,310

DNA-binding proteins obtained from ChIP-seq data. **b** Among the 1,310 proteins, 302

with low contribution scores were replaced with an equal number of DNA-binding

proteins selected based on gene co-expression data. **c** Similarly, 325 low-contribution

DNA-binding proteins were replaced with RNA-binding proteins selected based on gene

co-expression data. **d** DNA-binding sites based on co-expression data showed higher

DeepLIFT contribution scores. A and C: Binding sites of DNA-binding proteins derived

from ChIP-seq data. B and D: Binding sites derived from co-expression data. A and B:

Positive contribution scores. C and D: Absolute values of negative contribution scores

(Wilcoxon signed-rank test, $P < 2.2 \times 10^{-16}$). **e** Example of the relationship between the

contribution scores of a regulator and its expression levels across tissues. A linear

 regression line with 95% confidence intervals is shown.

Figure 3. Comparison of predicted target genes with experimental data.

The x-axis represents the rank of contribution scores of DNA/RNA-binding sites used to predict gene expression levels. Scores were ranked from 1 (highest positive contribution, left) to 1,310 (lowest negative contribution, right) and binned into 50-rank intervals. Rank 1 corresponds to ranks 1–50, rank 2 to ranks 51–100, and rank 26 to ranks 1,251–1,310. DNA/RNA-binding sites were categorized as being located in the promoters or transcribed regions of genes experimentally bound by the DNA/RNA-binding protein shown in each graph. These sites showed higher or lower contribution scores than those across all ChIP-seq binding sites. The y-axis indicates the number of sites within each 50-rank bin. Positive contribution scores are associated with gene activation, while negative scores correspond to gene repression.

Figure 4. Number of ov ... **ation terms in the target genes of DNA/RNA-binding** ...

Functional enrichment analysis was performed on (1) co-expressed genes contributing to gene expression prediction in the deep learning model, (2) all co-expressed genes with a z-score > 2, and (3) randomly selected genes matched in number to the total co-expressed genes. Gene Ontology (GO) functional annotations from the three categories—biological process, molecular function, and cellular component—were combined for the analysis. *P*-values from Fisher's exact test were adjusted using the false discovery rate (FDR) method. The target genes of each DNA/RNA-binding protein were found to include functionally associated genes.

Tables

Table 1. Common functional annotations between DNA/RNA-binding proteins and

525 **their putative regulatory target genes.**

526 Functional annotations of DNA/RNA-binding proteins were obtained from the UniProt

527 database. Annotations significantly enriched in their putative regulatory target genes were

528 identified using the PANTHER database. Gray lines separate individual annotation

529 entries.

ZKSCAN5
DNA-binding transcription factor activity
C2H2 zinc finger transcription factor
RNA polymerase II cis-regulatory region
sequence-specific DNA binding
metal ion binding

ZBTB34
C2H2 zinc finger transcription factor
DNA-binding transcription activator activity,
RNA polymerase II-specific
RNA polymerase II transcription regulatory
region sequence-specific DNA binding

ZNF557
DNA-binding transcription activator activity,
RNA polymerase II-specific
RNA polymerase II transcription regulatory
region sequence-specific DNA binding
ion binding, metal ion binding

ZNF446
C2H2 zinc finger transcription factor
extracellular space

ZC3H4
chromatin/chromatin-binding, or -regulatory
protein
transcription regulator activity
531 metal ion binding

POLR2G
RNA polymerase II transcription regulatory
region sequence-specific DNA binding
general transcription factor
translational protein

RPLP0
large ribosomal subunit
ribonucleoprotein complex
cytosolic large ribosomal subunit

SPEN
mRNA binding
chromatin binding
transcription coregulator activity

PKM
kinase
glycolytic process
ATP metabolic process

AKAP8
negative regulation of RNA biosynthetic
process
Circadian clock system
Cry-Per complex

Table 2. Functional enrichments detected in the regulatory target genes of RNA-

binding proteins.

Functional enrichment analysis was performed using the PANTHER database. This table

lists functional annotations that were statistically enriched in the regulatory target genes

of RNA-binding proteins after deep learning based on gene co-expression data, but not

enriched in the complete set of co-expressed genes (z-score > 2) including the RNA-

binding protein. 'Reference' indicates the total number of human genes. 'Analyzed' refers

to the number of regulatory target genes analyzed. 'Expected' is the number of genes with

the annotation expected by chance, based on its frequency in the reference set.

'Over/Under' indicates overrepresentation or underrepresentation. 'Fold Enrichment' is the

ratio of observed to expected counts. 'Raw P-value' represents the P-value from Fisher's

exact test, and 'FDR' indicates the Benjamini–Hochberg false discovery rate.

AKAP8							

PANTHER Pathways

PANTHER Pathways	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
Circadian clock system (P00015)	10	3	0.03	+	> 100	2.60E-06	4.16E-04

PANTHER GO-Slim Biological Process

PANTHER GO-Slim Biological Process	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
circadian regulation of gene expression (GO:0032922)	16	3	0.05	+	65.55	1.20E-05	1.27E-02
response to light stimulus (GO:0009416)	21	3	0.06	+	49.94	2.82E-05	1.99E-02
circadian rhythm (GO:0007623)	21	3	0.06	+	49.94	2.82E-05	1.49E-02
rhythmic process (GO:0048511)	24	3	0.07	+	43.7	4.27E-05	1.81E-02

GO cellular component complete

GO cellular component complete	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
Cry-Per complex (GO:1990512)	3	2	0.01	+	> 100	2.40E-05	4.77E-02

PKM

PANTHER Protein Class

PANTHER Protein Class	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
aldolase (PC00044)	5	2	0.1	+	21.02	3.48E-03	5.68E-02
kinase (PC00137)	134	9	2.55	+	3.53	1.07E-03	2.62E-02

PANTHER GO-Slim Molecular Function

PANTHER GO-Slim Molecular Function	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
enzyme inhibitor activity (GO:0004857)	107	8	2.04	+	3.93	1.01E-03	4.68E-02
kinase activity (GO:0016301)	562	23	10.7	+	2.15	7.08E-04	3.57E-02

PANTHER GO-Slim Biological Process

PANTHER GO-Slim Biological Process	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
endoplasmic reticulum tubular network organization (GO:0071786)	12	3	0.23	+	13.14	1.32E-03	3.68E-02
regulation of cellular component biogenesis (GO:0044087)	139	10	2.65	+	3.78	3.31E-04	1.05E-02

Acknowledgements

Supercomputing resources were provided by the Human Genome Center, the Institute of Medical Science, the University of Tokyo, and the ROIS National Institute of Genetics.

Author contributions

N.O. designed and conducted the research, and wrote and revised the manuscript.

Supplementary data

Supplementary data are available at

Conflict of interests

None declared.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI [16K00387], a start-up research grant from Osaka University, the KIOXIA Young Scientist Research Grant at Waseda University, and the Grant for Basic Science Research Projects from the Sumitomo Foundation. All grants were awarded to N.O.

References

1   Tasaki, S., Gaiteri, C., Mostafavi, S. & Wang, Y. Deep learning decodes the principles of differential gene expression. *Nat Mach Intell* **2**, 376-386, doi:10.1038/s42256-020-0201-6 (2020).

2   Osato, N. & Hamada, M. Systematic discovery of directional regulatory motifs associated with human insulator sites. *bioRxiv* (2025).

3   Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-range interactions. *Nature methods* **18**, 1196-1203, doi:10.1038/s41592-021-01252-x (2021).

4   Agarwal, V. *et al.* Massively parallel characterization of transcriptional regulatory elements. *Nature*, doi:10.1038/s41586-024-08430-9 (2025).

5   Obayashi, T., Kodate, S., Hibara, H., Kagaya, Y. & Kinoshita, K. COXPRESdb v8: an animal gene coexpression database navigating from a global view to detailed investigations.

Nucleic acids research **51**, D80-D87, doi:10.1093/nar/gkac983 (2023).

6   Yang, S. *et al.* COEXPEDIA: exploring biomedical hypotheses via co-expressions
associated with medical subject headings (MeSH). *Nucleic acids research* **45**, D389-D396,
doi:10.1093/nar/gkw868 (2017).

7   Zhu, Q. *et al.* Targeted exploration and analysis of large cross-platform human
transcriptomic compendia. *Nature methods* **12**, 211-214, 213 p following 214,
doi:10.1038/nmeth.3249 (2015).

8   Miller, H. E. & Bishop, A. J. R. Correlation AnalyzeR: functional predictions from gene
co-expression correlations. *BMC Bioinformatics* **22**, 206, doi:10.1186/s12859-021-
04130-7 (2021).

9   Raina, P. *et al.* GeneFriends: gene co-expression databases and tools for humans and
model organisms. *Nucleic acids research* **51**, D145-D158, doi:10.1093/nar/gkac1031
(2023).

10   Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global
discovery of conserved genetic modules. *Science (New York, N.Y.)* **302**, 249-255,
doi:10.1126/science.1087447 (2003).

11   Lemoine, G. G., Scott-Boyer, M. P., Ambroise, B., Perin, O. & Droit, A. GWENA: gene
co-expression networks analysis and extended modules characterization in a single
Bioconductor package. *BMC Bioinformatics* **22**, 267, doi:10.1186/s12859-021-04179-4
(2021).

12   Warde-Farley, D. *et al.* The GeneMANIA prediction server: biological network integration
for gene prioritization and predicting gene function. *Nucleic acids research* **38**, W214-220,
doi:10.1093/nar/gkq537 (2010).

13   Wong, A. K., Krishnan, A. & Troyanskaya, O. G. GIANT 2.0: genome-scale integrated
analysis of gene networks in tissues. *Nucleic acids research* **46**, W65-W70,
doi:10.1093/nar/gky408 (2018).

14   McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory regions.
Nature biotechnology **28**, 495-501, doi:10.1038/nbt.1630 (2010).

15   Tak Leung, R. W., Jiang, X., Chu, K. H. & Qin, J. ENPD - A Database of Eukaryotic
Nucleic Acid Binding Proteins: Linking Gene Regulations to Proteins. *Nucleic acids*
research **47**, D322-D329, doi:10.1093/nar/gky1112 (2019).

16   Shrikumar, A., Greenside, P. & Kundaje, A. in *Proceedings of the 34th International*
Conference on Machine Learning Vol. 70 (eds Precup Doina & Teh Yee Whye) 3145--
3153 (PMLR, Proceedings of Machine Learning Research, 2017).

17   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nature genetics* **25**, 25-29, doi:10.1038/75556 (2000).

18   Gene Ontology, C. *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**,
doi:10.1093/genetics/iyad031 (2023).

19   Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all.
Protein Sci **31**, 8-22, doi:10.1002/pro.4218 (2022).

20   UniProt, C. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic acids*
research **53**, D609-D617, doi:10.1093/nar/gkae1010 (2025).

21   Tan, X. *et al.* Novel Insights into the Circadian Rhythms Based on Long Noncoding and
Circular RNA Profiling. *Int J Mol Sci* **25**, doi:10.3390/ijms25021161 (2024).

22   Poelmans, G., Franke, B., Pauls, D. L., Glennon, J. C. & Buitelaar, J. K. AKAPs integrate
genetic findings for autism spectrum disorders. *Transl Psychiatry* **3**, e270,
doi:10.1038/tp.2013.48 (2013).

23   Snel, B., Lehmann, G., Bork, P. & Huynen, M. A. STRING: a web-server to retrieve and
display the repeatedly occurring neighbourhood of a gene. *Nucleic acids research* **28**,
3442-3444, doi:10.1093/nar/28.18.3442 (2000).

24   Szklarczyk, D. *et al.* The STRING database in 2023: protein-protein association networks
and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids*
research **51**, D638-D646, doi:10.1093/nar/gkac1000 (2023).

25   Tian, W. *et al.* Bioinformatics analysis of the role of aldolase A in tumor prognosis and
immunity. *Sci Rep* **12**, 11632, doi:10.1038/s41598-022-15866-4 (2022).

26   Zhang, F. *et al.* Elevated transcriptional levels of aldolase A (ALDOA) associates with cell
cycle-related genes in patients with NSCLC and several solid tumors. *BioData Min* **10**, 6,
doi:10.1186/s13040-016-0122-4 (2017).

27   Zlacka, J. & Zeman, M. Glycolysis under Circadian Control. *Int J Mol Sci* **22**,
doi:10.3390/ijms222413666 (2021).

28   Yang, Y., Pu, J. & Yang, Y. Glycolysis and chemoresistance in acute myeloid leukemia.
Heliyon **10**, e35721, doi:10.1016/j.heliyon.2024.e35721 (2024).

29   Puram, R. V. *et al.* Core Circadian Clock Genes Regulate Leukemia Stem Cells in AML.
Cell **165**, 303-316, doi:10.1016/j.cell.2016.03.015 (2016).

30   Lee, Y. Roles of circadian clocks in cancer pathogenesis and treatment. *Exp Mol Med* **53**,
1529-1538, doi:10.1038/s12276-021-00681-0 (2021).

31   Ladomery, M. Multifunctional proteins suggest connections between transcriptional and
post-transcriptional processes. *Bioessays* **19**, 903-909, doi:10.1002/bies.950191010
(1997).

32   Kim, S., Yu, N. K. & Kaang, B. K. CTCF as a multifunctional protein in genome regulation
and gene expression. *Exp Mol Med* **47**, e166, doi:10.1038/emm.2015.33 (2015).

33   Osato, N. Characteristics of functional enrichment and gene expression level of human

655   putative transcriptional target genes. *BMC Genomics* **19**, 957, doi:10.1186/s12864-017-
656   4339-5 (2018).

657   34   Burley, S. K. & Kamada, K. Transcription factor complexes. *Curr Opin Struct Biol* **12**, 225-
658   230, doi:10.1016/s0959-440x(02)00314-7 (2002).

659   35   Goos, H. *et al.* Human transcription factor protein interaction networks. *Nat Commun* **13**,
660   766, doi:10.1038/s41467-022-28341-5 (2022).

661   36   Berenson, A. *et al.* Paired yeast one-hybrid assays to detect DNA-binding cooperativity
662   and antagonism across transcription factors. *Nat Commun* **14**, 6570, doi:10.1038/s41467-
663   023-42445-6 (2023).

664   37   Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic*
665   *acids research* **41**, D991-995, doi:10.1093/nar/gks1193 (2013).

666   38   Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene
667   expression and hybridization array data repository. *Nucleic acids research* **30**, 207-210,
668   doi:10.1093/nar/30.1.207 (2002).

669   39   Luo, Y. *et al.* New developments on the Encyclopedia of DNA Elements (ENCODE) data
670   portal. *Nucleic acids research* **48**, D882-D889, doi:10.1093/nar/gkz1062 (2020).

671   40   Mudge, J. M. *et al.* GENCODE 2025: reference gene annotation for human and mouse.
672   *Nucleic acids research* **53**, D966-D975, doi:10.1093/nar/gkae1078 (2025).

673   41   Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments.
674   *Bioinformatics (Oxford, England)* **28**, 2184-2185, doi:10.1093/bioinformatics/bts356
675   (2012).

676   42   Kolmykov, S. *et al.* GTRD: an integrated view of transcription regulation. *Nucleic acids*
677   *research* **49**, D104-D111, doi:10.1093/nar/gkaa1057 (2021).

678   43   Hinrichs, A. S. *et al.* The UCSC Genome Browser Database: update 2006. *Nucleic acids*
679   *research* **34**, D590-598, doi:10.1093/nar/gkj144 (2006).

680   44   Van Nostrand, E. L. *et al.* A large-scale binding and functional map of human RNA-
681   binding proteins. *Nature* **583**, 711-719, doi:10.1038/s41586-020-2077-3 (2020).