Title

Pathway redistribution reveals a shared signaling backbone and context-dependent regulatory modules in RNA-binding protein networks

Naoki Osato [1*] and Kengo Sato [1]

[1] School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan.

E-mail: naokiosato11@gmail.com

*Corresponding author.

Abstract

Understanding how regulatory architectures are reorganized across cellular contexts remains a central challenge in functional genomics. Here, we integrate co-expression–derived candidate regulatory interactions with interpretable deep learning to generate gene-level contribution scores and introduce ΔNES (normalized enrichment score difference) to quantify pathway redistribution between cellular states. Because gene expression reflects the combined effects of multiple regulatory inputs, contribution scores capture relative regulatory influence rather than transcriptional abundance itself.

Applying this framework to neural progenitor cells and K562 leukemia cells, we identify systematic redistribution of functional modules across multiple RNA-binding proteins, including PKM, HNRNPK, and NELFE. Neural System– and Immune System–associated modules are differentially positioned along the ΔNES spectrum, indicating context-dependent redistribution of regulatory influence rather than isolated pathway activation events.

At the pathway level, Signal Transduction consistently forms a shared signaling backbone across proteins and cellular contexts, while modules related to neuronal functions, immune responses, and developmental processes exhibit context-dependent redistribution. Subpathway analysis further reveals convergence on receptor-mediated signaling processes, including FGFR/RTK-, IRS-, and MAPK-related pathways.

These redistribution patterns are preserved under alternative DeepLIFT background settings despite polarity changes in contribution–expression correlations, indicating that pathway-level contrasts arise from stable rank-structure differences rather than background-dependent score artifacts.

Together, our findings demonstrate that contribution score–based pathway

ranking reveals a conserved signaling backbone alongside context-dependent functional

modules, providing a framework for interpreting regulatory architecture beyond

expression-centric analyses.

Keywords

Computational biology, Deep learning, Regulation of gene expression, DNA/RNA-

binding proteins, Protein function prediction, ChatGPT, GSEA

Introduction

Understanding how regulatory architectures are organized and reconfigured

across cellular contexts remains a central challenge in molecular biology [1-10]. Nucleic

acid-binding proteins (NABPs), including DNA-binding proteins and RNA-binding

proteins (RBPs), play key roles in gene regulation; however, their functional

characterization across diverse cell types remains incomplete. While experimental

approaches such as ChIP-seq and eCLIP can identify binding sites, these methods are not

easily scalable and often fail to capture indirect or context-dependent regulatory

interactions [11], particularly for RBPs lacking well-defined binding motifs [12-14].

Computational approaches, including deep learning models, have improved the

prediction of gene expression by integrating regulatory features derived from binding data.

However, these methods typically rely on experimentally defined binding sites or

sequence features and do not directly reveal how individual NABPs contribute to gene

expression programs at the systems level. In particular, it remains unclear how regulatory

influence is organized across pathways and how it is reconfigured between cellular

contexts.

Gene co-expression analysis provides a complementary perspective by capturing

coordinated expression patterns across diverse biological conditions, reflecting both

direct and indirect regulatory relationships [15-23]. Integrating such information into

interpretable deep learning frameworks offers the potential to quantify regulatory

influence at the gene level without requiring prior knowledge of binding motifs or direct

interaction data.

Here, we integrate co-expression–derived regulatory interactions with

interpretable deep learning to compute gene-level contribution scores and introduce

ΔNES (normalized enrichment score difference) to quantify pathway redistribution

between cellular contexts. Rather than focusing on individual regulatory interactions, this

framework enables the analysis of regulatory architecture at the pathway level.

Applying this approach to neural progenitor cells and K562 leukemia cells, we

find that regulatory influence is not organized as distinct pathway sets, but instead is

structured around a shared signaling backbone, while functional modules are redistributed

in a context-dependent manner. This conceptual framework provides a systems-level

perspective on gene regulation, revealing how common signaling networks are

differentially utilized across cellular states.

Results

Gene co-expression features enhance expression prediction accuracy

Replacing low-contribution DBP inputs with gene co-expression data led to a

substantial increase in model performance   (**Fig. 1**). In human foreskin fibroblasts

(HFFs), substituting 302 out of 1,310 DBPs with co-expression profiles improved the

correlation between predicted and actual gene expression from 0.70 to 0.80 (**Fig. 2a, b**).

These co-expression inputs included both NABP-encoding and non-NABP mRNAs, with

gene symbols for DNA- and RNA-binding proteins obtained from the ENPD (Eukaryotic

Nucleic acid Binding Protein) database[24], and co-expression profiles sourced from

COXPRESdb [15]. A subsequent experiment replaced 325 low-contribution DBPs with

co-expression interactions involving RBPs, resulting in a further increase in prediction

accuracy ($r = 0.81$; **Fig. 2c**). Collectively, these findings demonstrate that integrating co-

expression interactions into the model enhances its ability to capture gene regulatory

dynamics beyond direct binding site information.

Contribution scores derived from co-expression–based NABP–gene interactions

exhibited broader distributions and higher overall magnitudes than those obtained from

ChIP-seq–based inputs across all four cell types examined (Wilcoxon signed-rank test, p

$< 10^{-2}$; **Fig. 2d** and **Supplementary Fig. S2c**; **Supplementary Table S1**). Each pair of

violin plots represents absolute contribution scores for co-expression–based sites (right

plot in each pair) and the corresponding ChIP-seq sites not included in that set (left plot).

A–P correspond to specific cell types and NABP classes as follows: **B**, DBP co-

expression–based sites in neural progenitor cells (NPCs); **A**, NPC ChIP-seq sites

excluding **B**. **D**, RBP co-expression–based sites in NPCs; **C**, NPC ChIP-seq sites

excluding **D**. **F**, DBP co-expression–based sites in HepG2 cells; **E**, HepG2 ChIP-seq sites

excluding **F**. **H**, RBP co-expression–based sites in K562 cells; **G**, K562 ChIP-seq sites

excluding **H**. **J**, DBP co-expression–based sites in HMEC cells; **I,** HMEC ChIP-seq sites

excluding **J**. **L**, RBP co-expression–based sites in HMEC cells; **K**, HMEC ChIP-seq sites

excluding **L**. **N**, DBP co-expression–based sites in human foreskin fibroblasts (HFFs); **M**,

HFF ChIP-seq sites excluding **N**. **P**, RBP co-expression–based sites in HFFs; **O**, HFF

ChIP-seq sites excluding **P**. Corresponding violin plots generated using an amplified

background (input × 2) are shown in **Supplementary Figure S2 c**. However, the nature

of this difference depended on the class of nucleic acid–binding protein. For DNA-

binding proteins (DBPs), contribution score distributions differed significantly between

ChIP-seq–derived and co-expression–derived regulatory sites, yet their mean and median

values were often comparable across cell types. This indicates that, for DBPs, co-

expression–based feature selection primarily reshapes the distributional structure of

contribution scores without uniformly shifting their central tendency. In contrast, for

RNA-binding proteins (RBPs), regulatory sites inferred from gene co-expression

consistently showed substantially higher DeepLIFT contribution scores than those

derived from experimental binding data alone. This pattern was observed across all

examined cell types, including K562, NPCs, HMECs, and HFFs, and was reflected by

marked increases in both mean and median contribution scores. These results suggest that

co-expression–based inference preferentially enriches for RBP-associated regulatory

interactions that contribute more directly to gene expression prediction, whereas

experimental binding data for RBPs may include a larger fraction of interactions that are

weakly coupled to transcriptomic output. These observations are consistent with

previous reports showing that incorporating RNA-binding protein– and miRNA-

associated features improves gene expression prediction accuracy, as exemplified by the

DEcode framework[25]. Importantly, these trends were robust to DeepLIFT background

definition: comparable differences between ChIP-seq– and co-expression–derived targets

were observed under both background scaling schemes (input $\times$ 0.5 and input $\times$ 2). Thus,

while background scaling influences the sign and absolute scale of contribution scores, it

does not alter the fundamental distinction between binding-based and co-expression–

based regulatory inference.

To assess the influence of background selection on DeepLIFT-derived contribution

scores, we compared results obtained using a reduced background (input $\times$ 0.5; **Fig. 2e**)

and an amplified background (input $\times$ 2; **Fig. 2f**) for the same representative DNA- and

RNA-binding proteins. Under the reduced background, contribution scores tended to

show negative correlations with gene expression, whereas the amplified background

produced predominantly positive correlations. These patterns appeared approximately

mirrored around zero, consistent with DeepLIFT's definition as a relative contribution

measure with respect to the chosen reference. Importantly, despite this inversion in sign,

the relative ranking of genes and the overall cell type–specific structure of contribution

score distributions remained highly consistent across background settings. In both cases,

genes with larger-magnitude contribution scores were preferentially associated with

higher or lower expression levels depending on the background, indicating that

DeepLIFT captures relative regulatory importance rather than absolute activation or

repression. A conceptual model explaining how input definition and background scaling

give rise to these patterns is provided in **Supplementary Fig. S4**. Notably, background-

dependent differences were more pronounced for RNA-binding proteins (RBPs) (**Fig.**

2e,f). For the same representative RBPs, scatter plots generated using the reduced

background (input $\times$ 0.5) exhibited more distinct, partially separated distributions across

the four cell types, whereas under the amplified background (input $\times$ 2), data points from

different cell types overlapped more extensively. This suggests that the reduced-

background configuration enhances sensitivity to cell type–specific regulatory variation

for RBPs. In contrast, DNA-binding proteins (DBPs) generally showed stronger

correlations with gene expression under both background conditions, consistent with their

more direct transcriptional regulatory roles. Together, these results underscore the

importance of background-aware and protein class–specific interpretation of DeepLIFT

contribution scores.

Experimental binding data support model-inferred regulatory targets

To determine whether DeepLIFT-derived regulatory contribution scores reflect

biologically meaningful nucleic acid–binding protein (NABP) activity, we systematically

compared model-inferred regulatory targets with experimentally validated ChIP-seq and

eCLIP binding data across multiple genomic regions, threshold definitions, and cellular

contexts (**Fig. 3**). Experimentally validated binding sites are consistent with deep

learning–derived regulatory targets. Genes with ChIP-seq or eCLIP peaks for a given

NABP exhibited disproportionately high or low importance ranks, indicating that

experimentally confirmed binding sites are associated with stronger regulatory influence

in gene expression prediction models. This relationship supports the biological validity

of DeepLIFT-based contribution scores, which quantify the importance of specific

NABPs in driving gene expression changes.

To evaluate this association, we analyzed ChIP-seq data for 32 DBPs in HepG2

cells and eCLIP data for 29 RBPs in K562 cells—two cell lines with the most

comprehensive publicly available NABP binding site datasets from ENCODE and related

studies. A gene was considered associated with an NABP if a ChIP-seq peak was located

within ±5 kb of its transcription start site (TSS) or if an eCLIP peak was mapped to its

transcribed region. As illustrated in **Fig. 3a** and **Supplementary Fig. S1**, NABPs with

experimentally validated binding sites consistently showed higher absolute contribution

score values across their predicted target genes, reinforcing their regulatory relevance and

corresponding with prior interpretations of DeepLIFT scores[25,26].

Model-derived importance scores reliably predict DBP binding sites, as evidenced

by significant overlap with ChIP-seq peaks (**Fig. 3b, Supplementary Table S2** and **S3,**

Supplementary Fig. S3a). To evaluate this correspondence, we analyzed ChIP-seq data

for 32 DNA-binding proteins (DBPs) and quantified how frequently high-scoring gene

regions—defined by DeepLIFT contribution scores—coincided with experimental

binding evidence. Overlap ratios were calculated as the proportion of high-contribution

regions that intersected with ChIP-seq peaks, considering both promoter-proximal (±5

kb) and distal (±100 kb) regions around transcription start sites (TSSs). To account for

the functional heterogeneity and abundance of ChIP-seq peaks, we applied a precision-

at-K approach, focusing on the top-ranked or top-scoring gene regions most likely to

reflect regulatory relevance. Four thresholds were used to define high-contribution

regions: two based on contribution score rank (top/bottom 20 or 50) and two based on

absolute score magnitude (≥ 0.01 or ≤ -0.01; ≥ 0.001 or ≤ -0.001). A control analysis

using randomly selected gene regions of matched size was conducted in parallel. In all

tested conditions, the overlap between ChIP-seq peaks and high-contribution regions

exceeded that of the random control, supporting the model's ability to prioritize

functionally relevant regulatory sites.

Similarly, as illustrated in **Fig. 3c**, **Supplementary Table S4**, and **Supplementary**

Fig. S3b, the model-derived importance scores consistently prioritize biologically

relevant RBP–target interactions, as evidenced by the cumulative overlap between high-

contribution gene regions and eCLIP peaks, which was substantially greater than that

observed in the random control. We assessed this by analyzing eCLIP data for 29 RNA-

binding proteins (RBPs) in K562 cells, comparing high-contribution regions—defined

using the same thresholds as for DBPs—to eCLIP peaks mapped to transcribed regions

(from TSS to TES). Collectively, these results demonstrate that the model effectively

captures functionally meaningful NABP–target associations for both DNA- and RNA-

binding proteins.

Significant differences in the predicted regulatory targets of several NABPs were

observed between K562 and HepG2 cells based on DeepLIFT contribution scores (**Fig.**

3d). To assess cell type–specific regulatory activity, we compared the DeepLIFT scores

of genes harboring NABP binding sites, as defined by ChIP-seq or eCLIP. Because our

deep learning model identifies target candidates from NABP-associated co-expressed

gene sets, not all predicted targets necessarily reflect direct binding; rather, they represent

genes that participate in shared or downstream regulatory programs. RPLP0 exhibited

nearly identical score distributions across the two cell types, whereas ETF1, NELFE, and

CAMTA2 showed marked divergence. These differences indicate that the model captures

and adjusts NABP-dependent regulatory relationships in a cell type–specific manner.

Deep learning uncovers functionally enriched NABP target genes beyond co-expression networks

Deep learning predictions revealed that nucleic acid–binding proteins (NABPs)

possess regulatory target genes with coherent and biologically specific functional

enrichment, extending well beyond patterns detectable from co-expression alone (**Tables**

1, 2 and **Supplementary Note 2**). Across twelve representative NABPs[27,28], predicted

target sets consistently showed significant enrichment for biologically related Gene

Ontology (GO) terms, whereas randomly sampled gene sets of matched size displayed no

such enrichment (**Fig. 4**). Notably, several enriched annotations were absent from the

broader pools of co-expressed genes (Pearson's z-score > 2), indicating that the model

selectively extracts biologically relevant signals from large co-expression neighborhoods.

Functional enrichment across GO Slim, Reactome, and PANTHER Pathways further

confirmed this specificity (**Table 1**)[29]. A ChatGPT-based gene set interpretation method

independently supported these results, providing functional summaries and literature

consistent with GO enrichment and UniProt annotations [30] (**Table 2** and **Supplementary**

Note 2)[31].

Cell type–dependent functional specificity revealed by deep learning–

refined NABP targets

Deep learning refinement further exposed cell type–specific biological functions

associated with regulatory targets of several RBPs. For example, the predicted targets of

PKM showed distinct pathway enrichments between K562 cancer cells and neural

progenitor cells (NPCs). Pathways such as *Pentose phosphate pathway*, *Glycolysis*, and

Hypoxia response mediated by HIF activation were significantly enriched in K562 but

not in NPCs, whereas *Integrin signaling pathway*, *Glycogen metabolism*, and *Glutamate*

neurotransmitter release cycle were specifically enriched in NPCs using a background

defined as 0.5× the original input feature values[32-41] (**Fig. 5**, **Table 3**, **Supplementary**

Table S5). ChatGPT-based interpretation produced consistent functional assignments

(**Supplementary Note 2**). Notably, these cell type–dependent associations align with

known biology in cancer cells and NPCs.

Additional RBPs exhibited similar specificity. Predicted target genes of ILF2 were

enriched for *G2/M DNA replication checkpoint* in K562 cells but not in NPCs[42-44] ; targets

of EIF2S2 were associated with *Serine and glycine biosynthesis* in NPCs but not in

K562[45,46] ; and targets of DNAJC2 were enriched for the *Ubiquitin proteasome pathway*

in K562 but not in NPCs[47,48] **(Fig. 5**, **Table 3**, **Supplementary Table S5)**. ChatGPT-

based inference again recapitulated these cell type–dependent functional patterns

(**Supplementary Note 2**), consistent with previous reports.

Together, these results demonstrate that the deep learning–based refinement of

NABP target predictions enables systematic detection of cell type–dependent functions,

revealing regulatory specificity not captured by co-expression alone.

Functional enrichment analysis of NABP target genes using

contribution score–based GSEA

Gene set enrichment analysis (GSEA) based on DeepLIFT-derived contribution

scores revealed robust, cell type–dependent functional redistribution among NABP

regulatory targets. Rather than ranking genes by expression levels, we ranked predicted

targets according to their relative regulatory contributions within each cell type, enabling

pathway-level interpretation based on shifts in the positional distribution of functionally

related genes along contribution-ranked lists, independent of absolute score magnitude or

sign (**Fig 6a**). Using this framework, PKM-associated pathways exhibited pronounced

redistribution between neural progenitor cells (NPCs) and K562 leukemia cells. Multiple

Reactome pathways showed large differences in normalized enrichment scores (ΔNES =

NES_K562 $-$ NES_NPC), indicating substantial divergence in inferred regulatory

coupling across cellular contexts. Pathways related to synaptic organization and neuronal

system functions were enriched among positive ΔNES values (K562-biased), whereas

immune-related pathways were preferentially represented among negative ΔNES values

(NPC-biased). These contrasts were consistently captured across pathways ranked by

ΔNES (**Fig. 6b**, **Table 4**, and **Supplementary Fig. S5**). Importantly, ΔNES reflects

redistribution of contribution-ranked pathway positioning rather than direct activation or

repression of lineage programs. Conceptually, ΔNES quantifies the relative positional

shift of pathway-associated genes along contribution-ranked regulatory landscapes

between cellular contexts.

Because DeepLIFT contribution scores depend on background definition, we

evaluated alternative reference configurations (input × 0.5 and input × 2). Although

background scaling altered the polarity of contribution–expression correlations and the

absolute direction of individual NES values, relative ΔNES-based pathway differences

between NPC and K562 were preserved across configurations (**Fig. 6**; **Supplementary

Tables S7–S9**). The amplified background (input × 2) increased the number of genes with

non-negligible contribution scores and generally yielded higher NES magnitudes,

reflecting increased sensitivity without altering the relative redistribution patterns. These

observations support the robustness of ΔNES as a comparative metric driven by stable

rank-structure differences rather than background-dependent score polarity. Extending

this analysis beyond PKM, similar category-level shifts were observed for additional

RNA-binding proteins, including HNRNPK and NELFE (**Supplementary Tables S10–

S13**). Across all three proteins, Neural System–annotated pathways were significantly

overrepresented among high ΔNES terms, whereas Immune System–annotated pathways

were enriched among low ΔNES terms (one-sided Fisher's exact test, $p < 0.001$;

Supplementary Table S14). This reproducible enrichment pattern indicates systematic

redistribution of regulatory influence between NPC and K562 contexts that is not

restricted to a single protein, and remains consistent when evaluated at both pathway and

functional module levels.

Notably, Signal Transduction emerged as the dominant top-level Reactome

category across the ΔNES spectrum for all three RBPs (**Supplementary Table S14**).

Subpathway analysis revealed that this enrichment was driven primarily by broadly

utilized regulatory modules, including FGFR/RTK signaling, phospholipase C–mediated

cascades, WNT/BMP pathways, and receptor-associated second-messenger systems. At

the level of exact Reactome pathway identity, overlap across RBPs was limited, with only

a single pathway (FGFR1C ligand binding and activation) shared among PKM,

HNRNPK, and NELFE, and a small number of pairwise overlaps such as NF-κB

activation and signaling between PKM and NELFE (**Supplementary Table S16**).

Consistent with this observation, pairwise comparison using the Jaccard similarity index

revealed strong similarity between HNRNPK and NELFE ($J \approx 0.65$), whereas PKM

shared only limited overlap with either protein (PKM–HNRNPK: $J \approx 0.13$; PKM–

NELFE: $J \approx 0.15$), indicating divergence at the level of individual pathway identity.

However, grouping pathways into functional modules revealed a markedly different

structure. Despite limited exact overlap, all three RBPs converged on common receptor-

proximal signaling modules, particularly within the FGFR signaling axis and associated

phospholipase C cascades, as well as ligand–receptor and MAPK-related signaling

processes. These findings indicate that RBPs share a conserved signaling backbone at the

functional level, even when individual pathway identities differ. Furthermore, PKM

exhibited a broader and more diversified signaling profile, incorporating additional

modules related to Rho GTPase signaling, Notch signaling, and neuronal-associated

pathways, whereas bottom-ranked pathways were largely restricted to general signaling

processes such as GPCR, WNT/BMP, and ligand-binding receptor pathways, consistent

with leukemia/hematopoietic signaling states in K562 cells. Together, these results

support a hierarchical model in which a shared signaling backbone is differentially

extended by cell type–biased functional modules, rather than being defined by exact

pathway overlap.

To enable systematic interpretation, ΔNES-ranked pathways were grouped into

functional modules based on shared biological processes (see **Methods** and

Supplementary Table S15). This module-level organization revealed a clear separation

between pathways with positive ΔNES values (relatively enriched in K562) and those

with negative ΔNES values (relatively enriched in NPC) (**Fig. 6c**). Pathways with positive

ΔNES were predominantly associated with signaling and interaction-related modules,

including neuronal system–related signaling, metabolic processes, and stemness-

associated programs. In contrast, pathways with negative ΔNES were enriched for

immune system–related modules and developmental programs, such as organogenesis

and lineage specification. Despite this divergence, a subset of signaling pathways—

including receptor tyrosine kinase (RTK), MAPK, and WNT-related pathways—was

consistently observed across RBPs, forming a shared signaling backbone. These findings

indicate that RBPs do not primarily regulate distinct sets of cell-type-specific genes, but

instead modulate common signaling networks while redistributing functional outputs

across cellular contexts.

Consistent with established roles of PKM2 in metabolic rewiring, nuclear

transcriptional regulation, and proliferative state maintenance [49,50], the observed ΔNES

redistribution is compatible with context-dependent repositioning of metabolic and

neuronal regulatory modules between leukemia and neural progenitor states [36]. HNRNPK

and NELFE, which coordinate transcriptional and post-transcriptional regulatory

processes during development and malignancy, displayed parallel redistribution patterns,

further supporting a shared regulatory architecture modulated across cellular contexts

(**Supplementary Note**: Functional interpretation of ΔNES-based pathway redistribution

across PKM, HNRNPK, and NELFE). Together, these findings demonstrate that

contribution score–based GSEA captures reproducible, category-level functional

redistribution across cell types while revealing a conserved upstream signaling backbone

shared among multiple RNA-binding proteins. This framework provides an interpretable

and scalable strategy for dissecting context-dependent regulatory organization beyond

expression-centric or threshold-dependent analyses.

To further investigate the gene-level composition of enriched pathways, we

examined leading-edge genes identified by GSEA for ΔNES-ranked Reactome pathways.

Notably, many leading-edge genes were classified as shared functional genes rather than

strictly cell-type-specific genes, even within pathways associated with neuronal or

immune systems (**Supplementary Table S17**). These results suggest that RBPs

predominantly modulate broadly shared gene networks while contributing to context-

dependent functional differences at the pathway level.

Discussion

Our findings demonstrate that RNA-binding proteins (RBPs) regulate gene

expression programs through redistribution of regulatory influence across a shared

signaling backbone, rather than by controlling entirely distinct pathway sets. Across

multiple RBPs, including PKM, HNRNPK, and NELFE, Signal Transduction

consistently forms a conserved signaling backbone, while functional modules associated

with neuronal and immune processes are differentially positioned depending on cellular

context. This hierarchical organization indicates that regulatory rewiring occurs through

redistribution across common signaling networks.

Importantly, ΔNES captures this redistribution as shifts in the positional

distribution of pathway-associated genes along contribution-ranked lists, rather than

reflecting direct pathway activation or repression. Because gene expression represents the

combined effects of multiple regulatory inputs, contribution scores quantify relative

regulatory influence, enabling detection of subtle but systematic differences in regulatory

architecture across cell types.

At the pathway level, the shared signaling backbone is characterized by broadly

utilized receptor-mediated and growth factor–associated pathways, including RTK,

MAPK, and WNT-related signaling. Although individual pathway identities vary across

RBPs, functional module–level analysis reveals convergence on common upstream

signaling processes. This structural convergence distinguishes our findings from

conventional enrichment-based analyses, which typically emphasize pathway identity

rather than organization.

In contrast, modules associated with neuronal functions and immune responses

exhibit context-dependent redistribution, indicating that RBPs modulate how shared gene

networks are functionally deployed across cellular states. This model contrasts with

transcription factors, which often determine cellular identity through selective activation

of more restricted gene sets. Instead, RBPs appear to regulate functional allocation within

broadly shared signaling and metabolic networks.

The robustness of these patterns across alternative DeepLIFT background

settings further supports the interpretation that pathway-level contrasts arise from stable

rank-structure differences rather than background-dependent artifacts. While co-

expression–based inference does not directly establish causal regulatory relationships, it

captures functional coupling at the systems level, providing a scalable framework for

analyzing regulatory architecture in the absence of experimentally defined binding

information.

Looking forward, expanding this framework to a broader range of cell types and

regulatory factors will enable systematic characterization of pathway redistribution across

diverse biological contexts. Comparative analysis across transcription factors and RNA-

binding proteins may further clarify their distinct and complementary roles in shaping

regulatory architectures. Extending the framework to inter-individual variation and

disease-associated transcriptomes, as demonstrated in related deep learning studies of

gene expression regulation, may provide insight into the regulatory basis of phenotypic

variation and disease susceptibility. Such analyses could facilitate the identification of

early disease-associated shifts in regulatory architecture and support applications in

diagnosis and preventive medicine.

Furthermore, by linking predicted regulatory targets to pathway-level

redistribution patterns, this framework may contribute to the reconstruction of gene

regulatory networks and enable in silico prediction of transcriptional responses to

perturbations. These capabilities could support the identification of candidate therapeutic

targets and guide strategies for modulating gene expression programs in a controlled

manner.

419   Together, our results establish contribution score–based pathway redistribution

420 as a framework for uncovering conserved signaling architectures alongside context-

421 dependent functional modules, providing a systems-level perspective on gene regulation

422 beyond expression-centric analyses.

425 **Methods**

426 **Predictive modeling of gene expression across diverse cell types using**

427 **deep learning and NABP binding information**

428   To predict gene expression across multiple human cell types, we adapted a

429 previously established deep learning framework that integrates DNA-binding site data for

430 transcription factors and other regulatory proteins. In this implementation, regulatory

431 elements located both proximally and distally (up to ±1 Mb) from transcription start sites

432 (TSSs) were included as input features (**Fig. 1**)[25,51]. To minimize the effects of cellular

433 heterogeneity in bulk tissue, RNA-seq datasets were selected at the individual cell-type

434 level. Publicly available RNA-seq data were retrieved for four functionally distinct cell

435 types: human foreskin fibroblasts (HFFs), mammary epithelial cells (HMECs), neural

436 progenitor cells (NPCs), and either HepG2 hepatoma cells or K562 leukemia cells,

437 depending on the class of regulatory protein analyzed[52,53]. Gene expression values

438 (FPKM) were computed using the RSeQC toolkit based on GENCODE v19 annotations,

439 and subsequently $\log_2$-transformed after adding a small offset to avoid undefined values.

440 Following the approach described by Tasaki et al.[25,51], expression fold changes were

441 calculated relative to the median expression level across the four cell types. For each

442 model iteration, three common cell types (HFFs, HMECs, and NPCs) were used, and

either HepG2 (for DNA-binding protein analysis) or K562 (for RNA-binding protein

analysis) was included to match the availability of ChIP-seq and eCLIP validation

datasets. These cell types were chosen to represent diverse biological lineages and

regulatory contexts. For specific analyses requiring direct comparison of DeepLIFT

contribution scores across cell types—such as the evaluation of cell type–dependent

regulatory differences for the same nucleic acid-binding protein (e.g., **Fig. 3d**)—models

were trained using a four-cell-type configuration comprising HFFs, NPCs, HepG2, and

K562. This unified configuration enabled direct comparison of contribution score

distributions between HepG2 and K562 while maintaining a shared background of

common cell types.

Extended regulatory features and DeepLIFT-compatible model design

To capture distal regulatory signals beyond promoter regions, we integrated DNA-

binding site data from putative enhancers into a deep learning framework for gene

expression prediction. Building upon our earlier work[25,51], this model extends the

architecture originally proposed by Tasaki et al. (2020)[25,51], which focused on promoter-

proximal regions and included RNA-binding protein and miRNA inputs. In our previous

study, aimed at identifying insulator-associated DNA-binding proteins, we deliberately

excluded RNA-binding proteins and miRNAs to focus on DNA-centered regulatory

elements. Instead, we combined DNA-binding sites from both promoter (-2 kb to +1 kb)

and enhancer-like regions (±1 Mb from the transcription start site [TSS]) into a unified

input matrix, enabling DeepLIFT analysis under the high-accuracy "genomics-default"

mode (**Supplementary Fig. S6**). In this study, we use DeepLIFT to compute attribution

values for each gene. We refer to these values as DeepLIFT scores, also described as

importance scores or contribution scores in the context of feature attribution.

Binding site coordinates were obtained from the GTRD database (v20.06)[54], and

regulatory associations were inferred using cis-eQTLs from GTEx v8. A ChIP-seq peak

that overlapped a 100-bp window centered on an expression-associated SNP was

considered linked to the corresponding transcript. Trans-eQTLs were excluded, and

coordinates were converted to the hg19 assembly using UCSC LiftOver[55]. To limit

feature dimensionality while maximizing genomic coverage, we computed peak lengths

for each DNA-binding protein (DBP) in 100-bp bins across ±1 Mb around the TSS, then

retained the maximum peak value within each 10-kb window, yielding up to 200 enhancer

bins. These were concatenated with 30 promoter-region bins to form a 230-bin unified

input layer. This single-input-layer structure was necessary because DeepLIFT's

genomics-default configuration ("RevealCancel rule on dense layers and Rescale rule on

conv layers") does not support models with multiple input layers. Concatenating all DBP-

related features into one matrix thus maintained compatibility while enhancing

interpretability.

Model training followed the protocol described by Tasaki et al. and applied in our

prior work[25,51]. Specifically, 2,705 genes from chromosome 1 were used for testing, while

22,251 genes from other chromosomes were randomly assigned to the training set and

2,472 to the validation set. Models were implemented in TensorFlow 2.6.0 and trained

on an NVIDIA RTX 3080 GPU, typically completing in 2 to 4 hours. Feature importance

was assessed using DeepLIFT, which assigned contribution scores to input features

indicating their relative influence on gene expression prediction.

Predicting NABP regulatory targets using gene co-expression–informed substitutions

490
491

To uncover regulatory targets of NABPs without relying on direct binding site

assays, we implemented a co-expression–driven modification of the deep learning input.

Specifically, DNA-binding proteins with low contribution scores in the original model

were replaced with NABPs whose regulatory interactions were inferred from gene co-

expression data. Of the 1,310 DBPs initially included, 302 were substituted with other

DBPs, and in a separate RNA-binding protein analysis, 325 were replaced with RNA-

binding proteins (RBPs) identified from the ENPD database[24]. Putative NABP–target

relationships were established using co-expression profiles provided by COXPRESdb[15],

where a Pearson's z-score > 2 between a NABP-encoding gene and a non-NABP gene

indicated a strong co-expression link. These co-expression pairs served as proxies for

regulatory interactions and were used to generate new input features for the model. To

integrate these proxies into the deep learning framework, hypothetical NABP binding

sites were assigned at the transcription start site (TSS) and 2 kb upstream of each co-

expressed gene. The model was then retrained using this modified input, and DeepLIFT

was applied to calculate feature-level importance scores (hereafter referred to as

contribution scores), which quantify the influence of each NABP–gene interaction on the

model's predicted expression output. These contribution scores represent feature-wise

attributions derived from the trained deep learning model and do not correspond directly

to measured gene expression levels, which reflect the integrated effects of multiple

regulatory inputs. This approach enables the functional characterization of NABPs with

unknown or uncharacterized binding motifs by leveraging gene co-expression as a proxy

for regulatory targeting.

Validation of predicted NABP targets using experimentally determined binding

To assess the biological relevance of the predicted regulatory targets, we compared deep learning–inferred NABP–gene associations with experimentally validated binding data. Specifically, we evaluated the spatial overlap between predicted target genes and ChIP-seq peaks for 32 DNA-binding proteins (within ±5 kb of transcription start sites [TSSs]) as well as eCLIP peaks for 29 RNA-binding proteins (within the transcribed region from TSS to transcription end site [TES]). Gene structure annotations were derived from the GENCODE v19 release[56], and binding site data were obtained from the ENCODE portal and prior publications (**Supplementary Table S6**)[57]. This overlap analysis provides orthogonal validation for the predicted targets and supports the utility of gene co-expression–based modeling for inferring NABP–target relationships.

Prioritization of NABP regulatory targets based on importance score rankings

To identify the primary regulatory targets of each NABP, genes were ranked based on the importance scores assigned to NABP binding sites by the deep learning model. Genes with binding sites consistently ranked in the top five or bottom five across all NABPs were selected as putative targets, reflecting high positive or negative regulatory influence. These targets represent genes most strongly associated with a given NABP's regulatory activity. To ensure sufficient statistical power for downstream enrichment analysis, rank thresholds were adjusted for each NABP based on the number

of high-confidence predictions. While a contribution score rank cutoff of five was

sufficient in most cases, the threshold was relaxed in specific cases—for example, up to

rank ten for proteins, which initially yielded a smaller number of predicted targets.

Functional enrichment and AI-based interpretation confirm the biological

relevance of NABP regulatory targets. To evaluate the biological coherence of predicted

NABP regulatory targets, we performed functional enrichment analysis using the

Statistical Overrepresentation Test from the PANTHER database alongside custom in-

house scripts[29]. Gene Ontology (GO) terms from biological process, molecular function,

and cellular component categories were analyzed for enrichment across three gene sets:

(1) primary regulatory target genes identified by the model, (2) all co-expressed genes

(Pearson's z-score > 2), and (3) randomly selected genes of matched size. Enrichment

was assessed using Fisher's exact test with Benjamini–Hochberg false discovery rate

(FDR) correction, and FDR < 0.05 was considered statistically significant. For this

analysis, human gene symbols were mapped to Ensembl Gene IDs using the HGNC

database (https://www.genenames.org/), and GO annotations were obtained from the

Gene Ontology resource[27,28]. Target gene IDs served as the test set, while the full

expression dataset used in the deep learning model formed the reference background.

Gene set enrichment analysis (GSEA) was performed using the pre-ranked

GSEA algorithm implemented in GSEApy (v1.1.11). For each DNA-binding protein

(DBP) and RNA-binding protein (RBP), DeepLIFT contribution scores were obtained for

all genes from the trained deep learning models. Because raw DeepLIFT scores vary in

magnitude across genes and include a large number of zeros, scores were normalized

within each gene by scaling the values across all 1,310 DBPs/RBPs to a continuous range

from −1 to 1 using min–max normalization. Genes were then ranked according to these

normalized contribution values specific to the DBP or RBP of interest. The ranked lists

were used as input for GSEA with the Reactome (v2025.1) and Gene Ontology

(Molecular Function, Biological Process, and Cellular Component, 2021 releases) gene

set collections. Analyses were performed with 10,000 permutations, weighted enrichment

statistics (weight = 1), and a minimum gene-set size of 1 to retain all functional categories.

Enrichment significance was assessed using normalized enrichment scores (NES),

nominal *p*-values, and false-discovery rates (FDR). This procedure enabled the

identification of pathways enriched among genes with high positive or negative

DeepLIFT contributions, thereby revealing regulatory roles and cell type–specific

pathways associated with each nucleic acid–binding protein. Pairwise similarity between

Signal Transduction subpathway sets was quantified using the Jaccard similarity index,

defined as the size of the intersection divided by the size of the union of the compared

sets. Unlike conventional pathway activity methods such as GSVA [58] or AUCell [59], which

rank genes based on expression levels, our framework ranks genes using deep learning–

derived contribution scores, enabling pathway analysis in terms of inferred regulatory

influence.

To complement the statistical analyses, we applied a ChatGPT-based functional

inference method[31]. This approach used curated prompts—detailed in *Extended Data*

Tables 1 and 5—to predict functional roles based on gene sets, and retrieved relevant

literature using a modified version of the "4.Reference search and validation.ipynb"

script[31]. This script is publicly available on Code Ocean and GitHub

(https://doi.org/10.24433/CO.7045777.v1   ;

https://github.com/idekerlab/llm_evaluation_for_gene_set_interpretation).   The

ChatGPT-based results provided consistent and interpretable biological themes aligned

with the enrichment findings.

588   Definition of functional modules for Reactome pathways

589   To facilitate interpretation of pathway-level results, individual Reactome

pathways were grouped into higher-order functional modules based on shared biological

processes. Functional modules were manually curated by integrating pathway annotations,

pathway names, and known biological relationships among pathways. Pathways

belonging to the same top-level Reactome category were first examined and further

subdivided into functionally coherent modules (e.g., synaptic signaling, receptor tyrosine

kinase signaling, immune receptor signaling, and metabolic sub-pathways). This

grouping was performed to reduce redundancy among closely related pathways and to

enable more interpretable visualization of pathway enrichment patterns. Each pathway

was assigned to a single functional module, and a corresponding abbreviated module

name ("module short") was defined for compact representation in figures. The complete

mapping between Reactome pathways and functional modules is provided in

Supplementary Table S15. Although this module definition involves manual curation,

the grouping was guided by established biological knowledge and Reactome pathway

structure, ensuring consistency across categories and RBPs.

605   Statistical analysis

Functional enrichment was assessed using Fisher's exact test, with multiple testing

correction applied via the Benjamini–Hochberg false discovery rate (FDR) procedure.

Statistical significance was defined as FDR < 0.05. Distributions of DeepLIFT

contribution scores for nucleic acid-binding sites derived from gene co-expression data

and ChIP-seq data were compared using the Mann–Whitney U test (also known as the

Wilcoxon rank-sum test). A p-value < 0.05 was considered statistically significant. GSEA

(gene set enrichment analysis) was performed using DeepLIFT contribution scores of

DBPs and RBPs. The normalized enrichment score (NES) was used to quantify whether

regulatory target genes associated with specific biological functions were preferentially

located at the high or low end of the ranked gene list. Statistical significance was assessed

using permutation-based p-values, as implemented in the GSEA framework.

Data availability

RNA-seq data for human foreskin fibroblast (HFF) cells were obtained from the Gene

Expression Omnibus (GEO) under accession number GSM941744. RNA-seq data for

human mammary epithelial cells (HMEC) were retrieved from ENCODE via the UCSC

Genome   Browser   (http://genome.ucsc.edu/encode/)   under   the   file   name

wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam. RNA-seq data for human neural

progenitor cells (NPC) were obtained from GEO under accession number GSM915326.

RNA-seq data for HepG2 and K562 cells were obtained from the ENCODE portal under

accession numbers ENCFF670LIE and ENCFF584WML, respectively.   mRNA co-

expression   data   were   downloaded   from   COXPRESdb   v7

(https://zenodo.org/record/6861444). The list of nucleic acid-binding proteins was

obtained from the Eukaryotic Nucleic acid Binding Protein (ENPD) database

(http://qinlab.sls.cuhk.edu.hk/ENPD/). ChIP-seq data for transcription factors were

downloaded from the GTRD database (v20.06 release; https://gtrd.biouml.org). eQTL

data (GTEx_Analysis_v8_eQTL.tar) were obtained from the GTEx portal (v8 release;

https://gtexportal.org). eCLIP data for RNA-binding proteins were retrieved from the

ENCODE portal (accession ENCSR456FVU) and from Supplementary Data 2 of Van

Nostrand et al., *Nature* (2020), https://doi.org/10.1038/s41586-020-2077-3.

Code availability

The deep learning model implemented in this study is available at

https://github.com/reposit2/coexgene.

Figures

a **Step 1: Predicting Regulatory Targets of Nucleic Acid-Binding Proteins (NABPs)**

Input:
- Binding site data for DNA-binding proteins (DBPs) around each gene
- Regulatory target genes predicted based on gene co-expression data

Deep Learning Analysis:
- Predict gene expression levels in each cell type
- Assign importance scores (DeepLIFT) to NABPs for each gene

Infer regulatory targets of NABPs based on importance scores

Regulatory targets

Same functional role of genes

NABP

Step 2: Functional Characterization of Predicted NABP Targets

- Perform enrichment analysis (e.g., GO, PANTHER Protein Class, Pathways, Reactome) using tools such as PANTHER
- Apply a ChatGPT-based method to infer gene functions and retrieve supporting literature
- Integrate ontology-based and semantic approaches
- Conduct GSEA using importance scores of predicted regulatory targets

Identify functional roles of NABPs in a cell type–specific context

b 1,310 nucleic acid-binding proteins

Gene expression prediction model

ChIP-seq peaks | ChIP-seq | Gene co-expression-based sites | Input layer

Input layer → 1D convolution → 1D convolution → 1D convolution → 1D convolution → 1D convolution → 1D Global Average Pooling → Fully connected layer → Fully connected layer → Output layer

Feature extraction

1D Convolution layer

Fully-connected and output layers

25,071 genes
Gene A, Gene B, Gene C, Gene D ...

Fibroblasts
HMEC
NPC
HepG2 or K562

High Low

Output: gene expression levels across human cell types

ChIP-seq peaks of DNA-binding proteins

Gene expression level

A>G

A>G   eGene

eSNP : expression-associated SNP

TSS : Transcription start sites

eGene: expression-associated gene; a gene whose expression level is associated with an eSNP

Figure 1. Overview of the deep learning framework for predicting and functionally characterizing NABP regulatory targets.

a. This diagram outlines the overall strategy used to infer the functional roles of nucleic acid-binding proteins (NABPs) from gene expression data. Gene expression profiles from various human cell types were integrated with NABP–gene co-expression relationships derived from COXPRESdb. These inferred interactions were used to replace low-contribution DNA-binding proteins in a deep learning model originally trained on ChIP-seq–based binding site data. The modified model was trained to predict gene expression levels and identify regulatory target genes using DeepLIFT-derived importance scores. Functional annotations of the predicted target genes were assessed using conventional enrichment analyses and GSEA, along with a ChatGPT-based method for literature-guided interpretation.

b. Gene expression levels in human cells were predicted using the genomic locations of DNA-binding sites at promoters and distal elements.

During model training, a DNA-binding protein bound to a distal region overlapping an expression-associated SNP (eSNP) was linked to a potentially regulated gene (eGene) based on eQTL data. For each gene–protein pair, ChIP-seq peak locations were binned (indicated by green boxes) and formatted into a matrix as input for deep learning.

In co-expression–based models, binding sites were artificially assigned near transcription start sites (TSSs). The deep learning model consisted of 1D convolutional layers, global average pooling, and fully connected layers. It predicted gene expression levels across 25,071 genes and multiple human cell types.

Figure 2. Relationship between gene expression levels and contribution scores.

Gene expression levels (log$_2$ TPM) predicted by a deep learning model were compared

with observed expression levels using Spearman's rank correlation.

a–c. Schematic overview of input feature configurations used for gene expression

prediction.

a. DNA-binding sites of 1,310 proteins derived from ChIP-seq data.

b. Replacement of 302 low-contribution DNA-binding proteins with co-expression–

derived DNA-binding proteins.

c. Replacement of 325 DNA-binding proteins with RNA-binding proteins identified from

co-expression relationships.

d. Comparison of absolute DeepLIFT contribution scores for predicted regulatory sites

derived from gene co-expression versus ChIP-seq data, using a background defined as

0.5× the original input feature values. Each pair of violin plots represents co-expression–

based sites (right) and corresponding ChIP-seq sites not included in that set (left).

Contribution scores were significantly higher for co-expression–based sites (Wilcoxon

signed-rank test, p-values $< 10^{-6}$). Results obtained using an amplified background (input

× 2) are shown in **Supplementary Fig. S2c**.

e.f. Relationship between gene expression levels and DeepLIFT contribution scores for

representative DNA- and RNA-binding proteins across four cell types. Contribution

scores were computed using a background defined as input × 0.5 (**e**) or input × 2 (**f**). The

same set of proteins is shown in both panels, enabling direct comparison of background

effects. While background scaling alters the sign of the DeepLIFT–expression correlation,

the relative ordering of contribution scores and cell type–specific patterns are preserved.

For each protein, the Spearman correlation coefficient and corresponding *p*-value

displayed above the plot correspond to the cell type showing the strongest correlation

among the four analyzed cell types. Data points are color-coded by cell type, revealing

consistent distributional trends across all four cell types.

b

Category 1 (A1/B1) Category 3 (C1/D1) Random 1 (A1/B1) Random 3 (C1/D1)
Category 2 (A2/B2) Category 4 (C2/D2) Random 2 (A2/B2) Random 4 (C2/D2)

Cumulative number of DBPs with high-scoring gene regions overlapping ChIP-seq sites

promoter+enhancer regions

Proportion of high-scoring gene regions overlapping ChIP-seq sites

promoter regions

Proportion of high-scoring gene regions overlapping ChIP-seq sites

c

Category 1 (A1/B1) Random 1 (A1/B1)
Category 2 (A2/B2) Random 2 (A2/B2)
Category 3 (C1/D1) Random 3 (C1/D1)
Category 4 (C2/D2) Random 4 (C2/D2)

Cumulative number of RBPs with high-scoring gene regions overlapping eCLIP peaks

transcribed regions

Proportion of high-scoring gene regions overlapping eCLIP peaks

d

Figure 3. Comparison of predicted regulatory target genes with experimental ChIP-seq and eCLIP binding data. This figure illustrates the correspondence between gene-level regulatory contribution scores inferred by deep learning and experimentally identified nucleic acid-binding protein (NABP) binding sites, using a DeepLIFT background defined as 0.5× the original input feature values.

a. Rank distribution of DeepLIFT contribution scores for nucleic acid-binding sites validated by ChIP-seq (DNA-binding proteins) or eCLIP (RNA-binding proteins). Contribution scores were ranked from highest to lowest and grouped into equal-sized rank bins. Binding sites located within promoters or transcribed regions of experimentally bound genes showed non-random enrichment toward higher or lower contribution ranks compared with all binding sites.

b. Proportion of high-contribution genomic regions overlapping ChIP-seq binding sites in promoter-associated regions. The fractions of high-contribution regions

overlapping ChIP-seq peaks are shown for promoter + enhancer regions (±100 kb from

the transcription start site) and promoter-only regions (±5 kb). Multiple rank- and score-

based thresholds were applied to define high- and low-contribution regions (see

Supplementary Note and Tables S2 and S3).

c. Overlap between high-contribution regions and eCLIP peaks for RNA-binding

proteins. Cumulative distributions show the proportion of high-contribution gene regions

overlapping experimentally determined eCLIP peaks within transcribed regions (from

transcription start site to transcription end site), compared with random controls.

Corresponding analyses using an amplified DeepLIFT background (input × 2) produced

similar results and are shown in **Supplementary Figure S3**.

d. Comparison of DeepLIFT contribution scores between HepG2 and K562 cells

under different background definitions. Scatter plots display contribution scores for

predicted regulatory target genes of three representative NABPs (RPLP0, ETF1, and

NELFE), comparing K562 (x-axis) and HepG2 (y-axis). Absolute contribution scores

were $\log_2$-transformed for visualization, and only genes with experimentally validated

binding sites were included. For each NABP, results are shown for two background

configurations (input × 0.5 and input × 2).

Figure ... **annotation terms among target genes of** **nucleic** ...

Panels a ... se discovery rate (FDR)–adjusted *p*-values of <0.001 ... l enrichment analysis was conducted for three gene sets: (1) model-predicted co-expressed genes, (2) all co-expressed genes with a Pearson correlation *z*-score > 2, and (3) randomly selected genes of equal size. Gene Ontology (GO) terms from the categories of biological process, molecular function, and cellular component were combined. Fisher's exact test *p*-values were adjusted using the Benjamini–Hochberg method to control the false discovery rate (FDR).

Figure 5. Functional mapping of nucleic acid-binding proteins reveals context-dependent regulatory roles in cancer and neural progenitor cells.

A deep learning model trained on transcriptomic co-expression data was used to predict the regulatory influence of nucleic acid-binding proteins (NABPs) on gene expression. Contribution scores were computed to quantify the impact of each NABP across target genes, and the highest-ranking genes were designated as predicted regulatory targets. Functional interpretation was performed using a large language model–based inference method, which identified key biological pathways associated with each NABP.

a

b

Protein-protein interactions at synapses
ΔNES = NES_K562 – NES_NPC
ΔNES = 2.9,   ΔNES > 0 → K562-biased association

Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell
ΔNES = -2.8,   ΔNES < 0 → NPC-biased association

Figure 6. Pathway redistribution reveals a shared signaling backbone and context-dependent functional modules across RNA-binding proteins.

a. Conceptual framework of ΔNES-based pathway analysis. Gene-level contribution scores were computed using DeepLIFT and used to rank genes for each RNA-binding protein (RBP) and cellular context. Gene set enrichment analysis (GSEA) was applied to Reactome pathways, and pathway redistribution between cell types was quantified as ΔNES (NES_K562 − NES_NPC), reflecting shifts in the positional distribution of pathway-associated genes along contribution-ranked lists.

b. Representative GSEA enrichment plots illustrating pathway redistribution. Positive ΔNES values indicate relative enrichment in K562 cells, whereas negative ΔNES values indicate relative enrichment in neural progenitor cells (NPCs). These opposing

patterns demonstrate redistribution of pathway-associated genes across contribution-

ranked gene lists rather than direct pathway activation or repression. Other examples are

shown in **Supplementary Fig. S5**.

c. Functional module–level organization of ΔNES-ranked pathways across RBPs

(PKM, HNRNPK, and NELFE). A conserved signaling backbone is observed across all

RBPs, alongside context-dependent redistribution of functional modules. Top (ΔNES >

0) and bottom (ΔNES < 0) pathways were grouped into functional modules based on

shared biological processes. Numbers in the heatmap indicate the number of Reactome

pathways assigned to each functional module. Color scale indicates pathway counts or

enrichment distribution. Across all RBPs, broadly utilized signaling pathways—

including receptor tyrosine kinase (RTK), MAPK, and WNT-related signaling—form a

conserved signaling backbone shared across cellular contexts. In contrast, modules

associated with neuronal functions (e.g., synaptic signaling) and immune processes are

differentially enriched, indicating context-dependent functional redistribution. These

results support a hierarchical model in which RBPs regulate common signaling networks

while modulating cell type–specific functional outputs through redistribution of pathway-

associated genes.

Tables

Table 1. Functional roles inferred for nucleic acid-binding proteins: integration of

UniProt annotations, ChatGPT-based predictions, and confidence scores.

Functional annotations obtained from UniProt entries were systematically compared with

significantly enriched biological processes among the model-predicted regulatory targets,

as identified by the PANTHER Overrepresentation Test (see **Supplementary Table S5**).

Shared or overlapping functions between each nucleic acid-binding protein (NABP) and

its predicted targets are indicated. Functional roles inferred using a ChatGPT-based

interpretation method are highlighted in yellow, with confidence scores—defined as the

proportion of annotated genes within the gene set—provided in parentheses. Confidence

levels are categorized as high (0.87–1.00), medium (0.82–0.86), low (0.01–0.81), or none

(0). For methodological reference, see Hu M. et al., *Nat. Methods*, 2025

(https://doi.org/10.1038/s41592-024-02525-x)[31].

ZBTB34 (HepG2 cells)

C2H2 zinc finger transcription factor

RNA polymerase II transcription regulatory region sequence-specific DNA binding

regulation of transcription by RNA polymerase II

C2H2 zinc-finger–directed regulation of RNA polymerase II transcription (0.88)

ZC3H4 (HepG2 cells)

chromatin/chromatin-binding, or -regulatory protein

RNA processing factor

basal RNA polymerase II transcription machinery binding

Chromatin-regulated RNA polymerase II transcription and RNA processing (0.89)

ZKSCAN5 (HepG2 cells)

C2H2 zinc finger transcription factor

RNA polymerase II cis-regulatory region sequence-specific DNA binding

autophagy

macroautophagy

Transcription-coupled autophagy–lysosome remodeling and membrane trafficking (0.78)

ZNF446 (HepG2 cells)

DNA-binding transcription activator activity, RNA polymerase II-specific

protein import into peroxisome matrix

non-motile cilium assembly

Transcription factor–directed ciliogenesis with peroxisome import and genome maintenance (0.86)

ZNF557 (HepG2 cells)

DNA-binding transcription activator activity, RNA polymerase II-specific

mRNA polyadenylation factor

RNA methyltransferase

RNA processing factor

RNA metabolism protein

Transcription-coupled RNA processing and RNP biogenesis (0.90)

POLR2G (NPC cells)

regulation of transcription by RNA polymerase II

RNA processing factor

translation initiation factor

Mitochondrial translation

RNA processing–coupled cytosolic and mitochondrial translation (0.90)

RBM27 (K562 cells)

RNA processing factor

RNA splicing factor

RNA metabolism protein

chromatin/chromatin-binding, or -regulatory protein

SUMO is proteolytically processed

Ub-specific processing proteases

RNAPII-coupled RNA processing and proteostasis axis (0.86)

SPEN

chromatin/chromatin-binding, or -regulatory protein

non-receptor serine/threonine protein kinase

ubiquitin-protein ligase

Signaling by Rho GTPases

Chromatin remodeling–anchored RNAPII transcription coordinated by ubiquitin and Rho-kinase signaling (0.78)

PKM (K562 cells)

Hypoxia response via HIF activation

glucose 6-phosphate metabolic process

Disorders of transmembrane transporters

Membrane Trafficking

Hypoxia/HIF-1–orchestrated metabolic reprogramming with cytoskeletal–membrane remodeling (0.80)

PKM (NPC cells)

Cellular response to hypoxia

KEAP1-NFE2L2 (NRF2) pathway

actin or actin-binding cytoskeletal protein

Hypoxia-driven metabolic rewiring with KEAP1–NRF2 signaling and actin–endosomal remodeling (0.86)

DNAJC2 (K562 cells)

RNA processing factor

chaperone

proteasome-mediated ubiquitin-dependent protein catabolic process

proteasomal protein catabolic process

ribosome biogenesis

RNA processing–coupled ribosome biogenesis and translation with proteostasis surveillance (0.88)

EIF2S2 (NPC cells)

proteinogenic amino acid biosynthetic process

Serine glycine biosynthesis

Unfolded Protein Response (UPR)

NFE2L2 regulating anti-oxidant/detoxification enzymes

ATF4 activates genes in response to endoplasmic reticulum stress

Integrated stress response coordinating UPR, antioxidant defense, and amino-acid/secretory metabolism (0.86)

ILF2 (K562 cells)

RNA processing factor

nucleic acid binding

G2/M DNA replication checkpoint

MASTL Facilitates Mitotic Progression

Coupled RNA processing–nuclear export aligned with mitotic cell-cycle control (0.86)

Table 2. Functional characterization of predicted regulatory targets of nucleic

acid-binding proteins using a ChatGPT-based inference framework.

Representative biological functions were inferred for each NABP by applying a large

language model–based gene set interpretation method. Functional themes are

summarized in bold for each NABP, while numbered subcategories denote distinct

functional roles associated with individual gene subsets.

ZBTB34 (HepG2 cells)
C2H2 zinc-finger–directed regulation of RNA polymerase II transcription (0.88)
1. Sequence-specific transcriptional control (core signal)
2. Transcriptional modulators and repressors (adjacent layer)
3. Signaling inputs that gate transcriptional outputs
4. RNA- and chromatin-adjacent regulators
5. Contextual effectors shaped by the transcriptional program

ZC3H4 (HepG2 cells)
Chromatin-regulated RNA polymerase II transcription and RNA processing (0.89)
1. Sequence-specific control + chromatin remodeling
2. RNAPII initiation and elongation
3. Co-transcriptional RNA processing
4. RNP assembly and nuclear–cytoplasmic transport
5. Ribosome biogenesis and rRNA metabolism
6. Post-translational regulation of the transcriptional apparatus
7. Genome maintenance coupling
8. Growth-signal gating
9. Peripheral modules

ZKSCAN5 (HepG2 cells)
Transcription-coupled autophagy–lysosome remodeling and membrane trafficking (0.78)
1. Transcriptional control as the top-level driver
2. Phosphoinositide signaling primes autophagosome initiation
3. Core autophagy execution and cargo selection
4. Endosome–lysosome tethering and autophagosome fusion
5. Proteasome and ubiquitylation complement lysosomal turnover
6. Organelle interfaces that feed the degradative axis
7. Ciliary/centrosomal trafficking as a specialized downstream client
8. Stress-responsive transcriptional tuning

ZNF446 (HepG2 cells)
Transcription factor–directed ciliogenesis with peroxisome import and genome maintenance (0.86)
1. Transcriptional control and RNA maturation
2. Ciliogenesis and membrane/organelle logistics
3. Peroxisome matrix import and organization
4. Genome maintenance and telomere control
5. Metabolic and cofactor support
6. Growth and sensory context
7. Network integration

ZNF557 (HepG2 cells)
Transcription-coupled RNA processing and RNP biogenesis (0.90)
1. Core transcriptional control
2. Co-transcriptional RNA processing
3. RNA modification and quality control
4. RNP assembly and ribosome/tRNA/rRNA biogenesis
5. Broad TF superstructure
6. Coupled auxiliary programs (contextual, not dominant)
816 7. Network integration

PKM (K562 cells)
Hypoxia/HIF-1–orchestrated metabolic reprogramming with cytoskeletal–membrane remodeling (0.80)
1. Transcriptional hypoxia module setting the program
2. Metabolic rewiring toward glycolysis and redox support
3. Proteostasis and catabolic throughput to sustain the shift
4. Actin–cytoskeleton remodeling and focal adhesion dynamics
5. Vesicle trafficking and membrane remodeling to deliver the phenotype
6. mRNA processing and translation control coupled to stress
7. KEAP1–NFE2L2 (NRF2)–linked oxidative stress buffering (adjacent)

PKM (NPC cells)
Hypoxia-driven metabolic rewiring with KEAP1–NRF2 signaling and actin–endosomal remodeling (0.86)
1. Metabolic rewiring and hypoxia response
2. KEAP1–NRF2 cytoprotective axis
3. Actin/adhesion remodeling and endocytic–lysosomal traffic
4. Proteostasis, autophagy, and apoptosis gating
5. Integrated model

ILF2 (K562 cells)
Coupled RNA processing–nuclear export aligned with mitotic cell-cycle control (0.86)
1. Transcription initiation and elongation across Pol I/II/III
2. Co-transcriptional RNA maturation: splicing, 3′-end formation, RNA modification
3. mRNP assembly, nuclear export, and nucleocytoplasmic transport
4. Nuclear envelope/pore remodeling synchronized with mitosis
5. Post-transcriptional regulators and chromatin/epigenetic control
6. Secretory and membrane-traffic support + stress responses
7. Gene-set composition is concordant with these functions

ILF2 (NPC cells)
RNAP II/III–coupled RNA biogenesis and ribosome assembly (0.88)
1. Transcription initiation and elongation
2. Co-transcriptional RNA processing and RNP assembly
3. Nuclear export, surveillance, and decay
4. rRNA processing and ribosome biogenesis
5. tRNA metabolism and charging
6. Chromatin and repair context

DNAJC2 (K562 cells)
RNA processing–coupled ribosome biogenesis and translation with proteostasis surveillance (0.88)
1. RNA processing and export
2. Ribosome biogenesis and rRNA metabolism
3. Translation and initiation control
4. Proteostasis and stress-response coupling
5. Genome maintenance and cell-cycle gating
6. Organelle and mitochondrial support

Table 3. Functional enrichments among predicted regulatory target genes of nucleic

acid-binding proteins.

Functional enrichment analysis was performed using the PANTHER database. This table

provides selected examples of annotations that were statistically enriched among

regulatory target genes predicted by the deep learning model trained on gene co-

expression data. '**Reference**' indicates the total number of human genes; '**Analyzed**'

denotes the number of regulatory target genes examined; '**Expected**' represents the

number of genes expected to have the annotation by chance; '**Fold Enrichment**' reflects

the observed-to-expected ratio; '**Raw *p*-value**' reports the result of Fisher's exact test; and

'**FDR**' refers to the Benjamini–Hochberg false discovery rate.

DBP, RBP	Functional annotation	Enriched	Not Enriched	Analyzed / Reference	Expected	Fold Enrichment	Raw *p*-value	FDR
PKM	Pentose phosphate pathway (PC0762)	K562	NPC	3 / 9	0.15	20.24	3.E-04	1.E-02
PKM	Pentose phosphate pathway (R-HSA-71336)	K562	NPC	3 / 15	0.25	12.14	2.E-03	2.E-02
PKM	Glycolysis (PC00024)	K562	NPC	4 / 20	0.33	12.14	3.E-04	1.E-02
PKM	Hypoxia response via HIF activation (P00030)	K562	NPC	4 / 30	0.49	8.09	1.E-03	4.E-02
PKM	response to hypoxia (GO:0001666)	K562	NPC	5 / 18	0.30	16.86	8.E-06	4.E-04
ILF2	G2/M DNA replication checkpoint (R-HSA-69478)	K562	NPC	2 / 5	0.06	30.95	2.E-03	4.E-02
POLR2G	ALKBH3 mediated reversal of alkylation damage (R-HAS-112126)	K562	NPC	2 / 4	0.11	17.80	5.E-03	5.E-02
DNAJC2	Ubiquitin proteasome pathway (P00060)	K562	NPC	8 / 58	0.60	13.26	1.E-07	2.E-05
PKM	Integrin signalling pathway (P00034)	NPC	K562	12 / 192	2.70	4.45	2.E-05	1.E-03
PKM	Glycogen metabolism (R-HSA-8982491)	NPC	K562	4 / 24	0.34	11.87	3.E-04	6.E-03
PKM	Glutamate Neurotransmitter Release Cycle (R-HSA-210500)	NPC	K562	4 / 23	0.32	12.38	3.E-04	5.E-03
EIF2S2	Serine glycine biosynthesis (PC02776)	NPC	K562	3 / 6	0.09	33.63	6.E-05	1.E-02

Table 4. Cell type‒dependent enrichment of functional pathways identified by

GSEA

NES denotes the normalized enrichment score obtained from gene set enrichment

analysis (GSEA). ΔNES represents the difference in NES between K562 and NPC

(ΔNES = NES_{K562} − NES_{NPC}). This table summarizes pathway enrichment results for

PKM, shown as a representative RNA-binding protein consistent with **Fig. 6**. The

twenty Reactome pathways with the largest positive and negative ΔNES values are

listed in the upper and lower sections, respectively. Pathways with large positive ΔNES

values were predominantly annotated to high-level Reactome categories related to

neural development and differentiation, whereas pathways with large negative ΔNES

values were enriched for categories associated with immune regulation, cell cycle

control, and genome maintenance, reflecting contrasting functional associations

between NPCs and K562 cells. Corresponding analyses performed using an amplified

DeepLIFT background (input × 2) yielded qualitatively consistent pathway-level

contrasts, with generally higher enrichment scores due to an increased number of

predicted PKM-associated target genes (**Supplementary Table S8** and **S9**).

	Reactome pathway description	Reactome pathway ID	NES K562	NES NPC	Δ NES K562 - NPC	Reactome top pathway description	Reactome top pathway ID
Pathways associated with neural cell differentiation	Protein-protein interactions at synapses	R-HSA-6794362	1.71	-1.16	2.87	Neuronal System	R-HSA-112316
	Attachment and Entry	R-HSA-9694614	1.40	-1.36	2.76	Disease	R-HSA-1643685
	Golgi-to-ER retrograde transport	R-HSA-8856688	1.58	-1.17	2.75	Vesicle-mediated transport	R-HSA-5653656
	Glycosaminoglycan-protein linkage region biosynthesis	R-HSA-1971475	1.17	-1.58	2.75	Metabolism	R-HSA-1430728
	Defective EXT2 causes exostoses 2	R-HSA-3656237	1.39	-1.35	2.74	Disease	R-HSA-1643685
	Defective B4GALT7 causes EDS, progeroid type	R-HSA-3560783	1.12	-1.60	2.72	Disease	R-HSA-1643685
	Phase 0 - rapid depolarisation	R-HSA-5576892	1.46	-1.24	2.70	Muscle contraction	R-HSA-397014
	Asparagine N-linked glycosylation	R-HSA-446203	1.55	-1.13	2.68	Metabolism of proteins	R-HSA-392499
	HS-GAG degradation	R-HSA-2024096	1.17	-1.50	2.68	Metabolism	R-HSA-1430728
	Diseases associated with glycosaminoglycan metabolism	R-HSA-3560782	1.06	-1.60	2.66	Disease	R-HSA-1643685
	Regulation of CDH1 posttranslational processing and trafficking to plasma membrane	R-HSA-9768727	1.34	-1.31	2.65	Cell-Cell communication	R-HSA-1500931
	LGI-ADAM interactions	R-HSA-5682910	1.66	-0.94	2.60	Developmental Biology	R-HSA-1266738
	Extracellular matrix organization	R-HSA-1474244	1.37	-1.23	2.60	Extracellular matrix organization	R-HSA-1474244
	Synaptic adhesion-like molecules	R-HSA-8849932	1.23	-1.36	2.60	Neuronal System	R-HSA-112316
	Neuronal System	R-HSA-112316	1.57	-1.00	2.56	Neuronal System	R-HSA-112316
	Signaling by Rho GTPases, Miro GTPases and RHOBTB3	R-HSA-9716542	1.48	-1.06	2.54	Signal Transduction	R-HSA-162582
	Smooth Muscle Contraction	R-HSA-445355	1.45	-1.08	2.54	Muscle contraction	R-HSA-397014
	Ion channel transport	R-HSA-983712	1.33	-1.15	2.48	Transport of small molecules	R-HSA-382551
	Receptor-type tyrosine-protein phosphatases	R-HSA-388844	1.60	-0.88	2.48	Neuronal System	R-HSA-112316
	Respiratory syncytial virus (RSV) attachment and entry	R-HSA-9820960	1.38	-1.09	2.48	Disease	R-HSA-1643685
Pathways associated with cancer cells	Binding and Uptake of Ligands by Scavenger Receptors	R-HSA-2173782	-1.18	1.46	-2.64	Vesicle-mediated transport	R-HSA-5653656
	WNT ligand biogenesis and trafficking	R-HSA-3238698	-1.10	1.56	-2.66	Signal Transduction	R-HSA-162582
	Complement cascade	R-HSA-166658	-1.09	1.56	-2.66	Immune System	R-HSA-168256
	Role of phospholipids in phagocytosis	R-HSA-2029485	-1.32	1.34	-2.67	Immune System	R-HSA-168256
	Nucleotide Excision Repair	R-HSA-5696398	-1.28	1.39	-2.67	DNA Repair	R-HSA-73894
	Translation	R-HSA-72766	-1.14	1.60	-2.73	Metabolism of proteins	R-HSA-392499
	Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell	R-HSA-198933	-1.52	1.23	-2.75	Immune System	R-HSA-168256
	Transcription-Coupled Nucleotide Excision Repair (TC-NER)	R-HSA-6781827	-1.44	1.32	-2.76	DNA Repair	R-HSA-73894
	Creation of C4 and C2 activators	R-HSA-166786	-1.13	1.68	-2.81	Immune System	R-HSA-168256
	CD22 mediated BCR regulation	R-HSA-5690714	-1.11	1.72	-2.83	Immune System	R-HSA-168256
	Initial triggering of complement	R-HSA-166663	-1.23	1.61	-2.84	Immune System	R-HSA-168256
	Aerobic respiration and respiratory electron transport	R-HSA-1428517	-1.23	1.67	-2.90	Metabolism	R-HSA-1430728
	Response to metal ions	R-HSA-5660526	-1.23	1.69	-2.92	Cellular responses to stimuli	R-HSA-8953897
	FCGR activation	R-HSA-2029481	-1.18	1.78	-2.95	Immune System	R-HSA-168256
	Mitotic G1 phase and G1/S transition	R-HSA-453279	-1.24	1.72	-2.96	Cell Cycle	R-HSA-1640170
	M Phase	R-HSA-68886	-1.32	1.73	-3.05	Cell Cycle	R-HSA-1640170
	Mitotic Metaphase and Anaphase	R-HSA-2555396	-1.27	1.81	-3.08	Cell Cycle	R-HSA-1640170
	DNA Replication	R-HSA-69306	-1.27	1.82	-3.09	DNA Replication	R-HSA-69306
	Scavenging of heme from plasma	R-HSA-2168880	-1.44	1.67	-3.11	Vesicle-mediated transport	R-HSA-5653656
	Metallothioneins bind metals	R-HSA-5661231	-1.05	2.61	-3.66	Cellular responses to stimuli	R-HSA-8953897

Acknowledgements

Supercomputing resources were provided by the Human Genome Center, the Institute of

Medical Science, the University of Tokyo, and the ROIS National Institute of Genetics.

Author contributions

N.O. designed and conducted the research, and wrote and revised the manuscript. K.S.

revised the manuscript.

Supplementary data

Supplementary data are available at

Conflict of interests

None declared.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS)

KAKENHI [Grant Number 16K00387], a start-up research grant from Osaka University,

the KIOXIA Young Scientist Research Grant at Waseda University, the Grant for Basic

Science Research Projects from the Sumitomo Foundation, the Research Grant from the

CASIO Science Promotion Foundation, and the Research Grant from the Okawa

Foundation for Information and Telecommunications awarded to N.O., JST CREST

[Grant Number JPMJCR23N1], and the GteX Program Japan [Grant Number

JPMJGX23B4] awarded to K.S.

Supplementary Note and Tables

Supplementary Note for Figure 3. Definition of contribution score–

based regions, thresholds, and enrichment analyses

Contribution score ranking and binning

For each nucleic acid-binding protein (NABP), DeepLIFT contribution scores were

computed for all predicted gene-level regulatory regions and ranked from highest positive

to lowest negative values. Ranked scores were divided into consecutive bins of 50 ranks,

yielding 26 bins per NABP (rank 1: 1–50; rank 2: 51–100; …; rank 26: 1,251–1,310),

unless otherwise specified.

Definition of high- and low-contribution regions

High- and low-contribution regions were defined using both rank-based and score-

based thresholds to assess the robustness of enrichment results. Rank-based thresholds

included upper and lower rank cutoffs (e.g., top or bottom 20 and 50 ranks), while score-

based thresholds were defined using absolute contribution score cutoffs (e.g., ≥ 0.01 or $\geq$

0.001). All threshold combinations used in the analyses are summarized in **Tables S2** and

S3.

Genomic regions used for overlap analyses

For DNA-binding proteins, ChIP-seq overlap analyses were conducted within two

genomic contexts: promoter-only regions defined as ± 5 kb from the transcription start site

(TSS), and extended promoter–enhancer regions defined as ± 100 kb from the TSS. For

RNA-binding proteins, eCLIP overlap analyses were restricted to transcribed regions

spanning from the TSS to the transcription end site (TES), including both exonic and

intronic intervals.

Random controls

Random control regions were generated by sampling genomic intervals matched

for size and genomic context, excluding annotated binding sites, to estimate the expected

overlap by chance. All enrichment and overlap analyses were repeated using these

controls for comparison.

Background configuration for DeepLIFT analyses

Unless otherwise noted, DeepLIFT contribution scores were computed using a

background defined as $0.5\times$ the original input feature values. Analyses using an

alternative background configuration (input $\times$ 2) yielded qualitatively similar results and

are presented in **Supplementary Figures S2** and **S3.**

Cell type configurations used for model training and comparative analyses

To accommodate differences in experimental data availability and analytical

objectives, multiple cell type configurations were used across analyses. For the primary

analyses of DNA-binding proteins (DBPs), models were trained using expression data

from HFFs, HMECs, NPCs, and HepG2, enabling validation against ChIP-seq datasets

available for HepG2 cells. For RNA-binding protein (RBP) analyses, models were trained

using HFFs, HMECs, NPCs, and K562 to facilitate validation using eCLIP datasets

available for K562 cells. For analyses requiring direct comparison of regulatory

contribution patterns between HepG2 and K562 for the same nucleic acid-binding

proteins (e.g., **Fig. 3d**), a unified four–cell-type configuration comprising HFFs, NPCs,

HepG2, and K562 was employed. This configuration enabled direct comparison of

DeepLIFT contribution score distributions across these two cell types while maintaining

a shared transcriptomic background. In all cases, cell type selection was driven by

analytical requirements and the availability of independent experimental validation data,

and no validation datasets were used as direct inputs during model training.

Gene co-expression–based predictions align with ChIP-seq–defined

DNA-binding targets

High-ranking predicted binding regions frequently coincide with ChIP-seq peaks,

particularly in promoter–enhancer intervals. Deep learning–derived DNA-binding site

predictions based on gene co-expression data show substantial concordance with ChIP-

seq results, especially when broader genomic regions encompassing promoters and

putative enhancers are considered. When overlap was assessed within ±100 kb of

transcription start sites (TSSs)—a window supported by prior enhancer–gene mapping

studies[60,61]—the mean and median overlap ratios reached 0.70 and 0.77, respectively

(Ratio A2 / B2; **Supplementary Tables S2 and S3**). In contrast, overlap was lower for

promoter regions alone (±5 kb), with corresponding mean and median values of 0.35 and

0.34. Several DNA-binding proteins, including AKAP8, POGK, SPEN, THAP7, ZNF142,

and ZNF691, showed notably high promoter-region overlap ratios (>0.5), suggesting

potential direct regulatory activity. Nevertheless, not all ChIP-seq peaks necessarily

reflect functionally relevant transcriptional binding events. For RNA-binding proteins,

overlap was examined within the transcribed regions of RNA molecules, and the mean

and median overlap ratios were 0.70 and 0.76, respectively (Ratio A2 / B2;

Supplementary Table S4). These findings indicate that model-inferred high-scoring

sites do not universally coincide with experimentally determined binding regions,

underscoring the complexity and context dependence of regulatory interactions.

955 Functional interpretation of ΔNES-based pathway redistribution across

956 PKM, HNRNPK, and NELFE

957   To contextualize the ΔNES-based redistribution patterns described in the main

958 text, we examined established biological roles of PKM, HNRNPK, and NELFE in

959 relation to the pathway categories identified by contribution score–based GSEA.

961 PKM

PKM2, the predominant isoform in proliferative and stem-like states, is widely

recognized as a central mediator of oncogenic metabolic rewiring and has been shown to

exert non-canonical nuclear functions that coordinate transcriptional programs during

tumorigenesis [49,50]. Beyond its canonical glycolytic role, PKM2 regulates gene expression

through protein–protein interactions with transcriptional regulators and chromatin-

associated factors. Isoform switching between PKM2 and PKM1 accompanies metabolic

transitions during differentiation, including neuronal maturation, where PKM1-enriched

states are associated with oxidative metabolism and terminal differentiation [36]. PKM has

also been identified as a ribosome-associated factor capable of modulating mRNA

translation [62], and accumulating evidence suggests that metabolic enzymes, including

pyruvate kinase, can function as non-canonical RNA-binding proteins [63].

In addition to its roles in metabolism and transcriptional regulation, PKM2 has

emerged as a key regulator of immunometabolic processes. PKM2-dependent metabolic

reprogramming has been shown to modulate cytokine production, including both pro-

inflammatory and anti-inflammatory responses, and to promote immune cell proliferation

and activation [64-66]. These findings indicate that PKM2 links metabolic state to immune

signaling outputs, consistent with the enrichment of immune-associated pathways

observed in ΔNES-based analyses.

PKM2 is also tightly integrated with major signaling pathways. For example,

receptor tyrosine kinase signaling, including fibroblast growth factor receptor 1 (FGFR1),

directly phosphorylates PKM2, inducing a shift from an active tetrameric form to a less

active dimeric state and thereby promoting the Warburg effect [67]. In addition, PKM2 has

been functionally connected to NOTCH signaling, where coordinated regulation of

PKM2 and NOTCH1 influences tumor growth and metastasis [68], as well as to GPCR-

associated pathways, such as the GPR3–β-arrestin2–PKM2 axis, which regulates

glycolysis and liver pathophysiology [69]. PKM2 further contributes to TGF-β signaling by

stabilizing TGF-β receptor complexes, thereby promoting fibrotic responses [70].

Beyond these pathways, PKM2 participates in diverse context-dependent

processes, including lipid metabolism, angiogenesis, and metastatic progression. PKM2

has been implicated as a regulator of lipid metabolic programs, particularly in cardiac

tissues [71,72], and promotes tumor angiogenesis by modulating endothelial cell junction

dynamics and ATP production [73,74]. Oxidative modification of PKM2 has also been linked

to enhanced metastatic potential in cancer cells [75].

Together, these multifunctional properties position PKM2 at the interface of

metabolism, signaling, and gene regulation. This integrated regulatory capacity is

consistent with the ΔNES-inferred redistribution of neuronal, immune, metabolic, and signaling-associated pathways across neural progenitor and leukemia contexts, supporting a model in which PKM2 contributes to context-dependent repositioning of shared functional networks rather than driving strictly lineage-specific pathway activation [76-78].

HNRNPK

HNRNPK is a multifunctional RNA-binding protein that integrates transcriptional and post-transcriptional regulatory processes. Haploinsufficiency of HNRNPK predisposes to hematologic malignancies and disrupts proliferation and differentiation programs in myeloid lineages [79]. In stem and progenitor contexts, HNRNPK coordinates transcriptional activation of proliferation-associated genes, including MYC and growth factor signaling components, while simultaneously regulating RNA stability and processing [80]. These dual transcriptional and post-transcriptional roles support a model in which HNRNPK contributes to lineage specification and tumor-associated transcriptional remodeling. Accordingly, the redistribution of immune- and neural-associated pathway annotations across cellular contexts observed here is compatible with a broader function of HNRNPK in modulating differentiation–proliferation balance rather than directing a single lineage-specific program.

NELFE

NELFE is a subunit of the negative elongation factor (NELF) complex that regulates promoter-proximal RNA polymerase II pausing, a metazoan-specific

mechanism critical for rapid transcriptional responsiveness and developmental gene

regulation [81,82]. Oncogenic activation of NELFE has been shown to enhance MYC-

associated transcriptional programs and promote tumor progression [83]. In hematopoietic

differentiation, dynamic modulation of NELF protein abundance alters genome-wide

transcriptional pausing and controls granulocytic lineage commitment [84]. Furthermore,

NELF regulates progenitor expansion and stem cell pool maintenance in regenerative

contexts through modulation of p53-dependent transcriptional programs [85]. Collectively,

these findings indicate that NELFE functions as a context-dependent regulator of

transcriptional architecture, influencing lineage transitions through control of elongation

dynamics. The parallel ΔNES redistribution patterns observed for NELFE and HNRNPK

are therefore consistent with convergence of distinct RNA-binding proteins on shared

regulatory infrastructures governing differentiation and proliferative states.

1033

Shared signaling architecture

Across all three proteins, Signal Transduction emerged as the dominant top-level

Reactome category. Subpathway analysis revealed enrichment of broadly utilized

regulatory modules, including FGFR/RTK signaling, IRS-mediated cascades,

WNT/BMP pathways, and receptor-associated second-messenger systems. Although

neuronal receptor–related pathways were present, they constituted a minority of enriched

subpathways (**Fig. 6c**). This distribution argues against a simple reflection of neuron-

specific transcriptional signatures and instead supports the presence of a conserved

upstream signaling backbone that is differentially interpreted across cellular contexts.

Importantly, the ΔNES-based framework does not imply direct activation or

repression of lineage programs by individual proteins. Rather, it captures hierarchical

shifts in the positioning of pathway-associated genes along contribution-ranked lists, reflecting redistribution of inferred regulatory influence across cellular states. The convergence of redistribution patterns across metabolically (PKM), transcriptionally integrative (HNRNPK), and elongation-regulatory (NELFE) proteins supports the broader interpretation that contribution-based pathway ranking can reveal conserved regulatory scaffolds underlying context-dependent functional organization.

Supplementary Table S1. Co-expression–based selection enhances the contribution scores of NABP binding sites.

Binding sites identified through gene co-expression analysis exhibited higher absolute DeepLIFT contribution scores than those selected solely from ChIP-seq data without co-expression information. This trend was consistently observed for DNA-binding proteins and was significantly stronger for RNA-binding proteins across the analyzed cell types. For each group, the table presents the mean and median values of both positive and negative contribution scores. These results indicate that gene co-expression–based selection preferentially highlights functionally relevant regulatory interactions. The upper and lower sections of the table correspond to analyses performed using DeepLIFT background references scaled to input × 0.5 and input × 2, respectively.

DBP / RBP	Cell type	Co-expression–based predictions Upper row: mean values Lower row: median values	Non-co-expression–based predictions Upper row: mean values Lower row: median values
DBP	HepG2	5.3.E-03 2.8.E-03	3.7.E-03 1.9.E-03
RBP	K562	8.0.E-03 5.6.E-03	3.6.E-03 1.8.E-03
DBP	NPC	3.9.E-03 1.9.E-03	3.8.E-03 1.9.E-03
RBP	NPC	8.2.E-03 5.9.E-03	3.6.E-03 1.7.E-03
DBP	HMEC	4.0.E-03 1.8.E-03	3.8.E-03 1.8.E-03
RBP	HMEC	8.9.E-03 6.4.E-03	3.5.E-03 1.7.E-03
DBP	HFF	3.9.E-03 1.7.E-03	3.8.E-03 1.8.E-03
RBP	HFF	8.8.E-03 6.5.E-03	3.5.E-03 1.7.E-03

DBP / RBP	Cell type	Co-expression–based: mean (upper) median (lower)	Non-co-expression–based: mean (upper) median (lower)
DBP	HepG2	5.28.E-03 2.78.E-03	3.71.E-03 1.90.E-03
RBP	K562	7.95.E-03 5.60.E-03	3.56.E-03 1.78.E-03
DBP	NPC	3.88.E-03 1.87.E-03	3.82.E-03 1.87.E-03
RBP	NPC	8.20.E-03 5.90.E-03	3.56.E-03 1.73.E-03
DBP	HMEC	4.01.E-03 1.80.E-03	3.81.E-03 1.80.E-03
RBP	HMEC	8.92.E-03 6.37.E-03	3.55.E-03 1.68.E-03
DBP	HFF	3.89.E-03 1.70.E-03	3.82.E-03 1.79.E-03
RBP	HFF	8.76.E-03 6.49.E-03	3.55.E-03 1.71.E-03

1064

1065

1066

Supplementary Table S2. Enrichment of high-contribution promoter regions with experimentally validated DBP binding sites. (See Supplementary Excel file)

This table summarizes the proportion of promoter regions (±5 kb from transcription start sites [TSSs]) that were predicted to be functionally important by DeepLIFT contribution scores and validated by ChIP-seq data for the corresponding DNA-binding proteins (DBPs). Overlap ratios were calculated across four thresholds—two based on rank and two based on absolute score—to assess the robustness of the predictions.

High- and low-contribution regions were classified using the following thresholds for the 1,310 NABPs analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank ≤ 20 or ≥ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank ≤ 50 or ≥ 1,260)

Score-based thresholds:

(C1 and D1) contribution score ≥ 0.01 or ≤ –0.01

(C2 and D2) contribution score ≥ 0.001 or ≤ –0.001

Supplementary Table S3. Enrichment of high-contribution promoter and enhancer regions with ChIP-seq–validated DBP binding sites.

This table summarizes the proportion of predicted high-contribution regions— defined within ±100 kb of transcription start sites (TSSs), encompassing both promoter and distal enhancer regions—that overlap with ChIP-seq binding sites for the corresponding DNA-binding proteins (DBPs). Strong concordance between model-

derived predictions and ChIP-seq peaks across multiple thresholds highlights the

reliability of the contribution scoring method for identifying functionally relevant

regulatory sites.

High- and low-contribution regions were classified using the following thresholds

for the 1,310 NABPs analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank ≤ 20 or $\geq 1,290$)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank ≤ 50 or $\geq 1,260$)

Score-based thresholds:

(C1 and D1) contribution score ≥ 0.01 or ≤ -0.01

(C2 and D2) contribution score ≥ 0.001 or ≤ -0.001

1102

1103

Supplementary Table S4. Enrichment of high-contribution transcribed

regions with eCLIP–validated RBP binding sites.

This table summarizes the proportion of predicted high-contribution regions—

defined within the transcribed regions of genes (from transcription start site [TSS] to

transcription end site [TES])—that overlap with eCLIP peaks for the corresponding

RNA-binding proteins (RBPs). Strong concordance between model-based predictions

and eCLIP peaks across multiple thresholds underscores the reliability of the contribution

scoring method for identifying functionally relevant regulatory sites.

High- and low-contribution regions were classified using the following

thresholds applied to the 1,310 NABPs analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank ≤ 20 or ≥ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank ≤ 50 or ≥ 1,260)

Score-based thresholds:

(C1 and D1) contribution score ≥ 0.01 or ≤ –0.01

(C2 and D2) contribution score ≥ 0.001 or ≤ –0.001

1120

1121

References

1   Belz, G. T. & Nutt, S. L. Transcriptional programming of the dendritic cell network.

Nature reviews. Immunology **12**, 101–113 (2012). https://doi.org/10.1038/nri3149

2   Miller, J. C. *et al.* Deciphering the transcriptional network of the dendritic cell lineage.

Nature immunology **13**, 888–899 (2012). https://doi.org/10.1038/ni.2370

3   Ravasi, T. *et al.* An atlas of combinatorial transcriptional regulation in mouse and man.

Cell **140**, 744–752 (2010). https://doi.org/10.1016/j.cell.2010.01.044

4   Gertz, J. *et al.* Distinct properties of cell-type-specific and shared transcription factor

binding sites. *Molecular cell* **52**, 25–36 (2013).

https://doi.org/10.1016/j.molcel.2013.08.037

5   Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

https://doi.org/10.1016/j.cell.2018.01.029

6   Vaquerizas, J. M., Kummerfeld, S. K., Teichmann, S. A. & Luscombe, N. M. A census of

human transcription factors: function, expression and evolution. *Nature reviews. Genetics*

10, 252–263 (2009). https://doi.org/10.1038/nrg2538

7   Pierson, E. *et al.* Sharing and Specificity of Co-expression Networks across 35 Human

Tissues. *PLoS Comput Biol* **11**, e1004220 (2015).

https://doi.org/10.1371/journal.pcbi.1004220

8   Sonawane, A. R. *et al.* Understanding Tissue-Specific Gene Regulation. *Cell Rep* **21**,

1077–1088 (2017). https://doi.org/10.1016/j.celrep.2017.10.001

9   Saha, A. *et al.* Co-expression networks reveal the tissue-specific regulation of transcription

and splicing. *Genome research* **27**, 1843–1858 (2017).

https://doi.org/10.1101/gr.216721.116

10   Boyle, A. P. *et al.* Comparative analysis of regulatory information and circuits across

distant species. *Nature* **512**, 453–456 (2014). https://doi.org/10.1038/nature13668

11   Kribelbauer, J. F., Rastogi, C., Bussemaker, H. J. & Mann, R. S. Low-Affinity Binding Sites
and the Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev Biol*
35, 357–379 (2019). https://doi.org/10.1146/annurev-cellbio-100617-062719

12   Ray, D. *et al.* RNA-binding proteins that lack canonical RNA-binding domains are rarely
sequence-specific. *Sci Rep* **13**, 5238 (2023). https://doi.org/10.1038/s41598-023-32245-
9

13   Zeke, A. *et al.* Deep structural insights into RNA-binding disordered protein regions.
Wiley Interdiscip Rev RNA **13**, e1714 (2022). https://doi.org/10.1002/wrna.1714

14   Zaharias, S. *et al.* Intrinsically disordered electronegative clusters improve stability and
binding specificity of RNA-binding proteins. *J Biol Chem* **297**, 100945 (2021).
https://doi.org/10.1016/j.jbc.2021.100945

15   Obayashi, T., Kodate, S., Hibara, H., Kagaya, Y. & Kinoshita, K. COXPRESdb v8: an
animal gene coexpression database navigating from a global view to detailed investigations.
Nucleic acids research **51**, D80–D87 (2023). https://doi.org/10.1093/nar/gkac983

16   Yang, S. *et al.* COEXPEDIA: exploring biomedical hypotheses via co-expressions
associated with medical subject headings (MeSH). *Nucleic acids research* **45**, D389–D396
(2017). https://doi.org/10.1093/nar/gkw868

17   Zhu, Q. *et al.* Targeted exploration and analysis of large cross-platform human
transcriptomic compendia. *Nature methods* **12**, 211–214, 213 p following 214 (2015).
https://doi.org/10.1038/nmeth.3249

18   Miller, H. E. & Bishop, A. J. R. Correlation AnalyzeR: functional predictions from gene
co-expression correlations. *BMC Bioinformatics* **22**, 206 (2021).
https://doi.org/10.1186/s12859-021-04130-7

19   Raina, P. *et al.* GeneFriends: gene co-expression databases and tools for humans and
model organisms. *Nucleic acids research* **51**, D145–D158 (2023).
https://doi.org/10.1093/nar/gkac1031

20   Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global
discovery of conserved genetic modules. *Science (New York, N.Y.)* **302**, 249–255 (2003).
https://doi.org/10.1126/science.1087447

21   Lemoine, G. G., Scott-Boyer, M. P., Ambroise, B., Perin, O. & Droit, A. GWENA: gene
co-expression networks analysis and extended modules characterization in a single
Bioconductor package. *BMC Bioinformatics* **22**, 267 (2021).
https://doi.org/10.1186/s12859-021-04179-4

22   Warde-Farley, D. *et al.* The GeneMANIA prediction server: biological network integration
for gene prioritization and predicting gene function. *Nucleic acids research* **38**, W214–
220 (2010). https://doi.org/10.1093/nar/gkq537

23   Wong, A. K., Krishnan, A. & Troyanskaya, O. G. GIANT 2.0: genome-scale integrated
analysis of gene networks in tissues. *Nucleic acids research* **46**, W65–W70 (2018).
https://doi.org/10.1093/nar/gky408

24   Tak Leung, R. W., Jiang, X., Chu, K. H. & Qin, J. ENPD - A Database of Eukaryotic
Nucleic Acid Binding Proteins: Linking Gene Regulations to Proteins. *Nucleic acids*
research **47**, D322–D329 (2019). https://doi.org/10.1093/nar/gky1112

25   Tasaki, S., Gaiteri, C., Mostafavi, S. & Wang, Y. Deep learning decodes the principles of
differential gene expression. *Nat Mach Intell* **2**, 376–386 (2020).
https://doi.org/10.1038/s42256-020-0201-6

26   Shrikumar, A., Greenside, P. & Kundaje, A. in *Proceedings of the 34th International*
Conference on Machine Learning Vol. 70 (eds Precup Doina & Teh Yee Whye) 3145—
3153 (PMLR, Proceedings of Machine Learning Research, 2017).

27   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nature genetics* **25**, 25–29 (2000). https://doi.org/10.1038/75556

28   Gene Ontology, C. *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224** (2023).
https://doi.org/10.1093/genetics/iyad031

29   Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all.
Protein Sci **31**, 8–22 (2022). https://doi.org/10.1002/pro.4218

30   UniProt, C. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic acids*
research **53**, D609–D617 (2025). https://doi.org/10.1093/nar/gkae1010

31   Hu, M. *et al.* Evaluation of large language models for discovery of gene set function.
Nature methods **22**, 82–91 (2025). https://doi.org/10.1038/s41592-024-02525-x

32   Zahra, K., Dey, T., Ashish, Mishra, S. P. & Pandey, U. Pyruvate Kinase M2 and Cancer:
The Role of PKM2 in Promoting Tumorigenesis. *Front Oncol* **10**, 159 (2020).
https://doi.org/10.3389/fonc.2020.00159

33   Zhang, Z. *et al.* PKM2, function and expression and regulation. *Cell Biosci* **9**, 52 (2019).
https://doi.org/10.1186/s13578-019-0317-8

34   Alquraishi, M. *et al.* Pyruvate kinase M2: A simple molecule with complex functions. *Free*
Radic Biol Med **143**, 176–192 (2019).
https://doi.org/10.1016/j.freeradbiomed.2019.08.007

35   Azoitei, N. *et al.* PKM2 promotes tumor angiogenesis by regulating HIF-1alpha through
NF-kappaB activation. *Mol Cancer* **15**, 3 (2016). https://doi.org/10.1186/s12943-015-
0490-2

36   He, P. *et al.* PKM2 is a key factor to regulate neurogenesis and cognition by controlling
lactate homeostasis. *Stem Cell Reports* **20**, 102381 (2025).
https://doi.org/10.1016/j.stemcr.2024.11.011

37   Bonar, N. A. & Petersen, C. P. Integrin suppresses neurogenesis and regulates brain tissue
assembly in planarian regeneration. *Development* **144**, 784–794 (2017).
https://doi.org/10.1242/dev.139964

38   Porcheri, C., Suter, U. & Jessberger, S. Dissecting integrin-dependent regulation of neural
stem cell proliferation in the adult brain. *J Neurosci* **34**, 5222–5232 (2014).
https://doi.org/10.1523/JNEUROSCI.4928-13.2014

39   Zheng, X. *et al.* Metabolic reprogramming during neuronal differentiation from aerobic
glycolysis to neuronal oxidative phosphorylation. *eLife* **5** (2016).
https://doi.org/10.7554/eLife.13374

40   Agostini, M. *et al.* Metabolic reprogramming during neuronal differentiation. *Cell Death*
Differ **23**, 1502–1514 (2016). https://doi.org/10.1038/cdd.2016.36

41   Saez, I. *et al.* Neurons have an active glycogen metabolism that contributes to tolerance to
hypoxia. *J Cereb Blood Flow Metab* **34**, 945–955 (2014).
https://doi.org/10.1038/jcbfm.2014.33

42   Matthews, H. K., Bertoli, C. & de Bruin, R. A. M. Cell cycle control in cancer. *Nat Rev*
Mol Cell Biol **23**, 74–88 (2022). https://doi.org/10.1038/s41580-021-00404-3

43   Dominguez-Brauer, C. *et al.* Targeting Mitosis in Cancer: Emerging Strategies. *Molecular*
cell **60**, 524–536 (2015). https://doi.org/10.1016/j.molcel.2015.11.006

44   Levine, M. S. & Holland, A. J. The impact of mitotic errors on cell proliferation and
tumorigenesis. *Genes Dev* **32**, 620–638 (2018). https://doi.org/10.1101/gad.314351.118

45   Huang, X. *et al.* D-Serine regulates proliferation and neuronal differentiation of neural
stem cells from postnatal mouse forebrain. *CNS Neurosci Ther* **18**, 4–13 (2012).
https://doi.org/10.1111/j.1755-5949.2011.00276.x

46   El-Hattab, A. W. Serine biosynthesis and transport defects. *Mol Genet Metab* **118**, 153–
159 (2016). https://doi.org/10.1016/j.ymgme.2016.04.010

47   Kaida, A. & Iwakuma, T. Regulation of p53 and Cancer Signaling by Heat Shock Protein
40/J-Domain Protein Family Members. *Int J Mol Sci* **22** (2021).
https://doi.org/10.3390/ijms222413527

48   Sterrenberg, J. N., Blatch, G. L. & Edkins, A. L. Human DNAJ in cancer and stem cells.
Cancer Lett **312**, 129–142 (2011). https://doi.org/10.1016/j.canlet.2011.08.019

49   Yang, W. *et al.* ERK1/2-dependent phosphorylation and nuclear translocation of PKM2
promotes the Warburg effect. *Nat Cell Biol* **14**, 1295–1304 (2012).
https://doi.org/10.1038/ncb2629

50   Israelsen, W. J. & Vander Heiden, M. G. Pyruvate kinase: Function, regulation and role in
cancer. *Semin Cell Dev Biol* **43**, 43–51 (2015).
https://doi.org/10.1016/j.semcdb.2015.08.004

51   Osato, N. & Hamada, M. Systematic discovery of directional regulatory motifs associated
with human insulator sites. *bioRxiv* (2025).

52   Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic*
acids research **41**, D991–995 (2013). https://doi.org/10.1093/nar/gks1193

53   Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene
expression and hybridization array data repository. *Nucleic acids research* **30**, 207–210
(2002). https://doi.org/10.1093/nar/30.1.207

54   Kolmykov, S. *et al.* GTRD: an integrated view of transcription regulation. *Nucleic acids*
research **49**, D104–D111 (2021). https://doi.org/10.1093/nar/gkaa1057

55   Hinrichs, A. S. *et al.* The UCSC Genome Browser Database: update 2006. *Nucleic acids*
research **34**, D590–598 (2006). https://doi.org/10.1093/nar/gkj144

56   Mudge, J. M. *et al.* GENCODE 2025: reference gene annotation for human and mouse.
Nucleic acids research **53**, D966–D975 (2025). https://doi.org/10.1093/nar/gkae1078

57   Van Nostrand, E. L. *et al.* A large-scale binding and functional map of human RNA-
binding proteins. *Nature* **583**, 711–719 (2020). https://doi.org/10.1038/s41586-020-
2077-3

58   Hanzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for
microarray   and   RNA-seq   data.   *BMC   Bioinformatics*   **14**,   7   (2013).
https://doi.org/10.1186/1471-2105-14-7

59   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nature*
methods **14**, 1083–1086 (2017). https://doi.org/10.1038/nmeth.4463

60   Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-
range   interactions.   *Nature   methods*   **18**,   1196–1203   (2021).
https://doi.org/10.1038/s41592-021-01252-x

61   Gasperini, M. *et al.* A Genome-wide Framework for Mapping Gene Regulation via
Cellular   Genetic   Screens.   *Cell*   **176**,   1516   (2019).
https://doi.org/10.1016/j.cell.2019.02.027

62   Kejiou, N. S. *et al.* Pyruvate Kinase M (PKM) binds ribosomes in a poly-ADP ribosylation
dependent manner to induce translational stalling. *Nucleic acids research* **51**, 6461–6478
(2023). https://doi.org/10.1093/nar/gkad440

63   Castello, A. *et al.* Insights into RNA biology from an atlas of mammalian mRNA-binding
proteins. *Cell* **149**, 1393–1406 (2012). https://doi.org/10.1016/j.cell.2012.04.031

64   Toller-Kawahisa, J. E. *et al.* Metabolic reprogramming of macrophages by PKM2 promotes
IL-10   production   via   adenosine.   *Cell   Rep*   **44**,   115172   (2025).
https://doi.org/10.1016/j.celrep.2024.115172

65   Yang, P. *et al.* Pyruvate kinase M2 accelerates pro-inflammatory cytokine secretion and

cell proliferation induced by lipopolysaccharide in colorectal cancer. *Cell Signal* **27**, 1525–
1532 (2015). https://doi.org/10.1016/j.cellsig.2015.02.032

66   Liu, Z., Le, Y., Chen, H., Zhu, J. & Lu, D. Role of PKM2-Mediated Immunometabolic
Reprogramming on Development of Cytokine Storm. *Front Immunol* **12**, 748573 (2021).
https://doi.org/10.3389/fimmu.2021.748573

67   Kachel, P. *et al.* Phosphorylation of pyruvate kinase M2 and lactate dehydrogenase A by
fibroblast growth factor receptor 1 in benign and malignant thyroid tissue. *BMC Cancer*
15, 140 (2015). https://doi.org/10.1186/s12885-015-1135-y

68   Wang, J. *et al.* Down-regulation of NOTCH1 and PKM2 can inhibit the growth and
metastasis of colorectal cancer cells. *Am J Transl Res* **14**, 5455–5465 (2022).

69   Dong, T. *et al.* Activation of GPR3-beta-arrestin2-PKM2 pathway in Kupffer cells
stimulates glycolysis and inhibits obesity and liver pathogenesis. *Nat Commun* **15**, 807
(2024). https://doi.org/10.1038/s41467-024-45167-5

70   Gao, J., Zhao, Y., Li, T., Gan, X. & Yu, H. The Role of PKM2 in the Regulation of
Mitochondrial Function: Focus on Mitochondrial Metabolism, Oxidative Stress, Dynamic,
and Apoptosis. PKM2 in Mitochondrial Function. *Oxid Med Cell Longev* **2022**, 7702681
(2022). https://doi.org/10.1155/2022/7702681

71   Liu, Z., Le, Y. & Lu, D. PKM2: A novel helmsman of lipid metabolism. *Cell Signal* **134**,
111967 (2025). https://doi.org/10.1016/j.cellsig.2025.111967

72   Lee, K. C. Y. *et al.* PKM2 is a key regulator of cardiac lipid metabolism in mice.
Mitochondrion **85**, 102070 (2025). https://doi.org/10.1016/j.mito.2025.102070

73   Li, L., Zhang, Y., Qiao, J., Yang, J. J. & Liu, Z. R. Pyruvate kinase M2 in blood circulation
facilitates tumor growth by promoting angiogenesis. *J Biol Chem* **289**, 25812–25821
(2014). https://doi.org/10.1074/jbc.M114.576934

74   Gomez-Escudero, J. *et al.* PKM2 regulates endothelial cell junction dynamics and
angiogenesis via ATP production. *Sci Rep* **9**, 15022 (2019).
https://doi.org/10.1038/s41598-019-50866-x

75   He, D. *et al.* Methionine oxidation activates pyruvate kinase M2 to promote pancreatic
cancer metastasis. *Molecular cell* **82**, 3045–3060 e3011 (2022).
https://doi.org/10.1016/j.molcel.2022.06.005

76   Zhang, X., Lei, Y., Zhou, H., Liu, H. & Xu, P. The Role of PKM2 in Multiple Signaling
Pathways Related to Neurological Diseases. *Mol Neurobiol* **61**, 5002–5026 (2024).
https://doi.org/10.1007/s12035-023-03901-y

77   Venkatesh, H. S. *et al.* Electrical and synaptic integration of glioma into neural circuits.
Nature **573**, 539–545 (2019). https://doi.org/10.1038/s41586-019-1563-y

78   Yu, L., Chen, X., Sun, X., Wang, L. & Chen, S. The Glycolytic Switch in Tumors: How

Many   Players   Are   Involved?   *J   Cancer*   **8**,   3430–3440   (2017).
https://doi.org/10.7150/jca.21125

79   Gallardo, M. *et al.* hnRNP K Is a Haploinsufficient Tumor Suppressor that Regulates
Proliferation and Differentiation Programs in Hematologic Malignancies. *Cancer Cell* **28**,
486–499 (2015). https://doi.org/10.1016/j.ccell.2015.09.001

80   Li, J. *et al.* HNRNPK maintains epidermal progenitor function through transcription of
proliferation genes and degrading differentiation promoting mRNAs. *Nat Commun* **10**,
4198 (2019). https://doi.org/10.1038/s41467-019-12238-x

81   Adelman, K. & Lis, J. T. Promoter-proximal pausing of RNA polymerase II: emerging roles
in   metazoans.   *Nature   reviews.   Genetics*   **13**,   720–731   (2012).
https://doi.org/10.1038/nrg3293

82   Abuhashem, A., Garg, V. & Hadjantonakis, A. K. RNA polymerase II pausing in
development: orchestrating   transcription.   *Open   Biol*   **12**,   210220   (2022).
https://doi.org/10.1098/rsob.210220

83   Dang, H. *et al.* Oncogenic Activation of the RNA Binding Protein NELFE and MYC
Signaling in Hepatocellular Carcinoma. *Cancer Cell* **32**, 101–114 e108 (2017).
https://doi.org/10.1016/j.ccell.2017.06.002

84   Liu, X. *et al.* Dynamic Change of Transcription Pausing through Modulating NELF
Protein Stability Regulates Granulocytic Differentiation. *Blood Adv* **1**, 1358–1367 (2017).
https://doi.org/10.1182/bloodadvances.2017008383

85   Robinson, D. C. L. *et al.* Negative elongation factor regulates muscle progenitor expansion
for efficient myofiber repair and stem cell pool repopulation. *Dev Cell* **56**, 1014–1029
e1017 (2021). https://doi.org/10.1016/j.devcel.2021.02.025

1350