Title

Cell type–specific functions of nucleic acid-binding proteins revealed by

deep learning on co-expression networks

Naoki Osato [1*]

[1] School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan.

E-mail: naokiosato11@gmail.com

*Corresponding author.

Abstract

Nucleic acid-binding proteins (NABPs) exhibit cell type–specific regulatory functions, yet their target genes and roles remain incompletely defined due to limitations in current experimental methods. Here, we present a deep learning approach that integrates gene co-expression correlations to predict NABP regulatory targets and infer their functions across diverse cellular contexts, without requiring binding motif information. By replacing low-informative input features in a transcriptome-based prediction model with co-expression-derived interactions, we achieved improved accuracy in gene expression prediction. Functional enrichment analysis revealed biologically coherent annotations aligned with known NABP roles. In parallel, we employed a ChatGPT-assisted interpretation framework that provided functional predictions based on semantic patterns among gene sets—an approach that may offer interpretability even when gene sets are modest in size. Predictions supported by both statistical enrichment and ChatGPT-based inference were less likely to reflect spurious associations, reinforcing their biological relevance. Together, this integrative framework combines deep learning, co-expression networks, and generative AI to uncover both established and previously uncharacterized NABP functions in a cell type–specific manner.

Keywords

Computational biology, Deep learning, Gene expression, Gene regulation, Protein function prediction

Introduction

Understanding the cell type–specific regulatory functions of nucleic acid-binding proteins (NABPs), including both DNA-binding proteins (DBPs) and RNA-binding proteins (RBPs), remains a fundamental challenge in molecular biology [1-10]. Although experimental assays such as ChIP-seq and ChIA-PET can map DBP binding sites, they require high-quality antibodies and are not easily scalable across large sets of proteins or conditions. The challenge is even greater for RBPs, many of which lack well-defined binding motifs due to their reliance on intrinsically disordered regions (IDRs) for RNA recognition, rather than sequence-specific contacts [11-13]. Low-affinity DBP binding, which plays key roles in enhancer function and paralog specificity, is also difficult to interpret, especially in the context of phase-separated transcriptional condensates [14]. Conventional methods like eQTL mapping or DNase-seq require prior knowledge of binding motifs and struggle to resolve such indirect or context-dependent interactions [15-17]. While massively parallel reporter assays (MPRAs) and large language model–based architectures such as Enformer have improved our ability to relate sequence to function [18,19], they do not reveal which NABPs mediate the predicted regulatory effects. Thus, new strategies are needed to infer NABP–target relationships and their functional roles, especially in cases where binding motifs are unknown or direct binding evidence is lacking.

Deep learning has substantially improved the prediction of gene expression by modeling regulatory inputs from NABPs, including transcription factors and microRNAs (miRNA) [20]. For example, models incorporating ChIP-seq–derived NABP binding sites in promoter regions and predicted miRNA-binding sites in transcripts have achieved high correlation (r = 0.82) between predicted and observed expression levels. In a previous

study, we extended this approach by incorporating binding site data for DNA-binding

proteins within 1 Mb of transcription start sites (TSSs) in open chromatin, excluding

miRNAs and RBPs. To facilitate accurate computation of importance scores (also known

as contribution scores) and to streamline model interpretability, we unified the input

representation by excluding a previous input layer for RNA-binding proteins and

miRNAs [21]. While this modification slightly reduced predictive accuracy (r = 0.70), it

enabled broader coverage of distal regulatory elements and laid the foundation for

downstream functional analysis based on importance scores. These scores are later

leveraged to identify regulatory targets and infer NABP functions.

Gene co-expression analysis, a well-established method since the 2000s, offers a

complementary perspective by identifying genes with coordinated expression across

biological contexts [22-30]. These patterns, derived from transcriptomic datasets across

diverse cell types, can capture both direct and indirect regulatory associations—including

those involving NABPs with unknown or hard-to-characterize binding features. In this

study, we integrated gene co-expression data into a deep learning framework to assess its

utility for predicting gene expression and identifying context-specific regulatory targets

of NABPs. This combination provides a flexible means to infer functional relationships

even when direct binding data are lacking.

To test the effectiveness of this approach, we substituted the binding site features

of ~300 NABPs that contributed little to previous deep learning predictions with gene co-

expression–based interactions. The modified model successfully predicted gene

expression levels while assigning DeepLIFT-based importance scores to NABP–gene

interactions. Many predicted targets aligned with ChIP-seq or eCLIP-validated binding

partners and showed higher contribution scores than background interactions. Functional

enrichment analyses of ten representative NABPs revealed consistent correspondence

with known annotations in UniProt [31], and several additional functions were supported

by independent publications. To enhance interpretability, we also applied a ChatGPT-

based method to infer gene functions and retrieve supporting literature. The results were

consistent with enrichment-based findings and highlighted biologically meaningful

associations such as circadian regulation (AKAP8) and glycolysis (PKM).

This study provides a computational strategy to uncover the functional roles and

regulatory targets of NABPs without requiring experimentally defined binding sites. By

integrating gene co-expression profiles with deep learning, the framework captures both

direct and indirect interactions often missed by standard methods. This approach enables

the systematic and scalable identification of NABP functions, offering a new avenue for

exploring transcriptional regulation in a cell type–specific context.

Results

Gene co-expression features improve expression prediction accuracy

Replacing low-contribution DBP inputs with gene co-expression data led to a

substantial increase in model performance (**Fig. 1**). In human foreskin fibroblasts

(HFFs), substituting 302 out of 1,310 DBPs with co-expression profiles improved the

correlation between predicted and actual gene expression from 0.70 to 0.80 (**Fig. 2a, b**).

These co-expression inputs included both NABP-encoding and non-NABP mRNAs,

drawn from the ENPD (Eukaryotic Nucleic acid Binding Protein) database and related

motif studies [32]. A subsequent experiment replaced 325 low-contribution DBPs with co-

expression interactions involving RBPs, resulting in a further improvement in prediction

accuracy (r = 0.81; **Fig. 2c**). Collectively, these findings demonstrate that integrating co-

expression interactions into the model enhances its ability to capture gene regulatory

dynamics beyond direct binding site information.

Contribution scores calculated from co-expression–based NABP–gene interactions

showed broader variability and notably higher mean and median values than those of the

original feature set (**Fig. 2d**, **Supplementary Table S1**). Correlation analysis further

revealed consistent relationships between these contribution scores and $\log_2$-transformed

transcript expression levels across cell types (**Supplementary Fig. S2**). Example plots

demonstrated that the model adjusted its predictions in a cell type–specific manner, with

contribution scores reflecting the strength of inferred NABP regulatory influence **(Fig.**

2e).

Experimental binding data support model-inferred regulatory targets

Experimentally validated binding sites align with deep learning–inferred regulatory

targets. Genes with ChIP-seq or eCLIP peaks for a given NABP exhibited

disproportionately high or low importance ranks, indicating that experimentally

confirmed binding sites correspond to stronger regulatory influence in gene expression

prediction models. This relationship supports the biological validity of DeepLIFT-based

contribution scores, which quantify the importance of specific NABPs in driving gene

expression changes.

To evaluate this association, we analyzed ChIP-seq data for 32 DBPs in HepG2

cells and eCLIP data for 29 RBPs in K562 cells—two cell lines with the most

comprehensive publicly available NABP binding site datasets from ENCODE and other

studies. A gene was considered associated with an NABP if a ChIP-seq peak was located

within ±5 kb of its transcription start site (TSS) or if an eCLIP peak fell within its

transcribed region. As illustrated in **Fig. 3a** and **Supplementary Fig. S1**, NABPs with

experimentally mapped binding sites consistently showed greater absolute contribution

scores across their predicted target genes, reinforcing their regulatory relevance and

aligning with prior interpretations of DeepLIFT scores [20,33].

Model-derived importance scores reliably predict DBP binding sites, as evidenced

by significant overlap with ChIP-seq peaks. To evaluate this correspondence, we

analyzed ChIP-seq data for 32 DNA-binding proteins (DBPs) and quantified how

frequently high-scoring gene regions—defined by DeepLIFT contribution scores—

coincided with experimental binding evidence. Overlap ratios were calculated as the

proportion of high-contribution regions that intersected with ChIP-seq peaks, considering

both promoter-proximal (±5 kb) and distal (±100 kb) regions around transcription start

sites (TSSs). To account for the functional heterogeneity and abundance of ChIP-seq

peaks, we applied a precision-at-K approach, focusing on the top-ranked or top-scoring

gene regions most likely to reflect regulatory relevance. Four thresholds were used to

define high-contribution regions: two based on contribution score rank (top/bottom 20 or

50) and two based on absolute score magnitude (≥ 0.01 or ≤ -0.01; ≥ 0.001 or ≤ -0.001).

A control analysis using randomly selected gene regions of matched size was performed

in parallel. In all tested conditions, the overlap between ChIP-seq peaks and high-

contribution regions exceeded that of the random control, supporting the model's ability

to prioritize functionally relevant regulatory sites (**Fig. 3b**).

Deep learning uncovers functionally enriched NABP target genes

beyond co-expression networks

Model-predicted regulatory targets of nucleic acid-binding proteins (NABPs)

display consistent functional enrichment patterns, highlighting the biological specificity

of the deep learning approach (**Table 1, 2** and **Supplementary Note**). To evaluate the

functional coherence of these predicted targets, we conducted Gene Ontology (GO)

enrichment analysis on ten representative NABPs [34,35]. The selected gene sets

demonstrated statistically significant enrichment for functionally related GO terms, in

contrast to randomly sampled gene sets of equivalent size, which lacked such enrichment

(**Fig. 4**). Importantly, comparison with all co-expressed genes (Pearson's $z > 2$) revealed

that the model-selected subsets contained enriched annotations not detectable in the

broader co-expression profiles. This suggests that the deep learning model filters co-

expression signals to identify a biologically focused set of target genes. Functional

enrichment was confirmed using the PANTHER Overrepresentation Test across multiple

annotation resources, including GO Slim, Reactome, and PANTHER Pathways (**Table**

1) [36]. To complement these results, we applied a ChatGPT-based gene set interpretation

method (**Table 2** and **Supplementary Note**) [37], which inferred functions and retrieved

literature consistent with the GO-based findings and UniProt annotations [31]. Notably,

AKAP8 was linked not only to its known role in the mitotic cell cycle (UniProt) but also

to circadian clock regulation, supported by literature associating AKAP8 with molecular

clock genes and melatonin biosynthesis (**Fig. 5** and **6**; **Supplementary Table S4** and

Supplementary Note) [38]. For this analysis, AKAP8's top 50 predicted targets were

defined by ranking all genes according to their model-derived NABP contribution scores,

and selecting those for which AKAP8 received the highest score in K562 cells. These

results demonstrate that integrating co-expression–guided deep learning with large

language model–based interpretation yields robust insights into both canonical and

previously unrecognized biological functions of NABPs.

Deep learning–based refinement of NABP–target predictions enhances the

detection of biologically specific functional annotations. To evaluate how training

influences functional specificity, we compared GO term enrichment between co-

expressed gene sets and predicted regulatory targets for ten NABPs. Although the co-

expressed gene sets were larger and exhibited broader enrichment, several functionally

informative annotations were uniquely or more significantly enriched among the deep

learning–derived target genes **(Table 3**, **Supplementary Table S4 and S5)**. For instance,

circadian rhythm–related functions linked to AKAP8 and kinase- or glycolysis-related

annotations associated with PKM emerged only after model training (**Table 3** and

Supplementary Table S4) [39-42]. For PKM, we defined 332 predicted targets by ranking

all genes according to NABP contribution scores and selecting those for which PKM

ranked within the top two contributors. These results suggest that the model captures

biologically meaningful signals beyond mere expression correlations. Importantly, both

PKM and AKAP8 are RNA-binding proteins whose functional predictions were derived

from a deep learning model trained on gene expression data from K562 cells—a human

erythroleukemia cell line widely used in leukemia research. This contextual alignment

lends biological credibility to the inferred functions. ALDOA, a glycolysis-associated

partner of PKM, and AKAP8, linked to circadian gene networks, have both been

implicated in leukemia metabolism and clock regulation, underscoring the relevance of

these functions to K562 biology [38,43-46]. Furthermore, the ChatGPT-based functional

inference retrieved consistent biological themes—"circadian clock" for AKAP8 and

"glycolysis" for PKM—reinforcing the deep learning–based predictions through

independent, literature-derived validation (**Fig. 5** and **6**) [37].

In neuronal cells, however, AKAP8 exhibited minimal predicted regulatory activity,

in contrast to its circadian-linked role in cancer. The model identified very few AKAP8

targets in the neuronal context, which limited downstream functional annotation by both

PANTHER and ChatGPT-based interpretation. This was based on a contribution score–

based ranking across all genes, where only four genes had AKAP8 ranked as the top

contributor, and two genes had AKAP8 ranked among the six lowest-scoring NABPs.

These findings suggest that AKAP8 may lack broad transcriptional influence in neurons,

reinforcing its potential role as a context-dependent regulator.

PKM exhibits distinct regulatory profiles in cancer and neuronal cells, reflecting

divergent metabolic adaptations. In K562 cancer cells, PKM is associated with a

Warburg-like metabolic reprogramming, as indicated by the significant enrichment of

aldolase activity among its predicted targets, notably including ALDOC and TALDO1

(**Supplementary Table S4**) [47,48]. TALDO1, a key enzyme in the non-oxidative branch of

the pentose phosphate pathway, contributes to nucleotide biosynthesis and redox balance,

supporting the proliferative and survival needs of cancer cells. This metabolic shift aligns

with the activation of the hypoxia-inducible factor (HIF) pathway, involving HIF1A,

VEGFA, BNIP3, and DDIT4, which collectively enhance glycolysis, angiogenesis, and

adaptation to hypoxic conditions (**Supplementary Note**) [49,50]. In contrast, in neuronal

cells, ALDOC—but not TALDO1 or ALDOA—was among the predicted PKM-

regulated genes, and aldolase activity was not significantly enriched (**Supplementary**

Table S4) [51]. For this analysis, 326 predicted targets were selected based on NABP

contribution scores, where PKM ranked among the top two contributors per gene.   This

suggests that neuronal PKM (predominantly PKM1) maintains a stable glycolytic flux to

support basal ATP production, rather than engaging in metabolic reprogramming [52,53].

Furthermore, although several HIF-responsive genes (e.g., EGLN1, BNIP3, MIR210HG)

were among the predicted targets, HIF1A itself was not, indicating a context-dependent

modulation of the HIF axis in neurons. Collectively, these findings suggest that PKM is

repurposed across cell types, integrating core metabolic programs with distinct

transcriptional partners tailored to tissue-specific functional demands.

Finally, the integration of deep learning-based co-expression modeling with AI-

assisted literature mining enabled functional predictions even for less-characterized

regulators such as AKAP8. This framework facilitated both hypothesis generation and

validation across cell types, and highlights the utility of scalable, interpretable tools for

deciphering context-specific gene regulatory logic.

250 Discussion

These findings address a long-standing challenge in molecular biology: the

systematic functional characterization of nucleic acid-binding proteins (NABPs) across

diverse cellular contexts. While traditional methods rely heavily on experimental binding

data and known motifs, our approach leverages gene co-expression and machine learning

to circumvent these limitations. By capturing both direct and indirect regulatory signals,

it expands the scope of functional prediction for NABPs that lack conventional binding

site information.

Gene expression correlations alone can serve as a powerful basis for uncovering

the regulatory roles of nucleic acid-binding proteins (NABPs), even in the absence of

experimental binding data. Leveraging transcriptomic profiles from diverse public RNA-

seq datasets, our deep learning framework inferred regulatory targets of NABPs based

solely on co-expression patterns. Unlike binding site assays such as ChIP-seq or eCLIP—

which are technically demanding and limited by the availability of high-quality

antibodies—this computational approach is scalable and applicable across NABPs with

poorly characterized binding motifs. Functional enrichment analysis of the predicted

target genes revealed coherent biological roles, many of which aligned with existing

annotations in public databases. Additionally, several novel functional associations—

absent from UniProt or GO but supported by published literature—were uncovered.

These findings were further corroborated using a ChatGPT-assisted literature mining

method, which retrieved consistent biological themes and references. Together, these

results underscore the biological relevance of NABP–target predictions based on gene

expression correlations and demonstrate the value of integrating deep learning with AI-

assisted interpretation. This strategy offers a scalable path toward elucidating NABP

functions and discovering new regulatory mechanisms across diverse cellular contexts.

In cancer cells, AKAP8 appears to serve as a molecular integrator of transcriptional

regulation under environmental stress. Our analyses, combined with recent reports,

suggest that AKAP8 may support hypoxic adaptation through its interaction with DDX5,

a multifunctional RNA helicase known to promote HIF-1 transcriptional activity (**Fig. 6**)

[54]. DDX5 enhances the recruitment of HIF-1α/β heterodimers to hypoxia-responsive

elements (HREs), thus facilitating the expression of genes involved in glycolysis,

angiogenesis, and cell survival under low oxygen conditions [55]. Notably, AKAP8 has

been shown to physically interact with DDX5 in lung carcinoma cells, forming a

chromatin-associated complex that modulates R-loop stability and transcriptional

dynamics [56]. Given that DDX5 levels decrease under hypoxia, yet their re-expression

restores aberrant R-loop accumulation, AKAP8 may function as a stabilizing co-factor

that compensates for transcriptional stress in hypoxic tumor microenvironments. In

addition to hypoxia-associated functions, AKAP8 is implicated in circadian rhythm

regulation. In our study, AKAP8 was identified as a potential regulator of circadian genes

in K562 cancer cells, and ChatGPT-based inference retrieved functional annotations such

as "circadian clock" and "rhythm." Prior studies have linked the disruption of circadian

timing systems to tumor progression [57], suggesting that AKAP8 may contribute to

temporal dysregulation in cancer. Thus, AKAP8 may play dual roles by coupling

chromatin-associated RNA processing to both metabolic and temporal regulatory circuits

in malignancies. Taken together, these findings position AKAP8 as a context-dependent

regulatory hub that coordinates hypoxia response and circadian control through

interaction with DDX5 and possibly other nuclear partners. Targeting such

multifunctional scaffolds may offer new strategies for disrupting tumor cell plasticity and

transcriptional resilience in the face of environmental stress.

A key feature distinguishing cancer cells from non-proliferative cells such as

neurons is their ability to reprogram core metabolic pathways to support growth under

hypoxic and nutrient-limited conditions. Our analysis supports the presence of a PKM2–

HIF-1−ALDOC/TALDO1 regulatory axis in cancer cells (**Fig. 6**) [58-61], which coordinates

glycolytic and pentose phosphate pathway gene expression to enable metabolic flexibility.

Notably, although ALDOA is a well-established glycolytic enzyme and a known PKM

interactor in human and mouse (STRING database), it was not among the PKM co-

expressed genes in our K562 transcriptomic data (Pearson's z-score < 2), whereas

ALDOC—typically regarded as a brain-specific isoform—emerged as a strongly co-

expressed and PKM-regulated gene. This suggests that ALDOC may substitute or

complement ALDOA in hematopoietic contexts. In agreement with this, transcriptomic

profiles from the Human Protein Atlas indicate that both ALDOA and ALDOC are

expressed in K562 cells, raising the possibility that PKM2 may preferentially regulate

ALDOC under hypoxia, while ALDOA may operate independently or through non-PKM-

mediated control. Functionally, ALDOC contributes to anaerobic glycolysis and has been

implicated in chemoresistance and tumor cell survival, while TALDO1, a key enzyme in

the non-oxidative branch of the pentose phosphate pathway, facilitates carbon flux

recycling and supports NADPH and nucleotide biosynthesis—functions essential for

redox balance and proliferation under stress. These roles align with their upregulation in

various cancer models under hypoxic conditions. In contrast, neuronal cells

predominantly express the PKM1 isoform, which sustains a high-efficiency glycolytic

flux coupled to mitochondrial oxidative phosphorylation. In our neuronal cell model,

ALDOC—but not ALDOA or TALDO1—was predicted as a PKM-regulated gene, and

no significant enrichment for aldolase activity was detected. This underscores the

metabolic priorities of neurons, which emphasize consistent ATP supply over

biosynthetic plasticity, and exhibit minimal reliance on the HIF-1 pathway. The absence

of HIF1A among PKM targets in neurons further supports the notion of a transcriptionally

uncoupled hypoxic response in these cells. Together, these findings reveal how PKM

isoform usage and gene network context shape distinct metabolic wiring across cell types.

In cancer, PKM2 engages in metabolic reprogramming through the selective regulation

of genes like ALDOC and TALDO1 to support hypoxia-adaptive and proliferative states.

In neurons, PKM1 maintains homeostatic energy production under physiological

conditions. The selective engagement of ALDOC in both contexts—but through

divergent regulatory logic—highlights a potential metabolic node of interest, while the

lack of ALDOA co-expression with PKM suggests a context-dependent separation of

biochemical and transcriptional control. These insights may inform future strategies to

exploit tumor-selective metabolic dependencies.

Contribution scores derived from gene expression correlations capture both direct

and indirect regulatory interactions, offering insights beyond what ChIP-seq alone can

provide. While ChIP-seq detects thousands of DNA-binding events per protein, many of

these sites do not contribute directly to gene expression regulation and may reflect

multifunctional roles of transcription factors [62,63]. Consequently, ChIP-seq–derived

binding maps and DeepLIFT-based predictions show only partial overlap, underscoring

both methodological differences and the inherent complexity of regulatory networks. In

this study, gene co-expression analysis revealed fewer target genes than ChIP-seq but

identified subsets that are likely to be more functionally relevant. Our previous work on

insulator-associated factors linked such proteins to enhancer–promoter insulation and

chromatin organization, including phase separation and loop formation [21]. That study

incorporated eQTL data to anchor binding sites to expression outcomes. Here, by contrast,

we focused on NABPs with unknown motifs and excluded motif-based or chromatin

accessibility data such as DNase-seq or ATAC-seq, to assess the capacity of gene

expression correlations to reveal regulatory roles. Notably, the distribution of

contribution scores for co-expression–based inputs included both high and low values,

implying that gene expression patterns reflect not only direct binding but also indirect

regulation mediated through broader network effects. This highlights the value of

contribution scores as indicators of both proximal and distal regulatory influences, even

when binding motifs are unknown or experimental binding data are unavailable.

Functional roles of NABPs can be inferred even from small gene sets, provided

that the target genes exhibit coherent functional patterns. In this study, ten NABPs

showed functional enrichments consistent with known annotations, supporting the

validity of the deep learning–based predictions. These enrichments were identified using

Fisher's exact test with false discovery rate (FDR) correction via the PANTHER database,

which typically requires at least 50 genes for robust statistical significance. For proteins

with fewer than 50 predicted targets, enrichment was harder to detect, although functional

coherence was still apparent in some cases. Relaxing the ranking thresholds increased the

number of target genes and enhanced statistical power. Alternatively, more stringent

thresholds could be retained if the contribution scores were strong, offering flexibility

depending on analysis goals. Additional features—such as gene expression levels or co-

expression correlations—could further refine enrichment analyses, particularly for

borderline cases. Notably, enrichment analysis of both high- and low-ranking targets

revealed candidate novel functions for specific NABPs, suggesting that genes with large-

magnitude contribution scores are biologically informative. Moreover, a complementary

ChatGPT-based approach demonstrated utility in interpreting small gene sets. Unlike

standard enrichment tools, this method can infer biological roles even when gene counts

fall below conventional significance thresholds, provided that functionally related genes

are present. This capacity helps overcome statistical limitations and broadens the

interpretability of regulatory predictions in a cell type–specific context.

Strategic use of co-expression data enhances functional inference, while

indiscriminate replacement can diminish biological relevance. The selective integration

of gene co-expression data improved target prediction and functional coherence. In

contrast, replacing all input features with co-expression-derived interactions led to a

decrease in enriched annotations, and known functional roles were no longer recovered.

Excessive replacement of input features with gene co-expression data diminishes the

biological relevance of model predictions. When all NABP–target inputs were substituted

with co-expression-based interactions, the number of enriched functional annotations

declined, and previously identified functions from the UniProt database were no longer

recovered. This suggests that indiscriminate replacement may cause the model to capture

spurious or biologically irrelevant associations. A similar trend was observed in a prior

control analysis, where substituting true regulatory targets with randomly selected genes

led to a gradual loss of enrichment as randomness increased. Based on these observations,

the normalized count of enriched functional terms—adjusted by the number of predicted

targets—may serve as a useful proxy for evaluating the accuracy and biological validity

of NABP–target predictions [64]. These observations also raise intriguing questions about

how the deep learning model prioritizes informative input features during training. Future

work exploring the model's internal representation dynamics—such as how it

distinguishes signal from noise when presented with diverse regulatory interactions—

may provide valuable insights into the mechanisms of transcriptional regulation and

model interpretability. Investigating how the model converges on biologically

meaningful NABP–target relationships during training—possibly through attention

mechanisms or layer-wise relevance propagation—could deepen our understanding of

both gene regulation and model interpretability.

Cell type specificity and cooperative binding dynamics may underlie missing

regulatory signals for certain NABPs. While some NABPs showed no high- or low-

ranking target genes in HepG2 or K562 cells, this absence does not rule out regulatory

relevance. These proteins may exhibit cell type–specific regulatory activity detectable

only in other cellular contexts, where functional enrichment analysis could reveal distinct

biological roles. Moreover, many transcription factors and DNA-binding proteins act in

concert, forming multi-protein complexes that bind cooperatively to shared promoters [65-

67]. In such cases, individual components may not always yield strong contribution scores

due to distributed regulatory influence across the complex. This highlights the potential

of co-localization analysis as a complementary strategy to uncover cooperative regulatory

mechanisms and to disentangle the functional interplay of NABPs within transcriptional

networks.

This study establishes a scalable computational framework for uncovering the

functional roles of nucleic acid-binding proteins (NABPs) using gene expression

correlations and deep learning. By leveraging co-expression–based interactions and

contribution scoring, the model successfully identified both canonical and novel

regulatory associations across cell types—even in the absence of experimentally defined

binding motifs. The results highlight how expression-driven inference, complemented by

AI-assisted functional interpretation, can extend our understanding of NABP biology

beyond conventional binding site–centered approaches. Although the present analysis

focused on a representative subset of proteins and cell types, the framework is readily

adaptable to broader transcriptomic datasets and larger NABP cohorts. As such, it offers

a path forward for systematic functional annotation of NABPs, particularly those with

poorly characterized regulatory roles. Together, these findings lay the foundation for

future studies to decode complex gene regulatory architectures and integrate

transcriptomic and AI-driven predictions into functional genomics research at scale.

Methods

Predictive modeling of gene expression across diverse cell types using deep learning and NABP binding information

To predict gene expression across multiple human cell types, we adapted a previously established deep learning framework that integrates DNA-binding site data for transcription factors and other regulatory proteins. In this implementation, regulatory elements located both proximally and distally (up to ±1 Mb) from transcription start sites (TSSs) were included as input features (**Fig. 1**) [20,21]. To minimize the effects of cellular heterogeneity in bulk tissue, RNA-seq datasets were selected at the individual cell type level. Publicly available RNA-seq data were retrieved for four functionally distinct cell types: human foreskin fibroblasts (HFFs), mammary epithelial cells (HMECs), neural progenitor cells (NPCs), and either HepG2 hepatoma cells or K562 leukemia cells, depending on the class of regulatory protein analyzed [68,69]. Gene expression values (FPKM) were computed using the RSeQC toolkit based on GENCODE v19 annotations, and subsequently $\log_2$-transformed after adding a small offset to avoid undefined values. Following the approach described by Tasaki et al. [20,21], expression fold changes were calculated relative to the median expression level across the four cell types. For each model iteration, three common cell types (HFFs, HMECs, and NPCs) were used, and either HepG2 (for DNA-binding protein analysis) or K562 (for RNA-binding protein analysis) was included to match the availability of ChIP-seq and eCLIP validation datasets. These cell types were chosen to represent diverse biological lineages and regulatory contexts.

Extended regulatory features and DeepLIFT-compatible model design

To capture distal regulatory signals beyond promoter regions, we integrated DNA-

binding site data from putative enhancers into a deep learning framework for gene

expression prediction. Building upon our earlier work [20,21], this model extends the

architecture originally proposed by Tasaki et al. (2020) [20,21], which focused on promoter-

proximal regions and included RNA-binding protein and miRNA inputs. In our previous

study, aiming to identify insulator-associated DNA-binding proteins, we intentionally

excluded RNA-binding proteins and miRNAs to focus on DNA-centric regulatory

elements. Instead, we combined DNA-binding sites from both promoter (-2 kb to $+1$ kb)

and enhancer-like regions (± 1 Mb from the transcription start site [TSS]) into a single

input matrix, enabling DeepLIFT analysis under the high-accuracy "genomics-default"

mode.

Binding site coordinates were obtained from the GTRD database (v20.06) [70], and

regulatory associations were inferred using cis-eQTLs from GTEx v8. A ChIP-seq peak

overlapping a 100-bp window centered on an expression-associated SNP was considered

linked to the corresponding transcript. Trans-eQTLs were excluded, and coordinates were

converted to the hg19 assembly using UCSC LiftOver [71]. To limit feature dimensionality

while maximizing genomic coverage, we computed peak lengths for each DNA-binding

protein (DBP) in 100-bp bins across ± 1 Mb around the TSS, then selected the maximum

peak value across each 10-kb window, yielding up to 200 enhancer bins. These were

concatenated with 30 promoter-region bins to form a 230-bin unified input layer. This

single-layer structure was necessary because DeepLIFT's genomics-default

configuration ("RevealCancel rule on dense layers and Rescale rule on conv layers") does

not support models with multiple input layers. Concatenating all DBP-related features

into one matrix thus preserved compatibility while enhancing interpretability.

Model training followed the protocol described by Tasaki et al. and reused in our

prior work [20,21]. Specifically, 2,705 genes from chromosome 1 were used for testing,

while 22,251 genes from other chromosomes were randomly assigned to training and

2,472 to validation. Models were implemented in TensorFlow 2.6.0 with training on an

NVIDIA RTX 3080 GPU, typically completing within 2–4 hours. Feature importance

was assessed using DeepLIFT, which assigned contribution scores to input features

reflecting their impact on gene expression prediction.

484   Predicting NABP regulatory targets using gene co-expression–informed

485   substitutions

To uncover regulatory targets of NABPs without relying on direct binding site

assays, we implemented a co-expression–driven modification of the deep learning input.

Specifically, DNA-binding proteins with low contribution scores in the original model

were replaced with NABPs whose regulatory interactions were inferred from gene co-

expression data. Of the 1,310 DBPs initially included, 302 were substituted with other

DBPs, and in a separate RNA-binding protein analysis, 325 were replaced with RNA-

binding proteins (RBPs) identified from the ENPD database [32]. Putative NABP–target

relationships were established using co-expression profiles provided by COXPRESdb [22],

where a Pearson's z-score > 2 between a NABP-encoding gene and a non-NABP gene

indicated a strong co-expression link. These co-expression pairs served as proxies for

regulatory interactions and were used to generate new input features for the model. To

integrate these proxies into the deep learning framework, hypothetical NABP binding

sites were assigned at the transcription start site (TSS) and 2 kb upstream of each co-

expressed gene. The model was then retrained using this modified input, and DeepLIFT

was applied to calculate importance scores, evaluating the influence of each NABP–gene

interaction on predicted expression levels. This approach enables the functional

characterization of NABPs with unknown or uncharacterized binding motifs by

leveraging gene co-expression as a proxy for regulatory targeting.

Prioritization of NABP regulatory targets based on importance score rankings

To identify the primary regulatory targets of each NABP, genes were ranked

based on the importance scores assigned to NABP binding sites by the deep learning

model. Genes with binding sites consistently ranked in the top five or bottom five across

all NABPs were selected as putative targets, reflecting high positive or negative

regulatory influence. These targets represent genes most strongly associated with a given

NABP's regulatory activity. To ensure sufficient statistical power for downstream

enrichment analysis, rank thresholds were adjusted for each NABP depending on the

number of high-confidence predictions. While a cutoff of five was sufficient in most cases,

thresholds were relaxed to include up to eight genes for certain proteins such as RPLP0,

which initially yielded a smaller target set.

Functional enrichment and AI-based interpretation confirm the biological

relevance of NABP regulatory targets. To evaluate the biological coherence of predicted

NABP regulatory targets, we performed functional enrichment analysis using the

Statistical Overrepresentation Test from the PANTHER database alongside custom in-

house scripts [36]. Gene Ontology (GO) terms from biological process, molecular function,

and cellular component categories were analyzed for enrichment across three gene sets:

(1) primary regulatory target genes identified by the model, (2) all co-expressed genes

(Pearson's z-score > 2), and (3) randomly selected genes of matched size. Enrichment

was assessed using Fisher's exact test with Benjamini–Hochberg false discovery rate

(FDR) correction, and FDR < 0.05 was considered statistically significant. For this

analysis, human gene symbols were mapped to Ensembl Gene IDs using the HGNC

database (https://www.genenames.org/), and GO annotations were obtained from the

Gene Ontology resource [34,35]. Target gene IDs served as the test set, while the full

expression dataset used in the deep learning model formed the reference background. To

complement the statistical analysis, we applied a ChatGPT-based functional inference

method [37]. This approach used curated prompts—detailed in *Extended Data Tables 1 and*

5—to predict functional roles based on gene sets, and retrieved relevant literature using

a modified version of the "4.Reference search and validation.ipynb" script [37]. This script

is   publicly   available   on   Code   Ocean   and   GitHub

(https://doi.org/10.24433/CO.7045777.v1   ;

https://github.com/idekerlab/llm_evaluation_for_gene_set_interpretation).   The

ChatGPT-based results provided consistent and interpretable biological themes aligned

with the enrichment findings.

Validation of predicted NABP targets using experimentally determined

binding

To assess the biological relevance of the predicted regulatory targets, we

compared deep learning–inferred NABP–gene associations with experimentally

validated binding data. Specifically, we evaluated the spatial overlap between predicted

target genes and ChIP-seq peaks for 32 DNA-binding proteins (within ±5 kb of

transcription start sites [TSSs]) as well as eCLIP peaks for 29 RNA-binding proteins

(within the transcribed region from TSS to transcription end site [TES]). Gene structure

annotations were derived from the GENCODE v19 release [72], and binding site data were

obtained from the ENCODE portal and prior publications (**Supplementary Table S6**) [73].

This overlap analysis provides orthogonal validation for the predicted targets and

supports the utility of gene co-expression–based modeling for inferring NABP–target

relationships.

Statistical analysis

Functional enrichment was assessed using Fisher's exact test, with multiple testing

correction applied via the Benjamini–Hochberg false discovery rate (FDR) procedure.

Statistical significance was defined as FDR < 0.05. Distributions of DeepLIFT

contribution scores for nucleic acid-binding sites derived from gene co-expression data

and ChIP-seq data were compared using the Mann–Whitney U test (also known as the

Wilcoxon rank-sum test). A P-value < 0.05 was considered statistically significant.

Data availability

RNA-seq data for human foreskin fibroblast (HFF) cells were obtained from the Gene

Expression Omnibus (GEO) under accession number GSM941744. RNA-seq data for

human mammary epithelial cells (HMEC) were retrieved from ENCODE via the UCSC

Genome Browser (http://genome.ucsc.edu/encode/) under the file name

wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam. RNA-seq data for human neural

progenitor cells (NPC) were obtained from GEO under accession number GSM915326.

RNA-seq data for HepG2 and K562 cells were obtained from the ENCODE portal under

accession numbers ENCFF670LIE and ENCFF584WML, respectively. mRNA co-

expression data were downloaded from COXPRESdb

(https://zenodo.org/record/6861444). The list of nucleic acid-binding proteins was

obtained from the ENPD (Eukaryotic Nucleic acid Binding Protein) database

(http://qinlab.sls.cuhk.edu.hk/ENPD/). ChIP-seq data for transcription factors were

downloaded from the GTRD database (v20.06 release; https://gtrd.biouml.org). eQTL

data (GTEx_Analysis_v8_eQTL.tar) were obtained from the GTEx portal (v8 release;

https://gtexportal.org). eCLIP data for RNA-binding proteins were retrieved from the

ENCODE portal (ENCSR456FVU) and from Supplementary Data 2 of Van Nostrand et

al., *Nature* (2020), https://doi.org/10.1038/s41586-020-2077-3.

Code availability

The deep learning model used in this study is available at

https://github.com/reposit2/coexgene.

Figures

a **Step 1: Predicting Regulatory Targets of Nucleic Acid-Binding Proteins (NABPs)**

Input:
- Binding site data for DNA-binding proteins (DBPs) around each
- Regulatory target genes predicted based on gene co-expression data

Deep Learning Analysis:
- Predict gene expression levels in each cell type
- Assign importance scores (DeepLIFT) to NABPs for each gene

Infer regulatory targets of NABPs based on importance scores

Regulatory targets

Same functional role of genes

NABP

Step 2: Functional Characterization of Predicted NABP Targets

- Perform enrichment analysis (e.g., GO, PANTHER Protein Class, Pathways, Reactome) using tools like PANTHER
- Apply a ChatGPT-based method to infer functions and retrieve supporting literature
- Integrate ontology-based and semantic approaches

Identify functional roles of NABPs in a cell type–specific context

Figure 1. Overview of the deep learning framework for predicting and functionally

characterizing NABP regulatory targets.

a. This diagram outlines the overall strategy used to infer the functional roles of nucleic

acid-binding proteins (NABPs) from gene expression data. Gene expression profiles from

various human cell types were integrated with NABP–gene co-expression relationships

derived from COXPRESdb. These inferred interactions were substituted for low-

contribution DNA-binding proteins in a deep learning model originally trained on ChIP-

seq–based binding site data. The modified model was trained to predict gene expression

levels and identify regulatory target genes using DeepLIFT-derived importance scores.

Functional annotations of the predicted target genes were assessed using both

conventional enrichment analysis and a ChatGPT-based method for literature-guided

interpretation.

b. Gene expression levels in human cells were predicted using the genomic locations of

DNA-binding sites at promoters and distal elements.

During model training, a DNA-binding protein bound to a distal region overlapping an

expression-associated SNP (eSNP) was linked to a potentially regulated gene (eGene)

based on eQTL data. For each gene–protein pair, ChIP-seq peak locations were binned

(green boxes) and formatted into a matrix as input for deep learning.

In co-expression–based models, binding sites were located near transcription start sites

(TSSs). The deep learning model consisted of 1D convolutional layers, global average

pooling, and fully connected layers. It predicted gene expression levels across 25,071

genes and multiple human cell types.

Figure 2. Relationship between gene expression levels and contribution scores.

Gene expression levels ($\log_2$ TPM) predicted by a deep learning model were compared

with actual expression levels in HFF cells using Spearman's rank correlation.

a. Input included DNA-binding sites of 1,310 proteins obtained from ChIP-seq data.

b. Among these, 302 proteins with low contribution scores were replaced with an equal

number of DNA-binding proteins selected based on gene co-expression.

c. Similarly, 325 DNA-binding proteins were replaced with RNA-binding proteins

selected via co-expression data.

d. DeepLIFT contribution scores were significantly higher for binding sites derived from

gene co-expression data compared to those from ChIP-seq data (Wilcoxon signed-rank

test, $P < 2.2 \times 10^{-16}$). A and B show distributions of positive contribution scores, while C

and D show absolute values of negative contribution scores.

e. Representative examples of the correlation between a regulator's DeepLIFT

contribution score and the expression levels of its predicted target genes. Linear

regression lines with 95% confidence intervals are shown.

Figure 3. Comparison of predicted regulatory target genes with experimental ChIP-

seq and eCLIP binding data. This figure illustrates how predicted gene-level regulatory

contributions derived from deep learning align with experimentally identified nucleic

acid-binding protein targets across genomic regions.

a. Distribution of contribution scores across ChIP-seq–identified binding sites. The

x-axis indicates the rank of contribution scores assigned to nucleic acid-binding sites used

to predict gene expression levels. Scores were ranked from 1 (highest positive

contribution, left) to 1,310 (lowest negative contribution, right) and binned into 50-rank

intervals. Rank 1 corresponds to ranks 1–50, rank 2 to ranks 51–100, and rank 26 to ranks

1,251–1,310. nucleic acid-binding sites were categorized as being located in the

promoters or transcribed regions of genes experimentally bound by the nucleic acid-

binding protein shown in each graph. These sites showed higher or lower contribution

scores than those across all ChIP-seq binding sites. The y-axis indicates the number of

sites within each 50-rank bin. Positive contribution scores are indicative of gene

activation, whereas negative scores reflect gene repression.

b. Proportion of high-contribution gene regions overlapping ChIP-seq binding sites

in promoter+enhancer and promoter regions, relative to all high-contribution

regions. Labels indicating "promoter+enhancer regions" and "promoter regions" are

shown above the corresponding panels. The proportion represents the number of high-

contribution promoter+enhancer regions (within ±100 kb of the transcription start site

[TSS]; left panel) and promoter regions (within ±5 kb of the TSS; right panel) that overlap

with ChIP-seq binding sites, divided by the total number of high-contribution regions in

each category. To define high- and low-contribution regions, multiple thresholds were

applied to the 1,310 DNA-binding proteins analyzed: **Rank-based thresholds** — (A1

and B1): rank ≥ 20 or ≤ 1,290; (A2 and B2): rank ≥ 50 or ≤ 1,260. **Score-based thresholds**

Figure ... **annotation terms among target genes of**

nucleic ...

... **and** ... sted *p*-value < 0.001 and < 0.01, respectively.

... **unctio** ... hree gene sets: model-predicted co-expressed

genes, all co-expressed genes with a Pearson correlation *z*-score > 2, and randomly

670 selected genes of equal size. Gene Ontology (GO) terms from biological process,

671 molecular function, and cellular component categories were combined. Fisher's exact test

672 *p*-values were FDR-adjusted.

AKAP8 (K562 cells)
1. Circadian Control of Metabolic Regulation
This system includes multiple components of the circadian clock (*PER2, PER3, CRY2, NR1D1*), which regulate daily oscillations of gene expression in metabolic tissues. These transcriptional regulators control the rhythmic expression of key metabolic enzymes and transporters, thereby orchestrating the timing of energy storage, mobilization, and hormonal signaling across the light-dark cycle. Their co-presence with nutrient transporters and signaling proteins suggests functional integration in time-dependent metabolic homeostasis.

PKM (K562 cells)
1. Hypoxia Sensing and Adaptive Response
This system prominently features components regulated by or regulating **hypoxia-inducible factors (HIFs).** *HIF1A*, *EGLN1*, *EGLN3*, *MIR210HG*, *VEGFA*, *BNIP3*, and *DDIT4* are core mediators of the transcriptional and metabolic response to low oxygen conditions. *EGLNs* hydroxylate HIF1A under normoxia, while *HIF1A* activates genes that enhance glycolytic flux, angiogenesis (*VEGFA*), and survival during hypoxic stress (*BNIP3*, *MIR210HG*). *DDIT4* connects hypoxia to mTOR inhibition, reducing energy expenditure. These factors establish a functional axis adapting metabolism and signaling to oxygen availability.
2. Metabolic Reprogramming Toward Glycolysis
Glycolytic genes including *PFKP*, *PFKL*, *ENO1*, *ENO2*, *ENO3*, *ALDOC*, *LDHA*, and *HK2* are key effectors of the hypoxia-induced metabolic shift. HIF1A upregulates these enzymes, favoring glycolysis and lactate production over mitochondrial respiration. This shift is supported by regulatory components like *PDK1*, *PDK3*, and *SLC16A3* (lactate transporter), which further suppress pyruvate entry into mitochondria and facilitate acid-base balance under anaerobic metabolism.

Figure 5. Functional mapping of nucleic acid-binding proteins reveals context-dependent regulatory roles in cancer and neuronal cells.

A deep learning model trained on transcriptomic co-expression data was used to predict the regulatory influence of nucleic acid-binding proteins (NABPs) on gene expression. Contribution scores were computed to quantify the impact of each NABP across target genes, and top-ranking genes were designated as predicted regulatory targets. Functional interpretation was performed using a large language model–based inference method, which identified key biological pathways associated with each NABP. The resulting schematic highlights the distinct functional roles of AKAP8 and PKM in K562 leukemia cells—circadian regulation and metabolic reprogramming under hypoxia, respectively—compared with their more limited or divergent activities in neuronal cells.

Hypoxia Adaptation / Tumor Survival

Figure 6. Integrated hypoxia-responsive regulatory network orchestrated by

AKAP8 and PKM in cancer cells. This model illustrates how two nucleic acid-binding

proteins (NABPs)—AKAP8 and PKM—regulate distinct but converging transcriptional

›programs to support tumor adaptation under hypoxic stress. AKAP8 acts through

DDX5 to influence R-loop resolution [74], circadian gene expression, and stabilization of

HIF1A transcriptional activity. In parallel, PKM modulates HIF1A-driven metabolic

reprogramming, promoting glycolysis via ALDOC and redox homeostasis via

TALDO1. Predicted regulatory links were derived from contribution scores computed

by a deep learning framework and are supported by published functional studies in

cancer models.

Tables

Table 1. Functional roles inferred for nucleic acid-binding proteins: integration of

UniProt annotations, ChatGPT-based predictions, and confidence scores.

Functional annotations derived from UniProt entries were compared with significantly

enriched biological processes in the model-predicted regulatory targets, identified via the

PANTHER Overrepresentation Test (see Supplementary Table S4). Shared or

overlapping functions between each nucleic acid-binding protein (NABP) and its

predicted targets are shown. Functional roles inferred using a ChatGPT-based

interpretation method are highlighted in yellow, with parentheses indicating confidence

scores (proportion of annotated genes within the gene set). Confidence is categorized as

high (0.87–1.00), medium (0.82–0.86), low (0.01–0.81), or none (0). For reference, see

Hu M. et al., *Nat. Methods*, 2025 (https://doi.org/10.1038/s41592-024-02525-x) [37].

AKAP8

negative regulation of RNA biosynthetic process

Circadian clock system

Cry-Per complex

Circadian-regulated energy metabolism and insulin signaling (0.74)

PKM

kinase

glycolytic process

ATP metabolic process

Hypoxia-adaptive metabolic remodeling and cell migration (0.79)

POLR2G

RNA polymerase II transcription regulatory region sequence-specific DNA binding

general transcription factor

translational protein

Ribosome biogenesis and RNA maturation at the interface of translation and nuclear export (0.91)

RPLP0

large ribosomal subunit

ribonucleoprotein complex

cytosolic large ribosomal subunit

Mitochondrial-linked ribosome function and RNA-guided surveillance in translational homeostasis (0.84)

SPEN

mRNA binding

chromatin binding

transcription coregulator activity

Chromatin-associated transcriptional regulation and nuclear RNA export integration (0.83)

ZBTB34

C2H2 zinc finger transcription factor

DNA-binding transcription activator activity, RNA polymerase II-specific

RNA polymerase II transcription regulatory region sequence-specific DNA binding

Transcriptional regulation by C2H2 zinc finger proteins and signaling modulation (0.72)

ZC3H4

chromatin/chromatin-binding, or -regulatory protein

transcription regulator activity

metal ion binding

RNA-binding protein–mediated transcriptional integration and pre-mRNA processing (0.89)

ZKSCAN5

DNA-binding transcription factor activity

C2H2 zinc finger transcription factor

RNA polymerase II cis-regulatory region sequence-specific DNA binding

metal ion binding

RNA processing, ribonucleoprotein assembly, and transcription-coupled quality control (0.83)

ZNF446

C2H2 zinc finger transcription factor

extracellular space

Transcription-coupled RNA processing and mitochondrial RNA metabolism (0.81)

ZNF557

DNA-binding transcription activator activity, RNA polymerase II-specific

RNA polymerase II transcription regulatory region sequence-specific DNA binding

ion binding, metal ion binding

Transcriptional regulation and RNA metabolic coordination via zinc finger proteins and RNA-associated complexes (0.88)

Table 2. Functional characterization of predicted regulatory targets of nucleic

acid-binding proteins using a ChatGPT-based inference framework.

Representative biological functions were inferred for each NABP by applying a large

language model–assisted gene set interpretation method. Bold labels summarize the

predominant functional theme for each NABP, while numbered subcategories denote

distinct functional roles associated with individual gene subsets.

AKAP8 (K562 cells)
Circadian-regulated energy metabolism and insulin signaling (0.74)
1. Circadian Control of Metabolic Regulation
2. Insulin Signaling and Glucose Transport
3. Fatty Acid and Lipid Homeostasis
4. Metabolite Transport and Mitochondrial Function
5. Signal Integration and Vesicle Trafficking
6. Developmental and Structural Modulators

PKM (K562 cells)
Hypoxia-adaptive metabolic remodeling and cell migration (0.79)
1. Hypoxia Sensing and Adaptive Response
2. Metabolic Reprogramming Toward Glycolysis
3. Extracellular Matrix Remodeling and Migration
4. Vesicular and Membrane Trafficking
5. Mitochondrial Function and ROS Homeostasis
6. Transcriptional and Epigenetic Regulation

PKM (neuronal cells)
Hypoxia-driven metabolic and structural reprogramming (0.87)
1. Hypoxia Sensing and Transcriptional Response
2. Glycolytic Shift and Mitochondrial Modulation
3. Cytoskeletal Remodeling and Motility Enhancement
4. Endocytosis, Vesicle Transport, and Membrane Dynamics
5. RNA Processing and Translation Control
6. Ubiquitination and Proteostasis
7. Stress Signaling and Nuclear Regulation

POLR2G (K562 cells)
Ribosome biogenesis and RNA maturation at the interface of translation and nuclear export (0.91)
1. Cytoplasmic and Nuclear Ribosomal Component Assembly
2. Ribosomal RNA Processing and Modification
3. Translational Initiation and Ribosome Recycling
4. Proteasome-Linked Ribosome Quality Control
5. Nuclear Export, mRNP Surveillance, and RNA Chaperoning

RPLP0 (K562 cells)
Mitochondrial-linked ribosome function and RNA-guided surveillance in translational homeostasis (0.84)
1. Ribosomal Structural Components and Translational Core
2. RNA Modification and Surveillance by Small Non-Coding RNAs
3. RNA Surveillance and Nuclear Export Quality Control
4. Mitochondrial and Redox-Linked Translational Interface
5. Membrane Trafficking and Endoplasmic Reticulum Integration
6. Zinc Finger Regulators and DNA Repair Coupling

SPEN (K562 cells)
Chromatin-associated transcriptional regulation and nuclear RNA export integration (0.83)
1. Transcriptional Regulation and Chromatin Remodeling
2. RNA Binding, Processing, and Export
3. Protein Quality Control and Ubiquitin-Mediated Regulation
4. Signal-Responsive Transcription Integration
5. RNA Surveillance and Stability
6. Nuclear Structural and Trafficking Components

ZBTB34 (HepG2 cells)
Transcriptional regulation by C2H2 zinc finger proteins and signaling modulation (0.72)
1. Transcriptional Regulation by Zinc Finger Proteins
2. Signaling and Transcriptional Crosstalk
3. Modulators of Chromatin and RNA Stability
4. Functional Integration and Cell Fate Control

ZC3H4 (HepG2 cells)
RNA-binding protein–mediated transcriptional integration and pre-mRNA processing (0.89)
1. RNA-Binding Protein Network and Pre-mRNA Processing
2. Transcription-Coupled Regulatory Scaffold
3. Spliceosome Assembly and Dynamic Coordination
4. mRNA Export and Quality Control
5. Epigenetic and Histone Regulatory Interface
6. RNA Modification and Stability Regulation
7. Nucleolar and Ribosome Biogenesis Linkage

ZKSCAN5 (HepG2 cells)
RNA processing, ribonucleoprotein assembly, and transcription-coupled quality control (0.83)
1. RNA Processing and mRNA Surveillance
2. Transcriptional Regulation and Chromatin Remodeling
3. Ribosome Biogenesis and Mitochondrial Translation
4. Protein Quality Control and Ubiquitin-Dependent Regulation
5. Nucleocytoplasmic Transport and Organelle Coordination
6. ZNF and ZBTB Family-Mediated Regulatory Modules

ZNF446 (HepG2 cells)
Transcription-coupled RNA processing and mitochondrial RNA metabolism (0.81)
1. Transcriptional Regulation and Initiation
2. Splicing and RNA Processing
3. Mitochondrial RNA Metabolism and Translation
4. RNA Transport and Nuclear Export
5. Ubiquitin-Proteasome and RNA-Protein Complex Quality Control
6. Chromatin-Linked Transcription Modulators

ZNF557 (HepG2 cells)
Transcriptional regulation and RNA metabolic coordination via zinc finger proteins and RNA-associated complexes (0.88)
1. Transcriptional Regulation by Zinc Finger Proteins
2. General Transcription Machinery and RNA Polymerase Assembly
3. RNA Splicing, Processing, and Surveillance
4. rRNA Processing and Ribosome Biogenesis
5. Epigenetic and Chromatin-Level Regulation
6. Coupling with RNA Modification and Translation Control

Table 3. Functional enrichments among predicted regulatory target genes of nucleic

acid-binding proteins.

Functional enrichment analysis was conducted using the PANTHER database. This table

lists annotations that were statistically enriched in regulatory target genes predicted by a

deep-learning model trained on gene co-expression data. These annotations were not

enriched in the full set of co-expressed genes (Pearson's z-score > 2) that included the

nucleic acid-binding protein itself. 'Reference' refers to the total number of human genes.

'Analyzed' denotes the number of regulatory target genes examined. 'Expected' is the

number of genes expected to have the annotation by chance. 'Over/Under' indicates

whether the annotation is overrepresented or underrepresented. 'Fold Enrichment' is the

observed-to-expected ratio. 'Raw P-value' is the result of Fisher's exact test. 'FDR'

indicates the Benjamini–Hochberg false discovery rate.

AKAP8							
PANTHER Pathways							
PANTHER Pathways	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
Circadian clock system (P00015)	10	3	0.03	+	> 100	2.60E-06	4.16E-04
PANTHER GO-Slim Biological Process							
PANTHER GO-Slim Biological Process	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
circadian regulation of gene expression (GO:0032922)	16	3	0.05	+	65.55	1.20E-05	1.27E-02
response to light stimulus (GO:0009416)	21	3	0.06	+	49.94	2.82E-05	1.99E-02
circadian rhythm (GO:0007623)	21	3	0.06	+	49.94	2.82E-05	1.49E-02
rhythmic process (GO:0048511)	24	3	0.07	+	43.7	4.27E-05	1.81E-02
GO cellular component complete							
GO cellular component complete	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
Cry-Per complex (GO:1990512)	3	2	0.01	+	> 100	2.40E-05	4.77E-02
PKM							
PANTHER Protein Class							
PANTHER Protein Class	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
aldolase (PC00044)	5	2	0.1	+	21.02	3.48E-03	5.68E-02
kinase (PC00137)	134	9	2.55	+	3.53	1.07E-03	2.62E-02
PANTHER GO-Slim Molecular Function							
PANTHER GO-Slim Molecular Function	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
enzyme inhibitor activity (GO:0004857)	107	8	2.04	+	3.93	1.01E-03	4.68E-02
kinase activity (GO:0016301)	562	23	10.7	+	2.15	7.08E-04	3.57E-02
PANTHER GO-Slim Biological Process							
PANTHER GO-Slim Biological Process	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
endoplasmic reticulum tubular network organization (GO:0071786)	12	3	0.23	+	13.14	1.32E-03	3.68E-02
regulation of cellular component biogenesis (GO:0044087)	139	10	2.65	+	3.78	3.31E-04	1.05E-02

Acknowledgements

741 Supercomputing resources were provided by the Human Genome Center, the Institute of

742 Medical Science, the University of Tokyo, and the ROIS National Institute of Genetics.

Author contributions

745 N.O. designed and conducted the research, and wrote and revised the manuscript.

Supplementary data

Supplementary data are available at

Conflict of interests

None declared.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI [16K00387], a start-up research grant from Osaka University, the KIOXIA Young Scientist Research Grant at Waseda University, and the Grant for Basic Science Research Projects from the Sumitomo Foundation. All grants were awarded to N.O.

Supplementary Note and Tables

Gene co-expression–based predictions align with ChIP-seq–defined DNA-binding targets

High-ranking predicted binding regions frequently coincide with ChIP-seq peaks, particularly in promoter–enhancer intervals. Deep learning–derived DNA-binding site predictions based on gene co-expression data demonstrate substantial concordance with ChIP-seq data, particularly when broader genomic regions encompassing promoters and putative enhancers are considered. When overlap was assessed within ±100 kb of transcription start sites (TSSs)—a window supported by prior enhancer–gene mapping studies [19,75]—the mean and median overlap ratios reached 0.69 and 0.76, respectively (Ratio A2 / B2; Supplementary Tables S2 and S3). In contrast, overlap was lower for promoter regions alone (±5 kb), with corresponding mean and median values of 0.33 and 0.34. Some DNA-binding proteins, including FBXL19, POGK, SPEN, THAP7, ZNF142, and ZNF800, showed notably high promoter-region overlap ratios (>0.5), indicating potential direct regulatory activity. Nevertheless, not all ChIP-seq peaks necessarily reflect functionally relevant transcriptional binding events. Consequently, model-inferred high-scoring sites do not universally coincide with experimentally determined binding regions, underscoring the complexity and context dependence of regulatory interactions.

Context-specific predictions reveal known and emerging functional roles of NABPs

The deep learning–based predictions successfully recapitulate both established and potentially novel biological functions of NABPs. For example, AKAP8—previously annotated in UniProt with canonical cell cycle functions (e.g., "GO:0000278 mitotic cell cycle" and "GO:0044839 cell cycle G2/M phase transition")—was also predicted to be associated with circadian clock regulation in this study. Supporting literature retrieved from PubMed indicates a potential role for AKAP8 in circadian rhythm control: one study identified AKAP8 as highly co-expressed with known clock genes like *CRY1*, while another reported its involvement in melatonin biosynthesis, a process disrupted in individuals with autism spectrum disorders [76,77].

PKM (pyruvate kinase M) was functionally linked to glycolytic regulation, consistent with literature-based protein–protein interaction data, including its experimentally validated interaction with ALDOA in both human and mouse systems [39,40]. However, co-expression analysis in our dataset revealed a notable context-specific pattern: ALDOA and ALDOB were not significantly co-expressed with PKM (Pearson's *z*-score < 2), whereas ALDOC emerged as a strongly co-expressed gene prior to deep learning model training. This result suggests that ALDOC may serve as a context-dependent effector of PKM-regulated glycolysis, particularly in hematopoietic settings such as K562 cells. Indeed, ALDOC was not only among the predicted regulatory targets of PKM, but also appeared in gene sets annotated with hypoxia-responsive functions involving HIF1A. Moreover, transcriptomic data from the Human Protein Atlas confirmed that both ALDOA and ALDOC are expressed in K562 cells, reinforcing the possibility that PKM may selectively regulate ALDOC in specific cellular contexts despite the known biochemical role of ALDOA in glycolysis. These findings exemplify

how deep learning–guided predictions can refine our understanding of tissue-specific

metabolic wiring and highlight functional divergence within enzyme families.

Supplementary Table S1. Co-expression–based selection enhances the

importance scores of NABP binding sites.

Binding sites identified through gene co-expression exhibited higher absolute

DeepLIFT contribution scores compared to those selected solely from ChIP-seq data

without co-expression information. This trend was observed for both DNA-binding

proteins in HepG2 cells and RNA-binding proteins in K562 cells. For each group, the

table reports the mean and median of positive and negative contribution scores. These

findings suggest that gene co-expression–based selection prioritizes functionally

impactful regulatory interactions.

DNA-binding proteins (HepG2 cell)			
Co-expression–based		Non-co-expression–based	
Plus score		Plus score	
Mean	0.005673	Mean	0.003391
Median	0.003829	Median	0.001893
Minus score		Minus score	
Mean	-0.017000	Mean	-0.002502
Median	-0.002286	Median	-0.001583

RNA-binding proteins (K562 cell)			
Co-expression–based		Non-co-expression–based	
Plus score		Plus score	
Mean	0.004916	Mean	0.002962
Median	0.002422	Median	0.001689
Minus score		Minus score	
Mean	-0.011987	Mean	-0.002485
Median	-0.001521	Median	-0.001414

Supplementary Table S2. Enrichment of high-contribution promoter

regions with experimentally validated DBP binding sites. (See

Supplementary Excel file)

This table summarizes the proportion of promoter regions (±5 kb from transcription

start sites) that were both predicted to be functionally important by DeepLIFT

contribution scores and experimentally supported by ChIP-seq data for the same DNA-

binding proteins (DBPs). Overlap ratios were calculated across four thresholds—two

based on rank and two on absolute score—to assess the robustness of the predictions.

To classify high and low contributions, multiple thresholds were applied to the 1,310

DNA-binding proteins analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank $\leq$ 20 or $\geq$ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank $\leq$ 50 or $\geq$ 1,260)

Score-based thresholds:

(C1 and D1) contribution score $\geq$ 0.01 or $\leq$ –0.01

(C2 and D2) contribution score $\geq$ 0.001 or $\leq$ –0.001

Supplementary Table S3. Enrichment of high-contribution promoter

and enhancer regions with ChIP-seq–validated DBP binding sites.

This table presents the overlap ratios between predicted high-contribution regions

(defined within ±100 kb of transcription start sites, including distal enhancers) and ChIP-

seq binding sites for corresponding DBPs. High concordance between model-based

predictions and ChIP-seq peaks across multiple thresholds highlights the reliability of the

contribution scoring method in identifying functionally relevant regulatory sites.

To classify high and low contributions, the following thresholds were applied to

the 1,310 DNA-binding proteins analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank $\leq$ 20 or $\geq$ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank $\leq$ 50 or $\geq$ 1,260)

Score-based thresholds:

(C1 and D1) contribution score ≥ 0.01 or ≤ -0.01

(C2 and D2) contribution score ≥ 0.001 or ≤ -0.001

References

1　　Belz, G. T. & Nutt, S. L. Transcriptional programming of the dendritic cell network.
Nature reviews. Immunology **12**, 101-113, doi:10.1038/nri3149 (2012).

2　　Miller, J. C. *et al.* Deciphering the transcriptional network of the dendritic cell lineage.
Nature immunology **13**, 888-899, doi:10.1038/ni.2370 (2012).

3　　Ravasi, T. *et al.* An atlas of combinatorial transcriptional regulation in mouse and man.
Cell **140**, 744-752, doi:10.1016/j.cell.2010.01.044 (2010).

4　　Gertz, J. *et al.* Distinct properties of cell-type-specific and shared transcription factor
binding sites. *Molecular cell* **52**, 25-36, doi:10.1016/j.molcel.2013.08.037 (2013).

5　　Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650-665,
doi:10.1016/j.cell.2018.01.029 (2018).

6　　Vaquerizas, J. M., Kummerfeld, S. K., Teichmann, S. A. & Luscombe, N. M. A census of
human transcription factors: function, expression and evolution. *Nature reviews. Genetics*
10, 252-263, doi:10.1038/nrg2538 (2009).

7　　Pierson, E. *et al.* Sharing and Specificity of Co-expression Networks across 35 Human
Tissues. *PLoS Comput Biol* **11**, e1004220, doi:10.1371/journal.pcbi.1004220 (2015).

8　　Sonawane, A. R. *et al.* Understanding Tissue-Specific Gene Regulation. *Cell Rep* **21**,
1077-1088, doi:10.1016/j.celrep.2017.10.001 (2017).

9　　Saha, A. *et al.* Co-expression networks reveal the tissue-specific regulation of transcription
and splicing. *Genome research* **27**, 1843-1858, doi:10.1101/gr.216721.116 (2017).

10　　Boyle, A. P. *et al.* Comparative analysis of regulatory information and circuits across
distant species. *Nature* **512**, 453-456, doi:10.1038/nature13668 (2014).

11　　Ray, D. *et al.* RNA-binding proteins that lack canonical RNA-binding domains are rarely
sequence-specific. *Sci Rep* **13**, 5238, doi:10.1038/s41598-023-32245-9 (2023).

12　　Zeke, A. *et al.* Deep structural insights into RNA-binding disordered protein regions.
Wiley Interdiscip Rev RNA **13**, e1714, doi:10.1002/wrna.1714 (2022).

13　　Zaharias, S. *et al.* Intrinsically disordered electronegative clusters improve stability and

binding specificity of RNA-binding proteins. *J Biol Chem* **297**, 100945,
doi:10.1016/j.jbc.2021.100945 (2021).

14   Kribelbauer, J. F., Rastogi, C., Bussemaker, H. J. & Mann, R. S. Low-Affinity Binding Sites
and the Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev Biol*
35, 357-379, doi:10.1146/annurev-cellbio-100617-062719 (2019).

15   Neph, S. *et al.* Circuitry and dynamics of human transcription factor regulatory networks.
Cell **150**, 1274-1286, doi:10.1016/j.cell.2012.04.040 (2012).

16   Marbach, D. *et al.* Tissue-specific regulatory circuits reveal variable modular
perturbations across complex diseases. *Nature methods* **13**, 366-370,
doi:10.1038/nmeth.3799 (2016).

17   Fagny, M. *et al.* Exploring regulation in tissues with eQTL networks. *Proc Natl Acad Sci*
U S A **114**, E7841-E7850, doi:10.1073/pnas.1707375114 (2017).

18   Agarwal, V. *et al.* Massively parallel characterization of transcriptional regulatory elements.
Nature, doi:10.1038/s41586-024-08430-9 (2025).

19   Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-
range interactions. *Nature methods* **18**, 1196-1203, doi:10.1038/s41592-021-01252-x
(2021).

20   Tasaki, S., Gaiteri, C., Mostafavi, S. & Wang, Y. Deep learning decodes the principles of
differential gene expression. *Nat Mach Intell* **2**, 376-386, doi:10.1038/s42256-020-0201-
6 (2020).

21   Osato, N. & Hamada, M. Systematic discovery of directional regulatory motifs associated
with human insulator sites. *bioRxiv* (2025).

22   Obayashi, T., Kodate, S., Hibara, H., Kagaya, Y. & Kinoshita, K. COXPRESdb v8: an
animal gene coexpression database navigating from a global view to detailed investigations.
Nucleic acids research **51**, D80-D87, doi:10.1093/nar/gkac983 (2023).

23   Yang, S. *et al.* COEXPEDIA: exploring biomedical hypotheses via co-expressions
associated with medical subject headings (MeSH). *Nucleic acids research* **45**, D389-D396,
doi:10.1093/nar/gkw868 (2017).

24   Zhu, Q. *et al.* Targeted exploration and analysis of large cross-platform human
transcriptomic compendia. *Nature methods* **12**, 211-214, 213 p following 214,
doi:10.1038/nmeth.3249 (2015).

25   Miller, H. E. & Bishop, A. J. R. Correlation AnalyzeR: functional predictions from gene
co-expression correlations. *BMC Bioinformatics* **22**, 206, doi:10.1186/s12859-021-
04130-7 (2021).

26   Raina, P. *et al.* GeneFriends: gene co-expression databases and tools for humans and
model organisms. *Nucleic acids research* **51**, D145-D158, doi:10.1093/nar/gkac1031

(2023).

27   Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global
discovery of conserved genetic modules. *Science (New York, N.Y.)* **302**, 249-255,
doi:10.1126/science.1087447 (2003).

28   Lemoine, G. G., Scott-Boyer, M. P., Ambroise, B., Perin, O. & Droit, A. GWENA: gene
co-expression networks analysis and extended modules characterization in a single
Bioconductor package. *BMC Bioinformatics* **22**, 267, doi:10.1186/s12859-021-04179-4
(2021).

29   Warde-Farley, D. *et al.* The GeneMANIA prediction server: biological network integration
for gene prioritization and predicting gene function. *Nucleic acids research* **38**, W214-220,
doi:10.1093/nar/gkq537 (2010).

30   Wong, A. K., Krishnan, A. & Troyanskaya, O. G. GIANT 2.0: genome-scale integrated
analysis of gene networks in tissues. *Nucleic acids research* **46**, W65-W70,
doi:10.1093/nar/gky408 (2018).

31   UniProt, C. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic acids*
research **53**, D609-D617, doi:10.1093/nar/gkae1010 (2025).

32   Tak Leung, R. W., Jiang, X., Chu, K. H. & Qin, J. ENPD - A Database of Eukaryotic
Nucleic Acid Binding Proteins: Linking Gene Regulations to Proteins. *Nucleic acids*
research **47**, D322-D329, doi:10.1093/nar/gky1112 (2019).

33   Shrikumar, A., Greenside, P. & Kundaje, A. in *Proceedings of the 34th International*
Conference on Machine Learning Vol. 70   (eds Precup Doina & Teh Yee Whye) 3145--
3153 (PMLR, Proceedings of Machine Learning Research, 2017).

34   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nature genetics* **25**, 25-29, doi:10.1038/75556 (2000).

35   Gene Ontology, C. *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**,
doi:10.1093/genetics/iyad031 (2023).

36   Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all.
Protein Sci **31**, 8-22, doi:10.1002/pro.4218 (2022).

37   Hu, M. *et al.* Evaluation of large language models for discovery of gene set function.
Nature methods **22**, 82-91, doi:10.1038/s41592-024-02525-x (2025).

38   Farshadi, E., van der Horst, G. T. J. & Chaves, I. Molecular Links between the Circadian
Clock and the Cell Cycle. *J Mol Biol* **432**, 3515-3524, doi:10.1016/j.jmb.2020.04.003
(2020).

39   Snel, B., Lehmann, G., Bork, P. & Huynen, M. A. STRING: a web-server to retrieve and
display the repeatedly occurring neighbourhood of a gene. *Nucleic acids research* **28**,
3442-3444, doi:10.1093/nar/28.18.3442 (2000).

40   Szklarczyk, D. *et al.* The STRING database in 2023: protein-protein association networks
and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids*
research **51**, D638-D646, doi:10.1093/nar/gkac1000 (2023).

41   Tian, W. *et al.* Bioinformatics analysis of the role of aldolase A in tumor prognosis and
immunity. *Sci Rep* **12**, 11632, doi:10.1038/s41598-022-15866-4 (2022).

42   Zhang, F. *et al.* Elevated transcriptional levels of aldolase A (ALDOA) associates with cell
cycle-related genes in patients with NSCLC and several solid tumors. *BioData Min* **10**, 6,
doi:10.1186/s13040-016-0122-4 (2017).

43   Zlacka, J. & Zeman, M. Glycolysis under Circadian Control. *Int J Mol Sci* **22**,
doi:10.3390/ijms222413666 (2021).

44   Yang, Y., Pu, J. & Yang, Y. Glycolysis and chemoresistance in acute myeloid leukemia.
Heliyon **10**, e35721, doi:10.1016/j.heliyon.2024.e35721 (2024).

45   Puram, R. V. *et al.* Core Circadian Clock Genes Regulate Leukemia Stem Cells in AML.
Cell **165**, 303-316, doi:10.1016/j.cell.2016.03.015 (2016).

46   Lee, Y. Roles of circadian clocks in cancer pathogenesis and treatment. *Exp Mol Med* **53**,
1529-1538, doi:10.1038/s12276-021-00681-0 (2021).

47   Liberti, M. V. & Locasale, J. W. The Warburg Effect: How Does it Benefit Cancer Cells?
Trends Biochem Sci **41**, 211-218, doi:10.1016/j.tibs.2015.12.001 (2016).

48   Vander Heiden, M. G., Cantley, L. C. & Thompson, C. B. Understanding the Warburg
effect: the metabolic requirements of cell proliferation. *Science (New York, N.Y.)* **324**,
1029-1033, doi:10.1126/science.1160809 (2009).

49   Semenza, G. L. Hypoxia-inducible factors: mediators of cancer progression and targets
for cancer therapy. *Trends Pharmacol Sci* **33**, 207-214, doi:10.1016/j.tips.2012.01.005
(2012).

50   Keith, B., Johnson, R. S. & Simon, M. C. HIF1alpha and HIF2alpha: sibling rivalry in
hypoxic tumour growth and progression. *Nat Rev Cancer* **12**, 9-22, doi:10.1038/nrc3183
(2011).

51   Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science (New*
York, N.Y.) **347**, 1260419, doi:10.1126/science.1260419 (2015).

52   Israelsen, W. J. & Vander Heiden, M. G. Pyruvate kinase: Function, regulation and role in
cancer. *Semin Cell Dev Biol* **43**, 43-51, doi:10.1016/j.semcdb.2015.08.004 (2015).

53   Bluemlein, K. *et al.* No evidence for a shift in pyruvate kinase PKM1 to PKM2 expression
during tumorigenesis. *Oncotarget* **2**, 393-400, doi:10.18632/oncotarget.278 (2011).

54   Wang, X. & Liu, L. AKAP8 interacting with DDX5 to regulate R-loop balance involves in
lung carcinoma cell growth. *bioRxiv*, doi:10.1101/2023.11.28.568985 (2024).

55   Shirai, Y., Suwa, T., Kobayashi, M., Koyasu, S. & Harada, H. DDX5 enhances HIF‐1

activity by promoting the interaction of HIF‑1α with HIF‑1β and recruiting the
resulting heterodimer to its target gene loci. *Biology of the Cell* **116**,
doi:10.1111/boc.202300077 (2023).

56   Leszczynska, K. B. *et al.* Hypoxia‑mediated regulation of DDX5 through decreased
chromatin accessibility and post‑translational targeting restricts R‑loop accumulation.
Molecular Oncology **17**, 1173-1191, doi:10.1002/1878-0261.13431 (2023).

57   Farshadi, E., van der Horst, G. T. J. & Chaves, I. Molecular Links between the Circadian
Clock and the Cell Cycle. *Journal of Molecular Biology* **432**, 3515-3524,
doi:10.1016/j.jmb.2020.04.003 (2020).

58   Ran, J. *et al.* Hypoxia regulates glycolysis through the HIF-1α/BMAL1/ALDOC axis to
reduce oxaliplatin sensitivity in colorectal cancer. *Journal of Cancer* **16**, 2503-2515,
doi:10.7150/jca.108582 (2025).

59   De Vitis, C. *et al.* ALDOC- and ENO2- driven glucose metabolism sustains 3D tumor
spheroids growth regardless of nutrient environmental conditions: a multi-omics analysis.
Journal of Experimental & Clinical Cancer Research **42**, doi:10.1186/s13046-023-02641-
0 (2023).

60   Minchenko, O. H., Garmash, I. A. & Minchenko, D. O. Inhibition of IRE1 modifies
hypoxic regulation of G6PD, GPI, TKT, TALDO1, PGLS and RPIA genes expression in
U87   glioma   cells.   *The   Ukrainian   Biochemical   Journal* **89**,   38-49,
doi:10.15407/ubj89.01.038 (2017).

61   Obeidat, N. *et al.* Effects of cyclic acute and chronic hypoxia on the expression levels of
metabolism related genes in a pancreatic cancer cell line. *Biomedical Reports* **17**,
doi:10.3892/br.2022.1564 (2022).

62   Ladomery, M. Multifunctional proteins suggest connections between transcriptional and
post-transcriptional   processes.   *Bioessays* **19**,   903-909,   doi:10.1002/bies.950191010
(1997).

63   Kim, S., Yu, N. K. & Kaang, B. K. CTCF as a multifunctional protein in genome regulation
and gene expression. *Exp Mol Med* **47**, e166, doi:10.1038/emm.2015.33 (2015).

64   Osato, N. Characteristics of functional enrichment and gene expression level of human
putative transcriptional target genes. *BMC Genomics* **19**, 957, doi:10.1186/s12864-017-
4339-5 (2018).

65   Burley, S. K. & Kamada, K. Transcription factor complexes. *Curr Opin Struct Biol* **12**, 225-
230, doi:10.1016/s0959-440x(02)00314-7 (2002).

66   Goos, H. *et al.* Human transcription factor protein interaction networks. *Nat Commun* **13**,
766, doi:10.1038/s41467-022-28341-5 (2022).

67   Berenson, A. *et al.* Paired yeast one-hybrid assays to detect DNA-binding cooperativity

and antagonism across transcription factors. *Nat Commun* **14**, 6570, doi:10.1038/s41467-
023-42445-6 (2023).

68   Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic*
acids research **41**, D991-995, doi:10.1093/nar/gks1193 (2013).

69   Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene
expression and hybridization array data repository. *Nucleic acids research* **30**, 207-210,
doi:10.1093/nar/30.1.207 (2002).

70   Kolmykov, S. *et al.* GTRD: an integrated view of transcription regulation. *Nucleic acids*
research **49**, D104-D111, doi:10.1093/nar/gkaa1057 (2021).

71   Hinrichs, A. S. *et al.* The UCSC Genome Browser Database: update 2006. *Nucleic acids*
research **34**, D590-598, doi:10.1093/nar/gkj144 (2006).

72   Mudge, J. M. *et al.* GENCODE 2025: reference gene annotation for human and mouse.
Nucleic acids research **53**, D966-D975, doi:10.1093/nar/gkae1078 (2025).

73   Van Nostrand, E. L. *et al.* A large-scale binding and functional map of human RNA-
binding proteins. *Nature* **583**, 711-719, doi:10.1038/s41586-020-2077-3 (2020).

74   Young Cheon, N., Kim, S. & Yil Lee, J. in *Gene Expression*   Ch. Chapter 3,
(IntechOpen, 2022).

75   Gasperini, M. *et al.* A Genome-wide Framework for Mapping Gene Regulation via
Cellular Genetic Screens. *Cell* **176**, 1516, doi:10.1016/j.cell.2019.02.027 (2019).

76   Tan, X. *et al.* Novel Insights into the Circadian Rhythms Based on Long Noncoding and
Circular RNA Profiling. *Int J Mol Sci* **25**, doi:10.3390/ijms25021161 (2024).

77   Poelmans, G., Franke, B., Pauls, D. L., Glennon, J. C. & Buitelaar, J. K. AKAPs integrate
genetic findings for autism spectrum disorders. *Transl Psychiatry* **3**, e270,
doi:10.1038/tp.2013.48 (2013).

1052