Title

Cell type–specific functions of nucleic acid-binding proteins revealed by

deep learning on co-expression networks

Naoki Osato [1*] and Kengo Sato [1]

[1] School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan.

E-mail: naokiosato11@gmail.com

*Corresponding author.

Abstract

Nucleic acid-binding proteins (NABPs) play central roles in gene regulation, yet their functional targets and regulatory programs remain incompletely characterized due to the limited scope and context specificity of experimental binding assays. Here, we present a deep learning framework that integrates gene co-expression–derived interactions with contribution-based model interpretation to infer NABP regulatory influence across diverse cellular contexts, without relying on predefined binding motifs or direct binding evidence.

Replacing low-informative binding-based features with co-expression–derived interactions significantly improved gene expression prediction accuracy. Model-inferred regulatory targets showed strong and reproducible concordance with independent ChIP-seq and eCLIP datasets, exceeding random expectations across multiple genomic regions and threshold definitions. Functional enrichment and gene set enrichment analyses revealed coherent, cell type–specific regulatory programs, including cancer-associated pathways in K562 cells and differentiation-related processes in neural progenitor cells.

Notably, we demonstrate that DeepLIFT-derived contribution scores capture relative regulatory importance in a background-dependent but biologically robust manner, enabling systematic identification of context-dependent NABP regulatory roles. Together, this framework provides a scalable strategy for functional annotation of NABPs and highlights the utility of combining expression-driven inference with interpretable deep learning to dissect gene regulatory architectures at scale.

Keywords

Computational biology, Deep learning, Regulation of gene expression, DNA/RNA-binding proteins, Protein function prediction, Reactome, ChatGPT, GSEA

Introduction

Understanding the cell type–specific regulatory functions of nucleic acid-binding proteins (NABPs), including both DNA-binding proteins (DBPs) and RNA-binding proteins (RBPs), remains a fundamental challenge in molecular biology [1-10]. Although experimental assays such as ChIP-seq and ChIA-PET can map DBP binding sites, they require high-quality antibodies and are not easily scalable across large sets of proteins or conditions. The challenge is even greater for RBPs, many of which lack well-defined binding motifs due to their reliance on intrinsically disordered regions (IDRs) for RNA recognition, rather than sequence-specific contacts [11-13]. Interpreting low-affinity DBP binding, which contributes to enhancer activity and paralog-specific regulation, is particularly difficult—especially in the context of phase-separated transcriptional condensates [14]. Conventional methods like eQTL mapping or DNase-seq require prior motif knowledge and struggle to resolve such indirect or context-dependent interactions [15-17]. While massively parallel reporter assays (MPRAs) and large language model–based architectures such as Enformer have advanced our ability to relate sequence to function [18,19], they do not reveal which NABPs mediate the predicted regulatory effects. Thus, novel strategies are required to infer NABP–target relationships and functional roles, particularly when motif information is unavailable or direct binding evidence is missing.

Deep learning has substantially improved the prediction of gene expression by modeling regulatory inputs from NABPs, including transcription factors and microRNAs

(miRNAs) [20]. For example, models incorporating ChIP-seq–derived NABP binding sites

in promoter regions and predicted miRNA-binding sites in transcripts have achieved a

high correlation ($r = 0.82$) between predicted and observed expression levels. In a

previous study, we extended this approach by incorporating binding site data for DNA-

binding proteins within 1 Mb of transcription start sites (TSSs) in open chromatin, while

excluding miRNAs and RBPs. To facilitate accurate computation of importance scores

(also known as contribution scores) and to improve model interpretability, we unified the

input representation by excluding a previous input layer for RNA-binding proteins and

miRNAs [21]. While this modification slightly reduced predictive accuracy ($r = 0.70$), it

enabled broader coverage of distal regulatory elements and laid the foundation for

downstream functional analysis based on importance scores. These scores are later

leveraged to identify regulatory targets and infer NABP functions.

Gene co-expression analysis, a well-established method since the 2000s, offers a

complementary perspective by identifying genes with coordinated expression across

biological contexts [22-30]. These patterns, derived from transcriptomic datasets across

diverse cell types, can capture both direct and indirect regulatory associations—including

those involving NABPs with uncertain or difficult-to-characterize binding features. In

this study, we integrated gene co-expression data into a deep learning framework to assess

its utility for predicting gene expression and identifying context-specific regulatory

targets of NABPs. This combination provides a flexible means to infer functional

relationships even when direct binding data are lacking.

To test the effectiveness of this approach, we substituted the binding site features

of ~300 NABPs that contributed little to previous deep learning predictions with gene co-

expression–based interactions. The modified model successfully predicted gene

expression levels while assigning DeepLIFT-based importance scores to NABP–gene

interactions. Many predicted targets overlapped with ChIP-seq or eCLIP-validated

binding partners and showed higher importance scores than background interactions. In

this study, we use DeepLIFT to compute attribution values for each gene. We refer to

these values as DeepLIFT scores, also described as importance scores or contribution

scores in the context of feature attribution. Functional enrichment analyses of twelve

representative NABPs revealed consistent correspondence with known annotations in

UniProt [31], and several additional functions were supported by independent publications.

To enhance interpretability, we also applied a ChatGPT-based method to infer gene

functions and retrieve supporting literature. The results were consistent with enrichment-

based findings and highlighted biologically meaningful associations, highlighting cell

type–specific processes such as neural cell differentiation, as well as cancer-related

functions and pathways in neural progenitor and K562 cells.

This study provides a computational strategy to uncover the functional roles and

regulatory targets of NABPs without requiring experimentally defined binding sites. By

integrating gene co-expression profiles with deep learning, the framework captures both

direct and indirect interactions often missed by conventional methods. This approach

enables the systematic and scalable identification of NABP functions, providing a new

avenue for exploring transcriptional regulation in a cell type–specific manner.

Results

Gene co-expression features enhance expression prediction accuracy

Replacing low-contribution DBP inputs with gene co-expression data led to a

substantial increase in model performance (**Fig. 1**). In human foreskin fibroblasts

(HFFs), substituting 302 out of 1,310 DBPs with co-expression profiles improved the

correlation between predicted and actual gene expression from 0.70 to 0.80 (**Fig. 2a, b**).

These co-expression inputs included both NABP-encoding and non-NABP mRNAs, with

gene symbols for DNA- and RNA-binding proteins obtained from the ENPD (Eukaryotic

Nucleic acid Binding Protein) database[32], and co-expression profiles sourced from

COXPRESdb [22]. A subsequent experiment replaced 325 low-contribution DBPs with

co-expression interactions involving RBPs, resulting in a further increase in prediction

accuracy (r = 0.81; **Fig. 2c**). Collectively, these findings demonstrate that integrating co-

expression interactions into the model enhances its ability to capture gene regulatory

dynamics beyond direct binding site information.

Contribution scores derived from co-expression–based NABP–gene interactions

exhibited broader distributions and higher overall magnitudes than those obtained from

ChIP-seq–based inputs across all four cell types examined (Wilcoxon signed-rank test, p

$< 2.2 \times 10^{-16}$; **Fig. 2d** and **Supplementary Fig. S2c**; **Supplementary Table S1**). Each

pair of violin plots represents absolute contribution scores for co-expression–based sites

(right plot in each pair) and the corresponding ChIP-seq sites not included in that set (left

plot). **A**–**P** correspond to specific cell types and NABP classes as follows: **B**, DBP co-

expression–based sites in neural progenitor cells (NPCs); **A**, NPC ChIP-seq sites

excluding **B**. **D**, RBP co-expression–based sites in NPCs; **C**, NPC ChIP-seq sites

excluding **D**. **F**, DBP co-expression–based sites in HepG2 cells; **E**, HepG2 ChIP-seq sites

excluding **F**. **H**, RBP co-expression–based sites in K562 cells; **G**, K562 ChIP-seq sites

excluding **H**. **J**, DBP co-expression–based sites in HMEC cells; **I**, HMEC ChIP-seq sites

excluding **J**. **L**, RBP co-expression–based sites in HMEC cells; **K**, HMEC ChIP-seq sites

excluding **L**. **N**, DBP co-expression–based sites in human foreskin fibroblasts (HFFs); **M**,

HFF ChIP-seq sites excluding **N**. **P**, RBP co-expression–based sites in HFFs; **O**, HFF

ChIP-seq sites excluding **P**. Corresponding violin plots generated using an amplified

background (input × 2) are shown in **Supplementary Figure S2 c**. However, the nature

of this difference depended on the class of nucleic acid–binding protein. For DNA-

binding proteins (DBPs), contribution score distributions differed significantly between

ChIP-seq–derived and co-expression–derived regulatory sites, yet their mean and median

values were often comparable across cell types. This indicates that, for DBPs, co-

expression–based feature selection primarily reshapes the distributional structure of

contribution scores without uniformly shifting their central tendency. In contrast, for

RNA-binding proteins (RBPs), regulatory sites inferred from gene co-expression

consistently showed substantially higher DeepLIFT contribution scores than those

derived from experimental binding data alone. This pattern was observed across all

examined cell types, including K562, NPCs, HMECs, and HFFs, and was reflected by

marked increases in both mean and median contribution scores. These results suggest that

co-expression–based inference preferentially enriches for RBP-associated regulatory

interactions that contribute more directly to gene expression prediction, whereas

experimental binding data for RBPs may include a larger fraction of interactions that are

weakly coupled to transcriptomic output.   These observations are consistent with

previous reports showing that incorporating RNA-binding protein– and miRNA-

associated features improves gene expression prediction accuracy, as exemplified by the

DEcode framework [20]. Importantly, these trends were robust to DeepLIFT background

definition: comparable differences between ChIP-seq– and co-expression–derived targets

were observed under both background scaling schemes (input × 0.5 and input × 2). Thus,

while background scaling influences the sign and absolute scale of contribution scores, it

does not alter the fundamental distinction between binding-based and co-expression–

based regulatory inference.

To assess the influence of background selection on DeepLIFT-derived contribution

scores, we compared results obtained using a reduced background (input × 0.5; **Fig. 2e**)

and an amplified background (input × 2; **Fig. 2f**) for the same representative DNA- and

RNA-binding proteins. Under the reduced background, contribution scores tended to

show negative correlations with gene expression, whereas the amplified background

produced predominantly positive correlations. These patterns appeared approximately

mirrored around zero, consistent with DeepLIFT's definition as a relative contribution

measure with respect to the chosen reference. Importantly, despite this inversion in sign,

the relative ranking of genes and the overall cell type–specific structure of contribution

score distributions remained highly consistent across background settings. In both cases,

genes with larger-magnitude contribution scores were preferentially associated with

higher or lower expression levels depending on the background, indicating that

DeepLIFT captures relative regulatory importance rather than absolute activation or

repression. A conceptual model explaining how input definition and background scaling

give rise to these patterns is provided in **Supplementary Fig. S4**. Notably, background-

dependent differences were more pronounced for RNA-binding proteins (RBPs). For the

same representative RBPs, scatter plots generated using the reduced background (input ×

0.5) exhibited more distinct, partially separated distributions across the four cell types,

whereas under the amplified background (input × 2), data points from different cell types

overlapped more extensively. This suggests that the reduced-background configuration

enhances sensitivity to cell type–specific regulatory variation for RBPs. In contrast,

DNA-binding proteins (DBPs) generally showed stronger correlations with gene

expression under both background conditions, consistent with their more direct

transcriptional regulatory roles. Together, these results underscore the importance of

background-aware and protein class–specific interpretation of DeepLIFT contribution

scores.

Experimental binding data support model-inferred regulatory targets

To determine whether DeepLIFT-derived regulatory contribution scores reflect

biologically meaningful nucleic acid–binding protein (NABP) activity, we systematically

compared model-inferred regulatory targets with experimentally validated ChIP-seq and

eCLIP binding data across multiple genomic regions, threshold definitions, and cellular

contexts (**Fig. 3**). Experimentally validated binding sites are consistent with deep

learning–derived regulatory targets. Genes with ChIP-seq or eCLIP peaks for a given

NABP exhibited disproportionately high or low importance ranks, indicating that

experimentally confirmed binding sites are associated with stronger regulatory influence

in gene expression prediction models. This relationship supports the biological validity

of DeepLIFT-based contribution scores, which quantify the importance of specific

NABPs in driving gene expression changes.

To evaluate this association, we analyzed ChIP-seq data for 32 DBPs in HepG2

cells and eCLIP data for 29 RBPs in K562 cells—two cell lines with the most

comprehensive publicly available NABP binding site datasets from ENCODE and related

studies. A gene was considered associated with an NABP if a ChIP-seq peak was located

within ±5 kb of its transcription start site (TSS) or if an eCLIP peak was mapped to its

transcribed region. As illustrated in **Fig. 3a** and **Supplementary Fig. S1**, NABPs with

experimentally validated binding sites consistently showed higher absolute contribution

score values across their predicted target genes, reinforcing their regulatory relevance and

corresponding with prior interpretations of DeepLIFT scores [20,33].

Model-derived importance scores reliably predict DBP binding sites, as evidenced

by significant overlap with ChIP-seq peaks (**Fig. 3b, Supplementary Table S2** and **S3**).

To evaluate this correspondence, we analyzed ChIP-seq data for 32 DNA-binding

proteins (DBPs) and quantified how frequently high-scoring gene regions—defined by

DeepLIFT contribution scores—coincided with experimental binding evidence. Overlap

ratios were calculated as the proportion of high-contribution regions that intersected with

ChIP-seq peaks, considering both promoter-proximal (±5 kb) and distal (±100 kb) regions

around transcription start sites (TSSs). To account for the functional heterogeneity and

abundance of ChIP-seq peaks, we applied a precision-at-K approach, focusing on the top-

ranked or top-scoring gene regions most likely to reflect regulatory relevance. Four

thresholds were used to define high-contribution regions: two based on contribution score

rank (top/bottom 20 or 50) and two based on absolute score magnitude (≥ 0.01 or $\leq -$

0.01; ≥ 0.001 or ≤ -0.001). A control analysis using randomly selected gene regions of

matched size was conducted in parallel. In all tested conditions, the overlap between

ChIP-seq peaks and high-contribution regions exceeded that of the random control,

supporting the model's ability to prioritize functionally relevant regulatory sites.

Similarly, as illustrated in **Fig. 3c** and **Supplementary Table S4**, the model-

derived importance scores consistently prioritize biologically relevant RBP–target

interactions, as evidenced by the cumulative overlap between high-contribution gene

regions and eCLIP peaks, which was substantially greater than that observed in the

random control. We assessed this by analyzing eCLIP data for 29 RNA-binding proteins

(RBPs) in K562 cells, comparing high-contribution regions—defined using the same

thresholds as for DBPs—to eCLIP peaks mapped to transcribed regions (from TSS to

TES). Collectively, these results demonstrate that the model effectively captures

functionally meaningful NABP–target associations for both DNA- and RNA-binding

proteins.

Significant differences in the predicted regulatory targets of several NABPs were

observed between K562 and HepG2 cells based on DeepLIFT contribution scores (**Fig.**

3d). To assess cell type–specific regulatory activity, we compared the DeepLIFT scores

of genes harboring NABP binding sites, as defined by ChIP-seq or eCLIP. Because our

deep learning model identifies target candidates from NABP-associated co-expressed

gene sets, not all predicted targets necessarily reflect direct binding; rather, they represent

genes that participate in shared or downstream regulatory programs. RPLP0 exhibited

nearly identical score distributions across the two cell types, whereas ETF1, NELFE, and

CAMTA2 showed marked divergence. These differences indicate that the model captures

and adjusts NABP-dependent regulatory relationships in a cell type–specific manner.

Deep learning uncovers functionally enriched NABP target genes

beyond co-expression networks

Deep learning predictions revealed that nucleic acid–binding proteins (NABPs)

possess regulatory target genes with coherent and biologically specific functional

enrichment, extending well beyond patterns detectable from co-expression alone (**Tables**

1, 2 and **Supplementary Note**). Across twelve representative NABPs [34,35], predicted

target sets consistently showed significant enrichment for biologically related Gene

Ontology (GO) terms, whereas randomly sampled gene sets of matched size displayed no

such enrichment (**Fig. 4**). Notably, several enriched annotations were absent from the

broader pools of co-expressed genes (Pearson's *z*-score > 2), indicating that the model

selectively extracts biologically relevant signals from large co-expression neighborhoods.

Functional enrichment across GO Slim, Reactome, and PANTHER Pathways further

confirmed this specificity (**Table 1**) [36]. A ChatGPT-based gene set interpretation method

independently supported these results, providing functional summaries and literature

consistent with GO enrichment and UniProt annotations [31] (**Table 2** and **Supplementary**

Note) [37].

Cell type–dependent functional specificity revealed by deep learning–

refined NABP targets

Deep learning refinement further exposed cell type–specific biological functions

associated with regulatory targets of several RBPs. For example, the predicted targets of

PKM showed distinct pathway enrichments between K562 cancer cells and neural

progenitor cells (NPCs). Pathways such as *Pentose phosphate pathway*, *Glycolysis*, and

Hypoxia response mediated by HIF activation were significantly enriched in K562 but

not in NPCs, whereas *Integrin signaling pathway*, *Glycogen metabolism*, and *Glutamate*

neurotransmitter release cycle were specifically enriched in NPCs using a background

defined as 0.5× the original input feature values[38-47] (**Fig. 5**, **Table 3**, **Supplementary**

Table S5). ChatGPT-based interpretation produced consistent functional assignments

(**Supplementary Note**). Notably, these cell type–dependent associations align with

known biology in cancer cells and NPCs.

Additional RBPs exhibited similar specificity. Predicted target genes of ILF2 were

enriched for *G2/M DNA replication checkpoint* in K562 cells but not in NPCs[48-50] ; targets

of EIF2S2 were associated with *Serine and glycine biosynthesis* in NPCs but not in

K562[51,52] ; and targets of DNAJC2 were enriched for the *Ubiquitin proteasome pathway*

in K562 but not in NPCs[53,54] (**Fig. 5**, **Table 3**, **Supplementary Table S5**). ChatGPT-

based inference again recapitulated these cell type–dependent functional patterns

(**Supplementary Note**), consistent with previous reports.

Together, these results demonstrate that the deep learning–based refinement of

NABP target predictions enables systematic detection of cell type–dependent functions,

revealing regulatory specificity not captured by co-expression alone.

285   Functional enrichment analysis of NABP target genes using
286   contribution score–based GSEA

Gene set enrichment analysis (GSEA) revealed robust, cell type–dependent functional

differences among NABP regulatory target genes, providing clearer biological insight

than threshold-based enrichment approaches. Rather than ranking genes by expression

levels, we ranked predicted target genes according to their DeepLIFT contribution scores,

enabling functional interpretation based on relative regulatory importance within each

cell type. This framework allows pathway enrichment to be assessed by shifts in the

positioning of functionally related genes along the ranked list, independent of absolute

score magnitude or sign. Using this contribution score–based GSEA, we identified

pronounced differences in PKM-associated pathway enrichment between neural

progenitor cells (NPCs) and K562 leukemia cells (**Fig. 6**). Notably, multiple Reactome

pathways exhibited large differences in normalized enrichment scores ($\Delta\text{NES} = \text{NES}_{\text{K562}}$

$-\ \text{NES}_{\text{NPC}}$), indicating substantial divergence in the inferred regulatory associations of

PKM across these two cellular contexts. For example, pathways related to synaptic

organization and neural functions, such as Protein–protein interactions at synapses,

showed strong enrichment differences between NPCs and K562, whereas pathways

associated with immune signaling and cancer-related processes were preferentially

enriched in K562 cells. These contrasting patterns were consistently captured across

multiple pathways ranked by ΔNES (**Table 4**). Importantly, because DeepLIFT

contribution scores depend on the choice of background reference, the absolute direction

of enrichment (positive or negative NES) may invert under different background

definitions. Accordingly, we focused on relative differences in NES between cell types

rather than the absolute sign of enrichment. Consistent pathway-level differences between

NPCs and K562 were observed under both background configurations (input × 0.5, **Fig.**

6; input × 2, **Supplementary Table S8**), supporting the robustness of cell type–specific

functional contrasts inferred from PKM contribution scores. Notably, the amplified

background configuration (input × 2) tended to yield a larger number of genes with non-

negligible contribution scores, resulting in stronger overall enrichment scores and higher

NES values across pathways (**Supplementary Tables S7** and **S9**). This effect reflects

increased sensitivity in detecting PKM-associated regulatory targets under the input × 2

background, while preserving the relative enrichment differences between cell types.

Together, these results demonstrate that GSEA based on deep learning–derived

contribution scores effectively captures cell type–specific NABP functional associations

that are not readily detected by expression-based or threshold-dependent analyses. By

emphasizing relative enrichment differences across cellular contexts, this approach

provides a robust framework for dissecting context-dependent regulatory programs and

linking NABPs to biologically meaningful pathways.

Discussion

These findings address a long-standing challenge in molecular biology: the

systematic functional characterization of nucleic acid-binding proteins (NABPs) across

diverse cellular contexts. While traditional methods rely heavily on experimental binding

data and known motifs, our approach utilizes gene co-expression and machine learning

to circumvent these limitations. By capturing both direct and indirect regulatory signals,

it broadens the scope of functional prediction for NABPs that lack conventional binding

site information.

Gene expression correlations alone can serve as a powerful basis for uncovering

the regulatory roles of nucleic acid-binding proteins (NABPs), even in the absence of

experimental binding data. Leveraging transcriptomic profiles from diverse public RNA-

seq datasets, our deep learning framework inferred regulatory targets of NABPs based

solely on co-expression patterns. Unlike binding site assays such as ChIP-seq or eCLIP—

which are technically demanding and limited by the availability of high-quality

antibodies—this computational approach is scalable and applicable across NABPs with

poorly characterized binding motifs. Functional enrichment analysis of the predicted

target genes revealed biologically meaningful roles, many of which were consistent with

existing annotations in public databases. In addition, several previously unannotated

functional associations—not captured in UniProt or GO but supported by published

studies—were uncovered. GSEA further demonstrated that genes contributing to specific

cell type–associated pathways exhibited systematic shifts toward higher or lower

DeepLIFT scores, indicating differential regulatory influence across cellular contexts.

These findings were corroborated by a ChatGPT-assisted literature mining approach,

which independently recovered concordant biological themes and supporting references.

Together, these results underscore the biological relevance of NABP target predictions

derived from deep learning–based co-expression modeling and highlight the value of

integrating such predictions with AI-assisted interpretation. This strategy provides a

scalable framework for elucidating NABP functions and discovering previously

unrecognized regulatory mechanisms across diverse cell types.

Contribution scores derived from gene expression correlations capture both direct

and indirect regulatory interactions, offering insights beyond what ChIP-seq alone can

provide. While ChIP-seq detects thousands of DNA-binding events per protein, many of

these sites do not contribute directly to gene expression regulation and may reflect the

multifunctionality of transcription factors [55,56]. Consequently, ChIP-seq–derived binding

maps and DeepLIFT-based predictions show only partial overlap, underscoring both

methodological differences and the inherent complexity of regulatory networks. In this

study, gene co-expression analysis revealed fewer target genes than ChIP-seq but

identified subsets that are likely to be more functionally relevant. Our previous work on

insulator-associated factors linked such proteins to enhancer–promoter insulation and

chromatin organization, including phase separation and loop formation [21]. That study

incorporated eQTL data to anchor binding sites to expression outcomes. Here, by contrast,

we focused on NABPs with unknown motifs and excluded motif-based or chromatin

accessibility data such as DNase-seq or ATAC-seq, to assess the capacity of gene

expression correlations to reveal regulatory roles. Notably, the distribution of

contribution scores for co-expression–based inputs included both high and low values,

implying that gene expression patterns reflect not only direct binding but also indirect

regulation mediated through broader network effects. This highlights the value of

contribution scores as indicators of both proximal and distal regulatory influences, even

when binding motifs are unknown or experimental binding data are unavailable. The cell

type–specific differences in DeepLIFT contribution score distributions observed in **Fig.**

3d suggest that the regulatory roles of individual nucleic acid-binding proteins are not

fixed but are dynamically reshaped across cellular contexts. Notably, while some proteins

such as RPLP0 exhibited highly conserved contribution patterns between HepG2 and

K562 cells, others, including ETF1 and NELFE, displayed pronounced divergence,

indicating cell type–dependent shifts in their inferred regulatory influence. These findings

support the notion that NABPs participate in distinct regulatory programs depending on

cellular state and lineage, and that DeepLIFT-based analyses can sensitively capture such

context-dependent regulatory reorganization. This context dependence is particularly

relevant for interpreting NABP function in disease versus developmental states, where

subtle shifts in regulatory importance may precede overt transcriptional changes.

Functional roles of NABPs can be inferred even from small gene sets, provided

that the target genes exhibit shared or functionally related patterns. In this study, twelve

NABPs showed functional enrichments consistent with known annotations, supporting

the validity of the deep learning–based predictions. These enrichments were identified

using Fisher's exact test with false discovery rate (FDR) correction via the PANTHER

database, which typically requires at least 50 genes for robust statistical significance. For

proteins with fewer than 50 predicted targets, enrichment may be harder to detect

depending on the functional composition of the gene set. Relaxing the ranking thresholds

increased the number of target genes and enhanced statistical power. Alternatively, more

stringent thresholds could be retained if the contribution scores were sufficiently strong,

offering flexibility depending on analysis goals. Additional features—such as gene

expression levels or co-expression correlations—could further refine enrichment analyses,

particularly for borderline cases. Notably, enrichment analysis of both high- and low-

ranking targets revealed candidate novel functions for specific NABPs, suggesting that

genes with large-magnitude contribution scores are biologically informative. Moreover,

a complementary ChatGPT-based approach demonstrated utility in interpreting small

gene sets. Unlike standard enrichment tools, this method can infer biological roles even

when gene counts fall below conventional significance thresholds, provided that genes

with related functions are present. This capacity helps overcome statistical limitations and

broadens the interpretability of regulatory predictions in a cell type–specific context.

GSEA based on DeepLIFT scores provides a complementary view of pathway-level

regulation by assessing how genes associated with the same pathway are distributed along

the model's importance landscape, rather than relying on gene expression levels. This

approach revealed striking differences between normal and disease-associated cell types.

For example, genes enriched in neural progenitor cell (NPC) pathways tended to cluster

at higher or lower DeepLIFT scores in NPCs, whereas their score distributions shifted

substantially in K562 cells, indicating altered regulatory programs between the two cell

states. Such analyses offer a powerful means to uncover context-dependent regulatory

mechanisms across diverse cell types, conditions, and developmental stages.

Strategic use of co-expression data enhances functional inference, while

indiscriminate replacement can diminish biological relevance. The selective integration

of gene co-expression data improved target prediction and the functional interpretability

of the model. In contrast, replacing all input features with co-expression-derived

interactions led to a decrease in enriched annotations, and known functional roles were

no longer recovered. Excessive replacement of input features with gene co-expression

data further diminished the biological relevance of model predictions. When all NABP–

target inputs were substituted with co-expression-based interactions, the number of

enriched functional annotations declined, and previously identified functions from the

UniProt database were no longer recovered. This suggests that indiscriminate

replacement may cause the model to capture non-informative or biologically spurious

associations. A similar trend was observed in a prior control analysis, where substituting

true regulatory targets with randomly selected genes led to a gradual loss of enrichment

as randomness increased. Based on these observations, the normalized count of enriched

functional terms—adjusted by the number of predicted targets—may serve as a useful

proxy for evaluating the accuracy and biological validity of NABP–target predictions [57].

These findings also raise intriguing questions about how the deep learning model

prioritizes informative input features during training. Future work exploring the model's

internal representation dynamics—such as how it distinguishes signal from noise when

presented with diverse regulatory interactions—may provide valuable insights into the

mechanisms of transcriptional regulation and model interpretability. Investigating how

the model converges on biologically meaningful NABP–target relationships during

training could deepen our understanding of both gene regulation and model

interpretability.

Cell type specificity and cooperative binding dynamics may underlie the absence

of detectable regulatory signals for certain NABPs. While some NABPs showed no

high- or low-ranking target genes in HepG2 or K562 cells, such absence does not rule out

regulatory relevance. These proteins may exhibit cell type–specific regulatory activity

detectable only in other cellular contexts, where functional enrichment analysis could

reveal distinct biological roles. Moreover, many transcription factors and DNA-binding

proteins act in concert, forming multi-protein complexes that bind cooperatively to shared

promoters [58-60]. In such cases, individual components may not always yield strong

contribution scores due to their distributed regulatory influence across the complex. This

highlights the potential of co-binding site analysis as a complementary strategy to uncover

cooperative regulatory mechanisms and to disentangle the functional interplay of NABPs

within transcriptional networks.

Beyond characterizing cell type–specific regulatory programs, our framework also

suggests a potential application to preventive and health-oriented research. By

systematically scoring "cleaning" pathways—including proteostasis, DNA damage repair,

RNA quality control, immune surveillance, and metabolic stress responses—based on the

predicted regulatory influence of nucleic acid–binding proteins, it becomes possible to

estimate the integrity of cellular maintenance systems before overt disease onset. Such

pathway-level scores may capture early functional weakening of protective mechanisms

that are not yet reflected by pathological phenotypes or single-gene alterations. This

perspective aligns with observations from cancer-resistant species and long-lived

individuals, in which enhanced cellular maintenance, quality control, and stress

adaptation are thought to contribute to disease avoidance rather than treatment. In this

context, deviations in cleaning pathway scores could serve as quantitative indicators of

declining cellular robustness, potentially enabling risk stratification or early detection of

disease predisposition. While the present study focuses on neural progenitor and cancer

cell models, the same analytical framework could, in principle, be extended to

longitudinal human data or population-scale transcriptomic profiles, providing a

foundation for preventive or health-monitoring applications that complement

conventional disease-centered analyses.

This study establishes a scalable computational framework for uncovering the

functional roles of nucleic acid-binding proteins (NABPs) by integrating gene

expression–based associations with deep learning–derived contribution scores. By

leveraging co-expression–driven interactions rather than relying solely on experimentally

defined binding motifs, the approach enables systematic inference of both canonical and

previously uncharacterized regulatory relationships across cell types. Collectively, our

results demonstrate that expression-driven contribution patterns, when coupled with

pathway-level analyses and AI-assisted functional interpretation, can capture biologically

meaningful differences between normal and disease-associated cellular states. This

capability suggests that the framework may be particularly valuable for identifying early

shifts in regulatory and cellular maintenance pathways that precede overt disease

phenotypes, thereby complementing traditional binding-centric or mutation-focused

analyses. Although the present study focused on a representative subset of NABPs and

cell types, the framework is readily extensible to larger protein cohorts, additional cell

states, and diverse transcriptomic datasets. In this way, it provides a foundation for

systematic functional annotation of NABPs, including those with poorly characterized

roles, and offers a path toward integrating transcriptomic signals and AI-driven

predictions into large-scale functional genomics and disease-relevant research.

Methods

489 **Predictive modeling of gene expression across diverse cell types using**

490 **deep learning and NABP binding information**

491   To predict gene expression across multiple human cell types, we adapted a

492 previously established deep learning framework that integrates DNA-binding site data for

493 transcription factors and other regulatory proteins. In this implementation, regulatory

494 elements located both proximally and distally (up to ± 1 Mb) from transcription start sites

495 (TSSs) were included as input features (**Fig. 1**) [20,21]. To minimize the effects of cellular

496 heterogeneity in bulk tissue, RNA-seq datasets were selected at the individual cell-type

497 level. Publicly available RNA-seq data were retrieved for four functionally distinct cell

498 types: human foreskin fibroblasts (HFFs), mammary epithelial cells (HMECs), neural

499 progenitor cells (NPCs), and either HepG2 hepatoma cells or K562 leukemia cells,

500 depending on the class of regulatory protein analyzed [61,62]. Gene expression values

501 (FPKM) were computed using the RSeQC toolkit based on GENCODE v19 annotations,

502 and subsequently log$_2$-transformed after adding a small offset to avoid undefined values.

503 Following the approach described by Tasaki et al. [20,21], expression fold changes were

504 calculated relative to the median expression level across the four cell types. For each

505 model iteration, three common cell types (HFFs, HMECs, and NPCs) were used, and

506 either HepG2 (for DNA-binding protein analysis) or K562 (for RNA-binding protein

507 analysis) was included to match the availability of ChIP-seq and eCLIP validation

508 datasets. These cell types were chosen to represent diverse biological lineages and

509 regulatory contexts. For specific analyses requiring direct comparison of DeepLIFT

510 contribution scores across cell types—such as the evaluation of cell type–dependent

511 regulatory differences for the same nucleic acid-binding protein (e.g., **Fig. 3d**)—models

512 were trained using a four-cell-type configuration comprising HFFs, NPCs, HepG2, and

K562. This unified configuration enabled direct comparison of contribution score

distributions between HepG2 and K562 while maintaining a shared background of

common cell types.

Extended regulatory features and DeepLIFT-compatible model design

To capture distal regulatory signals beyond promoter regions, we integrated DNA-

binding site data from putative enhancers into a deep learning framework for gene

expression prediction. Building upon our earlier work [20,21], this model extends the

architecture originally proposed by Tasaki et al. (2020) [20,21], which focused on promoter-

proximal regions and included RNA-binding protein and miRNA inputs. In our previous

study, aimed at identifying insulator-associated DNA-binding proteins, we deliberately

excluded RNA-binding proteins and miRNAs to focus on DNA-centered regulatory

elements. Instead, we combined DNA-binding sites from both promoter (-2 kb to +1 kb)

and enhancer-like regions (±1 Mb from the transcription start site [TSS]) into a unified

input matrix, enabling DeepLIFT analysis under the high-accuracy "genomics-default"

mode. In this study, we use DeepLIFT to compute attribution values for each gene. We

refer to these values as DeepLIFT scores, also described as importance scores or

contribution scores in the context of feature attribution.

Binding site coordinates were obtained from the GTRD database (v20.06) [63], and

regulatory associations were inferred using cis-eQTLs from GTEx v8. A ChIP-seq peak

that overlapped a 100-bp window centered on an expression-associated SNP was

considered linked to the corresponding transcript. Trans-eQTLs were excluded, and

coordinates were converted to the hg19 assembly using UCSC LiftOver [64]. To limit

feature dimensionality while maximizing genomic coverage, we computed peak lengths

for each DNA-binding protein (DBP) in 100-bp bins across ±1 Mb around the TSS, then

retained the maximum peak value within each 10-kb window, yielding up to 200 enhancer

bins. These were concatenated with 30 promoter-region bins to form a 230-bin unified

input layer. This single-input-layer structure was necessary because DeepLIFT's

genomics-default configuration ("RevealCancel rule on dense layers and Rescale rule on

conv layers") does not support models with multiple input layers. Concatenating all DBP-

related features into one matrix thus maintained compatibility while enhancing

interpretability.

Model training followed the protocol described by Tasaki et al. and applied in our

prior work [20,21]. Specifically, 2,705 genes from chromosome 1 were used for testing,

while 22,251 genes from other chromosomes were randomly assigned to the training set

and 2,472 to the validation set. Models were implemented in TensorFlow 2.6.0 and

trained on an NVIDIA RTX 3080 GPU, typically completing in 2 to 4 hours. Feature

importance was assessed using DeepLIFT, which assigned contribution scores to input

features indicating their relative influence on gene expression prediction.

Predicting NABP regulatory targets using gene co-expression–informed

substitutions

To uncover regulatory targets of NABPs without relying on direct binding site

assays, we implemented a co-expression–driven modification of the deep learning input.

Specifically, DNA-binding proteins with low contribution scores in the original model

were replaced with NABPs whose regulatory interactions were inferred from gene co-

expression data. Of the 1,310 DBPs initially included, 302 were substituted with other

DBPs, and in a separate RNA-binding protein analysis, 325 were replaced with RNA-

binding proteins (RBPs) identified from the ENPD database [32]. Putative NABP–target

relationships were established using co-expression profiles provided by COXPRESdb [22],

where a Pearson's z-score > 2 between a NABP-encoding gene and a non-NABP gene

indicated a strong co-expression link. These co-expression pairs served as proxies for

regulatory interactions and were used to generate new input features for the model. To

integrate these proxies into the deep learning framework, hypothetical NABP binding

sites were assigned at the transcription start site (TSS) and 2 kb upstream of each co-

expressed gene. The model was then retrained using this modified input, and DeepLIFT

was applied to calculate importance scores, evaluating the influence of each NABP–gene

interaction on predicted expression levels. This approach enables the functional

characterization of NABPs with unknown or uncharacterized binding motifs by

leveraging gene co-expression as a proxy for regulatory targeting.

574   Prioritization of NABP regulatory targets based on importance score

575   rankings

576   To identify the primary regulatory targets of each NABP, genes were ranked

based on the importance scores assigned to NABP binding sites by the deep learning

model. Genes with binding sites consistently ranked in the top five or bottom five across

all NABPs were selected as putative targets, reflecting high positive or negative

regulatory influence. These targets represent genes most strongly associated with a given

NABP's regulatory activity. To ensure sufficient statistical power for downstream

enrichment analysis, rank thresholds were adjusted for each NABP based on the number

of high-confidence predictions. While a contribution score rank cutoff of five was

sufficient in most cases, the threshold was relaxed in specific cases—for example, up to

rank ten for proteins, which initially yielded a smaller number of predicted targets.

Functional enrichment and AI-based interpretation confirm the biological

relevance of NABP regulatory targets. To evaluate the biological coherence of predicted

NABP regulatory targets, we performed functional enrichment analysis using the

Statistical Overrepresentation Test from the PANTHER database alongside custom in-

house scripts [36]. Gene Ontology (GO) terms from biological process, molecular function,

and cellular component categories were analyzed for enrichment across three gene sets:

(1) primary regulatory target genes identified by the model, (2) all co-expressed genes

(Pearson's z-score > 2), and (3) randomly selected genes of matched size. Enrichment

was assessed using Fisher's exact test with Benjamini–Hochberg false discovery rate

(FDR) correction, and FDR < 0.05 was considered statistically significant. For this

analysis, human gene symbols were mapped to Ensembl Gene IDs using the HGNC

database (https://www.genenames.org/), and GO annotations were obtained from the

Gene Ontology resource [34,35]. Target gene IDs served as the test set, while the full

expression dataset used in the deep learning model formed the reference background.

Gene set enrichment analysis (GSEA) was performed using the pre-ranked

GSEA algorithm implemented in GSEApy (v1.1.11). For each DNA-binding protein

(DBP) and RNA-binding protein (RBP), DeepLIFT contribution scores were obtained for

all genes from the trained deep learning models. Because raw DeepLIFT scores vary in

magnitude across genes and include a large number of zeros, scores were normalized

within each gene by scaling the values across all 1,310 DBPs/RBPs to a continuous range

from −1 to 1 using min–max normalization. Genes were then ranked according to these

normalized contribution values specific to the DBP or RBP of interest. The ranked lists

were used as input for GSEA with the Reactome (v2025.1) and Gene Ontology

(Molecular Function, Biological Process, and Cellular Component, 2021 releases) gene

set collections. Analyses were performed with 10,000 permutations, weighted enrichment

statistics (weight = 1), and a minimum gene-set size of 1 to retain all functional categories.

Enrichment significance was assessed using normalized enrichment scores (NES),

nominal *p*-values, and false-discovery rates (FDR). This procedure enabled the

identification of pathways enriched among genes with high positive or negative

DeepLIFT contributions, thereby revealing regulatory roles and cell type–specific

pathways associated with each nucleic acid–binding protein.

To complement the statistical analyses, we applied a ChatGPT-based functional

inference method [37]. This approach used curated prompts—detailed in *Extended Data*

Tables 1 and 5—to predict functional roles based on gene sets, and retrieved relevant

literature using a modified version of the "4.Reference search and validation.ipynb" script

[37]. This script is publicly available on Code Ocean and GitHub

(https://doi.org/10.24433/CO.7045777.v1   ;

https://github.com/idekerlab/llm_evaluation_for_gene_set_interpretation).   The

ChatGPT-based results provided consistent and interpretable biological themes aligned

with the enrichment findings.

Validation of predicted NABP targets using experimentally determined

binding

To assess the biological relevance of the predicted regulatory targets, we

compared deep learning–inferred NABP–gene associations with experimentally

validated binding data. Specifically, we evaluated the spatial overlap between predicted

target genes and ChIP-seq peaks for 32 DNA-binding proteins (within ±5 kb of

transcription start sites [TSSs]) as well as eCLIP peaks for 29 RNA-binding proteins

(within the transcribed region from TSS to transcription end site [TES]). Gene structure

annotations were derived from the GENCODE v19 release [65], and binding site data were

obtained from the ENCODE portal and prior publications (**Supplementary Table S7**) [66].

This overlap analysis provides orthogonal validation for the predicted targets and

supports the utility of gene co-expression–based modeling for inferring NABP–target

relationships.

Statistical analysis

Functional enrichment was assessed using Fisher's exact test, with multiple testing

correction applied via the Benjamini–Hochberg false discovery rate (FDR) procedure.

Statistical significance was defined as FDR < 0.05. Distributions of DeepLIFT

contribution scores for nucleic acid-binding sites derived from gene co-expression data

and ChIP-seq data were compared using the Mann–Whitney U test (also known as the

Wilcoxon rank-sum test). A p-value < 0.05 was considered statistically significant. GSEA

(gene set enrichment analysis) was performed using DeepLIFT contribution scores of

DBPs and RBPs. The normalized enrichment score (NES) was used to quantify whether

regulatory target genes associated with specific biological functions were preferentially

located at the high or low end of the ranked gene list. Statistical significance was assessed

using permutation-based p-values, as implemented in the GSEA framework.

Data availability

RNA-seq data for human foreskin fibroblast (HFF) cells were obtained from the Gene

Expression Omnibus (GEO) under accession number GSM941744. RNA-seq data for

human mammary epithelial cells (HMEC) were retrieved from ENCODE via the UCSC

Genome Browser (http://genome.ucsc.edu/encode/) under the file name

wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam. RNA-seq data for human neural

progenitor cells (NPC) were obtained from GEO under accession number GSM915326.

RNA-seq data for HepG2 and K562 cells were obtained from the ENCODE portal under

accession numbers ENCFF670LIE and ENCFF584WML, respectively. mRNA co-

expression data were downloaded from COXPRESdb v7

(https://zenodo.org/record/6861444). The list of nucleic acid-binding proteins was

obtained from the Eukaryotic Nucleic acid Binding Protein (ENPD) database

(http://qinlab.sls.cuhk.edu.hk/ENPD/). ChIP-seq data for transcription factors were

downloaded from the GTRD database (v20.06 release; https://gtrd.biouml.org). eQTL

data (GTEx_Analysis_v8_eQTL.tar) were obtained from the GTEx portal (v8 release;

https://gtexportal.org). eCLIP data for RNA-binding proteins were retrieved from the

ENCODE portal (accession ENCSR456FVU) and from Supplementary Data 2 of Van

Nostrand et al., *Nature* (2020), https://doi.org/10.1038/s41586-020-2077-3.

Code availability

The deep learning model implemented in this study is available at

https://github.com/reposit2/coexgene.

Figures

a **Step 1: Predicting Regulatory Targets of Nucleic Acid-Binding Proteins (NABPs)**

Input:
- Binding site data for DNA-binding proteins (DBPs) around each gene
- Regulatory target genes predicted based on gene co-expression data

Deep Learning Analysis:
- Predict gene expression levels in each cell type
- Assign importance scores (DeepLIFT) to NABPs for each gene

Infer regulatory targets of NABPs based on importance scores

Regulatory targets

Same functional role of genes

NABP

Step 2: Functional Characterization of Predicted NABP Targets

- Perform enrichment analysis (e.g., GO, PANTHER Protein Class, Pathways, Reactome) using tools such as PANTHER
- Apply a ChatGPT-based method to infer gene functions and retrieve supporting literature
- Integrate ontology-based and semantic approaches
- Conduct GSEA using importance scores of predicted regulatory targets

Identify functional roles of NABPs in a cell type–specific context

b 1,310 nucleic acid-binding proteins

Gene expression prediction model

Input layer → 1D convolution → 1D convolution → 1D convolution → 1D convolution → 1D convolution → 1D Global Average Pooling → Fully connected layer → Fully connected layer → Output layer

Feature extraction

Input layer

1D Convolution layer

Fully-connected and output layers

ChIP-seq peaks

ChIP-seq

Gene co-expression-based sites

ChIP-seq peaks of DNA-binding proteins

Gene expression level

A>G

A>G   eGene

eSNP : expression-associated SNP

TSS : Transcription start sites

eGene: expression-associated gene; a gene whose expression level is associated with an eSNP

25,071 genes

Gene A Gene B Gene C Gene D ...

Fibroblasts
HMEC
NPC
HepG2 or K562

High   Low

Output: gene expression levels across human cell types

Figure 1. Overview of the deep learning framework for predicting and functionally characterizing NABP regulatory targets.

a. This diagram outlines the overall strategy used to infer the functional roles of nucleic acid-binding proteins (NABPs) from gene expression data. Gene expression profiles from various human cell types were integrated with NABP–gene co-expression relationships derived from COXPRESdb. These inferred interactions were used to replace low-contribution DNA-binding proteins in a deep learning model originally trained on ChIP-seq–based binding site data. The modified model was trained to predict gene expression levels and identify regulatory target genes using DeepLIFT-derived importance scores. Functional annotations of the predicted target genes were assessed using conventional enrichment analyses and GSEA, along with a ChatGPT-based method for literature-guided interpretation.

b. Gene expression levels in human cells were predicted using the genomic locations of DNA-binding sites at promoters and distal elements.

During model training, a DNA-binding protein bound to a distal region overlapping an expression-associated SNP (eSNP) was linked to a potentially regulated gene (eGene) based on eQTL data. For each gene–protein pair, ChIP-seq peak locations were binned (indicated by green boxes) and formatted into a matrix as input for deep learning.

In co-expression–based models, binding sites were artificially assigned near transcription start sites (TSSs). The deep learning model consisted of 1D convolutional layers, global average pooling, and fully connected layers. It predicted gene expression levels across 25,071 genes and multiple human cell types.

Figure 2. Relationship between gene expression levels and contribution scores.

Gene expression levels (log$_2$ TPM) predicted by a deep learning model were compared

with observed expression levels using Spearman's rank correlation.

a–c, Schematic overview of input feature configurations used for gene expression

prediction.

(**a**) DNA-binding sites of 1,310 proteins derived from ChIP-seq data.

(**b**) Replacement of 302 low-contribution DNA-binding proteins with co-expression–

derived DNA-binding proteins.

(**c**) Replacement of 325 DNA-binding proteins with RNA-binding proteins identified

from co-expression relationships.

d, Comparison of absolute DeepLIFT contribution scores for predicted regulatory sites

derived from gene co-expression versus ChIP-seq data, using a background defined as

0.5× the original input feature values. Each pair of violin plots represents co-expression–

based sites (right) and corresponding ChIP-seq sites not included in that set (left).

Contribution scores were significantly higher for co-expression–based sites (Wilcoxon

signed-rank test, p < 2.2 × 10^{-16}). Results obtained using an amplified background (input

× 2) are shown in **Supplementary Fig. S2c**.

e,f, Relationship between gene expression levels and DeepLIFT contribution scores for

representative DNA- and RNA-binding proteins across four cell types. Contribution

scores were computed using a background defined as input $\times$ 0.5 (**e**) or input $\times$ 2 (**f**). The

same set of proteins is shown in both panels, enabling direct comparison of background

effects. While background scaling alters the sign of the DeepLIFT–expression correlation,

the relative ordering of contribution scores and cell type–specific patterns are preserved.

b

d

Figure 3. Comparison of predicted regulatory target genes with experimental ChIP-seq and eCLIP binding data. This figure illustrates the correspondence between gene-level regulatory contribution scores inferred by deep learning and experimentally identified nucleic acid-binding protein (NABP) binding sites, using a DeepLIFT background defined as 0.5× the original input feature values.

a. Rank distribution of DeepLIFT contribution scores for nucleic acid-binding sites validated by ChIP-seq (DNA-binding proteins) or eCLIP (RNA-binding proteins). Contribution scores were ranked from highest to lowest and grouped into equal-sized rank bins. Binding sites located within promoters or transcribed regions of experimentally bound genes showed non-random enrichment toward higher or lower contribution ranks compared with all binding sites.

b. Proportion of high-contribution genomic regions overlapping ChIP-seq binding sites in promoter-associated regions. The fractions of high-contribution regions

overlapping ChIP-seq peaks are shown for promoter + enhancer regions (±100 kb from

the transcription start site) and promoter-only regions (±5 kb). Multiple rank- and score-

based thresholds were applied to define high- and low-contribution regions (see

Supplementary Note and Tables S2 and S3).

c. Overlap between high-contribution regions and eCLIP peaks for RNA-binding

proteins. Cumulative distributions show the proportion of high-contribution gene regions

overlapping experimentally determined eCLIP peaks within transcribed regions (from

transcription start site to transcription end site), compared with random controls.

Corresponding analyses using an amplified DeepLIFT background (input × 2) produced

similar results and are shown in **Supplementary Figure S3**.

d. Comparison of DeepLIFT contribution scores between HepG2 and K562 cells

under different background definitions. Scatter plots display contribution scores for

predicted regulatory target genes of three representative NABPs (RPLP0, ETF1, and

NELFE), comparing K562 (x-axis) and HepG2 (y-axis). Absolute contribution scores

were $\log_2$-transformed for visualization, and only genes with experimentally validated

binding sites were included. For each NABP, results are shown for two background

configurations (input × 0.5 and input × 2).

Figure [...] annotation terms among target genes of [...] nucleic [...]

Panels a [...] se discovery rate (FDR)–adjusted *p*-values of [...] 0.001 [...] l enrichment analysis was conducted for three gene sets: (1) model-predicted co-expressed genes, (2) all co-expressed genes with a Pearson correlation *z*-score > 2, and (3) randomly selected genes of equal size. Gene Ontology (GO) terms from the categories of biological process, molecular function, and cellular component were combined. Fisher's exact test *p*-values were adjusted using the Benjamini–Hochberg method to control the false discovery rate (FDR).

Figure 5. Functional mapping of nucleic acid-binding proteins reveals context-dependent regulatory roles in cancer and neural progenitor cells.

A deep learning model trained on transcriptomic co-expression data was used to predict the regulatory influence of nucleic acid-binding proteins (NABPs) on gene expression. Contribution scores were computed to quantify the impact of each NABP across target genes, and the highest-ranking genes were designated as predicted regulatory targets. Functional interpretation was performed using a large language model–based inference method, which identified key biological pathways associated with each NABP.

a

Protein-protein interactions at synapses

Glycosaminoglycan-protein linkage region biosynthesis

Gap junction assembly

Synaptic adhesion-like molecules

Figure 6. Cell type–specific pathway associations inferred from contribution score–based GSEA.

GSEA (gene set enrichment analysis) was performed using DeepLIFT contribution scores of nucleic acid-binding proteins (NABPs). This analysis revealed that genes with high or low contribution scores for a given NABP are preferentially associated with biological pathways relevant to each cell type.

The x-axis represents genes ranked by their DeepLIFT contribution scores for each NABP, with higher-scoring genes placed toward the left. PKM is shown here as a representative RNA-binding protein to illustrate contribution score–based pathway analysis. The y-axis indicates the running enrichment score, which increases when pathway-associated genes are concentrated among genes with higher contribution scores and decreases when they are enriched among genes with lower contribution scores. The normalized enrichment score (NES) summarizes the strength and direction of this enrichment across pathways.

For example, genes annotated to "Protein–protein interactions at synapses" were preferentially enriched among genes with lower PKM contribution scores in NPCs but among genes with higher contribution scores in K562 cells, indicating a marked cell type– specific difference in the inferred association between PKM and this pathway. Conversely, genes associated with "Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell" showed contrasting enrichment patterns between K562 and NPCs, highlighting differential pathway associations inferred from PKM contribution scores across the two cell types.

Tables

Table 1. Functional roles inferred for nucleic acid-binding proteins: integration of

UniProt annotations, ChatGPT-based predictions, and confidence scores.

Functional annotations obtained from UniProt entries were systematically compared with

significantly enriched biological processes among the model-predicted regulatory targets,

as identified by the PANTHER Overrepresentation Test (see **Supplementary Table S5**).

Shared or overlapping functions between each nucleic acid-binding protein (NABP) and

its predicted targets are indicated. Functional roles inferred using a ChatGPT-based

interpretation method are highlighted in yellow, with confidence scores—defined as the

proportion of annotated genes within the gene set—provided in parentheses. Confidence

levels are categorized as high (0.87–1.00), medium (0.82–0.86), low (0.01–0.81), or none

(0). For methodological reference, see Hu M. et al., *Nat. Methods*, 2025

(https://doi.org/10.1038/s41592-024-02525-x) [37].

ZBTB34 (HepG2 cells)
C2H2 zinc finger transcription factor
RNA polymerase II transcription regulatory region sequence-specific DNA binding
regulation of transcription by RNA polymerase II
C2H2 zinc-finger–directed regulation of RNA polymerase II transcription (0.88)

ZC3H4 (HepG2 cells)
chromatin/chromatin-binding, or -regulatory protein
RNA processing factor
basal RNA polymerase II transcription machinery binding
Chromatin-regulated RNA polymerase II transcription and RNA processing (0.89)

ZKSCAN5 (HepG2 cells)
C2H2 zinc finger transcription factor
RNA polymerase II cis-regulatory region sequence-specific DNA binding
autophagy
macroautophagy
Transcription-coupled autophagy–lysosome remodeling and membrane trafficking (0.78)

ZNF446 (HepG2 cells)
DNA-binding transcription activator activity, RNA polymerase II-specific
protein import into peroxisome matrix
non-motile cilium assembly
Transcription factor–directed ciliogenesis with peroxisome import and genome maintenance (0.86)

ZNF557 (HepG2 cells)
DNA-binding transcription activator activity, RNA polymerase II-specific
mRNA polyadenylation factor
RNA methyltransferase
RNA processing factor
RNA metabolism protein
Transcription-coupled RNA processing and RNP biogenesis (0.90)

POLR2G (NPC cells)
regulation of transcription by RNA polymerase II
RNA processing factor
translation initiation factor
Mitochondrial translation
RNA processing–coupled cytosolic and mitochondrial translation (0.90)

RBM27 (K562 cells)
RNA processing factor
RNA splicing factor
RNA metabolism protein
chromatin/chromatin-binding, or -regulatory protein
SUMO is proteolytically processed
Ub-specific processing proteases
RNAPII-coupled RNA processing and proteostasis axis (0.86)

SPEN
chromatin/chromatin-binding, or -regulatory protein
non-receptor serine/threonine protein kinase
ubiquitin-protein ligase
Signaling by Rho GTPases
Chromatin remodeling–anchored RNAPII transcription coordinated by ubiquitin and Rho-kinase signaling (0.78)

PKM (K562 cells)
Hypoxia response via HIF activation
glucose 6-phosphate metabolic process
Disorders of transmembrane transporters
Membrane Trafficking
Hypoxia/HIF-1–orchestrated metabolic reprogramming with cytoskeletal–membrane remodeling (0.80)

PKM (NPC cells)
Cellular response to hypoxia
KEAP1-NFE2L2 (NRF2) pathway
actin or actin-binding cytoskeletal protein
Hypoxia-driven metabolic rewiring with KEAP1–NRF2 signaling and actin–endosomal remodeling (0.86)

DNAJC2 (K562 cells)
RNA processing factor
chaperone
proteasome-mediated ubiquitin-dependent protein catabolic process
proteasomal protein catabolic process
ribosome biogenesis
RNA processing–coupled ribosome biogenesis and translation with proteostasis surveillance (0.88)

EIF2S2 (NPC cells)
proteinogenic amino acid biosynthetic process
Serine glycine biosynthesis
Unfolded Protein Response (UPR)
NFE2L2 regulating anti-oxidant/detoxification enzymes
ATF4 activates genes in response to endoplasmic reticulum stress
Integrated stress response coordinating UPR, antioxidant defense, and amino-acid/secretory metabolism (0.86)

ILF2 (K562 cells)
RNA processing factor
nucleic acid binding
G2/M DNA replication checkpoint
MASTL Facilitates Mitotic Progression
Coupled RNA processing–nuclear export aligned with mitotic cell-cycle control (0.86)

Table 2. Functional characterization of predicted regulatory targets of nucleic

acid-binding proteins using a ChatGPT-based inference framework.

Representative biological functions were inferred for each NABP by applying a large

language model–based gene set interpretation method. Functional themes are

summarized in bold for each NABP, while numbered subcategories denote distinct

functional roles associated with individual gene subsets.

ZBTB34 (HepG2 cells)
C2H2 zinc-finger–directed regulation of RNA polymerase II transcription (0.88)
1. Sequence-specific transcriptional control (core signal)
2. Transcriptional modulators and repressors (adjacent layer)
3. Signaling inputs that gate transcriptional outputs
4. RNA- and chromatin-adjacent regulators
5. Contextual effectors shaped by the transcriptional program

ZC3H4 (HepG2 cells)
Chromatin-regulated RNA polymerase II transcription and RNA processing (0.89)
1. Sequence-specific control + chromatin remodeling
2. RNAPII initiation and elongation
3. Co-transcriptional RNA processing
4. RNP assembly and nuclear–cytoplasmic transport
5. Ribosome biogenesis and rRNA metabolism
6. Post-translational regulation of the transcriptional apparatus
7. Genome maintenance coupling
8. Growth-signal gating
9. Peripheral modules

ZKSCAN5 (HepG2 cells)
Transcription-coupled autophagy–lysosome remodeling and membrane trafficking (0.78)
1. Transcriptional control as the top-level driver
2. Phosphoinositide signaling primes autophagosome initiation
3. Core autophagy execution and cargo selection
4. Endosome–lysosome tethering and autophagosome fusion
5. Proteasome and ubiquitylation complement lysosomal turnover
6. Organelle interfaces that feed the degradative axis
7. Ciliary/centrosomal trafficking as a specialized downstream client
8. Stress-responsive transcriptional tuning

ZNF446 (HepG2 cells)
Transcription factor–directed ciliogenesis with peroxisome import and genome maintenance (0.86)
1. Transcriptional control and RNA maturation
2. Ciliogenesis and membrane/organelle logistics
3. Peroxisome matrix import and organization
4. Genome maintenance and telomere control
5. Metabolic and cofactor support
6. Growth and sensory context
7. Network integration

ZNF557 (HepG2 cells)
Transcription-coupled RNA processing and RNP biogenesis (0.90)
1. Core transcriptional control
2. Co-transcriptional RNA processing
3. RNA modification and quality control
4. RNP assembly and ribosome/tRNA/rRNA biogenesis
5. Broad TF superstructure
6. Coupled auxiliary programs (contextual, not dominant)
844   7. Network integration

PKM (K562 cells)
Hypoxia/HIF-1–orchestrated metabolic reprogramming with cytoskeletal–membrane remodeling (0.80)
1. Transcriptional hypoxia module setting the program
2. Metabolic rewiring toward glycolysis and redox support
3. Proteostasis and catabolic throughput to sustain the shift
4. Actin–cytoskeleton remodeling and focal adhesion dynamics
5. Vesicle trafficking and membrane remodeling to deliver the phenotype
6. mRNA processing and translation control coupled to stress
7. KEAP1–NFE2L2 (NRF2)–linked oxidative stress buffering (adjacent)

PKM (NPC cells)
Hypoxia-driven metabolic rewiring with KEAP1–NRF2 signaling and actin–endosomal remodeling (0.86)
1. Metabolic rewiring and hypoxia response
2. KEAP1–NRF2 cytoprotective axis
3. Actin/adhesion remodeling and endocytic–lysosomal traffic
4. Proteostasis, autophagy, and apoptosis gating
5. Integrated model

ILF2 (K562 cells)
Coupled RNA processing–nuclear export aligned with mitotic cell-cycle control (0.86)
1. Transcription initiation and elongation across Pol I/II/III
2. Co-transcriptional RNA maturation: splicing, 3′-end formation, RNA modification
3. mRNP assembly, nuclear export, and nucleocytoplasmic transport
4. Nuclear envelope/pore remodeling synchronized with mitosis
5. Post-transcriptional regulators and chromatin/epigenetic control
6. Secretory and membrane-traffic support + stress responses
7. Gene-set composition is concordant with these functions

ILF2 (NPC cells)
RNAP II/III–coupled RNA biogenesis and ribosome assembly (0.88)
1. Transcription initiation and elongation
2. Co-transcriptional RNA processing and RNP assembly
3. Nuclear export, surveillance, and decay
4. rRNA processing and ribosome biogenesis
5. tRNA metabolism and charging
6. Chromatin and repair context

DNAJC2 (K562 cells)
RNA processing–coupled ribosome biogenesis and translation with proteostasis surveillance (0.88)
1. RNA processing and export
2. Ribosome biogenesis and rRNA metabolism
3. Translation and initiation control
4. Proteostasis and stress-response coupling
5. Genome maintenance and cell-cycle gating
6. Organelle and mitochondrial support

Table 3. Functional enrichments among predicted regulatory target genes of nucleic

acid-binding proteins.

Functional enrichment analysis was performed using the PANTHER database. This table

provides selected examples of annotations that were statistically enriched among

regulatory target genes predicted by the deep learning model trained on gene co-

expression data. '**Reference**' indicates the total number of human genes; '**Analyzed**'

denotes the number of regulatory target genes examined; '**Expected**' represents the

number of genes expected to have the annotation by chance; '**Fold Enrichment**' reflects

the observed-to-expected ratio; '**Raw *p*-value**' reports the result of Fisher's exact test; and

'**FDR**' refers to the Benjamini–Hochberg false discovery rate.

DBP, RBP	Functional annotation	Enriched	Not Enriched	Analyzed / Reference	Expected	Fold Enrichment	Raw *p*-value	FDR
PKM	Pentose phosphate pathway (PC0762)	K562	NPC	3 / 9	0.15	20.24	3.E-04	1.E-02
PKM	Pentose phosphate pathway (R-HSA-71336)	K562	NPC	3 / 15	0.25	12.14	2.E-03	2.E-02
PKM	Glycolysis (PC00024)	K562	NPC	4 / 20	0.33	12.14	3.E-04	1.E-02
PKM	Hypoxia response via HIF activation (P00030)	K562	NPC	4 / 30	0.49	8.09	1.E-03	4.E-02
PKM	response to hypoxia (GO:0001666)	K562	NPC	5 / 18	0.30	16.86	8.E-06	4.E-04
ILF2	G2/M DNA replication checkpoint (R-HSA-69478)	K562	NPC	2 / 5	0.06	30.95	2.E-03	4.E-02
POLR2G	ALKBH3 mediated reversal of alkylation damage (R-HAS-112126)	K562	NPC	2 / 4	0.11	17.80	5.E-03	5.E-02
DNAJC2	Ubiquitin proteasome pathway (P00060)	K562	NPC	8 / 58	0.60	13.26	1.E-07	2.E-05
PKM	Integrin signalling pathway (P00034)	NPC	K562	12 / 192	2.70	4.45	2.E-05	1.E-03
PKM	Glycogen metabolism (R-HSA-8982491)	NPC	K562	4 / 24	0.34	11.87	3.E-04	6.E-03
PKM	Glutamate Neurotransmitter Release Cycle (R-HSA-210500)	NPC	K562	4 / 23	0.32	12.38	3.E-04	5.E-03
EIF2S2	Serine glycine biosynthesis (PC02776)	NPC	K562	3 / 6	0.09	33.63	6.E-05	1.E-02

Table 4. Cell type−dependent enrichment of functional pathways identified by

GSEA

NES denotes the normalized enrichment score obtained from gene set enrichment

analysis (GSEA). ΔNES represents the difference in NES between NPC and K562. This

table summarizes the pathway enrichment results for PKM, presented as an example

RBP consistent with **Figure 6**. The seventeen Reactome pathways with the highest and

lowest ΔNES values are listed in the upper and lower sections, respectively. Pathways

with high ΔNES values were associated with neural cell differentiation, whereas those

with low ΔNES values were associated with cancer-related functions.

	Reactome pathway description	Reactome Pathway ID	NES		Δ NES
			K562	NPC	K562 - NPC
Pathways associated with neural cell differentiation	Protein-protein interactions at synapses	R-HSA-6794362	1.79	-1.11	2.89
	Glycosaminoglycan-protein linkage region biosynthesis	R-HSA-1971475	1.23	-1.56	2.79
	Golgi-to-ER retrograde transport	R-HSA-8856688	1.62	-1.15	2.77
	Attachment and Entry	R-HSA-9694614	1.39	-1.37	2.76
	Defective EXT2 causes exostoses 2	R-HSA-3656237	1.39	-1.36	2.75
	Defective B4GALT7 causes EDS, progeroid type	R-HSA-3560783	1.13	-1.60	2.73
	Phase 0 - rapid depolarisation	R-HSA-5576892	1.45	-1.24	2.69
	Gap junction assembly	R-HSA-190861	1.62	-1.06	2.68
	Diseases associated with glycosaminoglycan metabolism	R-HSA-3560782	1.07	-1.61	2.68
	Extracellular matrix organization	R-HSA-1474244	1.46	-1.21	2.67
	Synaptic adhesion-like molecules	R-HSA-8849932	1.37	-1.24	2.61
	Neuronal System	R-HSA-112316	1.63	-0.96	2.59
	Degradation of the extracellular matrix	R-HSA-1474228	1.50	-1.08	2.59
	IRE1alpha activates chaperones	R-HSA-381070	1.25	-1.31	2.56
	Smooth Muscle Contraction	R-HSA-445355	1.51	-1.04	2.54
	FGFR2c ligand binding and activation	R-HSA-190375	1.29	-1.23	2.52
	Receptor-type tyrosine-protein phosphatases	R-HSA-388844	1.62	-0.88	2.50
	Muscle contraction	R-HSA-397014	1.57	-0.92	2.49
	LGI-ADAM interactions	R-HSA-5682910	1.53	-0.97	2.49
	Respiratory syncytial virus (RSV) attachment and entry	R-HSA-9820960	1.39	-1.1	2.49
Pathways associated with cancer cells	Activation of anterior HOX genes in hindbrain development during early embryogenesis	R-HSA-5617472	-1.11	1.50	-2.62
	Role of LAT2/NTAL/LAB on calcium mobilization	R-HSA-2730905	-1.12	1.50	-2.62
	FCERI mediated MAPK activatio	R-HSA-2871796	-0.92	1.72	-2.64
	WNT ligand biogenesis and trafficking	R-HSA-3238698	-1.10	1.56	-2.66
	Role of phospholipids in phagocytosis	R-HSA-2029485	-1.31	1.38	-2.69
	Nucleotide Excision Repair	R-HSA-5696398	-1.30	1.41	-2.71
	Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell	R-HSA-198933	-1.52	1.24	-2.75
	Transcription-Coupled Nucleotide Excision Repair (TC-NER)	R-HSA-6781827	-1.47	1.33	-2.79
	Creation of C4 and C2 activators	R-HSA-166786	-1.13	1.70	-2.83
	CD22 mediated BCR regulation	R-HSA-5690714	-1.11	1.72	-2.83
	Initial triggering of complement	R-HSA-166663	-1.23	1.64	-2.87
	FCGR activation	R-HSA-2029481	-1.17	1.79	-2.96
	Scavenging of heme from plasma	R-HSA-2168880	-1.26	1.74	-3.00
	Mitotic G1 phase and G1/S transition	R-HSA-453279	-1.27	1.73	-3.01
	Response to metal ions	R-HSA-5660526	-1.29	1.77	-3.06
	Mitotic Metaphase and Anaphase	R-HSA-2555396	-1.27	1.79	-3.06
	M Phase	R-HSA-68886	-1.27	1.84	-3.11
	DNA Replication	R-HSA-69306	-1.30	1.94	-3.23
	Synthesis of DNA	R-HSA-69239	-1.29	1.98	-3.28
	Metallothioneins bind metals	R-HSA-5661231	-1.04	2.61	-3.65

Acknowledgements

873 Supercomputing resources were provided by the Human Genome Center, the Institute of

874 Medical Science, the University of Tokyo, and the ROIS National Institute of Genetics.

Author contributions

N.O. designed and conducted the research, and wrote and revised the manuscript. K.S.

revised the manuscript.

Supplementary data

Supplementary data are available at

Conflict of interests

None declared.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS)

KAKENHI [Grant Number 16K00387], a start-up research grant from Osaka University,

the KIOXIA Young Scientist Research Grant at Waseda University, the Grant for Basic

Science Research Projects from the Sumitomo Foundation, the Research Grant from the

CASIO Science Promotion Foundation, and the Research Grant from the Okawa

Foundation for Information and Telecommunications awarded to N.O., JST CREST

[Grant Number JPMJCR23N1], and the GteX Program Japan [Grant Number

JPMJGX23B4] awarded to K.S.

Supplementary Note and Tables

Supplementary Note for Figure 3. Definition of contribution score–based regions, thresholds, and enrichment analyses

Contribution score ranking and binning

For each nucleic acid-binding protein (NABP), DeepLIFT contribution scores were computed for all predicted gene-level regulatory regions and ranked from highest positive to lowest negative values. Ranked scores were divided into consecutive bins of 50 ranks, yielding 26 bins per NABP (rank 1: 1–50; rank 2: 51–100; …; rank 26: 1,251–1,310), unless otherwise specified.

Definition of high- and low-contribution regions

High- and low-contribution regions were defined using both rank-based and score-based thresholds to assess the robustness of enrichment results. Rank-based thresholds included upper and lower rank cutoffs (e.g., top or bottom 20 and 50 ranks), while score-based thresholds were defined using absolute contribution score cutoffs (e.g., ≥ 0.01 or ≥ 0.001). All threshold combinations used in the analyses are summarized in **Tables S2** and **S3.**

Genomic regions used for overlap analyses

For DNA-binding proteins, ChIP-seq overlap analyses were conducted within two genomic contexts: promoter-only regions defined as ±5 kb from the transcription start site (TSS), and extended promoter–enhancer regions defined as ±100 kb from the TSS. For RNA-binding proteins, eCLIP overlap analyses were restricted to transcribed regions

spanning from the TSS to the transcription end site (TES), including both exonic and

intronic intervals.

Random controls

Random control regions were generated by sampling genomic intervals matched

for size and genomic context, excluding annotated binding sites, to estimate the expected

overlap by chance. All enrichment and overlap analyses were repeated using these

controls for comparison.

Background configuration for DeepLIFT analyses

Unless otherwise noted, DeepLIFT contribution scores were computed using a

background defined as $0.5\times$ the original input feature values. Analyses using an

alternative background configuration (input $\times$ 2) yielded qualitatively similar results and

are presented in **Supplementary Figures S2** and **S3.**

Cell type configurations used for model training and comparative analyses

To accommodate differences in experimental data availability and analytical

objectives, multiple cell type configurations were used across analyses. For the primary

analyses of DNA-binding proteins (DBPs), models were trained using expression data

from HFFs, HMECs, NPCs, and HepG2, enabling validation against ChIP-seq datasets

available for HepG2 cells. For RNA-binding protein (RBP) analyses, models were trained

using HFFs, HMECs, NPCs, and K562 to facilitate validation using eCLIP datasets

available for K562 cells. For analyses requiring direct comparison of regulatory

contribution patterns between HepG2 and K562 for the same nucleic acid-binding

proteins (e.g., **Fig. 3d**), a unified four–cell-type configuration comprising HFFs, NPCs,

HepG2, and K562 was employed. This configuration enabled direct comparison of

DeepLIFT contribution score distributions across these two cell types while maintaining

a shared transcriptomic background. In all cases, cell type selection was driven by

analytical requirements and the availability of independent experimental validation data,

and no validation datasets were used as direct inputs during model training.

Gene co-expression–based predictions align with ChIP-seq–defined DNA-binding targets

High-ranking predicted binding regions frequently coincide with ChIP-seq peaks,

particularly in promoter–enhancer intervals. Deep learning–derived DNA-binding site

predictions based on gene co-expression data show substantial concordance with ChIP-

seq results, especially when broader genomic regions encompassing promoters and

putative enhancers are considered. When overlap was assessed within ±100 kb of

transcription start sites (TSSs)—a window supported by prior enhancer–gene mapping

studies [19,67]—the mean and median overlap ratios reached 0.70 and 0.77, respectively

(Ratio A2 / B2; **Supplementary Tables S2 and S3**). In contrast, overlap was lower for

promoter regions alone (±5 kb), with corresponding mean and median values of 0.35 and

0.34. Several DNA-binding proteins, including AKAP8, POGK, SPEN, THAP7, ZNF142,

and ZNF691, showed notably high promoter-region overlap ratios (>0.5), suggesting

potential direct regulatory activity. Nevertheless, not all ChIP-seq peaks necessarily

reflect functionally relevant transcriptional binding events. For RNA-binding proteins,

overlap was examined within the transcribed regions of RNA molecules, and the mean

and median overlap ratios were 0.70 and 0.76, respectively (Ratio A2 / B2;

Supplementary Table S4). These findings indicate that model-inferred high-scoring

sites do not universally coincide with experimentally determined binding regions,

underscoring the complexity and context dependence of regulatory interactions.

Supplementary Table S1. Co-expression–based selection enhances the

contribution scores of NABP binding sites.

Binding sites identified through gene co-expression analysis exhibited higher

absolute DeepLIFT contribution scores than those selected solely from ChIP-seq data

without co-expression information. This trend was consistently observed for DNA-

binding proteins and was significantly stronger for RNA-binding proteins across the

analyzed cell types. For each group, the table presents the mean and median values of

both positive and negative contribution scores. These results indicate that gene co-

expression–based selection preferentially highlights functionally relevant regulatory

interactions. The upper and lower sections of the table correspond to analyses performed

using DeepLIFT background references scaled to input × 0.5 and input × 2, respectively.

DBP / RBP	Cell type	Co-expression–based predictions. Upper row: mean values. Lower row: median values	Non-co-expression–based predictions. Upper row: mean values. Lower row: median values
DBP	HepG2	5.3.E-03	3.7.E-03
		2.8.E-03	1.9.E-03
RBP	K562	8.0.E-03	3.6.E-03
		5.6.E-03	1.8.E-03
DBP	NPC	3.9.E-03	3.8.E-03
		1.9.E-03	1.9.E-03
RBP	NPC	8.2.E-03	3.6.E-03
		5.9.E-03	1.7.E-03
DBP	HMEC	4.0.E-03	3.8.E-03
		1.8.E-03	1.8.E-03
RBP	HMEC	8.9.E-03	3.5.E-03
		6.4.E-03	1.7.E-03
DBP	HFF	3.9.E-03	3.8.E-03
		1.7.E-03	1.8.E-03
RBP	HFF	8.8.E-03	3.5.E-03
		6.5.E-03	1.7.E-03

DBP / RBP	Cell type	Co-expression–based: mean (upper) median (lower)	Non-co-expression–based: mean (upper) median (lower)
DBP	HepG2	5.28.E-03	3.71.E-03
		2.78.E-03	1.90.E-03
RBP	K562	7.95.E-03	3.56.E-03
		5.60.E-03	1.78.E-03
DBP	NPC	3.88.E-03	3.82.E-03
		1.87.E-03	1.87.E-03
RBP	NPC	8.20.E-03	3.56.E-03
		5.90.E-03	1.73.E-03
DBP	HMEC	4.01.E-03	3.81.E-03
		1.80.E-03	1.80.E-03
RBP	HMEC	8.92.E-03	3.55.E-03
		6.37.E-03	1.68.E-03
DBP	HFF	3.89.E-03	3.82.E-03
		1.70.E-03	1.79.E-03
RBP	HFF	8.76.E-03	3.55.E-03
		6.49.E-03	1.71.E-03

Supplementary Table S2. Enrichment of high-contribution promoter regions with experimentally validated DBP binding sites. (See Supplementary Excel file)

This table summarizes the proportion of promoter regions (±5 kb from transcription start sites [TSSs]) that were predicted to be functionally important by DeepLIFT contribution scores and validated by ChIP-seq data for the corresponding DNA-binding proteins (DBPs). Overlap ratios were calculated across four thresholds—two based on rank and two based on absolute score—to assess the robustness of the predictions.

High- and low-contribution regions were classified using the following thresholds for the 1,310 NABPs analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank ≤ 20 or ≥ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank ≤ 50 or ≥ 1,260)

Score-based thresholds:

(C1 and D1) contribution score ≥ 0.01 or ≤ –0.01

(C2 and D2) contribution score ≥ 0.001 or ≤ –0.001

Supplementary Table S3. Enrichment of high-contribution promoter and enhancer regions with ChIP-seq–validated DBP binding sites.

This table summarizes the proportion of predicted high-contribution regions—defined within ±100 kb of transcription start sites (TSSs), encompassing both promoter and distal enhancer regions—that overlap with ChIP-seq binding sites for the corresponding DNA-binding proteins (DBPs). Strong concordance between model-

derived predictions and ChIP-seq peaks across multiple thresholds highlights the
reliability of the contribution scoring method for identifying functionally relevant
regulatory sites.

High- and low-contribution regions were classified using the following thresholds
for the 1,310 NABPs analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank $\leq$ 20 or $\geq$ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank $\leq$ 50 or $\geq$ 1,260)

Score-based thresholds:

(C1 and D1) contribution score $\geq$ 0.01 or $\leq$ –0.01

(C2 and D2) contribution score $\geq$ 0.001 or $\leq$ –0.001

1023

1024

Supplementary Table S4. Enrichment of high-contribution transcribed
regions with eCLIP–validated RBP binding sites.

This table summarizes the proportion of predicted high-contribution regions—
defined within the transcribed regions of genes (from transcription start site [TSS] to
transcription end site [TES])—that overlap with eCLIP peaks for the corresponding
RNA-binding proteins (RBPs). Strong concordance between model-based predictions
and eCLIP peaks across multiple thresholds underscores the reliability of the contribution
scoring method for identifying functionally relevant regulatory sites.

High- and low-contribution regions were classified using the following
thresholds applied to the 1,310 NABPs analyzed:

Rank-based thresholds:

(A1 and B1) proteins ranked in the top or bottom 20 (i.e., rank $\leq$ 20 or $\geq$ 1,290)

(A2 and B2) proteins ranked in the top or bottom 50 (i.e., rank $\leq$ 50 or $\geq$ 1,260)

Score-based thresholds:

(C1 and D1) contribution score $\geq$ 0.01 or $\leq$ –0.01

(C2 and D2) contribution score $\geq$ 0.001 or $\leq$ –0.001

1041

1042

1043   **References**

1044

1   Belz, G. T. & Nutt, S. L. Transcriptional programming of the dendritic cell network.
Nature reviews. Immunology **12**, 101-113 (2012). https://doi.org/10.1038/nri3149

2   Miller, J. C. *et al.* Deciphering the transcriptional network of the dendritic cell lineage.
Nature immunology **13**, 888-899 (2012). https://doi.org/10.1038/ni.2370

3   Ravasi, T. *et al.* An atlas of combinatorial transcriptional regulation in mouse and man.
Cell **140**, 744-752 (2010). https://doi.org/10.1016/j.cell.2010.01.044

4   Gertz, J. *et al.* Distinct properties of cell-type-specific and shared transcription factor
binding   sites.   *Molecular*   *cell*   **52**,   25-36   (2013).
https://doi.org/10.1016/j.molcel.2013.08.037

5   Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650-665 (2018).
https://doi.org/10.1016/j.cell.2018.01.029

6   Vaquerizas, J. M., Kummerfeld, S. K., Teichmann, S. A. & Luscombe, N. M. A census of
human transcription factors: function, expression and evolution. *Nature reviews. Genetics*
10, 252-263 (2009). https://doi.org/10.1038/nrg2538

7   Pierson, E. *et al.* Sharing and Specificity of Co-expression Networks across 35 Human
Tissues.   *PLoS*   *Comput*   *Biol*   **11**,   e1004220   (2015).
https://doi.org/10.1371/journal.pcbi.1004220

8   Sonawane, A. R. *et al.* Understanding Tissue-Specific Gene Regulation. *Cell Rep* **21**,
1077-1088 (2017). https://doi.org/10.1016/j.celrep.2017.10.001

9   Saha, A. *et al.* Co-expression networks reveal the tissue-specific regulation of transcription
and   splicing.   *Genome*   *research*   **27**,   1843-1858   (2017).
https://doi.org/10.1101/gr.216721.116

10   Boyle, A. P. *et al.* Comparative analysis of regulatory information and circuits across

distant species. *Nature* **512**, 453-456 (2014). https://doi.org/10.1038/nature13668

11 Ray, D. *et al.* RNA-binding proteins that lack canonical RNA-binding domains are rarely
sequence-specific. *Sci Rep* **13**, 5238 (2023). https://doi.org/10.1038/s41598-023-32245-
9

12 Zeke, A. *et al.* Deep structural insights into RNA-binding disordered protein regions.
Wiley Interdiscip Rev RNA **13**, e1714 (2022). https://doi.org/10.1002/wrna.1714

13 Zaharias, S. *et al.* Intrinsically disordered electronegative clusters improve stability and
binding specificity of RNA-binding proteins. *J Biol Chem* **297**, 100945 (2021).
https://doi.org/10.1016/j.jbc.2021.100945

14 Kribelbauer, J. F., Rastogi, C., Bussemaker, H. J. & Mann, R. S. Low-Affinity Binding Sites
and the Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev Biol*
35, 357-379 (2019). https://doi.org/10.1146/annurev-cellbio-100617-062719

15 Neph, S. *et al.* Circuitry and dynamics of human transcription factor regulatory networks.
Cell **150**, 1274-1286 (2012). https://doi.org/10.1016/j.cell.2012.04.040

16 Marbach, D. *et al.* Tissue-specific regulatory circuits reveal variable modular
perturbations across complex diseases. *Nature methods* **13**, 366-370 (2016).
https://doi.org/10.1038/nmeth.3799

17 Fagny, M. *et al.* Exploring regulation in tissues with eQTL networks. *Proc Natl Acad Sci*
U S A **114**, E7841-E7850 (2017). https://doi.org/10.1073/pnas.1707375114

18 Agarwal, V. *et al.* Massively parallel characterization of transcriptional regulatory elements.
Nature (2025). https://doi.org/10.1038/s41586-024-08430-9

19 Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-
range interactions. *Nature methods* **18**, 1196-1203 (2021).
https://doi.org/10.1038/s41592-021-01252-x

20 Tasaki, S., Gaiteri, C., Mostafavi, S. & Wang, Y. Deep learning decodes the principles of
differential gene expression. *Nat Mach Intell* **2**, 376-386 (2020).
https://doi.org/10.1038/s42256-020-0201-6

21 Osato, N. & Hamada, M. Systematic discovery of directional regulatory motifs associated
with human insulator sites. *bioRxiv* (2025).
https://doi.org/https://doi.org/10.1101/2024.01.20.573595

22 Obayashi, T., Kodate, S., Hibara, H., Kagaya, Y. & Kinoshita, K. COXPRESdb v8: an
animal gene coexpression database navigating from a global view to detailed investigations.
Nucleic acids research **51**, D80-D87 (2023). https://doi.org/10.1093/nar/gkac983

23 Yang, S. *et al.* COEXPEDIA: exploring biomedical hypotheses via co-expressions
associated with medical subject headings (MeSH). *Nucleic acids research* **45**, D389-D396
(2017). https://doi.org/10.1093/nar/gkw868

24   Zhu, Q. *et al.* Targeted exploration and analysis of large cross-platform human
transcriptomic compendia. *Nature methods* **12**, 211-214, 213 p following 214 (2015).
https://doi.org/10.1038/nmeth.3249

25   Miller, H. E. & Bishop, A. J. R. Correlation AnalyzeR: functional predictions from gene
co-expression correlations. *BMC Bioinformatics* **22**, 206 (2021).
https://doi.org/10.1186/s12859-021-04130-7

26   Raina, P. *et al.* GeneFriends: gene co-expression databases and tools for humans and
model organisms. *Nucleic acids research* **51**, D145-D158 (2023).
https://doi.org/10.1093/nar/gkac1031

27   Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global
discovery of conserved genetic modules. *Science (New York, N.Y.)* **302**, 249-255 (2003).
https://doi.org/10.1126/science.1087447

28   Lemoine, G. G., Scott-Boyer, M. P., Ambroise, B., Perin, O. & Droit, A. GWENA: gene
co-expression networks analysis and extended modules characterization in a single
Bioconductor package. *BMC Bioinformatics* **22**, 267 (2021).
https://doi.org/10.1186/s12859-021-04179-4

29   Warde-Farley, D. *et al.* The GeneMANIA prediction server: biological network integration
for gene prioritization and predicting gene function. *Nucleic acids research* **38**, W214-220
(2010). https://doi.org/10.1093/nar/gkq537

30   Wong, A. K., Krishnan, A. & Troyanskaya, O. G. GIANT 2.0: genome-scale integrated
analysis of gene networks in tissues. *Nucleic acids research* **46**, W65-W70 (2018).
https://doi.org/10.1093/nar/gky408

31   UniProt, C. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic acids*
research **53**, D609-D617 (2025). https://doi.org/10.1093/nar/gkae1010

32   Tak Leung, R. W., Jiang, X., Chu, K. H. & Qin, J. ENPD - A Database of Eukaryotic
Nucleic Acid Binding Proteins: Linking Gene Regulations to Proteins. *Nucleic acids*
research **47**, D322-D329 (2019). https://doi.org/10.1093/nar/gky1112

33   Shrikumar, A., Greenside, P. & Kundaje, A. in *Proceedings of the 34th International*
Conference on Machine Learning Vol. 70 (eds Precup Doina & Teh Yee Whye) 3145--
3153 (PMLR, Proceedings of Machine Learning Research, 2017).

34   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nature genetics* **25**, 25-29 (2000). https://doi.org/10.1038/75556

35   Gene Ontology, C. *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224** (2023).
https://doi.org/10.1093/genetics/iyad031

36   Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all.
Protein Sci **31**, 8-22 (2022). https://doi.org/10.1002/pro.4218

1140	37	Hu, M. *et al.* Evaluation of large language models for discovery of gene set function.
1141		*Nature methods* **22**, 82-91 (2025). https://doi.org/10.1038/s41592-024-02525-x
1142	38	Zahra, K., Dey, T., Ashish, Mishra, S. P. & Pandey, U. Pyruvate Kinase M2 and Cancer:
1143		The Role of PKM2 in Promoting Tumorigenesis. *Front Oncol* **10**, 159 (2020).
1144		https://doi.org/10.3389/fonc.2020.00159
1145	39	Zhang, Z. *et al.* PKM2, function and expression and regulation. *Cell Biosci* **9**, 52 (2019).
1146		https://doi.org/10.1186/s13578-019-0317-8
1147	40	Alquraishi, M. *et al.* Pyruvate kinase M2: A simple molecule with complex functions. *Free*
1148		*Radic Biol Med* **143**, 176-192 (2019).
1149		https://doi.org/10.1016/j.freeradbiomed.2019.08.007
1150	41	Azoitei, N. *et al.* PKM2 promotes tumor angiogenesis by regulating HIF-1alpha through
1151		NF-kappaB activation. *Mol Cancer* **15**, 3 (2016). https://doi.org/10.1186/s12943-015-
1152		0490-2
1153	42	He, P. *et al.* PKM2 is a key factor to regulate neurogenesis and cognition by controlling
1154		lactate homeostasis. *Stem Cell Reports* **20**, 102381 (2025).
1155		https://doi.org/10.1016/j.stemcr.2024.11.011
1156	43	Bonar, N. A. & Petersen, C. P. Integrin suppresses neurogenesis and regulates brain tissue
1157		assembly in planarian regeneration. *Development* **144**, 784-794 (2017).
1158		https://doi.org/10.1242/dev.139964
1159	44	Porcheri, C., Suter, U. & Jessberger, S. Dissecting integrin-dependent regulation of neural
1160		stem cell proliferation in the adult brain. *J Neurosci* **34**, 5222-5232 (2014).
1161		https://doi.org/10.1523/JNEUROSCI.4928-13.2014
1162	45	Zheng, X. *et al.* Metabolic reprogramming during neuronal differentiation from aerobic
1163		glycolysis to neuronal oxidative phosphorylation. *eLife* **5** (2016).
1164		https://doi.org/10.7554/eLife.13374
1165	46	Agostini, M. *et al.* Metabolic reprogramming during neuronal differentiation. *Cell Death*
1166		*Differ* **23**, 1502-1514 (2016). https://doi.org/10.1038/cdd.2016.36
1167	47	Saez, I. *et al.* Neurons have an active glycogen metabolism that contributes to tolerance to
1168		hypoxia. *J Cereb Blood Flow Metab* **34**, 945-955 (2014).
1169		https://doi.org/10.1038/jcbfm.2014.33
1170	48	Matthews, H. K., Bertoli, C. & de Bruin, R. A. M. Cell cycle control in cancer. *Nat Rev*
1171		*Mol Cell Biol* **23**, 74-88 (2022). https://doi.org/10.1038/s41580-021-00404-3
1172	49	Dominguez-Brauer, C. *et al.* Targeting Mitosis in Cancer: Emerging Strategies. *Molecular*
1173		*cell* **60**, 524-536 (2015). https://doi.org/10.1016/j.molcel.2015.11.006
1174	50	Levine, M. S. & Holland, A. J. The impact of mitotic errors on cell proliferation and
1175		tumorigenesis. *Genes Dev* **32**, 620-638 (2018). https://doi.org/10.1101/gad.314351.118

1176   51   Huang, X. *et al.* D-Serine regulates proliferation and neuronal differentiation of neural
1177   stem cells from postnatal mouse forebrain. *CNS Neurosci Ther* **18**, 4-13 (2012).
1178   https://doi.org/10.1111/j.1755-5949.2011.00276.x

1179   52   El-Hattab, A. W. Serine biosynthesis and transport defects. *Mol Genet Metab* **118**, 153-
1180   159 (2016). https://doi.org/10.1016/j.ymgme.2016.04.010

1181   53   Kaida, A. & Iwakuma, T. Regulation of p53 and Cancer Signaling by Heat Shock Protein
1182   40/J-Domain Protein Family Members. *Int J Mol Sci* **22** (2021).
1183   https://doi.org/10.3390/ijms222413527

1184   54   Sterrenberg, J. N., Blatch, G. L. & Edkins, A. L. Human DNAJ in cancer and stem cells.
1185   *Cancer Lett* **312**, 129-142 (2011). https://doi.org/10.1016/j.canlet.2011.08.019

1186   55   Ladomery, M. Multifunctional proteins suggest connections between transcriptional and
1187   post-transcriptional processes. *Bioessays* **19**, 903-909 (1997).
1188   https://doi.org/10.1002/bies.950191010

1189   56   Kim, S., Yu, N. K. & Kaang, B. K. CTCF as a multifunctional protein in genome regulation
1190   and gene expression. *Exp Mol Med* **47**, e166 (2015).
1191   https://doi.org/10.1038/emm.2015.33

1192   57   Osato, N. Characteristics of functional enrichment and gene expression level of human
1193   putative transcriptional target genes. *BMC Genomics* **19**, 957 (2018).
1194   https://doi.org/10.1186/s12864-017-4339-5

1195   58   Burley, S. K. & Kamada, K. Transcription factor complexes. *Curr Opin Struct Biol* **12**, 225-
1196   230 (2002). https://doi.org/10.1016/s0959-440x(02)00314-7

1197   59   Goos, H. *et al.* Human transcription factor protein interaction networks. *Nat Commun* **13**,
1198   766 (2022). https://doi.org/10.1038/s41467-022-28341-5

1199   60   Berenson, A. *et al.* Paired yeast one-hybrid assays to detect DNA-binding cooperativity
1200   and antagonism across transcription factors. *Nat Commun* **14**, 6570 (2023).
1201   https://doi.org/10.1038/s41467-023-42445-6

1202   61   Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic*
1203   *acids research* **41**, D991-995 (2013). https://doi.org/10.1093/nar/gks1193

1204   62   Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene
1205   expression and hybridization array data repository. *Nucleic acids research* **30**, 207-210
1206   (2002). https://doi.org/10.1093/nar/30.1.207

1207   63   Kolmykov, S. *et al.* GTRD: an integrated view of transcription regulation. *Nucleic acids*
1208   *research* **49**, D104-D111 (2021). https://doi.org/10.1093/nar/gkaa1057

1209   64   Hinrichs, A. S. *et al.* The UCSC Genome Browser Database: update 2006. *Nucleic acids*
1210   *research* **34**, D590-598 (2006). https://doi.org/10.1093/nar/gkj144

65   Mudge, J. M. *et al.* GENCODE 2025: reference gene annotation for human and mouse.

Nucleic acids research **53**, D966-D975 (2025). https://doi.org/10.1093/nar/gkae1078

66 Van Nostrand, E. L. *et al.* A large-scale binding and functional map of human RNA-
binding proteins. *Nature* **583**, 711-719 (2020). https://doi.org/10.1038/s41586-020-
2077-3

67 Gasperini, M. *et al.* A Genome-wide Framework for Mapping Gene Regulation via
Cellular   Genetic   Screens.   *Cell*   **176**,   1516   (2019).
https://doi.org/10.1016/j.cell.2019.02.027

1219