Title

Functional prediction of DNA/RNA-binding proteins by deep learning

from gene expression correlations

Naoki Osato [1*]

[1] School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan.

E-mail: naokiosato11@gmail.com

*Corresponding author.

Abstract

Deep learning methods have been developed to predict gene expression levels in

different cell and tissue types from DNA-binding sites of proteins including transcription

factors identified by ChIP-seq assay. However, experimental identification of

DNA/RNA-binding sites of proteins and functional characterization of proteins are

difficult for some proteins. Gene co-expression correlation has been identified from gene

expression data among cells and tissues, and includes the interactions between a

DNA/RNA-binding protein and other genes. To predict the functional roles of

DNA/RNA-binding proteins based on the interactions between a DNA/RNA-binding

protein and other genes, I used the interactions to predict gene expression levels by

partially replacing the input data in the deep leaning analysis. The correlation coefficient

between actual and predicted gene expression levels increased from 0.70 to 0.80,

indicating that gene co-expression data contribute to predict gene expression levels. Some

of the functional annotations statistically overrepresented in the predicted target genes

correspond to the known and associated functions of the DNA/RNA-binding proteins.

Furthermore, functional annotations that are not registered in a database of functional

annotations of genes have been confirmed in published studies. It contributes to predict

the functional roles of DNA/RNA-binding proteins.

Keywords

Computational biology, Deep learning, Gene expression, Gene regulation, Protein

function prediction

Introduction

Deep learning analysis has improved the accuracy of predicting gene expression

levels from the binding sites of DNA/RNA-binding proteins including transcription

factors and other regulatory factors (Tasaki, et al., 2020). Transcription factors bind to

promoters around transcription start sites (TSSs) and enhancers, which are located far

upstream and downstream of genes and TSSs. The method predicts gene expression levels

in 53 human tissues based on the experimental and predicted data of binding sites of

DNA/RNA-binding proteins and miRNAs in promoter and mRNA-transcribed regions.

The correlation coefficient between actual and predicted gene expression levels was 0.82.

In my previous study, I modified the method to include the data of DNA-binding proteins

bound at open chromatin regions in up to 1 Mbp upstream and downstream regions from

TSSs, some of which regions would be associated with enhancers, and removed the

binding sites of RNA-binding proteins and miRNAs, the correlation coefficient was 0.70

(Osato and Hamada, 2025).

The binding sites of DNA/RNA-binding proteins are identified by experimental

assays such as ChIP-seq and ChIA-pet. These assays require high-quality antibodies of

the proteins, but it is difficult to obtain high-quality antibodies for some proteins. For

some DNA/RNA-binding proteins, it may be impractical to determine the binding

DNA/RNA sequences by experimental analysis with the time, cost, and labor available.

Large Language Model (LLM)-based deep learning models, such as Enformer, will

predict DNA sequences that are important for predicting gene expression levels (Avsec,

et al., 2021). Massively parallel reporter assays detected the activity of hundreds of

thousands of sequences (Agarwal, et al., 2025). However, it would be difficult to identify

which DNA/RNA-binding proteins bind to the DNA sequences.

Analysis of gene co-expression correlation between cell and tissue types suggested

regulatory interactions among genes, including mRNAs encoding DNA/RNA-binding proteins. Gene expression data by RNA-seq assay and the results of gene co-expression correlation analysis are available in databases (Miller and Bishop, 2021; Obayashi, et al., 2023; Raina, et al., 2023; Yang, et al., 2017; Zhu, et al., 2015). Gene co-expression network analysis has been reported (Lemoine, et al., 2021; Obayashi, et al., 2023; Raina, et al., 2023; Stuart, et al., 2003; Warde-Farley, et al., 2010; Wong, et al., 2018). The correlation of gene expression would be regulated by direct and/or non-direct interactions of molecules. Some of the molecules may be elusive to detect by experimental analysis and difficult to predict by other computational analyses. The correlation of gene expression is potentially associated with regulations that have not been identified.

In this study, to investigate the functional annotations enriched in the putative regulatory target genes of a DNA/RNA-binding protein, I used gene co-expression data to extract the interactions between a DNA/RNA-binding protein and other genes. The DNA-binding sites of about 300 proteins with a low contribution score in my previous deep learning model were replaced with those of other DNA/RNA-binding proteins based on gene co-expression data. Deep learning learned to predict gene expression levels and selected interactions between a DNA-binding site and a gene that contribute to the prediction. The regulatory target genes of a transcription factor include functionally related genes (McLean, et al., 2010). Thus, the functional enrichment of genes is potentially discovered, although the prediction of regulatory target genes may not be precise enough to achieve a research purpose. I compared the overlap between the predicted regulatory target genes of a DNA/RNA-binding proteins and the DNA/RNA-binding sites by experimental assays such as ChIP-seq and eCLIP. Then, I performed functional enrichment analyses of the predicted regulatory target genes of ten DNA/RNA-

binding proteins and found that the enriched functional annotations corresponded to the

known functional annotations in the UniProt database. Furthermore, I discovered that

some functional annotations enriched in the putative regulatory target genes of a

DNA/RNA-binding protein were not described in the UniProt database, but were

investigated in published papers.

These analyses would be useful to identify the functional roles and regulatory target

genes of DNA/RNA-binding proteins without using experimental data from protein-

binding assays, and are expected to predict the direct and non-direct interactions between

DNA/RNA-binding proteins and their regulatory target genes, where some interactions

may be difficult to study by experimental analysis. This will contribute to the discovery

of novel interactions and functions, and reduce the cost of experimental analysis.

Results

To examine the effect of the interaction data of co-expressed genes on the

prediction of gene expression levels, among 1,310 DNA-binding proteins in the input

data of deep learning, 302 DNA-binding proteins with low contribution scores

(DeepLIFT scores) were replaced with the interaction data of co-expressed genes

encoding DNA-binding proteins and the mRNAs other than DNA/RNA-binding proteins.

The DNA-binding proteins encoded by the co-expressed genes were randomly selected

from the gene symbols of DNA-binding proteins in the ENPD database (Tak Leung, et

al., 2019), and in the other databases of DNA-binding proteins used for the analysis of

DNA-binding motifs. Interestingly, the correlation coefficient between actual and

predicted gene expression levels increased from 0.70 to 0.80 in human foreskin fibroblast

(HFF) cells using co-expression data (**Fig. 2a, b**). Then, among 1,310 DNA-binding

proteins, 325 DNA-binding proteins with low contribution scores were replaced with the

interaction data of co-expressed genes encoding RNA-binding proteins and other mRNAs.

The RNA-binding proteins encoded by the co-expressed genes were randomly selected

from the gene symbols of RNA-binding proteins in the ENPD database. The correlation

coefficient increased from 0.70 to 0.81 using co-expression data as well (**Fig. 2c**). These

results suggest that the interactions of co-expression of genes contribute to the prediction

of gene expression levels.

The contribution scores of the input data selected from the gene expression

correlations were distributed over a wide range between high and low, but the mean and

median of the scores were larger than the original (**Fig. 2d and Supplementary Table**

S1). I examined the distribution of Spearman's correlations between contribution scores

and gene expression levels (log2-TPMs, transcripts per million) (**Supplementary Fig.**

S2), and plotted example relationships between the contribution scores of a DNA/RNA-

binding protein and the expression levels of the regulatory target genes across cell types

(**Fig. 2e**). Based on the interactions between co-expressed genes and a DNA/RNA-

binding protein, deep learning adjusted the prediction model of gene expression levels in

each cell type.

To investigate whether the regulatory target genes of a DNA/RNA-binding protein

selected based on their contribution scores are associated with the DNA/RNA-binding

sites of the protein, I used the DNA-binding sites by ChIP-seq assay and the RNA-binding

sites by eCLIP assay around the genes, where a DNA-binding site of the protein by ChIP-

seq assay within ±5 kb of TSSs or an RNA-binding site of the protein by eCLIP assay in

mRNA-transcribed regions of a gene (ChIP-seq data of 32 DNA-binding proteins in

HepG2 cell and eCLIP data of 29 RNA-binding proteins in K562 cell). The genes with a

DNA-binding site of the protein or an RNA-binding site of the protein corresponded to high positive or low negative contribution scores of the regulatory target genes of the DNA/RNA-binding protein derived from the deep leaning of gene co-expression data (**Fig. 3 and Supplementary Fig. S1**). The high and low contribution scores would be associated with the regulatory functions of activation and repression of gene expression from the description of the contribution score (DeepLIFT score) on the GitHub webpage of DeepLIFT (Shrikumar, et al., 2017) and a previous study using DeepLIFT (Tasaki, et al., 2020).

The transcriptional target genes of a transcription factor include associated functions of genes. The functional enrichment of genes would be useful to predict the function of a transcription factor and to confirm that the transcriptional target genes are not randomly selected. For the functional enrichment analysis of regulatory target genes of DNA/RNA-binding proteins, I used the statistical overrepresentation test of the gene list analysis of Panther DB (https://pantherdb.org/) (Thomas, et al., 2022). The test uses functional annotation databases such as PANTHER Pathways, Protein Class, GO (Gene Ontology)-Slim, GO, and Reactome Pathways. The functional annotations enriched in putative regulatory target genes of a DNA/RNA-binding protein were compared with those of the DNA/RNA-binding protein obtained from the UniProt database (e.g. https://www.uniprot.org/uniprotkb/Q9Y2L8/entry) (UniProt, 2025). The ten examples of DNA/RNA-binding proteins are shown in **Table 1** and **Supplementary Table S4**. The putative functional annotations of the DNA/RNA-binding proteins correspond to and are associated with those in the UniProt database and publications. Notably, among the ten examples, AKAP8 has cell cycle-related functional annotations such as "GO:0000278 mitotic cell cycle" and "GO:0044839 cell cycle G2/M phase transition" in the UniProt

database (https://www.uniprot.org/uniprotkb/O43823/entry). In this study, AKAP8 was

suggested to be involved in circadian clock systems. By searching Google and the

PubMed database, I found the following papers showing that AKAP8 is involved in

circadian rhythms: "But the genes, such as NFIL3, AKAP8, and SALL3, highly correlated

with known molecular clock genes (CRY1) can be identified as the candidate circadian

genes." and "AKAP5 and AKAP8 are both involved in the biosynthesis of the circadian

hormone melatonin, which is markedly disturbed in people with ASDs." (Poelmans, et

al., 2013; Tan, et al., 2024). These results demonstrate that the method and analysis in this

study detected known and novel functional annotations associated with DNA/RNA-

binding proteins.

The deep learning analysis changed the contribution (i.e., weights in the neural

network) of a DNA/RNA-binding protein to the prediction of the expression levels of the

regulatory target genes of the DNA/RNA-binding protein during the learning phase. To

investigate the effect of learning on the functional enrichment analysis of the target genes,

I performed the functional enrichment analysis of all co-expressed genes (z score > 2) of

the ten DNA/RNA-binding proteins using Panther DB, and compared the functional

annotations enriched in all co-expressed genes of a DNA/RNA-binding protein and those

of the regulatory target genes of the DNA/RNA-binding protein in the deep learning

analysis shown in **Table 2** and **Supplementary Table S5**. Although the number of

functional annotations enriched in all co-expressed genes of a DNA/RNA-binding protein

increased due to the increase in genes, some functional annotations were only or highly

enriched in the regulatory target genes of the DNA/RNA-binding protein after learning.

Notably, as in the case of AKAP8 functional annotations, circadian rhythm-related

functional annotations were detected only after learning (**Table 2** and annotations

highlighted in yellow in the 'S2_AKAP8' tab of **Supplementary Table S4**). As for PKM

functional annotations, kinase etc. were identified only after learning (**Table 2** and

'S2_PKM' in **Supplementary Table S4**). PKM interacts with ALDOA (Snel, et al., 2000;

Szklarczyk, et al., 2023) (STRING database interaction, mouse PKM https://string-

db.org/network/10090.ENSMUSP00000034834 ; human PKM also has an interaction

with ALDOA in the database), and I found research papers supporting that the functional

annotation "aldolase (PC00044)" is associated with PKM involved in glycolysis:

"Aldolase A (ALDOA) is an enzyme that plays an important role in glycolysis and

gluconeogenesis, which is closely related to tumor metabolism" (Tian, et al., 2022; Zhang,

et al., 2017). These functional annotations were enriched after learning. Thus, they would

be involved in the functions in K562 cell where the regulatory target genes of an RNA-

binding protein were selected. Circadian clocks and PKM would be associated with

human acute myeloid leukemia (AML) cells via glycolysis (Lee, 2021; Puram, et al.,

2016; Yang, et al., 2024; Zlacka and Zeman, 2021). These results suggest that combining

gene co-expression data with deep learning to predict gene expression levels would reveal

the functional associations of DNA/RNA-binding proteins in a cell or tissue type.

Discussion

The expression levels of human genes in cell types have been predicted using deep

learning based on the DNA-binding sites of proteins. Here, I applied this method to

predict unknown interactions between a DNA/RNA-binding protein and genes based on

gene expression correlations between cell types. Gene expression correlations were

statistically analyzed using gene expression data from RNA-seq experiments in cell and

tissue types. Experimental analysis of DNA/RNA-binding sites by ChIP-seq, eCLIP

assays, etc. is more time-consuming, expensive, and laborious than RNA-seq experiments.

Experimental analysis of DNA/RNA-binding sites is still difficult for some DNA/RNA-

binding proteins. Deep learning analysis of DNA/RNA-binding proteins based on

correlations of their expression levels with other genes predicted regulatory target genes

of the proteins. Regulatory target genes include associated functions of genes. Enrichment

analysis of the functional annotations of the regulatory target genes revealed the

functional associations and roles of the DNA/RNA-binding proteins. The predicted

functional annotations correspond to the known functional annotations of the proteins.

Furthermore, functional annotations not described in a database were also discovered and

confirmed in published studies.

When comparing DNA-binding sites predicted based on the contribution scores for

predicting gene expression levels with DNA-binding sites by ChIP-seq assay, the two

types of DNA-binding sites may not overlap completely. Although tens of thousands of

DNA-binding sites of proteins are identified by ChIP-seq assay, some of them may not

be directly involved in the regulation of gene expression. Some transcription factors are

involved in multiple functions (Ladomery, 1997) (Kim, et al., 2015). The number of co-

expressed genes of a DNA/RNA-binding protein with z score $\geq$ 2 was about less than

1,500, and was less than the number of genes with ChIP-seq peaks of the DNA/RNA-

binding proteins, but the most of the co-expressed genes with a statistical significance

would be involved in the regulation of gene expression. Our previous analysis of

insulator-associated DNA-binding proteins revealed that 97 transcription factors and

DNA-binding proteins would be associated with insulator functions that affect enhancer-

promoter interactions (Osato and Hamada, 2025). Some of them would be important for

the formation of chromatin loops, transcriptional condensates and/or phase separation,

which are different mechanisms to affect gene expression from the direct regulation of

genes by transcription factors. In the deep learning analysis of insulator-associated DNA-

binding proteins, we used eQTL data to find DNA-binding sites of proteins that affect

gene expression levels. However, in this study, I did not use the DNA-binding motif

sequences of proteins and open chromatin regions by DNase-seq or ATAC-seq assay to

identify their DNA-binding sites because the purpose of this study is to investigate DNA-

binding proteins whose DNA-binding motifs are unknown. Moreover, the association

between eSNP and eGene of eQTL data may include indirect interactions. The

contribution scores of DNA-binding sites of proteins selected based on gene expression

correlation showed higher or lower scores, suggesting that indirect interactions between

DNA-binding proteins and other genes would affect the scores.

As for the analysis of functional enrichment of regulatory target genes of a DNA-

binding protein, ten DNA/RNA-binding proteins correspond to the known functional

annotations of the DNA/RNA-binding proteins, suggesting that the number of predicted

regulatory target genes is sufficient to identify the functional annotations of the proteins

by a statistical test. The identification of functional enrichments of regulatory target genes

of a protein seemed to require more than one hundred of genes by a Fisher's exact test

adjusted with a false discovery rate (FDR) in Panther DB. The number of regulatory target

genes of some proteins was less than a hundred, and their functional enrichments seemed

to be difficult to identify. One of the next steps of this study would be to improve the

method for identifying the functional enrichments of a small number of genes by

considering other information such as gene expression levels and correlation of gene

expression.

When I replaced all the input data of the deep learning analysis with the interactions

between DNA/RNA-binding proteins and other genes based on gene expression

correlation, the number of functional enrichments of predicted regulatory target genes of

a DNA/RNA-binding protein was reduced, and the result of the functional enrichment

analysis did not include the same functional annotations as the known functional

annotations of the DNA/RNA-binding protein in the UniProt database. Thus, the high

replacement ratio of the input data would lead to learning error-prone interactions

between a DNA/RNA-binding protein and regulatory target genes. In my previous study,

when the regulatory target genes of a DNA-binding protein were replaced by randomly

selected genes, the number of functional enrichments decreased as the number of

randomly selected genes increased. The number of functional enrichments reflects the

accuracy of the regulatory target genes (Osato, 2018).

The DNA-binding proteins were ranked according to the contribution scores of

their DNA-binding sites for each gene. When the rank of a DNA-binding protein is higher

or equal to 50 or lower or equal to 1,260 out of a total of 1,310, the functional enrichments

of putative target genes appeared to be identified by the statistical test in Panther DB.

Some DNA/RNA-binding proteins did not have high or low-ranking putative regulatory

target genes in HepG2 cells. These proteins may have high or low-ranking target genes

in other cell types, and thus functional enrichment analysis of target genes could be

performed to reveal their functional roles. Some transcription factors and DNA-binding

proteins would bind to each other to form the protein complex and regulate gene

expression levels (Burley and Kamada, 2002) (Goos, et al., 2022) (Berenson, et al., 2023).

These proteins would bind to the same promoters of genes and the ranking of each protein

in the complex may not always be high or low. Analysis of the co-localization of

DNA/RNA-binding proteins would reveal the functional roles of the protein complex.

The deep learning model, model parameters, and analysis methods used in this

study could be adapted and improved for the purpose of this study. I examined a small

number of DNA/RNA-binding proteins in a few cell types in this study, and would be

able to predict the functional roles of a larger number of DNA/RNA-binding proteins in

cell and tissue types. The results of the analysis would be useful to understand and study

the biological and molecular functions of unknown functions of DNA/RNA-binding

proteins and their regulatory target genes.

Methods

Deep learning analysis

I modified the deep learning method for predicting gene expression levels among

cell and tissue types based on the DNA-binding sites of transcription factors and other

regulators, and included the DNA-binding sites of transcription factors in regions far

upstream and downstream of TSSs (Osato and Hamada, 2025; Tasaki, et al., 2020) (**Fig.**

1). To reduce the heterogeneity of cells in a tissue, I used gene expression data of cell

types. The gene expression data of different cell types were obtained from public

databases: human foreskin fibroblasts (HFFs) (GSM941744), human mammary epithelial

cells (HMECs) (Encyclopedia of DNA Elements at UCSC

http://genome.ucsc.edu/encode/,

'wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam') (Barrett, et al., 2013; Edgar, et

al., 2002), neural progenitor cells (NPCs) (GSM915326, ENCFF529SIO), human

hepatoma cell line (HepG2) (ENCFF670LIE), and human leukemia cell line (K562)

(ENCFF584WML). FPKM values of gene expression levels were calculated based on

GENCODE V19 gene annotations in the ENCODE portal database

(https://www.encodeporoject.org/) (Luo, et al., 2020; Mudge, et al., 2025) using the

RSeQC tool (http://rseqc.sourceforge.net/) (Wang, et al., 2012). The FPKM values were

transformed to log2-fold with the addition of 0.25 to avoid negative infinity results. Log2-

fold changes relative to the median log2-FPKM across the four cell types were calculated

according to the original deep learning method (transcriptome data processing as

described by Tasaki et al.) (Osato and Hamada, 2025; Tasaki, et al., 2020).

The DNA-binding sites overlapping with open chromatin regions in putative

enhancers located outside of promoter regions were added as input data in the deep

learning method. To identify the DNA-binding sites of DNA-binding proteins (DBPs) in

promoter and enhancer regions, ChIP-seq data of the DBPs were obtained from GTRD

database release v20.06 (https://gtrd.biouml.org) (Kolmykov, et al., 2021). The binding

site of a DBP that regulates the expression level of each transcript was selected based on

eQTL data (GTEx_Analysis_v8_eQTL.tar) from GTEx portal release v8

(https://gtexportal.org). The DNA-binding site located within a 100-bp region around a

single-nucleotide polymorphism (SNP) of an eQTL was supposed to regulate the

expression of the transcript corresponding to the eQTL. I did not use *trans*-eQTL data,

where the SNP of an eQTL was not located on the same chromosome as the transcribed

region of the eQTL. The genomic locations of the eQTLs were converted to the

corresponding locations in the hg19 version of the human genome using the LiftOver tool

(http://hgdownload.soe.ucsc.edu/downloads.html#liftover) (Hinrichs, et al., 2006).

The original model of the input data of DNA-binding sites in promoter regions was

used for the deep learning analysis: the lengths of a DBP ChIP-seq peak in each 100-base

bin in a region between 2 kbp upstream and 1 kbp downstream of the TSS of each gene

or transcript (**Fig. 1**). To add the input data of DNA-binding sites of DBPs in more

upstream and downstream regions of a TSS and reduce the number of bins, I first

computed the length of a DBP ChIP-seq peak in a 100-base bin and then selected the

maximum length in each interval of 100 bins in a region between 1 megabase upstream

and downstream of a TSS. This step outputted the maximum 200 longest DBP ChIP-seq

peaks for each transcript. I modified the original model and extended it to include the

data.

In the training of the modified deep learning model, I used the same test, training,

and validation data as in the original method: all 2,705 genes encoded on chromosome 1

as test data, training data (22,251 genes), and validation data (2,472 genes) randomly

selected from the other genes (Osato and Hamada, 2025; Tasaki, et al., 2020). The

DeepLIFT tool was used to calculate the contribution scores of DNA-binding sites of

proteins, which are important for predicting gene expression levels (Shrikumar, et al.,

2017). The input data for promoters and putative enhancer regions were concatenated into

a total of 230 bins of the lengths of 1,310 DBP ChIP-seq peaks, consisting of 30 bins for

promoters and 200 bins for putative enhancers, because the DeepLIFT tool 'RevealCancel

rule on dense layers and the Rescale rule on conv layers (genomics default)', with high

scoring accuracy, cannot be applied to a neural network with multiple input layers.

Ubuntu (version 20.04), Python (version 3.8.10), TensorFlow (version 2.6.0), NVIDIA

CUDA toolkit (version 11.4.2), GeForce RTX 3080 GPU with 10 GB memory, and Intel

Core i7-10700 CPU with 128 GB memory were used for training and prediction, and the

original scripts were modified to run on newer versions of the software. The training

method was the same as the original, and the training time was two to four hours.

Functional prediction of DNA/RNA-binding proteins

To predict the regulatory target genes of DNA/RNA-binding proteins without using

data from protein binding assays such as ChIP-seq or eCLIP, among 1,310 DNA-binding

proteins in the input data of deep learning, 302 DNA-binding proteins with lower

contribution scores for predicting gene expression levels were replaced by other DNA-

binding proteins. For the analysis of RNA-binding proteins, 325 DNA-binding proteins

were replaced by RNA-binding proteins. The gene symbols of RNA-binding proteins

were obtained from the ENPD database (Tak Leung, et al., 2019). The interactions

between replaced DNA/RNA-binding proteins and genes regulated by the proteins were

estimated based on the co-expression pairs of genes between cells and tissues obtained

from the COEXPRESdb (Obayashi, et al., 2023). Gene co-expression is the similarity of

expression profiles between a pair of genes. Functionally related genes and genes with

the same regulation are generally expressed together between cell and tissue types. If a

gene encoding a DNA/RNA-binding protein shows a similar expression pattern with

other genes, they are potentially regulated by the protein and functionally related. The co-

expression interactions with z score > 2 between the genes encoding other DNA/RNA-

binding proteins and the genes other than DNA/RNA-binding proteins were used as input

data, replacing those with lower contribution scores. The binding sites of the DNA- and

RNA-binding protein were hypothetically located at 2 kb upstream of the TSSs and TSSs

of the co-expressed genes, respectively. Deep learning and the DeepLIFT tool with the

new input data calculated the contribution scores of the binding sites of DNA/RNA-

binding proteins for predicting gene expression levels.

For each DNA/RNA-binding protein, I examined the rank of the contribution score

of the DNA/RNA-binding site by comparing it with the contribution scores of the other

DNA/RNA-binding proteins for each gene. According to the tendency of high or low

ranks (i.e., high positive or low negative contribution scores) of DNA/RNA-binding sites

of a protein among its regulatory target genes, I selected regulatory target genes with high

or low ranks, where the rank within the top five or bottom five for each gene, of

DNA/RNA-binding sites of the protein as the primary target genes regulated by the

protein. The rank threshold for selecting primary target genes was adjusted for each

DNA/RNA-binding protein to detect functional enrichments. To obtain functional

enrichments associated with the DNA/RNA-binding protein, the rank threshold was

restricted within the top five or bottom five. The functional enrichment analysis of the

primary target genes regulated by the DNA/RNA-binding protein was performed using

'Statistical overrepresentation test' of Panther DB (https://pantherdb.org/) (Thomas, et al.,

2022). The Ensemble Gene IDs of the primary target genes were used as the analyzed

gene set, and the Ensemble Gene IDs in the gene expression data for deep leaning were

used as the reference gene set. Fisher's exact test was performed with adjustment for the

Benjamini-Hochberg false discovery rate.

To examine the binding sites of ChIP-seq data and eCLIP data around the

regulatory target genes of DNA/RNA-binding proteins predicted by deep learning of gene

co-expression data, I selected the DNA-binding sites within ±5 kbp of a TSS of a gene by

ChIP-seq assay or the RNA-binding sites between the transcription start and end of a gene

by eCLIP assay based on the Human GENCODE v19 gene annotation (Mudge, et al.,

2025). For this analysis, ChIP-seq data of 32 DNA-binding proteins and eCLIP data of

29 RNA-binding proteins were obtained from the ENCODE portal and the paper

(**Supplementary Table S6**) (Van Nostrand, et al., 2020). I also examined the ratios of the

number of genes with high and low contribution scores of the DNA/RNA-binding site

around genes with the DNA/RNA-binding site by ChIP-seq or eCLIP assays.

Figures

Figure 1. Overview of prediction of gene expression levels. The output data of gene expression levels in human cells were predicted from the input data of genomic locations of DNA-binding sites of proteins at promoters and more distant regions. In the machine learning phase, a DNA-binding protein bound to a distant region overlapping an eSNP was associated with a gene (eGene) that was potentially regulated by the DNA-binding protein based on the eSNP and eGene of the eQTL data. Based on the association of each gene, the locations of ChIP-seq peaks of DNA-binding proteins were binned (green box) and summarized in a matrix as input data.

Figure 2. Comparison of actual and predicted gene expression levels.

The predicted log2-TPM (transcripts per million) values of genes were compared with the actual expression levels in HFF cells using Spearman's rank correlation. **a** The input data for deep learning consists of the DNA-binding sites of 1,310 DNA-binding proteins from ChIP-seq data. **b** Among 1,310 proteins, 302 DNA-binding proteins with low contribution scores were replaced by the same number of DNA-binding proteins selected based on gene expression correlation data. **c** Among 1,310 proteins, 325 DNA-binding proteins with low contribution scores were replaced by the same number of RNA-binding proteins selected based on gene expression correlation data. **d** DNA-binding sites based on co-expression data showed higher contribution (DeepLIFT) scores. A and C: Binding sites of DNA-binding proteins were obtained from ChIP-seq data. B and D: Binding sites based on co-expression data between genes. A and B: Positive contribution scores. C and D: Absolute values of negative contribution scores. (Wilcoxon signed-rank test, $P <$ 2.2e-

16) **e** Example relationships between the predictive contributions of a regulator and its

expression levels across tissues. A line of best fit based on linear regression was depicted

with 95 percent confidence intervals.

Figure 3. Comparison of predicted target genes with experimental data. The *x* axis

shows the rank of the contribution score of a DNA/RNA-binding site for predicting a

gene expression level. Contribution scores of DNA/RNA-binding sites were ranked from

1 (high positive contribution scores, left end) to 1,310 (low negative contribution scores,

right end), binned with fifty rank intervals. Rank 1 on the *x* axis represents 1 to 50 ranks,

rank 2 represents 51 to 100 ranks, and the last rank 26 represents 1,251 to 1,310 ranks.

The DNA/RNA-binding sites were hypothetically located in promoters or genes bound

by the DNA/RNA-binding protein of each graph. The DNA/RNA-binding sites located

in promoters or genes bound by the DNA/RNA-binding protein show higher or lower

contribution scores than the DNA-binding sites of the ChIP-seq data as a whole. The *y*

axis shows the number of contribution scores in each bin of fifty rank intervals. Positive

contribution scores would correspond to activation and negative scores would correspond

to repression of gene expression.

Tables

Table 1. Common functional annotations between a DNA/RNA-binding protein and

its putative regulatory target genes.

The functional annotations of DNA/RNA-binding proteins were obtained from the

UniProt database. The functional annotations statistically enriched in their putative

regulatory target genes were derived from Panther DB. Gray lines separate each

annotation.

ZKSCAN5

DNA-binding transcription factor activity

C2H2 zinc finger transcription factor

RNA polymerase II cis-regulatory region sequence-specific DNA binding

metal ion binding

ZBTB34

C2H2 zinc finger transcription factor

DNA-binding transcription activator activity, RNA polymerase II-specific

RNA polymerase II transcription regulatory region sequence-specific DNA binding

ZNF557

DNA-binding transcription activator activity, RNA polymerase II-specific

RNA polymerase II transcription regulatory region sequence-specific DNA binding

ion binding, metal ion binding

ZNF446

C2H2 zinc finger transcription factor

extracellular space

ZC3H4

chromatin/chromatin-binding, or -regulatory protein

transcription regulator activity

metal ion binding

POLR2G

RNA polymerase II transcription regulatory region sequence-specific DNA binding

general transcription factor

translational protein

RPLP0

large ribosomal subunit

ribonucleoprotein complex

cytosolic large ribosomal subunit

SPEN

mRNA binding

chromatin binding

transcription coregulator activity

PKM

kinase

glycolytic process

ATP metabolic process

AKAP8

negative regulation of RNA biosynthetic process

Circadian clock system

Cry-Per complex

Table 2. Functional enrichments detected in the regulatory target genes of an RNA-

binding protein.

The functional enrichment analysis was performed by Panther DB. The table shows the

functional annotations statistically enriched in the regulatory target genes of an RNA-

binding protein after deep learning based on co-expressed genes, but not statistically

enriched in all the co-expressed genes (z score > 2) including the RNA-binding protein.

'Reference' is the number of human genes, 'Analyzed' is the number of human regulatory

target genes of an RNA-binding protein, 'Expected' is the number of genes with the

functional annotation (e.g. Circadian clock system) estimated from the ratio of genes with

the functional annotation in the reference genes, 'Over/Under' stands for

overrepresentation and underrepresentation, 'Fold Enrichment' is the ratio of the number

of analyzed genes with the functional annotation to the expected number of genes, 'raw

P-value' is the P-value of Fisher's exact test, and FDR is the Benjamini-Hochberg false

discovery rate of the test.

AKAP8							
PANTHER Pathways							
PANTHER Pathways	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
Circadian clock system (P00015)	10	3	0.03	+	> 100	2.60E-06	4.16E-04
PANTHER GO-Slim Biological Process							
PANTHER GO-Slim Biological Process	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
circadian regulation of gene expression (GO:0032922)	16	3	0.05	+	65.55	1.20E-05	1.27E-02
response to light stimulus (GO:0009416)	21	3	0.06	+	49.94	2.82E-05	1.99E-02
circadian rhythm (GO:0007623)	21	3	0.06	+	49.94	2.82E-05	1.49E-02
rhythmic process (GO:0048511)	24	3	0.07	+	43.7	4.27E-05	1.81E-02
GO cellular component complete							
GO cellular component complete	Reference (17130)	Analyzed (49)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
Cry-Per complex (GO:1990512)	3	2	0.01	+	> 100	2.40E-05	4.77E-02
PKM							
PANTHER Protein Class							
PANTHER Protein Class	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
aldolase (PC00044)	5	2	0.1	+	21.02	3.48E-03	5.68E-02
kinase (PC00137)	134	9	2.55	+	3.53	1.07E-03	2.62E-02
PANTHER GO-Slim Molecular Function							
PANTHER GO-Slim Molecular Function	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
enzyme inhibitor activity (GO:0004857)	107	8	2.04	+	3.93	1.01E-03	4.68E-02
kinase activity (GO:0016301)	562	23	10.7	+	2.15	7.08E-04	3.57E-02
PANTHER GO-Slim Biological Process							
PANTHER GO-Slim Biological Process	Reference (17130)	Analyzed (326)	Expected	Over/ Under	Fold Enrichment	raw P-value	FDR
endoplasmic reticulum tubular network organization (GO:0071786)	12	3	0.23	+	13.14	1.32E-03	3.68E-02
regulation of cellular component biogenesis (GO:0044087)	139	10	2.65	+	3.78	3.31E-04	1.05E-02

Acknowledgements

Supercomputing resources were provided by the Human Genome Center of the Institute

of Medical Science at the University of Tokyo and ROIS National Institute of Genetics.

Author contributions

N.O. designed and performed the research, and wrote and revised the manuscript.

Supplementary data

Supplementary data are available at

Conflict of interests

None declared.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS)

KAKENHI [16K00387], Osaka University start-up research grant, KIOXIA young

scientist research grant at Waseda University; and Sumitomo basic science research

projects to NO.

Data availability

Human foreskin fibroblast (HFF) cell RNA-seq data were obtained from GSM941744 in

the Gene Expression Omnibus (GEO) database. The RNA-seq data of human mammary

epithelial   cells   (HMEC)   were   found   on   ENCODE   at   UCSC

http://genome.ucsc.edu/encode/ at

'wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam'. Human neural progenitor cell

(NPC) RNA-seq data were obtained from GSM915326. HepG2 and K562 RNA-seq data

were obtained from ENCFF670LIE.bam and ENCFF584WML.bam in the ENCODE

portal (https://www.encodeproject.org/). Co-expression data of mRNAs were

downloaded from COXPRESdb (https://zenodo.org/record/6861444/files/Hsa-u.v22-

05.G16651-S245698.combat_pca.subagging.z.d.zip). The list of DNA/RNA-binding

proteins were obtained from ENPD database (http://qinlab.sls.cuhk.edu.hk/ENPD/).

ChIP-seq data of TFs were downloaded from the v20.06 release of the GTRD database

(https://gtrd.biouml.org). eQTL data (GTEx_Analysis_v8_eQTL.tar) were downloaded

from the v8 release of the GTEx portal (https://gtexportal.org). eCLIP data of RNA-

binding proteins were obtained from ENCSR456FVU in the ENCODE portal and from

Supplementary Data 2 in the paper (Van Nostrand EL et al. *Nature* 2020 doi:

10.1038/s41586-020-2077-3).

References

Agarwal, V.*, et al.* Massively parallel characterization of transcriptional regulatory elements.
Nature 2025.
Avsec, Z.*, et al.* Effective gene expression prediction from sequence by integrating long-range
interactions. *Nature methods* 2021;18(10):1196-1203.
Barrett, T.*, et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic acids*
research 2013;41(Database issue):D991-995.
Berenson, A.*, et al.* Paired yeast one-hybrid assays to detect DNA-binding cooperativity and
antagonism across transcription factors. *Nat Commun* 2023;14(1):6570.
Burley, S.K. and Kamada, K. Transcription factor complexes. *Curr Opin Struct Biol*

2002;12(2):225-230.

Edgar, R., Domrachev, M. and Lash, A.E. Gene Expression Omnibus: NCBI gene expression and

hybridization array data repository. *Nucleic acids research* 2002;30(1):207-210.

Goos, H*., et al.* Human transcription factor protein interaction networks. *Nat Commun*

2022;13(1):766.

Hinrichs, A.S*., et al.* The UCSC Genome Browser Database: update 2006. *Nucleic acids research*

2006;34(Database issue):D590-598.

Kim, S., Yu, N.K. and Kaang, B.K. CTCF as a multifunctional protein in genome regulation and

gene expression. *Exp Mol Med* 2015;47(6):e166.

Kolmykov, S*., et al.* GTRD: an integrated view of transcription regulation. *Nucleic acids research*

2021;49(D1):D104-D111.

Ladomery, M. Multifunctional proteins suggest connections between transcriptional and post-

transcriptional processes. *Bioessays* 1997;19(10):903-909.

Lee, Y. Roles of circadian clocks in cancer pathogenesis and treatment. *Exp Mol Med*

2021;53(10):1529-1538.

Lemoine, G.G*., et al.* GWENA: gene co-expression networks analysis and extended modules

characterization in a single Bioconductor package. *BMC Bioinformatics* 2021;22(1):267.

Luo, Y*., et al.* New developments on the Encyclopedia of DNA Elements (ENCODE) data portal.

Nucleic acids research 2020;48(D1):D882-D889.

McLean, C.Y*., et al.* GREAT improves functional interpretation of cis-regulatory regions. *Nature*

biotechnology 2010;28(5):495-501.

Miller, H.E. and Bishop, A.J.R. Correlation AnalyzeR: functional predictions from gene co-

expression correlations. *BMC Bioinformatics* 2021;22(1):206.

Mudge, J.M*., et al.* GENCODE 2025: reference gene annotation for human and mouse. *Nucleic*

acids research 2025;53(D1):D966-D975.

Obayashi, T*., et al.* COXPRESdb v8: an animal gene coexpression database navigating from a

global view to detailed investigations. *Nucleic acids research* 2023;51(D1):D80-D87.

Osato, N. Characteristics of functional enrichment and gene expression level of human putative

transcriptional target genes. *BMC Genomics* 2018;19(Suppl 1):957.

Osato, N. and Hamada, M. Systematic discovery of directional regulatory motifs associated with

human insulator sites. *bioRxiv* 2025.

Poelmans, G*., et al.* AKAPs integrate genetic findings for autism spectrum disorders. *Transl*

Psychiatry 2013;3(6):e270.

Puram, R.V*., et al.* Core Circadian Clock Genes Regulate Leukemia Stem Cells in AML. *Cell*

2016;165(2):303-316.

Raina, P*., et al.* GeneFriends: gene co-expression databases and tools for humans and model

organisms. *Nucleic acids research* 2023;51(D1):D145-D158.

Shrikumar, A., Greenside, P. and Kundaje, A. Learning Important Features Through Propagating

Activation Differences. In: Doina, P. and Yee Whye, T., editors, *Proceedings of the 34th*

International Conference on Machine Learning. Proceedings of Machine Learning Research:

PMLR; 2017. p. 3145--3153.

Snel, B.*, et al.* STRING: a web-server to retrieve and display the repeatedly occurring

neighbourhood of a gene. *Nucleic acids research* 2000;28(18):3442-3444.

Stuart, J.M.*, et al.* A gene-coexpression network for global discovery of conserved genetic modules.

Science (New York, N.Y.) 2003;302(5643):249-255.

Szklarczyk, D.*, et al.* The STRING database in 2023: protein-protein association networks and

functional enrichment analyses for any sequenced genome of interest. *Nucleic acids research*

2023;51(D1):D638-D646.

Tak Leung, R.W.*, et al.* ENPD - A Database of Eukaryotic Nucleic Acid Binding Proteins: Linking

Gene Regulations to Proteins. *Nucleic acids research* 2019;47(D1):D322-D329.

Tan, X.*, et al.* Novel Insights into the Circadian Rhythms Based on Long Noncoding and Circular

RNA Profiling. *Int J Mol Sci* 2024;25(2).

Tasaki, S.*, et al.* Deep learning decodes the principles of differential gene expression. *Nat Mach*

Intell 2020;2(7):376-386.

Thomas, P.D.*, et al.* PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci*

2022;31(1):8-22.

Tian, W.*, et al.* Bioinformatics analysis of the role of aldolase A in tumor prognosis and immunity.

Sci Rep 2022;12(1):11632.

UniProt, C. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic acids research*

2025;53(D1):D609-D617.

Van Nostrand, E.L.*, et al.* A large-scale binding and functional map of human RNA-binding

proteins. *Nature* 2020;583(7818):711-719.

Wang, L., Wang, S. and Li, W. RSeQC: quality control of RNA-seq experiments. *Bioinformatics*

(Oxford, England) 2012;28(16):2184-2185.

Warde-Farley, D.*, et al.* The GeneMANIA prediction server: biological network integration for

gene prioritization and predicting gene function. *Nucleic acids research* 2010;38(Web Server

issue):W214-220.

Wong, A.K., Krishnan, A. and Troyanskaya, O.G. GIANT 2.0: genome-scale integrated analysis of

gene networks in tissues. *Nucleic acids research* 2018;46(W1):W65-W70.

Yang, S.*, et al.* COEXPEDIA: exploring biomedical hypotheses via co-expressions associated with

medical subject headings (MeSH). *Nucleic acids research* 2017;45(D1):D389-D396.

Yang, Y., Pu, J. and Yang, Y. Glycolysis and chemoresistance in acute myeloid leukemia. *Heliyon*

2024;10(15):e35721.

Zhang, F*., et al.* Elevated transcriptional levels of aldolase A (ALDOA) associates with cell cycle-

related genes in patients with NSCLC and several solid tumors. *BioData Min* 2017;10:6.

Zhu, Q*., et al.* Targeted exploration and analysis of large cross-platform human transcriptomic

compendia. *Nature methods* 2015;12(3):211-214, 213 p following 214.

Zlacka, J. and Zeman, M. Glycolysis under Circadian Control. *Int J Mol Sci* 2021;22(24).