Methylation pseudotime analysis for label-free profiling of the temporal chromatin landscape with long-read sequencing

Annie Trinh[1,2,3], Navied Akhtar[2,3,4], Kwadwo Bonsu[2,5], Nandor Laszik[2,3,4], Asia Mendelevich[6], Tanye Wen[2,3,4], Julien L. P. Morival[2,3,4], Katelyn E. Diune[3], Mitchell Frazeur[3], Justin E. Vega[3], Alexander A. Gimelbrant[6], Elizabeth L. Read[2,5,7,], Timothy L. Downing[1,2,3,4,7,8,9,*]

1. Department of Microbiology and Molecular Genetics, University of California-Irvine, Irvine, CA 92697, USA
2. The NSF-Simons Center for Multiscale Cell Fate Research, University of California-Irvine, Irvine, CA 92697, USA
3. Edwards Lifesciences Cardiovascular Innovation and Research Center, University of California-Irvine, Irvine, CA 92697, USA
4. Department of Biomedical Engineering, University of California-Irvine, Irvine, CA 92697, USA
5. Department of Chemical and Biomolecular Engineering, University of California-Irvine, Irvine, CA 92697, USA
6. Altius Institute for Biomedical Sciences, Seattle, WA, USA
7. Center for Complex Biological Systems, University of California-Irvine, Irvine, CA 92697, USA
8. Sue and Bill Gross Stem Cell Research Center, University of California-Irvine, Irvine, CA 92697, USA
9. Chao Family Comprehensive Cancer Center, University of California-Irvine, Irvine, CA 92697, USA

*Corresponding author: tim.downing@uci.edu

Abstract

Faithful epigenetic inheritance across cell divisions is essential to maintaining cell identity and involves numerous epigenetic modifications, whose roles in establishing chromatin architecture are less understood. Technological approaches to temporally order epigenetic modifications throughout the cell cycle often face limitations in sequence resolution and rely on potentially damaging mitotic labeling or conversion steps. Herein, we present Methylation Pseudotime Analysis Through read-level Heterogeneity (MPATH), a label- and conversion-free method to infer post-replication DNA strand maturity from methylation patterns across single molecules. We use MPATH to temporally order hydroxymethylation throughout mitotic inheritance, revealing that CpGs within cis-regulatory elements undergo transitions between methylation states at sub-cell-cycle timescales. When applied to long reads generated by NOMe-seq, MPATH uncovered relationships between nucleosome occupancy and DNA maturity. Finally, extension of MPATH to phased reads reveals allele-specific trends in pseudotime distribution associated with X chromosome activity. Our findings suggest that when coupled with multimodal sequencing strategies, MPATH could provide valuable insights into chromatin restoration dynamics.

Main

While individual cells of a multicellular organism carry the same genetic code, the establishment and maintenance of unique cell identities is made possible by epigenetic regulation of gene expression[1,2]. To maintain cell identity following genome replication, the epigenetic landscape must be faithfully restored to nascent DNA[3]. Chromatin re-establishment is an orchestrated process involving numerous epigenetic modifications and modifiers, whose functional roles in collaboratively shaping chromatin architecture are not well understood[4,5]. Given that the epigenome needs to be de- and reassembled during each cycle of replication, the ordering of DNA modifications across sub-cell-cycle timescales could inform the collaborative processes underlying epigenetic crosstalk.

While time-lapse microscopy and mass spectrometry have enabled tracking of epigenetic modifications throughout time[6–9], these approaches often lack base-pair resolution, which may be valuable for understanding sequence-specific contexts of epigenetic inheritance. To achieve temporal resolution of epigenetic modifications through sequencing, we previously applied mitotic labeling in human embryonic stem cells (hESCs) to track CpG methylation on nascent DNA and reported a global delay in CpG remethylation following genome replication, which can be attributed to kinetic heterogeneity across CpG sites[10,11]. Other studies that employed mitotic labeling and bisulfite sequencing reported that replication-uncoupled CpG methylation can be impeded by nucleosome occupancy, suggesting that the replication-associated dynamics of epigenetic modifiers and modifications could inform how epigenetic marks propagate and influence each other throughout time[12]. While current replication-coupled sequencing approaches typically capture epigenomic data at length scales typically under 250 base pairs (bp), long-read sequencing has uncovered correlations between CpG methylation and nucleosome positioning that are otherwise masked at bulk resolution[13]. Also importantly, mitotic labeling with nucleoside analogs can be cytotoxic and influence cell identity[14–16], while bisulfite conversion can induce unwanted DNA degradation[17–19].

Herein, we present Methylation Pseudotime Analysis Through read-level Heterogeneity (MPATH), a label- and conversion-free method for determining DNA strand maturity with respect to genome replication. When coupled with other multimodal sequencing strategies, MPATH enables the temporal ordering of epigenetic modifications across sub-cell-cycle timescales with long-read and single-molecule resolution. We first demonstrated with BrdU labeling that nascent and mature chromatin can be distinguished by intramolecular CpG methylation patterns across individual long reads generated by nanopore sequencing. Then, we leveraged these patterns to infer chromatin maturity in unlabeled long reads. Benchmarking results demonstrated that

MPATH can recapitulate previously observed CpG remethylation dynamics without DNA labeling,

eliminating the need for nucleoside analogs and bisulfite conversion. We used MPATH to

temporally order CpG hydroxymethylation across replication-associated timescales in hESCs and

identified a subset of CpGs that exhibit a putative switch-like behavior between

hydroxymethylated and unhydroxymethylated states at cis-regulatory elements. When applied to

existing long-read Nucleosome Occupancy and Methylome-sequencing (NOMe-seq) datasets,

MPATH revealed an enrichment of mononucleosomes in nascent chromatin which was lost over

time. We also applied MPATH to phased long reads from Abl.1 mouse B cells and observed that

X chromosome alleles can progress differently across replication-associated timescales

according to activity. Our findings suggest that when coupled with ectopic and intrinsic DNA

modification mapping strategies, MPATH can dissect the molecular underpinnings of dynamic,

multi-factor chromatin restoration.

Results

Nanopore long-read sequencing enables detection and tracking of nascent strand DNA

methylation across sub-cell-cycle timescales

Following genome replication, the dynamics of epigenetic modifications underlying

chromatin maturation are yet to be fully elucidated (Fig. 1a). Given that global CpG methylation

is restored with a pronounced and highly variable delay in hESCs (Supplemental Fig. 1a), we

sought to leverage intramolecular CpG methylation patterns toward developing a label-free

method that would enable the temporal ordering of individual long reads (Fig. 1b). We posited

that such a method, when coupled with DNA modification mapping strategies, would be effective

toward profiling the dynamics of epigenetic modifications and modifiers across replication-

associated timescales. To demonstrate the utility of nanopore long-read sequencing in detecting

cell cycle-associated methylation states, we induced cell cycle arrest in HUES64 hESCs using 2

µg/mL nocodazole. Comparison of read-level methylation fraction (fraction of CpGs that are

methylated across single reads) to bulk methylation fraction (average bulk methylation of the

individual CpGs corresponding to those in a given read, derived from whole genome bisulfite

sequencing, or WGBS) revealed an increasing shift in read-level methylation states upon cell

cycle arrest, reflecting the suitability of the nanopore long-read sequencing platform for detecting

cell cycle-associated methylation states (Fig. 1c).

To characterize how nascent and mature chromatin differ in their intramolecular CpG

methylation states, we applied mitotic labeling in hESCs and performed nanopore sequencing

(Fig. 1d). Briefly, HUES64 hESCs were pulsed with 50 µM BrdU to label replicating DNA and

immediately collected (0h timepoint) and after 16h chase (corresponding to nascent and mature
chromatin, respectively) before sequencing. BrdU-containing reads and CpG methylation were
detected using nucleotide analog- and methylation-sensitive basecalling, respectively (Fig. 1e,
Supplemental Fig. 1b,c, and Supplemental Table 1) (Methods). On average, sequenced reads
generated 1.3x coverage per sample, with an average read length of 5.8 kb (Supplemental Table
1). We observed that CpGs from BrdU-containing reads collected at the 0h chase timepoint were
hypomethylated relative to the 16h chase timepoint and bulk CpG methylation levels obtained
from WGBS, supporting previous observations of a global delay in nascent DNA methylation (Fig.
1f). While increasing the pulse duration enabled more BrdU integration and therefore more BrdU-
containing reads, it also resulted in a tradeoff in our ability to capture nascent (versus mature)
DNA methylation levels (Fig. 1f).

To obtain a higher fraction of BrdU-labeled reads while still capturing nascent methylation,
we performed a second pulse-chase experiment using a 10-minute pulse time with a 10-fold
increase in BrdU concentration (500 µM) (Supplemental Table 1). Notably, cell viability remained
largely unchanged across a range of BrdU concentrations up to 100-fold across this short pulse
duration (Supplemental Fig. 1d). Additionally, BrdU integration did not appear to strongly favor
particular genomic regions (Supplemental Fig. 1e). Across both replicates that were pulsed with
500 µM BrdU for 10 minutes, we generated an average of 2x coverage per sample, with an
average N50 read length of 7.6 kb (Supplemental Table 1). Replicates captured a similar delay in
nascent DNA methylation levels (23.6% and 24.5% difference in bulk methylation of BrdU-
containing reads between 0 and 16h chase timepoints) (Fig. 2a).

Nascent and mature chromatin can be distinguished by intramolecular DNA methylation
patterns
Upon confirmation that BrdU-labeled nascent chromatin presented an increased
proportion of unmethylated CpG sites relative to mature chromatin at both bulk and single-
molecule resolution (Fig. 2a,b), we hypothesized that nascent and mature chromatin can be
distinguished by intramolecular CpG methylation patterns alone. We sought to devise read-level
metrics that could capture not only the proportion of methylated CpGs in a given read, but also
other features of CpG methylation patterns related to intramolecular heterogeneity. We first
computed read-level metrics previously used for quantitatively measuring epigenetic (methylation
ratio and nearest-neighbor Pearson correlation) or genetic (edit distance) differences between
DNA molecules, in addition to previously unexplored uniformity metrics (Fig. 2c, Supplemental
Note 1) (Methods). The degree of separation achieved by each read-level metric was quantified

using a squared Hellinger distance, D_H^2 (Methods) with edit distance having the highest D_H^2 value

(Fig. 2d). The distributions of read-level methylation between 0 and 16h reads suggested that

mature chromatin is highly methylated, whereas nascent chromatin exhibits widespread variability

in methylation ratios (Fig. 2e). To improve the separation between our nascent and mature read

populations, we developed a read-level edit distance metric (which computes single nucleotide

edit operations required to convert one sequence string to another[20]) that reports how similar the

methylation distribution of an individual read is to the corresponding methylation distribution in

bulk WGBS data. Edit distance appeared to reduce long-tailed variability in mature reads when

compared to distributions of the methylation ratio (Fig. 2f). Previous studies have used Pearson

correlations between nearest-neighbor CpG methylation states to quantify single-molecule

methylation heterogeneity[13]; however, this metric only modestly separated nascent and mature

reads in our data (Fig. 2g, Supplemental Fig. 2a).

To further describe read-level heterogeneity in CpG methylation across single molecules,

we used a simple matching coefficient (SMC) to calculate the proportion of matched CpG pairs in

each read (Supplemental Fig. 2b). Uniformity metrics were then derived from the SMC to compare

the proportions of matched and unmatched CpG pairs within individual reads (Fig. 2c,h). In

previous work, we reported that neighboring CpG sites exhibited correlated and faster

remethylation kinetics, which could be attributed to enzymatic processivity of DNA

methyltransferase 1 (DNMT1)[11]. To address the possibility that enzymatic processivity could

impact intramolecular CpG methylation patterns associated with nascent DNA timescales, we

also adapted these metrics to account for genomic distance between CpG pairs and methylation

states of nearest neighbors (Supplemental Fig. 2c-h).

Toward pseudotime reconstruction of epigenetic modifications across replication-

associated timescales, we used principal component analysis (PCA) to cluster our labeled reads

according to read-level heterogeneity in CpG methylation, as quantified by linear combinations of

the various read-level metrics (Fig. 2i and Supplemental Table 2). Together, our first and second

principal components explained ~79% of variance within the data, with substantial separation

between 0 and 16h chase reads, as indicated by a squared Hellinger distance of $D_H^2 = 0.447$

associated with principal component 1 (PC1) (Fig. 2i,j). Notably, PCA coefficients suggested most

metrics were weighted similarly in the calculation of PC1 (Fig. 2k).

To determine if PC1 is more effective at predicting chromatin maturity than other individual

read-level metrics, we divided the distributions of 0h and 16h chase reads into 10 equal bins

across PC1, as well as each of the 3 read-level metrics with the highest D_H^2 values: Uniformity,

Methylation Ratio, and 1-Edit Distance (Supplemental Fig. 2i-l, Supplemental Table 3). We then

calculated the proportion of reads in each bin that were derived from either the 0h or 16h chase
timepoints and found that, compared to PC1, 1-Edit Distance showed an overrepresentation of
0h chase reads across a majority of the bins, including bins with higher 1-Edit Distance values
(Supplemental Fig. 2j,l). This suggested that edit distance is less reliable for distinguishing mature
chromatin and supports the use of PC1 as a tool for pseudotime reconstruction. Notably, a
subpopulation of reads in our PCA that exhibited lower PC2 values did not appear to be temporally
dynamic (according to separation by PC1) (Fig. 2i). This subpopulation appeared to be
characterized by nearest-neighbor and short-range intramolecular CpG methylation patterns (Fig.
2k), suggesting that temporal ordering of regions with said features using MPATH could be
challenging.
214
MPATH recapitulates post-replication CpG methylation dynamics in unlabeled DNA
To investigate the genomic context of reads characterized by low PC2 values, we
performed an odds ratio analysis using H1 hESC genomic features annotated via chromHMM
(Fig. 3a,b, Supplemental Fig. 3a-b, Supplemental Table 4)[21]. We found that reads with low PC2
values enriched for genomic regions that exhibited hESC-specific hypomethylation, such as
promoters and enhancers (Fig. 3a,b, Supplemental Fig. 3a-c, Supplemental Table 4). This finding
corroborated with previous observations that genomic regions which are actively engaged in
hESCs (e.g., hESC-specific transcription factor binding sites) show invariant bulk methylation
levels across replication-associated timescales[10]. Therefore, we concluded that the separation of
nascent and mature reads could be improved by triaging reads that appear temporally stagnant
according to PC1, marked by lower PC2 values. To isolate our temporally dynamic subpopulation
of reads and increase the accuracy of identifying nascent and mature reads, we fitted lines defined
by linear inequalities based on BrdU-labeled data to maximize the percentages of true-positive
reads in the nascent and mature bins (Fig. 3c) (Methods). This filter resulted in a squared Hellinger
distance of $D_H^2 = 0.482$. The nascent and mature read bins exhibited accuracies of 98.7% (i.e.,
98.7% of labeled reads in the nascent bin were correctly identified as nascent 0h reads) and
91.9%, respectively (Fig. 3c).
We postulated that the linear combinations of read-level metrics which comprise PC1 and
PC2 can be applied to unlabeled reads from cells not treated with BrdU in order to infer nascent
and mature read populations. To this end, we sequenced a total of >30 million reads from
unlabeled HUES64 hESCs, generating >60x coverage with an N50 read length of 9.3 kb
(Supplemental Table 1). We computed read-level metrics for all reads and applied the linear
combinations comprising PC1 and PC2 derived from our labeled datasets, followed by triaging of

temporally stagnant reads (Fig. 3a,c). We name this pseudotime reconstruction method

Methylation Pseudotime Analysis Through read-level Heterogeneity (MPATH). After filtering,

MPATH enabled downstream analysis of approximately 9.5 million reads (Supplemental Table

1). To benchmark the ability of MPATH to accurately profile the dynamics of CpG remethylation

without DNA labeling, we compared the methylation difference between nascent and mature

reads from our unlabeled long-read data set (from which nascent and mature read populations

were inferred using MPATH) and publicly available Repli-Bisulfite Sequencing (Repli-BS) data

(GSE82045)[10], in which nascent and mature read populations corresponded to 0 and 16h chase

timepoints, respectively. To do this, we first identified differentially methylated regions (DMRs)

between 0 and 16h chase data in the Repli-BS study, ordered the DMRs by increasing methylation

difference, and grouped them into 10 bins called temporally dynamic tiles, as previously

described[22]. This gave us a set of tiles that were ordered according to the magnitude of their time

delay in CpG remethylation. We then computed the methylation difference between the

corresponding tiled regions captured in our inferred nascent and mature long reads. We observed

a high correlation between the methylation differences calculated from inferred versus BrdU-

labeled reads, as indicated by an R-squared value of 0.7799 (Fig. 3d), highlighting the ability of

MPATH to recapitulate post-replication CpG remethylation dynamics in the absence of DNA

labeling.

To investigate the ability of MPATH to capture previously observed CpG density-

dependent trends in methylation dynamics, we ranked CpGs according to their local density (+/-

48 bp) and calculated the local methylation difference between inferred nascent and mature reads

(Methods). In concordance with previous observations that neighboring CpG sites exhibit faster

remethylation kinetics, we found that the methylation difference between nascent and mature

reads was inversely correlated with CpG density in both inferred and Repli-BS datasets (Fig.

3e)[10,11].

We then postulated that PC1 does not only separate nascent and mature read populations

but also represents a continuum associated with chromatin maturity, from which a more explicit

pseudotime reconstruction could be derived. To test this hypothesis, we applied the linear

combinations of read-level metrics associated with PC1 and PC2 to long-read sequences

generated from previous BrdU pulse optimization experiments (Fig. 1e,f and Fig. 3f). We observed

that the 0h chase reads obtained from various BrdU pulse durations exhibited a positive

correlation between BrdU pulse duration and PC1 value (i.e., 0h chase reads appeared more

mature with increasing BrdU pulse time), thus confirming the capability of MPATH to temporally

order individual long reads across continuous replication-associated timescales (Fig. 3f).

MPATH enables simultaneous ordering of CpG methylation and hydroxymethylation

Methylated CpGs (5mCG) can be oxidized to hydroxymethylated CpGs (5hmCG) by ten-eleven translocation (TET) enzymes, which leads to active DNA demethylation through base excision repair[23–26]. Given that methylation states can vary during differentiation and in disease, the dynamics of methylation/demethylation turnover have been extensively studied across cell state transitions (e.g., over the course of embryonic development, differentiation, and post-enzymatic deletion)[27–29]. Knockouts of epigenetic writers DNMT3A/B and TETs (5mCG and 5hmCG, respectively) have also suggested that there is active enzymatic competition between DNMTs and TETs at genomic regions termed cDKO-DMRs (differentially methylated regions associated with knockout of *de novo* DNA methyltransferases DNMT3A and 3B)[30]. However, despite previous biochemical analyses suggesting that the removal and production of 5hmCG is replication-dependent[31], the dynamics of methylation/demethylation turnover during maintenance and throughout the cell cycle are not well understood. Although the post-replication maintenance of 5mCG has been extensively studied, emerging technologies that simultaneously measure 5mCG and 5hmCG have revealed that both modifications are coupled with the cell cycle and generally increase over time[9], while combinatorial quantification of these marks revealed asymmetry in 5hmCG turnover rates within CpG dyads[32]. While these approaches have enabled deeper insights into the interplay between 5mCG and 5hmCG, the dynamic coordination of these marks across sub-cell-cycle timescales remains unclear.

Furthermore, 5mCG and 5hmCG cannot be distinguished by traditional bisulfite sequencing methods, in which both marks are protected from conversion to uracil[33,34]. While oxidative bisulfite sequencing (oxBS-seq) and enzymatic methyl sequencing (EM-seq) can distinguish between these two base modifications, these technologies still rely on chemical or enzymatic conversion steps that require separate sample preparations[35,36]. In contrast, basecalling algorithms designed for nanopore sequencing can distinguish between 5mCG and 5hmCG through unique electrical traces without the need for conversion, enabling simultaneous profiling of both modifications within a single molecule[37].

Our pseudotime metrics were calculated based on a 2-state methylation caller, which only distinguishes 5mCG from unmethylated CpGs (CG). To distinguish 5mCG, 5hmCG, and CG marks across pseudotime, we used a 3-state methylation caller. We generated 1-kb tiles across the genome and intersected them with individual long reads from our unlabeled HUES64 dataset to generate 1-kb "single molecules" (Methods). We then calculated the modification ratio and modification uniformity across 10,000 randomly sampled 1-kb single molecules using both

methylation callers and ordered them according to pseudotime (Fig. 4a,b) (Methods). While the
genome-wide levels of 5mCG and 5hmCG appear to increase following replication, 5hmCG
increased to only a fraction of the level observed for 5mCG (0.209 according to the 2-state caller
and 0.264 according to the 3-state caller) (Fig. 4a). Interestingly, we found that the 2-state
methylation caller overestimates 5mCG modification ratios relative to the 3-state methylation
caller (which accounts for the presence of 5hmCG), perhaps due to the 2-state caller's inability to
distinguish 5hmCG from 5mCG. As expected, we observed a pronounced rise in 5mCG uniformity
across pseudotime when calculated based on the 2-state methylation caller. However, the 3-state
methylation caller showed a thwarted rise in 5mCG uniformity, while 5hmCG exhibited a rapid
loss in uniformity (Fig. 4b). Notably, this observation suggests that the deposition of 5hmCG is
intimately interspersed among 5mCG marks. Thus, while the propagation of 5mCG is believed to
occur *en bloc* via neighbor-guided mechanisms[11,38–40], conversion of these marks into 5hmCG
appeared to occur sporadically throughout the genome during DNA maturation.
319

MPATH identifies a subset of CpGs that alternate between methylation states at sub-cell-
cycle timescales

Given that 5hmCG turnover has been shown to be cell-cycle-dependent through bulk
biochemical analyses in mouse embryonic stem cells[31], we posited that the base-pair, single-
molecule, and temporal resolution provided by MPATH would enable us to investigate
methylation/demethylation turnover throughout sub-cell-cycle timescales in hESCs in a locus-
specific manner. We filtered for CpGs with at least 10x coverage in our unlabeled dataset (36,234
CpGs or ~11% of total CpGs captured). Each captured CpG was assigned a PC1 value based on
the read it was derived from. We then grouped CpGs into 10 pseudotime bins spanning all PC1
values. We calculated the proportion of CpGs methylated (5mCG ratio) or hydroxymethylated
(5hmCG ratio) in each pseudotime bin and clustered CpGs based on their magnitude of
hydroxymethylation across all pseudotime bins (Fig. 4c,d) (Methods). We observed that
approximately half (51.8%) of the CpG sites never experienced hydroxymethylation across
pseudotime. However, of the 17,437 CpGs that did experience hydroxymethylation, we identified
two subpopulations that appeared to be either highly or intermediately hydroxymethylated across
time (Fig. 4c,d). Interestingly, for highly hydroxymethylated CpGs, methylation and
hydroxymethylation marks co-emerge across the majority of pseudotime (Fig. 4c).
Hydroxymethylated CpGs, in general, enriched for cis-regulatory elements and repetitive regions,
which corroborates their previously reported genomic locations[30,41]. Notably, highly
hydroxymethylated CpGs enriched specifically for promoters and enhancers, as well as cDKO-

DMRs, which were previously identified as sites of DNMT3A/B and TET competition (Fig. 4e,

Supplemental Table 5)[30]. In contrast, intermediately hydroxymethylated CpGs enriched for

repetitive regions (Fig. 4f, Supplemental Table 5).

To compare the global dynamics of hydroxymethylation across pseudotime between the

two hydroxymethylated clusters, we plotted bulk hydroxymethylation ratios for each pseudotime

bin and found that while the intermediately hydroxymethylated cluster (teal) exhibited a gradual

accumulation of CpG hydroxymethylation across time, the highly hydroxymethylated cluster (red)

did not, suggesting that highly hydroxymethylated CpGs may fluctuate in hydroxymethylation

levels during DNA maturation (Fig. 4g,h). Analysis of highly hydroxymethylated CpGs at discrete

pseudotime bins 3, 6, and 9 (where only CpGs captured across all three bins were considered)

revealed a statistically significant increase, followed by decrease in bulk 5hmCG ratio (Fig. 4i).

This was in stark contrast to the monotonically increasing levels of 5hmCG observed in

intermediately hydroxymethylated CpGs (Fig. 4j). Thus, these observations revealed a subset of

CpGs with greater 5hmCG levels, which were enriched for promoters and enhancers, that actively

transition between methylation states at sub-cell-cycle timescales.

Extension of MPATH to existing NOMe-seq data enables the temporal ordering of

chromatin accessibility

The propagation of CpG methylation is often associated with nucleosome occupancy and

chromatin condensation[42–44]. However, the relationships between these epigenetic features

throughout mitotic inheritance are not well understood. To track chromatin accessibility across

sub-cell-cycle timescales and investigate the dynamic coordination between CpG methylation and

nucleosome occupancy, we applied MPATH to publicly available long-read NOMe-seq data

targeting the *H19/IGF2* imprinted locus in H9 hESCs (with >100x coverage) (GSE183760)[45].

Because the NOMe-seq long reads ranged from 73-84 kb, we created 10 kb sliding windows (with

50% overlap) across the targeted locus, which enabled us to fragment the long reads and apply

MPATH to a total of 1,934 read fragments (excluding the imprinting control region, or ICR), which

were more similar in length to our labeled reads from which MPATH was established (Fig. 5a).

We binned the 10 kb read fragments according to their inferred chromatin maturity, which resulted

in a total of 4 read subpopulations (i.e., PC1 bins) ranging from nascent reads (T1 bin) to mature

reads (T4 bin) (Methods and Fig. 5a).

Next, we sought to utilize MPATH to temporally profile chromatin accessibility (measured

through exogenous GpC methylation) at the *H19/IGF2* locus within the NOMe-seq data set. We

identified regions within read fragments that were accessible (GpC methylated) and inaccessible

(GpC unmethylated) to the M.CviPI GpC methyltransferase and observed that the distribution of

inaccessible runs was characterized by run lengths reminiscent of nucleosomes (Fig. 5b). For

example, the first peak occurs at a run length of approximately 130 bp, which is reminiscent of

the length of DNA wrapped around a mononucleosome (147 bp), whereas successive peaks

seem indicative of closely packed dinucleosomes, trinucleosomes, etc. Interestingly, separating

the read fragments according to PC1 bin revealed that T1 (nascent) read fragments were most

enriched for mononucleosomes (Fig. 5b). This result supported previous observations that

nascent chromatin is generally accessible and gradually condenses as the epigenetic landscape

is restored[46]. Accordingly, T4 (mature) read fragments were least enriched for mononucleosomes.

To validate the results obtained from coupling MPATH with NOMe-seq, we performed a

replication-coupled assay for transposase-accessible chromatin followed by sequencing (Repli-

ATAC sequencing), which applies mitotic labeling to map chromatin accessibility across

replication-associated timescales (Supplemental Fig. 4a)[47]. The distributions of insert size

presented similar trends (i.e., the enrichment of mononucleosomes within nascent reads from the

0h chase timepoint) (Fig. 5c). We also found that MPATH could identify temporal heterogeneity

within sub-nucleosomal chromatin accessibility patterns, which could potentially be attributed to

the dynamics of various chromatin-associated factors (including transcription factors) throughout

replication (Supplemental Fig. 4b).

Binning of read fragments also enabled us to investigate if various sites across a single

DNA molecule can exhibit different chromatin maturation dynamics in their local epigenetic

landscape. To this end, we calculated a weighted sum of PC1 bins for each individual read,

defined as the sum of the proportions of read fragments assigned to each weighted PC1 bin,

otherwise referred to as a PC1 weighted sum (Supplemental Note 1). We show that the PC1

weighted sum is directly proportional to the average PC1 across all 10 kb fragments of an

individual long read and clustered the long reads according to this metric (Fig. 5d,e). Interestingly,

we found that while 237 long reads (or 41.4% of total long reads) were composed of read

fragments derived from only one PC1 bin, 336 long reads (58.6%) exhibited variation in PC1

values assigned to their corresponding read fragments, suggesting that during chromatin

maturation, temporal heterogeneity can exist in single molecules across large genomic regions

(Fig. 5e).

To investigate the coordinated dynamics of nucleosome occupancy and CpG methylation

across sub-cell-cycle timescales, we first isolated NOMe-seq long reads generated from a 73 kb

targeted region within the *H19/IGF2* locus[45] and clustered captured GpCs according to their read

coverage to ensure fair evaluations of locus-specific trends in chromatin accessibility

(Supplemental Fig. 4c-e). For Cluster 5, which encompasses the ICR, we observed a decrease
in short inaccessible run lengths reminiscent of mononucleosomes throughout pseudotime (Fig.
5f, Supplemental Fig. 4f-j). Increasing chromatin inaccessibility in this cluster also appeared to
correlate with increasing CpG methylation uniformity, suggesting that nucleosome compaction
and CpG methylation uniformity are coupled across sub-cell-cycle timescales (Fig. 5g).

Phasing of long-read nanopore data reveals allele-specific trends in
pseudotime distribution associated with X chromosome activity

Previous studies have suggested that alleles can exhibit different dynamics during
genome replication[48–50]. In the context of X chromosome inactivation (XCI), the inactive X
chromosome (X_i) undergoes rapid and uniform replication during late S-phase of the cell cycle[51,52].
Thus, we posited that MPATH could discern X chromosome activity based on allelic differences
in pseudotime distribution. We sequenced genomic DNA from Abl.1 mouse B cells to >40x
coverage (N50 read length: 16.1 kb) and phased reads at a given 1 Mbp locus on the X
chromosome (chrX:103,530,980-104,531,001) into two haplotypes (Fig. 6a, Methods). We
determined that haplotype 1 was derived from X_i due to the unmethylated state of the *Xist*
promoter (Fig. 6a). We then created 20 kb windows across the locus and identified windows at
which both haplotypes exhibited similar bulk CpG methylation levels (<15% bulk CpG methylation
difference) to mitigate the influence of XCI-associated methylation differences on pseudotime
reconstruction (Fig. 6b). Then, we applied MPATH to all reads within these windows and found
that reads derived from X_i exhibited a more uniform distribution of PC1, which could reflect more
uniform (and rapid) chromatin maturation kinetics associated with X_i (Fig. 6c)[51]. In contrast,
phased reads derived from an autosome (chr10) exhibited very similar distributions of PC1, thus
suggesting that alleles can progress differently across replication-associated timescales in the
case of monoallelic expression (Fig. 6d).

Discussion

In this study, we presented a novel label- and conversion-free method that leverages
intramolecular CpG methylation patterns for methylation pseudotime analysis of single DNA
molecules. We demonstrated that when coupled with DNA modification mapping strategies,
MPATH can be particularly valuable for temporally ordering epigenetic modifications without the
need for cytotoxic nucleoside analogs or chemical conversion steps. Although actively engaged
regions exhibit stagnant mean methylation levels throughout the cell cycle[10], applying MPATH to
long-read sequences enabled the temporal ordering of these elements using surrounding regions

within individual reads. Coupling MPATH with a 3-state methylation caller enabled unprecedented
temporal resolution into DNA methylation/demethylation turnover during stem cell maintenance,
revealing a subset of CpGs which alternate between hydroxymethylated and
unhydroxymethylated states at sub-cell-cycle timescales within cis-regulatory elements,
potentially reflecting competition between DNA methylation writer and eraser enzymes[28,30,53–55].
We also showed that retroactive application of MPATH could provide temporal context to an
existing long-read NOMe-seq dataset, revealing patterns of CpG methylation and nucleosome
occupancy that were coordinated in time. These findings support the potential of MPATH as a
framework for investigating the extent to which these coordinated patterns are shaped through
post-replication epigenetic crosstalk. Finally, extension of MPATH to phased reads revealed how
alleles can progress differently through replication-associated timescales according to X
chromosome activity. Therefore, by providing temporal context to phased long-read datasets,
MPATH offers a novel approach for studying the mechanistic underpinnings of monoallelic gene
expression.

MPATH utilizes linear combinations of discrete read-level metrics generated based on
PCA to infer chromatin maturity. In contrast to other trajectory inference methods that utilize
machine learning or dynamical modeling[56–59], the advantages of our approach include the ability
to benchmark our pseudotime scores against "ground truth" labeled data and identify specific
features of methylation patterning that drive pseudotime reconstruction. Many multimodal
sequencing strategies have already been adapted to long-read sequencing platforms, which will
support diverse applications of MPATH[60–62]. Future efforts will benefit from extending MPATH to
other cell types and synthetic cellular systems. We posit that MPATH will be especially useful for
deciphering epigenetic crosstalk and understanding chromatin dynamics during replication and
cell state transitions at sub-cell-cycle time resolution.

Methods

Cell culture

HUES64 hESCs were cultured on Geltrex (Thermo Fisher Scientific, A1413302) coated tissue
culture plates with mTeSR (STEMCELL Technologies) media. Cells were incubated at 37°C with
5% CO_2. Media was changed daily and cells were passaged upon reaching 80% confluency using
ReleSR (STEMCELL Technologies). v-Abl pro-B clonal cell line Abl.1, derived previously from
129S1/SvImJ × CAST/EiJ F1 female mice[63], were cultured in RPMI medium (Gibco) containing
15% FBS (Sigma), 1X L-Glutamine (Gibco), 1X Penicillin/Streptomycin (Gibco), and 0.1% β-
mercaptoethanol (Sigma).

Cell cycle arrest

hESCs were treated with 2 µg/ml nocodazole for 16 h, collected, and washed twice with DPBS (Gibco, 14190144) prior to genomic DNA extraction by phenol-chloroform extraction.

BrdU labeling

To label nascent DNA, 5-bromo-2′-deoxyuridine (BrdU) (Sigma-Aldrich, B5002) was added to the cell culture medium at a final concentration of 50 µM or 500 µM. Cells were incubated in labeled medium for 10 minutes at 37°C with 5% CO_2 (for BrdU pulse optimization, incubation times of 20, 30, and 60 minutes were also used). Next, nascent samples were washed once with DPBS (Gibco, 14190144), and cells were collected using Accutase (STEMCELL Technologies), pelleted, and fixed in 75% ethanol by adding 100% ethanol dropwise to the cell resuspension until a final ethanol concentration of 75% was reached. In contrast, mature samples were washed three times with DPBS, and further incubated in fresh mTeSR. At 16 hours of incubation, mature samples were collected, pelleted, and fixed in 75% ethanol as previously described.

Assessing cell viability

Collected cells were stained using the LIVE/DEAD™ Fixable Violet Dead Cell Stain Kit (Thermo Fisher Scientific, L34955) according to the manufacturer's instructions. Samples were analyzed via flow cytometry using a NovoCyte 3000 (Agilent Technologies).

Genomic DNA isolation

For samples that were arrested with nocodazole or labeled with 50 µM BrdU, cells were lysed via incubation in SDS-PK buffer (0.5% wt/vol) sodium dodecyl sulfate (SDS), 50 mM Tris-HCl, 0.01 M EDTA, 1 M NaCl and 0.2 mg/ml proteinase K) at 56 °C for 2 h. Genomic DNA was isolated using phenol-chloroform extraction and isopropanol precipitation. For samples labeled with 500 µM BrdU, genomic DNA was isolated using the Puregene Cell Kit (8 x 10^8) (Qiagen, 158043) or Monarch® Genomic DNA Purification Kit (NEB, T3010S) according to the manufacturer's instructions. Abl.1 B cells were collected on day 5, and debris were removed using Histopaque. DNA was prepared using Sigma GenElute kit, and then enriched for longer fragments using Circulomics SRE XS kit.

DNA shearing for nanopore sequencing

509 For 500 µM BrdU labeled samples and unlabeled samples, genomic DNA was sheared to
510 approximately 6-8 kb using g-TUBEs (Covaris, SKU 520079) by spinning 2-3 µg of unfragmented
511 gDNA at 7,200 rpm for 1 min in a Pico 21 Microcentrifuge (Thermo Scientific). Then, tubes were
512 inverted and spun down under the same conditions.
513

Nanopore sequencing
For samples labeled with 50 µM BrdU, libraries were prepared from purified and fragmented
genomic DNA using a Ligation Sequencing Kit (Oxford Nanopore Technologies, SQK-LSK110)
according to the manufacturer's instructions. Briefly, DNA repair and end-prep were performed
using NEBNext® FFPE DNA Repair Mix (NEB, M6630) and NEBNext® Ultra II End Repair/dA-
tailing Module (NEB, E7546). Adapter ligation and clean-up were performed using the NEBNext®
Quick Ligation Module (NEB, E6056). Long Fragment Buffer from the Ligation Sequencing Kit
was used in the clean-up step to enrich for DNA fragments of >3 kb. Libraries were loaded onto
R9.4.1 flow cells (Oxford Nanopore Technologies, FLO-MIN106D) and sequenced using a
MinION Mk1b device (Oxford Nanopore Technologies) for 16 hours. Minimum read length was
set to 200 bp. For Abl.1 B cells, libraries were prepared similarly, loaded onto R9.4.1 flow cells,
and sequenced using a PromethION P2 solo device (Oxford Nanopore Technologies). For
samples labeled with 500 µM BrdU and samples not labeled with BrdU, libraries were prepared
using a Ligation Sequencing Kit (Oxford Nanopore Technologies, SQK-LSK114) according to the
manufacturer's instructions, similarly to previously described. Libraries for samples labeled with
500 µM BrdU were loaded onto R10.4.1 flow cells (Oxford Nanopore Technologies, FLO-MIN114)
and sequenced using a MinION Mk1b device for 32 hours. Libraries generated from unlabeled
samples were loaded onto R10.4.1 flow cells (Oxford Nanopore Technologies, FLO-PRO114M)
and sequenced on a PromethION P2 solo device (Oxford Nanopore Technologies) for 72 hours.
Minimum read length was set to 1000 bp. All sequencing data was collected using MinKNOW
v23.07.5.
535

Preprocessing of nanopore data
For BrdU-labeled samples loaded on R9.4.1 flow cells (Oxford Nanopore Technologies, FLO-
MIN106D), basecalling on raw current signals was performed using Guppy v6.5.7. DNA
sequences were aligned to the hg19 human reference genome using Minimap2 v2.24 and
chimeric reads were removed using samtools v1.15.1. To identify BrdU-containing reads,
DNAscent v3.1.2[64] was used to detect the BrdU base analog within individual reads. CpG
methylation was called using nanopolish v0.13.2[65]. For BrdU-labeled samples loaded on R10.4.1

flow cells (Oxford Nanopore Technologies, FLO-MIN114) and unlabeled samples loaded on
R10.4.1 flow cells (Oxford Nanopore Technologies, FLO-PRO114M), basecalling and alignment
to the hg19 human reference genome were performed using Dorado v0.6.2. Chimeric reads were
removed using samtools v1.15.1. To identify BrdU-containing reads, DNAscent v4.0.1 was used
to detect the nucleoside analog within individual reads. CpG methylation was called using Dorado
v0.6.2. Nanoplot v1.41.6 was used to assess and visualize read-level statistics such as number
of reads, read length N50, mean and median read length, and mean and median read quality. For
mouse B cells raw nanopore reads were basecalled and CpG methylation detection was done
using Guppy v6.3.9 (basecall_model=2021-05-05_dna_r9.4.1_promethion_384_dd219f32).
Reads were aligned to the mm10 mouse reference genome using Minimap2 v2.22 with default
parameters.

Calculating read-level metrics
For each individual read, average BrdU is defined as the average of the probabilities that each
thymidine is BrdU, as determined from DNAscent v4.0.1. A mock-treated sample with no BrdU
was used to determine a threshold average BrdU value, above which a read is determined to be
BrdU-containing. Methylation ratio is defined as the ratio of total methylated CpGs to total CpGs
within the read. The uniformity of a read is defined as the proportion of matched CpG pairs minus
the proportion of unmatched CpG pairs within the read. Detailed information and equations for all
read-level metrics, including metrics that are adapted to account for nearest neighbors or genomic
distance, can be found in the Supplemental material (Supplemental Note 1).

Dimensionality reduction for methylation pseudotime analysis
Read-level metrics were integrated into a principal component analysis (PCA) to enable
methylation pseudotime analysis. PCA was conducted using MATLAB (R2023b).

Applying linear combinations generated by MPATH to unlabeled data
Each principal component in the PCA on labeled data was characterized by a linear combination
of the input read-level metrics, where each individual read-level metric was associated with a
unique coefficient for each principal component. Read-level metrics were calculated for reads
derived from unlabeled data. Then, the coefficients of the linear combinations comprising the
principal components derived from labeled data were applied to the read-level metrics calculated
from unlabeled data.

Sourcing genomic annotations

All annotated genomic regions were previously inferred through chromHMM[21] and downloaded

from the UCSC Table Browser[66]. Promoter regions were defined as 2 kb upstream and 500 bp

downstream of transcription start sites (TSS). Intergenic regions were determined using the

BEDTools "subtract" function[67]. The BEDTools "intersect" function was used to determine which

regions were captured by sequenced long reads.

Enrichment analyses

Odds ratio (OR) analyses were performed to determine the strength of the association of read

populations of interest or CpGs of interest with particular genomic regions. For enrichment

analysis of low PC2 reads, OR was calculated as $\frac{a/c}{b/d}$, where a = number of reads that are

categorized as low PC2 and capture the genomic region of interest, b = number of reads that are

categorized as low PC2 and do not capture the genomic region of interest, c = number of reads

that are not categorized as low PC2 reads and capture the genomic region of interest,

and d = number of reads that are not categorized as low PC2 reads and do not capture the

genomic region of interest. For enrichment analysis of hydroxymethylation clusters, a = number

of CpGs within a given cluster that occur within the genomic region of interest, b = number of

CpGs within a given cluster that occur outside of the genomic region of interest, c = number of

CpGs that are not within a given cluster that occur within the genomic region of interest,

and d = number of CpGs that are not within a given cluster that occur outside of the genomic

region of interest. The logarithmic OR value (log(OR)) was calculated for each genomic region of

interest. The Fisher's exact test was used to determine the statistical significance of ORs.

Heatmaps were created using the R package pheatmap v1.0.12.

Binning read populations across pseudotime

Three linear inequalities across PC1 and PC2 defined the population of reads that can be

analyzed by MPATH (Supplemental Note 1). These linear inequalities were manually optimized

to encapsulate reads that are temporally dynamic, as determined by the placement of BrdU-

labeled reads. Two linear inequalities were then curated to define nascent and mature reads

within this population. The latter linear inequalities were manually optimized to maximize the

proportion of true nascent reads in the nascent bin and true mature bins in the mature bin. To

create 4 bins across pseudotime, the population of reads that were not categorized as nascent or

mature were split into 2 more bins according to the PC1 value associated with the peak of the

BrdU 10 minute pulse, 0h chase read distribution across PC1 (Figure 3F).

Quantifying CpG hydroxymethylation across pseudotime

CpG hydroxymethylation was called on long reads derived from unlabeled hESCs using Dorado v0.6.2, as described above. Each read was assigned a PC1 value using MPATH as described above. To profile CpG hydroxymethylation across pseudotime globally, we generated 1 kb tiles of the hg19 human reference genome and intersected this with long reads in order to create 1 kb single molecules. We calculated the methylation ratio and uniformity across 10,000 randomly sampled 1 kb single molecules (Supplemental Note 1) and plotted these values against the PC1 value corresponding to the read from which the 1 kb single molecule was derived as a smooth curve using the R package ggplot2 v3.5.1. All possible PC1 values from this data set were categorized into 10 equal-sized bins. To profile CpG hydroxymethylation across pseudotime locus-specifically, we first filtered for CpGs with $\geq$10x coverage. For each of these CpGs, we calculated the proportion of reads within each PC1 bin that capture that CpG in a hydroxymethylated or methylated state. This data was plotted as a heatmap using pheatmap v1.0.12. Clusters of hydroxymethylated CpGs were determined by first categorizing CpGs into CpGs that experience hydroxymethylation vs no hydroxymethylation. Then, the hydroxymethylated cluster was divided evenly into 2 sub-clusters, where the first cluster is considered highly hydroxymethylated and the second cluster is considered intermediately hydroxymethylated.

Processing publicly available NOMe-seq data

Raw nanopore reads from NOMe-seq conducted on H9 hESCs were downloaded in fast5 format from the National Center for Biotechnology Information (NCBI) under Bioproject ID PRJNA510783. Basecalling, CpG methylation calling, GpC methylation calling, and alignment to the hg19 human reference genome were all performed using Dorado v0.6.2. Chimeric reads were removed using samtools v1.15.1. To avoid ambiguous methylation outputs, CpGs and GpCs in a GCG context were excluded for downstream analysis (26.4% of CpGs and 18.5% of GpCs).

Inaccessible Run Lengths

We first created 10 kb windows of the hg38 human reference genome and intersected them with the CpG and GpC methylation data from NOMe-seq long reads to create 10 kb single molecules. The start and end coordinates of runs within a molecule were defined as the midpoint between the first or last methylated GpC and nearest unmethylated GpC. Inaccessible runs were characterized by runs in which all GpCs were unmethylated.

Repli-ATAC sequencing

HUES64 hESCs were grown to ~70% confluency in a 6-well plate. Different chase timepoints
were assigned to different wells. To label nascent DNA, cell culture medium was replaced with 2
mL/well of mTeSR containing 50 µM BrdU. Cells were incubated in labeled medium for 1 hour at
37°C with 5% CO_2. Next, 0h chase samples were washed once with mTeSR, and cells were
collected using Accutase. For later chase timepoints, two washes with mTeSR were done before
incubating the cells at 37°C with 5% CO_2 until the chase timepoint was reached. Then, cells were
collected using Accutase. Collected cells were resuspended in 1 mL ice-cold DPBS buffer and
100K cells were transferred to and pelleted in a 1.5 mL Eppendorf tube at 4°C. The pellet was
resuspended in 50 uL cold lysis buffer (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.1%
IGEPAL CA-630). Nuclei were spun down immediately at 500 x g for 10 minutes at 4°C,
supernatant was discarded, and the pellet was stored on ice.

To initiate the transposition reaction, the nuclei were resuspended in transposition reaction mix:
50 µL TD (2x reaction buffer, Illumina, FC-121-1030), 5 µL TDE1 (Nextera Tn5 Transposase,
Illumina, FC-121-1030), and 45 µL nuclease-free water (Invitrogen, 10977-015). The transposition
reaction was incubated on a thermomixer at 37°C for 30 minutes. Then, tagmented DNA was
purified using the MinElute PCR Purification kit (Qiagen, 28004). To amplify transposed DNA
fragments from each sample, the following was combined in a 0.2 mL PCR tube: 20 µL transposed
DNA, 5 µL nuclease-free water, 25 µL NEBNext High-Fidelity 2x PCR Master Mix (NEB, M0541).
Samples were placed in a thermal cycler for 1 cycle of 72°C for 5 minutes to allow for extension
of both ends of the primer after transposition. DNA was purified using the MinElute PCR
Purification kit.

10 µL of each sample was used for BrdU immunoprecipitation (IP) and 10 µL was stored aside at
4°C as a non-IP control. To denature the DNA in the IP sample, 490 µL of TE buffer was added
and the sample was incubated at 95°C for 5 minutes on a thermomixer before cooling in an ice
bath. The denatured DNA was collected and added to 60 µL of 10x IP buffer (28.5 mL nuclease-
free water, 5 mL of 1 M sodium phosphate, pH 7.0, 14 mL of 5 M NaCl, and 2.5 mL of 10% wt/vol
Triton X-100). 40 µL of 12.5 µg/mL anti-BrdU antibody (BD Biosciences, 555627) was added, and
the sample was incubated for 1 hour at room temperature with constant rotation. The sample was
centrifuged at 17,000 x g for 5 minutes at 4°C and the supernatant was completely removed. 750
µL of ice-cold 1x IP buffer was added before centrifuging and removing the supernatant again.

The pellet was then resuspended in 200 μL digestion buffer (50 mM Tris-HCl, pH 8.0, 10 mM

EDTA, 0.5% SDS) with 0.25 mg/mL Proteinase K and incubated overnight at 37°C, then for 60

minutes at 56°C. Next, the sample was purified using AMPure XP Beads (Beckman Coulter,

A63881) (2x bead volume) and resuspended in 20 μL Elution Buffer. To amplify the transposed

DNA fragments, the following was combined in a 0.2 mL PCR tube: 20 μL transposed DNA, 2.5

μL 25 μM Custom Nextera PCR Primer 1, 2.5 μL 25 μM Custom Nextera PCR Primer 2, 25 μL

NEBNext High-Fidelity 2x PCR Master Mix. The following PCR conditions were used: 1 cycle of

72°C for 5 minutes, 98°C for 30 seconds, followed by 5 cycles of 98°C for 10 seconds, 63°C for

30 seconds, 72°C for 1 minute. The PCR reaction was monitored using qPCR to stop amplification

before saturation, in order to reduce GC and size bias[68]. Finally, the amplified library was purified

using the MinElute PCR Purification kit.

Repli-ATAC sequencing data processing

Fastq files were trimmed using Trim Galore v0.4.4[69] and aligned using BWA[70]. Samtools v1.15.1

was used to remove mitochondrial DNA, duplicate reads, and unmapped reads. BamTools[71] was

used to remove reads containing more than 4 mismatches and reads with insert size greater than

2 kb. Peak calling was performed using MACS2[72]. To quantify insert sizes, the

CollectInsertSizeMetrics function of Picard[73] was used with M=0.5 for each bam file. Histogram

counts were merged across 2 replicates and plotted as a density of all reads as normalization.

Phasing long-read data

To phase the long reads from the publicly available NOMe-seq data set (GSE183760), reads that

capture the imprinting control region (ICR) at the *H19/IGF2* locus were assigned an MPATH score

according to their intramolecular CpG methylation patterns, excluding the CpGs within the ICR.

To determine if a read was derived from the maternal or paternal allele, we first calculated the

fraction of methylated CpGs in the ICR for each read. The ICR was considered to be methylated

if the fraction of methylated CpGs was >0.5 and unmethylated if the fraction of methylated CpGs

was <0.5. Reads with methylated ICRs were assigned to the paternal allele while reads with

unmethylated ICRs were assigned to the maternal allele, in corroboration with the known

methylation states of the paternal and maternal ICRs for the *H19/IGF2* locus. Genomic variants

for the 129S1/SvImJ and CAST/EiJ mouse strains were obtained from The Jackson Laboratory

database. Homozygous SNP sites from each strain were used to construct a VCF file representing

the F1 cross. Read phasing was performed using Whatshap v2.0.

Statistical analyses

Statistical analyses were performed using GraphPad Prism v9.3.1 for Windows (linear regression

models) and in R v4.0.3 (Fisher's exact test).

Data availability

Raw sequencing data and processed data for this study are available through the Gene

Expression Omnibus. Data associated with Figures 1-5 can be accessed at GSE290978

(Nanopore) and GSE290974 (ATAC-seq). Raw data in POD5 or FAST5 format for Figures 1-5

are available upon request. Both raw and processed data associated with Figure 6 can be

accessed at GSE286139.

Code availability

Source code for analysis is available at https://github.com/downinglab/mpath/.

Acknowledgements

We thank M. Byun for insightful discussions and M. Boemo for sharing a pre-release version of

DNAscent that helped toward preliminary analyses for this study. This work was partially

supported by NSF grants (EF2022182 and DMS1763272), a grant from the Simons Foundation

(594598, QN), and an NIH New Innovator Award (DP2) Grant DP2CA250382-01 to T.L.D. A.T.

was supported by a training grant from the California Institute for Regenerative Medicine under

Award Number EDUC4-12822. The content is solely the responsibility of the authors and does

not necessarily represent the official views of the California Institute for Regenerative Medicine.

Author contributions

T.L.D., E.L.R., A.A.G., and A.T. conceptualized this study and supervised all experiments and

data analysis. A.T., N.A., J.L.P.M., K.D., M.P.F., and J.V. performed experiments. A.T., N.A., K.B.,

N.L., A.M., T.W., and M.P.F. processed and analyzed data. All authors contributed to data

interpretation. A.T. wrote the original draft of the manuscript. T.L.D., E.L.R., A.A.G., and A.T.

reviewed and edited the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

1.   Barrero, M. J., Boué, S. & Izpisúa Belmonte, J. C. Epigenetic Mechanisms that Regulate
Cell Identity. *Cell Stem Cell* **7**, 565–570 (2010).

2.   Nashun, B., Hill, P. W. & Hajkova, P. Reprogramming of cell fate: epigenetic memory
and the erasure of memories past. *EMBO J* **34**, 1296–1308 (2015).

3.   Stewart-Morgan, K. R., Petryk, N. & Groth, A. Chromatin replication and epigenetic cell
memory. *Nat Cell Biol* **22**, 361–371 (2020).

4.   Biswas, S. & Rao, C. M. Epigenetic tools (The Writers, The Readers and The Erasers)
and their implications in cancer therapy. *European Journal of Pharmacology* **837**, 8–24 (2018).

5.   Gillette, T. G. & Hill, J. A. Readers, writers, and erasers: chromatin as the whiteboard of
heart disease. *Circ Res* **116**, 1245–1253 (2015).

6.   Saxton, M. N., Morisaki, T., Krapf, D., Kimura, H. & Stasevich, T. J. Live-cell imaging
uncovers the relationship between histone acetylation, transcription initiation, and nucleosome
mobility. *Sci. Adv.* **9**, eadh4819 (2023).

7.   Lungu, C., Pinter, S., Broche, J., Rathert, P. & Jeltsch, A. Modular fluorescence
complementation sensors for live cell detection of epigenetic signals at endogenous genomic
sites. *Nat Commun* **8**, 649 (2017).

8.   Fujimori, T. *et al.* Single-cell chromatin state transitions during epigenetic memory
formation. *bioRxiv* 2023.10.03.560616 (2023) doi:10.1101/2023.10.03.560616.

9.   Stewart-Morgan, K. R. *et al.* Quantifying propagation of DNA methylation and
hydroxymethylation with iDEMS. *Nat Cell Biol* **25**, 183–193 (2023).

10.   Charlton, J. *et al.* Global delay in nascent strand DNA methylation. *Nat Struct Mol Biol*
25, 327–332 (2018).

11.   Busto-Moner, L. *et al.* Stochastic modeling reveals kinetic heterogeneity in post-
replication DNA methylation. *PLoS Comput Biol* **16**, e1007195 (2020).

12.   Ming, X. *et al.* Kinetics and mechanisms of mitotic inheritance of DNA methylation and
their roles in aging-associated methylome deterioration. *Cell Res* **30**, 980–996 (2020).

13.   Kerr, L., Kafetzopoulos, I., Grima, R. & Sproul, D. Genome-wide single-molecule
analysis of long-read DNA methylation reveals heterogeneous patterns at heterochromatin that
reflect nucleosome organisation. *PLoS Genet* **19**, e1010958 (2023).

14.   Haskins, J. S. *et al.* Evaluating the Genotoxic and Cytotoxic Effects of Thymidine
Analogs, 5-Ethynyl-2′-Deoxyuridine and 5-Bromo-2′-Deoxyurdine to Mammalian Cells. *IJMS* **21**,
6631 (2020).

15.   Schneider, L. & d'Adda Di Fagagna, F. Neural stem cells exposed to BrdU lose their
global DNA methylation and undergo astrocytic differentiation. *Nucleic Acids Research* **40**,
5332–5342 (2012).

16.   Desai, R. V. *et al.* A DNA repair pathway can regulate transcriptional noise to promote
cell fate transitions. *Science* **373**, eabc6506 (2021).

17.   Olova, N. *et al.* Comparison of whole-genome bisulfite sequencing library preparation
strategies identifies sources of biases affecting DNA methylation data. *Genome Biol* **19**, 33
(2018).

18.   Tanaka, K. & Okamoto, A. Degradation of DNA by bisulfite treatment. *Bioorganic &*
Medicinal Chemistry Letters **17**, 1912–1915 (2007).

19.   Grunau, C., Clark, S. J. & Rosenthal, A. Bisulfite genomic sequencing: systematic
investigation of critical experimental parameters. *Nucleic Acids Res* **29**, E65-65 (2001).

20.   Kluin, R. J. C. *et al.* XenofilteR: computational deconvolution of mouse and human reads
in tumor xenograft sequence data. *BMC Bioinformatics* **19**, 366 (2018).

21.   Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and
characterization. *Nat Methods* **9**, 215–216 (2012).

22.   Morival, J. L. Understanding the molecular and functional consequences of epigenome
dynamics in cell fate, aging, and disease. *UC Irvine.* doi:ProQuest ID:
Morival_uci_0030D_17136. Merritt ID: ark:/13030/m53550hn.

23.   Zhang, X., Zhang, Y., Wang, C. & Wang, X. TET (Ten-eleven translocation) family
proteins: structure, biological functions and applications. *Sig Transduct Target Ther* **8**, 297
(2023).

24.   Zhang, M. *et al.* Ten-eleven translocation 1 mediated-DNA hydroxymethylation is
required for myelination and remyelination in the mouse brain. *Nat Commun* **12**, 5091 (2021).

25.   Santiago, M., Antunes, C., Guedes, M., Sousa, N. & Marques, C. J. TET enzymes and
DNA hydroxymethylation in neural development and function — How critical are they?
Genomics **104**, 334–340 (2014).

26.   Kohli, R. M. & Zhang, Y. TET enzymes, TDG and the dynamics of DNA demethylation.
Nature **502**, 472–479 (2013).

27.   Seisenberger, S. *et al.* The Dynamics of Genome-wide DNA Methylation
Reprogramming in Mouse Primordial Germ Cells. *Molecular Cell* **48**, 849–862 (2012).

28.   Rulands, S. *et al.* Genome-Scale Oscillations in DNA Methylation during Exit from
Pluripotency. *Cell Systems* **7**, 63-76.e12 (2018).

29.   Ginno, P. A. *et al.* A genome-scale map of DNA methylation turnover identifies site-
specific dependencies of DNMT and TET activity. *Nat Commun* **11**, 2680 (2020).

30.   Charlton, J. *et al.* TETs compete with DNMT3 activity in pluripotent cells at thousands of
methylated somatic enhancers. *Nat Genet* **52**, 819–827 (2020).

31.   Otani, J. *et al.* Cell Cycle-Dependent Turnover of 5-Hydroxymethyl Cytosine in Mouse
Embryonic Stem Cells. *PLoS ONE* **8**, e82961 (2013).

32.   Chialastri, A., Sarkar, S., Schauer, E. E., Lamba, S. & Dey, S. S. Combinatorial
quantification of 5mC and 5hmC at individual CpG dyads and the transcriptome in single cells
reveals modulators of DNA methylation maintenance fidelity. *Nat Struct Mol Biol* **31**, 1296–1308
(2024).

33.   Kernaleguen, M. *et al.* Whole-Genome Bisulfite Sequencing for the Analysis of Genome-
Wide DNA Methylation and Hydroxymethylation Patterns at Single-Nucleotide Resolution. in
Epigenome Editing (eds. Jeltsch, A. & Rots, M. G.) vol. 1767 311–349 (Springer New York, New
York, NY, 2018).

34.   Derbala, D., Garnier, A., Bonnet, E., Deleuze, J.-F. & Tost, J. Whole-Genome Bisulfite
Sequencing Protocol for the Analysis of Genome-Wide DNA Methylation and
Hydroxymethylation Patterns at Single-Nucleotide Resolution. in *Epigenome Editing* (eds.
Jeltsch, A. & Rots, M. G.) vol. 2842 353–382 (Springer US, New York, NY, 2024).

35.   Booth, M. J. *et al.* Oxidative bisulfite sequencing of 5-methylcytosine and 5-
hydroxymethylcytosine. *Nat Protoc* **8**, 1841–1851 (2013).

36.   Vaisvila, R. *et al.* Enzymatic methyl sequencing detects DNA methylation at single-base
resolution from picograms of DNA. *Genome Res* **31**, 1280–1289 (2021).

37.   Liu, Y. *et al.* DNA methylation-calling tools for Oxford Nanopore sequencing: a survey
and human epigenome-wide evaluation. *Genome Biol* **22**, 295 (2021).

38.   Hermann, A., Goyal, R. & Jeltsch, A. The Dnmt1 DNA-(cytosine-C5)-methyltransferase
Methylates DNA Processively with High Preference for Hemimethylated Target Sites. *Journal of*
Biological Chemistry **279**, 48350–48359 (2004).

39.   Svedružić, Ž. M. & Reich, N. O. Mechanism of Allosteric Regulation of Dnmt1's
Processivity. *Biochemistry* **44**, 14977–14988 (2005).

40.   Ren, H., Taylor, R. B., Downing, T. L. & Read, E. L. Locally correlated kinetics of post-
replication DNA methylation reveals processivity and region specificity in DNA methylation
maintenance. *J. R. Soc. Interface.* **19**, 20220415 (2022).

41.   Marion-Poll, L. *et al.* DNA methylation and hydroxymethylation characterize the identity
of D1 and D2 striatal projection neurons. *Commun Biol* **5**, 1321 (2022).

42.   Ren, W. *et al.* Direct readout of heterochromatic H3K9me3 regulates DNMT1-mediated

maintenance DNA methylation. *Proc Natl Acad Sci USA* **117**, 18439–18447 (2020).

43.   Buitrago, D. *et al.* Impact of DNA methylation on 3D genome structure. *Nat Commun* **12**,

3243 (2021).

44.   Collings, C. K. & Anderson, J. N. Links between DNA methylation and nucleosome

occupancy in the human genome. *Epigenetics & Chromatin* **10**, 18 (2017).

45.   Battaglia, S. *et al.* Long-range phasing of dynamic, tissue-specific and allele-specific

regulatory elements. *Nat Genet* **54**, 1504–1513 (2022).

46.   Ostrowski, M. S. *et al.* The single-molecule accessibility landscape of newly replicated

mammalian chromatin. *Cell* S0092867424012558 (2024) doi:10.1016/j.cell.2024.10.039.

47.   Stewart-Morgan, K. R., Reverón-Gómez, N. & Groth, A. Transcription Restart

Establishes Chromatin Accessibility after DNA Replication. *Molecular Cell* **75**, 284-297.e6

(2019).

48.   Mostoslavsky, R. *et al.* Asynchronous replication and allelic exclusion in the immune

system. *Nature* **414**, 221–225 (2001).

49.   Singh, N. *et al.* Coordination of the random asynchronous replication of autosomal loci.

Nat Genet **33**, 339–341 (2003).

50.   Gimelbrant, A. A. & Chess, A. An epigenetic state associated with areas of gene

duplication. *Genome Res.* **16**, 723–729 (2006).

51.   Poonperm, R. *et al.* Replication dynamics identifies the folding principles of the inactive

X chromosome. *Nat Struct Mol Biol* **30**, 1224–1237 (2023).

52.   Hansen, R. Role of late replication timing in the silencing of X-linked genes. *Human

Molecular Genetics* **5**, 1345–1353 (1996).

53.   Gu, T. *et al.* DNMT3A and TET1 cooperate to regulate promoter epigenetic landscapes

in mouse embryonic stem cells. *Genome Biol* **19**, 88 (2018).

54.   Zhang, X. *et al.* DNMT3A and TET2 compete and cooperate to repress lineage-specific

transcription factors in hematopoietic stem cells. *Nat Genet* **48**, 1014–1023 (2016).

55.   Chao, L. *et al.* Competitive binding of TET1 and DNMT3A/B cooperates the DNA

methylation pattern in human embryonic stem cells. *Biochimica et Biophysica Acta (BBA) -

Gene Regulatory Mechanisms* **1865**, 194861 (2022).

56.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381–386 (2014).

57.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat

Methods* **14**, 979–982 (2017).

58.   Weiler, P., Van Den Berge, K., Street, K. & Tiberi, S. A Guide to Trajectory Inference

and RNA Velocity. in *Single Cell Transcriptomics* (eds. Calogero, R. A. & Benes, V.) vol. 2584

269–292 (Springer US, New York, NY, 2023).

59.   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

60.   Lee, I. *et al.* Simultaneous profiling of chromatin accessibility and methylation on human

cell lines with nanopore sequencing. *Nat Methods* **17**, 1191–1199 (2020).

61.   Yue, X. *et al.* Simultaneous profiling of histone modifications and DNA methylation via

nanopore sequencing. *Nat Commun* **13**, 7939 (2022).

62.   Altemose, N. *et al.* DiMeLo-seq: a long-read, single-molecule method for mapping

protein–DNA interactions genome wide. *Nat Methods* **19**, 711–723 (2022).

63.   Zwemer, L. M. *et al.* Autosomal monoallelic expression in the mouse. *Genome Biol* **13**,

R10 (2012).

64.   Boemo, M. A. DNAscent v2: detecting replication forks in nanopore sequencing data

with deep learning. *BMC Genomics* **22**, 430 (2021).

65.   Simpson, J. T. *et al.* Detecting DNA cytosine methylation using nanopore sequencing.

Nat Methods **14**, 407–410 (2017).

66.   Karolchik, D. The UCSC Table Browser data retrieval tool. *Nucleic Acids Research* **32**,

493D – 496 (2004).

67.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

68.   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.

Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin,

DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213–1218 (2013).

69.   F. Krueger. Trim Galore.

70.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

71.   Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P. & Marth, G. T.

BamTools: a C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* **27**,

1691–1692 (2011).

72.   Gaspar, J. M. Improved peak-calling with MACS2. Preprint at

https://doi.org/10.1101/496521 (2018).

73.   Picard.

Figure 1. Nanopore long read sequencing enables detection and tracking of nascent strand DNA methylation across sub-cell-cycle timescales.

Figure 2. Nascent and mature chromatin can be distinguished by intramolecular DNA methylation patterns.

Figure 3. MPATH recapitulates post-replication CpG methylation dynamics in unlabeled DNA.

Figure 4. MPATH enables simultaneous ordering of CpG methylation and hydroxymethylation.

Figure 5. Coupling MPATH with NOMe-seq enables tracking of nucleosome occupancy across sub-cell-cycle timescales.

Figure 6. MPATH reveals allele-specific trends in pseudotime distribution associated with X chromosome activity.

Figure 1. Capturing the dynamics of CpG methylation via long read sequencing.

Figure 2. Separation of nascent vs mature chromatin via individual read-level metrics.

Figure 3. Enrichment analyses of labeled temporally stagnant reads.

a 0h Chase

high PC2 | low PC2
1_Active_Promoter
2_Weak_Promoter
3_Poised_Promoter
9_Txn_Transition
4_Strong_Enhancer
12_Polycomb_repressed
6_Weak_Enhancer
8_Insulator
11_Weak_Txn
7_Weak_Enhancer
5_Strong_Enhancer
14_Repetitive-CNV
15_Repetitive-CNV
10_Txn_Elongation
13_Heterochrom-lo

-1   0   1

b 16h Chase

high PC2 | low PC2
3_Poised_Promoter
1_Active_Promoter
2_Weak_Promoter
12_Polycomb_repressed
6_Weak_Enhancer
4_Strong_Enhancer
5_Strong_Enhancer
7_Weak_Enhancer
11_Weak_Txn
8_Insulator
10_Txn_Elongation
9_Txn_Transition
14_Repetitive-CNV
13_Heterochrom-lo
15_Repetitive-CNV

-0.6   0   0.6

Figure 4. Temporal profiling of nucleosome occupancy using MPATH and publicly available NOMe-seq data.

Main Figures

Figure 1. ONT long-read sequencing enables detection and tracking of nascent strand DNA methylation across sub-cell-cycle timescales. a) Illustrative schematic of chromatin maturation across S-phase. CpG sites are represented as circles (black: methylated, white: unmethylated). Different temporal dynamics are shown for transcription factors in red and orange. Purple nucleosomes are shown to exhibit shifts in positioning throughout time. **b)** Illustrative schematic of methylation pseudotime analysis. Individual long reads can be temporally ordered according to their intramolecular CpG methylation patterns. **c)** Scatter plots comparing read-level methylation fraction for individual long reads to corresponding bulk reference methylation fraction derived from WGBS between control and nocodazole-treated HUES64 hESCs. **d)** Schematic workflow of BrdU pulse-chase experiment, followed by ONT long-read sequencing and detection of modified bases. **e)** Hexbin plots depicting methylation ratio and average BrdU content across individual long reads generated from pulse optimization experiments. Red dashed line corresponds to an average BrdU threshold derived from a mock-treated negative control. Reads with average BrdU content above this threshold are considered BrdU-positive. **f)** Bar graph depicting bulk methylation percentage of BrdU-positive reads from BrdU-labeled experimental conditions, as calculated by total methylated CpGs divided by total CpGs. Gray diamond represents the bulk methylation percentage for reads generated by WGBS that capture the corresponding CpGs in BrdU-positive long reads from a given condition. Black diamond represents the bulk methylation percentage for reads generated by WGBS that capture the corresponding CpGs in all long reads from a given condition.

Figure 2. Nascent and mature chromatin can be distinguished by intramolecular CpG methylation patterns. a) Bar graph depicting bulk methylation percentage of BrdU-positive reads from BrdU-labeled experimental conditions, as calculated by total methylated CpGs divided by total CpGs. Gray diamond represents the bulk methylation percentage for reads generated by WGBS that capture the corresponding CpGs in BrdU-positive long reads from a given condition. Black diamond represents the bulk methylation percentage for reads generated by WGBS that capture the corresponding CpGs in all long reads from a given condition. **b)** Distributions of CpG methylation across individual BrdU-containing reads in the 0h and 16h chase conditions. Reads are not shown aligned to the genome. **c)** Illustrative schematic depicting the differences in how CpG methylation ratio and uniformity are calculated. CpG sites within reads are represented as circles (black: methylated, white: unmethylated). Matrices represent CpG pairs that are correlated

in methylation status (black tiles) or uncorrelated in methylation status (white tiles). **d)**
Distributions of methylation ratio for reads derived from the 0h (n = 848) and 16h (n = 975) chase
timepoints. **e)** Distributions of 1-edit distance for reads derived from the 0h (n = 848) and 16h (n
= 975) chase timepoints. **f)** Distributions of nearest-neighbor Pearson correlation R for reads
derived from the 0h (n = 836) and 16h (n = 822) chase timepoints. **g)** Distributions of uniformity
for reads derived from the 0h (n = 848) and 16h (n = 975) chase timepoints. For **d-g**, light blue
density plot corresponds to 0h chase and dark blue density plot corresponds to 16h chase. **h)** Bar
plot depicting the squared Hellinger distance between 0h and 16h read distributions
corresponding to each given metric. **i)** PCA based on intramolecular CpG methylation patterns
across individual reads (dots). Density plots across PC1 and PC2 are also shown. **j)** Total amount
of variance that can be explained by each principal component. **k)** Heatmap representing the
coefficients associated with each metric in a given principal component's corresponding linear
combination.

Figure 3. Methylation pseudotime analysis through read-level heterogeneity (MPATH)
captures the temporal dynamics of CpG remethylation without DNA labeling. a) PCA plot
displaying 0h and 16h chase clusters, as well as unlabeled reads (n = 10,000) to which the same
linear combinations from the original PCA were applied. Dashed line separates high PC2 and low
PC2 read subpopulations. **b)** Heatmap depicting enrichment of high PC2 and low PC2 populations
for certain genomic regions annotated by chromHMM. **c)** PCA plot annotated to show the
optimized inferred nascent and mature read clusters. **d)** Scatter plot showing the correlation in
methylation difference percentage between inferred versus labeled nascent and mature read
populations. Dashed lines represent linear regression models applied to the corresponding group.
Temporally dynamic tiles were generated from Repli-BS data. The y-axis value corresponding to
each temporally dynamic tile was obtained by calculating the bulk CpG methylation percentage
of inferred nascent and mature reads that capture the CpGs in a given temporally dynamic tile.
For the randomly sampled condition, reads that capture the CpGs in a given temporally dynamic
tile were randomly sampled, regardless of inferred post-replication maturity. **e)** Scatter plots and
associated linear regression models for methylation difference percentage between labeled
versus inferred nascent and mature read populations, as it relates to CpG density. 100 bp tiles
were created for each individual CpG in the hg19 reference genome and ranked by CpG density
across the tile. **f)** PCA scatter plot depicting the BrdU-containing reads from previous pulse
optimization experiments. Linear combinations of read-level metrics associated with the original

PC1 and PC2 were applied to the read populations. Density plots across PC1 and PC2 are also
shown.

Figure 4. MPATH enables simultaneous profiling of CpG methylation and
hydroxymethylation across replication-associated timescales. a) Loess smooth curves of
CpG methylation ratio and CpG hydroxymethylation ratio, which were calculated for 10,000
randomly sampled 1 kb single molecules using 2 state and 3 state methylation callers. PC1 values
for each 1 kb single molecule corresponds to the PC1 value of the read from which the 1 kb single
molecule was derived. **b)** Loess smooth curves of CpG methylation uniformity and CpG
hydroxymethylation uniformity, which were calculated for 10,000 randomly sampled 1 kb single
molecules using 2 state and 3 state methylation callers. PC1 values for each 1 kb single molecule
corresponds to the PC1 value of the read from which the 1 kb single molecule was derived. **c)**
Heatmaps displaying the CpG methylation ratio and hydroxymethylation ratio for individual CpGs
(n = 36,234) across 10 pseudotime bins. Each row represents an individual CpG that is captured
at least 10x across pseudotime. CpGs were ordered according to their total hydroxymethylation
level across pseudotime and positions are retained between both heatmaps. Red, teal, and beige
bars present 3 distinct clusters of CpGs that exhibit high hydroxymethylation, intermediate
hydroxymethylation, and no hydroxymethylation across pseudotime, respectively. **d)** Proportion
of CpGs that are methylated, hydroxymethylated, and unmethylated in each cluster of the
heatmap. **e)** Bar graph depicting enrichment of the highly hydroxymethylated CpG cluster for
cDKO-DMRs and certain genomic regions annotated by chromHMM (n = 8,718 CpGs). **f)**. Bar
graph depicting enrichment of the intermediately hydroxymethylated CpG cluster for cDKO-DMRs
and certain genomic regions annotated by chromHMM (n = 8,718 CpGs). **g-h)** Loess smooth
curves of hydroxymethylation ratio of CpGs from the **g)** highly hydroxymethylated CpG cluster (n
= 8,718) and **h)** intermediately hydroxymethylated CpG cluster (n = 8,718) across each of the 10
pseudotime bins depicted in panel **c**. **i-j)** Bar plots displaying the hydroxymethylation ratio of CpGs
that are captured in pseudotime bins 3, 6, and 9 from the **i)** highly hydroxymethylated CpG cluster
(n = 2,120) and **j)** intermediately hydroxymethylated CpG cluster (n = 1,577). Error bars represent
standard error and p values are derived from a two-sided Mann-Whitney U test.

1009

Figure 5. Coupling MPATH with NOMe-seq enables temporal ordering of chromatin
accessibility. a) Scatter plots representing the original long reads from GSE183760 (left) and
individual 10 kb fragments derived from those long reads (right). Each dot represents an individual
read of 10 kb fragment. Reads and fragments were transformed using the linear combinations

comprising the original PC1 and PC2. PC1 was divided into 4 bins T1-T4, ordered from most
nascent to most mature. **b)** Density plots showing the distributions of inaccessible run lengths for
10 kb fragments that fall into each of the PC1 bins (T1-T4). T1, n = 7,866. T2, n = 5,550. T3, n =
5,810. T4, n = 2,402. **c)** Density plots showing the distributions of insert size associated with each
chase timepoint in Repli-ATAC. 0hr, n = 10,863,954. 1hr, n = 22,901,076. 4hr, n = 37,219,193.
16hr, n = 40,564,395. **d)** Scatter plot and linear regression model depicting the correlation
between an average of PC1 values from individual 10 kb fragments comprising a long read versus
a weighted sum of PC1 bins for individual 10 kb fragments comprising a long read. **e)** Heatmap
depicting the proportion of 10 kb fragments comprising an individual long read that fall into each
PC1 bin. Each row represents an individual long read. Reads are ordered by their weighted sum
of PC1 bins. **f)** Scatter plot and Loess smooth curve depicting the proportion of inaccessible run
lengths from individual 10 kb fragments that are 0-200 bp long, plotted across the PC1 weighted
sum of the 10 kb fragment. Only 10 kb fragments from Cluster 5 are shown. **g)** Scatter plot and
linear regression model depicting the correlation between inaccessible run length and CpG
methylation uniformity across 10 kb fragments from Cluster 5. R^2 was calculated from mean CpG
methylation uniformity values. Error bars represent standard error. p value was derived from an
F-test.

1031

Figure 6. MPATH reveals allele-specific trends in pseudotime distribution associated with
genomic imprinting and X chromosome inactivation. a) Browser window depicting phased
long reads from Abl.1 mouse B cells across chrX:103,468,194-103,497,066. Colored according
to methylation status of CpG sites (blue: unmethylated, red: methylated). Haplotype 1 is derived
from the inactive X, as shown through lack of CpG methylation at the *Xist* promoter (shaded blue
region). **b)** Bar plot depicting the difference in bulk CpG methylation between reads derived from
haplotype 1 vs 2 across 51 20-kb windows. Bars that are blue represent windows at which the
difference in bulk CpG methylation between reads derived from haplotype 1 vs 2 is ≤ 15%. **c)**
Density plot depicting the distribution of PC1 values assigned to reads derived from haplotype 1
(n = 637) vs haplotype 2 (n = 600) within filtered windows across the X chromosome region (chrX:
103,530,980-104,531,001) at which the difference in bulk CpG methylation between reads
derived from haplotype 1 vs 2 is ≤ 15%. **d)** Density plot depicting the distribution of PC1 values
assigned to downsampled reads derived from haplotype 1 (n = 637) vs haplotype 2 (n = 600)
within a region of the autosome chr10.

1046

1047 # Supplemental Figures

1048

Figure 1. Capturing the temporal dynamics of CpG methylation via long read sequencing.
a) Representative browser track depicting bulk CpG methylation percentage across BrdU[+] reads
(generated via Repli-BS) that are derived from the 0h chase (nascent) and 16h chase (mature).
b-c) Bioinformatics analysis pipelines for identifying BrdU-containing reads and calling CpG
methylation for **b)** R9.4.1 and **c)** R10.4.1 flow cells from Oxford Nanopore Technologies. **d)** Bar
plot depicting % live cells following treatment of HUES64 hESCs with different BrdU
concentrations and chase times. Data is shown for 2 biological replicates and was measured
through LIVE/DEAD Stain. **e)** Bar plot displaying percentage of total bases sequenced in BrdU-
containing reads vs all reads that overlap with annotated genomic regions. Data is shown for both
chase timepoints across 2 biological replicates.

Figure 2. Separation of nascent vs mature chromatin via individual read-level metrics.
a-h) Distributions of **a)** nearest-neighbor Pearson correlation p values, **b)** simple matching
coefficient (SMC) across total reads, **c)** nearest-neighbor SMC, **d)** uniformity of CpGs <100bp
apart, **e)** uniformity of CpGs 100bp-1kb apart, **f)** uniformity of CpGs 1kb-5kb apart, **g)** uniformity
of CpGs 5kb-10kb apart, and **h)** nearest-neighbor uniformity for reads derived from the 0h (n =
848 reads) and 16h (n = 975 reads) chase timepoints. **i-l)** Stacked bar plots depicting the
proportion of BrdU-labeled reads within each **i)** methylation ratio, **j)** 1-edit distance, **k)** uniformity,
or **l)** PC1 bin that are derived from the 0h chase vs 16h chase. For each read-level metric, all
possible values were ordered least to greatest and divided into 10 bins of equal intervals.

Figure 3. Enrichment analyses of labeled and unlabeled temporally stagnant reads. a-b)
Heatmap depicting enrichment of high PC2 and low PC2 labeled read populations from the **a)** 0h
chase time point and **b)** 16h chase time point for certain genomic regions annotated by
chromHMM.

**Figure 4. Temporal profiling of nucleosome occupancy using MPATH and publicly
available NOMe-seq data. a)** Schematic illustrating the Repli-ATAC sequencing workflow in
HUES64 hESCs. **b)** Density plots showing the distributions of sub-nucleosomal inaccessible run
lengths (<100 bp) for 10 kb fragments that fall into each of the PC1 bins (T1-T4). **c)** Heatmap
depicting the coverage of GpCs across methylation pseudotime with single-molecule resolution.
Only data from the 73kb CRISPR-Cas9 targeted region were used. Only GpCs captured at least
30x are shown (n = 557). Each row represents an individual GpC and each column represents an

individual read. Reads are ordered left to right by PC1 weighted sum value (180 reads total) and

GpCs are linearly ordered top to bottom. GpCs are colored blue (methylated) or green

(unmethylated) if captured by a read and gray if uncaptured by a read. GpCs are clustered based

off of their coverage. **d)** Scatter plot depicting the coverage (percentage of reads that are NA for

a given GpC) of individual GpCs across the genomic coordinates captured by the 73kb CRISPR-

Cas9 targeted region. **e)** Box plot showing the distributions of coverage (percentage of reads that

are NA for a given Gaps) for GpCs in Clusters 1 (n = 105), 2 (n = 124), 3 (n = 126), 4 (n = 49),

and 5 (n = 153). **f-j)** Scatter plots and Loess smooth curves depicting the proportion of

inaccessible run lengths from individual 10 kb fragments that are **f)** 0-250bp long, **g)** 0-280bp

long, **h)** 100-200bp long, **i)** 100-250bp long, and **j)** 100-280bp long, plotted across the PC1

weighted sum of the 10 kb fragment. Only 10 kb fragments from Cluster 5 are shown.

1093