Harnessing Microbial Tools: *Escherichia coli* as a Vehicle for

Neuropeptide Functional Analysis in *Caenorhabditis elegans*

Elizabeth M. DiLoreto[1], Shruti Shastry[1], Emily J. Leptich[2], Douglas K. Reilly[1,3], Rachel

N. Arey[2], Jagan Srinivasan[1,4]*

[1] Department of Biology and Biotechnology, Worcester Polytechnic Institute, 100

Institute Drive, Worcester, MA, USA, 01609

[2] Department of Molecular and Cellular Biology, Baylor College of Medicine, 6450 E

Cullen St, Houston, TX, USA 77030

[3] Current Position: Department of Biology, Tufts University, Medford, MA, USA

[4] Neuroscience Program, Worcester Polytechnic Institute, Worcester, MA, USA

* Corresponding author email: jsrinivasan@wpi.edu

<u>Running head (50 characters)</u>

Testing neuropeptide function by feeding *E. coli t*o *C. elegans*

<u>Keywords</u>

Caenorhabditis elegans, neuropeptides, behavioral rescue, neuropeptide receptor

Abstract

Animals respond to changes in their environment and internal states via neuromodulation. Neuropeptides modulate neural circuits with flexibility because one gene can produce either multiple copies of the same neuropeptide or different neuropeptides. However, with this architectural complexity, the function of discrete and active neuropeptides is muddled. Here, we design a genetic tool that facilitates functional analysis of individual peptides. We engineered *Escherichia coli* bacteria to express active peptides, fed loss-of-function *Caenorhabditis elegans*, and rescued the activity of genes with varying lengths and functions: *pdf-1, flp-3, ins-6*, and *ins-22*. Some peptides were functionally redundant, while others exhibit unique and previously uncharacterized functions. We postulate our rescue-by-feeding approach can elucidate the functional landscape of neuropeptides, identifying the circuits and complex peptidergic pathways that regulate different behavioral and physiological processes.

Article summary

Studying individual neuropeptides opens new avenues for exploring neuromodulation at a finer resolution. The researchers developed a method to create DNA vectors that encode an endogenous peptide sequence flanked by sequences containing dibasic endopeptidase cleavage sites in *Caenorhabditis elegans*. The researchers transformed these vectors into bacteria and fed them to *C. elegans*, which restored wildtype behavior in neuropeptide loss-of-function mutants. The researchers also discovered that neuropeptides from the same gene perform distinct functions, a research area more ready to explore using the presented technology.

Graphical Abstract

Created in BioRender. DiLoreto, E. (2025) https://BioRender.com/ypbrtlk

Introduction

The modulation of neural circuits is essential for life, as appropriate responses to

environmental stimuli depend on precise neural tuning. Neuropeptides, which are

relatively short amino acid chains (less than 100 amino acids), play a crucial role in this

process by regulating neural circuits while also functioning as neurotransmitters or

neurohormones [1, 2]. They are widely used across the animal kingdom, even in

organisms that possess only secretory cells without neurons [3]. Neuropeptides can

also have conserved structure across species while serving a variety of functions. Oxytocin-like and vasopressin-like peptides are ancient neurohormones that regulate social attachment, lactation, and blood pressure in mammals [4-9]. However, in the Echinoderm sea star, *Asterias rubens*, the vasopressin-like and oxytocin-like neuropeptide ortholog, asterotocin, serves instead to relax muscles in the cardiac stomach during fictive feeding[9]. The *Caenorhabditis elegans* vasopressin ortholog, nematocin, interacts with serotonin and dopamine signaling to modulate gustatory associative memory and male mating behaviors [4, 10].

58 Neuropeptides act in conjunction with neurotransmitters, typically binding to G-protein coupled receptors, where they modulate synaptic activity more slowly and over a longer duration than neurotransmitters alone [2, 11]. Multiple modulators enable neurons to serve distinct functions within specific neural circuits [12]. Beyond the nervous system, neuropeptides also regulate immune and endocrine functions [13, 14]. In humans, there are varying reports of 90 or 131 neuropeptide precursor genes, yet all studies agree that there are likely more neuropeptides in the 900 other peptide genes [15-18]. Post-transcriptional modifications can give rise to an even greater number of functional neuropeptides, numbering in the thousands [15-19]. This complexity is even seen in a "simpler" model organism, the roundworm *Caenorhabditis elegans*. *C. elegans* is a microscopic nematode that displays robust behaviors driven by just over 300 neurons [20, 21]. The *C. elegans* genome encodes three classes of neuropeptides-FMRFamide-like peptides (FLP), insulin-like peptides (INS), and non-FLP/insulin neuropeptide-like peptides (NLP) [22-24]- and encodes over 300 individual neuropeptides through 160 identified genes that modulate the functional connectome [1, 19, 25, 26]. The complexity of the neuropeptide genome, combined with extra-synaptic neuropeptide signaling, makes elucidating the role of individual peptides difficult, as canonical studies often rely on null mutations and transgenic rescues [27-30]. In particular, full gene rescue restores complete preproproteins and makes discrimination of discrete peptide function difficult [1, 31]. Previous approaches to study individual endogenous neuropeptide function involved using synthetic peptides applied by soaking the nervous system of the worm [32] or injecting the peptide into the worm [33]. The use of synthetic peptides can be costly as pure peptide must be synthesized, often

outsourced, for every neuropeptide experiment. Interestingly, feeding of exogenous
peptides via *E. coli* has proven successful in manipulating *C. elegans* biological function
[34]. However, this type of *E. coli* feeding of peptides has not been accomplished with
endogenous neuropeptides.

Here, we report an approach that enables the manipulation of neuropeptide
activity by feeding neuropeptide macromolecules produced by bacteria, allowing for the
simultaneous delivery of individual neuropeptides to large populations of animals. Our
gain-of-function approach of feeding bacteria containing the neuropeptide
macromolecules (mRNA and peptides) provides ease of application and cost-saving
measures to deliver endogenous peptides to *C. elegans.* Neuropeptide rescue by
feeding, outlined here, expands on RNA interference (RNAi) feeding paradigms, which
have been successfully used to test gene function [35, 36], wherein a plasmid encoding
the RNA of interest is driven by isopropyl-β-D-thiogalactoside (IPTG)-induction.
Whereas RNAi employs paired T7 promoters facing one another to produce double-
stranded RNA, our protocol utilizes a single T7 promoter to generate mRNA encoding
the peptide of interest. Once the neuropeptide vector is assembled and transformed into
E. coli, delivery of the peptide simply requires feeding the worms. This method allows
for temporal control of peptide delivery, animals can be fed at any life stage, and the
number of animals treated is readily flexible.

The paradigm rescues individual, processed neuropeptides, leveraging the
genetic amenability of the *C. elegans* food source, *Escherichia coli*, to circumvent the
need for transgenic development and enable high-throughput rescue and elucidation of
individual neuropeptide function [37]. We demonstrate the application of this paradigm
by rescuing behaviors driven by neuropeptides synthesized from *pdf-1*, *ins-6*, and *ins-*
22, and expand upon the previously established behaviors of *flp-3* [38].

Materials and Methods

C. elegans Strains

C. elegans strains were maintained on nematode-growth media (NGM: 51 mM NaCl, 10
mM peptone, 51 mM agar, 1 mM $CaCl_2$, 1 mM $MgSO_4$, 25 mM KPO_4 (pH 6.0), 13 μM

Cholesterol) plates seeded with *E. coli* OP50 at 15 – 20°C, according to standard

procedures [39]. Strains used are listed in **Table S1**. The strains selected from previous

publications have validated the associated neuropeptide behavior. For *pdf-1* associated

assays, the strain UR954 (*pdf-1(tm1996)*III; *him-5(e1490)*V) creates a *pdf-1* loss of

function (lof) behavior by a 588 base pair gene deletion resulting in a splice donor

variant. The control strain for *pdf-1* behaviors was CB4088 *him-5(e1490)*V, which

creates a higher incidence of males by a point mutation (G to A) resulting in a splice

acceptor variant. *flp-3 lof* behavior was tested by JSR99 (*flp-3(pk361)*X; *him-*

*8(e1489)*IV) whose *flp-3* allele results in a 439 base pair deletion resulting in a

suspected frame shift. The *him-8* allele CB4088 (*him-8(e1489)*IV) selected results in a

higher incidence of males by a point mutation (G to A) resulting in a missense mutation.

The Bristol wildtype isolate (N2) was used as the control for the INS behavior tests

since they were based in hermaphrodite behavior. *ins-6 lof* behavior was tested by

IV302 (*ins-6(tm2416)*II; kyEx2595[*str-2*p::GCaMP2.2b + *unc-122*p::GFP]), where the

tm2416 allele results in a 316 base pair deletion that results in a splice donor variant.

ins-22 lof behavior was tested by RB2594 (*ins-22(ok3616)*III) which creates a ~700

base pair gene deletion resulting in a putative frame shift.

JSR188 was generated by mating UR954 and UR930 animals. UR930 (*pdfr-*

*1(ok3425)*III; *him-5(e1490)*V) results in *pdfr-1* loss of function by a 605 base pair

deletion resulting in a splice acceptor variant. The cross was monitored using validated

primers   from   Wormbase.org;   *pdf-1(tm1996)*   external   left

(CCGTGGAGGTACCAGGTTAA),   *pdf-1(tm1996)*   external   right

(TCAAGGTCTCGCCTCTAGGA),   *pdfr-1(ok3425)*   external   left

(CGTGGAATCATCGCTACCTT),   and   *pdfr-1(ok3425)*   external   right

(TTTATGCAGGCTTATTGCCC). The final generated strain was confirmed by

sequencing   using   the   primers   *pdf-1(tm1996)*   internal   left

(CCATTCCGATGAGTCCGTTG)   and   *pdfr-1(ok3425)*   internal   left

(TTTACTCCTTGACGGGAACG). To test the FLP-3 receptor FRPR-16 associated

behavior, JSR152 (*flp-3(pk361)*X; *frpr-16(gk5305*[*lox*p + *myo-2*p::GFP::*unc-54 3' UTR* +

*rps-27*p::neoR::*unc-54 3' UTR* + *lox*p]II)) was used, creating a *frpr-16 lof* mutation by

deleting 1685 base pairs, resulting in a splice acceptor variant. JSR166 was generated

by mating IV302 and HC196 animals. HC196 (*sid-1(qt9)*V) results in *sid-1* loss of

function single base pair mutation C to T resulting in the gain of a stop codon. The cross

was monitored using primers external left (TCCTTCGCCTGGAATTAGAT), external

right (AGCAAAATCCGTACATTCGC) and sequenced with an internal left primer

(AAACTTCCGACGGTTTATCC).

Neuropeptide Plasmid Design and Feeding

Peptide Plasmid Design and Generation

DNA sequences encoding individual peptides were identified via www.wormbase.org.

The endogenous processing patterns of the peptides selected are mapped in (**Figure**

S1). Sequences were flanked with the endogenous cleavage sites for the EGL-3

processing enzyme, which cleaves dibasic resides. Sequences encoding MRFGKR and

KRK-STOP codons were therefore placed prior to, and following the peptide codon

sequences, respectively. This construct lacks the signal peptide sequence found in

endogenous neuropeptides. Finally, Gateway Cloning sites attB1 and attB2 sites were

attached to the ends of the sequences. These final sequences (comprised of

attB1::MRFGKR::peptide::KRK-STOP::attB2) were ordered from Integrated DNA

Technologies (IDT) using their DNA Oligo and Ultramer DNA Oligo services, depending

on the size of the oligo ordered. Both forward and reverse sequences were ordered

(**Table S2**).

Lyophilized oligos were prepared following IDT Annealing Oligonucleotides

Protocol   (https://www.idtdna.com/pages/education/decoded/article/annealing-

oligonucleotides). In short, they were resuspended in Duplex Buffer (100 mM Potassium

Acetate; 30 mM HEPES, pH 7.5; available from IDT), preheated to 94 °C to a final

concentration of 40 µM. Complementary oligo sequences were then mixed in equimolar

ratios and placed in a thermocycler at 94 °C for 2 minutes prior to a stepwise cooling to

room temperature.

Annealed oligos were used to perform a BP reaction with pDONR p1-2 donor

vector to generate pENTRY clones (Gateway Cloning, ThermoFisher). Entry clones

were then recombined with pDEST-527 (a gift from Dominic Esposito (Addgene plasmid

11518)) in LR reactions generating expression clones. The control (SCRAMBLE) was

generated in an identical manner, with the sequence between the cleavage sites being

amplified from pL4440 (provided by Victor Ambros, University of Massachusetts Medical

School, MA) to produce a peptide that lacks sequence homology to any known peptide

(**Table S2**). Purified neuropeptide expression vectors were stored at -20 °C or -80 °C for

long-term storage.

For the *ins-22* experiments, the DNA coding sequence for the peptide was

identified as described above, without the Gateway Cloning sites. These sequences

were codon-optimized and cloned in pET-21a(+) cloning vector via XhoI/XhoI strategy

by GenScript (www.genscript.com). The SCRAMBLE control sequence and plasmid

were generated using the same cloning strategy by GenScript. Plasmids arrived

lyophilized and were re-suspended in nuclease-free H_2O to a concentration of 100

ng/µL and stored at -20 °C.

The plasmid maps displayed in **Figure S2a,b** were rendered by SnapGene®

software (from Dotmatics; available at snapgene.com).

Peptide Feeding Bacterial Preparation

The neuropeptide expression vectors were transformed into competent DH5α cells

(NEB® [New England Biolabs] 5-alpha Competent *E. coli* (High Efficiency) Cat#

C2987I). Peptide-expressing cultures on LB agar plates (10 g/L NaCl, 10 g/L Tryptone,

5 g/L Yeast Extract, 5 g/L Agar) containing 100 µg/mL ampicillin (AMP), were stored at 4

°C for up to 2 months. From this plate, single colonies were selected for growth in 5 mL

of LB media (10 g/L NaCl, 10 g/L Tryptone, 5 g/L Yeast Extract) containing ampicillin at

37 °C for 16 hours. The optical density of the grown cultures (OD_{600}) was adjusted to

1.0. 75 µL of 1.0 OD_{600} peptide-expressing culture was plated onto 60 mm NGM agar

plates containing 100 µg/mL AMP and 1 mM isopropyl-β-D-thiogalactoside (IPTG) at

room temperature for at least 8 hours prior to use, for no longer than 1 week.

Competent iOP50 (*E. coli*, rnc14::(delta)Tn10, lacz(gamma)A::T7pol camFRT)

obtained from *Caenorhabditis* Genetics Center was created using previously described

methods [40]. Briefly, iOP50 stocks were streaked out onto plates containing 50 µg/mL

tetracycline and LB and grown at 37°C overnight. Then, single colonies of iOP50 were isolated and used to inoculate 2.0 mL LB containing 50 µg/mL tetracycline and placed in a 37°C shaker overnight. 0.5 mL of each culture was used to inoculate 3.0 mL of LB medium without antibiotics and was grown for 2 hours at 37°C in a shaker. Each culture was then aliquoted in 0.75 mL increments into four 1.5 mL tubes and cooled on ice for 10 minutes. Next, the cells were harvested by centrifugation at 6,000 rpm for 5 minutes at 4°C. The resulting supernatant was discarded, and the cells were placed back on ice and resuspended in 1.0 mL 100 mM ice-cold $CaCl_2$ solution. Cells were kept on ice for 20 minutes and then harvested by centrifugation as in the previous step. After discarding the supernatant, the cells were resuspended in 150 µL of ice-cold $CaCl_2$ solution and placed on ice for immediate use in transformation.

Expression clones were transformed into competent iOP50 cultures using the New England Biolabs protocol (C2987H/C2987I). Single colonies of transformed cells were isolated and cultured overnight in 3.0 mL of LB medium containing 50 µg/mL tetracycline (TET) and 17 µg/mL chloramphenicol (CAM) at 37°C in a shaker. Cultures were frozen down as glycerol stocks and stored at -80°C. Successful transformation was verified by isolating plasmids from cultures using a Zyppy Plasmid Miniprep Kit (Zymo Research #D4019) and sending samples for whole-plasmid sequencing.

For plating the bacteria containing *PDF-1* and *FLP-3*, LB medium containing 50 µg/mL ampicillin (AMP), 50 µg/mL tetracycline (TET), and 17 µg/mL chloramphenicol (CAM) was inoculated with iOP50 expressing either the peptide of interest or the control peptide and cultured for 16-20 hours at 37°C in a shaker. 75 µL of bacteria expressing the peptide was plated onto 60 mm NGM plates containing 50 µg/mL AMP, 50 µg/mL TET, and 17 µg/mL CAM, and 1 mM IPTG. For all peptides, seeded plates were left to dry at least 8 hours at room temperature before plating worms and were used no longer than 1 week after plating.

For *In-22,* LB containing 50 µg/mL tetracycline and 50 µg/mL carbenicillin was inoculated with iOP50 expressing either the peptide of interest or the control peptide and cultured for 18-20 hours at 37°C in a shaker. 100 mm NGM plates containing 50 µg/mL tetracycline, 50 µg/mL carbenicillin, and 1 mM IPTG were plated with 1.0 mL of

the corresponding culture. Seeded plates were left to dry for at least 24 hours at room
temperature and were never stored for longer than 72 hours before use. Approximately
one hour before transferring worms to peptide or control plates (SCRAMBLE), 200 µL of
0.1 M IPTG was added to each plate and left to dry.

Peptide Feeding of Bacterial Cells or Supernatant Alone

To test if the bacterial cells expressing the peptide construct or the supernatant that the
cells were grown in was sufficient to restore neuropeptide-associated behavior, the cells
expressing the peptide were separated from the media in which they grew and might
have expressed macromolecules into. 5 mL of LB medium containing 100 µg/mL
ampicillin (AMP) with DH5α cells expressing the control SCRAMBLE construct or FLP-
3-9 was grown for 16 hours at 37 °C with shaking. At the same time, 5 mL of LB
containing OP50 *E. coli that did* not express the peptide was grown for 16 hours at 37
°C with shaking. Afterwards, the cultures were centrifuged at 3900 RPM (rotations per
minute) or 11,000 x G for 5 minutes at 4 °C to separate the cells from the media. The
supernatant from the DH5α *E. coli* cells expressing the peptide was mixed with the
OP50 cells not expressing the peptide, and the DH5α *E. coli* cells expressing the
peptide were mixed with the OP50 cell supernatant. The cultures were adjusted to have
an OD_{600} = 1.0. 75 µL of each culture (SCRAMBLE DH5α cells, SCRAMBLE DH5α
supernatant, FLP-3-9 DH5α cells, and FLP-3-9 DH5α supernatant) was plated onto 60
mm NGM agar plates containing 100 µg/mL AMP and 1 mM IPTG. The plates were
stored at room temperature, 20-22 °C, for at least 8 hours prior to use, for no longer
than 1 week.

Peptide Feeding

All animals were maintained on bacterial lawns of OP50 *E. coli*, at 20 °C, on NGM agar
plates until the start of experiments. Mutant or control animals were transferred onto
lawns of DH5α or iOP50 *E. coli*. Animals were reared on the peptide-expressing lawns
for at least 48 hours at 20 °C before assaying for rescue of mutant phenotype at the
appropriate developmental stage. See each section of the behavioral assay methods for
the exact peptide feeding parameters.

<u>Behavioral Assays</u>

Excursion Assay (pdf-1)

The mate searching behavior of *pdf-1* mutants in a *him-5* background was quantified in
a food leaving assay. *him-5* animals were used as wildtype control to ensure the
presence of males. *him-5* and *pdf-1* animals were fed bacteria expressing no peptide
(OP50), control peptide (SCRAMBLE), PDF-1A peptide, PDF-1B peptide, or 1:1 PDF-1A
and PDF-1B peptide (by mixing bacterial cultures expressing these peptides). Four to
six larval L4 hermaphrodites were passed onto peptide-containing plates. These worms
were grown at 20 °C for approximately 4 days, until the male progeny of the initial L4
worms had developed into young adults. The day prior to the assay, young adult male
pdf-1 or *him-5* worms were singled onto 60 mm plates containing lawns of either OP50
or appropriate peptide cultures.

To determine the mate searching effects of *pdf-1*, we performed a food-leaving
assay as described previously [41-43], with modifications. In this assay, individuals were
placed on a small food spot, and track patterns were scored at the indicated times.
Assay plates were prepared one day prior to assaying on 100 mm plates with 10 mL of
Leaving Assay media (25 mM KPO_4 (pH 6.0), 1 mM $MgSO_4$, 1 mM $CaCl_2$, 50 mM NaCl,
2.5 g/L Peptone, 17 g/L Agar). On center of plate, 7 µL of OP50 was added. In assays
using iOP50 assay plates, 7 µL of iOP50 containing the appropriate peptide (ex.
SCRAMBLE iOP50 assay plates made for animals fed SCRAMBLE iOP50) was added.
Plates were lightly covered and stored at room temperature overnight (~20^0C, < 40%
humidity).

On the day of the assay, a single male was plated on the assay plate and placed
in a dark, 20°C incubator. At 2, 6, and 24 hours after the worms are plated, plates were
scored for male worm tracks. Tracks were scored based on their distance from the
center of the plate [41]. "Never left food" indicates the absence of tracks outside the
food spot. "Minor excursion" indicates that the tracks never extended beyond 10 mm
from the food. "Major excursion" indicates the presence of tracks past the 10 mm
boundary. We quantified the food leaving behavior at three different time points (2, 6,
and 24 hours) [42].

We additionally quantified the mate searching defect of *pdf-1* as a Probability of
Leaving per hour (P_L/hr) [43]. In this method, we reduce the analysis of the leaving
behavior to a single value, by scoring the worms leaving the food lawn as Leavers
(traveling beyond 35 mm from the food source) or Non-Leavers (did not travel further
than 35 mm from center of food). Worm in the > 35 mm from the center of food bin were
scored as "Leavers" [43]. The probability of leaving per hour (Probability of Leaving (P_L))
was calculated using an R script developed in [42], first cited in [43], with assistance by
(Personal Communications to Barrios, 2020). P_L was calculated by the "hazard obtained
by fitting an exponential parametric survival model to the censored data using maximum
likelihood" [43].

Avoidance Assay (flp-3)

Avoidance drop assays were performed as described previously [38, 44]. This was
performed on *him-8* or *flp-3;him-8* loss of function male animals. Fifty to sixty L4 male
worms were picked by sex to isolate the males and stored at 20 °C for at least
5 hours to overnight to be assayed as young adults. One to four hours prior to the
assay, the lids of unseeded plates were tilted to allow any excess moisture to evaporate
off the plates. At the time of the assay, 10 or more males were transferred onto each of
the dried, unseeded NGM plates. A drop of either water or 1 µM ascr#8 (a nonvolatile
C. elegans mating pheromone) was placed on the tail of forward moving animals, and
their response was scored as either an avoidance response, or no response. The
avoidance index was calculated as:

Avoidance Index $= \frac{(\# \, of \, worms \, avoiding \, drop)}{total \, \# \, of \, worms \, tested}$

Chemotaxis Assay (ins-6)

A salt chemotaxis assay was performed as previously described in [45],[46], to test the
functionality of *ins-6* after rescue-by-feeding. Wildtype (N2 Bristol) and *ins-6* loss of
function worms were fed either a control peptide (SCRAMBLE) or INS-6 peptide. This
was done throughout the animals' lives by passing 4-6 L4 larval onto an NGM plate
containing 100 µg/µL AMP and 1 mM IPTG with 75 µL of peptide OD_{600} = 1.0. These

worms were grown at 20 °C for about 4 days until the progeny of the initial L4 worms
were young adults.

Chemotaxis plates were prepared the day prior to testing, by adding 10 mL of
agar (5 mM KPO_4 (pH 6.0), 1 µM $MgSO_4$, 1 µM $CaCl_2$, 20 g/L agar, and 8 g/L Difco
Nutrient Broth) to 100 mm petri dishes. To prepare the "high salt" plates, 750 mM NaCl
was added prior to plate pouring. The high salt plate were stored at 4°C for up to one
month, while the Chemotaxis Plates were made one day prior to testing. To create the
high and low salt plugs, the back end of a Pasteur pipette was used to punch a 5 mm
plug out of each plate. These 2 plugs were placed on opposite edges of the plate. A 10
mm radius was be drawn around the location of the plugs. Plates were stored, lightly
covered, overnight at room temperature (~20°C, < 40% humidity).

On the day of the assay, young adult worms, either control worms fed OP50 *E.*
coli or peptide-fed worms, were washed off peptide plates with M9. Worms were
allowed to settle by gravity for 4 minutes before removing the supernatant, washing the
worms. Worms were then washed with Chemotaxis Buffer (5 mM KPO_4 (pH 6.0), 1 mM
$MgSO_4$, 1 mM $CaCl_2$) three times.

To prepare the Chemotaxis plates for the assay, high and low salt plugs were
removed with forceps, and 1 µL of 0.5 M sodium azide (NaN_3) was added to the plage
where the plugs had rested. Approximately 30 µL of worms in Chemotaxis Buffer were
spotted onto the edge of the agar, between the location of the high and low salt
gradient. Worms were left to chemotax for 1 hour before scoring the number of worms
within the 10 mm radii of high and low salt areas using the below choice index to assess
the behavior of animals that had made a choice between the two salt location:

Choice Index $= \frac{(\# \, of \, worms \, high \, salt - \# \, of \, worms \, low \, salt)}{total \, \# \, of \, worms \, on \, both \, spots}$

Olfactory Chemotaxis Assay (ins-22)

Chemotaxis assays were performed using worms that were transferred at the L4 stage
to plates seeded with iOP50 containing a plasmid expressing either a control sequence
(SCRAMBLE) or the peptide of interest (INS-22). These experiments were performed on
wildtype (N2 Bristol) animals or *ins-22* loss of function animals. After 48 hours of

exposure, we tested the naïve chemosensory preference of Day 2 Adults using a protocol based on previously published assays [44]. Briefly, assays were performed on unseeded 100mm NGMs. On the back of each plate, two marks were made on opposite sides of the plate, approximately 5mm from the edge. 1 µL of sodium azide (Thermo Fisher) was placed on both spots and allowed to dry before adding 1 µL of 0.1% or 10% butanone (Sigma Aldrich) diluted in ethanol on one mark and ethanol on the other. Using M9 buffer, worms were washed off their plates and subsequently washed three times to eliminate any leftover bacteria that could impact baseline behavior. Then, worms were placed near the bottom center of the plate, equidistant between the two marks, and allowed to chemotax for an hour. Chemotaxis indices were calculated as follows:

$$\text{Chemotaxis Index} = \frac{(\# \, of \, worms \, butanone - \# \, of \, worms \, ethanol)}{total \, \# \, of \, worms}$$

Statistical Analysis

Power Analysis

Prior to the study, a post-hoc power analysis (ClinCalc LLC., Chicago, IL, USA) was performed using preliminary data from OP50 *E. coli* fed worms to determine the number of samples required to establish a difference between the wildtype controls and the neuropeptide loss of function mutants. This was performed using a 0.05 alpha rate.

For the excursion assay comparing *pdf-1 lof* animals in a *him-5* background compared to *him-5* males, a dichotomous power analysis was used to compare non-leavers (< 35 mm) to leavers (> 35 mm). n = 39 (where n = 1 is one male on one assay plate) was required for 100% powered experiment and 13 for power > 80%. A sample size of n = 50 samples per genotype and feeding condition was selected.

For avoidance behavior towards ascr#8 comparing wildtype control *him-8* animals to *flp-3 lof* males in a *him-8* background, n = 3 samples (n = 1 is one plate of ten worms) was appropriate to have a 100% powered test, using a continuous power test. A sample size of N = 10 was selected. Points plotted on graphs represent n = 1.

For the salt chemotaxis assay with *ins-6 lof* animals, n = 4 samples (n = 1 is one
plate of 100-200 worms) were required for power > 80% and 13 for 100% power. A
sample size of n = 10 was selected. Points plotted on graphs represent n = 1.

For chemotaxis behavior of *ins-22* animals, the behavior of SCRAMBLE fed
iOP50 animals towards 10% butanone was used to determine that 20 samples (n = 1 is
one plate of 100-200 worms) were required for 97.5% power and 11 samples were
required for power > 80%. A sample size of n > 15 was selected. Points plotted on
graphs represent n = 1.

Reproducibility

In all experiments, the investigators were blinded to genotype. Young adult animals
were tested in all behavioral assays. For the behavioral tests of *flp-3* and *ins-6*, the
assay was repeated over at least 3 days, with no more than 3 replicates in a day, until at
least 10 plates had been tested. For *ins-22*, the assays included testing 5 plates per
day, for a total of three biological replicates (n = 15). For *pdf-1,* the assay was
performed across at least two days per condition, with at least n = 50, wherein each
assay plate with one male is equal to n = 1. When possible, the results of an individual n
are displayed on the graphs.

Statistical Comparison

Statistical analysis was performed in GraphPad Prism (version 10.4.1). Prior to any
statistical comparison, the normality and frequency distribution were assessed for each
of the groups. If these tests showed normal distributions, a Student's unpaired two-
tailed t-test followed by Bonferroni's correction was applied; this was used for *ins-6* and
ins-22 results. If the data was non-parametric, as was often the case with the *flp-3* data,
a Mann-Whitney test was performed followed by Bonferroni's correction. For the *pdf-1*
tests where animals were fed different types of peptides, a One-way ANOVA with
Dunnett's post-hoc test was used to compare within genotype results. For all graphs,
the average was plotted with the standard error of the mean (SEM).

Results and Discussion

<u>Bacterial Delivery of Neuropeptide Vectors Rescues Behavior in Loss of Function</u>
<u>Mutants</u>

To demonstrate the efficacy of the neuropeptide delivery by bacterial feeding, we tested
the behavior of loss of function (*lof*) animals from each of the three neuropeptide
classes in *C. elegans*. Each of these behavioral assays was paired with the appropriate
wildtype control. For male animal behavior, *him-5* or *him-8* lines were used to produce a
higher incidence of males (him) than the wildtype population of 0.1% males [47, 48].
The choice of which *him* mutant to use depended on the genetic position of
neuropeptide gene being tested and other the position of other genes of interest used in
the initial studies establishing the associated neuropeptide behavior.

Male Excursion Behavior (pdf-1)

Wildtype males display a characteristic exploratory behavior when left on a lawn of food,
as well-fed males leave food in search of mates [42, 43]. The pigment dispersing factor
PDF-1 (also known as NLP-74) neuropeptide plays a significant role in male mate-
searching behavior, regulating neural circuits that control the choice between two
predominant male interests, finding food and finding mates [42, 49]. The *pdf-1* precursor
encodes two neuropeptides: PDF-1A (SNAELINGLIGMDLGKLSAVG-N terminus) and
PDF-1B (SNAELINGLLSMNLNKLSGAG-N terminus) [50, 51]. The males used in this
behavior were generated by a *him-5 lof* background mutation. *pdf-1 lof* males did not
leave food as readily as wildtype worms, suggesting that these worms do not display
exploratory behavior, as previously described (**Figure S2a,b**) [42]. When fed either
DH5α *E. coli* expressing individual PDF-1 peptide (or a 1:1 combination of both), *pdf-1*
lof had a higher P_L compared to SCRAMBLE fed *pdf-1 lof* males (**Figure 1a, S2c**).
However, PDF-1B and a 1:1 ratio of both peptides resulted in a significantly different P_L
from *pdf-1* males fed PDF-1A alone. While the P_L of peptide-fed males were significantly
higher than *pdf-1 lof* males fed SCRAMBLE, they were still significantly lower than
wildtype males fed SCRAMBLE (**Figure 1a**).

Pheromone Avoidance (flp-3)

C. elegans produce and release pheromones as a means of chemical communication to

signal a variety of features to other animals, such as food or mate presence [52]. One

such pheromone is ascaroside #8 (ascr#8) [53]. We have previously worked to

characterize the behavior of animals towards this hermaphrodite produced pheromone

and have found that wildtype hermaphrodites avoid ascr#8 while males are attracted to

it [38]. We have further identified that males deficient in the neuropeptide gene *flp-3* lose

their attraction towards ascr#8 and start to avoid the cue (**Figure S2d**) [38]. The males

of this behavior were created in a *him-8 lof* background. The *flp-3* neuropeptide

precursor creates 10 different neuropeptides, yet only 2 are involved in the male

behavior towards ascr#8 [38]. Both the second and the ninth peptide (FLP-3-2 and FLP-

3-9) produced by *flp-3* are involved with suppressing the basal avoidance behavior seen

in *flp-3 lof* males, yet only FLP-3-9 is necessary for restoring the attraction towards

ascr#8 [38, 54]. When we fed male *flp-3 lof* animals DH5α *E. coli* expressing FLP-3-9

(NPENDTPFGTMRFG-N terminus), we saw a significant suppression of avoidance

behavior towards ascr#8 (**Figure 1b**).

Salt Chemotaxis (ins-6)

Insulin and insulin-like peptides serve signaling functions in *Drosophila* and *C. elegans*

homologous to the human insulin-like growth factor (IGF), which regulates FOXO

activity [55]. In *C. elegans, ins-6* encodes only one processed INS-6 peptide

(VPAPGETRACGRKLISLVMAVCGDLCNPQEGKDIATECCGNQCSDDYIRSACCP-N

terminus) [56, 57], though the gene was originally postulated to encode two putative

proteins [1, 23]. The formation of a single longer peptide is common in the insulin-like

class of peptides [58]. *ins-6* functions in dauer formation [56, 58] and sensory

modulation of large fluctuations in salt concentration, with loss of *ins-6* causing

dysfunction of NaCl attraction [59]. While wildtype hermaphrodite worms were attracted

to high concentrations of salt (750 mM), there was a significant decrease in attraction in

ins-6 mutant animals (**Figure S2e**) [59]. Wildtype and *ins-6 lof* animals fed SCRAMBLE

peptide also had significantly different levels of salt chemotaxis (**Figure 1c**). Feeding of

DH5α *E. coli* expressing INS-6 peptide to *ins-6 lof* animals resulted in an in attraction to
high salt concentration (**Figure 1c**).

<u>Supernatant from Bacteria Expressing Peptide May Be Sufficient for Behavioral Rescue</u>
We tested whether the cells or the supernatant from a grown liquid culture expressing
the neuropeptide construct was sufficient for behavioral restoration (**Figure S3**). We
selected the *flp-3* associated avoidance behavior of ascr#8 to screen the different
culturing conditions. We found that while the behavior of *him-5* males fed any
combination of the initial culture of DH5α cells or supernatant expressing a peptide
construct (SCRAMBLE or FLP-3-9) did not alter their lack of avoidance towards ascr#8.
However, *flp-3;him-5 lof* males did exhibit reduced avoidance to ascr#8 when they were
fed a combination of the supernatant of DH5α cells expressing the FLP-3-9 peptide with
OP50 cells that did not express any peptide. These results suggest that the peptide
macromolecule of interest that restores the neuropeptide associated behavior is
excreted from the bacterial cells during the initial growth of the culture. This also
supports that the cells do not need to be induced with IPTG (which is only present in the
plates and not the liquid cultures) to express this macromolecule.

This lack of induction requirement might not entirely be surprising. Other researchers
using *E. coli* as a vehicle for expressing a designed construct have found it to have
leaky expression prior to induction, whether by IPTG or another induction agent [60, 61].
These studies found that while the amount of dsRNA or enzyme produced did increase
after induction, the uninduced state was sufficient for the creation of these
macromolecules. This would mean that the bacterial cells are able to create the
expression vector macromolecules regardless of the addition of IPTG.

<u>Different *E. coli* food sources lead to Neuropeptide Behavior Rescue</u>
While the previous experiments in this study have used DH5α *E. coli* expressing
peptides to shape neuropeptide related behaviors, there are cases when using different
E. coli strains may be desirable. Given the differences in nutrient composition of the
different *E. coli* strains and worms preferring more nutritious bacteria [62], expressing
peptides in HB101 or HB115 strains offers viable alternatives. To most closely compare
to other laboratory studies, the ability to use the widely used uracil deficient OP50 *E.*

coli strain as the neuropeptide delivery system is desirable [39]. Recently an IPTG
inducible strain of OP50 has been created, iOP50, which can be cultured to create
chemically competent cells [40]. All constructs presented in this section were
transformed into iOP50 before being fed to the *C. elegans*.

For testing male *pdf-1* associated leaving behavior, we first fed the animals on
iOP50 peptide plates before transferring them to a leaving assay plate that is seeded
with OP50 containing no peptide. Under these conditions, male *pdf-1 lof* animals fed
PDF-1 peptides from iOP50 left food at a higher rate than *pdf-1 lof* animals fed a
SCRAMBLE peptide from iOP50 (**Figure 2a**). However, the extent of food leaving by
males after being fed iOP50 with peptide was lower than observed when animals were
fed DH5α *E. coli* (**Figure 1a**). This may be because animals are tested over the course
of 24 hours and the level of peptide in their bodies may decrease over time. To address
this, we tested animals fed iOP50 with peptide and then transferred them to iOP50
assay plates seeded with the same peptide they were previously fed (**Figure 2b**). By
doing this, the trend of *pdf-1 lof* animals fed PDF-1 leaving food at a higher rate
continued, and the overall rates of leaving food also increased (**Figure 2b**).

flp-3 lof males exposed to ascr#8 after being fed FLP-3-9 in iOP50 lacked
avoidance to the cue while males fed the SCRAMBLE peptide continued to avoid it
(**Figure 2c**). These results matched the observed behavior of animals fed FLP-3-9 in
DH5α *E. coli* (**Figure 1b**). For *flp-3*, males are transferred directly from iOP50 peptide
plate onto an unseeded NGM assay plate for no more than 30 minutes during the
testing. Due to this, it was not necessary to reapply the iOP50 peptide during testing.

We also tested the delivery of peptide using a different neuropeptide expression
vector. To test the behavior associated with INS-22, a new backbone vector of pET-21
was procured to circumvent the use of Gateway Cloning and the pDest-527 vector
(**Figure S4a**). In this new neuropeptide expression vector, the peptide endogenous
peptide sequence flanked by the peptide cut sites was ordered as a complete plasmid in
a pET-21 vector (**Figure S2b**). This vector was used to test the behavior of INS-22
(MHTTTILICFFIFLVQVSTMDAHTDKYVRTLCGKTAIRNIANLCPPKPEMKGICSTGEYP
SITEYCSM-N terminus). Unlike INS-6, INS-22 antagonizes the DAF-2 neuropeptide

receptor [63, 64]. We discovered that animals lacking *ins-22* exhibit increased naive
chemotaxis towards a neutral concentration of 10% butanone compared to wildtype
animals (**Figure S4c**). This behavior persists in animals fed the SCRAMBLE peptide in
iOP50 *E. coli* (**Figure 2d**). When fed the INS-22 peptide, the *ins-22 lof* animals
displayed naive chemotaxis similar to wildtype animals, indicating that peptide feeding
rescued normal naive chemotaxis towards 10% butanone (**Figure 2d**).

<u>Neuropeptide Rescue by Feeding Requires the Cognate Neuropeptide Receptor</u>

To ensure that the observed behavior is resulting from the neuropeptide being fed to the
animal, activating the neuropeptide receptor, we tested the behavior of animals deficient
in the neuropeptide receptor. Pairing neuropeptides ligands to their cognate receptors is
an ongoing effort in peptidomic research. Recent efforts have uncovered many of the
individual neuropeptide pairings and the range over which these connections functions
[26, 65, 66]. For the current study, we were interested in studying the function of
receptors that have one peptide ligand. This would eliminate the class of insulin-like
peptides which bind to DAF-2 from this current study [23, 63].

PDF-1 peptides binds primarily to one neuropeptide receptor PDFR-1 [42, 67].
Though PDFR-1 also binds to peptides produced by *pdf-2* (also known as *nlp-37*), these
ligands are not involved in behavior of males leaving food to search for mates [42, 66].
pdfr-1 lof mutants in a *him-5* background do not leave food to search for mates (**Figure
S5a,b**), even when fed excess peptide (**Figure 3a, S5c**). The double mutant males of
the *pdfr-1* and *pdf-1* neuropeptide, partially left food to search for mates when fed PDF-
1A but not when fed the other peptides (**Figure 3a**). Though PDF-1A was able to
promote the male leaving behavior, we do not believe this was to a biologically
significant level since the rates of leaving food were lower than the rate than *pdf-1 lof*
males left food. This probability of leaving food of *pdf-1;pdfr-1* males was not to the rate
of *pdf-1* males leaving food when fed a SCRAMBLE peptide (**Figure 3a**). The absence
of food leaving to search for mates, even when receptor mutants are fed the
neuropeptide, suggests that there are no off-target effects of the neuropeptide impacting
other receptors.

The *flp-3* neuropeptide gene makes 10 different peptides that bind to 4 different

receptors [38, 66]. Neuropeptide receptors DMSR-1-2 and DMSR-7-1 bind to many of

the *flp* class of neuropeptides including *flp-3*, while FRPR-16-1 binds *flp-3* and *nlp-23*

peptides and NPR-10-1 binds *flp-3* and *nlp-23* peptides [66]. We have previously

identified that the receptors FRPR-16 and NPR-10 are involved in the *flp-3* male

behavior towards ascr#8 [38]. The FLP-3-9 neuropeptide binds activates the FRPR-16

receptor at lower concentrations than the NPR-10 receptor [38], so we elected this as

our neuropeptide receptor candidate for FLP-3-9. We have previously demonstrated

that without FRPR-16 males will avoid ascr#8 [38]. When we generated a double

mutant of *flp-3;fprpr-16* we observed avoidance of ascr#8 in animals fed OP50 *E. coli*

without peptide (**Figure S5a**). When fed FLP-3-9 peptide in DH5α *E. coli*, the avoidance

behavior is sustained (**Figure 3b**). These data shows that while FLP-3-9 binds to

multiple receptors, FRPR-16 loss of function is sufficient to alter male behavior towards

ascr#8.

Behavioral Impacts of Neuropeptide Overexpression Depend on the Peptide Identity

One question that arises with this feeding method is, how much neuropeptide makes its

way into the worms and what would happen if there were an overexpression of

neuropeptide? Quantifying the amount of neuropeptide produced endogenously is a

difficult open question in the peptidomics field. One indication of how much of a

particular neuropeptide is required is by looking at the neuropeptide precursor gene.

Some neuropeptides are encoded multiple times in the same gene, like *flp-6* encoding

KSAYMRFG-N terminus six times [68]. One method used to detect the presence of

neuropeptides is liquid chromatography mass spectrometry (LC-MS) [19, 69-71]. While

this method is extremely precise for detecting known and novel neuropeptides, it lacks

the ability to quantify the amount of each neuropeptide present *in vivo* (personal

communication with Sven Van Bael and Lisabet Timmermann, KU Leuven). One way to

test the effects of what is likely an overexpression of neuropeptide is by feeding the

neuropeptide-containing DH5α bacteria to wildtype animals.

We tested the appropriate wildtype controls for each previously assessed

behavior. We observed an increase in the leaving behavior of *him-5* male worms fed

PDF-1B and a decrease when fed PDF-1A in DH5α *E. coli* (**Figure 4a, S6a**). This
behavioral difference disappears when both peptides are fed in combination (**Figure**
4a). These data suggest that different PDF-1 peptides play varying roles in mating
behavior. Next, we examined the effects of this behavior using the iOP50 *E. coli*
delivery. When males were fed an excess of iOP50 peptide and subsequently placed on
an iOP50 assay plate, their rates of leaving food (**Figure 4b, S6b**) did not increase.
Wildtype animals fed PDF-1 peptides produced by iOP50 also did not show an
increased leaving rate of food when tested on an OP50-containing assay plate (**Figure**
S6c,d). In fact in both scenarios, the rates of leaving food decrease. This suggests that
there may be a precise balance of neuropeptide required for proper behavioral tuning.

When *him-8* male animals were fed FLP-3-9, their lack of ascr#8 avoidance was
not changed compared to animals fed SCRAMBLE (**Figure 4c**). Together these data
indicate that there are no obvious behavioral deficits across all neuropeptides being
delivered in excess. Wildtype males fed FLP-3-9 iOP50 *E. coli* continued their non-
avoidance of ascr#8 (**Figure 4d**).

We then tested different insulin ligands for their overexpression phenotypes.
When wildtype N2 animals was fed INS-6 in DH5α *E. coli*, there was no difference in the
choice index of high salt taxis (**Figure 4e**). These data suggest that INS-6
overexpression by microbial feeding does not impact the behavior of animals towards
high NaCl. There has been evidence that overexpression of some insulin like peptides
may impact other behaviors like dauer formation [55].

When wildtype animals were fed an excess of INS-22 iOP50 *E. coli*, their
behavior towards 10% butanone was decreased compared to animals fed SCRAMBLE
(**Figure 4f**). When exposed to a highly attractive concentration of 0.1% butanone,
wildtype animals fed INS-22 in iOP50 *E. coli* significantly reduced their attraction
towards the cue (**Figure 4g**). While differences in behavior towards varying
concentrations of butanone have been previously established [72, 73], the impact of
INS-22 on these behaviors has not been implicated in detail. Differences between the
overexpression effects of the *ins* peptides may be due to the different interactions with
the DAF-2 receptor, INS-6 is a strong agonist and INS-22 is an antagonist [63, 64].

Neuropeptide Feeding Does Not Utilize RNAi-based Mechanisms

Our method of neuropeptide study is most similar to RNA interference approaches,

leading us to hypothesize that the microbes could be delivering RNA to the *C. elegans*.

RNAi functions to knock down a gene by delivering dsRNA that interferes with the

endogenous mRNA of that gene [74-76]. RNAi was first performed in *C. elegans* [75,

76] and has since been accomplished by feeding, soaking, and injecting dsRNA [77].

The RNAi vector contains two T7 promoters going in opposite directions to create

dsRNA [74, 78], while our vector contains only 1, indicating that single stranded RNA

(ssRNA) should be produced. The RNAi vector is then contained in an *E. coli* strain that

has been modified to make more dsRNA, such as HT115 [79, 80]. RNA produced by

bacteria in the gut microbiota can be transported across the intestinal epithelium, as

identified by RNAi feeding [80]. This transport has also been seen with non-coding RNA

being transmitted from pathogenic bacteria to worms [81]. We hypothesize that unlike

RNAi, this method can be used to introduce larger neuropeptides encoded by over 30

base pairs [82].

To assess if our neuropeptide delivery system utilizes previously identified RNAi

mechanisms, we tested mutants deficient in the ability to transport RNA. We specifically

looked into *sid-1* (systemic RNA interference defective protein 1) loss of function

mutants. SID-1 selectively binds to dsRNA of varying (selecting towards longer) length

and not ssRNA or dsDNA [83-86]. We expect that our rescue by feeding approach

should not be effected by the loss of SID-1 since we predict that our method is either

relying on dsDNA, due to the selection of DH5α *E. coli* not containing T7 polymerase

[87]. We found that mutants lacking *sid-1* behave like wild-type animals when receiving

INS-6 peptide through feeding (**Figure S7a**). Additionally, we discovered that *sid-1;ins-6*

double mutants fed the SCRAMBLE peptide display the *ins-6 lof* phenotype (**Figure**

S7b). When fed INS-6, *sid-1;ins-6 lof* animals restored their NaCl attraction to in a

pattern similar to single mutant *ins-6* animals, suggesting that SID-1 is not involved in

our neuropeptide rescue by feeding method.

Conclusions

We present a novel method of neuropeptide rescue that utilizes bacterial expression to deliver individual peptides for behavioral rescue phenotypes. Our technology is advantageous over transgenic studies, as it allows for functional characterizing individual peptides. This tool readily allows for testing newly de-orphanized neuropeptide/neuropeptide receptor pairings [66]. This genetic tool is built off the principles of RNAi feeding techniques, though for a restoration rather than a reduction of function, to supply worms with the peptide of interest through their food source as previously shown for the scorpion venom peptide mBmKTX to alter lifespan and egg-laying behavior in *C. elegans* [34]. Based on our results in the present study, we believe that this method can be used to restore neuropeptide function within the different class of neuropeptides in *C. elegans.*

Our future investigation will be focused primarily on uncovering the mechanisms by which the bacteria create and deliver the neuropeptide macromolecule to *C. elegans*. We believe it unlikely that the *C. elegans* are processing the DNA. Even though DH5α *E. coli* is primarily used to make DNA, and does not contain T7 polymerase, it does contain *E. coli* RNA polymerase which could recognize additional promoter sites [88]. If the *E. coli* isn't producing the RNA, then that requires the *C. elegans* to be doing so. Like other eukaryotes, *C. elegans* have 3 types of RNA polymerases which each produce a distinct type of RNA all within the nucleus [89, 90]. For this rescue by feeding paradigm, we would expect RNA polymerase II to be producing the RNA since we expect ssRNA acting as mRNA to be produced and then fabricated into neuropeptides at the ribosome should *C. elegans* be the ones required to make the protein [90]. This would again require the neuropeptide vector to contain promoter sites to be recognized by RNA polymerase II [91-93]. If the *E. coli* is producing ssRNA, there are mechanisms by which this macromolecule can translocate into the nucleus, as it is becoming more open to investigation with the rise in mRNA vaccines [94].

Previous studies have shown that RNA produced by bacteria in the gut microbiota can be transported across the intestinal epithelium [80, 81], and our vector for neuropeptide delivery is modeled after RNAi feeding [38, 77, 78, 80]. Delivery of

exogenous scorpion venom peptide proved effective in changing *C. elegans* behavior

[34]. In humans, neuropeptides and neurotransmitters are produced by the gut

microbiome and transferred to the nervous system for use [95]. Our neuropeptide

delivery method may already make use of some mechanisms by which these other

techniques function or could be using novel mechanisms of delivery. Furthermore,

previous studies have shown that RNA uptake can alter transgenerational inheritance

[96], despite degradation rates. Our rescue of the exploratory behavior of *pdf-1 lof*

males even in the 24-hour timescale suggests that our feeding paradigm exploits similar

mechanisms, allowing peptide-encoding mRNAs to remain present throughout the

assay. We propose that if peptides were taken up, their degradation rates would likely

impede rescue efficiency at later timepoints, unlike what we observe in our experiments,

suggesting that the mechanism is similar to dsRNA feeding, with mRNA uptake driving

rescue rather than peptide uptake.

Based on our results, we propose that the neuropeptide rescue-by-feeding

strategy delivers mRNA that is ready for translation by the *C. elegans* cellular

machinery, rather than supplying *C. elegans* with fully translated and processed

peptides. The plasma membrane of a cell is an intricate complex of multiple lipid and

protein molecules. Small molecules with moderate polarity diffuse through the cell

membrane passively, but most metabolites and short peptides require specialized

membrane transporters for translocation [97]. Given the large number of neuropeptides

encoded in the genome of *C. elegans* [1], having specialized transporters for peptide

transport is not feasible [98]. The most substantial evidence for this statement is that the

processed INS-6 peptide is 54 amino acids in length and INS-22 is 68: the *C. elegans*

intestine expresses peptide transporters that only uptake smaller, inactive di- and tri-

amino acid peptide chains [98]. Thus, rescue of chemotaxis behavior by INS-6 peptide

feeding and INS-22 suggests that the mechanism of rescue may be similar to double

stranded RNA (dsRNA) uptake, rather than peptidergic transport across the intestinal

membrane [99, 100]. Future investigations will see if there is a limit to the size of

peptide that can be rescued.

Understanding neuropeptide function is essential from both the perspective of
regulating neuronal (synaptic and non-synaptic) channels of communication and from a
broader view of general neuronal functions throughout the brain. Revealing how a
specific neuropeptide acts at both the cellular and subcellular receptor levels is a critical
link in comprehending the role of neuromodulation in circuits. An enhanced
experimental pipeline for investigating peptide function will facilitate progress toward
understanding how neural circuit activity within the network is modulated by
neuropeptides in its various states, resulting in flexible decision-making during
behaviors.

Data Availability

Strains (**Table S1**) and plasmids (**Table S2**) are available upon request. The authors
affirm that all data necessary for confirming the conclusions of the article are present
within the article, figures, and tables. The raw data or a summary containing the
average, n, standard error measure, and associated statistical tests for all figures are
available upon request.

Acknowledgments

We thank Prof. Frank Schroeder, Cornell University, for providing ascr#8 and Prof.
Victor Ambros for reagents. We want to thank Dr. Arantza Barrios for help in executing
the leaving assay analysis in R. We extend our thanks to Dr. Sreekanth H. Chalasani
and Dr. Douglas Portman for their insights and updates to the method for testing the
phenotypes related to *ins-6* and *pdf-1*, respectively. pDest-527 was a gift from Dominic
Esposito (Addgene plasmid # 11518; http://n2t.net/addgene:11518 ;
RRID:Addgene_11518). We thank the *Caenorhabditis* Genetics Center for supplying
some strains; the CGC is funded by NIH Office of Research Infrastructure Programs
(P40 OD010440). The INS-22 strain was provided by the *C. elegans* Gene Knockout
Project at the Oklahoma Medical Research Foundation, which was part of the
International *C. elegans* Gene Knockout Consortium.

Funding

This work was funded by NIH R01DC016058 (JS), NIH DP2NS132372 (RNA), and NIH
F31AG081095 (EJL).

Conflict of Interest

EMD, DKR, and JS would like to disclose the filing of Utility Patent US-20220380430-A1
from May 20, 2022, which covers details of the neuropeptide vector and feeding.

Figure 1: Behavioral of neuropeptide associated behavior in loss of function mutant background

A) *pdf-1;him-5 lof* mutant males fed a SCRAMBLE peptide have a significantly lower probability of leaving food when compared to wildtype control *him-5* males ([####]p $\leq$ 0.0001, unpaired t-test). When fed any of the *pdf-1* produced peptides in a DH5α *E. coli* vehicle, the probability of leaving is significantly higher than the SCRAMBLE fed *pdf-1* males (****p $\leq$ 0.0001, One-way ANOVA with Dunnett's post-hoc test).

B) When exposed to a water solvent control (ascr#8 -), both *him-8* and *flp-3* males act

the same. When exposed to ascr#8 (ascr#8 +), wildtype *him-8* males do not avoid the

cue while *flp-3;him-8 lof* males do. When fed FLP-3-9 peptide in a DH5α *E. coli* vehicle,

the avoidance towards ascr#8 by *flp-3 lof* males is significantly decreased (****$p \leq$

0.0001, Mann-Whitney test with Bonferroni correction, new p-value ****$p \leq 0.00005$).

C) Wildtype N2 animals are more attracted to high 750 mM NaCl than *ins-6 lof* animals.

When fed INS-6 in a DH5α *E. coli* vehicle, the attraction of *ins-6 lof* animals to NaCl is

significantly increased (*$p \leq 0.05$, ***$p \leq 0.001$, unpaired t-test with Bonferroni

correction, new p-value *$p \leq 0.025$, ***$p \leq 0.0005$).

Figure 2: Neuropeptide rescue by feeding can be accomplished in different *E. coli* vehicles

A) *him-5* males fed a SCRAMBLE peptide in iOP50 *E. coli* leave food at a higher rate than *pdf-1;him-5 lof* males ($^{####}$p ≤ 0.0001, unpaired t-test). *pdf-1;him-5 m*ales fed PDF-1 peptides in iOP50 and then plated on an assay plate containing OP50 *E. coli* without peptide, do leave food at a higher rate than *pdf-1* SCRAMBLE iOP50 *E. coli* fed animals (****p ≤ 0.0001, One-way ANOVA with Dunnett's post-hoc test).

B) *him-5* males fed SCRAMBLE peptide in iOP50 *E. coli* and then placed on an assay plate seeded with SCRAMBLE peptide in iOP50 *E. coli* leave food at a higher rate than

pdf-1;him-5 lof males ($^{####}$p ≤ 0.0001, unpaired t-test). *pdf-1;him-5* males fed PDF-1

peptides in iOP50 and then plated on an assay plate containing iOP50 *E. coli*

expressing the corresponding peptide, leave food at a higher rate than *pdf-1;him-5 lof*

animals fed SCRAMBLE iOP50 *E. coli* and at a higher rate than animals on assay

plates without peptide (****p ≤ 0.0001, One-way ANOVA with Dunnett's post-hoc test).

C) When exposed to a water solvent control (ascr#8 -), both *him-8* and *flp-3;him-8*

males act the same. When exposed to ascr#8 (ascr#8 +), wildtype *him-8* males do not

avoid the cue while *flp-3 lof* males do. When fed FLP-3-9 peptide in iOP50 *E. coli*, the

avoidance towards ascr#8 by *flp-3 lof* males is significantly decreased (****p ≤ 0.0001,

Mann-Whitney test with Bonferroni correction, new p-value ****p ≤ 0.00005).

D) Animals deficient in *ins-22* show significantly stronger chemotaxis towards 10%

butanone than *ins-22 lof* animals fed INS-22. The *ins-22 lof* animals fed INS-22 have a

chemotaxis index no different than wildtype animals (^{ns}p ≥ 0.05, ***p ≤ 0.001, unpaired

t-test wildtype SCRAMBLE to INS-22, Mann-Whitney test comparisons to *ins-22* fed

SCRAMBLE with Bonferroni correction (new p-value ^{ns}p ≥ 0.025, ****p ≤ 0.00005).

Figure 3: Neuropeptide receptor required for neuropeptide feeding

A) *pdfr-1;him-5 lof* animals do not leave food more than 35mm, even if fed PDF-1 peptides in DH5α *E. coli*. Double mutants of *pdf-1;pdfr-1* in a *him-5* background also did not leave food at the rate of *pdf-1* animals fed SCRAMBLE, though *pdf-1;pdfr-1;him-5* animals fed PDF-1A DH5α *E. coli* do have a higher incidence of leaving food than

774   SCRAMBLE fed animals ($^{ns}p \geq 0.05$, ****$p \leq 0.0001$, One-way ANOVA within genotypes

775   with Dunnett's post hoc test).

776   **B)** Double mutant *flp-3;frpr-16* males in a *him-8* background fed bacteria expressing

777   FLP-3-9 avoid ascr#8 to the same level as SCRAMBLE DH5α *E. coli* fed animals ($^{ns}p \geq$

778   0.05, Mann-Whitney test with Bonferroni correction, new p-value $^{ns}p \geq 0.025$).

Figure 4: Behaviors resulting from overexpression of neuropeptides depend on the neuropeptide

A) *him-5* males fed PDF-1A in DH5α *E. coli* had a significant decrease in their probability of leaving food rate compared to animals fed SCRAMBLE peptide. PDF-1B *him-5* animals had a significant increase in P_L/hr when compared to SCRAMBLE fed animals. When PDF-1A and -1B were fed to animals in equal combination, their probability of leaving food was no different from SCRAMBLE fed animals (****$p \leq$ 0.0001, One-way ANOVA with Dunnett's post-hoc test).

B) *him-5* males fed PDF-1 peptides in iOP50 *E. coli* and then tested on iOP50 assay plates containing a corresponding peptide, do not have excursions leaving food more than 35 mm at higher rates than SCRAMBLE fed males (**$p \leq 0.01$, ****$p \leq 0.0001$, One-way ANOVA with Dunnett's post-hoc test).

C) *him-8* males exposed to ascr#8 had no difference in their avoidance behavior whether they were fed bacteria expressing SCRAMBLE or FLP-3-9 in DH5α *E. coli* ($^{ns}p \geq 0.05$, Mann-Whitney test).

D) *him-8* males fed SCRAMBLE or FLP-3-9 in iOP50 *E. coli* avoid ascr#8 at the same rate and behave the same towards the water solvent control as the previous studies ($^{ns}p \geq 0.05$, Mann-Whitney test).

E) Wildtype hermaphrodites fed SCRAMBLE or INS-6 peptide in DH5α *E. coli* displayed the same attraction towards high 750 mm NaCl ($^{ns}p \geq 0.05$, unpaired t-test test).

F) Wildtype hermaphrodites fed INS-22 peptide in iOP50 *E. coli* displayed slightly less attraction towards 10% butanone than animals fed SCRAMBLE (*$p \leq 0.05$, unpaired t-test).

G) Wildtype hermaphrodites fed INS-22 peptide in iOP50 *E. coli* were not as attracted to 0.1% butanone compared to animals fed SCRAMBLE (****$p \leq 0.0001$, unpaired t-test).

Supplemental Material

Figure S1: Endogenous Processing of Neuropeptides in *C. elegans*

Endogenous neuropeptide processing occurs inside neurons. From a neuropeptide precursor gene, an entire pre-propeptide is translated. We tested a representative

peptide from each neuropeptide class in *C. elegans.* The DNA sequences for the

neuropeptide precursor genes were from Wormbase.org and the putative unprocessed

protein sequence and AlphaFold protein structure were obtained from uniprot.org. Once

the pre-propeptide is translated, it is then processed by a series of enzymes. First the

signal peptide is removed. And then the individual peptides are cleaved, at sites

indicated by red arrows. We designed our neuropeptide constructs to contain these

convertase enzyme cleavage sites, so we would anticipate that our delivered

neuropeptide could be processed as demonstrated. Cleavage of individual peptides

happens by the proprotein convertase enzymes. *C. elegans* have four protein

convertases serving different functions in various tissues. Most common for

neuropeptide processing appears to be EGL-3/KPC-2 (PC2 homolog cleaves K/R) but

there are also, KPC-1 (cleaving K/R in dendrites), BLI-4/KPC-4 (cleaving K/R in the

cuticle), and AEX-5/KPC-3 (cleaving paired basic amino acids in the muscle and

intestine) (last three are PC1 homologs) [56, 101-103]. After protein cleavage creates

individual neuropeptides, the peptides can be post-translationally modified. The FLP

[68] and NLP peptides are often amidated [19] while INS peptides generally have

disulfide bonds running between their α and β chains [58, 63]. Once the neuropeptides

are processed, they are packaged into dense core vesicles for release at the synapse.

They can be released with or without neurotransmitters. After binding to GPCRs on

post-synaptic neurons, they slowly diffuse away from synapse and are broken down;

this occurs at a slower rate than neurotransmitter reuptake leading to longer lasting

neuropeptide effects [18, 104]. Created in BioRender. DiLoreto, E. (2025)

https://BioRender.com/dyorxwg

Figure S2: Neuropeptide gene deficient animals exhibit different behaviors than
they wildtype counterparts

A) *pdf-1;him-5* males fed OP50 without peptide have a significantly lower rate of leaving
food beyond 35 mm compared to wildtype *him-5* animals (****p $\leq$ 0.0001, unpaired t-
test).

B) The location of males can be traced across time and binned as animals who never
left food (0 mm), those that had a minor excursion (10 mm >), and those with a major
excursion (10 mm <). This is then converted to a percentage and plotted across time.
These values contribute to the probability of leaving rate. *pdf-1* animals fed OP50
without peptide have lower rates of major excursions compared to wildtype *him-5*
animals.

C) SCRAMBLE DH5α *E. coli* fed *pdf-1 lof* males have a lower percentage of animals
making major excursions off of food than *him-5* animals.

D) Compared to wildtype *him-8* males, *flp-3;him-8 lof* males fed OP50 *E. coli* without
peptide avoid ascr#8 at a significantly higher rate (****p $\leq$ 0.0001, Mann-Whitney test).

E) Compared to wildtype animals, *ins-6* animals fed OP50 *E. coli* without peptide have a
lower choice index towards high 750 mm NaCl (****p $\leq$ 0.0001, unpaired t-test).

Figure S3: Bacterial cell supernatant from DH5α cells liquid culture growth alone may be sufficient to restore neuropeptide behavior

him-8 lof males fed the supernatant or cells derived from fed DH5α bacterial culture expressing the SCRAMBLE or FLP-3-9 peptide did not alter the lack of avoidance towards ascr#8. *flp-3;him-8* males fed the supernatant from a DH5α culture expressing FLP-3-9 did exhibit a suppression of avoidance behavior. (comparisons between ascr#8 stimulus (+) conditions within genotypes, $^{ns}p \geq 0.05$, $*p \leq 0.05$, $**p \leq 0.01$, $****p \leq 0.0001$, One-way ANOVA with Tukey's multiple comparison)

863 **Figure S4: Plasmid maps of old Gateway cloning and new constructions**

864 **A)** pDest-527 vector used in Gateway cloning. Created in SnapGene

865 **B)** pET-21a(+) vector used in Genscript XhoI/XhoI construction. Created in SnapGene

866 **C)** Chemotaxis values of wildtype (N2 Bristol) and *ins-22 lof* animals towards 10%
867 butanone. (***$p \leq 0.001$, Mann-Whitney test).

Figure S5: Neuropeptide receptor *lof* animals exhibit defective behavior without peptide feeding

A) *pdf-1;pdfr-1 lof* in a *him-5* background do leave food greater than 35 mm at a higher

rate than *pdfr-1;him-5 lof* males (****p $\leq$ 0.0001, unpaired t-test).

B) Excursion rates of neuropeptide receptor mutants (*pdfr-1;him-5*) and

neuropeptide/neuropeptide receptor mutants (*pdf-1;pdfr-1;him-5*) fed OP50 *E. coli* do

not readily leave food or have a major excursion 10 mm <*.*

C) *pdf-1;him-5* neuropeptide receptor mutants fed PDF-1 peptides in DH5αa *E. coli* do

not have excursions off food at high rates like *him-5* males.

D) *flp-3;frpr-16* double mutants in a *him-8* background avoid ascr#8 at a higher rate than

the solvent control (****p $\leq$ 0.0001, Mann-Whitney test).

Figure S6: Overexpression of wildtype animals fed neuropeptide in DH5α or iOP50 *E. coli*

A) Most *him-5* males fed SCRAMBLE or PDF-1 peptides in DH5α *E. coli* made a major excursions traveling < 10 mm from food by 2 hours after plating.

B) *him-5* males do not remain on food at the rates of *pdf-1;him-5* animals fed PDF-1 peptides in iOP50 *E. coli* and tested on iOP50 assay plates containing peptides. *pdf-1;him-5* males fed PDF-1 peptides in iOP50 *E. coli* have excursions off food at higher rates than SCRAMBLE-fed animals.

C) *him-5* males fed PDF-1 peptides in iOP50 *E. coli* and then tested on OP50 assay plates not containing peptide, do not have excursions leaving food more than 35 mm at higher rates than SCRAMBLE fed males (*p ≤ 0.05, **p ≤ 0.01,****p ≤ 0.0001, One-way ANOVA with Dunnett's post-hoc test).

893 **D)** *him-5* males do not remain on food at the rates of *pdf-1;him-5* animals fed PDF-1
894 peptides in iOP50 *E. coli* and tested in OP50 assay plates not containing peptides.

Figure S7: Investigating putative neuropeptide uptake mechanisms

A) Choice index of OP50 fed animals moving towards 0 or 750 mM NaCl. Mutants
lacking *ins-6* showed reduced preference for 750 mM NaCl. ($^{ns}p \geq 0.05$, *$p \leq 0.05$, **p
≤ 0.01,****$p \leq 0.0001$, One-way ANOVA with Dunnett's multiple comparison).

B) Choice index of dsRNA transport mutant animals (*sid-1*) demonstrated that when fed
DH5α bacteria expressing SCRAMBLE peptide both *ins-6 lof* and *sid-1;ins-6 lof* animals
displayed defective NaCl attraction. After feeding of DH5α *E. coli* expressing INS-6,
there was no statistically significant difference in the behavior towards NaCl compared
to wildtype animals. (comparisons made within peptide feeding condition, $^{ns}p \geq 0.05$, *p
≤ 0.05, **$p \leq 0.01$,****$p \leq 0.0001$, One-way ANOVA with Dunnett's multiple
comparison).

Table S1: *C. elegans* Strains Used in Current Study

Strain Name	Genotype	Source Lab	Used in Paper

N2	Bristol Wildtype isolate	CGC	1c, S1e, 2d, S2c, 4e-g
CB1489	*him-8(e1489)*IV	CGC	1b, S1d, 2c, 3b, S3d, 4c,d. S5a
CB4088	*him-5(e1490)*V	CGC	1a, S1a-c, 2a,b, 3a, S3a-c, 4a, b, S4
HC196	*sid-1(qt9)*V	CGC	S5b,c
IV302	*ins-6(tm2416)*II; kyEx2595[*str-2*p::GCaMP2.2b + *unc-122*p::GFP]	Shreekanth Chalasani	1c, S1e, 4e
JSR99	*flp-3(pk361)*X; *him-8(e1489)*IV	Jagan Srinivasan	1b, s1d, 2c, 3b, S3d, 4c,d, S5a
JSR152	*flp-3(pk361)*X; *frpr-16(gk5305*[*lox*p + *myo-2*p::GFP::*unc-54 3' UTR* + *rps-27*p::neoR::*unc-54 3' UTR* + *lox*p]II)	Jagan Srinivasan	3b, s3d
JSR166	*ins-6(tm2416)*II; kyEx2595[*str-2*p::GCaMP2.2b + *unc-122*p::GFP]; *sid-1(qt9)*V	Jagan Srinivasan	S5b,c
JSR188	*pdf-1(tm1996)*III; *pdfr-1(ok3425)*III; *him-5(e1490)*V	Jagan Srinivasan	3a, S3a-c
UR930	*pdfr-1(ok3425)*III; *him-5(e1490)*V	Douglas Portman	3a, S3a-c
UR954	*pdf-1(tm1996)*III; *him-5(e1490)*V	Douglas Portman	1a, S1a-c, 2a,b, 3a, S3a-c, 4a, b, S4
RB2594	*ins-22(ok3616)*III	CGC	2d, S2c, 4f,g

911 **Table S2: Neuropeptide Oligo Sequences Ordered for Individual Peptide Feeding**

912 Neuropeptide DNA sequence indicated in lowercase.

Peptide	Animo Acid (C-x-N)	Sequence Ordered

SCRAMBLE	NSKLHRGGGRSRTSGSTGSMASHARGSPGLQ	For:GGGGACAAGTTTGTACAAAAAAGCAGGCTGGATGCGCTTTGGAAAACGTaattcgagctccaccgcggtggcggccgctctagaactagtggatccaccggttccatggctagccacgcgcgtggatcccccgggctgcaggAAACGTAAATAACACCCAGCTTTCTTGTACAAAGTGGTCCCC
		Rev:GGGGACCACTTTGTACAAGAAAGCTGGGTGTTATTTACGTTTcctgcagcccggggatccacgcgcgtggctagccatggaaccggtggatccactagttctagagcggccgccaccgcggtggagctcgaattACGTTTTCCAAAGCGCATCCAGCCTGCTTTTTTGTACAAACTTGTCCCC
PDF-1A	VQFVKRSNAELINGLIGMDLGKLSAVGKR	For:GGGGACAAGTTTGTACAAAAAAGCAGGCTGGGTTCAGTTCGTaaaacgcagcaacgccgagcttatcaacggactcatcggaatggatttgggaaaattgtcagctgtcggaAAACGCTGACACCCAGCTTTCTTGTACAAAGTGGTCCCC
		Rev:GGGGACCACTTTGTACAAGAAAGCTGGGTGTTATTTACGTTTtcagcgttttccgacagctgacaattttcccaaatccattccgatgagtccgttgataagctcggcgttgctgcgttttacgaactgaacACGTTTTCCAAAGCGCATCCAGCCTGCTTTTTTGTACAAACTTGTCCCC
PDF-1B	VQFVKRSNAELINGLLSMNLNKLSGAGRR	For:GGGGACAAGTTTGTACAAAAAAGCAGGCTGGGTTCAGTTCGTAAAACGCTcaaacgcggaacttatcaacggtcttctcagcatgaacctcaacAAATTGTCTGGAGCTGGTCGACGATGACACCCAGCTTTCTTGTACAAAGTGGTCCCC
		Rev:GGGGACCACTTTGTACAAGAAAGCTGGGTGTTATTTACGTTTtcatcgtcgaccagctccagacaaatttgttgaggttcatgctgagaagaccgttgataagttccgcgtttgagcgttttacgaactgaacACGTTTTCCAAAGCGCATCCAGCCTGCTTTTTTGTACAAACTTGTCCCC
FLP-3-9	NPENDTPFGTMRFG	For:GGGGACAAGTTTGTACAAAAAAGCAGGCTGGATGCGCTTTGGAAAACGTaatcctgagaacgacacaccattcggaacaatgagatttggaAAACGTAAATAACACCCAGCTTTCTTGTCAAAGTGGTCCCC
		Rev:GGGGACCACTTTGACAAGAAAGCTGGGTGTTATTTACGTTTtccaaatctcattgttccgaatggtgtgtcgttctcaggattACGTTTTCCAAAGCGCATCCAGCCTGCTTTTTTGTACAAACTTGTCCCC
INS-6	VPAPGETRACGRKLISLVMAVCGDLCNPQEGKDIATECCGNQCSDDYIRSACCP	For:GGGGACAAGTTTGTACAAAAAAGCAGGCTGGATGCGCTTTGGAAAACGTgttccagcaccaggagaaactcgtgcctgcggaagaaaactcatctctttagtcatggctgtctgtggagatctttgcaacccacaagaaggaaaggacattgcgactgaatgctgcggaaatcagtgttctgatgactacataagatctgcttgttgtccaAAACGTAAATAACACCCAGCTTTCTTGTACAAAGTGGTCCCC
		Rev:GGGGACCACTTTGTACAAGAAAGCTGGGTGTTATTTACGTTTtggacaacaagcagatcttatgtagtcatcagaacactgatttccgcagcattcagtcgcaatgtcctttccttcttgtgggttgcaaagatctccacagacagccatgactaaagagatgagttttcttccgcaggcacgagtttctcctggtgctggaacACGTTTTCCAAAGCGCATCCAGCCTGCTTTTTTGTACAAACTTGTCCCC
INS-22	MHTTTILICFFIFLVQVSTMDAHTDKYVRTLCGKTAIRNIANLCPPKPEMKGICSTGEYPSITEYCSM	ATGAGGTTTGGAAAAAGAatgcacactacaacgatcctgatttgttttttcatcttcttggtccaggttagcaccatggatgctcacaccgataaatacgtgcgcaccctgtgcggcaagaccgcaattcgtaatattgcgaacctgtgtccgccaaaaccggaaatgaagggcatctgctctactggtgagtatccgagcattaccgagtactgctccatgggttttagcgactcgcaaatcaagttcatgtgctgcgacaaccagAAACGTAAATAA

References

1. Li, C. and K. Kim, *Neuropeptides.* WormBook : the online review of *C. elegans*
biology, 2008: p. 1-36.
2. Dale Purves, G.J.A., David Fitzpatrick, William C Hall, Anthony-Samuel
LaMantia, Richard D Mooney, Michael L Platt, Leonard E White, *Neuroscience.* 6
ed. 2018, Sunderland, MA: Oxford University Press.
3. Yañez-Guerra, L.A., D. Thiel, and G. Jékely, *Premetazoan Origin of Neuropeptide*
Signaling. Mol Biol Evol, 2022. **39**(4).
4. Beets, I., et al., *Vasopressin/oxytocin-related signaling regulates gustatory*
associative learning in C. elegans. Science, 2012. **338**(6106): p. 543-5.
5. Gruber, C.W., *Physiology of invertebrate oxytocin and vasopressin*
neuropeptides. Experimental Physiology, 2014. **99**(1): p. 55-61.
6. Insel, T.R. and L.J. Young, *The neurobiology of attachment.* Nature Reviews
Neuroscience, 2001. **2**(2): p. 129-136.
7. Willets, J.M., et al., *Regulation of oxytocin receptor responsiveness by G protein-*
coupled receptor kinase 6 in human myometrial smooth muscle. Molecular
endocrinology (Baltimore, Md.), 2009. **23**(8): p. 1272-1280.
8. Lukas, M., et al., *The Neuropeptide Oxytocin Facilitates Pro-Social Behavior and*
Prevents Social Avoidance in Rats and Mice. Neuropsychopharmacology, 2011.
36(11): p. 2159-2168.
9. Odekunle, E.A., et al., *Ancient role of vasopressin/oxytocin-type neuropeptides*
as regulators of feeding revealed in an echinoderm. BMC Biology, 2019. **17**(1): p.
60.
10. Garrison, J.L., et al., *Oxytocin/vasopressin-related peptides have an ancient role*
in reproductive behavior. Science, 2012. **338**(6106): p. 540-3.
11. Frooninckx, L., et al., *Neuropeptide GPCRs in C. elegans.* Frontiers in
Endocrinology, 2012. **3**(167).
12. Nusbaum, M.P., D.M. Blitz, and E. Marder, *Functional consequences of*
neuropeptide and small-molecule co-transmission. Nat Rev Neurosci, 2017.
18(7): p. 389-403.
13. Souza-Moreira, L., et al., *Neuropeptides as pleiotropic modulators of the immune*
response. Neuroendocrinology, 2011. **94**(2): p. 89-100.
14. Nässel, D.R. and M. Zandawala, *Endocrine cybernetics: neuropeptides as*
molecular switches in behavioural decisions. Open Biol, 2022. **12**(7): p. 220174.
15. Tai, K.Y., et al., *Selected neuropeptide genes show genetic differentiation*
between Africans and non-Africans. BMC Genetics, 2020. **21**(1): p. 31.
16. *PeptidesDB.* 2008 April 30 2008 [cited 2022; Available from: /www.peptides.be.
17. Burbach, J.P.H., *Neuropeptides from concept to online database*
www.neuropeptides.nl. European Journal of Pharmacology, 2010. **626**(1): p. 27-
48.
18. Russo, A.F., *Overview of Neuropeptides: Awakening the Senses?* Headache,
2017. **57 Suppl 2**(Suppl 2): p. 37-46.
19. Van Bael, S., et al., *Mass spectrometric evidence for neuropeptide-amidating*
enzymes in Caenorhabditis elegans. J Biol Chem, 2018. **293**(16): p. 6052-6063.

20.　White, J.G., et al., *The structure of the nervous system of the nematode*
Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci, 1986. **314**(1165): p.
1-340.
21.　Sammut, M., et al., *Glia-derived neurons are required for sex-specific learning in*
C. elegans. Nature, 2015. **526**(7573): p. 385-390.
22.　Li, C., K. Kim, and L.S. Nelson, *FMRFamide-related neuropeptide gene family in*
Caenorhabditis elegans. Brain Res, 1999. **848**(1-2): p. 26-34.
23.　Pierce, S.B., et al., *Regulation of DAF-2 receptor signaling by human insulin and*
ins-1, a member of the unusually large and diverse C. elegans insulin gene
family. Genes Dev, 2001. **15**(6): p. 672-86.
24.　Nathoo, A.N., et al., *Identification of neuropeptide-like protein gene families in*
Caenorhabditiselegans and other species. Proc Natl Acad Sci U S A, 2001.
98(24): p. 14000-5.
25.　Hobert, O., *The neuronal genome of Caenorhabditis elegans.*, in *WormBook: The*
Online Review of C. elegans Biology. 2005-2018: Pasadena, CA.
26.　Watteyne, J., et al., *Neuropeptide signaling network of Caenorhabditis elegans:*
from structure to behavior. Genetics, 2024. **228**(3).
27.　Barrios, A., et al., *PDF-1 neuropeptide signaling modulates a neural circuit for*
mate-searching behavior in C. elegans. Nature Neuroscience, 2012. **15**(12): p.
1675-1684.
28.　Hussey, R., et al., *Oxygen-sensing neurons reciprocally regulate peripheral lipid*
metabolism via neuropeptide signaling in Caenorhabditis elegans. PLoS
genetics, 2018. **14**(3): p. e1007305-e1007305.
29.　Nath, R.D., et al., *C. elegans Stress-Induced Sleep Emerges from the Collective*
Action of Multiple Neuropeptides. Current biology, 2016. **26**(18): p. 2446-2455.
30.　Nelson, M.D., et al., *FRPR-4 Is a G-Protein Coupled Neuropeptide Receptor*
That Regulates Behavioral Quiescence and Posture in Caenorhabditis elegans.
PLoS One, 2015. **10**(11): p. e0142938.
31.　Van Bael, S., et al., *Identification of Endogenous Neuropeptides in the Nematode*
C. elegans Using Mass Spectrometry. Methods Mol Biol, 2018. **1719**: p. 271-291.
32.　Papaioannou, S., et al., *Role of a FMRFamide-like family of neuropeptides in the*
pharyngeal nervous system of Caenorhabditis elegans. Journal of Neurobiology,
2005. **65**(3): p. 304-319.
33.　Ohno, H., et al., *Luqin-like RYamide peptides regulate food-evoked responses in*
C. elegans. Elife, 2017. **6**.
34.　Xu, J., et al., *Feeding recombinant E. coli with GST-mBmKTX fusion protein*
increases the fecundity and lifespan of Caenorhabditis elegans. Peptides, 2017.
89: p. 1-8.
35.　Fraser, A.G., et al., *Functional genomic analysis of C. elegans chromosome I by*
systematic RNA interference. Nature, 2000. **408**(6810): p. 325-330.
36.　Kamath, R.S., et al., *Effectiveness of specific RNA-mediated interference through*
ingested double-stranded RNA in Caenorhabditis elegans. Genome Biol, 2001. **2**.
37.　Dirksen, P., et al., *CeMbio - The Caenorhabditis elegans Microbiome Resource.*
G3 (Bethesda), 2020. **10**(9): p. 3025-3039.
38.　Reilly, D.K., et al., *Distinct neuropeptide-receptor modules regulate a sex-specific*
behavioral response to a pheromone. Commun Biol, 2021. **4**(1): p. 1018.

39. Brenner, S., *The genetics of Caenorhabditis elegans.* Genetics, 1974. **77**(1): p.
71-94.
40. Neve, I.A.A., et al., *Escherichia coli Metabolite Profiling Leads to the*
Development of an RNA Interference Strain for Caenorhabditis elegans. G3
(Bethesda), 2020. **10**(1): p. 189-198.
41. Ryan, D.A., et al., *Sex, age, and hunger regulate behavioral prioritization through*
dynamic modulation of chemoreceptor expression. Curr Biol, 2014. **24**(21): p.
2509-17.
42. Barrios, A., et al., *PDF-1 neuropeptide signaling modulates a neural circuit for*
mate-searching behavior in C. elegans. Nat Neurosci, 2012. **15**(12): p. 1675-82.
43. Lipton, J., et al., *Mate searching in Caenorhabditis elegans: a genetic model for*
sex drive in a simple invertebrate. J Neurosci, 2004. **24**(34): p. 7427-34.
44. Hilliard, M.A., et al., *Worms taste bitter: ASH neurons, QUI-1, GPA-3 and ODR-3*
mediate quinine avoidance in Caenorhabditis elegans. EMBO J, 2004. **23**(5): p.
1101-11.
45. Leinwand, S.G. and S.H. Chalasani, *Neuropeptide signaling remodels*
chemosensory circuit composition in Caenorhabditis elegans. Nature
neuroscience, 2013. **16**(10): p. 1461-1467.
46. Bargmann, C.I. and H.R. Horvitz, *Chemosensory neurons with overlapping*
functions direct chemotaxis to multiple chemicals in C. elegans. Neuron, 1991.
7(5): p. 729-42.
47. Phillips, C.M., et al., *HIM-8 binds to the X chromosome pairing center and*
mediates chromosome-specific meiotic synapsis. Cell, 2005. **123**(6): p. 1051-63.
48. Meneely, P.M., et al., *Crossover distribution and frequency are regulated by him-*
5 in Caenorhabditis elegans. Genetics, 2012. **190**(4): p. 1251-66.
49. Reilly, D.K., *Neuromodulation of Sex-Specific Pheromone-Mediated Behaviors*, in
Biology and Biotechnology. 2020, Worcester Polytechnic Institute: Worcester,
MA. p. 519.
50. Janssen, T., et al., *Functional characterization of three G protein-coupled*
receptors for pigment dispersing factors in Caenorhabditis elegans. J Biol Chem,
2008. **283**(22): p. 15241-9.
51. Janssen, T., et al., *Discovery and characterization of a conserved pigment*
dispersing factor-like neuropeptide pathway in Caenorhabditis elegans. J
Neurochem, 2009. **111**(1): p. 228-41.
52. Muirhead, C.S. and J. Srinivasan, *Small molecule signals mediate social*
behaviors in C. elegans. J Neurogenet, 2020. **34**(3-4): p. 395-403.
53. Pungaliya, C., et al., *A shortcut to identifying small molecule signals that regulate*
behavior and development in Caenorhabditis elegans. Proc Natl Acad Sci U S A,
2009. **106**(19): p. 7708-13.
54. Salama, R., E. DiLoreto, and J. Srinivasan, *The Architect of Neurotransmission in*
C. elegans : How FLP-3 Neuropeptides' Structures Direct their Function.
MicroPubl Biol, 2023. **2023**.
55. Chen, Z., et al., *Two insulin-like peptides antagonistically regulate aversive*
olfactory learning in C. elegans. Neuron, 2013. **77**(3): p. 572-585.

56.   Hung, W.L., et al., *A Caenorhabditis elegans developmental decision requires*
1049   *insulin signaling-mediated neuron-intestine communication.* Development, 2014.
1050   **141**(8): p. 1767-79.
1051   57.   Hua, Q.-X., et al., *A divergent INS protein in Caenorhabditis elegans structurally*
1052   *resembles human insulin and activates the human insulin receptor.* Genes &
1053   development, 2003. **17**(7): p. 826-831.
1054   58.   Zheng, S., et al., *A functional study of all 40 Caenorhabditis elegans insulin-like*
1055   *peptides.* J Biol Chem, 2018. **293**(43): p. 16912-16922.
1056   59.   Leinwand, S.G. and S.H. Chalasani, *Neuropeptide signaling remodels*
1057   *chemosensory circuit composition in Caenorhabditis elegans.* Nat Neurosci,
1058   2013. **16**(10): p. 1461-7.
1059   60.   Gao, B. and Q. Sun, *Programming gene expression in multicellular organisms for*
1060   *physiology modulation through engineered bacteria.* Nature Communications,
1061   2021. **12**(1): p. 2689.
1062   61.   Carrillo Rincón, A.F., et al., *A dual-inducible control system for multistep*
1063   *biosynthetic pathways.* Journal of Biological Engineering, 2024. **18**(1): p. 68.
1064   62.   Shtonda, B.B. and L. Avery, *Dietary choice behavior in Caenorhabditis elegans.* J
1065   Exp Biol, 2006. **209**(Pt 1): p. 89-102.
1066   63.   Zhu, R. and I.D. Chin-Sang, *C. elegans insulin-like peptides.* Mol Cell Endocrinol,
1067   2024. **585**: p. 112173.
1068   64.   Chen, Y. and L.R. Baugh, *Ins-4 and daf-28 function redundantly to regulate C.*
1069   *elegans L1 arrest.* Dev Biol, 2014. **394**(2): p. 314-26.
1070   65.   Ripoll-Sánchez, L., et al., *The neuropeptidergic connectome of C. elegans.*
1071   Neuron, 2023. **111**(22): p. 3570-3589.e5.
1072   66.   Beets, I., et al., *System-wide mapping of peptide-GPCR interactions in C.*
1073   *elegans.* Cell Rep, 2023. **42**(9): p. 113058.
1074   67.   Hilbert, Z.A. and D.H. Kim, *PDF-1 neuropeptide signaling regulates sexually*
1075   *dimorphic gene expression in shared sensory neurons of C. elegans.* Elife, 2018.
1076   **7**.
1077   68.   Li, C. and K. Kim, *Family of FLP Peptides in Caenorhabditis elegans and Related*
1078   *Nematodes.* Frontiers in Endocrinology, 2014. **5**.
1079   69.   Van Bael, S., et al., *Identification of Endogenous Neuropeptides in the Nematode*
1080   *C. elegans Using Mass Spectrometry.* Methods in molecular biology (Clifton,
1081   N.J.), 2018. **1719**: p. 271-291.
1082   70.   Bael, S.V., *Neuropeptidomics in C. elegans*, in *Biochemistry and Biotechnology*.
1083   2018, KU Leuven: Betekom, Belgium. p. 216.
1084   71.   Van Bael, S., et al., *A Caenorhabditis elegans Mass Spectrometric Resource for*
1085   *Neuropeptidomics.* J Am Soc Mass Spectrom, 2018. **29**(5): p. 879-889.
1086   72.   Torayama, I., T. Ishihara, and I. Katsura, *Caenorhabditis elegans integrates the*
1087   *signals of butanone and food to enhance chemotaxis to butanone.* J Neurosci,
1088   2007. **27**(4): p. 741-50.
1089   73.   L'Etoile, N.D. and C.I. Bargmann, *Olfaction and odor discrimination are mediated*
1090   *by the C. elegans guanylyl cyclase ODR-1.* Neuron, 2000. **25**(3): p. 575-86.
1091   74.   Timmons, L., D.L. Court, and A. Fire, *Ingestion of bacterially expressed dsRNAs*
1092   *can produce specific and potent genetic interference in Caenorhabditis elegans.*
1093   Gene, 2001. **263**(1-2): p. 103-12.

75.   Fire, A., et al., *Potent and specific genetic interference by double-stranded RNA*
in Caenorhabditis elegans. Nature, 1998. **391**(6669): p. 806-11.
76.   Timmons, L. and A. Fire, *Specific interference by ingested dsRNA.* Nature, 1998.
395(6705): p. 854.
77.   Conte, D., Jr., et al., *RNA Interference in Caenorhabditis elegans.* Current
protocols in molecular biology, 2015. **109**: p. 26.3.1-26.3.30.
78.   Fraser, A.G., et al., *Functional genomic analysis of C. elegans chromosome I by*
systematic RNA interference. Nature, 2000. **408**(6810): p. 325-30.
79.   García, K., et al., *Inactivated E. coli transformed with plasmids that produce*
dsRNA against infectious salmon anemia virus hemagglutinin show antiviral
activity when added to infected ASK cells. Front Microbiol, 2015. **6**: p. 300.
80.   Kamath, R.S., et al., *Effectiveness of specific RNA-mediated interference through*
ingested double-stranded RNA in Caenorhabditis elegans. Genome Biol, 2001.
2(1): p. RESEARCH0002.
81.   Kaletsky, R., et al., *C. elegans interprets bacterial non-coding RNAs to learn*
pathogenic avoidance. Nature, 2020. **586**(7829): p. 445-451.
82.   Weber, F., et al., *Double-stranded RNA is produced by positive-strand RNA*
viruses and DNA viruses but not in detectable amounts by negative-strand RNA
viruses. J Virol, 2006. **80**(10): p. 5059-64.
83.   Shih, J.D. and C.P. Hunter, *SID-1 is a dsRNA-selective dsRNA-gated channel.*
RNA (New York, N.Y.), 2011. **17**(6): p. 1057-1065.
84.   Shih, J.D., et al., *The SID-1 double-stranded RNA transporter is not selective for*
dsRNA length. Rna, 2009. **15**(3): p. 384-90.
85.   *Cryo-EM structure reveals how SID-1 recognizes dsRNA that initiates systemic*
RNAi. Nat Struct Mol Biol, 2024. **31**(7): p. 1007-1008.
86.   Li, W., et al., *Systemic RNA Interference Deficiency-1 (SID-1) Extracellular*
Domain Selectively Binds Long Double-stranded RNA and Is Required for RNA
Transport by SID-1. J Biol Chem, 2015. **290**(31): p. 18904-13.
87.   Kostylev, M., et al., *Cloning Should Be Simple: Escherichia coli DH5α-Mediated*
Assembly of Multiple DNA Fragments with Short End Homologies. PLoS One,
2015. **10**(9): p. e0137466.
88.   Shimada, T., et al., *The whole set of constitutive promoters recognized by RNA*
polymerase RpoD holoenzyme of Escherichia coli. PLoS One, 2014. **9**(3): p.
e90447.
89.   Carter, R. and G. Drouin, *Structural differentiation of the three eukaryotic RNA*
polymerases. Genomics, 2009. **94**(6): p. 388-96.
90.   Girbig, M., A.D. Misiaszek, and C.W. Müller, *Structural insights into nuclear*
transcription by eukaryotic DNA-dependent RNA polymerases. Nature Reviews
Molecular Cell Biology, 2022. **23**(9): p. 603-622.
91.   Chen, R.A., et al., *The landscape of RNA polymerase II transcription initiation in*
C. elegans reveals promoter and enhancer architectures. Genome Res, 2013.
23(8): p. 1339-47.
92.   Bowman, E.A. and W.G. Kelly, *RNA polymerase II transcription elongation and*
Pol II CTD Ser2 phosphorylation: A tail of two kinases. Nucleus, 2014. **5**(3): p.
224-36.

93.   Zhu, Y., et al., *Quantitative analysis of transcription start site selection reveals*
control by DNA sequence, RNA polymerase II activity and NTP levels. Nat Struct
Mol Biol, 2024. **31**(1): p. 190-202.
94.   Sattar, S., et al., *Nuclear translocation of spike mRNA and protein is a novel*
feature of SARS-CoV-2. Front Microbiol, 2023. **14**: p. 1073789.
95.   Leeuwendaal, N.K., J.F. Cryan, and H. Schellekens, *Gut peptides and the*
microbiome: focus on ghrelin. Curr Opin Endocrinol Diabetes Obes, 2021. **28**(2):
p. 243-252.
96.   Seydoux, G. and A. Fire, *Soma-germline asymmetry in the distributions of*
embryonic RNAs in Caenorhabditis elegans. Development, 1994. **120**(10): p.
2823-34.
97.   Yang, N.J. and M.J. Hinner, *Getting across the cell membrane: an overview for*
small molecules, peptides, and proteins. Methods in molecular biology (Clifton,
N.J.), 2015. **1266**: p. 29-53.
98.   Meissner, B., et al., *Deletion of the intestinal peptide transporter affects insulin*
and TOR signaling in Caenorhabditis elegans. J Biol Chem, 2004. **279**(35): p.
36739-45.
99.   Rao, M. and S. Sockanathan, *Molecular mechanisms of RNAi: Implications for*
development and disease. Birth Defects Research Part C: Embryo Today:
Reviews, 2005. **75**(1): p. 28-42.
100.   Bernstein, E., et al., *Role for a bidentate ribonuclease in the initiation step of RNA*
interference. Nature, 2001. **409**(6818): p. 363-6.
101.   Kass, J., et al., *The EGL-3 proprotein convertase regulates mechanosensory*
responses of Caenorhabditis elegans. J Neurosci, 2001. **21**(23): p. 9265-72.
102.   Li, C. and K. Kim, *Neuropeptides*, in *WormBook: The Online Review of C.*
elegans Biology. 2018.
103.   Husson, S.J., et al., *Defective processing of neuropeptide precursors in*
Caenorhabditis elegans lacking proprotein convertase 2 (KPC-2/EGL-3): mutant
analysis by mass spectrometry. J Neurochem, 2006. **98**(6): p. 1999-2012.
104.   Mains, R. and B. Eipper, *The Neuropeptides*, in *Basic Neurochemistry:*
Molecular, Cellular and Medical Aspects., G. Siegel, et al., Editors. 1999,
Lippincott-Raven: Philadelphia.

1171