1

Mechanotransduction and Biophysical Regulation of Fibroblast Plasticity Under Prolonged Shear Stress

Neha Paddillaya[1,2], Anahita Jayaram[3], Paturu Kondaiah[4], Namrata Gundiah[2*]

[1]Department of Bioengineering, Indian Institute of Science, Bangalore.

[2]Department of Mechanical Engineering, Indian Institute of Science, Bangalore.

[3]Department of Life Sciences, School of Natural Sciences, Shiv Nadar Institution of Eminence, Delhi NCR.

[4]Sri Shankara Cancer Hospital and Research Centre, Bangalore

* For correspondence:

Namrata Gundiah,

Biomechanics Laboratory,

Mechanical Engineering Department,

Indian Institute of Science, Bangalore.

Email: namrata@iisc.ac.in; ngundiah@gmail.com

SUMMARY

Complex interactions between mechanical stimuli and biochemical signaling shape the
evolving fibrotic milieu. Fibroblasts contribute to this process, transitioning into
myofibroblasts on stiffer substrates, and under low shear in 3D environments. The time-
dependent effects of prolonged physiological shear stress on fibroblasts however remain
poorly understood. We demonstrate that sustained shear stress (1.2 Pa) over 12 hours
induces a temporal cascade of signaling and biophysical changes, driving fibroblast trans-
differentiation into myofibroblasts. Cells adopt a spindle-shaped morphology under shear,
actively remodeling actin and α-SMA to form elongated stress fibers. Myofibroblasts exhibit
increased vinculin and zyxin expression in mature adhesions that remain elevated post-
shear. Traction stresses progressively increased over the shear duration and persisted for
12 hours after removal of the stimulus, indicating mechanical memory. Post-shear,
deadhesion strengths increased significantly, accompanied by shifts in gene expressions
governing extracellular matrix homeostasis. Prolonged shear enhanced MSD, directionality,
and migration speeds, crucial for effective tissue repair. Finally, we demonstrate temporal
changes in the biochemical gene expressions mediating the shear-induced remodeling of
the extracellular matrix, cytoskeleton, and focal adhesions during the myofibroblast
transition. These findings provide insights into the mechanobiological factors driving
fibroblast-to-myofibroblast transitions which may inform the development of mechano-
therapeutic strategies.

KEYWORDS

24 Myofibroblast, mechanobiology, fibrosis, traction stress, directional migration, deadhesion

25 strength.

1 INTRODUCTION

Fibrotic scar formation is driven by the accumulation and activation of fibroblasts into

large fusiform-shaped myofibroblasts through a process of fibrogenesis[1,2]. Fibroblasts are

highly migratory following activation, demonstrate upregulation of proliferative and cytokine

markers, and increased production of extracellular matrix (ECM) proteins and matrix

metalloproteinases (MMPs)[3]. Synergistic interactions between the mechanical stimuli and

biochemical signaling are highly relevant in fibrosis due to ischemic cardiomyocyte death,

lung injury, cancers, or wound healing. For example, infiltrating fibroblasts in an ovine

aneurysm model of the myocardial tissue replace injured cardiomyocytes with randomly

oriented collagen, disrupting electrical activity, leading to ventricular arrhythmias (Figure 1A,

1B)[4,5]. The injured tissue subsequently undergoes significant remodeling due to collagen

crosslinking and tissue compaction during the proliferative stage of fibrosis, resulting in

increased stiffness[6–9]. Higher tissue stiffness, a hallmark of fibrotic and chronic inflammatory

disease, further activates quiescent fibroblasts into myofibroblasts. This activation alters

cellular mechanotransduction, reinforcing and amplifying the fibrotic phenotype (Figure 1C).

Mechanical forces are increasingly recognized as potent modulators of cellular

behavior, influencing morphology, gene expression, and differentiation[10]. This recognition

has led to a growing interest in understanding how mechanobiological cues contribute to the

initiation and progression of fibrosis. Excessive dynamic stretching stimulates myofibroblast

activation through the development of contractile actomyosin bundles, α-SMA expression,

and stretch-dependent signaling[11]. Myofibroblasts have well-defined adhesions and higher

expression of intracellular gap junctions[12]. Substrate stiffness also plays an important role in

the trans-differentiation of fibroblasts into myofibroblasts, also called fibroblast-to-

myofibroblast transition (FMT)[13,14]. Myofibroblast differentiation is mediated by TGF-β activity

through its binding to receptors on the cell membrane; this activity is elevated on stiffer

substrates and correlates with $\alpha 5$ integrin expression[13–15]. Recent studies show that cues

from the three-dimensional collagen network structure regulate myofibroblast differentiation

under static conditions[16]. Fluid shear stress (FSS) can also induce FMT: fibroblasts on 2D

substrates elongate and align in the direction of shear, and differentiate into myofibroblasts

under shear[17,18]. Low fluid shear (0.1-0.3 dyn/cm²) in 3D matrices induces alignment of the

underlying collagen fibers and facilitates FMT through the upregulation of TGF-β1[20].

Fibroblast migration is enhanced through the interplay between shear stress and

biochemical signaling pathways, including TGF-β, Rho/ROCK, FAK, and YAP, which

promote cytoskeletal reorganization and drive the formation of contractile stress fibers[21,22].

Despite advances in understanding FMT, the molecular mechanisms driving these

processes due to physiological FSS remain poorly understood[23]. Significant gaps also

remain in our understanding of how temporal sequences in mechanotransduction pathways

driving FMT are initiated and orchestrated under prolonged fluid flows. Finally, there is little

evidence to show if sheared cells retain a 'mechanical memory' of the transition once the

shear stimulus is removed.

Our study aims to address these questions using a custom microscope-mounted fluid

shear device (Figure 1D).[24–26] The instrument enables precise control of the magnitude and

duration of shear stress, permits real-time cell visualization, and provides a physiologically

relevant milieu to identify mechanotransduction pathways during FMT. We applied laminar

shear stress (1.2 Pa) to NIH3T3 fibroblasts and analyzed the temporal changes in traction

stresses, migration, adhesion strength, and gene expressions during shear over 12 hours,

and recovery post-shear for 12 hours. We demonstrate myofibroblast transitions induced

due to prolonged physiologically relevant FSS, marked by increased gene expressions

linked to focal adhesion, cytoskeletal, and ECM remodeling. Transformed cells exerted

higher tractions and had enhanced adhesion strengths which persisted 12 hours post-shear.

The interplay between shear stress and biochemical pathways promotes cytoskeletal

reorganization and drives the formation of contractile stress fibers, enhancing fibroblast

migrations. These results provide insights into fibrotic mechanisms, highlighting potential

pathways for therapeutic intervention.

RESULTS

Time-dependent adaption of cytoskeletal and focal adhesions to shear stress

We hypothesized that prolonged FSS induces stress fiber and focal adhesion

remodeling in fibroblasts. To study the effects of prolonged shear on cytoskeletal remodeling,

we subjected NIH3T3 fibroblasts to a constant shear stress of 1.2 Pa at 20 minutes, 1 hour,

3 hours, 6 hours, and 12 hours using a custom microscope-mounted fluid shear device

(Figure 1C; see Materials and Methods). The device features a 1° cone that rotates above a

cell-seeded substrate using a hard drive motor, controlled with an electronic speed

controller, generating a broad annular region of laminar shear stress (Figure 1D). This region

of uniform FSS, extending 2 to 12 mm away from the cone tip, allows simultaneous

assessment of a large number of cells in a single experiment over an extended duration. We

stopped the shear stress after 12 hours (Shear group) and monitored cells over 12 hours in

the absence of FSS (Recovery group). Cells were fixed at various time points, and stained

with rhodamine-phalloidin (red) to visualize actin, DAPI (blue) for the nucleus, and antibodies

against zyxin (green) and vinculin (cyan) to identify mature and dynamic focal adhesions

(Figure 2A). We used these images to assess time dependent changes in cell areas, aspect

ratios, and stress fiber lengths due to prolonged FSS and post-shear, and during recovery

(Figure 2B, C, D; see Materials and Methods).

Our data demonstrate distinct time-dependent changes in cell morphologies during

FSS application (Figure 2A). Following shear, cells show prominent stress fiber with zyxin

and vinculin localization. Cell areas gradually increased under FSS (Figure 2B), with no

significant differences at 20 minutes compared to Control (no shear). However, the aspect

ratios initially decreased before increasing steadily over time (Figure 2C). These results

suggest that a minimum stress duration is required to induce observable changes (Figure

S1). By 1 hour, cell areas increased significantly, indicating fibroblast remodeling, alignment,

and elongation in the shear direction ($p < 0.05$). Aspect ratios and cell areas plateaued

between 6 and 12 hours of FSS ($p < 0.05$) and remained unchanged in the Recovery group.

The stress fiber lengths also increased in a time-dependent manner, peaking to

89.12 ± 8.04 μm at 12 hours before marginally decreasing during recovery (Figure 2D;

see Materials and Methods). These findings suggest that prolonged FSS exposure induces cytoskeletal modifications that drive morphological changes.

We quantified changes in actin, zyxin, and vinculin intensities from the immunofluorescence images of cells (n=15) in the Control, Shear (12 hours), and Recovery (12 hours post shear) groups, respectively (Figure 2E-G; see Materials and Methods). Actin intensity was significantly higher in the Shear group compared to Control ($p<0.05$), indicating extensive cytoskeletal remodeling under FSS. These changes persisted 12 hours after the removal of the fluid stimulus. Vinculin and zyxin intensities, markers of mature and dynamic focal adhesions, also increased with shear duration, suggesting progressive strengthening of mature focal adhesions under fluid shear. The recovery group showed reduced vinculin intensity, compared to zyxin-positive regions that were unchanged after the removal of FSS (Figure 2F, G). The distribution of focal adhesion areas, fit to a log-normal model, showed smaller adhesions in the Control group compared to the Shear and Recovery groups that had larger focal adhesions (Figure 2H). The observed focal adhesion growth and increased stress fiber lengths under prolonged FSS indicate significant changes in biophysical properties that remain elevated for 12 hours post-shear.

Prolonged shear induces FMT, and enhances cell contractility and adhesion strength

How does stress fiber and focal adhesion remodeling alter the biophysical properties of cells? To investigate this, we seeded fibroblasts on bead-coated polyacrylamide substrates (10 kPa) and applied 1.2 Pa shear using the custom microscope-mounted device (see Materials and Methods). Figure 3A shows representative brightfield images alongside corresponding traction maps, calculated using a regularized Fourier Transform Traction Cytometry (Reg-FTTC) method[27], from the Control, Shear (12 hours), and Recovery (12 hours) groups, respectively. These data provide the first traction measurements of cells subjected to prolonged elevated FSS.

Maximum tractions and total strain energy densities increased steadily under shear (Figure 3B, 3C), with higher tractions localized at the polarized cell ends. Tractions

increased significantly after ~ 1 hour of shear ($p<0.05$) but showed no further differences between the 6 hours and 12 hours Shear groups ($p<0.05$). Surprisingly, tractions increased significantly and remained unchanged after the removal of the shear stimulus, indicating that increased contractile activity persisted during recovery ($p<0.05$). Corresponding strain energies also increased with prolonged FSS. Figure 3C shows a marginal increase from 20 minutes to 1 hour, followed by a pronounced increase between 3 and 12 hours. However, these values remained elevated post-shear in the Recovery group. Stable focal adhesions (Figure 2A) and stress fibers (Figure 2D) support a highly contractile state, which is associated with increased α-SMA expression — an indicator of the contractile phenotype in fibroblasts. α-SMA was significantly elevated under FSS and remained unchanged after shear removal ($p<0.05$; Figures 3D, E). Sustained tractions and cytoskeletal changes suggest retention of "mechanical memory" and a potential shift towards a myofibroblast-like phenotype, which may influence cellular remodeling and fibrosis progression.

We hypothesized that prolonged shear alters the adhesion strengths of transformed cells. To test this, we measured the critical adhesion strengths of cells in the Control, Shear, and Recovery groups using the device on fibronectin-coated substrates (see Materials and Methods). The FSS was steadily ramped from 1.2 Pa in increments of 0.2 Pa/ minute in these experiments until all cells detached from the substrate (see Materials and Methods section)[24,26]. The resulting sigmoidal graph (Figure 3F) shows the fraction of cells remaining on the substrate with each step shear increase. We quantified the critical deadhesion strengths (τ_{50}) using these data and obtained the stresses required to detach weakly adherent (τ_{20}) and strongly adherent cells (τ_{90}) from the substrate. The rightward shift in deadhesion curves post-shear indicates increased adhesion strength. Interestingly, significant deviations in the initial part of the deadhesion curve for the Recovery group suggest that ~15% of cells retained highly dynamic adhesions, similar to the Control group. In contrast, the Shear group exhibited a leftward shift relative to Control, suggesting weaker and more dynamic adhesions (Figure 3G).

We fit the deadhesion curves to a cell detachment model incorporating

experimentally obtained fibroblast area distributions (Equation 1) [24,26]. The model simulates

stochastic binding kinetics of integrin-mediated slip bonds, distributed around the periphery

of hemispherical cells. Figure 3G shows that the model effectively captures these

experimental results. However, the Recovery group exhibited significant deviations from the

model, particularly at τ_{10} and τ_{90} indicating high adhesive heterogeneity in the cell population

post-recovery. This heterogeneity likely arises from variations in myofibroblast transitions

and potential reversals[28]. These deadhesion dynamics highlight the complex interplay

between adhesion strength and cytoskeletal adaptation to prolonged shear stress.

FSS induces changes in gene expressions for cytoskeletal, focal adhesions, and ECM

proteins

To explore the signaling pathways driving fibroblast adaptation to prolonged FSS, we

analyzed mRNA levels of key genes in extracellular matrix (ECM) remodeling, focal

adhesion dynamics, and the cytoskeletal regulation across Control, Shear, and Recovery

groups. We selected primers targeting genes associated with actin dynamics (Actin, RhoA),

ECM (Collagen-1, Collagen-3, Fibronectin, MMP2, MMP9), focal adhesions (Vinculin, Talin,

FAK, IGTAV, IGTA5, IGTB1, IGTB3), and α-SMA, a myofibroblast differentiation marker (see

Materials and Methods). Gene expressions were normalized to GAPDH levels and the

Control group.

Figure 4A shows distinct time-dependent changes in gene expression profiles

corresponding to the ECM, triggered by prolonged FSS. Fibronectin expression significantly

increased, peaking between 3 and 12 hours of FSS, before reducing post-shear. Collagen-1

and Collagen-3 — key structural components of the ECM — were upregulated during shear

and in the recovery phase. Matrix metalloproteinases corresponding to MMP2 and MMP9

displayed distinct changes under shear: MMP2 levels remained elevated from 3 hours of

FSS through the post-shear phase, whereas MMP9 levels were higher post-shear. Focal

adhesion-related genes for Vinculin and Talin were minimally upregulated at 3 and 6 hours of

shear (Figure 4B), aligning with observations on maximal cell elongation and flow-induced alignment. Integrins IGTAV and IGTA5 showed a moderate increase during shear exposure but were significantly upregulated during recovery. IGTB3 exhibited a steady and gradual increase through the shear exposure, suggesting its involvement in stabilizing adhesions under prolonged FSS.

Shear stress significantly upregulated RhoA and FAK expressions by 3 hours (Figure 4C), with elevated levels persisting over 12 hours post-shear. Increased RhoA expression suggests active cytoskeletal reorganization under shear, consistent with the observed increases in traction stress and stress fiber formation. Similarly, FAK, a central mechanotransduction regulator, was elevated alongside actin and α-SMA, both of which increased steadily under shear. These data indicate cytoskeletal strengthening and myofibroblast transition which persisted during recovery, indicating the transformation of fibroblasts into myofibroblasts that have stable adhesions and prioritize ECM deposition. Our results highlight the complex regulatory mechanisms governing fibroblast adaptation to shear stress, emphasizing immediate and sustained signaling changes that drive fibroblast transformation under prolonged fluid shear.

Migration and directionality of fibroblasts

Fibroblast migration is essential in early fibrosis for tissue repair. In contrast, myofibroblasts are characterized by increased contractility, and ECM deposition to remodel tissues. We hypothesized that prolonged FSS enhances directed fibroblast migration but reduces myofibroblast motility. Figure 5A-C shows rose plots of migration behaviors in Control, Shear (12 hours), and Recovery (12 hours) conditions. Under FSS, fibroblasts migrated along their oriented directions (Movies 1-3). In contrast, cells in the Recovery group surprisingly exhibited minimal directional changes, migrating along straight paths (Figure 5C). Mean squared displacement (MSD) was significantly higher in the Shear group (Figure 5D), driven by cytoskeletal remodeling and dynamic focal adhesion formation under FSS (see Materials and Methods). In contrast, MSD values were significantly lower in the

Recovery group, suggesting stable adhesions, as myofibroblasts prioritize ECM deposition over migration. Thus, FSS enhances fibroblast mechanoadaptability, while myofibroblasts reinforce adhesion stabilization, shifting to a less migratory phenotype. The effective diffusivity (α), derived from the slope of the log(MSD) with log(time) data (Figure S3), was 0.81 in the Control group, indicating sub-diffusive migration without directional cues. In contrast, α increased to 1.26 in the Shear group, reflecting super-diffusive migration with persistent flow-aligned movement. The Recovery group had a lower α (=0.59), signifying constrained, sub-diffusive migration with loss of directional ability.

Directionality Ratios (DR) were significantly higher in the Shear group than in the Recovery group at 2 hours, indicating sustained directional persistence along specific directions post-shear (p<0.05; Figure 5E). However, myofibroblasts had significantly lower DR values within ~60 minutes post-shear. Migration speeds also differed significantly between all groups (p<0.05; Figure 5F). These differences highlight the role of FSS on cytoskeletal dynamics and adhesion strengths in fibroblasts, whereas myofibroblasts retained their mechanical memory of the transformation. Prolonged shear stress hence induces fibroblast-to-myofibroblast transformation, leading to stabilized adhesions, reduced motility, and elevated contractility (Figure 3B). These results hence show fibroblast mechano-adaptivity as a critical factor in tissue repair, remodeling, and the progression of fibrosis.

DISCUSSION

Fibroblasts are of mesenchymal origin and regulate the balance between ECM synthesis and breakdown to maintain tissue homeostasis[29]. Disruption of this fine balance in fibrotic disease stimulates the differentiation of fibroblasts into myofibroblasts due to chemokine signaling and mechanical activation[23]. While initially protective, FMT leads to excessive ECM deposition, tissue stiffening, and impaired organ function in fibrotic disease[30,31]. We investigated fibroblast adaptation to prolonged physiological shear stress (1.2 Pa), and its subsequent recovery post-shear, and uncovered key mechanisms in the

complex and nonlinear coupling between biochemical signaling and biophysical changes

driving FMT. Our study highlights four key findings: first, fibroblasts adopt an elongated

spindle-shaped myofibroblast morphology under shear that is linked to increased stress fiber

lengths and altered distributions of vinculin and zyxin along the cell periphery. These

changes remained unchanged 12 hours post-shear, emphasizing the presence of cellular

mechanical memory. Second, FSS induced cytoskeletal remodeling increases cell tractions

in myofibroblasts, and upregulates actin and α-SMA expressions, sustaining the contractile

phenotype post-shear. Third, prolonged FSS reduced the critical deadhesion strengths, and

induced significant, time-dependent changes in ECM-related genes (FIBRONECTIN, Col1,

Col3, MMP2, and MMP9) and cytoskeletal remodeling (Actin, α-SMA, RHO), indicative of

the myofibroblast transition[32,33]. Upregulation in RhoA, FAK, actin, and α-SMA expressions

due to shear highlight FMT-associated mechanotransduction in fibroblasts. Finally,

prolonged shear enhances fibroblast migrations (MSD, directionality, and speed),

demonstrating distinct mechanical adaptations that are crucial for effective tissue repair in

fibrosis.

The disruption of "function dictates form" provides a powerful framework for

understanding how tissues adapt to forces that drive disease progression. Polarized,

spindle-shaped fibroblasts aligned with flow due to increased stress fiber lengths[34]. This

cytoskeletal reorganization, mediated by actin and α-SMA, is accompanied by changes in

vinculin and zyxin expression. Shear-induced upregulation of α-SMA, a hallmark of FMT,

begins ~6 hours of shear exposure and persists post-shear. The accompanying focal

adhesion growth, reorganization, and reinforcement further support the transition[35,36].

Myofibroblasts have microfilament bundles that terminate at specialized mature adhesion

complexes, facilitating ECM attachment and contractility[23,29,37]. Stress fibers contribute to the

formation of mature adhesions, enabling cells to resist FSS by generating tractions[10,20,31].

Focal adhesion reinforcement and stress fiber formation are both critical in maintaining the

contractile myofibroblast phenotype[38,39]. Despite marginal decreases in stress fiber lengths

and vinculin expression post-shear, myofibroblasts retained the ability to generate elevated

tractions. Increased zyxin in fibroblasts (Figure 2G, 2H) further supports its role in

strengthening and stabilizing the cellular architecture[36,40,41].

Shear induced upregulation of RhoA and FAK regulates cytoskeletal dynamics

through biochemical signaling, enhancing cellular contractility and altering the tensional

homeostasis[33 42]. Elevated expression of ECM-related genes, such as fibronectin, collagen-

1/3, and matrix metalloproteinases (MMP2 and MMP9), in the Shear and Recovery groups

indicates active tissue remodeling, a hallmark of FMT[18]. MMP2 regulates ECM degradation

and remodeling[43], whereas MMP9 helps breakdown the ECM under stretch[44]. Notably,

MMP2 expression remained elevated through shear exposure and recovery, whereas MMP9

showed a biphasic response, peaking at 3 and 12 hours. These findings align with gene

expression changes observed during tissue remodeling in myocardial infarction, lung

fibrosis, and wound healing[29]. Focal adhesion remodeling and stress fiber reinforcement

during shear and post-shear altered the migration dynamics of fibroblasts. Directionality

ratios and migration speeds significantly increased in the Shear group, which exhibited sub-

diffusive migration compared to super-diffusive behaviors of Control cells (Figure 5D-F).

These results agree with reports that show fibroblasts lack directionality under static

conditions and have persistent anomalous migration[17]. Prolonged shear induced cellular

polarization, with cells predominantly migrating along the shear direction in our study.

There are some limitations of this study: first, collagen fibers align under shear,

creating tracks that facilitate cell elongation and migrations along the fiber direction. Figure

S4 shows fluorescence images of collagen-functionalized coverslips subjected to uniform

shear for 30 minutes in the device, demonstrating fiber alignment that may aid the

realignment responses of fibroblasts. However, isolating the individual contributions of shear

and collagen alignment remains challenging. Second, the observed mechanical memory of

cells at 12 hours post-shear may not persist over longer durations, warranting further

investigation. Third, the media used to cause fluid shear on cells may contain released

soluble factors that contribute to FMT. We did not however assay the media for the presence

of such chemicals. Finally, we did not investigate potential changes in proliferation and

apoptotic markers under shear. Understanding the mechanotransductive processes driving FMT is crucial for developing therapeutic strategies to mitigate fibrotic diseases, where excessive myofibroblast activity leads to pathological tissue remodeling and stiffening. Targeting myofibroblast mechanoadaptation offers a promising avenue for the management and treatment of fibrosis.

ACKNOWLEDGEMENTS

We acknowledge the Bioimaging Facility, Indian Institute of Science (IISc) for help with the use of the Andor Dragonfly 502 spinning disk confocal microscope. NG is grateful to DBT (BT/BR23724/BRB/10/1606/2017) and SERB (SPG/2021/002425) for project support. Protein quantification equipment in the Biomechanics lab was acquired through a generous grant from DST-FIST. We also thank Prof. Pramod Pullarkat (RRI, India) and Prof. Gautam Menon (Ashoka University, India) for useful discussions during the early stages of this work.

Author contributions

NP: Methodology, Software, Validation, Formal Analysis, Investigation, Writing – Original Draft, Data Visualization. AJ: Methodology, Software, and Validation of migration result. PK: Conceptualization, Methodology, Formal Analysis, Writing – Reviewing and Editing, Supervision. NG: Conceptualization, Methodology, Software, Formal Analysis, Resources, Data Curation, Writing – Original draft, Reviewing and Editing, Data Visualization, Supervision, Project Administration and Funding Acquisition.

Declaration of interests

The authors declare no competing interests.

Figure 1

Figure 1: Stress-Induced Fibroblast to Myofibroblast Transition (FMT). (A) Representative

histological sections of ovine myocardial tissue at 2 and 8 months post-infarction show

significant changes in the mid-wall region[4]. Green arrows indicate remodeled fibrotic tissue

characterized by a high density of fibroblasts. (B) Schematic shows the biophysical,

morphological, and biochemical drivers underlying FMT. (C) A custom microscope-mounted

fluid shear device was used to apply uniform FSS on fibroblasts over a prolonged duration.

(D) Flow visualization studies demonstrate a large annual region of laminar shear that

extends from 2 -12 mm from the 1° cone tip.

Figure 2: Time-Dependent Responses to Prolonged Shear Exposure. (A) Representative

immunofluorescence images of NIH3T3 fibroblast cells from Control (no shear), 6 and 12

hours shear, and 12 hours post-shear (Recovery) groups. Cells were stained for actin (red),

nucleus (blue), zyxin (green), and vinculin (cyan). Scale bar: 50µm (B) Variations in cell

areas (n=50), (C) Aspect ratios (n=50), and (D) Stress fiber lengths were measured at each

time point during shear and post-shear (n=15). Fluorescence intensities were quantified and

compared for (E) Actin, (F) Zyxin, and (G) Vinculin in the focal adhesions for Control, (12

hours) Shear, and (12 hours post-shear) Recovery groups (n=15 each group). Differences

between the non-significant groups alone are labeled; all other group comparisons show

statistically significant differences at $p<0.05$. (H) Probability density plots with log-normal fits

are shown for projected zyxin areas (µm²) from the images.

Figure 3: Cell Contractility and Adhesion Dynamics under FSS and post-shear. (A)

Representative brightfield images and corresponding traction stresses (Pa) are shown from

Control, Shear, and Recovery groups measured using the regularized Fourier Transform

Traction Cytometry (Reg-FTTC) method[27]. Scale bar: 50μm. (B) Maximum tractions

increased from 60 to 360 minutes during shear, and did not change after removing the shear

(n=15). (C) Strain energy increased with time due to shear and remained relatively

unchanged post-shear (n=15). (D) Immunofluorescence images of α-SMA (purple)

demonstrate activation of mechanotransduction pathways. Scale bar 50μm. (E)

Quantification of α-SMA expression across Control, Shear, and Recovery conditions with

fluorescence intensity (A.U.). (F) The de-adhesion strengths of cells are reported for cells in

Control (n=370), Shear (n=366), and Recovery (n=375) groups. Normalized cell number

followed sigmoidal behaviors (G) Clear differences are visible between the weakly adherent

(τ_{20}) shear stresses, critical deadhesion strengths (τ_{50}), and strongly adherent (τ_{90}) shear

stresses. Non-significant differences between the groups are indicated using n.s. All other

groups are statistically significant ($p<0.05$).

12 **Figure 4:** Gene Expression Changes under Prolonged Shear. Fold changes in gene

13 expression levels ($2^{-\Delta\Delta Ct}$) are shown during a steady increase in FSS and recovery phases.

14 Primers corresponding to (A) key extracellular matrix (ECM) constituents (Fibronectin,

15 Collagen I (Col1), Collagen III (Col3), MMP2, and MMP9). (B) Focal adhesion markers

including Talin, Vinculin, ITGA5, ITGAV, ITGB1, and ITGB3. (C) Signaling molecules (FAK

and RhoA) and (D) myofibroblast markers (Actin and α-SMA) show clear changes.

Figure 5: Cell Migration Dynamics under FSS and during Recovery. Migration tracks of cells

from (A) Control (n=87), (B) Shear (n=90), and (C) Recovery after removal of shear stimulus

(n=98) illustrate displacements over 2 hours that have different dynamics. (D) Mean squared

displacement (MSD) highlights differences in migrations between groups. (E) Directionality

ratios indicate changes in the orientation of cell migration. (F) Cell migration speeds

(µm/min) were significantly different in comparisons between the Control, Shear, and

Recovery groups (p<0.05).

Supplementary Figure: S1

Representative actin and DAPI stained images of fibroblasts at various points during shear

and recovery. Shear direction bottom to top. Scale 20um.

Supplementary Figure: S2

Brightfield and traction maps of representative cells at various points during shear and

recovery. Tractions, calculated using a regularized FTTC method, show higher values at the

polarized ends of the cell for cells under shear. Scale 20 µm.

3 **Supplementary Figure S3:** Mean Squared Displacement (MSD) variations with time are

4 shown in log-log plots for cells in Control (blue), Shear (orange), and Recovery (gray)

5 groups. The slopes to these data, α, are the effective diffusivity values that were used to

6 compare with various diffusive behaviors. R^2 values show the goodness of fit to the data.

1 **Supplementary Figure S4:** Collagen alignment on substrates under 1.2 Pa shear was

2 determined using 40 µg/ mL collagen I petridishes. Images are shown 12 hours post-shear.

3 Scale bar: 20 µm.

Supplementary videos:

Movie 1: Control cell migrations on fibronectin coated substrates show persistent

random migrations

Time-lapse video shows cell migrations in the Control group over 2 hours. Cells exhibit

active random migrations movement over time.

Movie 2: Sheared cells have directed migration on fibronectin-coated glass substrates

Time-lapse video shows fibroblast migrations over 2 hours following a 12 hour exposure to

1.2 Pa FSS. Cells have increased alignment and directional persistence along the shear

direction.

1   **Movie 3: Removal of FSS showed migrations along straight trajectories over 2 hour**

2   **period**

5   Transformed fibroblasts following 12 hours shear exposure were maintained for 12 hours

6   without shear, and migrations were quantified over 2 hours. Cells display reduced migration

7   speeds and move primarily along one direction.

MATERIALS AND METHODS

Cell Culture

NIH-3T3 mouse embryonic fibroblast cells at early passage numbers were

maintained in Dulbecco's Modified Eagle Medium (DMEM, Invitrogen) supplemented with

10% (v/v) fetal bovine serum (FBS, Invitrogen) and 1% (v/v) penicillin/streptomycin (Sigma-

Aldrich) in a humidified incubator at 37°C with 5% CO_2. Cells were passaged every 2–3 days

and harvested by washing thrice with phosphate-buffered saline (PBS) and detached from

the flasks using 0.25% trypsin-EDTA (Invitrogen) for experiments. Fibronectin (40 µg/ml)

was added to a coverslip at 37 °C for 1 hour in a humid chamber and washed thrice with

PBS for experiments. Cells were seeded at 5000 cells/ml on fibronectin coated coverslips

and allowed to attach and spread for 6 hours before starting the shear experiments.

Fluid shear studies using a custom microscope-mounted device

A custom fluid shear device was used for all studies (Figure 1D). The device was

placed on a live-cell epifluorescence microscope (Leica DMI 6000) equipped with a PECON

cage incubator for live-cell imaging during shear. Details of the device are provided in our

previous work[24,26]. Briefly, a 1° cone was rotated using a hard drive motor powered by a DC

supply and controlled *via* an Arduino-based electronic speed controller to modulate pulse

width and frequency (Figure 1D). Shear stresses, calculated using equations for Couette

flows generated in the device, are directly proportional to the angular speed of the motor.

The constant rotation of the cone produces a large annulus of laminar shear stress (Figure

1D). Flow visualization experiments, performed using fluorescently labeled beads (2 µm

size; 2 µgm/ml), were used to identify the regions of laminar flow in the device. An annular

laminar flow region of 2-12 mm, located away from the cone tip and ends of the Petri plate,

was used in subsequent analyses.

Cells were stained with Hoechst (Invitrogen) for 3 minutes in an incubator maintained

at 37 °C and 5% CO_2 and plated on fibronectin coated coverslips in the device. Uniform

1.2 Pa FSS was exerted on the cells and the experiments were stopped at various time

points during the application of shear for cell morphometry studies and to quantify changes

in the gene expressions. These intervals included 20 minutes, 1, 3, 6, and 12 hours. We also

quantified changes in the critical deadhesion strength of fibroblasts based on earlier

protocols[24,26] by ramping the shear stress by 0.2 Pa per minute using a custom Arduino

code. Images acquired using a CCD camera at each stress increment were used to count

the number of cells on the substrate based on the fluorescence of the Hoechst-stained

nuclei. The ratio of cells remaining (N/N_0) on the substrate showed sigmoidal dependence

with shear stress. The critical adhesion strength (τ_{50}) was defined as the shear stress at

which 50% of the cells remained adhered to the substrate. Shear stresses based on a 20%

cell fraction removed from the substrate, τ_{20}, was calculated to estimate weakly adherent

cells. τ_{90} value of shear stress was used to characterize strongly attached cells based on the

removal of 90% of cells from the substrate[24].

Experimental sigmoidal curves were fit to a cell detachment model based on the

statistical distribution of cell areas and adhesion strengths[24]. The cell was approximated as a

hemispherical shape which was attached to a substrate using a uniform distribution of focal

adhesions along the cell perimeter. This model considers heterogeneity in cell size and

adhesion strength to calculate the fraction of cells adherent on the substrate at each step

increase in shear stress due to Stoke's flow in a low Reynold's number moving fluid. Bell's

deterministic equations were used to model the molecular clusters as slip bonds which are

stable with an infinite lifetime until they experience a critical load. The fraction of cells

remaining on the substrate, $\Phi(\tau)$, is given as:

and
$$\Phi(\tau) = \frac{1}{2}\left[1 - \mathrm{erf}\left(\frac{\ln\left(\frac{\tau}{\tau_{50}}\right)}{\sqrt{2}s}\right)\right]$$
(1)

In this expression, the parameter s is proportional to the distribution width of attachment

points for cells, as determined from experimental data of cell spread areas. The term τ_{50}

represents the critical shear stress required to detach 50% of the cells from the substrate,

and $\Phi(\tau)$ denotes the fraction of adherent cells at a given shear stress τ. Data from

experiments were fit to the model using MATLAB (v R2019b, The Mathworks Inc. Natick,

MA).

Traction Force Microscopy (TFM)

We assessed changes in cell contractility under shear stress using traction force

microscopy (TFM), described in our earlier work[27]. Briefly, glass coverslips were cleaned and

prepared with polyacrylamide (PA) gels of 10 kPa stiffness using a mixture of acrylamide and

bisacrylamide in distilled water, combined with 10% APS and TEMED. An optimized

fluorescent bead density on top of the gel surface allowed the use of an epifluorescence

microscope for the traction studies. The PA gels were functionalized with sulpho-SANPAH

and coated with fibronectin (40 μg/ml) for cell attachment. NIH3T3 cells were seeded onto

the gels at 2000 cells/ml and allowed to attach for 6 hours in an incubator at 37 °C with 5%

CO_2. The substrate with cells on PA hydrogel was placed in the custom device and

subjected to 1.2 Pa shear stress for various durations (20 minutes, 1, 3, 6, 12 hours) and

during recovery (n=15 each group). Phase-contrast/ epifluorescent images of the cells and

beads were acquired using a Leica DMI6000B live-cell microscope equipped with a PECON

stage to obtain the cell boundary and deformed images of beads on the hydrogel surface

due to cell attachments. Cells were removed using trypsin-EDTA, and the referential bead

positions were acquired. Traction data were analyzed using a custom regularized Fourier

Transform Traction Cytometry (Reg-FTTC) MATLAB code[27].

Migration studies

NIH3T3 cells from Control (no shear), Shear (12 hours), and Recovery (12 hours)

were seeded at 5000 cells/mL on glass substrates and placed in an incubator for 6 hours to

permit cell attachment and spreading. We used an inverted live cell fluorescence

microscope (Leica DMI6000B, Leica Microsystems, Germany) with a 10X Phase Contrast

objective to capture time-lapse images of cells every 15 minutes over a 6 hour period in a

humid temperature-controlled PECON chamber maintained at 37°C and CO_2 levels at 5%. A

motorized stage was used to capture multiple positions (N~10) of each cell in the Control,

Shear, and Recovery groups. Phase contrast time-lapse images were converted to 8-bit and

drift-corrected using the Template Matching plugin in FIJI with bilinear interpolation[45].

Migrations of cells were tracked using a Manual Tracking Plugin after applying a centering

correction. Cells that migrated a distance greater than three cell lengths were included in the

analysis. Migration tracks were aligned to a common point of origin, and the overall

displacement, mean, and instantaneous cell speeds were calculated using the (x,y) positions

of cells at each time instant, t. Mean Square Displacement (MSD) and the directionality ratio

(DR) were calculated using the DiPer code[46] using overlapping time intervals. Based on this

method, the MSD(n), representing the displacement for step size, n, is given by

$$MSD(n) = \frac{1}{N-n+1}\sum_{i=0}^{N-n}\left[\left(x_{(i+n)\Delta t} - x_{i\Delta t}\right)^2 + \left(y_{(i+n)\Delta t} - y_{i\Delta t}\right)^2\right] \qquad (2)$$

where N is the total number of displacements step size n and Δt the minimal time interval

between points. Population averages of MSDs were plotted with time. The directionality ratio

was calculated as

$$\text{Directionality Ratio (DR)} = \langle\frac{d_t}{D_t}\rangle_C \qquad (3)$$

d_t represents the straight-line distance from the initial position of the cell to its position at

time t, providing a direct measure of displacement from the starting point. D_t denotes the

actual path length traveled by the cell from the start to its position at time t, which includes all

turns and deviations. The brackets indicate an average of these values across C cells at

time, t.

RNA Isolation and RT-qPCR

Cells were washed with PBS, and TRIzol® reagent was added to each Petri dish for

RNA extraction using the RNeasy Mini Kit (Qiagen). Two micrograms of RNA were used for

cDNA synthesis with a high-capacity cDNA reverse transcription kit (Applied Biosystems).

Quantitative real-time PCR (qRT-PCR) was conducted using PowerUp SYBR Green master

mix with 10 ng of cDNA as the template. Fold changes were calculated using the $2^{-\Delta\Delta CT}$

method and gene expression was normalized to GAPDH and no-shear controls. Data were

analyzed across three biological replicates for each Control, Shear, and Recovery group in

our study.

Immunofluorescence

Cultured cells were rinsed twice with chilled PBS (Fisher Scientific), fixed with 4%

paraformaldehyde for 15 minutes, permeabilized with 0.1% Triton X-100 for 3 minutes,

and blocked using 5% bovine serum albumin (BSA) with 5% fetal bovine serum (FBS) for

1 hour at room temperature for immunofluorescence staining. Specimens were incubated

for 1 hour at room temperature with primary anti-vinculin antibody (1:500, Sigma Aldrich)

followed by secondary Alexa Fluor 488-conjugated antibody (1:600, Molecular Probes,

Invitrogen), with Alexa Fluor 568-phalloidin (1:200, Invitrogen) to stain F-actin. Samples were

rinsed twice with PBS, nuclei were labeled with DAPI (1:500, Invitrogen) for 2 minutes,

rinsed, and mounted on coverslips with ProLong™ Gold Antifade Mountant (Invitrogen).

Confocal images were acquired using a Leica SP5 confocal microscope. Images were

analyzed using Fiji (ImageJ) to quantify cell spread area after aligning samples along the

shear direction.

Stress fibers were quantified using confocal images of cells stained for actin, and a

custom MATLAB code as described earlier[27]. This method involves segmenting the image to

identify actin fibers using intensity thresholding and morphological operations. The lengths of

individual stress fibers were calculated using the Hough transform of the segmented fibers

from each image. Approximately 5-10 fibers were identified in each cell, and 15 cell images

were analyzed for each group in the study.

Statistical Analysis

Data from experiments performed in three replicates are reported as mean ± SD.

Differences between groups were assessed using one-way analysis of variance (ANOVA)

with Bonferroni post-hoc tests for individual comparisons. A p-value of < 0.05 was

considered statistically significant. Non-significant differences are indicated as n.s.

References

1.   Talman, V. & Ruskoaho, H. Cardiac fibrosis in myocardial infarction-from repair and

remodeling to regeneration. Cell Tissue Res. 365, 563–581 (2016).

2.   Kuhn, C. & McDonald, J. A. The roles of the myofibroblast in idiopathic pulmonary

fibrosis. Ultrastructural and immunohistochemical features of sites of active

extracellular matrix synthesis. Am. J. Pathol. 138, 1257–1265 (1991).

3.   Shaw, T. J. & Martin, P. Wound repair: a showcase for cell plasticity and migration.

Curr. Opin. Cell Biol. 42, 29–37 (2016).

4.   Gundiah, N. Role of elastin and collagen in the passive mechanics of the circulatory

system. University of California, Berkeley. ProQuest, 3146867, (2004).

5.   Babu, A. R., Byju, A. G. & Gundiah, N. Biomechanical properties of human ascending

thoracic aortic dissections. J. Biomech. Eng. 137, (2015).

6.   Gabbiani, G. The myofibroblast in wound healing and fibrocontractive diseases. J.

Pathol. 200, 500–503 (2003).

7.   Liu, G. et al. miR-21 mediates fibrogenic activation of pulmonary fibroblasts and lung

fibrosis. J. Exp. Med. 207, 1589–1597 (2010).

8.   Hinz, B. Formation and function of the myofibroblast during tissue repair. J. Invest.

Dermatol. 127, 526–537 (2007).

9.   Shinde, A. V. & Frangogiannis, N. G. Fibroblasts in myocardial infarction: a role in

inflammation and repair. J. Mol. Cell. Cardiol. 70, 74–82 (2014).

10.   Hinz, B., Celetta, G., Tomasek, J. J., Gabbiani, G. & Chaponnier, C. Alpha-smooth

muscle actin expression upregulates fibroblast contractile activity. Mol. Biol. Cell 12,

2730–2741 (2001).

11.   Wang, J., Chen, H., Seth, A. & McCulloch, C. A. Mechanical force regulation of

myofibroblast differentiation in cardiac fibroblasts. Am. J. Physiol.-Heart Circ. Physiol.

(2003) doi:10.1152/ajpheart.00387.2003.

12. Hinz, B., Pittet, P., Smith-Clerc, J., Chaponnier, C. & Meister, J.-J. Myofibroblast development is characterized by specific cell-cell adherens junctions. Mol. Biol. Cell 15, 4310–4320 (2004).

13. Leask, A. & Abraham, D. J. TGF-beta signaling and the fibrotic response. FASEB J. 18, 816–827 (2004).

14. Verma, B. K., Chatterjee, A., Kondaiah, P. & Gundiah, N. Substrate stiffness modulates TGF-β activation and ECM-associated gene expression in fibroblasts. Bioengineering 10, 998 (2023).

15. Desmoulière, A., Geinoz, A., Gabbiani, F. & Gabbiani, G. Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. J. Cell Biol. 122, 103–111 (1993).

16. Seo, B. R. et al. Collagen microarchitecture mechanically controls myofibroblast differentiation. Proc Natl Acad Sci U S A 117, 11387–11398 (2020).

17. Lei, X. et al. The effect of fluid shear stress on fibroblasts and stem cells on plane and groove topographies. Cell Adhes. Migr. 14, 12–23 (2020).

18. D'Urso, M. & Kurniawan, N. A. Mechanical and physical regulation of fibroblast–myofibroblast transition: From cellular mechanoresponse to tissue pathology. Front. Bioeng. Biotechnol. 8, (2020).

19. Kouzbari, K. et al. Oscillatory shear potentiates latent TGF-β1 activation more than steady shear as demonstrated by a novel force generator. Sci. Rep. 9, 6065 (2019).

20. Ng, C. P., Hinz, B. & Swartz, M. A. Interstitial fluid flow induces myofibroblast differentiation and collagen alignment in vitro. J. Cell Sci. 118, 4731–4739 (2005).

21. Shi, Z.-D. & Tarbell, J. M. Fluid flow mechanotransduction in vascular smooth muscle cells and fibroblasts. Ann. Biomed. Eng. 39, 1608–1619 (2011).

22. Paddillaya, N. et al. Biophysics of cell-substrate interactions under shear. Front. Cell Dev. Biol. 7, 251 (2019).

23. Baum, J. & Duffy, H. S. Fibroblasts and myofibroblasts: what are we talking about? J. Cardiovasc. Pharmacol. 57, 376–379 (2011).

24. Paddillaya, N. et al. Cell adhesion strength and tractions are mechano-diagnostic features of cellular invasiveness. Soft Matter 18, 4378–4388 (2022).

25. Pullarkat, P. A., Vishavkarma, R., Gundiah, N. & Chahare, N. A microscope mountable fluid shear device, IN201641029893A, 2016. pg 275/677.

26. Maan, R., Rani, G., Menon, G. I. & Pullarkat, P. A. Modeling cell-substrate de-adhesion dynamics under fluid shear. Phys. Biol. 15, 046006 (2018).

27. Kulkarni, A. H., Ghosh, P., Seetharaman, A., Kondaiah, P. & Gundiah, N. Traction cytometry: regularization in the Fourier approach and comparisons with finite element method. Soft Matter 14, 4687–4695 (2018).

28. Doyle, A. D., Nazari, S. S. & Yamada, K. M. Cell–extracellular matrix dynamics. Phys. Biol. 19, 021002 (2022).

29. Younesi, F. S., Miller, A. E., Barker, T. H., Rossi, F. M. V. & Hinz, B. Fibroblast and myofibroblast activation in normal tissue repair and fibrosis. Nat. Rev. Mol. Cell Biol. 25, 617–638 (2024).

30. Wynn, T. A. & Ramalingam, T. R. Mechanisms of fibrosis: therapeutic translation for fibrotic disease. Nat. Med. 18, 1028–1040 (2012).

31. Hinz, B. The myofibroblast: paradigm for a mechanically active cell. J. Biomech. 43, 146–155 (2010).

32. Chiquet, M., Renedo, A. S., Huber, F. & Flück, M. How do fibroblasts translate mechanical signals into changes in extracellular matrix production? Matrix Biol. J. Int. Soc. Matrix Biol. 22, 73–80 (2003).

33. Chiquet, M., Gelman, L., Lutz, R. & Maier, S. From mechanotransduction to extracellular matrix gene expression in fibroblasts. Biochim. Biophys. Acta 1793, 911–920 (2009).

34. Balaban, N. Q. et al. Force and focal adhesion assembly: a close relationship studied using elastic micropatterned substrates. Nat. Cell Biol. 3, 466–472 (2001).

35. Riveline, D. et al. Focal contacts as mechanosensors: externally applied local mechanical force induces growth of focal contacts by an mDia1-dependent and ROCK-independent mechanism. J. Cell Biol. 153, 1175–1186 (2001).

36. Dan, L., Chua, C.-K. & Leong, K.-F. Fibroblast response to interstitial flow: A state-of-the-art review. Biotechnol. Bioeng. 107, 1–10 (2010).

37. Balestrini, J. L., Chaudhry, S., Sarrazy, V., Koehler, A. & Hinz, B. The mechanical memory of lung myofibroblasts. Integr. Biol. Quant. Biosci. Nano Macro 4, 410–421 (2012).

38. Hoffman, L. M., Jensen, C. C., Chaturvedi, A., Yoshigi, M. & Beckerle, M. C. Stretch-induced actin remodeling requires targeting of zyxin to stress fibers and recruitment of actin regulators. Mol. Biol. Cell 23, 1846–1859 (2012).

39. Nix, D. A. & Beckerle, M. C. Nuclear-cytoplasmic shuttling of the focal contact protein, zyxin: a potential mechanism for communication between sites of cell adhesion and the nucleus. J Cell Biol 138, 1139–1147 (1997).

40. Geiger, B., Spatz, J. P. & Bershadsky, A. D. Environmental sensing through focal adhesions. Nat. Rev. Mol. Cell Biol. 10, 21–33 (2009).

41. Chatterjee, A., Kondaiah, P. & Gundiah, N. Stress fiber growth and remodeling determines cellular morphomechanics under uniaxial cyclic stretch. Biomech. Model. Mechanobiol. 21, 553–567 (2022).

42. Parsons, J.T., Horwitz, A.R. & Schwartz, M.A. Cell adhesion: integrating cytoskeletal dynamics and cellular tension. Nat Rev Mol Cell Biol. Sep;11(9):633-43, (2010).

43. Overall, C. M. & Kleifeld, O. Tumour microenvironment - opinion: validating matrix metalloproteinases as drug targets and anti-targets for cancer therapy. Nat. Rev. Cancer 6, 227–239 (2006).

44. Wang, Y. et al. The role of matrix metalloproteinase 9 in fibrosis diseases and its molecular mechanisms. Biomed. Pharmacother. 171, 116116 (2024).

45. Roca-Cusachs, P., Sunyer, R. & Trepat, X. Mechanical guidance of cell migration: lessons from chemotaxis. Curr. Opin. Cell Biol. 25, 543–549 (2013).

46. Chen, K. et al. Role of boundary conditions in determining cell alignment in response to stretch. Proc. Natl. Acad. Sci. 115, 986–991 (2018).

47. Burridge, K. & Guilluy, C. Focal adhesions, stress fibers and mechanical tension. Exp. Cell Res. 343, 14–20 (2016).

48. Livne, A., Bouchbinder, E. & Geiger, B. Cell reorientation under cyclic stretching. Nat. Commun. 5, 3938 (2014).

49. Tomasek, J. J., Gabbiani, G., Hinz, B., Chaponnier, C. & Brown, R. A. Myofibroblasts and mechano-regulation of connective tissue remodelling. Nat. Rev. Mol. Cell Biol. 3, 349–363 (2002).

50. Piersma, B., Bank, R. A. & Boersema, M. Signaling in Fibrosis: TGF-β, WNT, and YAP/TAZ Converge. Front. Med. 2, 59 (2015).

51. Puleo, J. I. et al. Mechanosensing during directed cell migration requires dynamic actin polymerization at focal adhesions. J. Cell Biol. 218, 4215 (2019).

52. Fiji: an open-source platform for biological-image analysis, Nature Methods. https://www.nature.com/articles/nmeth.2019.

53. Gorelik, R. & Gautreau, A. Quantitative and unbiased analysis of directional persistence in cell migration. Nat. Protoc. 9, 1931–1943 (2014).