Genomic diversity and antimicrobial resistance of *Staphylococcus aureus* in Saudi Arabia: a nationwide study using whole-genome sequencing.

Mohammed S. Alarawi[*1,32,33], Musaad Altammami[4], Mohammed Abutarboush[4], Maxat Kulmanov[1,2,29,30,32], Dalal M. Alkuraithy[7], Senay Kafkas[1,32], Robert Radley[1], Marwa Abdelhakim[1,29,32], Hind Masfer Abdullah Aldakhil[1], Reema A. Bawazeer[12,14], Mohammed A. Alolayan[4], Basel M. Alnafjan[4], Abdulaziz A. Huraysi[4], Amani Almaabadi[4], Bandar A Suliman[5,6], Areej G Aljohani[24], Hassan A. Hemeg[6], Turki S Abujamel[8,9], Anwar M. Hashem[8,9], Ibrahim A Al-Zahrani[9], Mohammed S Abdoh[10], Haya I Hobani[8], Rakan F Felemban[11], Wafaa A Alhazmi[9], Pei-Ying Hong[3], Majed F. Alghoribi[12,13,14,15], Sameera Aljohani[12,13,15], Hanan Balkhy[17], Abdulrahman Alswaji[12,14], Maha Alzayer[12,14], Bassam Alalwan[15], Mai M. Kaaki[18], Sharif M. Hala[13,14,16,31], Omniya Ahmad Fallatah[13,14,16], Wesam Ahmad Bahitham[13,14,16], Samer Yahya Zakri[13,14,16], Mohammad A Alshehri[13,14,16], Nader Kameli[19], Abdullah Algaissi[19], Edrous Alamer[19], Abdulaziz Alhazmi[19], Amjad A.Shajri[20], Majid Ahmed Darraj[21], Bandar Kameli[22], O. O. Sufyani[23], Badreldin Rahama[20], Abrar A. Bakr[34], Fahad M. Alhoshani[4], Azzam A. Alquait[4], Ali Somily[26], Ahmed M. Albarrag[26], Lamia Alosaimi[27], Sumayh A. Aldakeel[27], Fayez S. Bahwerth[28], Tamir T Abdelrahman[37], Séamus Fanning[35,36], Essam J. Alyamani[4], Takashi Gojobori[1,32,33], Satoru Miyazaki[25], Mohammed B. Al-Fageeh[4], and Robert Hoehndorf[†1,2,29,30,32,33]

[1]Computational Bioscience Research Center, King Abdullah University of Science and Technology, KAUST, [2]SDAIA–KAUST Center of Excellence in Data Science and Artificial Intelligence, King Abdullah University of Science and Technology, KAUST, [3]Environmental Science and Engineering Program, Division of Biological and Environmental Science and Engineering, KAUST, [4]Wellness and Preventive Medicine Institute, Health Sector, King Abdulaziz City for Science and Technology (KACST), Riyadh, Saudi Arabia, [5]College of Applied Medical Sciences, Taibah University, Madinah, Saudi Arabia, [6]Department of Medical Laboratory Technology, College of Applied Sciences, Taibah University, Madinah, Saudi Arabia, [7]Department of biological sciences, college of

science, University of Jeddah, Jeddah, Saudi Arabia., [8]Vaccines and Immunotherapy Unit, King Fahd Medical Research Center, King Abdulaziz University, Jeddah 21589, Saudi Arabia, [9]Department of Medical Laboratory Sciences, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah 21589, Saudi Arabia, [10]Epidemiology department, public health Administration, king Abdullah Medical Complex, Jeddah 23816, Saudi Arabia, [11]Alnoor Specialist Hospital, Ministry of Health, Makkah, Saudi Arabia, [12]Infectious Diseases Research Department, King Abdullah International Medical Research Center (KAIMRC), Riyadh, Saudi Arabia., [13]King Abdullah International Medical Research Center (KAIMRC), [14]King Saud bin Abdulaziz University-Health Sciences Ministry of National Guard-Health Affairs (MNGHA), [15]Department of Pathology and Laboratory Medicine, King Abdulaziz Medical City (KAMC),, [16]Ministry of National Guard Health Affairs (MNGHA), Riyadh, Saudi Arabia., [17]World Health Organization, Geneva, Switzerland, [18]Medical Laboratory, King Abdulaziz Medical City (KAMC), Ministry of National Guard Health Affairs, Jeddah, Saudi Arabia, [19]Emerging and Epidemic Infectious Diseases Research Unit, Medical Research Center, Jazan University, Jazan 45142, Saudi Arabia, [20]Department of Medical Laboratories Technology, College of Applied Medical Sciences, Jazan University, Jazan, Saudi Arabia, [21]Department of Medicine,Faculty of Medicine, Jazan University, Jazan, Saudi Arabia, [22]Regional Laboratory & Central Blood Bank Jazan Health, [23]Saudi Public Health Authority, Vector-Borne Diseases Laboratory, Jazan 45142, Saudi Arabia, [24]BndrGene Medical Lab, Madinah, Saudi Arabia, [25]Faculty of Pharmaceutical Sciences, Department of Pharmacy, Tokyo University of Science, Noda, Chiba, Japan, [26]Department of pathology, college of Medicine, King Saud University and King Saud University Medical City, Riyadh, Saudi Arabia, [27]The National Center for Genomic Technology (NCGT), Life Science and Environment Research Institute, King Abdulaziz City for Science and Technology (KACST), Riyadh, Saudi Arabia, [28]Medical Microbiology Laboratory, Hera General Hospital, Makkah healthcare cluster, Makkah, Saudi Arabia., [29]KAUST Center of Excellence for Smart Health (KCSH), King Abdullah University of Science and Technology, 4700 KAUST, Thuwal 23955, Saudi Arabia, [30]KAUST Center of Excellence for Generative AI, King Abdullah University of Science and Technology, 4700 KAUST, Thuwal 23955, Saudi Arabia, [31]Biothreat Response Department, Public Health Laboratory, the Saudi Public Health Authority, [32]Computer, Electrical and Mathematical Sciences & Engineering (CEMSE) Division, King Abdullah University of Science and Technology, King Abdullah University of Science and Technology, 4700 KAUST, Thuwal, Saudi Arabia., [33]Biological and Environmental Sciences & Engineering (BESE) Division, King Abdullah University of Science and Technology, King Abdullah University of Science and Technology, 4700 KAUST, Thuwal, Saudi Arabia, [34]Advanced Diagnostics and Therapeutics Institute, Health Sector, King Abdulaziz City for Science and Technology (KACST), Riyadh, Saudi

Arabia, [35]UCD-Centre for Food Safety, University College Dublin, Belfield, Dublin D04 N2E5, Ireland, [36]nstitute

for Global Food Security (IGFS), The Queen's University of Belfast, 19 Chlorine Gardens, Belfast BT9 5DL,

Northern Ireland, United Kingdom, [37]King Faisal Specialist Hospital & Research Centre- Madina

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) surveillance in regions with mass gatherings presents unique challenges for public health systems. Saudi Arabia, hosting millions of pilgrims annually, provides a distinctive setting for studying how human mobility shapes bacterial populations, yet comprehensive genomic surveillance data from this region remains limited. Here, we present an integrated analysis of *S. aureus* isolates collected across seven Saudi Arabian regions, combining whole-genome sequencing with extensive antimicrobial susceptibility testing and standardized metadata following FAIR data principles. Our analysis revealed striking differences between pilgrimage and non-pilgrimage cities. Pilgrimage cities showed significantly higher genetic diversity and antimicrobial resistance rates, harboring numerous international strains including recognized clones from diverse geographic origins. Reported lineage dynamics is changing, expanding toward community clones. While genomic prediction of antimicrobial resistance showed high accuracy for some antibiotics, particularly beta-lactams, with varying performance for others, highlighting the necessity for phenotypic testing in clinical settings. Our findings demonstrate how mass gatherings drive bacterial population structures and emphasize the importance of integrated surveillance approaches in regions with significant global connectivity and travel.

Importance

Genomic data enables the tracking of pathogens by revealing clonal expansions within populations and identifying successful lineages. However, comprehensive national-level data from Saudi Arabia remains limited on a large scale. The adoption of FAIR principles and reproducible workflows ensures robust,

---

*Address correspondence to: mohammed.alarawi@kaust.edu.sa

†Address correspondence to: robert.hoehndorf@kaust.edu.sa

consistent analysis, fostering effective data sharing. The OneHealth approach's success depends on the integration and collaboration across diverse domains in today's digital landscape.

Introduction

Staphylococcus aureus, a Gram-positive bacterium characterized by its grape-like clustering morphology, is both a common human and animal commensal organism and a significant pathogen that has emerged as a leading cause of hospital-acquired infections. Notably, methicillin-resistant *S. aureus* (MRSA) accounts for approximately 25-50% of these infections. MRSA emerged shortly after methicillin's introduction in 1958, following the acquisition of a mobile genetic element known as the staphylococcal chromosomal cassette `mec` (`SCCmec`). This cassette harbors the `mecA` gene, which encodes PBP2A, an alternative penicillin-binding protein involved in cell-wall synthesis. The modified protein exhibits reduced affinity to β-lactam antibiotics, conferring broad-spectrum resistance to this antibiotic class [30, 32].

The global impact of MRSA has expanded significantly beyond healthcare settings, establishing distinct reservoirs within communities and livestock populations [34, 21]. Understanding the epidemiological distribution of MRSA lineages and their corresponding resistance phenotypes across communities, regions, and countries has become increasingly crucial from public health prospective. This knowledge directly informs the development of targeted mitigation strategies, encompassing evidence-based policies for antibiotic stewardship, systematic screening protocols, and comprehensive livestock management practices [5, 42]. The clinical significance of MRSA is particularly evident in its role as a major pathogen in post-surgical complications, community-onset infections, and foodborne illness outbreaks [34, 6].

Saudi Arabia presents a unique context for studying MRSA transmission and resistance patterns. As the largest country on the Arabian Peninsula by both area and population, it holds profound religious significance as the custodian of Islam's two holiest sites. This unique position draws millions of Muslim pilgrims annually to the Kingdom. Mass gatherings increase chances for microbial transmission and the introduction of novel clones and antimicrobial resistance patterns. Large cities are hypothesized to function as entry points and subsequent dissemination of imported clones due to population density and connectivity. Consequently, international clones are predicted to emerge initially in these entry points before spreading in other regions within Saudi Arabia. Smaller, less connected cities are expected to

display a less presence of these clones. The epidemiology of MRSA in Saudi Arabia has been documented through various investigations [6, 47, 2, 14], although most studies have been limited to individual healthcare facilities or specific geographical regions. These investigations range from comprehensive phenotypic characterization of MRSA isolates' antimicrobial susceptibility profiles [2, 7, 1] to more detailed molecular analyses of strain characteristics [47].

Despite extensive research on MRSA in Saudi Arabia, there remains a significant gap in comprehensive datasets that integrate both genotypic and phenotypic information across multiple geographical regions. One critical component to address this gap is the use of FAIR (Findable, Accessible, Interoperable, Reusable) data principles [54]. Implementation of FAIR increases the value of accumulated data by enabling easier reuse and interoperability. Interoperable data is valuable because it can be combined with other datasets and aid in elucidating molecular mechanisms of antimicrobial resistance and identifying novel therapeutic targets in addition to transmission burden. Moreover, combined genotype–phenotype datasets enable the development of machine learning approaches for predicting drug resistance patterns and pathogenicity profiles, while also informing the design of targeted studies aligned with the One Health approach.

To address this knowledge gap, we established a collection of 686 *S. aureus* isolates from diverse regions across Saudi Arabia, encompassing clinical specimens, community screening samples, and wastewater isolates. We conducted whole-genome sequencing on the isolates and performed extensive drug resistance phenotyping screening, creating a novel geospatial genotype–phenotype dataset. Our analysis reveals high genetic diversity, including several previously unidentified sequence types distributed across different regions. These novel sequence types exhibit elevated patterns of drug resistance. Significantly, we observed higher genetic diversity and increased levels of drug resistance in cities associated with mass gatherings (Jeddah, Makkah, Madinah), suggesting that mass gathering may serve as drivers of antimicrobial resistance evolution and transmission. These findings provide crucial insights for developing targeted intervention strategies and underscore the importance of genomic surveillance in regions experiencing regular mass gatherings.

Materials and Methods

Sample Selection and Sourcing

Our study includes a total of 686 *S. aureus* isolates collected from seven distinct regions across Saudi Arabia. The majority of samples were obtained from tertiary care hospitals, with the Western region contributing the largest proportion: Jeddah (275 isolates), Madinah (102 isolates), and Makkah (36 isolates). Additional regional collections include Riyadh in the Central region (158 isolates), Hail in the North (64 isolates), AlHasaa in the East (33 isolates), and Jazan in the South (18 isolates). To broaden the scope of our surveillance, we supplemented the hospital-sourced isolates with 60 samples collected through a community screening of healthy individuals in Jeddah city and 5 environmental wastewater samples from the same region. A detailed breakdown of sample distribution is provided in supplementary Table S1 . Sample selection within each hospital followed a random sampling approach, and we included both methicillin-resistant (MRSA) and methicillin-susceptible *S. aureus* (MSSA) isolates to ensure comprehensive representation of circulating strains.

Bacterial Identification and Susceptibility Testing

All isolates were initially enriched on tryptic soy agar (TSA) supplemented with 5% [w/v] NaCl (Sigma-Aldrich, Germany) to selectively culture *S. aureus*. Single colonies were used for subsequent bacterial identification, susceptibility testing, and glycerol stock archival. Throughout the identification process, the *S. aureus* ATCC 29213 strain served as an internal reference control. Phenotypic drug resistance profiling was performed using the VITEK® 2 System with GP ID cards for identification and AST-P580 cards for antimicrobial susceptibility testing (bioMérieux, France). Supplementary table S2 shows the antimicrobial agents and their concentrations of the AST-P580 card. All test results were digitally recorded and stored for analysis.

DNA Isolation

Genomic DNA isolation was performed using single colonies obtained either directly from Petri dishes or from enriched broth cultures following identification. For each sample, 2 mL of overnight culture was

pelleted for DNA extraction. Concurrently, we prepared archived samples by combining 500 µL molecular grade glycerol (Thermofisher, USA) with 500 µL of the culture for future investigations. Genomic DNA was isolated using the automated KingFisher MagMax DNA isolation kit protocol (Thermofisher, USA). DNA quality and quantity were assessed using both a nano-drop instrument (Thermofisher, USA) and Qubit dsDNA BR kit (Thermofisher, USA).

Library Preparation and DNA Sequencing

Genomic DNA libraries were constructed using 100 ng of input DNA with the QIASeq FX DNA library kit (Qiagen, Germany) following the manufacturer's protocol. Library quality was assessed using an Agilent Bioanalyzer system (Agilent, USA). Whole-genome sequencing was performed on the Illumina NovaSeq 6000 platform (Illumina, USA) using SP flow-cells, generating 150-base paired-end reads with average insert sizes ranging from 350 to 450 bp, and 50-100X sequence depth.

Genomic Analysis

Initial quality control and preprocessing of raw sequencing reads were perfomred using TrimGalore v0.4.4 for adapter trimming [9]. Taxonomic identification and contamination utilized Kraken v2.0.8 beta [55] and Mash v2.3 [39]. Genomes were assembled using SKESA v2.4.0 [49] and called variants using Snippy v4.6.0 against the reference genome `NC_007795.1`.

Annotation of the assembled genomes utilized prokka v1.14.6 [46] and Roary for pangenome construction v3.13.0 [40]. A neighbor-joining phylogenetic tree was built using FastTree2 2.1.10 [43] based on the core genome alignment. Assignment of multilocus sequence types (MLST) performed using ABRicate v1.0.1 with the MLST tool.

Multiple approaches were employed for antimicrobial resistance analysis. Drug resistance genes were identified using both the CARD v5.1.1 [3] and ResFinder v4.0 [18] databases via ABRicate v1.0.1 [52]. Additionally, DeepARG v1.0.2 [10], a deep learning approach, to enhance resistance gene detection. Virulence factors were identified using the Virulence Factor Database (VFDB) [22] through ABRicate v1.0.1.

Methicillin resistance mechanisms were characterized by screening for the `mecA` gene using the staphopia-Sccmec v1.0.0 typing tool [41]. The analysis was complemented using a local database contain-

ing whole cassette sequences (`mecA`, `mecB`, `mecC`), cassette recombinase genes (CCR) genes, and inser-
tion sequences for each SCC*mec* type. Additional `mecA` cassette detection was performed using minimap2
v2.24-r1122 [35] with a local database. The SCC*mec* type assignments were determined based on cassette
element arrangements following the workflow at `https://github.com/cdnstp/SCCmec_CLA`.
Finally, Protein A gene polymorphism (spa) typing was performed using spaTyper v4.6.1 [44] with its
associated database.

Reproducibility and Implementation of FAIR Data Principles

We implemented our entire analysis workflow using the Common Workflow Language (CWL) [23] to
ensure computational reproducibility. The two main analysis workflows are shown in Supplementary
Figures S1 and S2 .

We executed all analyses on a local Arvados system [The Arvados Authors et al.], with complete
workflow definitions and execution environments preserved. To promote Findability, Accessibility, In-
teroperability, and Reusability (FAIR), we have made all source code, workflow definitions, and con-
figuration files freely available at `https://github.com/bio-ontology-research-group/`
`mrsa-sequences`. The repository includes detailed documentation of computational requirements,
software versions, and usage instructions. Additionally, raw sequencing data were deposited in the Eu-
ropean Nucleotide Archive (ENA) under accession number `PRJEB59751`. Furthermore, the metadata
of our samples as well as the measured antimicrobial resistance phenotypes using the Resource Descrip-
tion Framework (RDF) [33] are made available in a public repository [36] and on a Github repository.
Supplementary Figure S3 describes the RDF data model.

Results

Genetic diversity and geographic distribution of *S. aureus* in Saudi Arabia

686 samples of *S. aureus* from multiple regions in Saudi Arabia were sequenced, following quality control
validation, and confirmation as *S. aureus* through whole-genome sequencing and phenotypic testing. The
isolates consist of methicillin-resistant (MRSA) and methicillin-susceptible (MSSA) strains, with MSSA

proportions varying significantly by region: from 3.9% in Madinah to 34.4% in Hail (Table 1).

Molecular typing through PubMLST [31] classification was performed to understand the evolutionary relationships and population structure of these isolates, as this knowledge is crucial for tracking transmission patterns and implementing targeted infection control measures. The classification revealed nine major clonal complexes, with CC5 (26.38%), CC97 (12.39%), and CC22 (9.48%) being the most prevalent (Supplementary Figures S4). 31.92% of isolates belonged to previously unassigned clonal complexes. Our analysis also identified 14 novel sequence types (Supplementary Table S3). Figure 1 provides an overview of the molecular diversity and relatedness of the samples included in our study, in addition to genetics distance based on core Single Nucleotide Variants (SNVs) minimum spanning tree (Supplementary Figure S5).

We observed distinct patterns of strain distribution aligned with pilgrimage routes in major cities. Cities along the Hajj and Umrah routes exhibited high genetic diversity: Jeddah (9 CCs, 43 STs), Madinah (9 CCs, 26 STs), and Makkah (8 CCs, 14 STs), likely reflecting the international convergence of travelers. In contrast, regions with less international traffic showed more limited diversity, particularly Jazan (4 CCs, 6 STs). Analysis of the SCC*mec* revealed region-specific patterns, with SCC*mec*V ranging from 23.4% in Hail to 61.1% in Jazan, and SCC*mec*IVa ranging from 19.6% in Riyadh to 45.5% in AlHasa (Table 1, Figure 2, and Supplementary Figure S6).

Several internationally recognized strains were observed in pilgrimage cities. The Bengal Bay clone (CC1-ST772-PVL positive) [38] and CC30-ST1482 [13] were detected primarily in Makkah and Jeddah, regions that host large numbers of international pilgrims. The South Pacific Clone (CC30-ST30) [53] was found across most cities.

Phenotypic antimicrobial resistance

Antimicrobial susceptibility testing was performed on all isolates (Table 2). We observed the highest resistance rates for benzylpenicillin across all regions, ranging from 88.9% in AlHasa to 100% in Makkah. Oxacillin resistance, mediated by mecA, showed considerable variation, from 60.3% in Hail to 96.6% in Madinah. Fusidic acid emerged as the next most prevalent resistance, with particularly high rates in Riyadh (70.5%) and Jeddah (70.9%) but notably lower in Hail (35.3%). Fluoroquinolone resistance varied

Figure 1: Phylogenomic structure and diversity of *S. aureus* across Saudi Arabia. Maximum likelihood tree constructed from core genome SNPs of 686 isolates. Major Clonal Complexes (CCs), grouping related Sequence Types, are highlighted (the dominant CC5, CC8, CC22, ST672). Coloured rings display the associated SCCmec type (inner ring) and geographical region of origin (outer ring), revealing the distribution patterns of major lineages. CC5 two branches (ST5, and ST6) and the distinct clustering reflecting national population structure influenced by regional and international factors. Key virulence/resistance markers (PVL, TSST, ACME, fusB/C) are indicated in the outermost tracks.

Figure 2: Stacked bar chart illustrating the relative proportion (%) of the top 10 most frequent *S. aureus* isolates Sequence types per region from Saudi Arabia. The chart highlights the greater ST diversity observed in major pilgrimage-associated cities (Jeddah, Madinah, Makkah) and other high-traffic locations like Riyadh, compared to regions like Jazan, which shows dominance by fewer STs (e.g., ST5, ST30). Notable STs like ST5, ST97, ST672, ST8, and ST22 show varying prevalence across regions, indicating complex population structures influenced by both local factors and international travel patterns regions,.

significantly by region, with levofloxacin resistance ranging from 14.7% in Hail to 45.7% in Madinah, and similar patterns for moxifloxacin (13.4% to 41.5%). Erythromycin resistance rates ranged from 19.4% in AlHasa to 37.9% in Madinah. We observed variable resistance to aminoglycosides, with tobramycin resistance ranging from 8.1% in Makkah to 32.5% in Jeddah, while gentamicin resistance showed similar patterns. Tetracycline resistance rates varied considerably, from 6.2% in AlHasa to 26.5% in Jeddah.

Several antibiotics maintained complete efficacy across all regions. We observed no resistance to linezolid, tigecycline, or teicoplanin. Vancomycin, often considered a last-resort antibiotic [17], showed nearly complete effectiveness, with only a single borderline case identified in a sample from Madinah with updated MIC break point. Similarly, nitrofurantoin and rifampicin maintained high effectiveness, with resistance rates not exceeding 1.7% in any region.

Comparison of overall differences of antimicrobial resistance was performed for locations associated with mass gatherings and other cities. Pilgrimage-associated cities (Madinah, Makkah, and Jeddah) showed significantly higher overall resistance rates compared to other locations (33.9% vs 27.6%, $p = 0.047$, one-tailed t-test).

Genotypic identification of antimicrobial resistance

While whole-genome sequencing data identify resistance-associated genotypes, linking these directly to observed phenotypes remains a complex process. We compared phenotypic resistance results to genetic predictions using three approaches: CARD [4], ResFinder [18], and the deep learning tool DeepARG [10]. Genetic predictions showed varying levels of concordance with phenotypic resistance across different antibiotic classes (Table 3).

Beta-lactam predictions demonstrated high accuracy, particularly for benzylpenicillin, where both CARD and DeepARG achieved identical high performance (precision 0.983, recall 0.980), while ResFinder showed higher precision (0.994) but lower recall (0.572). Oxacillin predictions showed strong agreement between CARD and DeepARG (precision 0.858, recall 0.976), while ResFinder did not provide predictions for this antibiotic. For fusidic acid, both CARD and ResFinder achieved accurate results (precision 0.937 and 0.940 respectively), while DeepARG could only make few predictions.

However, we observed substantial discrepancies for other antibiotic classes. Aminoglycoside resis-

tance predictions varied markedly across tools. For gentamicin and tobramycin, CARD and DeepARG showed low precision (0.172–0.221) but high recall (0.937), while ResFinder maintained more balanced performance (precision 0.752–0.785, recall 0.793–0.807). Similarly, predictions for macrolides and lincosamides showed moderate concordance, erythromycin and clindamycin predictions showed only low concordance with phenotypic observations.

To test for the presence of known and potentially novel mechanisms of antimicrobial resistance, we constructed a pangenome (Supplementary Figure S8) and performed a pangenome-wide association study using Scoary [20] (Supplementary Table S4) . The result of the bacterial GWAS identified known mechanisms of antimicrobial resistance. Genes such as *mecA* and *mecR1* were strongly associated with oxacillin and cefoxitin resistance. Similarly, the aminoglycoside resistance was determinated by *aacA-aphD* and *knt* for tobramycin; tetracycline resistance was shown to be mediated by efflux pump *tet(K)*; and the genes *ermC* and *msrA* were associated with resistance to macrolides and lincosamides (erythromycin, clindamycin).

However, non-canonical association were present for other drugs classes. For instance, fluoroquinolone resistance was associated with genes such as *hsdM* and *entC2*. Benzylpenicillin resistance associated predominantly with *fcl_1*. Fusidic acid resistance showed associations primarily with transport-related genes (*natA*, *macB*, and *dppB*).

Virulence and pathogenicity

The prevalence and distribution of toxin and virulence factors indicates regional variation (see Figure 1 and Supplementary Table S5). Panton-Valentine Leukocidin (PVL), a cytotoxin often associated with community-associated MRSA (CA-MRSA), was detected in 17% of isolates overall. However, its prevalence showed marked regional differences, being highest in Madinah (32.4%) and lowest in Hail (6.3%). Notably, PVL positivity was linked to specific lineages found predominantly in pilgrimage cities, including the internationally recognized CC1/ST772 (Bengal Bay clone).

Toxic shock syndrome toxin (TSST) was found in 10% of isolates and TSST rates showed an inverse geographic distribution pattern, ranging from 0% in Jazan to 26% in Hail, with intermediate rates in pilgrimage cities: Jeddah (8%), Makkah (8%), and Madinah (13%). Western cities (Jeddah, Madina, and

Makkah) harbored the majority of Novel (ACME)-positive isolates.

Core virulence factors displayed widespread but distinctive distribution patterns across our isolates. Adhesion factors showed varying prevalence: clumping factors (clfA, clfB) appeared in over 95% of isolates, while the collagen binding protein (cna) occurred exclusively in CC5. The majority of isolates (>90%) carried immune evasion genes (*sak*, *sbi*, *scn*) and biofilm formation genes (icaABCD cluster). Among toxin genes, *sea* dominated with 30% prevalence, followed by other enterotoxins (*seb*, *sec*, *sed*, *seh*).

Discussion

Impact of mass gatherings on MRSA diversity and resistance patterns

This study described the genetic epidemiology of *S. aureus* across Saudi Arabia, a country with over 32 million inhabitants of which 41% are immigrants [11]. As of 2023, Saudi Arabia hosts over 2 million pilgrims for Hajj and 27 million pilgrims for Umrah annually [12]. While previous work has investigated the effect of mass gatherings and pilgrimage on antimicrobial resistance in Saudi Arabia [28], our integrated genotype–phenotype dataset reveals several distinct patterns associated with mass gathering activities.

The impact of mass gatherings on MRSA epidemiology is particularly evident in cities along the Hajj and Umrah routes. Jeddah, Makkah, and Madina showed significantly higher genetic diversity and unique strain patterns compared to other regions, with Jeddah, the main entry point for pilgrims, exhibiting the highest number of novel sequence types. This pattern aligns with observations from other Gulf regions, where mass gatherings coincide with diverse strain distributions [48, 19]. This diversity also includes a prevalence of internationally recognized strains, such as the PVL-positive Bengal Bay clone (CC1-ST772) and the South Pacific clone (CC30-ST30), which are found in pilgrimage cities, indicating a link between pilgrimage routes and the importation of globally circulating lineages [16].

This findings presented suggest that mass gatherings influence not only genetic diversity but also antimicrobial resistance patterns. Pilgrimage cities showed significantly higher overall resistance rates compared to other locations, suggesting these sites may serve as hotspots for resistance transmission. This is particularly evident in the distribution of SCC*mec* types, which suggests a shift from hospital-associated

to community-associated MRSA, reflecting global trends [24, 37]. The concentrated presence of ACME-positive isolates in western pilgrimage cities further supports the role of international travel in strain dissemination.

The successful persistence of these strains in mass gathering-associated regions appears to be facilitated by multiple factors. The high prevalence of adaptive features, including biofilm formation genes and immune evasion clusters, alongside various resistance mechanisms, may enable these strains to be established in these high-traffic areas. These findings extend previous work on the impact of mass gatherings on bacterial populations [1], demonstrating how such events can facilitate the concurrent spread of both antimicrobial resistance determinants and virulence factors.

This unique demographic pattern and its influence on MRSA evolution and clonal expansion [34] presents both challenges and opportunities for public health interventions. Our findings suggest that large travel hub cities may serve as entry points for emerging strains and resistance patterns, potentially allowing early detection of concerning variants before they achieve wider distribution.

CC97 is detected as MSSA, and MRSA carrying SCCmecV associated with FusC gene. Newly identified ST8637 in Madinah is an indicator of active expansion of this linage and successful distribution in the community. The presence of CC97 in clinical samples indicates successful establishment in hospitals. More investigations are needed to understand the genetic basis of this adaptation and its implications for public health, especially as it ranks second in presence across the regions.

Interestingly, a significant increase in the proportion of novel sequence type assignments in Jeddah (9/14) was observed, and this the main entry point for international travelers to Hajj and Umrah, and provides a further indication that pilgrimage is one of the main drivers of drug resistance and diversity in Saudi Arabia.

Limitations of Genomic Prediction for Antimicrobial Resistance

These findings highlight the complex relationship between genotype and phenotype in antimicrobial resistance prediction. While genomic prediction tools showed high precision for certain antibiotics (over 0.98 for beta-lactams and fusidic acid), they performed poorly for other drug classes, particularly aminoglycosides. This variable performance reflects both the complexity of resistance mechanisms and the current

limitations of prediction methods.

The GWAS analysis revealed both expected and unexpected genetic associations. While we confirmed established resistance determinants such as *mecA_1* and *mecR1* for beta-lactams, and *ermC* and *msr(A)* for macrolide resistance — we also identified several unexpected associations requiring careful interpretation. For instance, fluoroquinolone resistance showed associations with *hsdM* and *entC2*, likely representing indirect relationships rather than causative mechanisms. Similarly, the association of fusidic acid resistance with transport-related genes (*natA*, *macB*, *dppB*) rather than expected target site mutations suggests potential limitations in our GWAS approach for detecting point mutations.

This discrepancy between genetic markers and phenotypic resistance highlights several challenges. In particular, resistance mechanisms may involve complex genetic interactions that are not captured by current prediction tools. These may include environmental factors and gene expression regulation which can contribute to resistance phenotypes independently of genetic markers. The convergence of diverse international strains in our study setting may facilitate the exchange of resistance determinants through mobile genetic elements, complicating genotype–phenotype relationships.

Given these limitations, we would suggest maintaining phenotypic testing alongside genomic surveillance for effective clinical decision-making [15]. Such a dual approach is especially crucial in regions experiencing high levels of international travel and strain diversity, where rapid evolution and transmission of resistance mechanisms may occur. Future improvements in prediction accuracy will likely require integration of multiple data types, including transcriptomics and proteomics, to better capture the complex determinants of antimicrobial resistance.

Data Availability and Reproducibility

A key contribution of this study is the comprehensive implementation of FAIR (Findable, Accessible, Interoperable, and Reusable) principles [54] in data sharing. We structured our metadata using the Resource Description Framework (RDF) [33], creating a semantically rich representation of both sequence data and phenotypic measurements. Our data model (Supplementary Figure S3) incorporates standard terminologies and ontologies including the NCBI Taxonomy [45] for species annotation, the ChEBI ontology [29] for antimicrobial agents, the NCI Thesaurus [26] and the GENEPIO ontology [27] for health status

descriptions, and the Unit Ontology (UO) [25] for standardized units, enhancing interoperability with existing datasets.

This structured approach to data sharing serves multiple purposes. First, it enables direct integration with other surveillance datasets through standardized terminology and relationships. The use of controlled vocabularies for key metadata elements — such as specimen sources, host characteristics, and antimicrobial susceptibility measurements — facilitates automated data integration and comparative analyses across different studies and geographical regions.

Second, this dataset reported is particularly valuable for developing and validating machine learning approaches for antimicrobial resistance prediction. By providing paired genomic and phenotypic data in a standardized format, along with detailed host and environmental metadata, we enable researchers to develop prediction models that can account for contextual factors beyond purely genetic determinants, and may be more adapted to the complex demographics found among pilgrims. The comprehensive antimicrobial susceptibility data, including both MIC values and categorical interpretations, further provides rich training and validation data for such models.

The complete computational reproducibility of our analysis, implemented through a workflow in the Common Workflow Language (CWL) [8], ensures that other researchers can not only access our data but also reproduce and build upon our analytical methods. By making our workflows, source code, and execution environments publicly available, we facilitate easier adaptation of our methods to new datasets and research questions.

Implementation of FAIR principles extends beyond technical accessibility to practical reusability. The standardized metadata format allows researchers to integrate our findings with their local surveillance data, potentially revealing new patterns of MRSA transmission and evolution. Furthermore, our dataset can serve as a baseline for tracking changes in MRSA populations, particularly in the context of mass gatherings and international travel, where standardized data collection and sharing are crucial for effective surveillance.

Future Work

Our findings as reported point to several key directions for future surveillance strategies, particularly focusing on early detection at pilgrimage entry points. The implementation of systematic sampling could be strategically enhanced by targeting major travel hubs. Specifically, wastewater surveillance at airports serving pilgrimage routes, particularly the international airports in Jeddah and Madinah, could provide early warning signals of emerging strains. Aircraft wastewater sampling from pilgrimage flights, which has proven effective for COVID-19 surveillance [50], could be adapted for MRSA monitoring, offering insights into strain importation patterns before pilgrims arrive at their destinations.

Such targeted approach could be complemented by environmental sampling at key congregation points along pilgrimage routes. Strategic sampling points could include ablution facilities in major mosques and shared sanitation facilities in pilgrim accommodations, providing a comprehensive picture of strain circulation during mass gatherings. Integration of these environmental samples with clinical surveillance would create a more complete understanding of MRSA transmission dynamics during mass gathering.

The application of long-read sequencing technologies to these samples could resolve complex genetic elements that are challenging to characterize with short-read data alone. This would be particularly valuable for understanding the structure of SCCmec cassettes and other mobile genetic elements that may be exchanged during mass gatherings.

Finally, establishing a real-time surveillance system that combines this strategic sampling with rapid genomic and phenotypic analysis would enable faster response to emerging resistant strains. Such a system could serve as a model for monitoring other pathogens in similar mass gathering settings, contributing to global pathogen surveillance efforts and early warning systems for emerging antimicrobial resistance.

Conclusion

In conclusion this study demonstrates how mass gatherings and regional factors jointly shape bacterial population structures and antimicrobial resistance patterns at a national scale. While pilgrimage cities exemplify the intersection of global human mobility with pathogen evolution, the distinct patterns we observed in non-pilgrimage regions highlight the importance of local healthcare practices and potential

livestock–human transmission routes in shaping MRSA populations. This comprehensive view, spanning both high-traffic pilgrimage sites and regional healthcare settings, provides evidence for a framework to understand pathogen dynamics in complex environments where international, community, and agricultural factors converge. Through standardized surveillance approaches and data sharing, our work establishes a foundation for monitoring antimicrobial resistance in the context of both global human mobility and regional One Health challenges, with implications for public health strategies across the Middle East and beyond.

1 Data Availability Statement

Sequencing data is available on the Sequence Read Archive (SRA) as a BioProject under accession number `PRJEB59751` The phenotypes for each samples are available on Zenodo under DOI `10.5281/zenodo.1425085` [36].

2 ETHICS APPROVAL

This work approved by the Institutional Biosafety and Bioethics Committee (IBEC) of King Abdullah University of Science and Technology under approval number `18IBEC14`.

3 FUNDING

This work was sponsored by King Abdulaziz City for Science and Technology under award number ETSC&KACST-KAUST-2018-05-27-01.

4 CONFLICTS OF INTEREST

The authors declare that no conflicts of interest exist.

References

[1] Ahmed, B. and Mashat, B. H. (2014). Prevalence of classical enterotoxin genes in staphylococcus aureus isolated from food handlers in makkah city kitchens. *Asian J. Sci. Tech*, 5(11):727–731.

[2] Al Yousef, S. and Taha, E. (2016). Methicillin-resistant staphylococcus aureus in saudi arabia: Genotypes distribution review. *Saudi Journal of Medicine and Medical Sciences*, 4(1):2.

[3] Alcock, B. P., Raphenya, A. R., Lau, T. T., Tsang, K. K., Bouchard, M., Edalatmand, A., Huynh, W., Nguyen, A.-L. V., Cheng, A. A., Liu, S., et al. (2020a). Card 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic acids research*, 48(D1):D517–D525.

[4] Alcock, B. P., Raphenya, A. R., Lau, T. T., Tsang, K. K., Bouchard, M., Edalatmand, A., Huynh, W., Nguyen, A.-L. V., Cheng, A. A., Liu, S., et al. (2020b). Card 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic acids research*, 48(D1):D517–D525.

[5] Algammal, A. M., Hetta, H. F., Elkelish, A., Alkhalifah, D. H. H., Hozzein, W. N., Batiha, G. E.-S., El Nahhas, N., and Mabrok, M. A. (2020). Methicillin-resistant staphylococcus aureus (mrsa): One health perspective approach to the bacterium epidemiology, virulence factors, antibiotic-resistance, and zoonotic impact. *Infection and Drug Resistance*, Volume 13:3255–3265.

[6] Aljeldah, M. M. (2020). Prevalence of Methicillin-Resistant Staphylococcus aureus (MRSA) in Saudi Arabia: A Systematic Review. *Journal of Pure and Applied Microbiology*, 14(1):37–46.

[7] Alkuraythi, D. M., Alkhulaifi, M. M., Binjomah, A. Z., Alarwi, M., Aldakhil, H. M., Mujallad, M. I., Alharbi, S. A., Alshomrani, M., Alshahrani, S. M., Gojobori, T., et al. (2023). Clonal flux and spread of staphylococcus aureus isolated from meat and its genetic relatedness to staphylococcus aureus isolated from patients in saudi arabia. *Microorganisms*, 11(12):2926.

[8] Amstutz, P., Crusoe, M. R., Tijanić, N., Chapman, B., Chilton, J., Heuer, M., Kartashov, A., Leehr, D., Ménager, H., Nedeljkovich, M., et al. (2016). Common workflow language, v1. 0.

[9] Andrews, S., Krueger, F., Segonds-Pichon, A., Biggins, L., Virk, B., Dalle-Pezze, P., Wingett, S., Saadeh, H., and Ahlfors, H. (2015). Trim galore. *Trim Galore*.

[10] Arango-Argoty, G., Garner, E., Pruden, A., Heath, L. S., Vikesland, P., and Zhang, L. (2018). Deeparg: a deep learning approach for predicting antibiotic resistance genes from metagenomic data. *Microbiome*, 6:1–15.

[11] Authority, G. S. (2022). Saudcensus: Population summary report.

[12] Authority, G. S. (2023). Umrah statistics 2023. `https://www.stats.gov.sa/sites/default/files/Umrah%20Statistics%202023%20EN.pdf`. [Accessed 23-11-2024].

[13] Bakthavatchalam, Y. D., Vasudevan, K., Rao, S., Varughese, S., Rupali, P., Gina, M., Zervos, M., Peter, J. V., and Veeraraghavan, B. (2021). Genomic portrait of community-associated methicillin-resistant staphylococcus aureus st772-sccmec v lineage from india. *Gene Reports*, 24:101235.

[14] Balkhy, H., Memish, Z., Almuneef, M., Cunningham, G., Francis, C., Fong, K., Nazeer, Z., and Tannous, E. (2007). Methicillin-resistant staphylococcus aureus: a 5-year review of surveillance data in a tertiary care hospital in saudi arabia. *Infection Control & Hospital Epidemiology*, 28(8):976–982.

[15] Bhattacharyya, R. P., Bandyopadhyay, N., Ma, P., Son, S. S., Liu, J., He, L. L., Wu, L., Khafizov, R., Boykin, R., Cerqueira, G. C., et al. (2019). Simultaneous detection of genotype and phenotype enables rapid and accurate antibiotic susceptibility determination. *Nature medicine*, 25(12):1858–1864.

[16] Blomfeldt, A., Larssen, K., Moghen, A., Gabrielsen, C., Elstrøm, P., Aamot, H., and Jørgensen, S. (2017). Emerging multidrug-resistant bengal bay clone st772-mrsa-v in norway: molecular epidemiology 2004–2014. *European Journal of Clinical Microbiology & Infectious Diseases*, 36:1911–1921.

[17] Boneca, I. G. and Chiosis, G. (2003). Vancomycin resistance: occurrence, mechanisms and strategies to combat it. *Expert Opinion on Therapeutic Targets*, 7(3):311–328.

[18] Bortolaia, V., Kaas, R. S., Ruppe, E., Roberts, M. C., Schwarz, S., Cattoir, V., Philippon, A., Allesoe, R. L., Rebelo, A. R., Florensa, A. F., et al. (2020). Resfinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy*, 75(12):3491–3500.

[19] Boswihi, S. S., Udo, E. E., and AlFouzan, W. (2020). Antibiotic resistance and typing of the methicillin-resistant staphylococcus aureus clones in kuwait hospitals, 2016–2017. *BMC microbiology*, 20:1–9.

[20] Brynildsrud, O., Bohlin, J., Scheffer, L., and Eldholm, V. (2016). Rapid scoring of genes in microbial pan-genome-wide association studies with scoary. *Genome biology*, 17:1–9.

[21] Butaye, P., Argudín, M. A., and Smith, T. C. (2016). Livestock-associated mrsa and its current evolution. *Current Clinical Microbiology Reports*, 3(1):19–31.

[22] Chen, L., Zheng, D., Liu, B., Yang, J., and Jin, Q. (2015). Vfdb 2016: hierarchical and refined dataset for big data analysis—10 years on. *Nucleic Acids Research*, 44(D1):D694–D697.

[23] Crusoe, M. R., Abeln, S., Iosup, A., Amstutz, P., Chilton, J., Tijanić, N., Ménager, H., Soiland-Reyes, S., Gavrilović, B., Goble, C., and Community, T. C. (2022). Methods included: standardizing computational reuse and portability with the common workflow language. *Communications of the ACM*, 65(6):54–63.

[24] David, M. Z., Cadilla, A., Boyle-Vavra, S., and Daum, R. S. (2014). Replacement of ha-mrsa by ca-mrsa infections at an academic medical center in the midwestern united states, 2004-5 to 2008. *PLoS One*, 9(4):e92760.

[25] Gkoutos, G. V., Schofield, P. N., and Hoehndorf, R. (2012). The units ontology: a tool for integrating units of measurement in science. *Database*, 2012(0):bas033–bas033.

[26] Golbeck, J., Fragoso, G., Hartel, F., Hendler, J., Oberthaler, J., and Parsia, B. (2003). The national cancer institute's thésaurus and ontology. *Journal of Web Semantics*, 1(1):75–80.

[27] Griffiths, E., Dooley, D., Graham, M., Van Domselaar, G., Brinkman, F. S. L., and Hsiao, W. W. L. (2017). Context is everything: Harmonization of critical food microbiology descriptors and metadata for improved food safety and surveillance. *Frontiers in Microbiology*, 8.

[28] Haseeb, A., Saleem, Z., Faidah, H. S., Saati, A. A., AlQarni, A., Iqbal, M. S., Alghamdi, S., Elrggal, M. E., AlGethamy, M., Radwan, R. M., Mahrous, A. J., Abuhussain, S. S. A., Khayyat, S. M.,

Ibrahim, K., Godman, B., and Sheikh, A. (2023). Threat of antimicrobial resistance among pilgrims with infectious diseases during hajj: Lessons learnt from covid-19 pandemic. *Antibiotics*, 12(8):1299.

[29] Hastings, J., Owen, G., Dekker, A., Ennis, M., Kale, N., Muthukrishnan, V., Turner, S., Swainston, N., Mendes, P., and Steinbeck, C. (2015). Chebi in 2016: Improved services and an expanding collection of metabolites. *Nucleic Acids Research*, 44(D1):D1214–D1219.

[30] Jevons, M. P. (1961). "Celbenin" - resistant Staphylococci. *BMJ*, 1(5219):124–125.

[31] Jolley, K. A., Bray, J. E., and Maiden, M. C. J. (2018). Open-access bacterial population genomics: Bigsdb software, the pubmlst.org website and their applications. *Wellcome Open Research*, 3:124.

[32] Katayama, Y., Ito, T., and Hiramatsu, K. (2000). A new class of genetic element, staphylococcus cassette chromosome mec , encodes methicillin resistance in staphylococcus aureus. *Antimicrobial Agents and Chemotherapy*, 44(6):1549–1555.

[33] Klyne, G. and Carroll, J. J. (2004). Resource description framework (rdf): Concepts and abstract syntax. W3C Recommendation.

[34] Lakhundi, S. and Zhang, K. (2018). Methicillin-Resistant Staphylococcus aureus: Molecular Characterization, Evolution, and Epidemiology. *Clinical Microbiology Reviews*, 31(4).

[35] Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*, 34(18):3094–3100.

[36] Maxat, K. (2024). Metadata of methicillin resistant staphylococcus aureus (mrsa) sequences in saudi arabia.

[37] Mediavilla, J. R., Chen, L., Mathema, B., and Kreiswirth, B. N. (2012). Global epidemiology of community-associated methicillin resistant staphylococcus aureus (ca-mrsa). *Current opinion in microbiology*, 15(5):588–595.

[38] Nadig, S., Velusamy, N., Lalitha, P., Kar, S., Sharma, S., and Arakere, G. (2012). Staphylococcus aureus eye infections in two indian hospitals: emergence of st772 as a major clone. *Clinical ophthalmology*, pages 165–173.

[39] Ondov, B. D., Treangen, T. J., Melsted, P., Mallonee, A. B., Bergman, N. H., Koren, S., and Phillippy, A. M. (2016). Mash: fast genome and metagenome distance estimation using minhash. *Genome Biology*, 17(1).

[40] Page, A. J., Cummins, C. A., Hunt, M., Wong, V. K., Reuter, S., Holden, M. T., Fookes, M., Falush, D., Keane, J. A., and Parkhill, J. (2015). Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*, 31(22):3691–3693.

[41] Petit III, R. A. and Read, T. D. (2020). Bactopia: a flexible pipeline for complete analysis of bacterial genomes. *Msystems*, 5(4):10–1128.

[42] Ploug, T., Holm, S., and Gjerris, M. (2015). The stigmatization dilemma in public health policy-the case of mrsa in denmark. *BMC Public Health*, 15(1).

[43] Price, M. N., Dehal, P. S., and Arkin, A. P. (2010). Fasttree 2–approximately maximum-likelihood trees for large alignments. *PloS one*, 5(3):e9490.

[44] Sanchez-Herrero, J. F. and Mjsull (2020). spatyper: Staphylococcal protein a (spa) characterization pipeline.

[45] Schoch, C. L., Ciufo, S., Domrachev, M., Hotton, C. L., Kannan, S., Khovanskaya, R., Leipe, D., Mcveigh, R., O'Neill, K., Robbertse, B., Sharma, S., Soussov, V., Sullivan, J. P., Sun, L., Turner, S., and Karsch-Mizrachi, I. (2020). Ncbi taxonomy: a comprehensive update on curation, resources and tools. *Database*, 2020.

[46] Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, 30(14):2068–2069.

[47] Senok, A., Ehricht, R., Monecke, S., Al-Saedan, R., and Somily, A. (2016). Molecular characterization of methicillin-resistant staphylococcus aureus in nosocomial infections in a tertiary-care facility: emergence of new clonal complexes in saudi arabia. *New microbes and new infections*, 14:13–18.

[48] Senok, A., Nassar, R., Celiloglu, H., Nabi, A., Alfaresi, M., Weber, S., Rizvi, I., Müller, E., Reissig, A., Gawlik, D., et al. (2020). Genotyping of methicillin resistant staphylococcus aureus from the united arab emirates. *Scientific reports*, 10(1):18551.

[49] Souvorov, A., Agarwala, R., and Lipman, D. J. (2018). Skesa: strategic k-mer extension for scrupulous assemblies. *Genome Biology*, 19(1).

[50] Tay, M., Lee, B., Ismail, M. H., Yam, J., Maliki, D., Gin, K. Y.-H., Chae, S.-R., Ho, Z. J. M., Teoh, Y. L., Ng, L. C., et al. (2024). Usefulness of aircraft and airport wastewater for monitoring multiple pathogens including sars-cov-2 variants. *Journal of Travel Medicine*, 31(5):taae074.

[The Arvados Authors et al.] The Arvados Authors, Amstutz, P., César, N., Clegg, T., Di Pentima, L., Kutyła, D., Li, J., Smith, S., Vandewege, W., Wait Zaranek, A., and Wait Zaranek, S. Arvados.

[52] Torsten, S. (2020). Abricate.

[53] Udo, E., O'brien, F., Al-Sweih, N., Noronha, B., Matthew, B., and Grubb, W. (2008). Genetic lineages of community-associated methicillin-resistant staphylococcus aureus in kuwait hospitals. *Journal of clinical microbiology*, 46(10):3514–3516.

[54] Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., et al. (2016). The fair guiding principles for scientific data management and stewardship. *Scientific data*, 3.

[55] Wood, D. E., Lu, J., and Langmead, B. (2019). Improved metagenomic analysis with kraken 2. *Genome Biology*, 20(1).

Geolocation	Total samples	Unique CCs	Unique ST	Sccmec types rate%	PVL rate %	TSST rate %
Alhasa (East)	33	8	15	IVa 45.5; V 30.3; IVc 12.1; MSSA 6.1; VI 6.1	24.2	6
Hail (North)	64	9	19	MSSA 34.4; IVa 28.1; V 23.4; VI 6.3; IVc 4.7; IIIa 3.1	6.3	26
Jazan (South)	18	4	6	V 61.1; IVa 27.8; IVd 11.1	22.2	0
Jeddah (West)	275	9	43	V 45.1; IVa 32.4; MSSA 10.2; IVc/IVd 3.6; VI 2.9; Undetermined 0.7; VI/Vc/IVh/IIIa 0.4	13.5	8
Madina (West)	102	9	26	V 34.3; IVa 33.3; IVc 13.7; VI 6.9; MSSA 3.9; IVg 2.0; I 2.0; Others 1.0	32.4	13
Makkah (West)	36	8	14	V 41.7; IVa 38.9; MSSA 13.9; Undetermined 2.8; IVc 2.8	13.9	8
Riyadh (Central)	158	9	29	V 41.8; IVa 19.6; MSSA 18.4; IVc 8.9; IVd 4.4; VI 3.2; IVb 1.9; IVg/IVh 0.6	19.0	7

Table 1: Summary of MRSA statistics across different geolocations, including total counts, unique clonal complexes (CCs), unique sequence types (STs), SCC*mec* types rate, Panton-Valentine leukocidin (PVL) rate, and toxic shock syndrome toxin (TSST) rate.

Antibiotic	Riyadh	Jeddah	Makkah	Medina	Hail	Alhasa
Benzylpenicillin	98.3	97.4	100.0	97.4	95.6	88.9
Clindamycin	19.3	23.1	18.9	25.9	20.6	19.4
Erythromycin	27.8	25.6	37.8	37.9	25.0	19.4
FusidicAcid	70.5	70.9	62.2	60.9	35.3	46.9
Gentamicin	18.2	31.6	0.0	22.4	20.6	13.9
Levofloxacin	27.3	23.9	40.5	45.7	14.7	25.0
Linezolid	0.0	0.0	0.0	0.0	0.0	0.0
Moxifloxacin	21.6	19.3	33.3	41.5	13.4	25.0
Nitrofurantoin	0.0	0.9	0.0	1.7	0.0	0.0
Oxacillin	81.8	90.6	86.5	96.6	60.3	93.9
Rifampicin	0.0	0.9	0.0	1.7	0.0	0.0
Teicoplanin	0.0	0.0	0.0	0.0	0.0	0.0
Tetracycline	20.5	26.5	10.8	20.9	7.4	6.2
Tigecycline	0.0	0.0	0.0	0.0	0.0	0.0
Tobramycin	26.7	32.5	8.1	27.8	20.6	13.9
Vancomycin	0.0	0.0	0.0	0.0	0.0	0.0

Table 2: Antibiotic resistance rates by sampling location in percent.

Antibiotic	CARD	Resfinder	DeepARG	Concordance
Benzylpenicillin	.983/.980/.982	.994/.572/.726	.983/.980/.982	High CARD-DeepARG
Cefoxitin	1.0/.004/.007	.984/.950/.967	-	Low
Oxacillin	.858/.976/.913	-	.858/.976/.913	High CARD-DeepARG
Gentamicin	.172/.937/.291	.752/.793/.772	.172/.937/.291	Low
Tobramycin	.220/.950/.357	.785/.807/.796	.221/.950/.358	Low CARD-DeepARG
Erythromycin	.283/1.0/.441	.776/.889/.828	.584/.895/.707	Moderate
Clindamycin	.446/.861/.588	.704/.826/.760	.378/.861/.525	Moderate
Tetracycline	.167/.962/.285	.853/.771/.810	.491/.800/.609	Low
Fusidic Acid	.937/.925/.931	.940/.845/.890	1.0/.005/.010	High CARD-Resfinder

Table 3: Concordance of antimicrobial resistance predictions with the observed resistance patterns using three tools (CARD Database, Resfinder 4.0, and DeepARG). Metrics shown as Precision/Recall/F-score. Dash (-) indicates that no predictions could be made by the tool for this antimicrobial agent.