Growing Minds, Integrating Senses: Neural and Computational Insights into Age-related Changes in Audio-Visual and Tactile-Visual Learning in Children

Nina Raduner[a,b,c,d,e], Carmen Providoli[a,c,d,e], Sarah V. Di Pietro[a,e], Maya Schneebeli[a,e], Iliana I. Karipidis[a,d], Ella Casimiro[e,f], Saurabh Bedi[e,f], Michael von Rhein[c,e], Nora M. Raschle[b,d,e], Christian C. Ruff[d,e,f], Silvia Brem[a,d,e]

Affiliations:
[a] Department of Child and Adolescent Psychiatry and Psychotherapy, University Hospital of Psychiatry, University of Zurich, Zurich, Switzerland
[b] Jacobs Center for Productive Youth Development, University of Zurich, Zurich, Switzerland
[c] Child Development Center, University Children's Hospital Zurich, Zurich, Switzerland
[d] Neuroscience Center Zurich, University and ETH Zurich, Zurich, Switzerland
[e] URPP Adaptive Brain Circuits in Development and Learning (AdaBD), University of Zurich, Zurich, Switzerland
[f] Department of Economics, Zurich Center for Neuroeconomics, University of Zurich, Zurich, Switzerland

Corresponding author
Prof. Dr. Silvia Brem
Neumünsterallee 9
8032 Zürich
Switzerland
silvia.brem@uzh.ch
+41 (0)58 384 67 11

Abstract

Multisensory processing and learning shape cognitive and language development, influencing how we perceive and interact with the world from an early age. While multisensory processes mature into adolescence, it remains poorly understood how age influences multisensory associative learning. This study investigated age-related effects on multisensory processing and learning during audio-visual and tactile-visual learning in 67 children (5.7–13 years) by integrating behavioural and neuroimaging data with computational methods. A reinforcement-learning drift diffusion model revealed that older children processed information faster and made more efficient decisions on multisensory associations. These age-related increases coincided with higher activity in brain regions associated with cognitive control, multisensory integration, and memory retrieval, specifically during audio-visual learning. Notably, the bilateral anterior insula exhibited heightened activation in response to lower reward prediction errors, indicative of increased sensitivity to negative feedback with development. Finally, reward prediction errors modulated activation in reward processing and cognitive control regions, with this modulation remaining modality-independent and largely stable across age. In conclusion, while children employ similar learning strategies, older children make decisions more efficiently and engage neural resources more strongly. Our findings reflect ongoing maturation of neural networks supporting multisensory learning in middle childhood, enabling more adaptive learning in later childhood.

Keywords

multisensory learning
audio-visual and tactile-visual
reward prediction error
computational modelling
fMRI
children

Highlights

- faster information processing in older children during a multisensory learning task
- increasing brain activation with age in visual, parietal, and frontal regions
- reward prediction error processing are independent of age and modality
- heightened response to negative reward prediction errors with increasing age

1 Introduction

Efficient processing and integration of sensory information from multiple modalities including vision, hearing, and touch, are essential for interacting with our surroundings and adapting to the complexities of our environment. On a daily basis, humans encounter numerous stimuli by various senses, requiring appropriate processing and integration to be accurately understood (Van Atteveldt et al., 2014). Multisensory processing describes the perception of stimuli in multisensory environments (Murray et al., 2016). Multisensory learning (MSL) is the process of acquiring associations between stimuli from different senses (Lauzon et al., 2022), while multisensory integration (MSI) occurs when the brain combines information from multiple senses to form a unified percept (Stein et al., 2020; Wallace et al., 2020). When stimuli from different modalities provide congruent information, they are integrated and perceived as originating from the same source. Research using electrophysiology, fMRI, and behavioural studies has shown that successful associative MSL leads to MSI (Butler et al., 2011; Lauzon et al., 2022). Since these processes are sequential (multisensory processing is needed for MSL which is also the basis for MSI), they are often studied in one framework. MSI is particularly significant in naturalistic learning environments and classrooms, where multisensory exposure is common (Barutchu et al., 2019). Moreover, MSI skills are associated with enhanced cognitive processes including memory performance and reading abilities (Barutchu et al., 2011, 2019; Birch & Belmont, 1965; Denervaud et al., 2020; Dionne-Dostie et al., 2015). MSI likely promotes learning by enabling the effective combination of sensory inputs for better encoding, storage, and retrieval of information. For example, MSI underpins language learning, production, and development by integrating auditory and visual information like sounds and lip movements (Wallace et al., 2020).

Given the importance of MSL and MSI in cognitive development, it is crucial to investigate MSL in childhood and how it is similar or different across different sensory modalities. However, most studies on MSL have focused on adults and the studies investigating MSL in children mainly concentrated on audio-visual speech sound mapping (e.g., Frei et al., 2025; Karipidis et al., 2021; Wang et al., 2020). Only a few investigations targeted the learning and processing of non-linguistic audio-visual integration (e.g., Altarelli et al., 2020; Batson et al., 2011; Pasqualotto et al., 2024; Seitz et al., 2007). Moreover, while MSL can encompass signals from all senses, minimal attention has been directed towards the interplay between touch and vision or any other combination of stimulus modalities (see meta-analysis from Gao et al., 2023).

Despite the growing interest in and the importance of MSL, relatively few studies have examined its neural underpinnings (e.g., Butler et al., 2011; Lauzon et al., 2022), with most research focusing on MSI or failing to distinguish between the two processes. The meta-analysis by Gao et al. (2023) further highlights that audio-visual integration has been widely studied, whereas only a few studies examined tactile-visual integration (such as Gentile et al., (2013, 2015); Limanowski & Blankenburg, (2016, 2017)). In the mentioned meta-analysis, audio-visual integration was associated with changes in neural activation of early sensory regions, subcortical areas, and higher association areas supporting a flexible neural pathway model for audio-visual integration. Such a model is supported by research linking MSI and learning to an extended cortico-striato-thalamic network (Lakatos et al., 2007; Murray et al., 2016; Tyll et al., 2011), including early sensory regions (Ghazanfar & Schroeder, 2006; Lakatos et al., 2007; Noesselt et al., 2007; Stein & Stanford, 2008). Moreover, reviews summarising brain regions involved in MSI identified similar regions to be relevant for these processes: the auditory cortex, visual cortex, somatosensory cortex, superior temporal cortex, and frontal regions (Murray et al., 2016; Richlan, 2019). Exploratory meta-analyses on the few existing

studies on tactile-visual integration reveal that inferior temporal areas, the fusiform gyrus, and
inferior parietal areas are associated with tactile-visual integration (Gao et al., 2023). This
highlights the need for further research on the brain areas underlying MSL, particularly during
childhood, when these abilities are still developing.
Although infants already demonstrate the ability to process and integrate multisensory
information (Patterson & Werker, 1999, 2003), a deeper understanding of how these abilities
develop over childhood is crucial for advancing learning research. MSI has not yet reached full
maturity in children younger than eight years old and maturation of MSI skill continues into
adolescence (Desjardins et al., 1997; Dionne-Dostie et al., 2015; Gori et al., 2008; Nardini et
al., 2008). Research suggests the temporal window for audio-visual integration is still immature
in 11-year-old children (Dionne-Dostie et al., 2015). According to Murray et al. (2016),
children's MSI abilities are initially more reliant on low-level stimulus feature processing but
gradually shift toward higher stimulus feature processing and therefore cortical involvement.
This process reflects the brain's developmental trajectory where subcortical regions mature
before cortical regions. This implies that younger children, who depend more on the processing
of low-level features through subcortical and primary sensory areas, might have limited MSI
capabilities for complex learning tasks requiring greater levels of cognitive control. With
children's increasing age, however, the integration of sensory information progressively
involves higher-order brain regions, such as the frontal cortex. This in turn allows for more
flexible and sophisticated MSI that is crucial for advanced learning, problem-solving, and
adaptation in dynamic environments. As a result, these maturation processes allow for more
complex multisensory learning and integration as children grow (Murray et al., 2016).
To better understand the developmental processes, computational modelling provides
a powerful tool for examining the cognitive mechanisms underlying MSL, deepening the neural
and developmental insights. One approach in computational modelling integrates a learning
process, reflecting trial-wise belief updating, and a choice process, translating computational
quantities (e.g., values) into behavioural or neural responses (Mathys et al., 2011). This study
employs a combined reinforcement learning model as the learning process and drift diffusion
model as the choice process (RLDDM; Pedersen et al., 2017) to investigate multisensory
processing and learning in children (see also Fraga-González et al., 2025; Frei et al., 2025).
Reinforcement learning, one of the core mechanisms for feedback learning, compares
observed outcomes (rewards) with expected outcomes (values) and adjusts expectations
based on the discrepancy – known as the reward prediction error (RPE; Niv & Schoenbaum,
2008; Subramanian et al., 2022). The Rescorla-Wagner (RW) model describes this process
by weighting the RPE with the learning rate to update values (Rescorla & Wagner, 1972). On
the other hand, the drift diffusion model quantifies the decision-making process through the
drift rate, decision boundary, and non-decision time (Pedersen et al., 2017; Ratcliff, 1978;
Ratcliff & Rouder, 1998). By combining these models, we can identify latent cognitive
processes like value updating, learning rates, and RPEs, and relate them to the brain activation
during learning.
Building on these computational frameworks, previous research has identified specific
neural correlates of the cognitive parameters involved in learning. RPEs are consistently linked
to stronger brain responses in the striatum, the medial frontal cortex, and hippocampus
reflecting their relation to learning and memory (Cohen & Ranganath, 2005; Fraga-González
et al., 2025; Garrison et al., 2013; McClure et al., 2003; O'Doherty et al., 2003; Pagnoni et al.,
2002). Similarly, value processing involves regions such as the ventral striatum, anterior
cingulate cortex, anterior insula, medial orbitofrontal cortex, supplementary motor area, and
inferior parietal lobe (Liu et al., 2011; R. P. Wilson et al., 2018). These regions are pivotal in
integrating motivational and emotional aspects of decision-making (Liu et al., 2011). The

learning rate has been linked to activation in the inferior frontal gyrus, anterior insula, dorsal anterior cingulate cortex, and ventral caudate (Chien et al., 2016; O'Reilly, 2013; Wu et al., 2017). By mapping neural correlates of computational parameters such as values, reward prediction errors, and learning rates, our study aims to computationally characterize the perceptual and learning processes underlying MSL in children and study their change during development.

To further explore these neural and cognitive processes, the current study investigated MSL in middle childhood, focusing on learning sensory associations of different modalities (audio-visual vs. tactile-visual) and on age-related changes therein. The primary aims were: 1) to **investigate modality-specific differences in children's multisensory processing and learning** by comparing audio-visual (AV) and tactile-visual (TV) associations, using computational modelling of behavioural performance and neural activity data, 2) to characterize the **age-related changes in MSL** by analysing age-related differences in processing speed, decision-making and associated neural activity, and 3) **to examine how children process feedback during MSL**, specifically looking at age-effects on RPE and value-based learning for audio-visual and tactile-visual learning. We hypothesised that younger children would rely more on subcortical and primary sensory regions, while older children would engage higher-order cortical areas during MSL, indicating the increased employment of neural resources (Murray et al., 2016). Accordingly, we anticipated neural activation in brain regions previously associated with value processing (Liu et al., 2011; R. P. Wilson et al., 2018), reward prediction error processing (Cohen & Ranganath, 2007; Fraga-González et al., 2025; Garrison et al., 2013; McClure et al., 2003; Pagnoni et al., 2002), and learning rate processing (Chien et al., 2016; O'Reilly, 2013; Wu et al., 2017). We expected potential age-dependent variations likely to be pronounced in higher-cortical areas such as prefrontal and cingulate regions (Yaple et al., 2020). With this study, we bridge the gap in the existing literature by investigating non-linguistic audio-visual and tactile-visual associative learning across middle childhood.

We employed a cross-sectional design with 67 children (aged 5.7–13.0 years) completing a multisensory learning task involving AV and TV stimuli. Computational modelling, incorporating reinforcement learning and drift diffusion frameworks, was applied to characterise the cognitive processes underpinning learning. Functional MRI was employed to pinpoint brain regions engaged in modality-general and modality-specific learning processes and to characterize age-related changes in the activation of involved brain networks. Parametric modulation of the BOLD signal with model parameters finally allowed us to examine the neural correlates underlying specific learning and decision processes, as well as potential age-related changes in the neural processing.

2 Material and methods

2.1 Participants

This study included 80 healthy, native (Swiss-)German-speaking children. Children with either neurological or psychiatric disorders or intellectual disabilities (except for specific learning disorders or ADHD) were not included to participate in this study. Following the exclusion of one subject due to technical problems during scanning, another two subjects had to be excluded due to excessive movement in all runs (see chapter 2.3) and a high number of omissions (see Supplementary Material A.6). Additionally, 10 children were excluded from the final analysis as they had only one AV or TV run with sufficient MR data quality (no excessive movement). Consequently, the final sample comprised 67 children (M = 8.35 ± 1.82 years, 36 girls) with at least one high-quality run in both the AV and TV modalities. There was no significant age difference between girls and boys (see Supplementary Material A.1). All children had estimated IQ scores above 80. All participants had normal or corrected to normal vision and normal hearing. Mean IQ, processing speed, and attention scores were in the normal range (see **Table 1**; a table summarizing the demographics of the group split in 3 age bins [5.7-8.0, 8.1-10.5; 10.6-13] can be found in the Supplementary Material A.1), indicating age-appropriate cognitive abilities for all subjects (see Supplementary Material A.1 for more information). A total of ten children had either a clinical diagnosis (attention deficit hyperactivity disorder, developmental dyslexia, or developmental language delay) and/or clinically significant values in the CBCL questionnaire (Döpfner et al., 2014; see Supplementary Material A.1 for more information). The following analyses were conducted with and without these ten participants (see Supplementary Material B.1). This approach ensures that the results are not biased by potential neurodevelopmental differences while also allowing for a more comprehensive analysis that includes all participants.

Oral consent was obtained from all children, while written consent was provided by a parent or guardian for all children. The study was approved by the local ethics committee of the Canton of Zurich and neighbouring cantons in Switzerland (BASEC-Nr: 2022-01368), and the participants received vouchers and gifts in appreciation of their participation.

Table 1

Demographic characteristics of study participants

	M (*SD*) [range]
Age [y]	8.35 (1.82) [5.7-13]
Sex [f:m]	36:31
Handedness[a] [l:a:r]	6:3:57
IQ (verbal and non-verbal)	106.5 (10.6)
Rapid automatized naming [T-value]	50.1 (7.45)
Processing speed [T-value]	50.12 (7.97)
Sustained attention median RT[b] [PR]	51.55 (29.43)

Note. The handedness of the subjects was determined using the Edinburgh Handedness Scale (Oldfield, 1971). IQ values were estimated using the Peabody Picture Vocabulary Test – fourth edition (PPVT-IV; Lenhard et al., 2015) and subtests from the Wechsler Intelligence Scale for Children – fifth edition (WISC-V; Wechsler, 2017) or the Wechsler Preschool and Primary Scale of Intelligence – fourth edition (WPPSI-IV; Wechsler, 2012). Rapid automatized naming and processing speed measured with IDS-2 (Grob & Hagmann-von Arx, 2018). Sustained attention measured with KiTAP (Zimmermann et al., 2002). [a] One child's data are missing. [b] Percentiles are missing for two children due to the absence of normative values for their age. y = years, f = female, m = male, l = left, a = ambidextrous, r = right, RT = reaction time, PR = percentiles.

2.2 Experimental design

During the neuroimaging session, participants completed an audio-visual and tactile-visual multisensory learning task (adapted from Frei et al. (2025)) embedded in a child-friendly story in the MRI scanner. The multisensory learning task is a two-alternative forced choice task where the participants had to learn multisensory associations between visual and auditory or visual and tactile stimuli based on feedback. Participants were presented with two symbols alongside either a sound or vibration pattern, representing a rocket's sound or engine trembling, respectively. Participants guessed which rocket the sound or trembling belonged to by pressing a corresponding button. They were told to learn associations over time through feedback, which was implicitly suggested to be probabilistic, and to collect as many stars as possible (rewarded after a correct response; incentive feedback) during each run. The task was presented using goggles (CinemaVision 20/20, Resonance Technology, Northridge, CA), MR-compatible over-ear headphones (MR confon GmbH, Magdeburg, Germany, http://www.mr-confon.de/de/), a tactile stimulation device (mini-PTS piezoelectric stimulator device from dancer design, http://www.dancerdesign.co.uk/) stimulating the right pinkie, and the PsychoPy® software (Version 2022.1.1, Peirce et al. (2019); https://www.psychopy.org/). To ensure that each child could perceive the stimuli clearly during the MR session, the sound volume was adjusted, and vision was corrected individually as needed.

Not all children were able to complete every task according to the experimental protocol and therefore runs had to be repeated or excluded due to excessive movement, lack of motivation (too many omissions, see Supplementary Material A.6), or fatigue. For the purposes of this analysis, only data from children who completed at least one audio-visual and one tactile-visual run with adequate data quality were included. The 67 participants included in these analyses finally showed the following distribution across the task modalities: 18 children completed two runs of each modality, 39 completed only one run of each, 5 completed two AV and one TV run, and 5 completed one AV and two TV runs.

As depicted in **Figure 1**, during the inter-trial interval (ITI), the task background consisted of a space image (dark background with stars), two blank rockets on either side of a larger star in the middle, acting as the fixation cross. These elements were constantly displayed during the whole task. During stimulus presentation, one auditory stimulus was presented simultaneously with two visual stimuli appearing on either rocket for a duration of 2 seconds (**Figure 1**). In TV runs, a tactile stimulation was presented instead of an auditory sound (see Supplementary Material A.2). The participants were then instructed to choose the matching symbol by pressing the left or right button on an MR-compatible two-button response box for the left or right choice, respectively, and their choice was indicated by a blue frame during the inter-stimulus interval (ISI). The participant's choice was recorded during stimulus presentation and the ISI. After the response, they received feedback on their choice for 1.8s, which was followed by an ITI before the next trial began. The feedback in the task was probabilistic, with correct (e.g., positive feedback after a correct choice) feedback in most trials, and some trials providing incorrect (e.g., negative feedback after a correct choice; see Supplementary Material A.3 for more information). Neutral feedback was given after no choice was observed (see Supplementary Figure A.2B). The ITI and ISI were gamma distributed and had a mean duration of 1.87s [0.76 – 3.45s] and 3.45s [2.28 – 4.83s], respectively. After the first six participants, we slightly changed the timing of the ISI and ITI to provide more decision time. For these participants, the mean durations were 2.06s [0.79 – 3.39s] and 2.67s [1.11 – 4.34s], respectively for the ITI and ISI. Each run consisted of 44 trials and total task duration was 401 seconds (384 seconds for the first 6 subjects).

In total, there were four visual stimuli and four auditory or tactile stimuli per run, resulting in 16 possible combinations of which four were matching combinations. For the MR runs, we prepared six sets of visual, three sets of auditory stimuli, and three sets of tactile stimuli (see Supplementary Material A.2). As visual stimuli, artificial symbols from the PseudoSloan font were used (Vildavski et al., 2021). The pseudo letters were chosen by the complexity scores provided by Vildavski et al. (2021). Each set was matched such that it included one stimulus with low ($M = 3.55$, $SD = 0.02$), two stimuli with medium ($M_1 = 4.27$, $SD_1 = 0.09$ and $M_2 = 4.49$, $SD_2 = 0.05$) and one stimulus with high complexity ($M = 5.16$, $SD = 0.05$). Additionally, we also balanced the selection of pseudo letters between sets. For the audio-visual run, environmental sounds were used as auditory stimuli, which were selected from the mixkit sound effects database (https://mixkit.co/free-sound-effects/). Here, we chose sounds that could not be easily identified or named. After selection, the sounds were cut to a 2 second duration and normalized using Audacity® recording and editing software (version 3.5.1; Team Audacity, 2024). The mean Shannon spectral entropy (Cover & Thomas, 2005; Misra et al., 2004; Obin & Liuni, 2012), a measure of sound complexity, was calculated for each stimulus using the spectralEntropy function (Signal Processing Toolbox, version 9.1) in MATLAB (version R2022b; MathWorks Inc, 2022). Each set of environmental sounds consisted of one sound with a low ($M = .327$, $SD = 0.03$), two with a medium ($M_1 = .476$, $SD_1 = 0.05$ and $M_2 = .564$, $SD_2 = 0.04$), and one with a high entropy value ($M = .705$, $SD = 0.04$). Lastly, we created three different sets for the tactile stimulation. Each set of tactile stimuli consisted of four different vibration patterns lasting for 2 seconds. Each set contained two stimulations with distinct stable frequencies (low complexity), one was increasing or decreasing (medium complexity), and one had an (inverted) u-shape (high complexity; see Supplementary Figure A.2C). Ensuring that the stimuli varied based on complexity within a set was done to increase the discriminability between stimuli. The following variables were pseudo-randomised for all participants: the order of tasks (AV and TV) in the MR scanner, the trial structures (the sequence of stimulus presentation), the visual, auditory, and tactile sets used, and the assignment of the stimuli to the four correct combinations for each run.

Each participant completed behavioural runs (one for each modality) in a behavioural assessment session 1 – 42 days ($M = 10.6$ days, $SD = 8.1$ days) before the MR session (pre runs). During the behavioural session, each participant was walked through 13 instruction trials for the AV run by the examiner, afterwards they completed 10-20 practice trials on their own until the participant understood how the task works. Then they completed the first full AV behavioural run with 44 trials. Afterwards, the 13 instruction trials were repeated for the TV runs to familiarise the participants with the tactile vibration. If necessary, the 10-20 practice trials were also repeated, until the participants understood the task. Finally, each participant competed a full TV behavioural run with 44 trials. This procedure ensured that the participants were familiar with the task design before the MR session. The performance from the pre runs was then used to adapt the task difficulty in the MR scanner to ensure that the task was neither too easy nor too difficult (see Supplementary Material A.3 for adaptation criterions). Before entering the MR scanner, the task instructions were repeated to each participant, and they completed another 10-20 practice trials with the response box used in the MR. Therefore, each participant completed at least 134 trials of the multisensory learning task before entering into the MR. Importantly the stimulus material used for the introduction, practice, and pre-runs was different from the material used in the fMRI runs. Task difficulty was either adapted by the trial structure or by the level of probabilistic feedback. Additionally, some of the stimuli triplet combinations were presented more often than others (see Supplementary Material A.3 for more information about trial structure, probabilistic feedback, and stimulation frequency).

Figure 1

Multisensory learning task

Note. Illustration of the multisensory learning task showing one trial for the audio-visual (AV) and tactile-visual (TV) task with presentation durations. Two rockets, each displaying a symbol, were presented, while either a sound (AV) or a vibration pattern (TV) was played. Children then determined which rocket the sound or vibration belonged to by button press. Their chosen rocket was highlighted with a blue frame. Feedback was afterwards provided via a yellow star in the centre of the screen, accompanied by a checkmark for correct choices. Incorrect selections were indicated by a grey star and a red cross. After omissions, a double sided, yellow arrow was displayed, indicating no choice, and reminding the participants to response faster in future trials. See supplementary material A.2 for a depiction of all feedback stimuli.

2.3 MRI data acquisition and preprocessing

MRI data was acquired on a Philips Achieva 3 Tesla scanner (Best, The Netherlands) using a 32-channel coil. T2* weighted functional images were acquired with a whole-brain gradient echoplanar image (EPI) sequence (five dummy scans followed by 291 dynamic scans, repetition time TR = 1.395 s, echo time TE = 35 ms, 44 slices, voxel size = 3.0 x 3.0 x 3.0 mm3, slice gap = 0.3 mm, matrix size 64 x 62 px, flip angle = 80°, multiband factor = 2, sofTone factor = 2, sensitivity-encoding (SENSE) reduction factor = 2). Due to changes in the timing of the task (see chapter 2.2), the EPI sequence of 6 participants only included 278 dynamic scans while all the other parameters were the same. T1-weighted anatomical images were obtained for co-registration using a magnetization-prepared rapid acquisition gradient echo (MPRAGE) sequence (TR = 6.96 ms, TE = 3.2 ms, aligned at AC-PC, flip angle = 9°, voxel size = 1.0 x 1.0 x 1.0 mm^3, field of view = 270x255 mm^2, number of slices = 176).

MATLAB (version R2020b, MathWorks Inc, 2020) toolbox SPM12 (7219, Penny et al., 2007) was used for preprocessing and whole-brain analysis, which included slice-time correction, realignment, co-registration, and segmentation. Using the Template-O-Matic toolbox (Wilke et al., 2008), a paediatric template was created for normalization (age matched structural data). Voxels were resampled to 3 x 3 x 3 mm^3 isotropic voxels. An 8-mm full width at half maximum (FWHM) Gaussian kernel was applied for smoothing. After preprocessing, we repaired volumes with a composite scan-to-scan motion above 1.5mm (Euclidean distance with a radius of 65 mm) with the ArtRepair toolbox (Mazaika et al., 2007), using linear interpolation between the nearest unrepaired scans. After this, we flagged volumes with and without movement surrounded by scans with motion above 1.5mm. Runs with more than 20% of flagged volumes were excluded from the final analyses, resulting in a total of 18 runs being omitted (movement was not correlated with age; see Supplementary Material B.9). Whenever possible, we attempted to repeat runs with excessive movement during the MR session. Flagged scans were accounted for by including one binary regressor of no interest for each flagged volume (scrubbing).

In addition, functional data were denoised using a standard denoising pipeline (Nieto-Castanon, 2020) in CONN (Whitfield-Gabrieli & Nieto-Castanon, 2012) release 22.v2407 (Nieto-Castanon & Whitfield-Gabrieli, 2022) and SPM (Penny et al., 2007) release 12.7771 including the regression of potential confounding effects characterized by white matter time-series (5 CompCor noise components), CSF time-series (5 CompCor noise components), movement regressors and their first order derivatives (12 components), session and task effects and their first order derivatives (6 factors), and linear trends (2 factors) within each functional run, followed by bandpass frequency filtering of the BOLD time-series (Hallquist et al., 2013) between 0.008 Hz and 0.09 Hz. CompCor (Behzadi et al., 2007; Chai et al., 2012) noise components within white matter and CSF were estimated by computing the average BOLD signal as well as the largest principal components orthogonal to the BOLD average within each subject's eroded segmentation masks. The 10 CompCor noise regressors for white matter and CSF were then added as regressors of no interest to the first-level GLMs.

2.4 Computational modelling of values, RPEs, and learning rate

For the learning process, we implemented the trial-wise updating mechanism using a Rescorla-Wagner (RW) model which is an established model for reinforcement learning (Rescorla & Wagner, 1972). To model basic multisensory processing involved in constructing the computational quantities of the mental model and the choice processes that translate them into responses, a drift diffusion model (DDM; Ratcliff, 1978) was implemented based on the approach outlined by Pedersen et al. (2017). This generative model allowed us to estimate parameters of the trial-wise learning process, inferred from the accuracy and reaction times of the responses from each trial (Pedersen et al., 2017). This approach permitted the calculation of trial-wise values for each multisensory association, drift rates, and RPEs.

The model used in this study is illustrated in **Figure 2**. The RW model is used to calculate the RPEs and update the values V associated with the responses to the current multisensory association presented in each trial. The RPE for a trial t is defined as the difference between the actual reward R and the expected reward (= value V):

$$RPE_t = (R_t - V_t) \tag{1}$$

Then, the value V for each trial t was updated by the RPE weighted by the learning rate η:

$$V_t = V_{t-1} + \eta * (R_{t-1} - V_{t-1}) \tag{2}$$

Therefore, the learning rate is a weighting factor that indicates the extent to which values are updated after each trial (for a more detailed interpretation, see Zhang et al. (2020)). In this study, only the value for the chosen response to the specific multisensory stimulus combination was updated.

Then for each trial t, the values for the left $\bar{V}_{left,t}$ and right $\bar{V}_{right,t}$ stimulus combination were normalised to ensure that the sum of both values was 1 as seen in formula 3. The value for the left stimulus $V_{left,t}$ combination denotes the reward expectation for the multisensory stimulus combination of the auditory/tactile stimulus and the visual symbol displayed on the left side in trial t. $V_{right,t}$ denotes the multisensory stimulus combination of the auditory/tactile stimulus and the visual symbol displayed on the right side in trial t.

$$\bar{V}_{left,t} = \frac{V_{left,t}}{(V_{left,t} + V_{right,t})}, \bar{V}_{right,t} = 1 - \bar{V}_{left,t} \tag{3}$$

These normalised values from the RW were then entered in the DDM to calculate the drift rate v by multiplying the difference between the left and right value with the drift weight v_{mod}:

$$v_t = \left(\bar{V}_{left,t} - \bar{V}_{right,t}\right) * v_{mod} \tag{4}$$

Therefore, the drift weight is a scaling factor and corresponds to the sensitivity to value differences (e.g., with a high drift weight even a small difference between the values leads to high drift rate; Pedersen et al. (2017)). Larger differences between the two expected values lead to higher drift rates and therefore faster information accumulation (e.g., when one option has a clearly higher reward expectation), while information accumulation is slower, when the options have comparable reward expectations. We used the wiener first passage time (WFPT) algorithm (Navarro & Fuss, 2009) to fit the choice behaviour and reaction time data:

$$RT(x) \sim WFPT[a, \tau, z, v(t)] \tag{5}$$

With this algorithm, the probability density for hitting the upper boundary (left choice) or lower boundary (right choice) at any given time point t could be calculated:

$$f(t|v, a, z) = \frac{\pi}{a^2} \exp\left(-vaz - \frac{v^2 t}{2}\right) \sum_{k=1}^{\infty} k \exp\left(-\frac{k^2 \pi^2 t}{2a^2}\right) \sin\left(k\pi z\right) \tag{6}$$

This returns the probability of choosing x given the observed reaction time and the expected reward. For further details, please refer to the paper by Pedersen et al. (2017).

Figure 2

Schematic of the reinforcement learning drift diffusion model

Note. Deterministic variables are represented by rectangular nodes, fitted parameters by round nodes with solid lines, fixed parameters by round nodes with dashed lines, and observed data by grey round nodes. The parameters are estimated for each run separately. Learning rates η modulate the trial-wise values $\bar{V}$ which are updated depending on the feedback. The difference between the value of the left and right stimulus pair is multiplied with the v_{mod} parameter to estimate the trial-wise drift rate v. Parameters are estimated for each run r. a = decision boundary, τ = nondecision time, z = starting point (fixed parameter), R = feedback, t = trial, p = audio-visual or tactile-visual stimulus pair, v = drift rate, RT = reaction time, r = run.

In our approach, the learning rate η was a free parameter whereas the initial values for all stimulus combinations were fixed to 0.5 (uniform unbiased initial values). The DDM included three additional free parameters: the non-decision time τ, the boundary separation a, and the drift weight v_{mod}. The starting point z in the DDM was fixed to $0.5 * a$, indicating no bias towards any of the decisions. The boundary separation indicates the amount of information required until a decision criterion is reached. The non-decision time represents the amount of time that

is irrelevant for the decision-making process, such as the time required for response execution.
The drift rate v indicates the rate of accumulation of information. In the approach used in this
study, free parameters (learning rate, boundary separation, non-decision time, and drift weight)
were estimated separately for each run of 44 trials. The estimated parameters were then used
to obtain trial-wise values for the RPE, values, and drift rate, which were used to capture the
dynamic nature of learning and decision making.
Prior to applying the model to the dataset, we conducted simulations to verify that all
model parameters could be recovered successfully (see Supplementary Material A.5). Then,
the validated model was used for parameter estimation on the collected behavioural data and
the estimated parameters were then used to identify corresponding neural processes (Farrell
& Lewandowsky, 2018; R. C. Wilson & Collins, 2019). This approach ensured that our models
accurately captured the underlying cognitive processes of multisensory learning. The model
fitting pipeline was set up in MATLAB using the Genetic Algorithm (ga) from the Global
Optimization Toolbox, in accordance with the recommendations set forth by R. C. Wilson &
Collins (2019).

2.5   Statistical analyses

2.5.1   Analyses of task performance and computational parameters

As detailed in the supplementary materials (A.6), preprocessing of the behavioural data
involved the exclusion of trials with omissions, and reaction times that were either too fast or
too slow, resulting in a final set of 7,594 valid trials for analysis.
To evaluate learning outcomes and differences between modalities, mean reaction
times for correct trials, mean accuracy scores (correct responses divided by the total number
of trials not counting omissions), and mean absolute drift rate were calculated for each bin and
modality, where one bin represented one-fourth of the task (11 trials). Reaction times,
accuracies, learning rate, non-decision time, boundary separation, and the absolute drift rate
were then analysed using linear mixed models (library 'lmerTest', Kuznetsova et al., 2017) with
subject as a random effect and modality and age as fixed effects. The model for the reaction
times, accuracies, and absolute drift rate also included bin as fixed effect. Post-hoc analyses
for significant main effects with more than one level were computed (library 'emmeans'; Lenth,
2024) and p-values were adjusted for multiple comparisons by the Tukey method. Analyses
were conducted using the R Statistical language (version 4.4.1; R Core Team, 2024). Effect
sizes (η^2_P) are interpreted using the following guidelines: values < .01 are considered very
small, values from .01 up to (but not including) .06 are small, values from .06 up to (but not
including) .14 are medium, and values > .14 are large (Field, 2013).

2.5.2   Whole-brain analysis

For the first-level analysis, we built a general linear model (GLM) using the individual
onsets of two regressors of interest (stimulus, feedback) and the two parametric modulators
for value and RPE. The value modulator was added on the stimulus regressor and included
the normalised value for the chosen stimulus combination. The RPE modulator was added on
the feedback regressor and included the trial-wise RPE values. We added 4 additional
regressors of no interest: 1) stimulus onset for RT outlier trials, 2) feedback onset for RT outlier
trials, 3) stimulus onset for omission trials, and 4) feedback onset for omission trials to the
GLM. The GLM also included six realignment parameters and one additional regressor of no
interest for each flagged scan, whenever available. This was done to account for movement
effects (scrubbing). The GLM included all available AV and TV runs for each subject and was
convolved with the canonical hemodynamic response function implemented in SPM12.

Whole-brain second-level one-sample t-tests investigated the activation during stimulus processing, feedback processing, value processing, and RPE processing. Additionally, age was added as an effect of interest to all contrasts. For the contrasts investigating stimulus processing, the learning rate was added as an additional effect of interest (if more than one run was included, the mean learning rate from all included runs was used). Further, to investigate main and interaction effects between age and modality (AV and TV), a mixed-effects ANCOVA was conducted. The design matrix included the scans for each condition and participant. Age, condition, and their interaction were included as regressors of interest, where age is a continuous and condition a factorial variable. The model accounted for variability within subjects by including one regressor per subject. These analyses additionally included the covariates sex and handedness to control for potential confounds. The effect of age, learning rate, and the described contrasts were used to investigate general brain activity during multisensory learning and how the activation differed between modalities (AV and TV) and in relation to age. Statistical parametric maps were thresholded at a voxel-wise $p < .001$ (uncorrected), with clusters corrected for multiple comparisons using a family-wise error (FWE) rate of $p < .05$, unless stated otherwise. With this threshold, the likelihood of Type I error rates in parametric cluster correction can be reduced to an acceptable level (see Eklund et al. (2016)).

2.5.3   Region of interest analyses

For region of interest (ROI) analyses, the same first-level outputs were used as for the whole-brain analyses. Masks for multisensory regions, value processing and RPE processing regions were created using neurosynth (www.neurosynth.org). We downloaded the uniformity test activation mask for each topic and then processed each mask to extract all the subclusters as individual masks (a detailed description of the processing can be found in the Supplementary Material A.7). With the code provided by MarsBaR (https://marsbar-toolbox.github.io/faq.html), we extracted the beta values corrected for the mean signal in each ROI (Pernet, 2014). We extracted the mean beta values over all subclusters and for each subcluster separately from each mask. The "multisensory network" mask was used with the AV stimulus and TV stimulus contrasts, the "value network" mask with the AV value and TV value contrasts, and the "reward prediction error network" mask with the AV RPE and TV RPE contrasts. In a next step, these beta values were then analysed using linear mixed models in R to investigate differences between modalities and the effect of age. First, we looked at the effects of modality and age and their interaction on the whole mask (beta value ~ modality × age). Then we applied the same linear mixed model on each subcluster. In the multisensory network ROI, the learning rate was included as an additional main effect of interest. Age and learning rate were centred to the mean and the modality factor was orthogonalized. To correct for multiple comparisons, all the p-values from each linear mixed model were extracted and corrected using the False Discovery Rate (FDR) procedure (Benjamini & Hochberg, 1995). The FDR correction was applied with a significance level of $\alpha = 0.05$. This allowed us to control for the proportion of false positives among the test results.

3 Results

3.1 Behavioural results

General Task Performance: the mean reaction times for AV and TV runs were 2.26s (*SD* = 0.37s) and 2.49s (*SD* = 0.35s), respectively. The median accuracy for AV and TV runs over all bins were 65.91% (*IQR* = 24.70%) and 65.12% (*IQR* = 23.58%), respectively. Bin-level accuracy inspection revealed that, with the exception of two participants, all individuals exceeded the chance threshold (≥ 63.64%) in at least two bins across all runs. Further accuracy at the end of the task (bin 4), showed a median of 72.73% correct for both AV and TV runs. As expected in a learning task, these results clearly exceeded chance level. To enable a valid comparison of learning across modalities, we did not exclude participants, who surpassed chance performance in only one of the two modalities. An overview including omissions and outliers can be found in the supplementary material B.2.

Linear mixed models were computed to examine the effect of sensory modality (AV vs TV), bin (1, 2, 3, 4) and age on reaction times and accuracy.

Reaction Time: Residuals from the linear mixed-effects model on raw RT showed no significant deviation from normality (DHARMa KS p = .105; dispersion p = .856), whereas the log-transformed model exhibited significant quantile and outlier deviations (DHARMa KS p < .001; dispersion p = .856). Accordingly, all inferential statistics reported below are based on the untransformed RT model. The best fitting LMM for reaction times (reaction time ~ modality + bin + age + modality:age + (1 | participant)) showed no significant main effect of modality ($F(1,464)$ = .06, p = .814; η^2_p = .0001, 95% CI [.00, 1.00]), but a significant main effect of bin ($F(3,464)$ = 8.83, p < .001; η^2_p = .05, 95% CI [.02, 1.00]), and age ($F(1,65)$ = 10.66, p = .002; η^2_p = .14, 95% CI [.04, 1.00]), and a significant interaction between modality and age ($F(1,464)$ = 5.59, p = .018; η^2_p = .01, 95% CI [.001, 1.00]). Adding the interactions between modality and bin, age and bin, and the three-fold interaction between modality, age, and bin did not significantly improve model fit (ps > .0854). Post-hoc pairwise comparisons between the bins can be seen in Table B2.2 in the Supplementary Material. The results show significantly faster RTs in the third and last bin compared to the first and second bin. RTs were also faster the older the children, but this effect was weaker in TV compared to AV runs (see **Figure 3A** and **C**).

Accuracy: For the accuracy, the best fitting LMM (accuracy ~ modality + bin + age + modality:age + (1 | participant)) showed significant main effects of modality ($F(1,464)$ = 6.70, p = .010; η^2_p = .01, 95% CI [.002, 1.00]) at the intercept (age = 0), bin ($F(3,464)$ = 13.85, p < .001; η^2_p = .08, 95% CI [.04, 1.00]), and age ($F(1,65)$ = 10.51, p = .002; η^2_p = .14, 95% CI [.03, 1.00]), and a significant interaction between modality and age ($F(1,464)$ = 8.79, p = .003; η^2_p = .02, 95% CI [.004, 1.00]). Adding the interactions between modality and bin, age and bin, and the three-fold interaction between modality, age, and bin did not significantly improve model fit (ps > .163). Post-hoc pairwise comparisons between the bins can be seen in Table B2.2 in the Supplementary Material. The results show significantly higher accuracies in the third and last bin compared to the first and second bin, and with increasing age of the children, although this effect is less strong in TV runs compared to AV runs (see **Figure 3B** and **D**). Although the uncentered model estimated a modality effect at the intercept (age = 0), this effect became non-significant when age is mean centred prior to fitting the LMM, while all other effects remained unchanged.

Figure 3

Reaction time and accuracy per modality and task bin

Note. Panel **A** shows the reaction times per task and bin of task on the x-axis. **B** shows the accuracy per task and bin on the x-axis. The dashed and dotted lines connect the mean values for reaction times and accuracies per bin for AV and TV runs, respectively. **C** shows the interaction effect between age and modality on the reaction times. **D** shows the interaction effect between age and modality on the accuracies. AV = audio-visual (in blue), TV = tactile-visual (in green).

3.2 Modelling results

Linear mixed models were conducted to examine the effect of modality (AV vs. TV) and age on the learning rate, the non-decision time, and the boundary separation. For the drift rate, we fitted the same model as for the reaction time and accuracy, with modality (AV vs. TV), bin (1, 2, 3, 4), and age as predictors.

Learning rate: Mean learning rates were 0.19 (*SD* = 0.26) in AV and 0.20 (*SD* = 0.26) in TV runs. The best fitting LMM included only an intercept (and random intercepts for participants) which was significantly greater than zero (β = 0.20, 95% CI [.15, .24], t(131) = 8.34, p < .001). Adding the main effect of either modality and age, or the interaction between the two, did not improve model fit (ps > .271). These results indicate that learning rates were significantly different from 0, but did not vary with age or differ between AV and TV learning.

Non-decision time: Mean non-decision times were 0.98 (*SD* = 0.40) in AV and 1.14 (*SD* = 0.44) in TV runs. The best fitting LMM (non-decision time ~ modality + age + (1 | ID)) showed a significant main effect of modality (F(1,66) = 9.61, p = .003; η^2_p = .13, 95% CI [.03, 1.00]) and age on (F(1,65) = 6.27, p = .015; η^2_p = .09, 95% CI [.01, 1.00]) the non-decision time. These results indicate significantly longer non-decision times and thus prolonged multisensory processing in TV compared to AV runs (see **Figure 4A**), and shorter non-decision times with increasing age (see **Figure 4B**). Adding the interaction between modality and age to the model, did not improve model fit (χ^2(1) = 0.66, p = .416), indicating no significant interaction effect.

Boundary separation: Mean boundary separation values were 2.46 (*SD* = 0.35) in AV
and 2.54 (*SD* = 0.41) in TV runs. The best fitting LMM included only an intercept (and random
intercepts for participants) which was significantly greater than zero (β = 2.5, 95% CI [2.43,
2.57], t(131) = 67.44, *p* < .001). Adding the main effect of either modality and age, or the
interaction between the two, did not improve model fit (*p*s > .188). These results indicate that
boundary separations were significantly different from 0, but did not vary with age or differ
between AV and TV learning.

Drift rate: Mean absolute drift rate values for the 1st, 2nd, 3rd, and last bin were 0.10 (*SD*
= 0.11), 0.33 (*SD* = 0.33), 0.48 (*SD* = 0.45), 0.59 (*SD* = 0.53), respectively, in AV and 0.09 (*SD*
= 0.09), 0.27 (*SD* = 0.25), 0.41 (*SD* = 0.37), 0.50 (*SD* = 0.44), respectively, in TV runs. Results
of the best fitting LMM on the drift rate (absolute drift rate ~ modality + bin + age + modality:age
+ bin:age + (1 | participant)) show a significant main effect of modality (F(1, 461) = 14.57, *p* <
.001; η^2_p = .03, 95% CI [.01, 1.00]) and age (F(1,65) = 8.29, *p* = .005; η^2_p = .11, 95% CI [.02,
1.00]), but not bin (F(3,461) = 0.17, *p* = .920; η^2_p = .001, 95% CI [0.00, 1.00]). There was also
a significant interaction between modality and age (F(1,461) = 20.84, *p* < .001; η^2_p = .04, 95%
CI [.02, 1.00]) and between bin and age (F(3,461) = 6.32, *p* < .001; η^2_p = .04, 95% CI [.01,
1.00]). This indicates higher absolute drift rates and thus faster evidence accumulation and
more efficient decision making in older children, but this age effect on decision efficiency is
stronger in AV compared to TV runs and in later compared to earlier bins (see **Figure 4C** and
D). Post-hoc pairwise comparisons between the bins can be seen in **Table 2**.
21

Figure 4

Modality and age effects on non-decision time, boundary separation, and drift rate

Note. **A** depicts the non-decision time and **B** the correlation between age and non-decision time for
AV and TV runs. **C** shows the mean absolute drift rate per modality and bin. The dashed and dotted
lines connect the mean values per bin for AV and TV runs, respectively. **D** plots the correlation
between mean absolute drift rates and age per bin and modality. AV = audio-visual (in blue), TV =
tactile-visual (in green).

Table 2

Post-hoc tests of the effect of bin on drift rate

Model	contrast	estimate	SE	df	t	p
Drift rate ~ bin	1st - 2nd	-0.20	0.03	461.00	-7.36	< .001***
	1st - 3rd	-0.35	0.03	461.00	-12.72	< .001***
	1st - 4th	-0.45	0.03	461.00	-16.33	< .001***
	2nd - 3rd	-0.15	0.03	461.00	-5.36	< .001***
	2nd - 4th	-0.25	0.03	461.00	-8.97	< .001***
	3rd - 4th	-0.10	0.03	461.00	-3.61	.002**

Note. Post-hoc pairwise comparisons for the effect of the specified factor on the dependent variable. Estimates represent the mean difference between conditions. SE = standard error, df = degrees of freedom, p values are Bonferroni-corrected for multiple comparisons.

3.3 Whole-brain fMRI results

To investigate the effects of modality and age on brain activation, we analysed the BOLD signal during both stimulus and feedback processing using a whole-brain approach. Voxel-wise analyses allowed for an unbiased exploration of task-related activation patterns across the entire brain, without prior assumptions. Additionally, we incorporated parametric modulations during these phases to examine how variables such as value and reward prediction errors (RPEs) influence BOLD responses.

3.3.1 Effects of age on stimulus and feedback processing

During AV and TV stimulus processing, age correlated positively with activation in an extended visual and parietal cluster, frontal regions, temporal regions, and the supplementary motor cortex. Brain activation in bilateral post-central, parietal regions, bilateral pre- and post-central regions, medial frontal regions, and bilateral hippocampus negatively correlated with age during AV stimulus processing. During TV stimulus processing, age correlated negatively with brain activation in temporal, hippocampal, and postcentral, parietal regions (see **Figure 5A**). The main effect of age showed visual regions and the inferior frontal gyrus was more active with increasing age of the children for both AV and TV runs. The right supramarginal gyrus was less active the older the children were (see **Figure 5B**). The interaction effect indicates that the bilateral auditory showed a stronger age effect in AV compared to TV runs. Bilateral sensory regions showed a stronger age effect in TV compared to AV runs (see **Figure 5C**). The cluster table and general brain activation during stimulus processing can be found in the Supplementary Material B.3.

In AV runs during feedback processing, age positively correlated with brain activation increases in temporal, visual, and frontal regions. In TV runs during feedback processing, age correlated positively with the brain activation in visual and frontal regions (see **Figure 5A**). Brain activation in the right pallidum, left postcentral gyrus, and right Rolandic operculum showed a negative correlation with age during AV feedback processing. No significant negative correlation with age was observed during TV feedback processing. The main effect of age revealed that a cluster in the right supramarginal gyrus showed lower activation with age in both AV and TV runs. There were no significant clusters showing higher activation with age in both AV and TV runs (see **Figure 5B**). The interaction effect between age and modality revealed that auditory regions and the right angular gyrus showed higher activation with age in AV runs. There were no significant clusters showing higher activation in older children during TV runs, but not AV runs (see **Figure 5C**; the Table B4 in the Supplementary Material shows all the clusters). The general brain activation during feedback processing can be found in the Supplementary Material B.4.

Figure 5

Age-effects on brain activation during stimulus and feedback processing

A Brain activation correlated with age for AV and TV runs

B Main effect of age for stimulus and feedback processing

C Interaction effect of age and modality for stimulus and feedback processing

3.3.2 Modulation of the BOLD signal by model parameters

Significant modulation of the activation was observed for the parametric modulation of the BOLD signal with reward prediction errors and value processing. Additionally, the BOLD signal correlated significantly with the learning rate but not with drift rate. These findings are further explored in paragraphs below.

Reward prediction error processing: Whole-brain analyses showed no significant differences in RPE processing between AV and TV runs. Therefore, for subsequent analyses AV and TV runs were pooled. RPE positively modulated the BOLD activity during feedback processing in the ventral striatum, the precuneus, and in medial frontal regions, and negatively in an extended cluster around the anterior insula, the supplementary motor cortex and superior frontal cortex, bilateral temporal regions, and the bilateral fusiform gyrus extending into the cerebellum. We observed a significant positive effect of age on the modulation of the BOLD activity by the RPE in the left middle frontal gyrus (see **Figure 6** and **Table 3**).

Value processing: Whole-brain analyses showed no significant clusters modulated by values, neither for AV runs, TV runs, nor both modalities combined. We also did not find a significant effect of age on value processing.

Neural correlates of the learning rate: The mean learning rate, calculated across runs for each subject, showed no correlation with brain activation during stimulus processing for neither AV or TV runs separately, nor for both modalities combined.

Table 3

Significant clusters showing a positive and negative modulation by RPEs during feedback processing

Contrast	Brain Area	MNI Coordinates			Cluster size	T-value	Peak-level p_{unc}	Cluster-level p_{FWE}
		x	y	z				
AV + TV	R Striatum	13	0	-15	1582	8.68	<.001	<.001
	L Precuneus	-5	-60	18	1508	7.51	<.001	<.001
	L Inferior Temporal Gyrus	-59	-60	-3	108	6.63	<.001	.013
	L Superior Frontal Gyrus	-17	30	45	353	5.59	<.001	<.001
	R Middle Occipital Gyrus	46	-75	33	149	5.21	<.001	.003
	R Posterior-Medial Frontal	7	18	63	3215	-9.25	<.001	<.001
	R Insula Lobe	37	15	-6	2270	-8.94	<.001	<.001
	L Cerebellum (Crus 1)	-41	-57	-33	666	-7.83	<.001	<.001
	R Supramarginal Gyrus	61	-45	30	697	-7.56	<.001	<.001
	R Cerebellum (VI)	31	-57	-30	734	-6.97	<.001	<.001
Effect of age	L Middle Frontal Gyrus	-29	18	57	129	4.48	<.001	.006

Figure 6

Effects of reward prediction errors on brain activation

Note. Panel **A** shows the brain activation during feedback processing positively and negatively modulated by RPEs for AV and TV runs combined. **B** shows the positive age effect on RPE processing. **C** shows the RPE processing during AV and **D** during TV runs. Warm colours show a positive modulation/age effect on brain activation, and cold colours a negative modulation. Cluster defining threshold *p* < .001 (unc.), *p*FWEc = .05, family-wise error corrected. L = left, R = right, RPE = reward prediction error, AV = audio-visual, TV = tactile-visual, ant. = anterior, med. = medial, suppl. = supplementary, sup. = superior, occip. = occipital, pos. = positive, neg. = negative, mod. = modulation.

2   ## 3.4   ROI analyses

3   To examine the effects of age and modality more closely in brain regions commonly
4   associated with multisensory integration, reward prediction error, and value processing, we
5   conducted ROI analyses in a-priori selected regions of the multisensory, the RPE, and the
6   value networks. These networks and the included regions used for these analyses are shown
7   in **Figure 7**. ROI analyses in the value network are presented in Supplementary Material B.8,
8   as whole-brain analyses revealed no significant activation within the value mask.
9

Figure 7

Depiction of meta-analytical masks used for ROI analyses

A Multisensory network

B Reward prediction error network

Note. Panel **A** shows multisensory clusters and **B** the RPE clusters derived from NeuroSynth.

3.4.1 Brain activation during stimulus presentation

We first examined how brain activation during multisensory stimulus processing was affected by age and modality in regions previously associated with multisensory integration. We observed a significant main effect of modality with higher brain activation for AV runs compared to TV runs across the whole multisensory network ROI and the bilateral planum temporale. In the left postcentral gyrus, right precentral gyrus, and left superior parietal lobe brain activation was higher for TV compared to AV runs. Age positively correlated with the brain activation across the whole network ROI and in all subclusters except the right inferior occipital cortex, right superior parietal lobe and left thalamus. The effect of age was stronger in TV runs compared to AV runs in the left postcentral gyrus and the right precentral gyrus. In the bilateral planum temporale, the age effect was stronger for AV compared to TV runs (see **Figure 8**). The learning rate positively correlated with brain activation across the whole network, in the right planum temporale, and the right postcentral gyrus (the table with the effects can be found in Supplementary Material B.6).

Figure 8

Beta values in the multisensory network ROI and distinct clusters

Note. This figure shows the mean beta values for AV and TV stimulus processing in the whole multisensory network ROI and in each separate cluster, per subject and modality. AV in blue, TV in green. *P*-values are FDR corrected for multiple comparisons. L = left, R = right, AV = audio-visual, TV = tactile-visual, ROI = region of interest.

1
2

3.4.2 Brain activation during reward prediction error processing

Here, we examined how the parametric modulation of brain activation by trial-wise reward prediction errors was influenced by age and modality. Age negatively correlated with the brain activation across the whole RPE network, in the bilateral anterior insula, the right precentral gyrus, and the bilateral supplementary motor cortex. The negative correlation with age in several subclusters indicates higher brain activation with age to lower RPEs. The effect of age across the whole network is most likely driven by the effects in these subclusters. The brain activation across the whole network and in the right striatum shows an effect of modality, which indicates a more positive modulation by RPEs in TV compared to AV runs (see **Figure 9**; the table with model effects can be found in Supplementary Material B.7).

Figure 9

Beta values in the reward prediction error network ROI and distinct clusters

Note. This figure shows the mean beta values for the RPE contrasts in the RPE Network ROI and each separate cluster per subject and modality. AV in blue, TV in green. *P*-values are FDR corrected for multiple comparisons. . L = left, R = right, AV = audio-visual, TV = tactile-visual, ROI = region of interest, suppl. = supplementary, RPE = reward prediction error.

4 Discussion

This study provides insights into the behavioural and neural processing underlying multisensory learning in children, focusing on audio-visual and tactile-visual associations. Three key findings emerged: First, tactile-visual associations were harder to learn than audio-visual associations as shown by performance data across ages. Second, despite considerable individual variability in learning performance, a clear developmental pattern emerged: Older-aged children demonstrated faster processing speed and greater decision-making efficiency, and this enhanced behavioural performance corresponded to higher activation in brain regions for sensory processing, cognitive control, MSI, and memory retrieval. This effect was comparable between AV and TV runs and aligns with known neural maturation processes that support multisensory learning as children develop. Third, while the core processes of reward-based learning and RPE processing, which were modality-independent, remained relatively stable across middle childhood, the older the children were, the higher their brain activity for more negative RPE, suggesting an increased sensitivity to negative feedback. This may indicate a more refined approach to evaluating rewards and punishments with development and suggests a developmental shift in how children process and utilize feedback information to guide their learning.

These findings reveal modality- and age-related alterations in behavioural and neural processes during multisensory learning, highlighting stable core learning processes and developmental shifts in neural engagement. They further emphasize the importance of accounting for age and modality in MSI research.

4.1 Modality-specific processing and learning across middle childhood

Behavioural analyses revealed significant learning during the multisensory task across both modalities. This improvement was shown by faster reaction times and higher accuracies in the second half of each experimental run. We only observed distinct differences in performance between the two sensory modalities in the reaction time. Tactile-visual runs consistently showed slower reaction times compared to audio-visual runs, indicating that tactile-visual multisensory processing was prolonged compared to audio-visual multisensory processing. This modality-specific difference in learning difficulty became more pronounced with increasing age. Computational modelling provided deeper insights into the learning and decision-making in multisensory learning, clarifying modality-specific and age-related changes. First, we discuss how the sensory modality influence learning processes and second how age affects learning across modalities.

The learning rate, which reflects the influence of RPEs on value updating (Zhang et al., 2020), showed similar updating of reward expectation regardless of sensory modality. Similarly, the boundary separation did not differ between modalities, indicating similar amount of information needed until a decision can be reached. However, other computational parameters differed significantly between modalities. Specifically, non-decision time was longer, and drift rates were lower in TV compared to AV learning. These findings align with slower reaction times for tactile-visual association learning, suggesting a slower accumulation process for decision-making in TV runs (Friedrich & Beste, 2019; Ratcliff & McKoon, 2008).

Consistent with the response time analyses, computational modelling revealed that processing speed improved with increasing age, driven by increased drift rates. Older children had higher drift rates in both audio-visual and tactile-visual modalities, with a more pronounced increase over time in audio-visual learning, reflecting faster and more efficient information accumulation during decision processes (Ratcliff & Rouder, 1998). Non-decision time

decreased with age, indicating that the additional processing demands of tactile encoding were
decreasing with age. Additionally, thresholds for accumulating information did not vary with
age. In summary, while learning rates were consistent across modalities and ages, modality
influenced non-decision time and drift rate, with tactile-visual pairings requiring more extensive
processing. In contrast, age enhanced processing speed and decision efficiency across
modalities. This suggests that maturation enhances children's ability to associate relevant
sensory information, improving speed and accuracy in multisensory learning tasks (Barutchu
et al., 2020). To investigate the neural underpinnings of these age-related behavioural
differences, we analysed brain activation patterns during stimulus and feedback processing at
the whole-brain level and in specific ROIs.

4.2 Age-related recruitment of higher-order brain regions during multisensory learning

Whole-brain ANCOVA analyses showed a positive correlation between age and brain
activation during stimulus processing in visual regions and the inferior frontal gyrus. This
suggests that older children may engage in more detailed visual analyses to integrate visual,
auditory, and tactile information during stimulus processing. The inferior frontal gyrus has been
linked to working memory, episodic memory, and attention (Boisgueheneuc et al., 2006;
Diveica et al., 2023; Li et al., 2013), indicating that older children engage these processes
more strongly during stimulus processing independent of modality. While younger children also
recruit higher-order regions for MSI, they may do so less efficiently than older children,
highlighting ongoing MSI network maturation.
Brain activation during stimulus processing increased with age in the occipital cortex,
precuneus, the supplementary motor area (SMA), middle frontal gyrus, and parietal cortex.
This increased neural activity reflects the increased engagement of neural resources in older
children to successfully perform the demanding learning task. Enhanced intraparietal sulcus
activity is linked to cross-modal integration (Sours et al., 2017), mnemonic processes during
both multimodal encoding and retrieval (Tibon et al., 2019), and attention (Boeken & Markett,
2023), indicating a more focused attention on relevant visual stimuli supporting encoding and
learning of multisensory associations. The angular gyrus, a cross-modal hub (Hirst et al., 2021;
Seghier, 2013; Tibon et al., 2019), integrates information from multiple sensory modalities,
which is crucial for forming multisensory associations. The superior parietal lobule, implicated
in attention processes, supports the selection, and focus on relevant sensory information
during learning (Alahmadi, 2021). Greater activity in the SMA, associated with motor planning
(Chung et al., 2005) and working memory (Bahlmann et al., 2009; Chung et al., 2005; Cona &
Semenza, 2017), suggests that older children may plan their responses to a greater extent
which corresponds to the higher accuracies in older children. Additionally, the increased middle
frontal gyrus activity, tied to working memory and cognitive control (Eichenbaum, 2017),
implies a more efficient knowledge access, retrieval, and utilisation during multisensory
learning. Modality-specific age-related increases were observed in the superior temporal lobe
during AV stimulus processing involved in auditory perception and short-term auditory memory
(Crinion et al., 2006; Joseph, 1990). This points towards a greater reliance of older children on
auditory memory to consolidate and retrieve learned associations. During TV stimulus
processing, modality-specific age-related increases in brain activation were observed in
bilateral sensory regions. Indicating that the older children the more are task-relevant sensory
regions recruited during multisensory stimulus processing and thus enhancing perception,
processing, and subsequently multisensory learning. In the previous chapter we described an
increased efficiency in multisensory learning with increasing age. The neural results described
here align with the behavioural results, suggesting that increased recruitment of sensory

regions as children grow older facilitates multisensory processing and subsequently enhances multisensory learning. This is supported by previous research that showed a positive correlation between age and BOLD activation in task-relevant brain regions in a developmental sample (Schapiro et al., 2004), enhanced recruitment of task-specific neural resources to preserve task performance when task demands increase (Wessa et al., 2013), and increased activation during a working memory task in high performing adults (Nagel et al., 2009).

Modality-specific age-related decreases in activation were observed in both AV and TV conditions in a widespread network comprising the thalamus and a cluster spanning the postcentral gyrus, superior parietal lobe, and precuneus. Interestingly, younger children exhibited higher activation in the bilateral middle temporal gyrus during TV processing, suggesting increased engagement in semantic processing (Rolls, 2021). This indicates that younger children may associate semantic meanings to tactile stimuli to enhance memory and retrieval. We hypothesised that younger children would rely more on subcortical and primary sensory regions, while older children would engage higher-order cortical areas during MSL, indicating the increased employment of neural resources (Murray et al., 2016). This can partly be confirmed by the positive correlation with age during AV and TV stimulus processing in higher order brain regions such as parietal and frontal regions. We also observed negative correlations with age in subcortical regions, such as the thalamus, and primary sensory regions, such as temporal and central regions, during multisensory processing. But our analyses highlight, that the age-related changes in our sample appear to be more complex, with positive and negative correlations in both primary sensory and high-order brain regions. These findings suggest that age-related changes in neural activation patterns vary depending on task demands and the specific brain regions involved.

The results of the whole-brain analyses were corroborated by ROI analyses, focusing on regions identified as relevant for multisensory processing and integration in a Neurosynth meta-analysis. Older children exhibited greater activation in most of the regions related to MSI during stimulus processing, although in auditory regions, this increase was limited to AV processing, and in sensory regions specific to TV processing. These results show that older children engage sensory processing, cognitive control, MSI, attention allocation and memory regions more extensively. This enhanced neural engagement likely reflects greater cognitive effort, improving performance in multisensory learning tasks.

4.3 Differential brain activation in AV and TV feedback processing with age

On a whole-brain level, brain activation during AV feedback processing positively correlated with age in the middle frontal gyrus. This region, anatomically connected to the superior and inferior frontal gyri, supports working memory, episodic memory, and attention (Boisgueheneuc et al., 2006; Diveica et al., 2023; Li et al., 2013). Modality specific analyses revealed a positive correlation between brain activation in auditory regions and age during visual feedback processing in AV runs, suggesting older children utilize auditory memory traces to evaluate feedback and adjust behaviour to a greater extent. Similarly, increased visual activation with age in AV and TV runs indicates an increased reliance on visual memory traces for the integration of multisensory information. The fusiform gyrus, crucial for visual attention (Chen et al., 2019; Ruiz-Rizzo et al., 2018), and letter and word form processing (Dehaene & Cohen, 2011; Thesen et al., 2012), facilitates the formation of audio-visual percepts, such as letter-speech sound associations (Karipidis et al., 2018, 2021; Pleisch et al., 2019). In contrast, age-related positive correlations during TV feedback processing were limited to visual regions, likely due to the increased difficulty of integrating tactile information, which limited age-related effects. This aligns with lower TV performance and suggests similar feedback processing and learning strategies across ages. The fMRI results further align with

the drift rate findings, showing more pronounced age-related changes in the AV than TV learning.

4.4 Modality-independent reward prediction error processing

Our findings revealed that that RPE processing was largely modality-independent on the whole-brain level, with RPEs positively modulating brain activity in the ventral striatum, known for encoding RPEs, as well as the medial frontal cortex and hippocampus, linked to learning and memory (Cohen & Ranganath, 2005; Fraga-González et al., 2025; Garrison et al., 2013; McClure et al., 2003; O'Doherty et al., 2003; Pagnoni et al., 2002). RPEs negatively modulated brain activation in regions typically associated with negative RPE signalling, such as the cingulate cortex, anterior insula, and frontal areas (Fouragnan et al., 2018). ROI analyses further confirmed the modality independent RPE processing. Age-related effects on RPE processing were found in the middle frontal gyrus in whole-brain analyses, which indicates an increased involvement of working memory, episodic memory, and attention (Boisgueheneuc et al., 2006; Diveica et al., 2023; Li et al., 2013) with development during RPE processing. ROI analyses revealed a negative correlation between age and brain activity in the bilateral anterior insula, right precentral gyrus and bilateral supplementary motor cortex. The findings of higher anterior insula activation with negative RPEs in older children are indeed consistent in previous findings in adolescents (Hauser et al., 2015; Yaple et al., 2020). This observation supports neurodevelopmental theories proposing that the anterior insula, as an immaturely connected cognitive-emotional hub in adolescence (Smith et al., 2014), biases decision-making towards affectively influenced choices, consistent with the concept of a dominant social-affective system in adolescent cognition (Crone & Dahl, 2012; Icenogle & Cauffman, 2021; Loureiro, 2020; Smith et al., 2014; Van Leijenhorst et al., 2010). For our MSL paradigm the findings thus suggests that older children exhibit heightened sensitivity to negative or unexpected feedback, processing it more extensively, thereby facilitating more effective learning adjustments and error correction. But our findings indicate that this sensitivity is not limited to the anterior insula but extends to other RPE regions as well. In summary, as development progresses children exhibit enhanced MSI, relying on a broader network of brain regions, particularly during audio-visual learning. While RPE processing is modality-independent and stable across ages, older children demonstrate increased sensitivity to negative feedback, suggesting more efficient learning from reward and punishment.

4.5 Limitations

While our study sheds light on how multisensory processing and learning evolve during middle childhood several limitations should be considered when interpreting these results.

First, individual differences in learning abilities, attention and motivation may have influenced performance, particularly among younger children. Eye-tracking in future studies could help assess attention and engagement more directly.

Second, the adaptive nature of the task, ensuring comparable demands across participants, may have masked subtle age-related differences in cognitive and sensory processing. The task's inherent difficulty, especially for younger children, may have limited performance and contributed to variability in learning.

Third, although we designed the tactile vibration patterns to be distinct, they were generally more difficult to differentiate than the auditory stimuli, likely explaining the lower accuracy values during tactile-visual learning.

Further, as a cross-sectional study, our research captures a snapshot of performance at a single developmental stage. Longitudinal studies would be beneficial to track the evolution

of these cognitive processes in individual children and facilitate the understanding the developmental trajectories involved in multisensory learning.

Next, we did not observe any significant correlations between brain activation and either estimated values or learning rate. This may, in part, reflect the nature of the task, which was not explicitly designed to manipulate value and may therefore lack the sensitivity needed to detect robust neural correlates of value processing. Additionally, learning rates were estimated as a single value per run and showed limited variability across participants, age, or modality (see Chapter 3.2). As a result, the between-subject variance of this parameter may have been insufficient to detect meaningful associations with brain activation.

Furthermore, although our findings indicate learning-related changes in both behaviour and model-derived parameters, and all children completed an extensive familiarisation and practice session with the tasks, we cannot entirely rule out the possibility that some observed effects may be attributable to task familiarisation rather than solely associative learning.

Further, ten participants in our sample either had a clinical diagnosis (ADHD, developmental language delay, or developmental dyslexia) and/or a clinically significant T-value on the CBCL/6-18R. Analyses without this subgroup were mostly identical for the drift rate findings, age effects on stimulus processing, neural RPE processing on the whole-brain and ROI-level. The main effect of age and the interaction effects between age and modality during feedback processing on the other hand showed some differences (see Supplementary Material B.1). While all effects, except the stronger age effect during AV feedback processing in the angular gyrus, of the whole sample could be replicated in the smaller group, there were some additional clusters in the smaller sample. Thus, these findings have to be verified in further studies.

Finally, the structure of our task is no suited to directly assess multisensory integration. Since each trial presents both a matching and a non-matching visual stimulus, it is difficult to isolate multisensory integration effects from learning effects, i.e. by contrasting congruent with incongruent multisensory pairs. A follow-up task after the multisensory learning task, specifically designed to test integration (e.g., by comparing learned matching versus non-matching combinations), would be necessary to make claims about multisensory integration of newly learned multisensory associations.

The present findings thus provide valuable insights into how multisensory processing and learning are influenced by modality and age during middle childhood. Understanding how multisensory processing and learning unfold across development lays the groundwork for studying and further understanding impairments in these processes (e.g., related to developmental language delays or other neurodevelopmental disorders).

4.6 Conclusion

Our findings show that typically developing children can learn new associations between abstract visual and auditory or tactile stimuli within an experimental session. Learning rates remained consistent across modalities and age groups, but modality-specific factors influenced decision-making. With increasing development, children exhibited more efficient multisensory processing and integration. This was accompanied by increased neural activation in regions linked to cognitive control, working memory, and MSI, indicating the usage of increased neural resources for MSL the older the children. Older children exhibited increased activation in the bilateral anterior insula, right precentral gyrus, and bilateral supplementary motor cortex in response to negative RPEs, suggesting a heightened sensitivity to negative feedback and a more nuanced learning process that integrates both positive and negative feedback for behavioural adaptation. In contrast, younger children in our study apparently relied more on basic learning mechanisms and seemed less able to flexibly adapt their

behaviour based on negative feedback. Overall, older children use advanced cognitive strategies and neural resources in multisensory learning tasks, while younger children engage higher-order MSI regions less efficiently, reflecting ongoing dynamics and developmental maturation in multisensory learning. To conclude, our findings indicate that with increasing age children are more efficient at integrating information across different sensory modalities to aid learning of multisensory associations, which they may utilize more effectively to guide their behaviour. This ability is essential for advanced learning, problem-solving, and adaptation in dynamic environments.

CRediT authorship contribution statement

Nina Raduner: conceptualisation, data curation, formal analysis, investigation, methodology, project administration, visualization, writing – original draft; **Carmen Providoli:** conceptualisation, data curation, investigation, project administration, writing – review and editing; **Sarah V. Di Pietro:** conceptualisation, data curation, investigation, project administration, supervision, writing – review and editing; **Maya Schneebeli:** methodology, supervision, writing – review and editing; **Iliana I. Karipidis:** conceptualisation, supervision, writing – review and editing; **Ella Casimiro:** conceptualisation, methodology; **Saurabh Bedi:** conceptualisation, methodology, writing – review and editing; **Michael von Rhein:** conceptualisation, funding acquisition, supervision, writing – review and editing; **Nora M. Raschle:** conceptualisation, funding acquisition, methodology, supervision, writing – review and editing; **Christian C. Ruff:** conceptualisation, funding acquisition, methodology, resources, supervision, writing – review and editing; **Silvia Brem:** conceptualisation, funding acquisition, methodology, project administration, resources, supervision, writing – review and editing.

Declaration of competing interest

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used ChatGPT, Perplexity, Gemini, and Quillbot in order improve language and readability. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

Acknowledgments

We are very grateful for all the families and children for participating in this study. Further, we thank K. Bremer, A. Bussmann, N. Ehrhardt, E. Hohl, K. Gelic, S. Gubler, L. Hämmerli, S. Ismail, R. Jocham, I.P. Johnson, A. Lütolf, X. Michael, M. Peter, S. Schädler, J. Schiesser, M.A. Steuer, P. Sting, L. Strickler, C. Volk, and R. Wombacher for their help in planning, recruitment, and data collection. We thank Y. Cao for his contribution to implementing the drift diffusion model, P. Stämpfli for his advice on the MRI protocol, G.S.P. Pamplona and A. Haugg for their help in the statistical analyses and R. Borbás, P. Dimanova and S. Scatolin for their valuable input.

Funding

This work was supported by the University Research Priority Program Adaptive Brain Circuits in Development and Learning (AdaBD) of the University of Zurich (project ChildBrainCircuits).

Data and code availability

The data (no raw or pre-processed neuroimaging data) and code for this publication are available on https://osf.io/cmzne/ (DOI 10.17605/OSF.IO/CMZNE).

References

Alahmadi, A. A. S. (2021). Investigating the sub-regions of the superior parietal cortex using functional magnetic resonance imaging connectivity. *Insights into Imaging*, *12*(1). https://doi.org/10.1186/s13244-021-00993-9

Altarelli, I., Dehaene-Lambertz, G., & Bavelier, D. (2020). Individual differences in the acquisition of non-linguistic audio-visual associations in 5 year olds. *Developmental Science*, *23*(4). https://doi.org/10.1111/desc.12913

Bahlmann, J., Schubotz, R. I., Mueller, J. L., Koester, D., & Friederici, A. D. (2009). Neural circuits of hierarchical visuo-spatial sequence processing. *Brain Research*, *1298*. https://doi.org/10.1016/j.brainres.2009.08.017

Barutchu, A., Crewther, S. G., Fifer, J., Shivdasani, M. N., Innes-Brown, H., Toohey, S., Danaher, J., & Paolini, A. G. (2011). The relationship between multisensory integration and IQ in children. *Developmental Psychology*, *47*(3), 877–885. https://doi.org/10.1037/a0021903

Barutchu, A., Fifer, J. M., Shivdasani, M. N., Crewther, S. G., & Paolini, A. G. (2020). The Interplay Between Multisensory Associative Learning and IQ in Children. *Child Development*, *91*(2), 620–637. https://doi.org/10.1111/cdev.13210

Barutchu, A., Toohey, S., Shivdasani, M. N., Fifer, J. M., Crewther, S. G., Grayden, D. B., & Paolini, A. G. (2019). Multisensory perception and attention in school-age children. *Journal of Experimental Child Psychology*, *180*, 141–155. https://doi.org/10.1016/j.jecp.2018.11.021

Batson, M. A., Beer, A. L., Seitz, A. R., & Watanabe, T. (2011). Spatial shifts of audio-visual interactions by perceptual learning are specific to the trained orientation and eye. *Seeing and Perceiving*, *24*(6). https://doi.org/10.1163/187847611X603738

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1). https://doi.org/10.1016/j.neuroimage.2007.04.042

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *57*(1). https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Birch, H. G., & Belmont, L. (1965). AUDITORY-VISUAL INTEGRATION, INTELLIGENCE AND READING ABILITY IN SCHOOL CHILDREN. *Perceptual and Motor Skills*, *20*, 295–305. https://doi.org/10.2466/pms.1965.20.1.295

Boeken, O. J., & Markett, S. (2023). Systems-level decoding reveals the cognitive and behavioral profile of the human intraparietal sulcus. *Frontiers in Neuroimaging*, *1*. https://doi.org/10.3389/fnimg.2022.1074674

Boisgueheneuc, F. Du, Levy, R., Volle, E., Seassau, M., Duffau, H., Kinkingnehun, S., Samson, Y., Zhang, S., & Dubois, B. (2006). Functions of the left superior frontal gyrus in humans: A lesion study. *Brain*, *129*(12). https://doi.org/10.1093/brain/awl244

Butler, A. J., James, T. W., & James, K. H. (2011). Enhanced multisensory integration and motor reactivation after active motor learning of audiovisual associations. *Journal of Cognitive Neuroscience*, *23*(11). https://doi.org/10.1162/jocn_a_00015

Chai, X. J., Castañán, A. N., Öngür, D., & Whitfield-Gabrieli, S. (2012). Anticorrelations in resting state networks without global signal regression. *NeuroImage*, *59*(2). https://doi.org/10.1016/j.neuroimage.2011.08.048

Chen, L., Wassermann, D., Abrams, D. A., Kochalka, J., Gallardo-Diez, G., & Menon, V. (2019). The visual word form area (VWFA) is part of both language and attention circuitry. *Nature Communications*, *10*(1). https://doi.org/10.1038/s41467-019-13634-z

Chien, S., Wiehler, A., Spezio, M., & Gläscher, J. (2016). Congruence of inherent and acquired values facilitates reward-based decision-making. *Journal of Neuroscience*, *36*(18). https://doi.org/10.1523/JNEUROSCI.3084-15.2016

Chung, G. H., Han, Y. M., Jeong, S. H., & Jack, C. R. (2005). Functional heterogeneity of the supplementary motor area. *American Journal of Neuroradiology*, *26*(7).

Cohen, M. X., & Ranganath, C. (2005). Behavioral and neural predictors of upcoming decisions. *Cognitive, Affective and Behavioral Neuroscience*, *5*(2). https://doi.org/10.3758/CABN.5.2.117

Cohen, M. X., & Ranganath, C. (2007). Reinforcement learning signals predict future decisions. *Journal of Neuroscience*, *27*(2). https://doi.org/10.1523/JNEUROSCI.4421-06.2007

Cona, G., & Semenza, C. (2017). Supplementary motor area as key structure for domain-general sequence processing: A unified account. In *Neuroscience and Biobehavioral Reviews* (Vol. 72). https://doi.org/10.1016/j.neubiorev.2016.10.033

Cover, T. M., & Thomas, J. A. (2005). Elements of Information Theory. In *Elements of Information Theory*. https://doi.org/10.1002/047174882X

Crinion, J. T., Warburton, E. A., Lambon-Ralph, M. A., Howard, D., & Wise, R. J. S. (2006). Listening to narrative speech after aphasic stroke: The role of the left anterior temporal lobe. *Cerebral Cortex*, *16*(8). https://doi.org/10.1093/cercor/bhj053

Crone, E. A., & Dahl, R. E. (2012). Understanding adolescence as a period of social-affective engagement and goal flexibility. *Nature Reviews Neuroscience*, *13*(9). https://doi.org/10.1038/nrn3313

Dehaene, S., & Cohen, L. (2011). The unique role of the visual word form area in reading. In *Trends in Cognitive Sciences* (Vol. 15, Issue 6). https://doi.org/10.1016/j.tics.2011.04.003

Denervaud, S., Gentaz, E., Matusz, P. J., & Murray, M. M. (2020). Multisensory Gains in Simple Detection Predict Global Cognition in Schoolchildren. *Scientific Reports*, *10*(1), 1394. https://doi.org/10.1038/s41598-020-58329-4

Desjardins, R. N., Rogers, J., & Werker, J. F. (1997). An exploration of why preschoolers perform differently than do adults in audiovisual speech perception tasks. *Journal of Experimental Child Psychology*, *66*(1), 85–110. https://doi.org/10.1006/jecp.1997.2379

Dionne-Dostie, E., Paquette, N., Lassonde, M., & Gallagher, A. (2015). Multisensory Integration and Child Neurodevelopment. *Brain Sciences*, *5*(1), 32–57. https://doi.org/10.3390/brainsci5010032

Diveica, V., Riedel, M. C., Salo, T., Laird, A. R., Jackson, R. L., & Binney, R. J. (2023). Graded functional organization in the left inferior frontal gyrus: evidence from task-free and task-based functional connectivity. *Cerebral Cortex*, *33*(23). https://doi.org/10.1093/cercor/bhad373

Döpfner, M., Plück, J., Kinnen, C., & Checklist, A. D. C. B. (2014). CBCL Handbuch-Schulalter. Manual zum Elternfragebogen über das Verhalten von Kindern und Jugendlichen. In *(CBCL/6-18R), zum Lehrerfragebogen über das Verhalten von Kindern und Jugendlichen (TRF/6-18R) und zum Fragebogen für Jugendliche (YSR/11-18R)*. Hogrefe Göttingen.

Eichenbaum, H. (2017). Memory: Organization and Control. *Annual Review of Psychology*, *68*. https://doi.org/10.1146/annurev-psych-010416-044131

Eklund, A., Nichols, T. E., & Knutsson, H. (2016). Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(28). https://doi.org/10.1073/pnas.1602413113

Farrell, S., & Lewandowsky, S. (2018). Computational modeling of cognition and behavior. In *Computational Modeling of Cognition and Behavior*. https://doi.org/10.1017/CBO9781316272503

Field, A. (2013). Discovering statistics using IBM SPSS statistics. In *Statistics* (Vol. 58).

Fouragnan, E., Retzler, C., & Philiastides, M. G. (2018). Separate neural representations of prediction error valence and surprise: Evidence from an fMRI meta-analysis. *Human Brain Mapping*, *39*(7). https://doi.org/10.1002/hbm.24047

Fraga-González, G., Haller, P., Willinger, D., Gehrig, V., Frei, N., & Brem, S. (2025). Functional brain activations correlated with association strength and prediction error during novel symbol-speech sound learning. *Imaging Neuroscience*. https://doi.org/10.1162/imag_a_00439

Frei, N., Willinger, D., Haller, P., Fraga-González, G., Pamplona, G. S. P., Haugg, A., Lutz, C. G., Coraj, S., Hefti, E., & Brem, S. (2025). Towards a mechanistic understanding of reading difficulties: Deviant audiovisual learning dynamics and network connectivity in children with poor reading skills. *The Journal of Neuroscience*, e1119242025. https://doi.org/10.1523/JNEUROSCI.1119-24.2025

Friedrich, J., & Beste, C. (2019). The impact of stimulus modality on the processing of conflicting sensory information during response inhibition. *Neuroscience*, *410*. https://doi.org/10.1016/j.neuroscience.2019.05.010

Gao, C., Green, J. J., Yang, X., Oh, S., Kim, J., & Shinkareva, S. V. (2023). Audiovisual integration in the human brain: a coordinate-based meta-analysis. *Cerebral Cortex*, *33*(9). https://doi.org/10.1093/cercor/bhac443

Garrison, J., Erdeniz, B., & Done, J. (2013). Prediction error in reinforcement learning: A meta-analysis of neuroimaging studies. In *Neuroscience and Biobehavioral Reviews* (Vol. 37, Issue 7). https://doi.org/10.1016/j.neubiorev.2013.03.023

Gentile, G., Björnsdotter, M., Petkova, V. I., Abdulkarim, Z., & Ehrsson, H. H. (2015). Patterns of neural activity in the human ventral premotor cortex reflect a whole-body multisensory percept. *NeuroImage*, *109*. https://doi.org/10.1016/j.neuroimage.2015.01.008

Gentile, G., Guterstam, A., Brozzoli, C., & Henrik Ehrsson, H. (2013). Disintegration of multisensory signals from the real hand reduces default limb self-attribution: An fMRI study. *Journal of Neuroscience*, *33*(33). https://doi.org/10.1523/JNEUROSCI.1363-13.2013

Ghazanfar, A. A., & Schroeder, C. E. (2006). Is neocortex essentially multisensory? In *Trends in Cognitive Sciences* (Vol. 10, Issue 6, pp. 278–285). https://doi.org/10.1016/j.tics.2006.04.008

Gori, M., Del Viva, M., Sandini, G., & Burr, D. C. (2008). Young Children Do Not Integrate Visual and Haptic Form Information. *Current Biology*, *18*(9). https://doi.org/10.1016/j.cub.2008.04.036

Grob, A., & Hagmann-von Arx, P. (2018). *IDS-2-Intelligence and Development Scales-2. Intelligenz-und Entwicklungsskalen für Kinder und Jugendliche-Testkoffer*. Hogrefe.

Hallquist, M. N., Hwang, K., & Luna, B. (2013). The nuisance of nuisance regression: Spectral misspecification in a common approach to resting-state fMRI preprocessing reintroduces noise and obscures functional connectivity. *NeuroImage*, *82*. https://doi.org/10.1016/j.neuroimage.2013.05.116

Hauser, T. U., Iannaccone, R., Walitza, S., Brandeis, D., & Brem, S. (2015). Cognitive flexibility in adolescence: Neural and behavioral mechanisms of reward prediction error processing in adaptive decision making during development. *NeuroImage*, *104*. https://doi.org/10.1016/j.neuroimage.2014.09.018

Hirst, R. J., Whelan, R., Boyle, R., Setti, A., Knight, S., O'Connor, J., Williamson, W., McMorrow, J., Fagan, A. J., Meaney, J. F., Kenny, R. A., De Looze, C., & Newell, F. N. (2021). Gray matter volume in the right angular gyrus is associated with differential patterns of multisensory integration with aging. *Neurobiology of Aging*, *100*. https://doi.org/10.1016/j.neurobiolaging.2020.12.004

Icenogle, G., & Cauffman, E. (2021). Adolescent decision making: A decade in review. In *Journal of Research on Adolescence* (Vol. 31, Issue 4). https://doi.org/10.1111/jora.12608

Joseph, R. (1990). *The Temporal Lobes*. https://doi.org/10.1007/978-1-4757-5969-3_7

Karipidis, I. I., Pleisch, G., Brandeis, D., Roth, A., Röthlisberger, M., Schneebeli, M., Walitza, S., & Brem, S. (2018). Simulating reading acquisition: The link between reading outcome and multimodal brain signatures of letter–speech sound learning in prereaders. *Scientific Reports*, *8*(1), 7121.

Karipidis, I. I., Pleisch, G., Di Pietro, S. V., Fraga-González, G., & Brem, S. (2021). Developmental Trajectories of Letter and Speech Sound Integration During Reading Acquisition. *Frontiers in Psychology*, *12*. https://doi.org/10.3389/fpsyg.2021.750491

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13, 1–26). https://doi.org/10.18637/JSS.V082.I13

Lakatos, P., Chen, C. M., O'Connell, M. N., Mills, A., & Schroeder, C. E. (2007). Neuronal Oscillations and Multisensory Interaction in Primary Auditory Cortex. *Neuron*, *53*(2), 279–292. https://doi.org/10.1016/j.neuron.2006.12.011

Lauzon, S., Abraham, A. E., Curcin, K., Butler, B. E., & Stevenson, R. A. (2022). The relationship between multisensory associative learning and multisensory integration. *Neuropsychologia*, *174*. https://doi.org/10.1016/j.neuropsychologia.2022.108336

Lenhard, A., Lenhard, W., Segerer, R., & Suggate, S. (2015). *Peabody Picture Vocabulary Test-4. Ausgabe: Deutsche Fassung*. Frankfurt am Main: Pearson Assessment.

Lenth, R. (2024). *emmeans: Estimated marginal means, aka least-squares means* (R package version 1.10. 2). https://CRAN.R-project.org/package=emmeans

Li, W., Qin, W., Liu, H., Fan, L., Wang, J., Jiang, T., & Yu, C. (2013). Subregions of the human superior frontal gyrus and their connections. *NeuroImage*, *78*. https://doi.org/10.1016/j.neuroimage.2013.04.011

Limanowski, J., & Blankenburg, F. (2016). Integration of visual and proprioceptive limb position information in human posterior parietal, premotor, and extrastriate cortex. *Journal of Neuroscience*, *36*(9). https://doi.org/10.1523/JNEUROSCI.3987-15.2016

Limanowski, J., & Blankenburg, F. (2017). Posterior parietal cortex evaluates visuoproprioceptive congruence based on brief visual information. *Scientific Reports*, *7*(1). https://doi.org/10.1038/s41598-017-16848-7

Liu, X., Hairston, J., Schrier, M., & Fan, J. (2011). Common and distinct networks underlying reward valence and processing stages: A meta-analysis of functional neuroimaging studies. In *Neuroscience and Biobehavioral Reviews* (Vol. 35, Issue 5, pp. 1219–1236). https://doi.org/10.1016/j.neubiorev.2010.12.012

Loureiro, R. J. (2020). Decision making in adolescents: A multifaceted construct. In *Journal of Human Growth and Development* (Vol. 30, Issue 2). https://doi.org/10.7322/JHGD.V30.10362

MathWorks Inc. (2020). *MATLAB, Version 9.9.0 (R2020b)*.

MathWorks Inc. (2022). *MATLAB version: 9.13.0 (R2022b)*. The MathWorks Inc Natick, Massachusetts.

Mathys, C., Daunizeau, J., Friston, K. J., & Stephan, K. E. (2011). A Bayesian foundation for individual learning under uncertainty. *Frontiers in Human Neuroscience*, *MAY*. https://doi.org/10.3389/fnhum.2011.00039

Mazaika, P., Whitfield-Gabrieli, S., Reiss, A., & Glover, G. (2007). Artifact repair for fMRI data from high motion clinical subjects. *Human Brain Mapping*, *47*(58), 70231–70238.

McClure, S. M., Berns, G. S., & Montague, P. R. (2003). Temporal prediction errors in a passive learning task activate human striatum. *Neuron*, *38*(2). https://doi.org/10.1016/S0896-6273(03)00154-5

Misra, H., Ikbal, S., Bourlard, H., & Hermansky, H. (2004). Spectral entropy based feature for robust ASR. *ICASSP, IEEE International Conference on Acoustics, Speech and Signal Processing - Proceedings*, *1*. https://doi.org/10.1109/icassp.2004.1325955

Murray, M. M., Lewkowicz, D. J., Amedi, A., & Wallace, M. T. (2016). Multisensory Processes: A Balancing Act across the Lifespan. *Trends in Neurosciences*, *39*(8), 567–579. https://doi.org/10.1016/j.tins.2016.05.003

Nagel, I. E., Preuschhof, C., Li, S. C., Nyberg, L., Bäckman, L., Lindenberger, U., & Heekeren, H. R. (2009). Performance level modulates adult age differences in brain activation during spatial working memory. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(52). https://doi.org/10.1073/pnas.0908238106

Nardini, M., Jones, P., Bedford, R., & Braddick, O. (2008). Development of Cue Integration in Human Navigation. *Current Biology*, *18*(9). https://doi.org/10.1016/j.cub.2008.04.021

Navarro, D. J., & Fuss, I. G. (2009). Fast and accurate calculations for first-passage times in Wiener diffusion models. In *Journal of Mathematical Psychology* (Vol. 53, Issue 4). https://doi.org/10.1016/j.jmp.2009.02.003

Nieto-Castanon, A. (2020). FMRI denoising pipeline. In *Handbook of functional connectivity Magnetic Resonance Imaging methods in CONN*. https://doi.org/10.56441/hilbertpress.2207.6600

Nieto-Castanon, A., & Whitfield-Gabrieli, S. (2022). CONN functional connectivity toolbox: RRID SCR_009550, release 22. In *CONN functional connectivity toolbox: RRID SCR_009550, release 22*. https://doi.org/10.56441/hilbertpress.2246.5840

Niv, Y., & Schoenbaum, G. (2008). Dialogues on prediction errors. In *Trends in Cognitive Sciences* (Vol. 12, Issue 7). https://doi.org/10.1016/j.tics.2008.03.006

Noesselt, T., Rieger, J. W., Schoenfeld, M. A., Kanowski, M., Hinrichs, H., Heinze, H. J., & Driver, J. (2007). Audiovisual temporal correspondence modulates human multisensory superior temporal sulcus plus primary sensory cortices. *Journal of Neuroscience*, *27*(42), 11431–11441. https://doi.org/10.1523/JNEUROSCI.2252-07.2007

Obin, N., & Liuni, M. (2012). On the generalization of Shannon entropy for speech recognition. *2012 IEEE Workshop on Spoken Language Technology, SLT 2012 - Proceedings*. https://doi.org/10.1109/SLT.2012.6424204

O'Doherty, J. P., Dayan, P., Friston, K., Critchley, H., & Dolan, R. J. (2003). Temporal difference models and reward-related learning in the human brain. *Neuron*, *38*(2). https://doi.org/10.1016/S0896-6273(03)00169-7

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

O'Reilly, J. X. (2013). Making predictions in a changing world-inference, uncertainty, and learning. *Frontiers in Neuroscience*, *7 JUN*. https://doi.org/10.3389/fnins.2013.00105

Pagnoni, G., Zink, C. F., Montague, P. R., & Berns, G. S. (2002). Activity in human ventral striatum locked to errors of reward prediction. *Nature Neuroscience*, *5*(2). https://doi.org/10.1038/nn802

Pasqualotto, A., Cochrane, A., Bavelier, D., & Altarelli, I. (2024). A novel task and methods to evaluate inter-individual variation in audio-visual associative learning. *Cognition*, *242*. https://doi.org/10.1016/j.cognition.2023.105658

Patterson, M. L., & Werker, J. F. (1999). Matching phonetic information in lips and voice is robust in 4.5-month-old infants. *Infant Behavior and Development*, *22*(2), 237–247. https://doi.org/10.1016/S0163-6383(99)00003-X

Patterson, M. L., & Werker, J. F. (2003). Two-month-old infants match phonetic information in lips and voice. *Developmental Science*, *6*(2), 191–196. https://doi.org/10.1111/1467-7687.00271

Pedersen, M. L., Frank, M. J., & Biele, G. (2017). The drift diffusion model as the choice rule in reinforcement learning. *Psychonomic Bulletin and Review*, *24*(4). https://doi.org/10.3758/s13423-016-1199-y

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, *51*(1). https://doi.org/10.3758/s13428-018-01193-y

Penny, W., Friston, K., Ashburner, J., Kiebel, S., & Nichols, T. (2007). Statistical Parametric Mapping: The Analysis of Functional Brain Images. In *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. https://doi.org/10.1016/B978-0-12-372560-8.X5000-1

Pernet, C. R. (2014). Misconceptions in the use of the General Linear Model applied to functional MRI: A tutorial for junior neuro-imagers. In *Frontiers in Neuroscience* (Issue 8 JAN). https://doi.org/10.3389/fnins.2014.00001

Pleisch, G., Karipidis, I. I., Brem, A., Röthlisberger, M., Roth, A., Brandeis, D., Walitza, S., & Brem, S. (2019). Simultaneous EEG and fMRI reveals stronger sensitivity to orthographic strings in the left occipito-temporal cortex of typical versus poor beginning readers. *Developmental Cognitive Neuroscience*, *40*. https://doi.org/10.1016/j.dcn.2019.100717

R Core Team, R. (2024). *R: A language and environment for statistical computing*. R foundation for statistical computing, Vienna, Austria. https://www.R-project.org/

Ratcliff, R. (1978). A theory of memory retrieval. *Psychological Review*, *85*(2), 59–108. https://doi.org/10.1037/0033-295X.85.2.59

Ratcliff, R., & McKoon, G. (2008). The diffusion decision model: Theory and data for two-choice decision tasks. In *Neural Computation* (Vol. 20, Issue 4). https://doi.org/10.1162/neco.2008.12-06-420

Ratcliff, R., & Rouder, J. N. (1998). Modeling Response Times for Two-Choice Decisions. *Psychological Science*, *9*(5). https://doi.org/10.1111/1467-9280.00067

Rescorla, R. A., & Wagner, A. R. (1972). A Theory of Pavlovian Conditioning: Variations in the Effectiveness of Reinforcement and Nonreinforcement. *Classical Conditioning II: Current Research and Theory*.

Richlan, F. (2019). The functional neuroanatomy of letter-speech sound integration and its relation to brain abnormalities in developmental dyslexia. In *Frontiers in Human Neuroscience* (Vol. 13). https://doi.org/10.3389/fnhum.2019.00021

Rolls, E. T. (2021). The temporal cortex. In *Brain Computations: What and How*. Oxford University Press.

Ruiz-Rizzo, A. L., Neitzel, J., Müller, H. J., Sorg, C., & Finke, K. (2018). Distinctive correspondence between separable visual attention functions and intrinsic brain networks. *Frontiers in Human Neuroscience*, *12*. https://doi.org/10.3389/fnhum.2018.00089

Schapiro, M. B., Schmithorst, V. J., Wilke, M., Byars, A. W., Strawsburg, R. H., & Holland, S. K. (2004). BOLD fMRI signal increases with age in selected brain regions in children. *NeuroReport*, *15*(17). https://doi.org/10.1097/00001756-200412030-00003

Seghier, M. L. (2013). The angular gyrus: Multiple functions and multiple subdivisions. In *Neuroscientist* (Vol. 19, Issue 1). https://doi.org/10.1177/1073858412440596

Seitz, A. R., Kim, R., Van Wassenhove, V., & Shams, L. (2007). Simultaneous and independent acquisition of multisensory and unisensory associations. *Perception*, *36*(10). https://doi.org/10.1068/p5843

Smith, A. R., Steinberg, L., & Chein, J. (2014). The role of the anterior insula in adolescent decision making. In *Developmental Neuroscience* (Vol. 36, Issues 3–4). https://doi.org/10.1159/000358918

Sours, C., Raghavan, P., Medina, A. E., Roys, S., Jiang, L., Zhuo, J., & Gullapalli, R. P. (2017). Structural and Functional Integrity of the Intraparietal Sulcus in Moderate and Severe Traumatic Brain Injury. *Journal of Neurotrauma*, *34*(7). https://doi.org/10.1089/neu.2016.4570

Stein, B. E., & Stanford, T. R. (2008). Multisensory integration: Current issues from the perspective of the single neuron. In *Nature Reviews Neuroscience* (Vol. 9, Issue 4, pp. 255–266). https://doi.org/10.1038/nrn2331

Stein, B. E., Stanford, T. R., & Rowland, B. A. (2020). Multisensory integration and the society for neuroscience: Then and now. In *Journal of Neuroscience* (Vol. 40, Issue 1). https://doi.org/10.1523/JNEUROSCI.0737-19.2019

Subramanian, A., Chitlangia, S., & Baths, V. (2022). Reinforcement learning and its connections with neuroscience and psychology. *Neural Networks*, *145*. https://doi.org/10.1016/j.neunet.2021.10.003

Team Audacity. (2024). *Audacity (r): free audio editor and recorder [computer program]* (3.5.1). http://audacity.sourceforge.net/

Thesen, T., McDonald, C. R., Carlson, C., Doyle, W., Cash, S., Sherfey, J., Felsovalyi, O., Girard, H., Barr, W., Devinsky, O., Kuzniecky, R., & Halgren, E. (2012). Sequential then interactive processing of letters and words in the left fusiform gyrus. *Nature Communications*, *3*. https://doi.org/10.1038/ncomms2220

Tibon, R., Fuhrmann, D., Levy, D. A., Simons, J. S., & Henson, R. N. (2019). Multimodal integration and vividness in the angular gyrus during episodic encoding and retrieval. *Journal of Neuroscience*, *39*(22). https://doi.org/10.1523/JNEUROSCI.2102-18.2018

Tyll, S., Budinger, E., & Noesselt, T. (2011). Thalamic influences on multisensory integration. In *Communicative and Integrative Biology* (Vol. 4, Issue 4). https://doi.org/10.4161/cib.15222

Van Atteveldt, N., Murray, M. M., Thut, G., & Schroeder, C. E. (2014). Multisensory integration: Flexible use of general operations. In *Neuron* (Vol. 81, Issue 6). https://doi.org/10.1016/j.neuron.2014.02.044

Van Leijenhorst, L., Zanolie, K., Van Meel, C. S., Westenberg, P. M., Rombouts, S. A. R. B., & Crone, E. A. (2010). What motivates the adolescent? brain regions mediating reward sensitivity across adolescence. *Cerebral Cortex*, *20*(1). https://doi.org/10.1093/cercor/bhp078

Vildavski, V., Lo Verde, L., Blumberg, G., Parsey, J., & Norcia, A. (2021). PseudoSloan: A perimetric-complexity and area-controlled font for vision and reading research. *Journal of Vision*, *21*(9). https://doi.org/10.1167/jov.21.9.2857

Wallace, M. T., Woynaroski, T. G., & Stevenson, R. A. (2020). Multisensory integration as a window into orderly and disrupted cognition and communication. In *Annual Review of Psychology* (Vol. 71). https://doi.org/10.1146/annurev-psych-010419-051112

Wang, F., Karipidis, I. I., Pleisch, G., Fraga-González, G., & Brem, S. (2020). Development of Print-Speech Integration in the Brain of Beginning Readers With Varying Reading Skills. *Frontiers in Human Neuroscience*, *14*(289). https://doi.org/10.3389/fnhum.2020.00289

Wechsler, D. (2012). Wechsler preschool and primary scale of intelligence—fourth edition. *The Psychological Corporation San Antonio, TX*.

Wechsler, D. (2017). Wechsler Intelligence Scale for Children–Fifth Edition (WISC-V)(dt. Bearbeitung durch F. Petermann). In *Frankfurt a. M.: Pearson Assessment*.

Wessa, M., Heissler, J., Schönfelder, S., & Kanske, P. (2013). Goal-directed behavior under emotional distraction is preserved by enhanced task-specific activation. *Social Cognitive and Affective Neuroscience*, *8*(3). https://doi.org/10.1093/scan/nsr098

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connectivity*, *2*(3). https://doi.org/10.1089/brain.2012.0073

Wilke, M., Holland, S. K., Altaye, M., & Gaser, C. (2008). Template-O-Matic: A toolbox for creating customized pediatric templates. *NeuroImage*, *41*(3). https://doi.org/10.1016/j.neuroimage.2008.02.056

Wilson, R. C., & Collins, A. G. E. (2019). Ten simple rules for the computational modeling of behavioral data. *ELife*, *8*. https://doi.org/10.7554/eLife.49547

Wilson, R. P., Colizzi, M., Bossong, M. G., Allen, P., Kempton, M., MTAC, & Bhattacharyya, S. (2018). The Neural Substrate of Reward Anticipation in Health: A Meta-Analysis of fMRI Findings in the Monetary Incentive Delay Task. *Neuropsychology Review*, *28*(4), 496–506. https://doi.org/10.1007/s11065-018-9385-5

Wu, X., Wang, T., Liu, C., Wu, T., Jiang, J., Zhou, D., & Zhou, J. (2017). Functions of learning rate in adaptive reward learning. *Frontiers in Human Neuroscience*, *11*. https://doi.org/10.3389/fnhum.2017.00592

Yaple, Z. A., Yu, R., & Arsalidou, M. (2020). Spatial migration of human reward processing with functional development: Evidence from quantitative meta-analyses. *Human Brain Mapping*, *41*(14). https://doi.org/10.1002/hbm.25103

Zhang, L., Lengersdorff, L., Mikus, N., Gläscher, J., & Lamm, C. (2020). Using reinforcement learning models in social neuroscience: Frameworks, pitfalls and suggestions of best practices. *Social Cognitive and Affective Neuroscience*, *15*(6), 695–707. https://doi.org/10.1093/scan/nsaa089

Zimmermann, P., Fimm, B., & Gondan, M. (2002). *Testbatterie zur aufmerksamkeitsprüfung für kinder:(KiTAP);[das schloß der geister]*. Psytest.