Comparative and reproducible three-dimension functional spaces in primate cortical connectomes

Xiu-Xia Xing [1,*], **Xi-Nian Zuo** [2,3,4,*]

[1] Department of Applied Mathematics, College of Mathematics, Faculty of Science, Beijing University of Technology, Beijing 100124, China.
[2] State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, No 19 Xinjiekouwai Street, Haidian District, Beijing, 100875, China.
[3] Developmental Population Neuroscience Research Center, IDG/McGovern Institute for Brain Research, Beijing Normal University, No 19 Xinjiekouwai Street, Haidian District, Beijing, 100875, China.
[4] National Basic Science Data Center, No 2 Dongshengnan Road, Haidian District, Beijing, 100190, China.
[*] Corresponding authors: Xi-Nian Zuo & Xiu-Xia Xig; Emails: xinian.zuo@bnu.edu.cn, xingxx@bjut.edu.cn

ABSTRACT

Intrinsic activity or spontaneous oscillations are believed to be essential for the brain to maintain flexible, adaptive, and re-
sponsive cognition to both internal and external demands. However, a functional space remains elusive for understanding
this inside-outside organization in the primate cortex. Here, we leverage comparative fMRI analysis to identify three con-
served and reproducible functional connectivity gradients as three axes of manifold (FMA), which span a low-dimensional
organizational space of wakeful primate connectomes. In humans, the primary FMA covers a sensorimotor-to-transmodal
gradient that supports signal detection for allostatic anticipation. The second FMA mirrors a representation-to-modulation
gradient that integrates attention and performance monitoring to compute prediction errors. The third FMA profiles a self-
to-goal gradient that links salience processing and episodic memory to update brain prediction by weighting errors and
contextual relevance. Marmosets exhibit a homologous tripartite structure, suggesting evolutionary conservation of pre-
dictive coding motifs, albeit with species-specific topological variations. These findings unify predictive coding, allostasis,
and intrinsic activity into a signal-error-salience predictive modeling framework, where sensory integration, error computa-
tion, and salience modulation interact to optimize adaptive responses. By bridging neurocomputational theory with empir-
ical connectomics, this work establishes a cross-species blueprint of brain organization, offering experimental evidences or
insights into conserved mechanisms of free energy principles and dark energy explorations. These FMAs highlight how evo-
lution refines neural architectures while preserving core computational principles, paving the way for mechanistic studies
of predictive dysfunction and evolutionary drivers of primate brain complexity.

INTRODUCTION

The organization of brain activity and its relationship to cognition and behavior has increasingly been approached through the lens
of manifolds of connectivity or gradients [1,2]. This mathematical framework offers a way to analyze and represent complex neural
data in spacetime, providing information on the underlying structures and dynamics of the organization of the primate brain [3]. The
concept of cortical gradients refers to continuous spatial variations in connectomic profiles across the cortex. The brain organizes its
connectivity along these gradients, with functional regions smoothly transitioning from one state to another [4]. By modeling these
gradients as manifolds, researchers are now able to uncover how different brain regions interact and how they change in response
to different cognitive states or behaviors, with implications for linking brain structure with function [5].

Previous research highlights how gradients in neuron density, connectivity, and gene expression shape primate cortical organiza-
tion and suggests that gradients play a crucial role in brain function and offer insight into the evolution and development of brain
networks between species [6,7]. The cortical gradients derived using functional magnetic resonance (fMRI) data characterize the main
axes of continuous variations in brain connectivity across the cortex, a function manifold reflecting spatially organized patterns, where
the cortical regions are hierarchically or topographically organized and related along the gradients. We note that the cortical gra-
dient methods only keep a set of the most strong functional connectivity (e.g., the top 10% connectivity) for consideration [8]. This
threshold-based choice has derived consistent gradients of cortical organization between functional connectivity in the human
brain and structural connectivity in the monkey brain [4]. Beyond the tenet behind such a threshold choice to approximate the un-
derlying structural connectivity, the wave-like perspective of brain function indicated that all connectivity are meaningful properties
of cortical wave propagation [9,10]. Thus, a function manifold from the full fMRI connectivity data is essential to understand the corti-
cal organization of primate brain function but remains elusive.

Comparative and reproducible analysis of the function manifolds has been difficult due to challenges of performing fMRI in wakeful

nonhuman primate brains. In this work, we used two large-scale open datasets of resting-state fMRI (rfMRI) measurements from

the Human Connectome Project (HCP) and the Chinese HCP (CHCP) to reconstruct the function manifold in the human cortical

organization. Meanwhile, the same manifold analysis was also performed for wakeful marmoset brains based on two open rfMRI

datasets from the National Institute of Health (NIH) and the Institute of Neuroscience (ION) at the Chinese Academy of Sciences.

RESULTS

Figure 1: Top three gradient maps derived from the complete HCP connectivity matrix as axes of the functional manifold in the human cortical connectome. The left column (A,C,E) depicts the maps in the left hemisphere, and the right column (B,D,F) depicts the maps in the right hemisphere. The gradient values are averaged and ranked along spiral trajectories from the green-end inside to the yellow-end outside according to 200 parcellating units of each hemisphere [11]. These parcels are colored and rendered onto the human cerebral cortex in terms of the fifteen canonical cortical networks defined in previous studies [12,13]: Somatomotor-A (SMOT-A), Somatomotor-B (SMOT-B), Premotor-Posterior Parietal Rostral (PM-PPr), Action-Mode Network (AMN), Salience/Parietal Memory Network (SAL/PMN), Dorsal Attention-A (dATN-A), Dorsal Attention-B (dATN-B), Frontoparietal Network-A (FPN-A), Frontoparietal Network-B (FPN-B), Default-Mode Network-A (DMN-A), Default-Mode Network-B (DMN-B), Language (LANG), Visual Central (VIS-C), Visual Peripheral (VIS-P), and Auditory (AUD).

The Mega-ICA Group-PCA (MIGP) method [14] is used to construct a group-level connectome matrix from preprocessed rfMRI data,

which are described in previous publications [15–17]. It is designed to efficiently reduce the dimensionality of large-scale fMRI datasets

while preserving meaningful functional connectivity patterns across individuals. MIGP generates dense connectomes from HCP,

CHCP, NIH and ION datasets, respectively. These complete connectivity matrices are analyzed (that is, without the sparsity thresh-

old) using BrainSpace toolbox [8], producing a series of gradients on the organization of the cortical connectivity. Specifically, the top

three gradients are reproducible between independent datasets and explain a large portion of the observed variances in complete

brain connectivity maps (HCP: 20.6%, 14.0%, 6.0%; CHCP: 20.3%, 12.2%, 6.6%; NIH: 46.8%, 19.2%, 6.5%; ION: 36.6%, 26.5%, 7.2%).

We thus take the three gradients as a low-dimensional approximation of the high-dimension cortical connectivity space or its low-

dimension geometry. From this perspective, these gradients form a three-dimensional manifold on the cortical connectivity geom-

etry of the human brain function, that is, a function manifold. To visualize the functional manifold, we rescale each gradient to the same range from -1 to 1 and plot each vertex on the cortical surface as a point in the 3D space of the function manifold spanned by the three gradients as axes. The function manifold forms the surface of a regular tetrahedron or a functional representation in the human (HCP/CHCP, Fig. S1A/S2A) but not marmoset (NIH/ION, Fig. S1B/S2B).

| SMOT-d | PreMOT | VIS-m | ACC | highVN1 | highVN3 | orbFP | DMN | phTP | FPN | highVN2 | FPO | AUD | VIS-l | SMOT-v |

Figure 2: Top three gradient maps derived from the complete NIH connectivity matrix as axes of the functional manifold in the marmoset cortical connectome. The left column (A,C,E) depicts the maps in the left hemisphere, and the right column (B,D,F) depicts the maps in the right hemisphere. The gradient values are averaged and ranked along spiral trajectories from the green-end inside to the yellow-end outside according to 96 parcels in each hemisphere[17]. These parcels are rendered onto the marmoset cerebral cortex in terms of the fifteen canonical cortical networks defined in [17] using the same colorset as in the human cortical network parcellation: dorsal somatomotor (SMOT-d), ventral somatomotor (SMOT-v), premotor (PreMOT), medial primary visual and MT/MST (VIS-l), lateral primary visual and MT/MST (VIS-m), auditory and insular cortex (AUD), anterior cingulate cortex (ACC), frontal pole (FPO), high-level 1 visual-related network (highVN1), high-level 2 visual-related network (highVN2), high-level 3 visual-related network (highVN3), orbital frontal (orbFP), frontoparietal-like network (FPN), default-mode-like network (DMN), the parahippocampus/temporal pole (phTP).

To further delineate anatomy of the function manifold in the human cortex, we extract mean rescaled gradient values of all the vertices within each of the 200 parcelling units[11] assigned to fifteen canonical cortical networks defined in DU15NET[12] for each hemisphere, and plot them along a spiral trajectory in ascending order. As shown in Fig. 1A/1B, the first FMA (FMA-I) starts inside the spiral from the frontoparietal network (FPN-B) and default-mode network (DMN-B and DMN-A), and ends outside the spiral at the premotor-posterior parietal rostral network (PM-PPr), action-mode network (AMN)[13], dorsal attention network (dATN-B), visual network (VIS-P). The second FMA (FMA-II, Fig. 1C/1D) starts inside the spiral from DMN-B and DMN-A, language network (LANG), and ends outside the spiral at FPN-A, salience and parietal memory network (SAL/PMN), dATN-A. The third FMA (FMA-III) starts inside the spiral from SAL/PMN, DMN-B, AMN, dATN-A, and ends outside the spiral at FPN-A and FPN-B, DMN-A (Fig. 1E/1F). All the findings reported here are reproducible between HCP and CHCP (Fig. 1, Fig. S3).

We delineate the anatomy of the function manifold in the marmoset cortex in a similar vein by extracting mean rescaled gradient values of all vertices within each of the 96 parcelling units assigned to fifteen canonical cortical networks defined in[17]. FMA-I (Fig.

2A/2B) starts inside the spiral from the high-level 2 visual-related network (highVN2), frontoparietal-like network (FPN), lateral pri-

mary visual and MT/MST visual-related network (VIS-l), default-mode-like network (DMN), and ends outside the spiral at the audi-

tory and insular cortex salience-related network (AUD), premotor network (preMOT). FMA-II (Fig. 2C/2D) starts inside the spiral from

the high-level 1 visual-related network (highVN1), highVN2, orbital frontal network (orbFO), and ends outside the spiral at the dor-

sal somatomotor network (SMOT-d), FPN. FMA-III (Fig. 2E/2F) starts inside the spiral from the frontal pole network (FPO), preMOT,

high-level 3 visual-related network (highVN3), AUD, and ends outside the spiral at the dorsal and ventral somatomotor networks

(SMOT-d and SMOT-v). All the findings reported here are reproducible between NIH and ION (Fig. 2, Fig. S4).

DISCUSSION

The brain operates as a highly dynamic and self-organizing system governed by the principles of free energy and predictive coding,

where it continuously minimizes prediction errors through the constant comparison of sensory input with internal models [18]. This

enables the brain to anticipate and optimize responses to the environment, supporting allostasis, the regulation of internal states

for adaptive survival [19]. The brain exhibits the feature of the cost of "dark energy" through spontaneous intrinsic neural activity [20] to

maintain homeostasis, prepare for future events and facilitate communication between functional networks, and supports efficient

cognitive processing [12]. These processes, from inside-outside interactions between the brain and the body to top-down predictions

and bottom-up sensory integration, form a unified system that allows the brain to remain flexible, adaptive, and responsive to both

internal and external demands. Our findings, for the first time, provide three reproducible dimensions of functional space in such a

complex system as a basis to characterize its inside-out coordination observed by using fMRI to primate connectomes.

Although the three axes (that is, gradient maps) of this functional space conceptually represent distinct aspects of characterizing

the inside-outside organization in the primate cortex [21], it is quite open and challenging to determine or interpret their exact roles

in achieving predictive models of brain function. In human cortex, FMA-I has been documented as the sensorimotor-association,

unimodal-transmodal or external-internal gradient for a core feature of the human brain organization to support a domain-general

motif [4,22]. We argue that it serves as a signal-detection motif to capture the predictable information flow along the external en-

vironment and the internal environment to serve the prediction of allostasis [23]. FMA-II has a similar spatial profile to the multiple

demand system or the representation-modulation system [22], forming an error-calculation motif to compute predictive errors by

combining attention regulation, goal maintenance, strategy selection, performance monitoring with mental content. FMA-III has

rarely been reported by previous gradient studies, although several recent work demonstrated some sparse evidence linking it to

the recognition process of goal-directed and self-directed functions related to salience learning and episodic memory [24–26]. We

speculate on this axis as a salience-generation motif to support updates of the prediction by providing weights on both prediction

signals and errors. In the marmoset cortex, we note that a similar structure of the outside-inside functional space was observed. We

argue that the three axes revealed form a signal-error-salience predictive model in marmosets similar to humans but with a differ-

ent shape as in Figs. S1 and S2. However, understanding their comparative insights raises challenges and opportunities for future

studies [27] because fMRI studies in awakened animals are much limited compared to those in humans [28,29].

Our work produces comparative and reproducible low-dimensional representations of primate functional connectomes at the group

level. This addresses functional references in the brain connectivity space for individualized FMA mapping methods. The spacetime

dual regression [30] on these references will achieve reliable and valid FMA measurements at the individual level.

MATERIALS AND METHODS

The wakeful resting-state fMRI data are from the Human Connectome Project (HCP, N = 1003) [15] and the Chinese HCP (CHCP, N

= 217) [16] for healthy human young adults while the NIH (N = 26) and the ION (N = 13) resource from the Marmoset Brain Mapping

Project [17]. All our analyses are based on group average preprocessed whole brain dense functional connectome (.dconn) data iden-

tifying temporal correlations between all cortical vertices and subcortical voxels. Specifically, each preprocessed individual data set

was temporally trimmed and variance normalization applied and submitted to the principal component analysis (PCA) at the group

level (MIGP) [14]. The output of the group PCA with the spatial eigenvectors weighted in the upper N (N = 4500 for HCP, 2000 for

CHCP, N = 1800 for NIH and ION) is then renormalized, reweighted and correlated to form the average dense connectomes at the

group level ($91,282 \times 91,282$ entries for the HCP/CHCP and $76,457 \times 76,457$ entries for the NIH/ION). Diffusion map embedding is

applied to identify a set of low-dimensional manifolds (i.e., gradients) capturing principal dimensions of spatial variation in connec-

tivity. In particular, this algorithm is implemented in BrainSpace toolbox[8] without thresholding functional connectivity to derive the

functional affinity or similarity of the functional connectivity matrix. The first three gradients span a 3D functional manifold in the

wakeful connectome organization.

ACKNOWLEDGMENTS

This work has been supported by the 2030 scientific and technological innovation - the major project of Brain Science and Brain-Inspired Intelligence Technology (2021ZD0200500) and the Interdisciplinary Brain Database for In vivo Population Imaging (ID-BRAIN) at the National Basic Science Data Center. The authors thank Dr. Cirong Liu from the Institute of Neuroscience, Chinese Academy of Sciences for valuable discussions during the early preparation of the manuscript and sharing the marmoset dataset, and the National Basic Science Data Center for informatic support.

AUTHOR CONTRIBUTIONS

Conception and Design: X-X.X., X-N.Z. Data Analysis: X-N.Z. Initial Drafting of the Manuscript: X-X.X. Critical Review and Editing of the Manuscript: X-X.X., X-N.Z.

AUTHOR COMPETING INTERESTS

Competing interests
The authors declare that they have no competing interests.

DATA AVAILABILITY

MIGP-derived data will be deposited in the Chinese Color Nest Project (CCNP) - Lifespan Brain-Mind Development data Community in the Science Data Bank (`https://ccnp.scidb.cn/en`).

CODE AVAILABILITY

All the codes will be deposited in the Chinese Color Nest Project (CCNP) - Lifespan Brain-Mind Development data Community in the Science Data Bank (`https://ccnp.scidb.cn/en`).

Figure 3: 3D space of the functional manifold in primate cortical connectomes. Three axes are the three gradients of functional connectivity (gradient 1: red, gradient 2: green, gradient 3: blue). Given a vertex on the cortical surface, we plot a point in the 3D space with its coordinates as its three gradient values (Top panel: HCP, Bottom panel: NIH).

Figure 4: 3D space of the functional manifold in primate cortical connectomes. Three axes are the three gradients of functional connectivity (gradient 1: red, gradient 2: green, gradient 3: blue). Given a vertex on the cortical surface, we plot a point in the 3D space with its coordinates as its three gradient values (Top panel: CHCP, Bottom panel: ION).

SMOT-A | PM-PPr | VIS-P | AMN | dATN-B | LANG | FPN-B | DMN-B | DMN-A | FPN-A | SAL/PMN | dATN-A | AUD | VIS-C | SMOT-B

Figure 5: Top three gradient maps derived from the complete CHCP connectivity matrix as axes of the functional manifold in the human cortical connectome. The left column (A,C,E) depicts the maps in the left hemisphere, and the right column (B,D,F) depicts the maps in the right hemisphere. The gradient values are averaged and ranked along spiral trajectories from the green-end inside to the yellow-end outside according to 200 parcellating units of each hemisphere [11]. These parcels are colored and rendered onto the human cerebral cortex in terms of the fifteen canonical cortical networks defined in previous studies [12,13]: Somatomotor-A (SMOT-A), Somatomotor-B (SMOT-B), Premotor-Posterior Parietal Rostral (PM-PPr), Action-Mode Network (AMN), Salience/Parietal Memory Network (SAL/PMN), Dorsal Attention-A (dATN-A), Dorsal Attention-B (dATN-B), Frontoparietal Network-A (FPN-A), Frontoparietal Network-B (FPN-B), Default-Mode Network-A (DMN-A), Default-Mode Network-B (DMN-B), Language (LANG), Visual Central (VIS-C), Visual Peripheral (VIS-P), and Auditory (AUD).

| SMOT-d | PreMOT | VIS-m | ACC | highVN1 | highVN3 | orbFP | DMN | phTP | FPN | highVN2 | FPO | AUD | VIS-l | SMOT-v |

Figure 6: Top three gradient maps derived from the complete ION connectivity matrix as axes of the functional manifold in the marmoset cortical connectome. The left column (A,C,E) depicts the maps in the left hemisphere, and the right column (B,D,F) depicts the maps in the right hemisphere. The gradient values are averaged and ranked along spiral trajectories from the green-end inside to the yellow-end outside according to 96 parcels in each hemisphere [17]. These parcels are rendered onto the marmoset cerebral cortex in terms of the fifteen canonical cortical networks defined in [17] using the same colorset as in the human cortical network parcellation: dorsal somatomotor (SMOT-d), ventral somatomotor (SMOT-v), premotor (PreMOT), medial primary visual and MT/MST (VIS-l), lateral primary visual and MT/MST (VIS-m), auditory and insular cortex (AUD), anterior cingulate cortex (ACC), frontal pole (FPO), high-level 1 visual-related network (highVN1), high-level 2 visual-related network (highVN2), high-level 3 visual-related network (highVN3), orbital frontal (orbFP), frontoparietal-like network (FPN), default-mode-like network (DMN), the parahippocampus/temporal pole (phTP).

REFERENCES

[1] Julia M Huntenburg, Pierre-Louis Bazin, and Daniel S Margulies. Large-scale gradients in human cortical organization. *Trends in Cognitive Sciences*, 22(1):21–31, 2018.

[2] Rogier B Mars, Richard E Passingham, and Saad Jbabdi. Connectivity fingerprints: From areal descriptions to abstract spaces. *Trends in Cognitive Sciences*, 22(11):1026–1037, 2018.

[3] Christopher Langdon, Mikhail Genkin, and Tatiana A Engel. A unifying perspective on neural manifolds and circuits for cognition. *Nature Reviews Neuroscience*, 24(6):363–377, 2023.

[4] Daniel S Margulies, Satrajit S Ghosh, Alexandros Goulas, Marcel Falkiewicz, Julia M Huntenburg, Georg Langs, Gleb Bezgin, Simon B Eickhoff, F Xavier Castellanos, Michael Petrides, Elizabeth Jefferies, and Jonathan Smallwood. Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of National Academy of Sciences of the United States of America*, 113(44):12574–12579, 2016.

[5] James C Pang, Kevin M Aquino, Marianne Oldehinkel, Peter A Robinson, Ben D Fulcher, Michael Breakspear, and Alex Fornito. Geometric constraints on human brain function. *Nature*, 618(7965):566–574, 2023.

[6] Valerie J Sydnor, Bart Larsen, Danielle S Bassett, Aaron Alexander-Bloch, Damien A Fair, Conor Liston, Allyson P Mackey, Michael P Milham, Adam Pines, David R Roalf, Jakob Seidlitz, Ting Xu, Armin Raznahan, and Theodore D Satterthwaite. Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. *Neuron*, 109(18):2820–2846, 2021.

[7] Hao-Ming Dong, Daniel S Margulies, Xi-Nian Zuo, and Avram J Holmes. Shifting gradients of macroscale cortical organization mark the transition from childhood to adolescence. *Proceedings of National Academy of Sciences of the United States of America*, 118(28):e2024448118, 2021.

[8] Reinder Vos de Wae, Oualid Benkarim, Casey Paquola, Sara Lariviere, Jessica Royer, Shahin Tavakol, Ting Xu, Seok-Jun Hong, Georg Langs, Sofie Valk, Bratislav Misic, Michael Milham, Daniel Margulies, Jonathan Smallwood, and Boris C Bernhardt. Brainspace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Communications Biology*, 3(1):103, 2020.

[9] Taylor Bolt, Jason S Nomi, Danilo Bzdok, Jorge A Salas, Catie Chang, B T Thomas Yeo, Lucina Q Uddin, and Shella D Keilholz. A parsimonious description of global functional brain organization in three spatiotemporal patterns. *Nature Neuroscience*, 25(8):1093–1103, 2022.

[10] Lyle Muller, Frederic Chavane, John Reynolds, and Terrence J Sejnowski. Cortical travelling waves: mechanisms and computational principles. *Nature Reviews Neuroscience*, 19(5):255–268, 2018.

[11] Alexander Schaefer, Ru Kong, Evan M Gordon, Timothy O Laumann, Xi-Nian Zuo, Avram J Holmes, Simon B Eickhoff, and B T Thomas Yeo. Local-global parcellation of the human cerebral cortex from intrinsic functional connectivity mri. *Cerebral Cortex*, 28(9):3095–3114, 2018.

[12] Jingnan Du, Lauren M DiNicola, Peter A Angeli, Noam Saadon-Grosman, Wendy Sun, Stephanie Kaiser, Joanna Ladopoulou, Aihuiping Xue, B T Thomas Yeo, Mark C Eldaief, and Randy L Buckner. Organization of the human cerebral cortex estimated within individuals: networks, global topography, and function. *Journal of Physiology*, 131(6):1014–1082, 2024.

[13] Nico U F Dosenbach, Marcus E Raichle, and Evan M Gordon. The brain's action-mode network. *Nature Reviews Neuroscience*, 26(3):158–168, 2025.

[14] Stephen M Smith, Aapo Hyvarinen, Gael Varoquaux, Karla L Miller, and Christian F Beckmann. Group-pca for very large fmri datasets. *NeuroImage*, 101:738–749, 2014.

[15] Matthew F Glasser, Stephen M Smith, Daniel S Marcus, Jesper L R Andersson, Edward J Auerbach, Timothy E J Behrens, Timothy S Coalson, Michael P Harms, Mark Jenkinson, Steen Moeller, Emma C Robinson, Stamatios N Sotiropoulos, Junqian Xu, Essa Yacoub, Kamil Ugurbil, and David C Van Essen. The human connectome project's neuroimaging approach. *Nature Neuroscience*, 19(9):1175–1187, 2016.

[16] Jianqiao Ge, Guoyuan Yang, Meizhen Han, Sizhong Zhou, Weiwei Men, Lang Qin, Bingjiang Lyu, Hai Li, Haobo Wang, Hengyi Rao, Zaixu Cui, Hesheng Liu, Xi-Nian Zuo, and Jia-Hong Gao. Increasing diversity in connectomics with the chinese human connectome project. *Nature Neuroscience*, 19(9):1175–1187, 2023.

[17] Xiaoguang Tian, Yuyan Chen, Piotr Majka, Diego Szczupak, Yonatan Sanz Perl, Cecil Chern-Chyi Yen, Chuanjun Tong, Furui Feng, Haiteng Jiang, Daniel Glen, Gustavo Deco, Marcello G P Rosa, Afonso C Silva, Zhifeng Liang, and Cirong Liu. An integrated resource for functional and structural connectivity of the marmoset brain. *Nature Communications*, 13(1):7416, 2022.

[18] Chang Sub Kim. Free energy and inference in living systems. *Interface Focus*, 13(3):20220041, 2023.

[19] Gyorgy Buzsaki. *The Brain from Inside Out*. Oxford University Press, 06 2019.

[20] Marcus E Raichle. Neuroscience. the brain's dark energy. *Science*, 14(5803):1249–1250, 2006.

[21] Claus C Hilgetag, Alexandros Goulas, and Jean-Pierre Changeux. A natural cortical axis connecting the outside and inside of the human brain. *Network Neuroscience*, 6(4):950–959, 2022.

[22] Jiahe Zhang, Olamide Abiose, Yuta Katsumi, Alexandra Touroutoglou, Bradford C Dickerson, and Lisa Feldman Barrett. Intrinsic functional connectivity is organized as three interdependent gradients. *Scientific Reports*, 9(1):15976, 2019.

[23] Yuta Katsumi, Jordan E Theriault, Karen S Quigley, and Lisa Feldman Barrett. Allostasis as a core feature of hierarchical gradients in the human brain. *Network Neuroscience*, 6(4):1010–1031, 2022.

[24] Annie Zheng, David F Montez, Scott Marek, Adrian W Gilmore, Dillan J Newbold, Timothy O Laumann, Benjamin P Kay, Nicole A Seider, Andrew N Van, Jacqueline M Hampton, Dimitrios Alexopoulos, Bradley L Schlaggar, Deanna J Sylvester, Chad M amd Greene, Joshua S Shimony, Steven M Nelson, Gagan S Wig, Caterina Gratton, Kathleen B McDermott, Marcus E Raichle, Evan M Gordon, and Nico U F Dosenbach. Parallel hippocampal-parietal circuits for self- and goal-oriented processing. *Proceedings of National Academy of Sciences of the United States of America*, 118(34):e2101743118, 2021.

[25] Young Hye Kwon, Joseph J Salvo, Nathan L Anderson, Donnisa Edmonds, Ania M Holubecki, Maya Lakshman, Kwangsun Yoo, B T Thomas Yeo, Kendrick Kay, Caterina Gratton, and Rodrigo M Braga. Situating the salience and parietal memory networks in the context of multiple parallel distributed networks using precision functional mapping. *Cell Reports*, 44(1):115207, 2025.

[26] Peter A Angeli, Lauren M DiNicola, Noam Saadon-Grosman, Mark C Eldaief, and Randy L Buckner. Specialization of the human hippocampal long axis revisited. *Proceedings of National Academy of Sciences of the United States of America*, 122(3):e2422083122, 2025.

[27] Marco Pagani, Daniel Gutierrez-Barragan, A. Elizabeth de Guzman, Ting Xu, and Alessandro Gozzi. Mapping and comparing fmri connectivity networks across species. *Communications Biology*, 6(1):1238, 2023.

[28] Francesca Mandino, Stella Vujic, Joanes Grandjean, and Evelyn M R Lake. Where do we stand on fmri in awake mice? *Cerebral Cortex*, 34(1):bhad478, 2024.

[29] Chuanjun Tong, Cirong Liu, Kaiwei Zhang, Binshi Bo, Ying Xia, Hao Yang, Yanqiu Feng, and Zhifeng Liang. Multimodal analysis demonstrating the shaping of functional gradients in the marmoset brain. *Nature Communications*, 13(1):6584, 2022.

[30] Hui-Jie Li, Yin-Shan Wang, Xuequan Su, and Xi-Nian Zuo. Functional brain network mapping with dual regression. *Science China Life Sciences*, 60(12):1450–1452, 2017.