Kinetic mechanisms for the sequence dependence of transcriptional errors

Tripti Midha[a], Anatoly B. Kolomeisky[a,b,c,d,1], and Oleg A. Igoshin[a,b,e,f,1]

[a]Center for Theoretical Biological Physics, Rice University, Houston TX 77005
[b]Department of Chemistry, Rice University, Houston, TX 77005
[c]Department of Chemical and Biomolecular Engineering, Rice University, Houston, TX 77005
[d]Department of Physics and Astronomy, Rice University, Houston, TX 77005
[e]Department of Bioengineering, Rice University, Houston, TX 77005
[f]Department of Biosciences, Rice University, Houston, TX 77005

The fidelity of template-dependent mRNA synthesis during transcription elongation is the primary determinant of accurate gene expression and the maintenance of functional RNA transcripts. However, the mechanisms governing transcription fidelity remain incompletely understood. While previous studies have characterized how error rates vary with nucleotide identity at upstream and downstream positions from the incorporation site, the comprehensive microscopic explanation of this sequence dependence has not been elucidated. In this study, we develop a novel theoretical approach that integrates transcription proofreading mechanisms and inhomogenous DNA sequence effects. Using first-passage analysis validated by Monte Carlo simulations, we quantitatively characterize nucleotide-specific error rates during RNA polymerase II transcription. The model accurately reproduces experimental error rates and predicts kinetic parameters influencing transcriptional fidelity. Analysis reveals nucleotide incorporation rates follow the hierarchy U<C<G<A, consistent with independent experimental observations. Notably, our model not only explains how the error rates depend on the nature of the base immediately downstream (+1) but also predicts that the identity of the nucleotide at the second downstream position (+2) also plays an important role. Pyrimidines at position +2 contribute to lower error rates than purines, whereas the third downstream base (+3) has no effect. These previously unreported correlations are corroborated by bioinformatic analysis of existing datasets. In addition, using the BRCA1 gene as an example, we explore the physiological implications of sequence-dependent error rates, identifying an increased likelihood of premature stop codon errors. These findings clarify how DNA sequence context modulates nucleotide incorporation kinetics, advancing our understanding of transcriptional fidelity and its functional consequences.

Transcription elongation | template-dependent error | error probability

Correspondence: [1]*To whom correspondence should be addressed. E-mails: tolya@rice.edu, igoshin@rice.edu*

Introduction

The high fidelity of the transcription elongation process by RNA polymerase (RNAP) is crucial for the faithful gene expression (1). During transcription elongation, RNAP facilitates the addition of ribonucleotides (A, U, G, C) complementary to the DNA nucleotides (A, C, G, T) to the growing RNA chain by creating phosphodiester bonds. The process follows the Watson-Crick (WC) base pairing rules, implying that the purines (A, G) and pyrimidines (C, U) on the mRNA must make complementary base pairings with respective pyrimidines (T, C) and purines (G, A) on the DNA (see Fig. 1A) (2, 3). For accurate mRNA synthesis, the RNAP must distinguish the correct WC pairings (A:T, G:C, C:G, and U:A), denoted as the right, R, over the incorrect WC pairings, denoted as wrong, W, in the mRNA chain (see Fig. 1A) (3).

Despite the inherent complexity of involved biochemical processes, RNAP achieves a remarkably low error rate, i.e., a fraction of incorrect pairings. In many biocatalytic processes, including transcription, the high accuracy arises due to out-of-equilibrium error-correcting mechanisms known as kinetic proofreading (KPR) (4, 5). KPR allows RNAP to kinetically discriminate between R and W substrates through proofreading steps (6, 7). RNAP possesses a spectrum of intrinsic proofreading tools, including pausing, backtracking, and endonucleolytic cleavage of dinucleotides, which do not rely on external energy sources (8–11). Together, these mechanisms enable to maintain high fidelity essential for cellular function (12–15).

Theoretically, the mechanisms of transcription elongation by RNAP have been well explored using different approaches such as Brownian ratchet, thermal ratchet (16, 17), and templated-copolymerization methods (7, 17–19). In the templated-copolymerization method, the elongation process is described using a homogenous Markov chain (18–22). Various other approaches, such as the Markov-chain copolymerization theory and linear-decoupling method, under steady-state conditions, have successfully predicted the fidelity and speed of the elongation process. However, all these methods considered the elongation process only as an R/W binary copolymerization process and, therefore, ignored the effects of nucleotide identity on the elongation and proofreading process. However, in reality, the DNA templates are heterogeneous, and each right and wrong pairing might have its own proofreading dynamics.

Recent experimental studies employing high-throughput techniques, such as circle sequencing, demonstrate that DNA template sequences strongly influence transcription error rates (23). While the overall trends are somewhat complex, it is clear that, compared to pyrimidines, purines at the next downstream position on mRNA (+1) increase the error rate at the current position (0) (23). Several recent studies on transcription elongation by the RNAP enzyme indicated that the incorporation rate of nucleotides follows the

order U<C<G<A, suggesting that the incorporation rate of pyrimidines is slower than that of purines (16, 24). Furthermore, purines, double-ring structured nucleotides, have wider stacking areas than single-ring pyrimidines. This could increase their interaction strength with the active site of RNAP and influence their incorporation kinetics (25, 26). However, there is a lack of a theoretical understanding of the effect of nucleotide-specific kinetic rates on transcription fidelity for inhomogeneous DNA templates. Therefore, it is unclear if these kinetic differences can fully explain the sequence-dependence of error.

The sequence dependence of errors for inhomogeneous templates in DNA replication has been considered by several theoretical approaches, such as iterated functions and first-passage methods (27–30). These theories captured the potential role of nucleotide-specific kinetic parameters in controlling template-dependent fidelity (27–30). However, unlike DNA replication, transcription elongation involves more complex proofreading mechanisms involving backtracking and dinucleotide cleavage. Furthermore, there are fewer precise kinetic measurements for sequence-dependent transcription parameters. These reasons make theories for sequence-dependence of transcriptional errors more difficult to formulate.

In this paper, we develop a theoretical framework based on first-passage analysis that computes the template-specific transcription error rate and makes testable predictions of the kinetic parameters affecting transcription fidelity. First, in the context of the coarse-grained two-letter base alphabet (purines and pyrimidines), we examine whether our theoretical model can capture the experimentally observed sequence-dependent error rates (Fig. 2). We then use our theoretical approach to predict how the transcription fidelity is affected by long-range downstream neighbors, by extending the theory and checking the predictions by reanalyzing the data in Ref.(23) (Fig. 3). The results are utilized to predict sequence-dependent variation in kinetic parameters governing nucleotide incorporation and cleavage (Fig. 4). Furthermore, we also perform bioinformatics analysis to examine the physiological implications of sequence-dependent errors (Fig. 5). By examining the impact of synonymous errors and errors generating premature stop codons, our study provides insights into how transcription fidelity impacts functionality of living systems. Our study bridges theoretical understanding with experimental findings to unravel the complex mechanisms of transcription elongation.

Results

A. Nucleotide-specific kinetics and template heterogeneity modulate transcription elongation error rates.
To analyze the effect of the chemical identity of the current base (0) and the chemical identity of the base immediately downstream (+1) on the error, we first employ an approximate biophysical model of transcription elongation (see Fig. S1, SI Appendix) with kinetic parameters as detailed in Table S1 (see Methods, SI Appendix). First, for simplicity, we aggregate the 4-letter base alphabet of nucleotides into the

Fig. 1. (A) (A) Watson-Crick (WC) base pairing rules between nucleotides on the DNA template and mRNA sequence during the transcription process with four nucleotides. (B) Watson-Crick (WC) base pairing rules in a transcription process with two components: purines (Pu), and pyrimidines (Py). (C) Schematic representation of the biophysical model for the transcription elongation process with effective forward, backward, and dinucleotide cleavage reactions.

Fig. 2. (A) Variation of error rate at position 0 as a function of the incorporation factor, γ, for four dimers occupying positions at 0 and +1 on mRNA for the model with two components. (B) Variation of the ratio of the error rates with purines at current position, 0, to the pyrimidines as a function of the incorporation factor, γ, for a fixed monomer at the next downstream position +1. (C) Ratio of the error rates with purines to the pyrimidines at the next downstream (+1) position as a function of the incorporation factor, γ, for a fixed monomer at the current position, 0. (D) Fitting (shaded region) of theoretical errors (red bars) from the model with first-order neighbor effects to the experimental data (blue bars). Theoretical error (red bars) at the optimal kinetic parameters for the unfitted (unshaded region) dimers aligns with the corresponding experimental data (blue bars). Error bars denote the standard error mean for the experimental data for the six samples (23).

2-letter description, i.e., we consider that both the DNA template and mRNA copy sequence consist of only two types of monomers: purines (Pu=A, G) and pyrimidines (Py =T, U/C). In that picture, it is also assumed that the correct (R) copying corresponds to associating Pu to Py, whereas the wrong (W) copying corresponds to associating Pu to Pu or Py to Py (see Fig. 1A, Fig.1B). It is important to note here that, as per the WC pairing rules, Pu:Pu and Py:Py bonds are always wrong, however, not all Pu:Py bonds are right (e.g.

U:G). The limitation of the two-letter alphabet requires us to assume Pu:Py bonds are always right pairing, and this assumption allows us to pinpoint the simple intuitive basis of the sequence-dependence of errors. A more precise theoretical model with all four bases will be analyzed in a later section.

The heterogeneity in the DNA template is introduced by assuming distinct incorporation rates of pyrimidines and purines, quantified by a non-dimensional scale-factor γ, i.e., $k_{ij}^{Py} = \gamma k_{ij}^{Pu}$. Here, the subscripts $i, j \in \{R, W\}$ are two consecutive nucleotides on the mRNA chain, and the superscript denotes the nature of the incorporating nucleotide j, i.e, whether it is Pu or Py (see Methods for details). Using the first-passage analysis detailed in the Methods section, we numerically compute the template-dependent error rate, η, at the polymerase current position, 0. That error can be compared with the experimental data from Ref.(23) aggregated in the 2-letter description.

To predict how the difference in the nucleotide-specific incorporation rates affects the transcriptional error, we use the first-passage approach to compute the error rates for the four possible (Pu-Pu, Pu-Py, Py-Pu, Py-Py) nucleotide pairs (dimers) in the consecutive positions $(0, +1)$ in the mRNA sequence. The results are shown in Fig. 2A as a function of γ. When $\gamma = 1$, which corresponds to a homogeneous DNA template, errors are the same for all dimers. For $\gamma > 1$, i.e., when the pyrimidines are incorporated faster than purines, the error η is highest for Pu-Py dimer and lowest for the Py-Pu dimer. Conversely, for $\gamma < 1$, η is highest for Py-Pu and lowest for the Pu-Py dimers. Monte Carlo simulations further confirm the accuracy of our predictions for heterogeneous DNA templates (Fig. 2A, lines vs symbols) (31). Notably, alternative approaches such as copolymerization(20) and linear decoupling(19) cannot correctly identify the error unless $\gamma = 1$, i.e., unless there is no sequence-dependence of polymerization rates. Thus, our theory predicts two effects: slower incorporation of a base at the current position increases the probability of error, whereas slower incorporation of a base at the next position decreases the error.

Notably, the effects of current-base and next-base kinetics appear to be independent of each other. To explore the impact of the chemical nature of incorporating nucleotides at the current position, we compute the ratio of errors for dimers with Pu at position 0 to those with Py at position 0 (Fig. 2B). As one can see, this ratio remains the same regardless of the nucleotide at the $+1$ position. Similarly, to examine the influence of the nucleotide identity at the downstream position, we compute the ratio of errors for dimers with Pu at position $+1$ to those with Py at position $+1$ (Fig. 2C). This ratio does not depend on the identity of the nucleotide at the position 0. Kinetic effects can be understood intuitively in the following way. For example, for $\gamma < 1$, the error rate is lower for purines than for pyrimidines in the current position because the incorporation of the wrong nucleotide (i.e. pyrimidine) is slower than that of the correct nucleotide (Fig. 2B). On the other hand, the error rate is lower for pyrimidines than the purines at the downstream base (Fig. 2C) because the

slower bases at the downstream position give enough time to the polymerase to correct the error by cleaving nucleotides in case of misincorporations, thereby increasing the accuracy. Furthermore, the independence of the effects of the current and downstream bases (collapsing of the curves in Fig. 2B and Fig. 2C) can also be explained theoretically (see SI Appendix).

To quantitatively determine how the kinetic parameters for purines and pyrimidines differ, we fit theoretical predictions for the errors to corresponding experimental data. Note that the experimental data is known for the 64 cases, i.e., the error rates for nucleotides A, C, G, and U at a position (say) 0 for given nucleotides at the upstream (-1), and the downstream (+1) positions. We group them into four cases (Pu-Py, Py-Py, Pu-Pu, Py-Pu) occupying positions 0 and +1, as detailed in the SI Appendix. Moreover, to enable efficient parameter fitting, we utilize the approximate analytical expression of the theoretical error rate obtained under the biologically relevant conditions as a function of various kinetic parameters (see Methods, Eq. 5). This approximation (Eq. 5) agrees well with the results from the first-passage analysis (see Fig. S2). Given the sensitivity of transcription elongation errors to nucleotide incorporation and cleavage rates (7), we fix most of the parameters using previously determined values(7) as summarized in Table S1 and focus on fitting only three critical parameters: the incorporation rate for purines (k_{RR}^{Pu}), the pyrimidine-to-purine incorporation rate ratio (γ) such that $k_{RR}^{Py} = \gamma k_{RR}^{Py}$, and pyrimidines to purines dinucleotide cleavage rates ratio, ϕ, such that $r_{Py} = \phi r_{Pu}$. Here, r_i denotes the dinucleotide cleavage rates of misincorporated nucleotide when the correct one is i =Pu or Py). Using the least-squares optimization method weighted by the inverse of the standard deviation of experimental data, we fit theoretical error rates for Pu-Py and Py-Py dimers (shaded region in Fig. 2D) to experimental error rates by minimizing their differences. The results, displayed in Fig. 2D, show that the model successfully reproduces experimental error rates for dimers: Pu-Py, Py-Py, at the optimal parameter values: $k_{RR}^{Pu} = 100 s^{-1}$, $\gamma = 0.65$, $\phi = 2.0$, obtained from the fit. Furthermore, error rates for the other two dimers not used in fitting, Pu-Pu and Py-Pu, are calculated using these parameters and fall within experimental uncertainty (unshaded region in Fig 2D). These findings indicate that pyrimidines are incorporated about 35% slower than purines (γ=0.65), but their cleavage rate is 2x higher. Note that a purely non-nucleotide-specific proofreading mechanism, i.e. when $\phi = 1$, fails to fully account for the experimental data (see Fig. S3).

B. Theoretical model and bioinformatics analysis reveal reduced transcription error with pyrimidines at the second downstream position on mRNA.
During the proofreading process, RNAP has two possibilities to correct the error as it pauses twice – after the misincorporation of a nucleotide and after incorporating the following one. Each pause can result in dinucleotide cleavage, correcting the error (10). At each pause, there is a competition between the error correction and incorporation of the next base(7). Thus,

sequence-dependence of incorporation rate couples the identity of the nucleotide at the second downstream position (+2) and the error rate. To predict these effects theoretically, we generalize our theory to account for second-order effects, i.e., we now consider the dependence of kinetic rates on the presence of errors in the two preceding bases and the nature of the incorporated bases (SI Appendix, Fig. S4). The error rate for the more advanced model can also be computed numerically using the first-passage approach or with an approximate analytical formula(see Eq. 6, and SI Appendix).

To quantify the influence of the second downstream nucleotide, we compute the ratios of the errors with Pu at +2 position, $\eta(\mathrm{Pu}@+2)$, to that of Py at +2 position, $\eta(\mathrm{Py}@+2)$, for all four possible dimers at preceding positions. Fig. 3A shows the results as a function of γ. We observed that for $\gamma < 1$ (as indicated in the previous section), these error ratios consistently exceed 1. Thus, the error rate is higher when purines are at the +2 position because the slower incorporation of pyrimidines at +2 allows more time for error correction. This effect is independent of the nucleotide identity at position 0 but is influenced by the identity at position +1. Specifically, this error ratio is greater when Py occupies position +1 than Pu. Consequently, the error is lowest for the Py-Py combination at positions +1 and +2 (SI Appendix, Fig. S5).

It is important to note that while the authors of Ref. (23) did not report the effect of the +2 position on error, their accumulated data allows us to quantify these effects and test our theoretical predictions. To this end, we conducted a bioinformatics analysis of experimental data from Ref. (23) (see Fig. S6A). In Fig. 3B, we plot the ratio of the experimental error rates with Pu at +2 position to those with Py at +2 position for all four possible dimers at positions (0,+1). Across all dimers, the ratio exceeds 1, indicating that pyrimidines at the second downstream position reduce the error as predicted by our theoretical results. Thus we demonstrate the influence of the second downstream base pair on the error.

Since RNAP cleaves only dinucleotides, it is expected that the identity of the nucleotide at the third downstream (+3) position from the site of incorporation (0) should not significantly impact the error rate. To test this hypothesis, we compute the ratio of the error with purines at +3 downstream position to the error with pyrimidines at the same location using Eq. 6. Indeed, the ratio is 1 if the identities of the nucleotides at the preceding positions are the same (see Fig. S7A). To validate these theoretical results we again performed bioinformatics analysis of the experimental errors as a function of the identity of the nucleotide at position +3. The results show that the ratio of the errors with purines and pyrimidines at +3 position is always 1 within the error bar range (see Fig. S7B). These results indicate that the transcriptional error rate has correlations only up to the second base, reflecting the processes utilized in proofreading (namely, dinucleotide cleavage).

C. Predictions of the long-range genetic-context dependent errors align with experimental data. Next, we asked if our more advanced theoretical model could explain

Fig. 3. (A) Variation in the ratio of the theoretical error rates with Pu at +2 position to that of Py at +2 position for various dimers at 0, +1 position with changes in the incorporation factor, γ. **(B)** Ratio of the experimental error rates with Pu at +2 position to that of Py at +2 position for various dimers at 0, +1 position.

all corresponding experimental data and predict which kinetic parameters are sequence-dependent. For this purpose, we fit the theoretical errors from Eq. (6) for four (Pu-Pu-Pu, Pu-Pu-Py, Py-Pu-Pu, Py-Pu-Pu) of the eight trimers occupying positions (0,+1,+2) to the corresponding experimental data. Note that η in Eq. 6 is the function of various kinetic parameters. However, we choose to optimize the same kinetic parameters as in Fig. 2 fit, i.e. the incorporation rate for purine k_{RRR}^{Pu}, γ such that $k_{RRR}^{\mathrm{Py}} = \gamma k_{RRR}^{\mathrm{Pu}}$, and the dinucleotide cleavage rate of misincorporation instead of correct Pu, denoted by, r_{Pu}. Here, in k_{RRR}^{y}, the superscript $y \in \{\mathrm{Pu}, \mathrm{Py}\}$ denotes the nature of the incorporating nucleotide whereas the subscript denotes a trimer in the mRNA chain. The remaining parameters are fixed as described in Table S2.

The results in Fig. 4A indicate that for the optimal kinetic parameters, the theoretical errors are in excellent agreement with the corresponding experimental data for all the fitted trimers: Pu-Pu-Pu, Pu-Pu-Py, Py-Pu-Pu, Py-Pu-Pu (shaded region in Fig. 4A) as well as for the remaining predicted trimers: Pu-Py-Pu, Pu-Py-Py, Py-Py-Pu, Py-Py-Pu (unshaded region in Fig. 4A). Furthermore, the estimate for the incorporation rate parameter k_{RRR}^{Py}, is found to be close to the experiments and the previously discussed first-order model (10).

One can see that our model naturally captures the effect of the downstream neighbors on the error rate, but the identity of the upstream neighbor doesn't directly affect the error (see Fig. S6B). This is in contrast to our bioinformatics analyses of the experimental data of Ref. (23) (see Fig. S6A). This is because, in our model, the incorporation rate of any nucleotide at the current position (0) depends on whether the nucleotide at the upstream position (-1) is right or wrong but not on its chemical identity. Thus, to account for the observed effects, the identity of the upstream base pair should also affect the incorporation rate. However, how the upstream base affects the incorporation rate is not known. Therefore, we choose the simplest implementation, in which the incorporation rates of any nucleotide at position 0 is multiplied by the factor ψ if its upstream (-1) neighbor is a pyrimidine. The resulting approximate expression for the second-order model with two components, $\eta(-1, 0, +1, +2)$ (Eq. 6), will also depend on parameter ψ. To estimate the parameter ψ, we use

Fig. 4. (A) Fitting of theoretical error rates $\eta_2^S(0,+1,+2)$ (red bars) depending on nucleotide identity at positions 0,+1,+2 to the corresponding experimental data (blue bars). (B) Fitting of theoretical error rates (red bars) $\eta_2^S(-1,0,+1,+2)$ depending on the monomers at the -1, 0,+1,+2 positions to the corresponding experimental data (blue bars). (C) Fitting of theoretical error rates $\eta_4^S(0,+1,+2)$ for the second-order biophysical model with four components to the corresponding experimental data. Inset shows the average percentage of error for fitted, experimental, and unfitted data.

the data on error rate depending on the identity of four positions $(-1, 0, +1, +2)$ and fit half of these values to our theoretical predictions (shaded region of Fig. 4B) using the non-linear weighted least square method. At the optimal value of $\psi \approx 0.88$, the fitted theoretical error (shaded region) and predicted theoretical error (unshaded region of Fig. 4B) are in reasonable agreement with the experimental data. Thus, we predict a slight 12% deceleration of the incorporation rate when the upstream (-1) nucleotide is a pyrimidine.

We next evaluate if our model generalized to a 4-letter description, i.e., with four monomers (A, T, C, G) on the DNA template binding with the corresponding nucleotides (U, A, G, C) on the mRNA sequence (see Fig. 1A), can explain the corresponding experimental data and predict the incorporation rates of individual nucleotides. For this purpose, we computed the error rates under the biologically relevant condition (see SI Appendix), as given in Eq. (7). The the-

oretical error rate is a function of various parameters (given in Table S3), but we choose to find the optimal value of the k_{RRR}^A for nucleotide A, and the ratios γ_1, γ_2, γ_3 describing relative incorporation rates for other nucleotides, such that $k_{RRR}^U = \gamma_1 k_{RRR}^A$, $k_{RRR}^C = \gamma_2 k_{RRR}^A$, and $k_{RRR}^G = \gamma_3 k_{RRR}^A$. We also find the optimal fit for the cleavage rates of nucleotides in place of correct A, C, G, and U, denoted by r_A, r_C, r_G, and r_U, respectively. Using the non-linear weighted least square approximation method, we fit the theoretical error for analytical expression of error rate under the biologically relevant conditions for every fourth trimer in the 64 combinations (shown by the shaded trimers in Fig. 4C) as given in Eq. 7, to the corresponding experimental data in Fig. 4C. The optimal values for the parameters obtained from the fitting are given in Table S4. At these optimal values, the prediction for the trimers not used for fitting (unshaded trimers in Fig. 4C) also agrees well with the experimental

data with an average relative percentage of the error to be approximately 22 %. The fitting results indicate that the incorporation rates of nucleotides are in the following order: U<C<G<A, which also agrees well with the experimental findings ([16], [24]). Moreover, our fitting analysis reveals that the cleavage rate for misincorporations replacing correct pyrimidines (C, U) is higher than that for misincorporations replacing correct purines (A, G), aligning with the previously discussed two-letter alphabet model. Note that for the fitting we assume that the cleavage rates depend on the nature of the correct nucleotide being replaced by any misincorporated nucleotide. The fitting results for the case when cleavage rates depend on the nature of the misincorporated nucleotide are shown in Fig. S8, SI Appendix.

D. Sequence-dependent error increases the likelihood of premature terminating codon errors. To investigate the physiological significance of the sequence-dependent error rate, we asked how these effects might influence proteins after translation. As an example, we decided to focus on the 22 coding exons (coding sequences (CDS)) of the BRCA-1 gene, transcribed by RNAP. The human BRCA-1 gene has several isoforms encoding proteins of different lengths. For instance, the BRCA1-201 encodes the full-length protein consisting of 1863 amino acids and plays a central role in DNA repair and tumor suppression. Transcriptional errors can have distinct effects on the proteins ([32], [33]). On the one hand, it can change a codon to that encoding for the same amino acid (e.g. UCU (serine) changes to UCC (serine)), resulting in synonymous errors; on the other hand, it can change it to a stop codon (e.g., CAA (Glutamine) changes to UAA (a stop codon)), resulting in premature termination. We first investigated the physiological implication for the sequence-dependent errors for different nucleotides at positions -1, 0, and +1. For this purpose, we examined the error probability, E, defined as the fraction of the BRCA1 transcripts affected by premature stop errors or synonymous errors. Assuming that the error rates for each nucleotide are independent, for each value of the error rate, we computed the sequence-dependent error probability $E_{gc}(-1,0,+1) = 1 - \Pi_j(1 - n_{ijk}\eta_{ijk})$, of the occurrence of at least one error in the gene. Here, the product runs over index j, representing each nucleotide in the sequence that leads to any error, i denotes the 5' (upstream) neighbor of j, and k denotes the 3' downstream neighbor of j. Parameters n_{ijk}, and η_{ijk}, respectively, denote the number of errors and the error rate of the nucleotide j in the sequence (i,j,k). We compare this error probability with a weighted average error rate for the BRCA-1 gene, given by $E_{avg} = 1 - \Pi_j(1 - n_j\eta_{avg}(-1,0,1))$. Here, $\eta_{avg}(-1,0,1)$ is computed as the weighted average of the sequence-dependent error rate ($\eta(-1,0,+1)$) of all nucleotides in the genome. We then found the normalized error probability, E_{avg}, of all isoforms of the BRCA1 gene by dividing the error probability by the length of the amino acid sequences. Figure 5A displays the ratio of the mean of these normalized error probabilities, $\frac{E_{gc}(-1,0,+1)}{E_{avg}(-1,0,+1)}$ for both premature stop errors and syn-

onymous errors, where η_{ijk} is taken from the experimental data. We found the ratio to be always greater than 1 but the effect is much stronger for the premature stop errors. It is also determined that this observation is statistically significant by computing their p-values, which are quite small as indicated (see Fig. 5A). A similar behavior on the ratio of the error probabilities is observed when the long-range genetic context (-1,0,+1,+2) dependent error rates are considered (Fig. 5B). This indicates that the sequence-dependent error, whether short-range or long-range, increases the likelihood of premature termination in the encoded protein.

Fig. 5. (A) Ratios of the error probability with experimental error depending on immediate downstream and upstream sequence to the error probability with non-sequence context dependent errors for the BRCA-1 gene. (B) Ratios of the error probability with experimental error depending on two immediate downstream neighbors and one upstream neighbor to the error probability with non-sequence context-dependent errors for the BRCA-1 gene.

Discussion

In this work, we developed a theoretical framework to quantify the error rate of RNAP by explicitly considering the sequence dependence of transcription kinetics. We first considered a simplified model with first-order neighbor effects with only two types of components: purines and pyrimidines, on both the mRNA sequence and the DNA template (Fig. 1). The sequence-dependent error rate was computed using the first-passage method, and it aligns well with the Monte Carlo simulations (Fig. 2). Further fitting of theoretical predictions against experimental data shows a good quantitative agreement, supporting the model's ability to capture the sequence-dependent errors observed in transcription (Fig. 2).

The analysis suggests that the incorporation rates of pyrimidines are slower than purines, whereas the cleavage rate of misincorporated purines is faster. To account for the effect of second-downstream (+2) neighbors on the error rate at the current position (0), we performed the analysis for a more advanced (second-order) biophysical model (Fig. 3). Our results demonstrated that the error rate at each site is significantly impacted by the chemical identity of the immediate downstream and the second downstream bases. Our findings reveal that pyrimidines at downstream positions enhance transcriptional accuracy due to their slower incorporation rates and associated proofreading processes. In contrast, the identity of the third downstream base does not have any impact. These predictions are validated by performing bioinformatics analyses of the published experimental data. When

extended from a two-component to the four components (A, C, G, U), our models also accurately predict nucleotide-specific incorporation and proofreading kinetics, as validated by the experimental data (16, 23, 24).

Furthermore, with bioinformatic analysis, using BRCA1 coding sequences as an example, we demonstrated that sequence-dependent errors might significantly increase error probabilities for premature stop errors. These findings underscore the functional implications of transcription fidelity and highlight the necessity of integrating sequence-dependent effects into transcription models to accurately predict broader physiological consequences.

Our theoretical approach to investigate transcription elongation, calibrated against experimental data, accurately predicts nucleotide-specific incorporation rates. The analysis reveals that purines (A, G) polymerize faster than pyrimidines (C, U), a trend supported by experimental findings (25, 26). Structurally, purines have a double-ringed configuration, which facilitates stronger base-stacking interactions, probably leading to faster rates of association compared to single-ringed pyrimidines where such interactions are weaker (25, 26). Experimental studies further suggest that polymerization rates correlate with the 3' terminal base composition, following the order U<C<G<A. Additionally, a sequence-dependent thermal ratchet model for transcription elongation supports this hierarchy, reinforcing that pyrimidine incorporates more slowly than purines (16, 24). Our advanced theoretical model with second-order neighbor effects and all four nucleotides predicts the same incorporation rate hierarchy (U<C<G<A) as in the experiment.

Our theoretical analysis of errors under biologically relevant conditions suggests that, in addition to nucleotide incorporation rates, the dinucleotide cleavage rate of misincorporated nucleotides must be sequence-dependent. This cleavage is the basis of proofreading and mostly occurs after incorporation. By fitting our theoretical model with either two or four components to the corresponding experimental data, we found that dinucleotide cleavage rate for misincorporations replacing correct pyrimidines is approximately twice as fast as the rate of cleavage for those replacing correct purines. This further agrees with experimental studies showing that errors occurring in place of pyrimidine (U) are removed at twice the rate of those replacing purine (G) (12). Furthermore, the dinucleotide cleavage mechanism introduces two proofreading checkpoints that enhance transcriptional accuracy (7). This suggests that transcriptional fidelity is influenced not only by the identity of the nucleotide at the immediate downstream position but also by the identity of the second downstream nucleotide (see Fig. 3). Our more advanced theoretical model with second-order neighbor effects reveals that error rates decrease when pyrimidines occupy the second downstream position and reach their lowest when two consecutive pyrimidines are present. Conversely, the highest error rates occur when two purines appear in succession. This is consistent with the fact that pyrimidines exhibit slower incorporation rates and higher proofreading efficiency compared to purines (12, 24). Thus, transcriptional accuracy is

highest when two pyrimidines are positioned consecutively, whereas error rates peak when two purines appear together. Experimental studies indicate that the incorporation rates of nucleotides are affected when the downstream neighbor to be incorporated is a pyrimidine (34). However, it is unknown to what extent the incorporation dynamics will be affected when pyrimidines are at the upstream position. The experimental data emphasize the role of both upstream and downstream neighbors in determining transcriptional error (23), and hence the upstream neighbor effects on the incorporation dynamics must be determined. Our results indicate that incorporation rates of nucleotides are reduced by a factor ψ when the upstream (-1) nucleotide is a pyrimidine. In particular, at the optimal value of $\psi \approx 0.88$, our advanced model accurately reproduces experimental observations. This suggests that pyrimidines at upstream positions slightly slow down the incorporation process, thus potentially enhancing proofreading capability and transcription fidelity.

Transcriptional errors, influenced by genetic context, can affect the error probability (35–38). We examined the physiological implications of sequence-dependent error rates by assessing their impact on error probabilities in the BRCA1 gene, a key tumor suppressor (32, 33). Premature stop codon errors can impair protein function (32), while synonymous errors may reduce translation efficiency (33). Our analysis revealed that sequence-dependent error rates result in higher error probabilities than non-sequence-specific errors. This suggests that context sequence plays a significant role in determining transcription errors. The effect is mild for synonymous errors but about 20% for premature termination errors. The higher error probability for premature termination can be attributed to the composition of stop codons UAA, UAG, and UGA, which all contain a pyrimidine at position 0 and purines at the +1 and +2 downstream positions. Since purines at these downstream positions are associated with the highest error rates, there is a higher chance of misincorporations changing other bases into U at position 0 and resulting in a stop codon, i.e. premature termination errors. These findings highlight the relationship between transcription fidelity and functional genomic outcomes, stressing the critical role of genetic context in biological information processing. In conclusion, this study bridges theoretical insights with experimental data, providing a foundation for predictive models of transcription fidelity. Our results collectively emphasize the importance of integrating both short and long-range correlations and sequence-specific kinetic parameters in models of transcription fidelity.

Methods

We consider a theoretical framework for a template-dependent transcription elongation process, in which the monomers are added to the growing end of the mRNA molecule according to the DNA template sequence. We denote the sequence of length L on the DNA template as $X_1 X_2 \cdots X_L$. The elements of the mRNA copy sequence are denoted by y_i for $i = \{1, 2 \cdots, L\}$. During elongation, each DNA base X_i pairs with an mRNA base y_i forming $X_i : y_i$

pairs. The accuracy of these pairings is characterized by a binary sequence α_i where $\alpha_i \in \{R, W\}$ indicates whether the pairing at position i is correct (R) or incorrect (W).

We first consider a biophysical model incorporating first-order neighbor effects, with both the DNA and mRNA sequences composed of two nucleotide classes: purines (Pu) and pyrimidines (Py),i.e., $X_i, y_i \in \{Pu, Py\}$. In this model, the kinetic rates for nucleotide addition (k_i), removal ($\bar{k}_i$), and dinucleotide cleavage (r_i) at position i, depend on the accuracy of both current and the previously incorporated nucleotide, $\alpha_i \alpha_{i-1}$, i.e., whether α_i, α_{i-1} are R or W (see Fig. S1A). Additionally, we assume that the rates k_i, and r_i also depend on the chemical identity of the incorporating nucleotide y_i, i.e., whether it is Pu or Py (see Fig. 1C, Fig. S1B). It should be noted that if $\alpha_i = R$, and $X_i = Pu$, then the corresponding incorporated nucleotide $y_i = Py$, if $\alpha_i = R$, and $X_i = Py$, then $y_i = Pu$, $\alpha_i = W$, and $X_i = Pu$, then $y_i = Pu$, and if $\alpha_i = W$, and $X_i = Py$, then $y_i = Py$. In our model, we consider the incorporation rates of pyrimidines and purines to be related by $k_{ij}^{Py} = \gamma k_{ij}^{Pu}$, where, the superscript reflects the nature of incorporating monomer y_j, i.e., Pu or Py.

In what follows, we adapt the approach developed in Ref. (29). One can describe the transcription elongation as a first-order Markov chain process from a reflecting boundary at the first position to the absorbing boundary at the last. We assume that the mRNA transcription proceeds with the preexisting initial seed R or W, i.e., either correctly paired with the 3' end of the DNA template or not. The chain grows as the monomer at position i is added and removed with the rates k_i and $\bar{k}_i$, respectively. However, as boundary conditions, the chain's initial seed and the last added monomer cannot be removed. Additionally, to account for proofreading, we consider dinucleotide cleavage with the rate r_i from all sites except $i = 1, 2$, and L. Note that in our model, the kinetic rates are the effective rates obtained by combining several reaction substeps (3, 7). We now write the temporal evolutions of probability, $p_{\alpha_1 \alpha_2 \cdots \alpha_i}(t)$, of the growing chain sequence $\alpha_1 \alpha_2 \cdots \alpha_i$ ($1 \le i \le L$) for the given DNA template $X_1 X_2, \cdots X_L$, at time t, for the bulk positions i ($3 \le i \le L-3$),

$$\frac{d}{dt} p_{\alpha_1 \alpha_2 \cdots \alpha_i} = \sum_{\alpha_{i+1}} \bar{k}_{\alpha_i \alpha_{i+1}} p_{\alpha_1 \cdots \alpha_{i+1}} + k_{\alpha_{i-1} \alpha_i}^{y_i} p_{\alpha_1 \cdots \alpha_{i-1}}$$
$$+ \sum_{\alpha_{i+1}} \sum_{\alpha_{i+2}} r_{\alpha_{i+1} \alpha_{i+2}}^{y_{i+2}} p_{\alpha_1 \cdots \alpha_{i+2}}$$
$$- (k_{\alpha_i R}^{y_{i+1}} + k_{\alpha_i W}^{y_{i+1}} + \bar{k}_{\alpha_{i-1} \alpha_i} + r_{\alpha_{i-1} \alpha_i}^{y_i}) p_{\alpha_1 \alpha_2 \cdots \alpha_i}, \quad (1)$$

Here, the summations runs for α_{i+1}, $\alpha_{i+2} \in R, W$. The temporal evolution equations for the boundary positions are provided in the SI Appendix. We are interested in finding the final sequence distribution of the nascent chain, i.e., the long-time limit $P_{\alpha_1 \alpha_2 \cdots \alpha_L} = p_{\alpha_1 \alpha_2 \cdots \alpha_L}(t \to \infty)$. For this, the following boundary conditions are assumed. The initial boundary condition is $p_{\alpha_1}(t = 0) = q_{\alpha_1}^{X_1}$, such that $q_R^{X_1} + q_W^{X_1} = 1$, $p_{\alpha_1 \cdots \alpha_i}(t = 0) = 0$ (for $i \ge 2$). We also assume the transcription will be completed fully to absorbing state L in the long time limit, i.e., $p_{\alpha_1 \cdots \alpha_i}(t \to \infty) = 0$ for $1 \le i < L$. One can then integrate Eq. 1, along with

the temporal evolution equations for the probabilities for the boundary positions (see the SI Appendix) for t varying from limit 0 to ∞, and this leads to the iteration relations as given in SI Appendix. They can be transformed into an intuitive form of a forward inhomogeneous Markov chain, as below

$$P_{\alpha_1 \cdots \alpha_L} = q_{\alpha_1} M_{\alpha_1 \alpha_2} M_{\alpha_2 \alpha_3} \cdots M_{\alpha_i \alpha_{i+1}} \cdots M_{\alpha_{L-1} \alpha_L},$$
$$M_{\alpha_i \alpha_{i+1}} = \Pi_{\alpha_i \alpha_{i+1}} g_{\alpha_{i+1}} / g_{\alpha_i}, 1 \le i < L-1 \text{ and}$$
$$M_{\alpha_{L-1} \alpha_L} = k_{\alpha_{L-1} \alpha_L}^{y_L} / g_{\alpha_{L-1}}, \quad (2)$$

The expressions for $g_{\alpha_{i+1}}$, $\Pi_{\alpha_i \alpha_{i+1}}$ are provided in SI appendix. Furthermore, following the approach of Ref. (29), we can introduce the 2×2 stochastic transfer matrix $\mathbf{M}_{i,i+1}$ with each row sum equal to 1, i.e., $\sum_{\alpha_{i+1} = R, W} M_{\alpha_i \alpha_{i+1}} = 1$, for $\alpha_i = R, W$. In this context, one can calculate the positional probability for R or W at any position, i, on mRNA sequence given the monomer X_i at position i on the DNA template by $(P_R^i, P_W^i) = (q_R^{X_1}, q_W^{X_1}) M_{\alpha_1 \alpha_2} M_{\alpha_2 \alpha_3} M_{\alpha_{i-1} \alpha_i}$. For the first-order process, with one right and wrong nucleotide for a given monomer at the DNA template, one can write $\mathbf{M}_{i-1,i}$, as

$$\mathbf{M}_{i-1,i} = \begin{pmatrix} M_{R_{i-1} R_i} & M_{R_{i-1} W_i} \\ M_{W_{i-1} R_i} & M_{W_{i-1} W_i} \end{pmatrix} \quad (3)$$

Equivalently, the probability distribution at site i is given as, $(P_R^i, P_W^i) = (P_R^{i-1}, P_W^{i-1}) M_{i-1,i}$ (30). In particular, one can obtain the error probability,

$$P_W^i = \eta^i = P_R^{i-1} M_{R_{i-1} W_i} + P_W^{i-1} M_{W_{i-1} W_i}. \quad (4)$$

We now obtain the approximate analytical expression for the error rate under the biologically relevant conditions inspired by the kinetic parameters of real RNAPs (3, 8, 12). The analytical expressions of error are utilized to fit the theoretical results to the experimental data.

For the biophysical model with first-order neighbor effects and two components (Fig. S1), the biologically relevant conditions are intuitive as follows:

1. $k_{RR}^y \gg k_{RW}^y$, implying the addition of R nucleotide is always much faster than that of W, irrespective of the identity y of the incorporating monomer. monomers.

2. $k_{WR}^y / k_{WW}^y \gg k_{RR}^y / k_{WW}^y$, implying that the successive additions of W are almost inhibited, taken to approximately zero.

3. $k_{RR}^y > \bar{k}_{RR}, \bar{k}_{WR}, r_{RR}$. This means that successive additions of R always dominate the backward and cleavage rate, irrespective of the identity y of the incorporating monomer.

4. $r_{WW}^y, r_{WR}^y, r_{RW}^y \gg r_{RR}$, which means the terminus containing Ws are likely to be proofread faster.

Under the biologically relevant conditions, we first approximate g_{α_i}'s. The condition $k_{\alpha_{L-1} R}^{y_L} \gg k_{\alpha_{L-1} W}^{y_L}$ implies $g_{\alpha_{L-1}} \simeq k_{\alpha_{L-1} R}^{y_L}$. Following the same logic, we obtain $g_{\alpha_i} \simeq k_{\alpha_i R}^{y_{i+1}}$, $\Pi_{\alpha_i \alpha_{i+1}} \simeq k_{\alpha_i \alpha_{i+1}}^{y_{i+1}} / (\bar{k}_{\alpha_i \alpha_{i+1}} + r_{\alpha_i \alpha_{i+1}}^{y_{i+1}} + k_{\alpha_{i+1} R}^{y_{i+2}})$.

Similarly, one can write the approximate expression for the individual elements of the matrix M_i (see the SI Appendix). Further, under the biologically relevant conditions, $M_{RR} \gg M_{RW}, M_{WR} \gg M_{WW}$, and $M_{RW} \gg M_{WW}$ (19, 29, 30). The analytical expression of the probability for a wrong nucleotide at any i^{th} position of the chain is thus given as $P_W^i \simeq M_{R_{i-1}W_i}$ (see Appendix C, Ref. (29)). For the first-order biophysical model with two components, we denote the approximate expression of error rate at any current site, 0 (say), depending on the nucleotide at the downstream position, +1, by η_2^F, where the superscript represents the first-order model and subscript represent the two components. That is,

$$\eta_2^F \simeq \frac{k_{RW}^{y_0}}{k_{RR}^{y_0}\left(1 + \frac{\bar{k}_{RW} + r_{RW}^{y_0}}{k_{WR}^{y_{+1}}}\right).} \tag{5}$$

Here, the denominator of the RHS reflects the dinucleotide cleavage and the first-order neighbor effects. Depending on the chemical identity of the nucleotides $y_0, y_1 \in \{$ Pu, Py $\}$, we obtain the theoretical error rate for four dimers (y_0-y_{+1}) (see Fig. 2D, and SI Appendix).

For the more advanced model with second-order neighbor effects and two components (Pu, Py), the rates k_i, $\bar{k}_i$, and r_i depend on the accuracy of 3 nucleotides, i.e. on α_{i-2}, α_{i-1}, α_i (see Fig. S4A). For the template-dependent kinetics, we consider that the rate for addition, k_i, also depends on the identity of the incorporating nucleotide $y_i =$ Pu, Py. Thus, the incorporation rate $k_{ijk}^{xy} = k_{ijk}$, if $y =$ Pu, and $k_{ijk}^{xy} = \gamma k_{ijk}$, if $y =$ Py. Here, $i, j, k = R, W$, $x =$ Pu or Py, is the identity of the previous nucleotide j, and y denotes the nature, i.e., Pu or Py, of the current incorporating nucleotide k. Additionally, we also consider the case when the identity of the preceding (upstream) nucleotide $y_{i-1} =$ Pu or Py, can also affect the incorporation rate k_i (SI Appendix, Fig. S4B). For this case, when upstream nucleotide identity effects are considered, the rate $k_{ijk}^{xy} = k_{ijk}$ if $x, y =$ Pu, $k_{ijk}^{xy} = \gamma k_{ijk}$ if $y =$ Py and $x =$ Pu, $k_{ijk}^{xy} = \psi k_{ijk}$ if $y =$ Pu and $x =$ Py, $k_{ijk}^{xy} = \gamma \psi k_{ijk}$ if $y =$ Py and $x =$ Py. We start the elongation process with the initial seed as dimer $\alpha_1 \alpha_2$ with probability $q_{\alpha_1\alpha_2}^{X_1 X_2}$, such that $\sum_{\alpha_1, \alpha_2 = R, W} q_{\alpha_1\alpha_2}^{X_1 X_2} = 1$. Following the same logic as for the first-order biophysical model, we can obtain the final sequence probability $P_{\alpha_1\alpha_2\cdots\alpha_L}$ for the second-order model as a forward inhomogeneous Markov chain (see SI Appendix for details). Under the biologically relevant condition (see the SI Appendix), the approximate expression of the error rate for the second-order biophysical model with two components, η_2^S, where the superscript represents the second-order model and subscript represents the two components, can be written as,

$$\eta_2^S \simeq \frac{k_{RRW}^{y_{-1}y_0}}{k_{RRR}^{y_{-1}y_0}\left[1 + \frac{\bar{k}_{RRW} + r_{RW}^{y_0}}{k_{RWR}^{y_0 y_{+1}}}\left(1 + \frac{\bar{k}_{RWR} + r_{WR}^{y_{+1}}}{k_{WRR}^{y_{+1}y_{+2}}}\right)\right]} \tag{6}$$

Here, the denominator in the RHS reflects the dinucleotide cleavage and the second-order neighbor effects on the error

rate. For the case with no upstream nucleotide identity effects, we calculate the error rate for 8 trimers ($y_0 - y_{+1} - y_{+2}$) depending on the chemical identity of the nucleotides $y_i \in \{$ Pu, Py $\}$ for $i = \{0, +1, +2\}$ (see Fig. 4A). For the case with upstream nucleotide identity effects, we calculate the for 16 tetramers ($y_{-1} - y_0 - y_{+1} - y_{+2}$) depending on the chemical identity of the nucleotides $y_i \in \{$ Pu, Py $\}$ for $i = \{-1, 0, +1, +2\}$ (see Fig. 4B).

To consider a more realistic case when both DNA and mRNA sequences consist of four components (A,C, G, T/U), for every correct pairing R (A:U, C:G, G:C, T:A) we need to account for the three wrong pairing possibilities W_j, for $j = 1$ to 3. The kinetic parameters for the addition and the removal of every correct nucleotide and their corresponding three wrong nucleotides for the second-order biophysical model are given in Table S3. Similarly to the second-order model with two components, one can obtain the error for the model with four elements. Here, we write the approximate analytical expression of the error rate for the second-order model with four components, η_4^S, where the superscript represents the second-order model and the subscript denotes the four components, under bio-relevant conditions (see the SI Appendix) as below.

$$\eta_4^S \simeq \sum_{j=1}^{3} \frac{k_{RRW_j}^{y_0}}{k_{RRR}^{y_0}\left[1 + \frac{\bar{k}_{RRW_j} + r_{RW}^{y_0}}{k_{RW_jR}^{y_{+1}}}\left(1 + \frac{\bar{k}_{RWR} + r_{WR}^{y_{+1}}}{k_{WRR}^{y_{+2}}}\right)\right]} \tag{7}$$

Using the above equation, one can obtain the error rate for the 64 trimers ($y_0 - y_{+1} - y_{+2}$) depending on the chemical identity of the nucleotides $y_i \in \{$ A, U, C, G $\}$ for $i = \{0, +1, +2\}$ (see Fig. 4C).

ACKNOWLEDGEMENTS

We sincerely thank the authors of Ref. (23) for generously providing the datasets from their study. This work was supported by the Center for Theoretical Biological Physics National Science Foundation (NSF) Grant PHY-2019745. OAI also acknowledges support from the Welch Foundation Grant C-1995. ABK also acknowledges the support from the Welch Foundation (C-1559).

Bibliography

1. Hojoong Kwak and John T Lis. Control of transcriptional elongation. *Annual review of genetics*, 47(1):483–508, 2013.

2. James D Watson and Francis HC Crick. Molecular structure of nucleic acids: a structure for deoxyribose nucleic acid. *Nature*, 171(4356):737–738, 1953.

3. Yulia Yuzenkova, Aleksandra Bochkareva, Vasisht R Tadigotla, Mohammad Roghanian, Savva Zorov, Konstantin Severinov, and Nikolay Zenkin. Stepwise mechanism for transcription fidelity. *BMC biology*, 8(1):1–15, 2010.

4. John J Hopfield. Kinetic proofreading: a new mechanism for reducing errors in biosynthetic processes requiring high specificity. *Proc. Natl. Acad. Sci. U. S. A*, 71(10):4135–4139, 1974.

5. Jacques Ninio. Kinetic amplification of enzyme discrimination. *Biochimie*, 57(5):587–595, 1975.

6. Pablo Sartori and Simone Pigolotti. Kinetic versus energetic discrimination in biological copying. *Physical review letters*, 110(18):188101, 2013.

7. Tripti Midha, Joel D Mallory, Anatoly B Kolomeisky, and Oleg A Igoshin. Synergy among pausing, intrinsic proofreading, and accessory proteins results in optimal transcription speed and tolerable accuracy. *The Journal of Physical Chemistry Letters*, 14(14):3422–3429, 2023.

8. Maria L Kireeva, Yuri A Nedialkov, Gina H Cremona, Yuri A Purtov, Lucyna Lubkowska, Francisco Malagon, Zachary F Burton, Jeffrey N Strathern, and Mikhail Kashlev. Transient reversal of rna polymerase ii active site closing controls fidelity of transcription elongation. *Molecular cell*, 30(5):557–566, 2008.

9. Marianna Orlova, Janet Newlands, Asis Das, Alex Goldfarb, and Sergei Borukhov. Intrinsic transcript cleavage activity of rna polymerase. *Proc. Natl. Acad. Sci. U. S. A*, 92(10):4596–4600, 1995.

10. Maria L Kireeva and Mikhail Kashlev. Mechanism of sequence-specific pausing of bacterial rna polymerase. *Proc. Natl. Acad. Sci. U. S. A*, 106(22):8900–8905, 2009.

11. Jasmin F Sydow, Florian Brueckner, Alan CM Cheung, Gerke E Damsma, Stefan Dengl, Elisabeth Lehmann, Dmitry Vassylyev, and Patrick Cramer. Structural basis of transcription: mismatch-specific fidelity mechanisms and paused rna polymerase ii with frayed rna. *Mol. Cell*, 34(6):710–721, 2009.

12. Nikolay Zenkin, Yulia Yuzenkova, and Konstantin Severinov. Transcript-assisted transcriptional proofreading. *Science*, 313(5786):518–520, 2006.

13. Katherine James, Pamela Gamba, Simon J. Cockell, and Nikolay Zenkin. Misincorporation by RNA polymerase is a major source of transcription pausing in vivo. *Nucleic Acids Res.*, 45(3):1105–1113, 10 2016.

14. Pamela Gamba and Nikolay Zenkin. Transcription fidelity and its roles in the cell. *Current Opin. Microbiol.*, 42:13–18, 2018.

15. Matthew J Thomas, Angelina A Platas, and Diane K Hawley. Transcriptional fidelity and proofreading by rna polymerase ii. *Cell*, 93(4):627–637, 1998.

16. Lu Bai, Robert M Fulbright, and Michelle D Wang. Mechanochemical kinetics of transcription elongation. *Physical review letters*, 98(6):068103, 2007.

17. Lu Zhang, Fátima Pardo-Avila, Ilona Christy Unarta, Peter Pak-Hang Cheung, Guo Wang, Dong Wang, and Xuhui Huang. Elucidation of the dynamics of transcription elongation by rna polymerase ii using kinetic network models. *Accounts of chemical research*, 49(4):687–694, 2016.

18. Wenbo Fu, Qiushi Li, Yongshun Song, Yaogen Shu, Zhongcan Ouyang, and Ming Li. Theoretical analysis of rna polymerase fidelity: a steady-state copolymerization approach. *Communications in Theoretical Physics*, 74(1):015601, 2021.

19. Tripti Midha, Anatoly B Kolomeisky, and Oleg A Igoshin. Linear-decoupling enables accurate speed and accuracy predictions for copolymerization processes. *The Journal of Physical Chemistry Letters*, 15(37):9361–9368, 2024.

20. Pierre Gaspard and David Andrieux. Kinetics and thermodynamics of first-order markov chain copolymerization. *The Journal of chemical physics*, 141(4):044908, 2014.

21. Pierre Gaspard. Growth and dissolution of macromolecular markov chains. *Journal of Statistical Physics*, 164:17–48, 2016.

22. Yao-Gen Shu, Yong-Shun Song, Zhong-Can Ou-Yang, and Ming Li. A general theory of kinetics and thermodynamics of steady-state copolymerization. *Journal of Physics: Condensed Matter*, 27(23):235105, 2015.

23. Claire Chung, Bert M Verheijen, Xinmin Zhang, Biao Huang, Aeowynn Coakley, Eric McGann, Emily Wade, Olivia Dinep-Schneider, Jessica LaGosh, Maria-Eleni Anagnostou, et al. The fidelity of transcription in human cells. *Proceedings of the National Academy of Sciences*, 120(5):e2210038120, 2023.

24. M Angela Parsons, Richard R Sinden, and Michael G Izban. Transcriptional properties of rna polymerase ii within triplet repeat-containing dna from the human myotonic dystrophy and fragile x loci. *Journal of Biological Chemistry*, 273(41):26998–27008, 1998.

25. Shuntaro Takahashi, Hiromichi Okura, Pallavi Chilka, Saptarshi Ghosh, and Naoki Sugimoto. Molecular crowding induces primer extension by rna polymerase through base stacking beyond watson–crick rules. *RSC advances*, 10(55):33052–33058, 2020.

26. Matthew J Burak, Kip E Guja, and Miguel Garcia-Diaz. Nucleotide binding interactions modulate dntp selectivity and facilitate 8-oxo-dgtp incorporation by dna polymerase lambda. *Nucleic acids research*, 43(16):8089–8099, 2015.

27. Pierre Gaspard. Template-directed copolymerization, random walks along disordered tracks, and fractals. *Physical review letters*, 117(23):238101, 2016.

28. Pierre Gaspard. Iterated function systems for dna replication. *Physical Review E*, 96(4):042403, 2017.

29. Qiu-Shi Li, Pei-Dong Zheng, Yao-Gen Shu, Zhong-Can Ou-Yang, and Ming Li. Template-specific fidelity of dna replication with high-order neighbor effects: A first-passage approach. *Physical Review E*, 100(1):012131, 2019.

30. Qiu-Shi Li, Yao-Gen Shu, Zhong-Can Ou-Yang, and Ming Li. Kinetic assays of dna polymerase fidelity: A theoretical perspective beyond michaelis-menten kinetics. *Physical Review E*, 104(1):014408, 2021.

31. Daniel T Gillespie. Exact stochastic simulation of coupled chemical reactions. *The journal of physical chemistry*, 81(25):2340–2361, 1977.

32. Laure Perrin-Vidoz, Olga M Sinilnikova, Dominique Stoppa-Lyonnet, Gilbert M Lenoir, and Sylvie Mazoyer. The nonsense-mediated mrna decay pathway triggers degradation of most brca1 mrnas bearing premature termination codons. *Human molecular genetics*, 11(23):2805–2814, 2002.

33. Fran Supek, Belén Miñana, Juan Valcárcel, Toni Gabaldón, and Ben Lehner. Synonymous mutations frequently act as driver mutations in human cancers. *Cell*, 156(6):1324–1335, 2014.

34. Shannon F Holmes and Dorothy A Erie. Downstream dna sequence effects on transcription elongation: allosteric binding of nucleoside triphosphates facilitates translocation via a ratchet motion. *Journal of Biological Chemistry*, 278(37):35597–35608, 2003.

35. Madeleine Oman, Aqsa Alam, and Rob W Ness. How sequence context-dependent mutability drives mutation rate variation in the genome. *Genome Biology and Evolution*, 14(3):evac032, 2022.

36. Lai Wei. Selection on synonymous mutations revealed by 1135 genomes of arabidopsis thaliana. *Evolutionary Bioinformatics*, 16:1176934320916794, 2020.

37. Wenli Jia and Paul G Higgs. Codon usage in mitochondrial genomes: distinguishing context-dependent mutation from translational selection. *Molecular Biology and Evolution*, 25(2):339–351, 2008.

38. Claire S Chung, Yi Kou, Sarah J Shemtov, Bert M Verheijen, Ilse Flores, Kayla Love, Ashley Del Dosso, Max A Thorwald, Yuchen Liu, Daniel Hicks, et al. Transcript errors generate amyloid-like proteins in human cells. *Nature Communications*, 15(1):8676, 2024.

Supporting Information Text

Biophysical model with first-order neighbor effects. Here, we write the temporal evolutions of probability, $p_{\alpha_1 \alpha_2 \cdots \alpha_i}(t)$, of the growing chain sequence $\alpha_1 \, \alpha_2 \, \cdots \, \alpha_i$ for $i = 1, 2, L-2, L-1, L$ as below

$$
\begin{aligned}
\frac{dp_{\alpha_1}}{dt} &= \sum_{\alpha_2 = R,W} \bar{k}_{\alpha_1 \alpha_2} p_{\alpha_1 \alpha_2} + \sum_{\alpha_2 = R,W} \sum_{\alpha_3 = R,W} r^{y_3}_{\alpha_2 \alpha_3} p_{\alpha_1 \alpha_2 \alpha_3} - (k^{y_2}_{\alpha_1 R} + k^{y_2}_{\alpha_1 W}) p_{\alpha_1} \\
\frac{dp_{\alpha_1 \alpha_2}}{dt} &= \sum_{\alpha_3 = R,W} \bar{k}_{\alpha_2 \alpha_3} p_{\alpha_1 \alpha_2 \alpha_3} + k^{y_2}_{\alpha_1 \alpha_2} p_{\alpha_1} + \sum_{\alpha_3 = R,W} \sum_{\alpha_4 = R,W} r^{y_4}_{\alpha_3 \alpha_4} p_{\alpha_1 \cdots \alpha_4} - (k^{y_3}_{\alpha_2 R} + k^{y_3}_{\alpha_2 W} + \bar{k}_{\alpha_1 \alpha_2}) p_{\alpha_1 \alpha_2},
\end{aligned}
$$

$$
\frac{dp_{\alpha_1 \alpha_2 \cdots \alpha_{L-2}}}{dt} = \sum_{\alpha_{L-1}} \bar{k}_{\alpha_{L-2} \alpha_{L-1}} p_{\alpha_1 \cdots \alpha_{L-1}} + k^{y_{L-2}}_{\alpha_{L-3} \alpha_{L-2}} p_{\alpha_1 \cdots \alpha_{L-3}} - (k^{y_{L-1}}_{\alpha_{L-2} R} + k^{y_{L-1}}_{\alpha_{L-2} W} + \bar{k}_{\alpha_{L-3} \alpha_{L-2}} + r^{y_{L-2}}_{\alpha_{L-3} \alpha_{L-2}}) p_{\alpha_1 \cdots \alpha_{L-2}},
$$

$$
\begin{aligned}
\frac{dp_{\alpha_1 \alpha_2 \cdots \alpha_{L-1}}}{dt} &= k^{y_{L-1}}_{\alpha_{L-2} \alpha_{L-1}} p_{\alpha_1 \alpha_2 \cdots \alpha_{L-2}} - (k^{y_L}_{\alpha_{L-1} R} + k^{y_L}_{\alpha_{L-1} W} + \bar{k}_{\alpha_{L-2} \alpha_{L-1}} + r^{y_{L-1}}_{\alpha_{L-2} \alpha_{L-1}}) p_{\alpha_1 \alpha_2 \cdots \alpha_{L-1}}, \\
\frac{dp_{\alpha_1 \alpha_2 \cdots \alpha_L}}{dt} &= k^{y_L}_{\alpha_{L-1} \alpha_L} p_{\alpha_1 \alpha_2 \cdots \alpha_{L-1}}.
\end{aligned}
\tag{S1}
$$

Here, we obtain the final sequence distribution, i.e., $P_{\alpha_1 \alpha_2 \cdots \alpha_L} = p_{\alpha_1 \alpha_2 \cdots \alpha_L}(t \to \infty)$ of the mRNA chain. For this purpose, we integrate the temporal evolution equations for $p_{\alpha_1 \alpha_2 \cdots \alpha_i}(t)$, $\forall i = 1$ to L, given by Eq. (S1), and Eq. 2 (main manuscript), under the boundary conditions as defined in the methods (main manuscript).

$$
\begin{aligned}
-q_{\alpha_1} &= \sum_{\alpha_2 = R,W} \bar{k}_{\alpha_1 \alpha_2} \Gamma_{\alpha_1 \alpha_2} + \sum_{\alpha_2} \sum_{\alpha_3} r^{y_3}_{\alpha_2 \alpha_3} \Gamma_{\alpha_1 \alpha_2 \alpha_3} - (k^{y_2}_{\alpha_1 R} + k^{y_2}_{\alpha_1 W}) \Gamma_{\alpha_1} \\
0 &= \sum_{\alpha_3 = R,W} \bar{k}_{\alpha_2 \alpha_3} \Gamma_{\alpha_1 \alpha_2 \alpha_3} + \sum_{\alpha_3} \sum_{\alpha_4} r^{y_4}_{\alpha_3 \alpha_4} \Gamma_{\alpha_1 \cdots \alpha_4} - (k^{y_3}_{\alpha_2 R} + k^{y_3}_{\alpha_2 W} + \bar{k}_{\alpha_1 \alpha_2}) \Gamma_{\alpha_1 \alpha_2}, \\
0 &= \sum_{\alpha_{i+1}} \bar{k}_{\alpha_i \alpha_{i+1}} \Gamma_{\alpha_1 \cdots \alpha_{i+1}} + k^{y_i}_{\alpha_{i-1} \alpha_i} \Gamma_{\alpha_1 \cdots \alpha_{i-1}} + \sum_{\alpha_{i+1}} \sum_{\alpha_{i+2}} r^{y_{i+2}}_{\alpha_{i+1} \alpha_{i+2}} \Gamma_{\alpha_1 \cdots \alpha_{i+2}} \\
&\quad - (k^{y_{i+1}}_{\alpha_i R} + k^{y_{i+1}}_{\alpha_i W} + \bar{k}_{\alpha_{i-1} \alpha_i} + r^{y_i}_{\alpha_{i-1} \alpha_i}) \Gamma_{\alpha_1 \alpha_2 \cdots \alpha_i}, \quad 3 \le i < L-2, \\
0 &= \sum_{\alpha_{L-1} = R,W} \bar{k}_{\alpha_{L-2} \alpha_{L-1}} \Gamma_{\alpha_1 \cdots \alpha_{L-1}} + k^{y_{L-2}}_{\alpha_{L-3} \alpha_{L-2}} \Gamma_{\alpha_1 \cdots \alpha_{L-3}} - \left(k^{y_{L-1}}_{\alpha_{L-2} R} + k^{y_{L-1}}_{\alpha_{L-2} W} + \bar{k}_{\alpha_{L-3} \alpha_{L-2}} + r^{y_{L-2}}_{\alpha_{L-3} \alpha_{L-2}} \right) \Gamma_{\alpha_1 \cdots \alpha_{L-2}} \\
0 &= k^{y_{L-1}}_{\alpha_{L-2} \alpha_{L-1}} \Gamma_{\alpha_1 \alpha_2 \cdots \alpha_{L-2}} - (k^{y_L}_{\alpha_{L-1} R} + k^{y_L}_{\alpha_{L-1} W} + \bar{k}_{\alpha_{L-2} \alpha_{L-1}} + r^{y_{L-1}}_{\alpha_{L-2} \alpha_{L-1}}) \Gamma_{\alpha_1 \cdots \alpha_{L-1}}, \\
P_{\alpha_1 \alpha_2 \cdots \alpha_L} &= k^{y_L}_{\alpha_{L-1} \alpha_L} \Gamma_{\alpha_1 \alpha_2 \cdots \alpha_{L-1}}.
\end{aligned}
\tag{S2}
$$

Here, $\Gamma_{\alpha_1 \cdots \alpha_i} = \int_0^\infty p_{\alpha_1 \cdots \alpha_i}(t) dt$. One can solve the above equations to obtain the following iteration relations:

$$
\begin{aligned}
P_{\alpha_1 \cdots \alpha_L}(t \to \infty) &= (q_{\alpha_1}/g_{\alpha_1}) \Pi_{\alpha_1 \alpha_2} \Pi_{\alpha_2 \alpha_3} \cdots \Pi_{\alpha_i \alpha_{i+1}} \cdots \Pi_{\alpha_{L-2} \alpha_{L-1}} k^{y_L}_{\alpha_{L-1} \alpha_L} \\
\Pi_{\alpha_i \alpha_{i+1}} &= k^{y_{i+1}}_{\alpha_i \alpha_{i+1}} / (\bar{k}_{\alpha_i \alpha_{i+1}} + r^{y_{i+1}}_{\alpha_i \alpha_{i+1}} + g_{\alpha_{i+1}}), 1 \le i \le L-2 \\
g_{\alpha_{i+1}} &= \sum_{\alpha_{i+2}} \left[\Pi_{\alpha_{i+1} \alpha_{i+2}} (r^{y_{i+2}}_{\alpha_{i+1} \alpha_{i+2}} + g_{\alpha_{i+2}}) - \sum_{\alpha_{i+3}} \Pi_{\alpha_{i+1} \alpha_{i+2}} \left(\Pi_{\alpha_{i+2} \alpha_{i+3}} r^{y_{i+3}}_{\alpha_{i+2} \alpha_{i+3}} \right) \right], 1 \le i \le L-4 \\
g_{\alpha_{L-2}} &= \sum_{\alpha_{L-1}} \left[\Pi_{\alpha_{L-2} \alpha_{L-1}} (r^{y_{L-1}}_{\alpha_{L-2} \alpha_{L-1}} + g_{\alpha_{L-1}}) \right], \\
g_{\alpha_{L-1}} &= \sum_{\alpha_L} k^{y_L}_{\alpha_{L-1} \alpha_L}
\end{aligned}
\tag{S3}
$$

The above relations are further utilized to compute the stochastic transfer matrix $\mathbf{M}_{i-1,i}$. For instance, under the bio-relevant conditions (see methods), we can obtain $M_{R_{i-1} W_i}$, which is nothing but an approximate probability of a wrong nucleotide P^i_W at position i, and is given as

$$
\eta^i \simeq M_{R_{i-1} W_i} \simeq \frac{k^{y_i}_{RW}}{k^{y_i}_{RR}} \Bigg/ \left(1 + \frac{\bar{k}_{RW} + r^{y_0}_{RW}}{k^{y_{i+1}}_{WR}} \right).
\tag{S4}
$$

Theoretical explanation for the identical error rate ratio for dimers in the first-order biophysical model with two elements. In this section, we mathematically justify the reason for the two identical error ratio curves in Fig. 2B, and Fig. 2C (main manuscript) for the first-order biophysical model with two elements. For this purpose, we first theoretically compute the error ratio: $\eta(\text{Pu@0})/\eta(\text{Py@0})$ (see Fig. 2B, main manuscript) and show that it is identical for either Pu at +1 position or Py at +1 position. We then compute the error ratio $\eta(\text{Pu@}+1)/\eta(\text{Py@}+1)$ (see Fig. 2C, main manuscript) and show that the ratio is identical for either Pu at 0 position or Py at 0 position. To show this, we write the analytical expression of error rate for dimers $y_0 - y_{+1}$, occupying positions $(0, +1)$, where $y_0, y_1 \in \{\text{ Pu, Py }\}$ for the first-order biophysical model (see methods, main manuscript),

$$\eta(y_0 - y_1) \quad \simeq \quad \frac{k_{RW}^{y_0}}{k_{RR}^{y_0}} \left/ \left(1 + \frac{\bar{k}_{RW} + r_{RW}^{y_0}}{k_{WR}^{y_{+1}}} \right) \right. . \tag{S5}$$

Explicitly, the error rate, $\eta(y_0 - y_1)$, for all possible dimers $y_0 - y_1$ at $(0,+1)$ positions can be written using Eq. S5 as below

$$\eta(\text{Pu-Pu}) \quad \simeq \quad \frac{\gamma k_{RW}}{k_{RR}} \left/ \left(1 + \frac{\bar{k}_{RW} + r_{RW}}{k_{WR}} \right) \right. . \tag{S6}$$

$$\eta(\text{Py-Pu}) \quad \simeq \quad \frac{k_{RW}}{\gamma k_{RR}} \left/ \left(1 + \frac{\bar{k}_{RW} + r_{RW}}{k_{WR}} \right) \right. . \tag{S7}$$

$$\eta(\text{Pu-Py}) \quad \simeq \quad \frac{\gamma k_{RW}}{k_{RR}} \left/ \left(1 + \frac{\bar{k}_{RW} + r_{RW}}{\gamma k_{WR}} \right) \right. . \tag{S8}$$

$$\eta(\text{Py-Py}) \quad \simeq \quad \frac{k_{RW}}{\gamma k_{RR}} \left/ \left(1 + \frac{\bar{k}_{RW} + r_{RW}}{\gamma k_{WR}} \right) \right. . \tag{S9}$$

Using the above equations, we obtain

$$\frac{\eta(\text{Pu-Pu})}{\eta(\text{Py-Pu})} = \frac{\eta(\text{Pu-Py})}{\eta(\text{Py-Py})} = \gamma^2, \tag{S10}$$

for non-nucleotide-specific dinucleotide cleavage rates. Whereas, in the case when the cleavage rate for purines and pyrimidines differ by a factor ϕ, we get

$$\frac{\eta(\text{Pu-Pu})}{\eta(\text{Py-Pu})} = \frac{\eta(\text{Pu-Py})}{\eta(\text{Py-Py})} = \frac{\gamma^2 (k_{WR} + \bar{k}_{RW} + \phi r_{\text{Pu}})}{(k_{WR} + \bar{k}_{RW} + r_{\text{Pu}})}. \tag{S11}$$

It is clear from Eqs. (S10)-(S11) that the ratio of the error rate with Pu @0 to Py @0 is the same whether there is a Pu or Py at the +1 position. This is the reason we get identical curves, as shown in Fig. 2B (main manuscript) for the case of non-nucleotide-specific proofreading. Similarly, we observe that

$$\frac{\eta(\text{Pu-Pu})}{\eta(\text{Pu-Py})} = \frac{\eta(\text{Py-Pu})}{\eta(\text{Py-Py})} = \frac{\gamma k_{WR} + \bar{k}_{RW} + r_{RW}}{\gamma(k_{WR} + \bar{k}_{RW} + r_{RW})}, \tag{S12}$$

for non-nucleotide-specific dinucleotide cleavage rates. This implies that the ratio of the error rate with Pu @+1 to Py @+1 is the same whether there is a Pu or Py at the 0 position. This is the reason for the identical two curves in Fig. 2C for the case of non-nucleotide-specific proofreading. Thus, it can be said that the effects of current-base and next-base kinetics are independent of each other.

Advanced model with second-order neighbor effects. For the theoretical model with second-order neighbor effects, we assume that the elongation process starts with the initial seed $\alpha_1 \alpha_2$, with probability $q_{\alpha_1 \alpha_2}$, such that $\sum_{\alpha_1 \alpha_2} q_{\alpha_1 \alpha_2} = 1$. The first-passage process description for the model assumes that the first dimer acts as a reflecting boundary, i.e., it can not be removed, and the last site is the absorbing boundary. Let $\alpha_1 \alpha_2 \cdots \alpha_i$ denote the growing $mRNA$, with probability $p_{\alpha_1 \alpha_2 \cdots \alpha_i}$ for a given DNA template $X_1 X_2 \cdots X_i$. We write the temporal evolution equation for these probabilities, as below,

$$\frac{dp_{\alpha_1 \alpha_2}}{dt} \quad = \quad \sum_{\alpha_3 = R, W} \bar{k}_{\alpha_1 \alpha_2 \alpha_3} p_{\alpha_1 \alpha_2 \alpha_3} + \sum_{\alpha_3} \sum_{\alpha_4} r_{\alpha_3 \alpha_4}^{y_4} p_{\alpha_1 \alpha_2 \cdots \alpha_4} - (k_{\alpha_1 \alpha_2 R}^{y_2 y_3} + k_{\alpha_1 \alpha_2 W}^{y_2 y_3}) p_{\alpha_1 \alpha_2},$$

$$\frac{dp_{\alpha_1 \alpha_2 \alpha_3}}{dt} \quad = \quad \sum_{\alpha_3 = R, W} \bar{k}_{\alpha_2 \alpha_3 \alpha_4} p_{\alpha_1 \alpha_2 \alpha_3 \alpha_4} + k_{\alpha_1 \alpha_2 \alpha_3}^{y_2 y_3} p_{\alpha_1 \alpha_2} + \sum_{\alpha_4} \sum_{\alpha_5} r_{\alpha_3 \alpha_4}^{y_4} p_{\alpha_1 \cdots \alpha_4} - (k_{\alpha_2 \alpha_3 R}^{y_3 y_4} + k_{\alpha_2 \alpha_3 W}^{y_3 y_4} + \bar{k}_{\alpha_1 \alpha_2 \alpha_3}) p_{\alpha_1 \alpha_2 \alpha_3},$$

$$\frac{dp_{\alpha_1\alpha_2\cdots\alpha_i}}{dt} = \sum_{\alpha_{i+1}} \bar{k}_{\alpha_{i-1}\alpha_i\alpha_{i+1}} p_{\alpha_1\alpha_2\cdots\alpha_{i+1}} + k^{y_{i-1}y_i}_{\alpha_{i-2}\alpha_{i-1}\alpha_i} p_{\alpha_1\alpha_2\cdots\alpha_{i-1}} + \sum_{\alpha_{i+1}}\sum_{\alpha_{i+2}} r^{y_{i+2}}_{\alpha_{i+1}\alpha_{i+2}} p_{\alpha_1\cdots\alpha_{i+2}}$$
$$- (k^{y_iy_{i+1}}_{\alpha_{i-1}\alpha_i R} + k^{y_iy_{i+1}}_{\alpha_{i-1}\alpha_i W} + \bar{k}_{\alpha_{i-2}\alpha_{i-1}\alpha_i} + r^{y_i}_{\alpha_{i-1}\alpha_i}) p_{\alpha_1\alpha_2\cdots\alpha_i},$$

$$\frac{dp_{\alpha_1\cdots\alpha_{L-2}}}{dt} = \sum_{\alpha_{L-1}} \bar{k}_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} p_{\alpha_1\cdots\alpha_{L-1}} + k^{y_{L-3}y_{L-2}}_{\alpha_{L-4}\alpha_{L-3}\alpha_{L-2}} p_{\alpha_1\cdots\alpha_{L-3}}$$
$$- (k^{y_{L-2}y_{L-1}}_{\alpha_{L-3}\alpha_{L-2} R} + k^{y_{L-2}y_{L-1}}_{\alpha_{L-3}\alpha_{L-2} W} + \bar{k}_{\alpha_{L-4}\alpha_{L-3}\alpha_{L-2}} + r^{y_{L-2}}_{\alpha_{L-3}\alpha_{L-2}}) p_{\alpha_1\cdots\alpha_{L-2}},$$

$$\frac{dp_{\alpha_1\alpha_2\cdots\alpha_{L-1}}}{dt} = k^{y_{L-2}y_{L-1}}_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} p_{\alpha_1\alpha_2\cdots\alpha_{L-2}} - (k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1} R} + k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1} W} + \bar{k}_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} + r^{y_{L-1}}_{\alpha_{L-2}\alpha_{L-1}}) p_{\alpha_1\cdots\alpha_{L-1}},$$

$$\frac{dp_{\alpha_1\alpha_2\cdots\alpha_L}}{dt} = k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1}\alpha_L} p_{\alpha_1\alpha_2\cdots\alpha_{L-1}}. \qquad [\text{S13}]$$

We now obtain the final sequence distribution, i.e., $P_{\alpha_1\alpha_2\cdots\alpha_L} = p_{\alpha_1\alpha_2\cdots\alpha_L}(t \to \infty)$ of the mRNA chain for the second-order model, under the following boundary conditions. The initial boundary condition is $p_{\alpha_1\alpha_2}(t=0) = q^{X_1X_2}_{\alpha_1\alpha_2}$, such that $q^{X_1X_2}_{RR} + q^{X_1X_2}_{RW} + q^{X_1X_2}_{WR} + q^{X_1X_2}_{WW} = 1$, $p_{\alpha_1\cdots\alpha_i}(t=0) = 0$ (for $i \geq 3$). We also assume the long time limits $p_{\alpha_1\cdots\alpha_i}(t \to \infty) = 0$ for $2 \leq i < L$. We next integrate the temporal evolutions equations for $p_{\alpha_1\alpha_2\cdots\alpha_i}(t)$, $\forall i = 1$ to L, given by set of Eqs. (S13), to obtain

$$-q_{\alpha_1\alpha_2} = \sum_{\alpha_3} \bar{k}_{\alpha_1\alpha_2\alpha_3} \Gamma_{\alpha_1\alpha_2\alpha_3} + \sum_{\alpha_3}\sum_{\alpha_4} r^{y_4}_{\alpha_3\alpha_4} \Gamma_{\alpha_1\alpha_2\alpha_3\alpha_4} - (k^{y_2y_3}_{\alpha_1\alpha_2 R} + k^{y_2y_3}_{\alpha_1\alpha_2 W}) \Gamma_{\alpha_1\alpha_2},$$

$$0 = \sum_{\alpha_3} \bar{k}_{\alpha_2\alpha_3\alpha_4} \Gamma_{\alpha_1\alpha_2\alpha_3\alpha_4} + k^{y_2y_3}_{\alpha_1\alpha_2\alpha_3} \Gamma_{\alpha_1\alpha_2} + \sum_{\alpha_4}\sum_{\alpha_5} r^{y_4}_{\alpha_3\alpha_4} \Gamma_{\alpha_1\alpha_2\alpha_3\alpha_4} - (k^{y_3y_4}_{\alpha_2\alpha_3 R} + k^{y_3y_4}_{\alpha_2\alpha_3 W} + \bar{k}_{\alpha_1\alpha_2\alpha_3}) \Gamma_{\alpha_1\alpha_2\alpha_3},$$

$$0 = \sum_{\alpha_{i+1}} \bar{k}_{\alpha_{i-1}\alpha_i\alpha_{i+1}} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{i+1}} + k^{y_{i-1}y_i}_{\alpha_{i-2}\alpha_{i-1}\alpha_i} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{i-1}} + \sum_{\alpha_{i+1}}\sum_{\alpha_{i+2}} r^{y_{i+2}}_{\alpha_{i+1}\alpha_{i+2}} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{i+2}}$$
$$- (k^{y_iy_{i+1}}_{\alpha_{i-1}\alpha_i R} + k^{y_iy_{i+1}}_{\alpha_{i-1}\alpha_i W} + \bar{k}_{\alpha_{i-2}\alpha_{i-1}\alpha_i} + r^{y_i}_{\alpha_{i-1}\alpha_i}) \Gamma_{\alpha_1\alpha_2\cdots\alpha_i}, \quad 4 \leq i < L-2,$$

$$0 = \sum_{\alpha_{L-1}} \bar{k}_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{L-1}} + k^{y_{L-3}y_{L-2}}_{\alpha_{L-4}\alpha_{L-3}\alpha_{L-2}} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{L-3}}$$
$$- (k^{y_{L-2}y_{L-1}}_{\alpha_{L-3}\alpha_{L-2} R} + k^{y_{L-2}y_{L-1}}_{\alpha_{L-3}\alpha_{L-2} W} + \bar{k}_{\alpha_{L-4}\alpha_{L-3}\alpha_{L-2}} + r^{y_{L-2}}_{\alpha_{L-3}\alpha_{L-2}}) \Gamma_{\alpha_1\alpha_2\cdots\alpha_{L-2}},$$

$$0 = k^{y_{L-2}y_{L-1}}_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{L-2}} - (k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1} R} + k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1} W} + \bar{k}_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} + r^{y_{L-1}}_{\alpha_{L-2}\alpha_{L-1}}) \Gamma_{\alpha_1\alpha_2\cdots\alpha_{L-1}},$$

$$P_{\alpha_1\alpha_2\cdots\alpha_L} = k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1}\alpha_L} \Gamma_{\alpha_1\alpha_2\cdots\alpha_{L-1}}. \qquad [\text{S14}]$$

Here,

$$\Gamma_{\alpha_1\cdots\alpha_i} = \int_0^\infty p_{\alpha_1\cdots\alpha_i}(t)dt.$$

One can solve the above equations to obtain the following iteration relations:

$$P_{\alpha_1\alpha_2\cdots\alpha_L}(t \to \infty) = (q_{\alpha_1\alpha_2}/g_{\alpha_1\alpha_2})\Pi_{\alpha_1\alpha_2\alpha_3}\Pi_{\alpha_2\alpha_3\alpha_4}\cdots\Pi_{\alpha_i\alpha_{i+1}\alpha_{i+2}}\cdots\Pi_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}} k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1}\alpha_L}$$

$$\Pi_{\alpha_i\alpha_{i+1}\alpha_{i+2}} = k^{y_{i+1}y_{i+2}}_{\alpha_i\alpha_{i+1}\alpha_{i+2}}/(\bar{k}_{\alpha_i\alpha_{i+1}\alpha_{i+2}} + r^{y_{i+1}}_{\alpha_i\alpha_{i+1}} + g_{\alpha_{i+1}\alpha_{i+2}}), 1 \leq i \leq L-2$$

$$g_{\alpha_{i+1}\alpha_{i+2}} = \sum_{\alpha_{i+3}} \left[\Pi_{\alpha_{i+1}\alpha_{i+2}\alpha_{i+3}}(r^{y_{i+2}}_{\alpha_{i+1}\alpha_{i+2}} + g_{\alpha_{i+2}\alpha_{i+3}}) - \sum_{\alpha_{i+4}} \Pi_{\alpha_{i+1}\alpha_{i+2}\alpha_{i+3}}\left(\Pi_{\alpha_{i+2}\alpha_{i+3}\alpha_{i+4}} r^{y_{i+4}}_{\alpha_{i+3}\alpha_{i+4}} \right) \right]$$
$$\text{for} 1 \leq i \leq L-4,$$

$$g_{\alpha_{L-3}\alpha_{L-2}} = \sum_{\alpha_{L-1}} \left[\Pi_{\alpha_{L-3}\alpha_{L-2}\alpha_{L-1}}(r^{y_{L-1}}_{\alpha_{L-2}\alpha_{L-1}} + g_{\alpha_{L-2}\alpha_{L-1}}) \right],$$

$$g_{\alpha_{L-2}\alpha_{L-1}} = \sum_{\alpha_L} k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1}\alpha_L} \qquad [\text{S15}]$$

Equivalently, the above equations can be transformed into an intuitive form of a forward inhomogeneous Markov chain, as below

$$P_{\alpha_1\cdots\alpha_L}(t \to \infty) = q_{\alpha_1\alpha_2} M_{\alpha_1\alpha_2\alpha_3} M_{\alpha_2\alpha_3\alpha_4} \cdots M_{\alpha_i\alpha_{i+1}\alpha_{i+2}} \cdots M_{\alpha_{L-2}\alpha_{L-1}\alpha_L},$$
$$M_{\alpha_i\alpha_{i+1}\alpha_{i+2}} = \Pi_{\alpha_i\alpha_{i+1}\alpha_{i+2}} g_{\alpha_{i+1}\alpha_{i+2}}/g_{\alpha_i\alpha_{i+1}}, 1 \leq i < L-2, \text{ and}$$
$$M_{\alpha_{L-2}\alpha_{L-1}\alpha_L} = k^{y_{L-1}y_L}_{\alpha_{L-2}\alpha_{L-1}\alpha_L}/g_{\alpha_{L-2}\alpha_{L-1}}. \qquad [\text{S16}]$$

Following the approach in Refs. (1, 2), one can obtain the second-order stochastic transfer matrix $\mathbf{M}_{i-2i-1i}$

$$\mathbf{M}_{i-2i-1i} = \begin{pmatrix} M_{R_{i-2}R_{i-1}R_i} & M_{R_{i-2}R_{i-1}W_i} & 0 & 0 \\ 0 & 0 & M_{R_{i-2}W_{i-1}R_i} & M_{R_{i-2}W_{i-1}W_i} \\ M_{W_{i-2}R_{i-1}R_i} & M_{W_{i-2}R_{i-1}W_i} & 0 & 0 \\ 0 & 0 & M_{W_{i-2}W_{i-1}R_i} & M_{W_{i-2}W_{i-1}W_i} \end{pmatrix}. \qquad [\text{S17}]$$

Next, to obtain the approximate expression of the elements of the above matrix, we define the biologically relevant conditions for the advanced model with second-order neighbor effects as intuitive below (1, 3):

1. $k^{xy}_{\alpha\beta R} \gg k^{xy}_{\alpha\beta W}$, implying the addition of R nucleotide is always much faster than that of W, where $\alpha, \beta = R, W$, and $x, y = \text{Pu}, \text{Py}$.

2. $k^{xy}_{\alpha W R}/k^{xy}_{\alpha W W} \gg k^{xy}_{RRR}/k^{xy}_{RRW}$, implying that the successive additions of W are almost inhibited, taken to approximately zero. In fact, $k^{xy}_{\alpha W W} \approx 0$.

3. $k^{xy}_{RRR} > \bar{k}_{RRR}, \bar{k}_{RWR}, r_{RR}, k^{xy}_{WRR}$. This means that successive additions of R always dominate the backward and cleavage rates, guaranteeing the transcriptional elongation.

4. $r^y_{WW}, r^y_{WR}, r^y_{RW} \gg r_{RR}$, which means the terminus containing W s are likely to be proofread faster.

Note that for the model with four components, where we have 3 possible W_i's for every R incorporated, the above state bio-relevant condition for every W_i, for $i = 1, 2, 3$. Under the above-stated biological relevant conditions, one can calculate $M_{\alpha_{i-2}\alpha_{i-1}\alpha_i}$ by the following iterations $g_{\alpha_i\alpha_{i+1}} \simeq k^{y_{i+1}y_{i+2}}_{\alpha_i\alpha_{i+1}R} + k^{y_{i+1}y_{i+2}}_{\alpha_i\alpha_{i+1}W} \simeq k^{y_{i+1}y_{i+2}}_{\alpha_i\alpha_{i+1}R}$, and $\Pi_{\alpha_{i-1}\alpha_i\alpha_{i+1}} = k^{y_iy_{i+1}}_{\alpha_{i-1}\alpha_i\alpha_{i+1}}/(\bar{k}_{\alpha_{i-1}\alpha_i\alpha_{i+1}} + r_{\alpha_i\alpha_{i+1}} + g_{\alpha_i\alpha_{i+1}})$. Also we get $M_{W_{i-2}W_{i-1}W_i}, M_{R_{i-2}W_{i-1}W_i}, M_{W_{i-2}R_{i-1}W_i} \ll M_{R_{i-2}R_{i-1}W_i} \ll 1$. In this approximation, the positional probability $P_{\alpha_{i-1}\alpha_i} = (P_{R_{i-1}R_i}, P_{R_{i-1}W_i}, P_{W_{i-1}R_i}, P_{W_{i-1}W_i})$ represented by the eigenvector, $(1 - 2M_{R_{i-2}R_{i-1}W_i}, M_{R_{i-2}R_{i-1}W_i}, M_{R_{i-2}R_{i-1}W_i}, 0)$, of the matrix $M_{i-2i-1i}$ can be obtained (2). Equivalently, the probability of a wrong nucleotide at position i, η, is given as $P^i_W = P_{R_{i-1}W_i} + P_{W_{i-1}W_i} \approx M_{R_{i-2}R_{i-1}W_i}$, where

$$M_{R_{i-2}R_{i-1}W_i} \simeq \frac{k^{y_{i-1}y_i}_{RRW}}{k^{y_{i-1}y_i}_{RRR}} \bigg/ \left[1 + \frac{\bar{k}_{RRW} + r^{y_i}_{RW}}{k^{y_iy_{i+1}}_{RWR}} \left(1 + \frac{\bar{k}_{RWR} + r^{y_{i+1}}_{WR}}{k^{y_{i+1}y_{i+2}}_{WRR}} \right) \right]. \qquad [\text{S18}]$$

Thus, the approximate analytical expression of the error for the advanced model with second-order neighbor effects is

$$\eta^S_2(y_{-1} - y_0 - y_{+1} - y_{+2}) \approx \frac{k^{y_{-1}y_0}_{RRW}}{k^{y_{-1}y_0}_{RRR}} \bigg/ \left[1 + \frac{\bar{k}_{RRW} + r^{y_0}_{RW}}{k^{y_0y_{+1}}_{RWR}} \left(1 + \frac{\bar{k}_{RWR} + r^{y_{+1}}_{WR}}{k^{y_{+1}y_{+2}}_{WRR}} \right) \right]. \qquad [\text{S19}]$$

Grouping of experimental data for 64 cases based on nucleotides at -1,0,1 positions into 4 cases. In this section, we describe how the experimental error rates from Ref. (4) for nucleotides A, C, G, and U at position 0 were systematically grouped based on the identity of the nucleotides at the upstream (-1) and downstream (+1) positions. The goal of this procedure is to categorize the error rates into four distinct cases, depending on whether the nucleotide at position 0 is a purine (A, G) or a pyrimidine (C, U), and whether the nucleotide at position +1 is a purine (Pu) or pyrimidine (Py). We first collected the error rates for each nucleotide i at position 0, considering every possible nucleotide j at position +1, while disregarding the identity of the nucleotide at position -1, for $i, j \in \{$ A, C, G, U $\}$. This resulted in 16 distinct cases, accounting for all combinations of the four nucleotides at positions 0 and +1. Next, we grouped the data based on whether the nucleotide at position +1 was a purine (A, G) or a pyrimidine (C, U). For each nucleotide i at position 0, we separately considered its error rates when the downstream nucleotide was either a purine or a pyrimidine, reducing the number of cases from 16 to 8. Finally, we categorized the data based on the type of nucleotide at position 0. This resulted in 4 cases: Pu-Pu, Pu-Py, Py-Pu, Py-Py occupying 0, +1 positions on the mRNA sequence.

Bioinformatics analysis of the experimental data for error rate depending on the bases at one upstream and two downstream positions. We group the experimental error rates from Ref. (4) for nucleotides A, C, G, and U at position 0 based on the identity of the nucleotides at the upstream (-1), first downstream (+1), and second downstream (+2) positions. This grouping will categorize the error rates for 256 cases arising from nucleotide identity at (-1,0,+1,+2) positions into 16 distinct cases. These cases arise depending on whether the nucleotides at the positions (-1,0,+1,+2) are purine (A, G) or pyrimidines (C, U). We first collected the error rates for each nucleotide at position 0, considering whether the nucleotide at position -1 is a purine or a pyrimidine, reducing the cases to 128. We then grouped the data based on whether the nucleotide at +1 is purine or pyrimidine and then based on whether the nucleotide at +2 position is purine or pyrimidine. Finally, we categorized the data based on the type of nucleotide at position 0, resulting in error rates for 16 cases depending on whether there is a purine or a pyrimidine at -1, 0, +1, and +2.

Monte Carlo Simulations. To complement the numerical results for our theoretical model for two elements, purine (Pu) and pyrimidine (Py), we perform extensive Monte Carlo simulations (MCS). We employed the Gillespie algorithm (5) to simulate the growth of mRNA chain for a given DNA template sequence of length 1000. The mRNA sequence for the first-order neighbor effects starts with a monomer as an initial seed and grows according to the kinetic rates given in Table S1. For the process with second-order neighbor effects, the mRNA sequence is extended from a dimer as an initial seed. The probability of errors is directly inferred from the outcomes of 1000 simulations. A length of 1000 monomers is chosen to avoid any edge effects in computing the probability of errors.

A. References.

1. QS Li, PD Zheng, YG Shu, ZC Ou-Yang, M Li, Template-specific fidelity of dna replication with high-order neighbor effects: A first-passage approach. *Phys. Rev. E* **100**, 012131 (2019).
2. QS Li, YG Shu, ZC Ou-Yang, M Li, Kinetic assays of dna polymerase fidelity: A theoretical perspective beyond michaelis-menten kinetics. *Phys. Rev. E* **104**, 014408 (2021).
3. T Midha, AB Kolomeisky, OA Igoshin, Linear-decoupling enables accurate speed and accuracy predictions for copolymerization processes. *The J. Phys. Chem. Lett.* **15**, 9361–9368 (2024).
4. C Chung, et al., The fidelity of transcription in human cells. *Proc. Natl. Acad. Sci.* **120**, e2210038120 (2023).
5. DT Gillespie, Exact stochastic simulation of coupled chemical reactions. *The journal physical chemistry* **81**, 2340–2361 (1977).

Fig. S1. Schematic diagram of the two-elements first-order biophysical model for transcription elongation process with dinucleotide cleavage and (A) without template-dependent kinetics (B) with template-dependent kinetics. Here, $i, j = R, W$, i denotes the initial monomer in the mRNA sequence, and y denotes the nature, i.e., Pu or Py of the current incorporating nucleotide j. The rate $k_{ij}^y = k_{ij}$, if y = Pu, and $k_{ij}^y = \gamma k_{ij}$, if y = Py.

Fig. S2. Comparison of numerical error from Eq. (4), methods (main manuscript) with the approximate analytical error rate from Eq. (S4) for two dimers: Pu-Pu, Pu-Py at the current and downstream positions of mRNA varying as a function of Py to Pu incorporation rate ratio, γ. Symbols and dashed lines respectively denote the numerical and approximate error result.

Fig. S3. Fitting (shaded region) of theoretical errors (red bars) from the model with first-order neighbor effects and equal cleavage rates for purine and pyrimidines to the experimental data (blue bars). Theoretical errors (red bars) for the unfitted (unshaded region) dimers not aligned with the corresponding experimental data (blue bars). Error bars denote the standard error mean for the experimental data for the six samples (4).

Fig. S4. Schematic diagram of the two-element second-order biophysical model for transcription elongation process with dinucleotide cleavage and (A) without template-dependent kinetics (B) with template-dependent kinetics, where ij denotes the starting monomer. In k_{ijk}^{xy}, $i, j, k = R, W$, x denotes the nature, i.e., Pu or Py, of the previous incorporated nucleotide j, and y denotes the nature, i.e., Pu or Py, of the current incorporating nucleotide k. For no upstream effects, the rate $k_{ijk}^{xy} = k_{ijk}$, if y = Pu, and $k_{ijk}^{xy} = \gamma k_{ijk}$, if y = Py. For the upstream effects, the rate $k_{ijk}^{xy} = k_{ijk}$, if y = Pu, x = Pu, $k_{ijk}^{xy} = \gamma k_{ijk}$, if y = Py, x = Pu, $k_{ijk}^{xy} = \psi k_{ijk}$, if y = Pu, x = Py, $k_{ijk}^{xy} = \gamma \psi k_{ijk}$, if y = Py, x = Py.

Fig. S5. Variation of error from the two-element advanced model with a Pu at position 0 and four possible dimers occupying positions +1 and +2 on mRNA as a function of the incorporation factor, γ.

Fig. S6. (A) Experimental error rates at a current position 0 depending on either purine or pyrimidine at positions -1, 0, +1, +2, obtained from the bioinformatics analysis of experimental data from Ref.(4). Error bars represent the SEM. (B) Theoretical error rates $\eta_2^S(-1, 0, +1, +2)$ at a current position 0 depending on either purine or pyrimidine at positions -1, 0, +1, +2, obtained from the advanced model with second-order neighbor effects, two elements and no upstream nucleotide identity effect on the incorporation rates.

Fig. S7. (A) Variation in the ratio of the theoretical error rate η_2^S with Pu at +3 position to that of Py at +3 position $\eta(\text{Pu}@+3)/(\eta_{\text{Py}}@+3)$ with either Pu or Py at the current position, 0. (B) Variation in the ratio of the experimental error rate with Pu at +3 position to that of Py at +3 position.

Fig. S8. Fitting of theoretical error rates $\eta_4^S(0,+1,+2)$ for the second-order biophysical model with four elements to the corresponding experimental data with cleavage rates depending on the nature of misincorporated nucleotide. Inset shows the average percentage of error for fitted, experimental, and unfitted data.

Table S1. Kinetic parameters for the first-order biophysical model with two elements: Pu and Py.

Parameter	Value s^{-1}	Whether fitted or fixed
k_{RR}^{Pu}	100	Fitted
k_{RR}^{Py}	$\gamma \times 100$	Fitted γ
k_{RW}^{Pu}	.007	Fixed
k_{RW}^{Py}	$\gamma \times .007$	Fixed
k_{WR}^{Pu}	.01	Fixed
k_{WR}^{Py}	$\gamma \times .01$	Fixed
k_{WW}^{Pu}	10^{-6}	Fixed
k_{WW}^{Py}	$\gamma \times 10^{-6}$	Fixed
$\bar{k}_{RR}$	5	Fixed
$\bar{k}_{RW}$	9.65×10^{-6}	Fixed
$\bar{k}_{WR}$	.00105	Fixed
r_{RR}	.001	Fixed
$r_{RW}^{\mathrm{Pu}}, r_{WR}^{\mathrm{Pu}}, r_{WW}^{\mathrm{Pu}}$	.038	Fixed
$r_{RW}^{\mathrm{Py}}, r_{WR}^{\mathrm{Py}}, r_{WW}^{\mathrm{Py}}$	$\phi \times .038$	Fitted ϕ

Table S2. Kinetic parameters for the advanced model with second-order neighbor effects and with two elements. For the upstream effect, the incorporation rate k_{iRk}^{xy} gets multiplied by the factor ψ, when previously incorporated nucleotide x is Py.

Parameter	Value s^{-1}	Fitted or Fixed
$k_{RRR}^{x\mathrm{Pu}}$	100	Fitted
$k_{RRR}^{x\mathrm{Py}}$	$\gamma \times 100$	Fitted γ
$k_{RRW}^{x\mathrm{Pu}}$	.007	Fixed
$k_{RRW}^{x\mathrm{Py}}$	$\gamma \times .007$	Fixed
$k_{RWR}^{x\mathrm{Pu}}$	0.01	Fixed
$k_{RWR}^{x\mathrm{Py}}$	$\gamma \times 0.01$	Fixed
$k_{WRR}^{x\mathrm{Pu}}$	0.2	Fixed
$k_{WRR}^{x\mathrm{Py}}$	$\gamma \times 0.2$	Fixed
$\bar{k}_{RRR}$	5	Fixed
$\bar{k}_{RRW}$	9.65×10^{-6}	Fixed
$\bar{k}_{RWR}$	.00105	Fixed
$\bar{k}_{WRR}$	.000105	Fixed
r_{RR}	.001	Fixed
$r_{RW}^{\mathrm{Pu}}, r_{WR}^{\mathrm{Pu}}, r_{WW}^{\mathrm{Pu}}$	.038	Fixed
$r_{RW}^{\mathrm{Pu}}, r_{WR}^{\mathrm{Pu}}, r_{WW}^{\mathrm{Pu}}$	.038	Fitted

Table S3. Fixed values of the kinetic parameters for the computation of approximate theoretical error for the theoretical model with second-order neighbor effects and with four elements.

Parameter	Values^{-1}	Role
$k_{RW_jR}^{\mathrm{A}}$	.01	Addition rate of correct nucleotide A after any misincorporation at the previous position
$k_{RW_jR}^{\mathrm{U}}$	$\gamma_1 \times .02$	Addition of correct nucleotide U after any misincorporation at the previous position
k_{RWR}^{C}	$\gamma_2 \times .02$	Addition of correct nucleotide C after any misincorporation at the previous position
k_{RWR}^{G}	$\gamma_3 \times .02$	Addition of correct nucleotide G after any misincorporation at the previous position
$\bar{k}_{RWR}$	.00105	Removal rate of any correct nucleotide after any misincorporation
k_{WRR}^{A}	.2	Addition of correct nucleotide A after the dimer WR occupying the two previous position
k_{WRR}^{U}	$\gamma_1 \times .2$	Addition of correct nucleotide U after the dimer WR occupying the two previous position
k_{WRR}^{C}	$\gamma_2 \times .2$	Addition of correct nucleotide C after the dimer WR occupying the two previous position
k_{WRR}^{G}	$\gamma_3 \times .2$	Addition of correct nucleotide G after the dimer WR occupying the two previous position
$k_{RRW_j}^{\mathrm{U}}$	.0004	Addition rate of any misincorporation instead of U
$k_{RRW_j}^{\mathrm{A}}$	.002	Addition rate of any misincorporation instead of A
$k_{RRW_j}^{\mathrm{G}}$	.0008	Addition rate of any misincorporation instead of G
$k_{RRW_j}^{\mathrm{C}}$	.001	Addition rate of any misincorporation instead of C
$\bar{k}_{RRW}$	9.65×10^{-6}	Removal rate of any misincorporation

Table S4. Optimal kinetic parameters for the theoretical model with second-order neighbor effects and four elements obtained from fitting experimental data from Ref. (4).

Parameter	Value	Role
k_{RRR}^{A}	$80\ s^{-1}$	Incorporation rate of A on mRNA given T on the DNA template
γ_1	0.58	Incorporation rate ratio for U:A
γ_2	0.63	Incorporation rate ratio for C:A
γ_3	0.94	Incorporation rate ratio for G:A
r_{A}	$.012 s^{-1}$	Dinucleotide cleavage rate of misincorporations instead of correct A
r_{U}	$.06 s^{-1}$	Dinucleotide cleavage rate of misincorporations instead of correct U
r_{C}	$.038 s^{-1}$	Dinucleotide cleavage rate of misincorporations instead of correct C
r_{G}	$.01 s^{-1}$	Dinucleotide cleavage rate of misincorporations instead of correct G